Discovery of a novel simian pegivirus in common marmosets (*Callithrix jacchus*) with lymphocytic enteritis

Heffron AS[1], Lauck M[1], Somsen ED[1], Townsend EC[1], Bailey AL[2], Eickhoff J[3], Newman CM[1], Kuhn J[4], Mejia A[5], Simmons HA[5], O'Connor DH[1].

Affiliations

[1] Department of Pathology and Laboratory Medicine, School of Medicine and Public Health, University of Wisconsin-Madison, Madison, Wisconsin, United States of America

[2] Department of Pathology and Immunology, Washington University School of Medicine, St. Louis, Missouri, United States of America

[3] Department of Biostatistics & Medical Informatics, University of Wisconsin-Madison, Madison, WI, United States of America

[4] Integrated Research Facility at Fort Detrick, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Fort Detrick, Frederick, Maryland, United States of America

[5] Wisconsin National Primate Research Center, University of Wisconsin-Madison, Madison, Wisconsin, United States of America

Abstract

From 2010 to 2015, 73 common marmosets (*Callithrix jacchus*) housed at the Wisconsin National Primate Research Center (WNPRC) were diagnosed postmortem with lymphocytic enteritis. We used unbiased deep-sequencing to screen the blood of deceased enteritis-positive marmosets for the presence of RNA viruses. In five out of eight marmosets found to have lymphocytic enteritis, we discovered a novel pegivirus not present in ten subsequently deep-sequenced healthy marmosets. The novel virus, which we have named Southwest bike trail virus (SOBV), is most closely related to a strain of simian pegivirus A (68% nucleotide identity) that was previously isolated from a three-striped night monkey (*Aotus trivirgatus*). To determine the prevalence of this novel virus within the WNPRC marmoset colony, we screened 146 living animals and found an overall prevalence of 34% (50/146). Over the next four years, 85 of the 146 screened marmosets were examined histologically for lymphocytic enteritis. Out of these 85 animals, 27 SOBV-infected common marmosets had developed lymphocytic enteritis, compared to 42 uninfected common marmosets, indicating no association between this virus and development of enteritis ($p=0.820$). The novel pegivirus was also found in 2 of 32 (6.25%) healthy marmosets screened while in quarantine during the transfer from the New England Primate Research Center to the WNPRC.

Importance

Common marmosets (*Callithrix jacchus*) are a valuable model species. We discovered two variants of a novel pegivirus, which we named the Southwest bike trail virus (SOBV), in common marmosets which had postmortem histologic diagnosis of lymphocytic enteritis. We screened 146 live healthy marmosets in the Wisconsin National Primate Research Center colony and found 34% (50/146) of the animals were infected. SOBV was also present in 2 of 32 (6.25%) healthy marmosets from the New England Primate Research Center. These findings have implications for animal studies in which infection-free animals are desired, and they demonstrate the need for further investigations to increase understanding of this genus of viruses.

Introduction

Common marmosets (*Callithrix jacchus*) are a valuable model species due to their small body size, communal monogamous familial behavior, birth of hematopoietic chimeric litters, short parturition intervals, and status as members of a non-endangered primate species.[1-4] The utility of common marmosets in research has resulted in a recent increase in demand for these animals.[5] The Wisconsin National Primate Research Center (WNPRC) in Wisconsin, Madison, USA, houses a common marmoset colony typically consisting of about 240 common marmosets, which are used by researchers at the University of Wisconsin-Madison for groundbreaking research in neurological, neurobehavioral, and pharmacologic studies, and many others.[6-16]

From 2010 to 2015, 73 common marmosets housed at the WNPRC were euthanized due experimental end point, chronic, intractable diarrhea, or chronic severe weight loss, underwent necropsy with histology and were diagnosed with lymphocytic enteritis. Beyond the regrettable loss of animal life, common marmoset demise due to enteritis is harmful to both colony success and to the scientific studies to which these animals are assigned. Though lymphocytic enteritis is one of the most common causes of death in captive common marmosets,[17-22] the epizootic at WNPRC was associated with an unusually high disease incidence, prompting investigations into a possible infectious contributor. Unbiased deep-sequencing led to the discovery of two closely related variants of a novel pegivirus, most closely related to a variant of simian pegivirus A (SPgV-A) previously isolated from a three-striped night monkey (*Aotus trivirgatus*). This novel pegivirus was present in a subset of deceased common marmosets diagnosed postmortem with lymphocytic enteritis and not present in matched healthy controls. Pegiviruses, members of genus *Pegivirus (Amarillovirales: Flaviviridae)*, are ubiquitous in animal populations,[23-34] but their biological consequences are poorly understood. Given the importance of common marmosets as a model species and the disease burden caused by lymphocytic enteritis, we set out to characterize the possible link between these new viruses and the disease state.

Here, we report the discovery of two variants of a novel pegivirus in a captive common marmoset colony. We establish phylogenetic relationships with other known pegiviruses. Since this virus was discovered in common marmosets with lymphocytic enteritis and was absent in healthy controls, we measured the prevalence of the virus in the colony. We investigated the potential association between the virus and the occurrence of lymphocytic enteritis. We ultimately found no statistically significant association between infection status and disease. Still, we did find the virus to be highly prevalent (34%) in our colony, less prevalent (6.25%) in a comparison colony, and significantly associated with increasing age (p=0.03237). These findings have implications for animal studies in which infection-free animals are desired, and

83 they demonstrate the need for further investigations to increase understanding of this genus of
84 viruses.

Results

Captive common marmosets harbor a novel pegivirus

To examine the etiology of the unusually high rate of lymphocytic enteritis in deceased WNPRC
common marmosets, banked plasma samples from eight common marmosets diagnosed with
lymphocytic enteritis and from 10 healthy, live common marmosets to be used as controls were
screened by deep sequencing for the presence of DNA and RNA viruses. Five of eight
deceased marmosets with lymphocytic enteritis were infected with the previously undocumented
pegivirus. We propose this novel virus (BioProject accession number PRJNA613737), be
formally named the Southwest bike trail virus (SOBV). The ten healthy common marmoset
controls were found to be negative for this virus.
95
SOBV consists of a 9.8-kb-long contig that is highly similar to the genome of simian pegivirus A
(SPgV A) trivirgatus, a simian pegivirus previously discovered in a three-striped night monkey
(*Aotus trivirgatus*)[26] (Figure 1), with 68% nucleotide identity across the coding sequence when
aligned using ClustalW with an IUB cost matrix (gap extension cost, 6.66; gap open cost,
15.00). Four of the five marmosets infected with SOBV had variants of the virus having 98-99%
sequence identity, while one marmoset was infected with a variant with 88% sequence identity
to the others. We have named these variants SOBV-1 and -2.
103
104 Pairwise comparisons of nucleotide identity across the entire coding region further illustrate the
105 similarity of SOBV-1 and SOBV-2 and the divergence between these novel virus strains and the
106 next most closely-related viruses (Figure 2, Figure 3), most of which were simian pegiviruses.
107 Interestingly, a pegivirus isolate found in a bat, BPgV 737,[35] also shared a high degree of
108 similarity with the novel pegivirus and with similar simian pegiviruses.
109

Figure 1. A phylogenetic tree of newly discovered pegivirus Southwest bike trail virus (SOBV
strains 1 and 2) shows it is most closely related to pegiviruses found in other New World
monkeys. We generated maximum likelihood trees using MEGA6.06 (1,000 bootstrap
replicates, GTR+I+γ model) from codon-based alignments (via MAFFT); Bootstrap values of
less than 70 are omitted.
Abbreviations: HPgV = human pegivirus; SPgV = simian pegivirus; GBV-C = G.B. virus C; BPgV
= bat pegivirus; EqPgV = equine pegivirus; HePegi = hepegivirus; RPgV = rodent pegivirus
118

119
Figure 2. Sliding window similarity plots[36] show the relatedness of the amino acid sequences of
SOBV-2 and other closely related pegiviruses to SOBV-1. Dashed vertical lines indicate the
putative approximate start positions of inferred viral proteins, from left to right: E1, E2, NS2,
NS3, NS4A, NS4B, NS5A, and NS5B.[37]
Abbreviations: SPgV-A nigri = GBV-A-like virus recovered from *Saguinus nigricollis*; SPgV-A tri
= GBV-A-like virus recovered from *Aotus trivirgatus*; SPgV-A mx = GBV-A-like virus recovered
from *Saguinus mystax*; SPgV-A lab = GBV-A-like virus recovered from *Saguinus labiatus*; BPgV
= bat pegivirus recovered from *Eidolon helvum*
128

129
130 **Figure 3.** Sequence identity matrix based on amino acid alignment of the newly discovered
131 SOBV-1 and SOBV-2 (red box) compared to members of the 11 recognized pegivirus species
132 and of one* proposed species.[37]
133 *The classification of dolphin into species "Pegivirus L" has been suggested.[38]

Novel pegivirus infects up to 35% of a captive common marmoset colony

Having identified the novel pegivirus, we sought to determine its prevalence within the WNPRC common marmoset colony. We developed an RT-PCR assay to detect a conserved region of the putative helicase protein of SOBV and used this to screen plasma collected from 146 healthy live common marmosets in the WNPRC colony, confirming results through deep-sequencing of the amplicons. At the time of the initial screening in March–April 2014, 50 of the 146 (34.25%) healthy screened animals tested positive for SOBV. Nineteen of 60 females (31.67%) and 31 of 86 males (36.05%) tested positive at the time of screening. Sex was not associated with the likelihood of infection using univariate logistic regression (p=0.5834). Age at the time of screening was mildly associated with the likelihood of infection (p=0.03237), with the likelihood of positivity increasing with age (Figure 4).

WNPRC common marmosets screened for SOBV			
Age (years) at time of screening	Total number common marmosets	Number common marmosets infected	Number common marmosets non-infected
1	20	6	14
2	24	5	19
3	18	6	12
4	14	3	11
5	22	8	14
6	7	2	5
7	14	9	5
8	13	3	10
9	5	4	1
10	6	2	4
11	2	1	1
12	0	0	0
13	1	1	0

Figure 4. Prevalence of infection with Southwest bike trail virus (SOBV) in common marmosets at the WNPRC increases with age. One hundred forty-six live, healthy common marmosets in the WNPRC captive common marmoset colony were screened for SOBV using RT-PCR and deep sequencing methods. The likelihood of infection with these viruses was significantly statistically associated with increasing age (p=0.03237) using univariate logistic regression.

In November 2014, 82 common marmosets were transferred from the New England Primate Research Center (NEPRC) to the WNPRC. Samples from 32 NEPRC animals were collected while the animals were in quarantine. Two (6.25%) were found to be infected with SOBV when screened by RT-PCR in quarantine.

Presence of novel pegivirus is not statistically significantly associated with common marmoset lymphocytic enteritis

We next sought to determine whether infected animals were more likely to develop lymphocytic enteritis. Typical enteric architecture consists of slender, often branching villi, with short intestinal glands, small numbers of lymphocytes in the lamina propria, and prominent B cell aggregates dispersed throughout the length of the intestines (Figure 5, control). Lymphocytic enteritis was diagnosed as a disruption of this architecture, with lymphocytic infiltration that expands the lamina propria, resulting in widening and shortening of villi and hyperplasia of crypt epithelium (Figure 5, E1-E3). Cases varied in severity, with mild cases showing only slight

expansion of the lamina propria and advanced cases showing complete loss of villus architecture due to infiltration of the lamina propria with large numbers of CD3-positive lymphocytes. Eighty-five of the live WNPRC animals initially screened for SOBV in 2014 were euthanized for experimental end points or clinical illness between their screening and May 3, 2019. Sixty-nine (81.18%) of these animals were diagnosed by postmortem histological analysis with lymphocytic enteritis of the small intestine, or large intestine, or both. Two animals were removed from this analysis due to confounding factors (one animal had severe tissue autolysis, and the other animal had B cell lymphoma of the small and large intestines).

Figure 5. Representative photomicrographs show lymphocytic enteritis distorting normal intestinal structures in the duodenum, jejunum, and colon of common marmosets. Histology performed upon intestinal samples from 85 common marmosets. All intestinal sections were stained with hematoxylin and eosin (H&E). Intestinal sections from animals with enteritis were stained using B cell-specific and T cell-specific staining procedures (immunohistochemistry) with monoclonal antibodies to CD20 or CD79 (B cell markers) and CD3 (T cell marker), respectively.

186
187 Pegivirus infection was not found to be associated with an increased likelihood of developing
188 lymphocytic enteritis in the large intestine (p=0.196), small intestines (p=0.779), both small and
189 large intestines (p=0.0798), or either the large or small intestines (p=0.820) (Figure 6).
190

191
Figure 6. Infection with Southwest bike trail virus (SOBV) is not associated with the likelihood of
developing lymphocytic enteritis. Eighty-five common marmosets at the WNPRC, which had
been previously screened for SOBV by RT-PCR or deep-sequencing of plasma samples, were
examined postmortem for histological evidence of lymphocytic enteritis. Pegivirus infection was
not found to be associated with an increased likelihood of developing lymphocytic enteritis in the
large intestine (p=0.196), small intestines (p=0.779), both small and large intestines (p=0.0798),
or either the large or small intestines (p=0.820) using univariate logistic regression.

Discussion

199
We describe the discovery of a novel simian pegivirus first identified in common marmosets
diagnosed with lymphocytic enteritis. We show this pegivirus was prevalent in our colony during
a period of increased incidence of lymphocytic enteritis and that it was less prevalent in a
similar, healthy colony. The novel virus was not significantly associated with the likelihood of
developing lymphocytic enteritis, though prevalence of the virus increased with increasing age
in the common marmoset. With an average prevalence of 34%, SOBV appears common
throughout the WNPRC common marmoset colony.
207
Pegiviruses, the members of genus *Pegivirus (Amarillovirales*: *Flaviviridae)*, have single-
stranded, positive-sense RNA genomes and produce enveloped virions.[39] The first members of
the genus were identified about twenty years ago,[40] and since that time pegiviruses have been
found in many animal populations.[23-34,38] Pegiviruses have never been shown to be causative
agents of any disease or alteration in physiology,[41-58] though human pegivirus (HPgV; species
Pegivirus A) has been linked to incidence of various types of lymphoma,[59-66] though this remains
controversial.[67-71] HPgV has, intriguingly, been linked to improved outcomes in coinfection with
HIV-1[72-88] and Ebola virus,[87] leading some to propose the use of HPgV as a biotherapy.[89]
216
It is not known whether these viruses are unique to the captive common marmoset populations
or whether they are also present in wild populations.[90] Other pegiviruses have been discovered

in wild common marmosets in the 1990s,[91] but their prevalence has never been examined. The
prevalence of these viruses in our captive common marmoset population was quite high
compared to the best-studied pegivirus, HPgV, which is found in about 1–4% of human
populations.[92-99] The novel pegiviruses were most similar to a pegivirus discovered in a three-
striped night monkey (*Aotus trivirgatus*),[26] a monkey used in malaria research at other primate
research facilities.[100-102] Interestingly, SOBV was highly similar to several variants of a bat
pegivirus isolated from African straw-coloured fruit bats (*Eidolon helvum*). This was somewhat
surprising given that the bat pegivirus isolates fall into Pegivirus species B. However, bat
pegiviruses have been found to be prevalent (up to 5%) among wild bats around the globe and
to span the full diversity of pegivirus species.[35] Our findings suggest common ancestry or an
otherwise close relationship between these variants.
230
The routes of transmission of SOBV and of other simian pegiviruses have not been examined.
Human pegivirus transmission has been extensively studied and is known to be occur efficiently
through blood products or dialysis,[43,103-107] intravenous drug use and needle sticks,[104,108-110]
sexual intercourse,[104,108,111,112] and from mother to infant.[104,113-117] In humans, HPgV can replicate
at high titers in a host for more than a decade[103,118,119] with an unusually low mutation rate
compared to other RNA viruses,[120,121] with the host's production of anti-pegivirus antibodies
resulting in protective sterilizing immunity once the immune system recognizes and clears the
virus.[73,119] The duration of pegivirus infections in humans indicates SOBV may be similarly long-
lasting commensals of common marmosets, potentially impacting common marmoset lives in
unknown ways. Defining mechanisms of transmission will be important in preventing infection
and thereby allowing the study of these viruses' effects.
242
The high prevalence of this virus at the WNPRC raises essential considerations about potential
effects on common marmoset experiments. Facilities working with common marmosets should
prescreen the animals to establish the pegivirus infection status of animals in research. Future
investigations, perhaps involving the isolation of common marmosets for years at a time to
follow the natural history of long-term pegivirus infection in these animals, could examine the
long-term effects of infecting common marmosets with SOBV.
249
This study has several limitations. First, this study was observational in nature and could not
examine a causal link between viral positivity and the development of enteritis. Definitive
establishment of causation would require demonstrating that animals infected experimentally
develop lymphocytic enteritis. Second, not all of the animals initially screened were deceased at
the time of this analysis and future necropsies of these animals may contribute additional data
concerning the likelihood of enteritis development. Third, we were unable to culture this virus;
difficulties in culturing pegiviruses have been documented.[37,122] Finally, some animals in this
study may have cleared the virus before the samples we tested were collected. Consequently,
these animals could have been mistakenly classified as virus-naïve; others may have acquired
the virus after our initial screening. Serial testing of animals would have counteracted this
problem; however, this was a retrospective study, so serial testing was impossible.
Development of a SOBV-specific ELISA or other serodiagnosis tools would enable deeper
appropriate analyses of SOBV infection rates both prospectively and retrospectively.
263
264 In summary, this work describes the discovery of a novel simian pegivirus and investigates its
265 relationship with a widespread and devastating cause of common marmoset mortality. Our
266 study lays the groundwork for the future development of a nonhuman primate model system
267 using this natural infection as a potential model for studying the mechanisms of these enigmatic
268 viruses and providing a greater understanding of their genus as a whole.
269

Materials and methods

Animals

All animals in this study were common marmosets (*Callithrix jacchus* Linnaeus, 1758) housed at the Wisconsin National Primate Research Center (WNPRC) in Madison, WI, USA. The common marmoset colony at the WNPRC was established in 1960. The original animals were imported from northeastern Brazil, with the final importation occurring in the early 1970s. The average yearly population of the colony each year from 2010 to 2019 was approximately 240 animals, all of which were born in captivity. WNPRC animals screened were 41% (60 animals) female and 59% (86 animals) male. Age at the time of screening ranged from 0.82–12.82 years (mean 4.65+/-2.83 years, median 4.26 years).

The New England Primate Research Center (NEPRC), Southborough, MA, USA, was closed in 2015, resulting in a transfer of 82 common marmosets to WNPRC before closure in November 2014. Plasma samples were collected from these animals upon their arrival at WNPRC (November–December 2014) while quarantined in a separate building and location from the WNPRC marmoset colony. In the population initially from the NEPRC, 45 (55%) of the screened animals were female, and 37 (45%) were male. Age at the time of screening ranged from 0.65–10.66 years (mean 3.74+/-2.60, median 2.51 years) in this population.

Ethics

All common marmosets were cared for by WNPRC staff according to the regulations and guidelines outlined in the National Research Council's Guide for the Care and Use of Laboratory Animals, the Animal Welfare Act, the Public Health Service Policy on the Humane Care and Use of Laboratory Animals, and the recommendations of the Weatherall report (https://royalsociety.org/topics-policy/publications/2006/weatherall-report/). Per WNPRC standard operating procedures for animals assigned to protocols involving the experimental inoculation of infectious pathogens, environmental enhancement included constant visual, auditory, and olfactory contact with conspecifics, the provision of feeding devices that inspire foraging behavior, the provision and rotation of novel manipulanda, and enclosure furniture (i.e., perches, shelves). The common marmosets were housed socially in enclosures measuring 0.6m D × 0.9m W × 1.8 m H or 0.6m D × 1.2m W × 1.8 m H. The WNPRC maintains an exemption from the USDA for these enclosures as they do not meet the Animal Welfare Act regulations for floor space but greatly exceed height requirements as the species are arboreal. This study was approved by the University of Wisconsin-Madison College of Letters and Sciences and Vice Chancellor for Research and Graduate Education Centers Institutional Animal Care and Use Committee (animal protocol numbers G005401 and G005443).

Unbiased deep-sequencing

Samples from 18 common marmosets (8 deceased common marmosets diagnosed with lymphocytic enteritis through necropsy and 10 live, healthy common marmosets) from the WNPRC and 12 common marmosets (all live and healthy) from the NEPRC were screened for the presence of viruses using unbiased deep-sequencing. The live WNPRC common marmosets and the live NEPRC common marmosets were selected randomly for deep-sequencing.

DNA and RNA were isolated from plasma. Common marmoset plasma (1 ml/animal) was centrifuged at 5,000 x *g* for 5 min at 4°C. Supernatants were removed and filtered through a 0.45-µm filter, then centrifuged at maximum speed (20,817 g) for 5 min at 4°C. Supernatants were removed and incubated for 90 min at 37°C with a DNA/RNA digest cocktail consisting of 4 µl DNAfree DNAse (0.04 U/µl; Ambion, Austin, TX, USA), 6 µl Baseline Zero DNAse (0.1 U/µl, Epicentre Technologies, Madison, WI, USA), 1 µl Benzonase (1 U/µl, Sigma-Adrich, St. Louis, MO, USA), and 12 µl DNAse 10x buffer. Viral nucleic acids were then isolated using the Qiagen QIAamp MinElute Virus Spin Kit without the use of AW1 buffer or carrier RNA (Qiagen, Valencia, CA, USA). Random hexamers were used to prime cDNA synthesis (Life Technologies, Grand Island, NY, USA), followed by DNA purification using Ampure XP beads, as previously described[123,124]. Deep-sequencing libraries were prepared using the Nextera XT DNA Library Prep Kit (Illumina, San Diego, CA, USA) and sequenced on MiSeq (Illumina).

Viral sequence and phylogenetic analysis

Sequence data were analyzed using CLC Genomics Workbench 5.5 (CLC bio, Aarhus, Denmark). Low-quality reads (Phred <Q30) and short reads (<100 bp) were removed with CLC Genomics Workbench 7.1 (CLC bio, Aarhus, Denmark), and the remaining reads were assembled *de novo* using the MEGAHIT assembler. Assembled contiguous sequences (contigs) and singleton reads were queried against GenBank database nt using the basic local alignment search tools blastn. Nucleotide sequences were codon aligned individually for all known pegiviruses with complete genomes using ClustalW in the alignment editor program in MEGA v5.10 and edited manually. The best-fitting distance model of nucleotide substitution for each alignment was inferred using the maximum likelihood (ML) method with goodness of fit measured by the Bayesian information criterion in MEGA6.06. The best-fitting nucleotide substitution model for the phylogenetic alignments was inferred to be the GTR model with discrete gamma and invariant among-site rate variation.

Protein family analysis and putative protein predictions were performed using Pfam (http://pfam.xfam.org/). The nucleotide similarity of the novel pegivirus with related pegivirus lineages was determined across the polyprotein using SimPlot v3.5.1[36] following TranslatorX alignment (MAAFT) without Gblocks cleaning.

Screening for SOBV by RT-PCR

Plasma samples from 136 healthy WNPRC common marmosets were screened specifically for SOBV by RT-PCR. Twenty plasma samples collected from NEPRC animals were likewise screened by RT-PCR as they contained less than 100 µl.

Screening of the 136 animals was performed with samples from animals positive for SOBV by deep-sequencing as positive controls. RNA was isolated from 200–500 µl of plasma using the QIAamp Viral RNA Mini Kit (Qiagen). A primer set (forward primer: GGTGGTCCACGAGTGATGA; reverse primer: AGGTACCGCCTGGGGTTAG) targeting a region of the viral helicase which was conserved among the animals initially positive by deep sequencing was designed, resulting in a 615-bp amplicon. Viral RNA was reverse-transcribed and amplified using the SuperScript III High Fidelity One-Step RT-PCR kit (Invitrogen, Life Technologies, Carlsbad, CA, USA). The reverse transcription-PCR conditions were as follows: 50°C for 30 min; 94°C for 2 min; 40 cycles of 94°C for 15 s, 55°C for 30 s, and 68°C for 1 min; and 68°C for 5 min. Following PCR, amplicons were purified from excised gel slices (1% agarose) using the Qiagen MinElute Gel Extraction kit (Qiagen). Each amplicon was quantified

359 using Quant-IT HS reagents (Invitrogen), and approximately 1 ng of each was used in a
360 tagmentation reaction with the Nextera XT DNA Library Prep Kit. Final libraries representing
361 each amplicon were characterized for average length using a DNA high sensitivity chip on a
362 2100 bioanalyzer (Agilent Technologies, Loveland, CO, USA) and quantitated with Quant-IT HS
363 reagents. Libraries were sequenced on a MiSeq.

Postmortem diagnosis of lymphocytic enteritis

All animals humanely euthanized or found dead at the WNPRC undergo complete post mortem
examination (necropsy) with histology. Standard hematoxylin and eosin (H&E) stains are used
for histological examinations to determine whether normal tissue architecture and cellular
populations are present. In this study, immunohistochemical (IHC) CD3 and CD20 or CD79
staining was performed on any samples with lymphocytic enteritis to differentiate lymphocyte
populations (primarily T cells, B cells, or mixed T and B cells). Diagnosis of T-cell rich
lymphocytic enteritis was based on abnormal villus architecture of the intestines and IHC
staining.[21,125] If confounding factors hampered diagnosis (e.g., severe B cell lymphoma or
autolysis), the animal was removed from the analysis.

Statistical analysis

375 We used univariate logistic regression to evaluate the associations of SOBV viremia with
376 enteritis risk. Analyses were repeated to determine association with enteritis in small bowel only,
377 colon only, both small bowel and colon, and either small bowel or colon. All reported P-values
378 are two-sided and $P<0.05$ was used to define statistical significance. Statistical analyses were
379 conducted using R version 3.6.3 in RStudio version 1.1.383.

Data accessibility and management

381 Metagenomic sequencing data have been deposited in the Sequence Read Archive (SRA)
382 under Bioproject PRJNA613737. Derived data, analysis pipelines, and figures have been made
383 available for easy replication of these results at a publicly-accessible GitHub
384 (https://github.com/aheffron/SPgVwnprc_in_marmosets).
385
386

Acknowledgments

This work was supported in part through Laulima Government Solutions, LLC prime contract
with the US National Institute of Allergy and Infectious Diseases (NIAID) under Contract No.
HHSN272201800013C. J.H.K. performed this work as employees of Tunnell Government
Services (TGS), a subcontractor of Laulima Government Solutions, LLC under Contract No.
HHSN272201800013C. The views and conclusions contained in this document are those of the
authors and should not be interpreted as necessarily representing the official policies, either
expressed or implied, of the US Department of Health and Human Services or of the institutions
and companies affiliated with the authors. Access to the dolphin silhouette was generously
provided by Chris Huh (https://creativecommons.org/licenses/by-sa/3.0/).
397
398
399

References

1. Tardif S, Bales K, Williams L, et al. Preparing New World monkeys for laboratory research. *ILAR J.* 2006;47(4):307-315.

2. Abbott DH, Barnett DK, Colman RJ, Yamamoto ME, Schultz-Darken NJ. Aspects of common marmoset basic biology and life history important for biomedical research. *Comp Med.* 2003;53(4):339-350.

3. Mansfield K. Marmoset models commonly used in biomedical research. *Comp Med.* 2003;53(4):383-392.

4. Primate Factsheets: Common marmoset (Callithrix jacchus) Taxonomy, Morphology, & Ecology. 2005. http://pin.primate.wisc.edu/factsheets/entry/common_marmoset. Accessed 04 April 2020.

5. Servick K. Why are U.S. neuroscientists clamoring for marmosets? *Science Magazine.* 23 Oct. 2018, 2018.

6. Converse AK, Aubert Y, Allers KA, Sommer B, Abbott DH. Flibanserin-Stimulated Partner Grooming Reflects Brain Metabolism Changes in Female Marmosets. *J Sex Med.* 2015;12(12):2256-2266.

7. Smith AL, Freeman SM, Barnhart TE, et al. Initial investigation of three selective and potent small molecule oxytocin receptor PET ligands in New World monkeys. *Bioorg Med Chem Lett.* 2016;26(14):3370-3375.

8. Ausderau KK, Dammann C, McManus K, Schneider M, Emborg ME, Schultz-Darken N. Cross-species comparison of behavioral neurodevelopmental milestones in the common marmoset monkey and human child. *Dev Psychobiol.* 2017;59(7):807-821.

9. Vermilyea SC, Guthrie S, Meyer M, et al. Induced Pluripotent Stem Cell-Derived Dopaminergic Neurons from Adult Common Marmoset Fibroblasts. *Stem Cells Dev.* 2017;26(17):1225-1235.

10. Olson EJ, Shaw GC, Hutchinson EK, et al. Bone Disease in the Common Marmoset: Radiographic and Histological Findings. *Vet Pathol.* 2015;52(5):883-893.

11. Ziegler TE, Sosa ME, Colman RJ. Fathering style influences health outcome in common marmoset (Callithrix jacchus) offspring. *PLoS One.* 2017;12(9):e0185695.

12. Kropp J, Di Marzo A, Golos T. Assisted reproductive technologies in the common marmoset: an integral species for developing nonhuman primate models of human diseases. *Biol Reprod.* 2017;96(2):277-287.

13. Jones CA, Duffy MK, Hoffman SA, et al. Vocalization development in common marmosets for neurodegenerative translational modeling. *Neurol Res.* 2018;40(4):303-311.

14. Iwatsuki-Horimoto K, Nakajima N, Kiso M, et al. The Marmoset as an Animal Model of Influenza: Infection With A(H1N1)pdm09 and Highly Pathogenic A(H5N1) Viruses via the Conventional or Tracheal Spray Route. *Front Microbiol.* 2018;9:844.

15. Braun K, Schultz-Darken N, Schneider M, Moore CF, Emborg ME. Development of a novel postnatal neurobehavioral scale for evaluation of common marmoset monkeys. *Am J Primatol.* 2015;77(4):401-417.

16. Kraynak M, Flowers MT, Shapiro RA, Kapoor A, Levine JE, Abbott DH. Extraovarian gonadotropin negative feedback revealed by aromatase inhibition in female marmoset monkeys. *Am J Physiol Endocrinol Metab.* 2017;313(5):E507-E514.

17. Parambeth JC, Ross CN, Miller AD, et al. Serum Cobalamin and Folate Concentrations in Common Marmosets (Callithrix jacchus) with Chronic Lymphocytic Enteritis. *Comp Med.* 2019;69(2):135-143.

18. Chalmers DT, Murgatroyd LB, Wadsworth PF. A survey of the pathology of marmosets (Callithrix jacchus) derived from a marmoset breeding unit. *Lab Anim.* 1983;17(4):270-279.

19. David JM, Dick EJ, Jr., Hubbard GB. Spontaneous pathology of the common marmoset
(Callithrix jacchus) and tamarins (Saguinus oedipus, Saguinus mystax). *J Med Primatol.*
2009;38(5):347-359.
20. Tucker MJ. A survey of the pathology of marmosets (Callithrix jacchus) under
experiment. *Lab Anim.* 1984;18(4):351-358.
21. Ludlage E, Mansfield K. Clinical care and diseases of the common marmoset (Callithrix
jacchus). *Comp Med.* 2003;53(4):369-382.
22. Kaspareit J, Friderichs-Gromoll S, Buse E, Habermann G. Background pathology of the
common marmoset (Callithrix jacchus) in toxicological studies. *Exp Toxicol Pathol.*
2006;57(5-6):405-410.
23. Bukh J, Kim JP, Govindarajan S, et al. Experimental infection of chimpanzees with
hepatitis G virus and genetic analysis of the virus. *J Infect Dis.* 1998;177(4):855-862.
24. Birkenmeyer LG, Desai SM, Muerhoff AS, et al. Isolation of a GB virus-related genome
from a chimpanzee. *J Med Virol.* 1998;56(1):44-51.
25. Cheng Y, Zhang W, Li J, et al. Serological and histological findings in infection and
transmission of GBV-C/HGV to macaques. *J Med Virol.* 2000;60(1):28-33.
26. Sibley SD, Lauck M, Bailey AL, et al. Discovery and characterization of distinct simian
pegiviruses in three wild African Old World monkey species. *PLoS One.*
2014;9(2):e98569.
27. Adams NJ, Prescott LE, Jarvis LM, et al. Detection in chimpanzees of a novel flavivirus
related to GB virus-C/hepatitis G virus. *J Gen Virol.* 1998;79 (Pt 8):1871-1877.
28. Baechlein C, Grundhoff A, Fischer N, et al. Pegivirus Infection in Domestic Pigs,
Germany. *Emerg Infect Dis.* 2016;22(7):1312-1314.
29. Yang C, Wang L, Shen H, et al. Detection and genetic characterization of porcine
pegivirus in pigs in the United States. *Transbound Emerg Dis.* 2018;65(3):618-626.
30. Lei D, Ye Y, Lin K, et al. Detection and genetic characterization of porcine pegivirus from
pigs in China. *Virus Genes.* 2019;55(2):248-252.
31. Epstein JH, Quan PL, Briese T, et al. Identification of GBV-D, a novel GB-like flavivirus
from old world frugivorous bats (Pteropus giganteus) in Bangladesh. *PLoS Pathog.*
2010;6:e1000972.
32. Kapoor A, Simmonds P, Scheel TK, et al. Identification of rodent homologs of hepatitis C
virus and pegiviruses. *mBio.* 2013;4(2):e00216-00213.
33. Shi M, Lin XD, Vasilakis N, et al. Divergent Viruses Discovered in Arthropods and
Vertebrates Revise the Evolutionary History of the Flaviviridae and Related Viruses. *J*
Virol. 2016;90(2):659-669.
34. Tomlinson JE, Kapoor A, Kumar A, et al. Viral testing of 18 consecutive cases of equine
serum hepatitis: A prospective study (2014-2018). *J Vet Intern Med.* 2019;33(1):251-
257.
35. Quan PL, Firth C, Conte JM, et al. Bats are a major natural reservoir for hepaciviruses
and pegiviruses. *Proc Natl Acad Sci U S A.* 2013;110(20):8194-8199.
36. Lole KS, Bollinger RC, Paranjape RS, et al. Full-length human immunodeficiency virus
type 1 genomes from subtype C-infected seroconverters in India, with evidence of
intersubtype recombination. *J Virol.* 1999;73(1):152-160.
37. Virus Taxonomy: 2019 Release: Genus: Pegivirus. 2020. https://talk.ictvonline.org/ictv-
reports/ictv_online_report/positive-sense-rna-viruses/w/flaviviridae/363/genus-pegivirus.
Accessed 04 April 2020.
38. Rodrigues TCS, Subramaniam K, McCulloch SD, et al. Genomic characterization of a
novel pegivirus species from free-ranging bottlenose dolphins (Tursiops truncatus) in the
Indian River Lagoon, Florida. *Virus Res.* 2019;263:98-101.

39.   Smith DB, Becher P, Bukh J, et al. Proposed update to the taxonomy of the genera
Hepacivirus and Pegivirus within the Flaviviridae family. *J Gen Virol.* 2016;97(11):2894-
2907.
40.   Simons JN, Leary TP, Dawson GJ, et al. Isolation of novel virus-like sequences
associated with human hepatitis. *Nat Med.* 1995;1(6):564-569.
41.   Heringlake S, Tillmann HL, Cordes-Temme P, Trautwein C, Hunsmann G, Manns MP.
GBV-C/HGV is not the major cause of autoimmune hepatitis. *J Hepatol.* 1996;25(6):980-
984.
42.   Bralet MP, Roudot-Thoraval F, Pawlotsky JM, et al. Histopathologic impact of GB virus C
infection on chronic hepatitis C. *Gastroenterology.* 1997;112(1):188-192.
43.   Feucht HH, Fischer L, Sterneck M, Broelsch CE, Laufs R. GB virus C transmission by
blood products. *Lancet.* 1997;349(9049):435.
44.   Goeser T, Seipp S, Wahl R, Muller HM, Stremmel W, Theilmann L. Clinical presentation
of GB-C virus infection in drug abusers with chronic hepatitis C. *J Hepatol.*
1997;26(3):498-502.
45.   Kanda T, Yokosuka O, Imazeki F, et al. GB virus-C RNA in Japanese patients with
hepatocellular carcinoma and cirrhosis. *J Hepatol.* 1997;27(3):464-469.
46.   Hadziyannis SJ. Fulminant hepatitis and the new G/GBV-C flavivirus. *J Viral Hepat.*
1998;5(1):15-19.
47.   Rambusch EG, Wedemeyer H, Tillmann HL, Heringlake S, Manns MP. [Significance of
coinfection with hepatitis G virus for chronic hepatitis C--a review of the literature]. *Z
Gastroenterol.* 1998;36(1):41-53.
48.   Brown KE, Wong S, Young NS. Prevalence of GBV-C/HGV, a novel 'hepatitis' virus, in
patients with aplastic anaemia. *Br J Haematol.* 1997;97(2):492-496.
49.   Servant A, Bogard M, Delaugerre C, Cohen P, Deny P, Guillevin L. GB virus C in
systemic medium- and small-vessel necrotizing vasculitides. *Br J Rheumatol.*
1998;37(12):1292-1294.
50.   Viazov S, Alberts KR, Ross RS, Seemayer CA, Roggendorf M. Lack of association
between GBV-C infection and spontaneous abortion. *Eur J Clin Microbiol Infect Dis.*
1999;18(6):458-459.
51.   Misiani R, Mantero G, Bellavita P, et al. GB virus C infection in patients with type II
mixed cryoglobulinemia. *Ann Intern Med.* 1997;127(10):891-894.
52.   Crovatto M, Mazzaro C, Mishiro S, et al. GBV-C/HGV and HCV infection in mixed
cryoglobulinaemia. *Br J Haematol.* 1999;106(2):510-514.
53.   Liu F, Knight GB, Agnello V. Hepatitis C virus but not GB virus C/hepatitis G virus has a
role in type II cryoglobulinemia. *Arthritis Rheum.* 1999;42(9):1898-1901.
54.   Hardie D, Smuts H. Human pegivirus-1 in the CSF of patients with HIV-associated
neurocognitive disorder (HAND) may be derived from blood in highly viraemic patients. *J
Clin Virol.* 2017;91:58-61.
55.   Sauleda S, Rio J, Montalban X, Martinez-Caceres E, Esteban JI, Guardia J. Lack of
association between hepatitis G virus and multiple sclerosis. *J Neurol Neurosurg
Psychiatry.* 1998;64(2):283.
56.   Lamoril J, Andant C, Bogard C, et al. Epidemiology of hepatitis C and G in sporadic and
familial porphyria cutanea tarda. *Hepatology.* 1998;27(3):848-852.
57.   Font J, Tassies D, Garcia-Carrasco M, et al. Hepatitis G virus infection in primary
Sjogren's syndrome: analysis in a series of 100 patients. *Ann Rheum Dis.*
1998;57(1):42-44.
58.   Jones JF, Kulkarni PS, Butera ST, Reeves WC. GB virus-C--a virus without a disease:
we cannot give it chronic fatigue syndrome. *BMC Infect Dis.* 2005;5:78.

59. De Renzo A, Persico E, de Marino F, et al. High prevalence of hepatitis G virus infection
in Hodgkin's disease and B-cell lymphoproliferative disorders: absence of correlation
with hepatitis C virus infection. *Haematologica.* 2002;87(7):714-718; discussion 718.
60. Krajden M, Yu A, Braybrook H, et al. GBV-C/hepatitis G virus infection and non-Hodgkin
lymphoma: a case control study. *Int J Cancer.* 2010;126(12):2885-2892.
61. Giannoulis E, Economopoulos T, Mandraveli K, et al. The prevalence of hepatitis C and
hepatitis G virus infection in patients with B cell non-Hodgkin lymphomas in Greece: a
Hellenic Cooperative Oncology Group Study. *Acta Haematol.* 2004;112(4):189-193.
62. Wiwanitkit V. Individuals with HGV-RNA are at high risk of B cell non-Hodgkin's
lymphoma development. *Asian Pac J Cancer Prev.* 2005;6(2):215-216.
63. Ellenrieder V, Weidenbach H, Frickhofen N, et al. HCV and HGV in B-cell non-Hodgkin's
lymphoma. *J Hepatol.* 1998;28(1):34-39.
64. Pavlova BG, Heinz R, Selim U, Tuchler H, Pittermann E, Eder G. Association of GB
virus C (GBV-C)/hepatitis G virus (HGV) with haematological diseases of different
malignant potential. *J Med Virol.* 1999;57(4):361-366.
65. Chang CM, Stapleton JT, Klinzman D, et al. GBV-C infection and risk of NHL among
U.S. adults. *Cancer Res.* 2014;74(19):5553-5560.
66. Fama A, Xiang J, Link BK, et al. Human Pegivirus infection and lymphoma risk and
prognosis: a North American study. *Br J Haematol.* 2018;182(5):644-653.
67. Collier JD, Zanke B, Moore M, et al. No association between hepatitis C and B-cell
lymphoma. *Hepatology.* 1999;29(4):1259-1261.
68. Arican A, Sengezer T, Bozdayi M, et al. Prevalence of hepatitis-G virus and hepatitis-C
virus infection in patients with non-Hodgkin's lymphoma. *Med Oncol.* 2000;17(2):123-
126.
69. Nicolosi Guidicelli S, Lopez-Guillermo A, Falcone U, et al. Hepatitis C virus and GBV-C
virus prevalence among patients with B-cell lymphoma in different European regions: a
case-control study of the International Extranodal Lymphoma Study Group. *Hematol
Oncol.* 2012;30(3):137-142.
70. Keresztes K, Takacs M, Horanyi M, Miltenyi Z, Illes A. HCV and HGV infection in
Hodgkin's disease. *Pathol Oncol Res.* 2003;9(4):222-225.
71. Persico M, De Renzo A, Persico E, Notaro R, Torella R, Rotoli B. Hepatitis G virus in
patients with Hodgkin's lymphoma. *Br J Haematol.* 1998;103(4):1206-1207.
72. Toyoda H, Fukuda Y, Hayakawa T, Takamatsu J, Saito H. Effect of GB virus C/hepatitis
G virus coinfection on the course of HIV infection in hemophilia patients in Japan. *J
Acquir Immune Defic Syndr Hum Retrovirol.* 1998;17(3):209-213.
73. Tillmann HL, Heringlake S, Trautwein C, et al. Antibodies against the GB virus C
envelope 2 protein before liver transplantation protect against GB virus C de novo
infection. *Hepatology.* 1998;28(2):379-384.
74. Lefrere JJ, Roudot-Thoraval F, Morand-Joubert L, et al. Carriage of GB virus C/hepatitis
G virus RNA is associated with a slower immunologic, virologic, and clinical progression
of human immunodeficiency virus disease in coinfected persons. *J Infect Dis.*
1999;179(4):783-789.
75. Yeo AE, Matsumoto A, Hisada M, Shih JW, Alter HJ, Goedert JJ. Effect of hepatitis G
virus infection on progression of HIV infection in patients with hemophilia. Multicenter
Hemophilia Cohort Study. *Ann Intern Med.* 2000;132(12):959-963.
76. Xiang J, Wunschmann S, Diekema DJ, et al. Effect of coinfection with GB virus C on
survival among patients with HIV infection. *N Engl J Med.* 2001;345(10):707-714.
77. Tillmann HL, Heiken H, Knapik-Botor A, et al. Infection with GB virus C and reduced
mortality among HIV-infected patients. *N Engl J Med.* 2001;345(10):715-724.
78. Williams CF, Klinzman D, Yamashita TE, et al. Persistent GB virus C infection and
survival in HIV-infected men. *N Engl J Med.* 2004;350(10):981-990.

79. Van der Bij AK, Kloosterboer N, Prins M, et al. GB virus C coinfection and HIV-1 disease
progression: The Amsterdam Cohort Study. *J Infect Dis.* 2005;191(5):678-685.
80. Tenckhoff S, Kaiser T, Bredeek F, et al. Role of GB virus C in HIV-1-infected and
hepatitis C virus-infected hemophiliac children and adolescents. *J Acquir Immune Defic*
Syndr. 2012;61(2):243-248.
81. Sahni H, Kirkwood K, Kyriakides TC, et al. GBV-C viremia and clinical events in
advanced HIV infection. *J Med Virol.* 2014;86(3):426-432.
82. Hollingsworth RC, Jameson CL, Minton JE, et al. GBV-C/HGV coinfection in HIV-1-
positive men: frequent detection of viral RNA in blood plasma but absence from seminal
fluid plasma. *J Med Virol.* 1998;56(4):321-326.
83. Goubau P, Liu HF, Goderniaux E, Burtonboy G. Influence of CD4+ lymphocyte counts
on GB virus C/hepatitis G virus carriership in HIV-positive individuals. *J Med Virol.*
1999;57(4):367-369.
84. Rey D, Vidinic-Moularde J, Meyer P, et al. High prevalence of GB virus C/hepatitis G
virus RNA and antibodies in patients infected with human immunodeficiency virus type 1.
Eur J Clin Microbiol Infect Dis. 2000;19(9):721-724.
85. Voirin N, Trepo C, Esteve J, et al. Effects of co-infection with hepatitis C virus and GB
virus C on CD4 cell count and HIV-RNA level among HIV-infected patients treated with
highly active antiretroviral therapy. *AIDS.* 2002;16(11):1556-1559.
86. Li C, Collini P, Danso K, et al. GB virus C and HIV-1 RNA load in single virus and co-
infected West African individuals. *AIDS.* 2006;20(3):379-386.
87. Lauck M, Bailey AL, Andersen KG, Goldberg TL, Sabeti PC, O'Connor DH. GB virus C
coinfections in west African Ebola patients. *J Virol.* 2015;89(4):2425-2429.
88. Zhang W, Chaloner K, Tillmann HL, Williams CF, Stapleton JT. Effect of early and late
GB virus C viraemia on survival of HIV-infected individuals: a meta-analysis. *HIV Med.*
2006;7(3):173-180.
89. Bhattarai N, Stapleton JT. GB virus C: the good boy virus? *Trends Microbiol.*
2012;20(3):124-130.
90. Porter AF, Pettersson JH-O, Wei-Shan C, et al. Metagenomic identification of diverse
animal hepaciviruses and pegiviruses. *bioRxiv.* 2020.
91. Bukh J, Apgar CL. Five new or recently discovered (GBV-A) virus species are
indigenous to New World monkeys and may constitute a separate genus of the
Flaviviridae. *Virology.* 1997;229(2):429-436.
92. Seifried E, Bialleck H, Weber H, et al. [Prevalence of hepatitis G virus genome in blood
donors]. *Beitr Infusionsther Transfusionsmed.* 1997;34:11-15.
93. Roth WK, Waschk D, Marx S, et al. Prevalence of hepatitis G virus and its strain variant,
the GB agent, in blood donations and their transmission to recipients. *Transfusion.*
1997;37(6):651-656.
94. Yoshikawa A, Fukuda S, Itoh K, et al. Infection with hepatitis G virus and its strain
variant, the GB agent (GBV-C), among blood donors in Japan. *Transfusion.*
1997;37(6):657-663.
95. Gutierrez RA, Dawson GJ, Knigge MF, et al. Seroprevalence of GB virus C and
persistence of RNA and antibody. *J Med Virol.* 1997;53(2):167-173.
96. Lara C, Halasz R, Sonnerborg A, Sallberg M. Detection of hepatitis G virus RNA in
persons with and without known risk factors for blood-borne viral infections in Sweden
and Honduras. *J Clin Microbiol.* 1998;36(1):255-257.
97. Gallian P, Rodrigues V, Cantaloube JF, et al. High prevalence of GB-C/hepatitis G virus
in a Brazilian population with helminth infection. *J Med Virol.* 1998;56(4):310-315.
98. Mphahlele MJ, Aspinall S, Spooner R, Carman WF. Age related prevalence of hepatitis
G virus in South Africans. *J Clin Pathol.* 1999;52(10):752-757.

99.   Corwin AL, Hyams KC, Kim JP, et al. Short report: evidence of worldwide transmission
of hepatitis G virus. *Am J Trop Med Hyg.* 1997;57(4):455-456.
100. Herrera S, Perlaza BL, Bonelo A, Arevalo-Herrera M. Aotus monkeys: their great value
for anti-malaria vaccines and drug testing. *Int J Parasitol.* 2002;32(13):1625-1635.
101. Hutt MS, Davies DR, Voller A. Malarial infections in Aotus trivirgatus with special
reference to renal pathology. II. P. falciparum and mixed malaria infections. *Br J Exp*
Pathol. 1975;56(5):429-438.
102. Schmidt LH. Plasmodium falciparum and Plasmodium vivax infections in the owl monkey
(Aotus trivirgatus). II. Responses to chloroquine, quinine, and pyrimethamine. *Am J Trop*
Med Hyg. 1978;27(4):703-717.
103. Linnen J, Wages J, Jr., Zhang-Keck ZY, et al. Molecular cloning and disease association
of hepatitis G virus: a transfusion-transmissible agent. *Science.* 1996;271(5248):505-
508.
104. Karayiannis P, Thomas HC. Current status of hepatitis G virus (GBV-C) in transfusion: is
it relevant? *Vox Sang.* 1997;73(2):63-69.
105. Stark K, Meyer CG, Tacke M, et al. Hepatitis G virus RNA and hepatitis G virus
antibodies in renal transplant recipients: prevalence and risk factors. *Transplantation.*
1997;64(4):608-612.
106. Kallinowski B, Ahmadi R, Seipp S, Bommer J, Stremmel W. Clinical impact of GB-C
virus in haemodialysis patients. *Nephrol Dial Transplant.* 1998;13(1):93-98.
107. Huang CH, Kao JH, Kuo YM, Tsai TJ, Hung KY, Chen DS. GB virus C/hepatitis G virus
infection in patients on continuous ambulatory peritoneal dialysis. *Nephrol Dial*
Transplant. 1998;13(11):2914-2919.
108. Fiordalisi G, Bettinardi A, Zanella I, et al. Parenteral and sexual transmission of GB virus
C and hepatitis C virus among human immunodeficiency virus-positive patients. *J Infect*
Dis. 1997;175(4):1025-1026.
109. Anastassopoulou CG, Paraskevis D, Sypsa V, et al. Prevalence patterns and genotypes
of GB virus C/hepatitis G virus among imprisoned intravenous drug users. *J Med Virol.*
1998;56(3):246-252.
110. Shibuya A, Takeuchi A, Sakurai K, Saigenji K. Hepatitis G virus infection from needle-
stick injuries in hospital employees. *J Hosp Infect.* 1998;40(4):287-290.
111. Kao JH, Chen W, Chen PJ, Lai MY, Lin RY, Chen DS. GB virus-C/hepatitis G virus
infection in prostitutes: possible role of sexual transmission. *J Med Virol.* 1997;52(4):381-
384.
112. Semprini AE, Persico T, Thiers V, et al. Absence of hepatitis C virus and detection of
hepatitis G virus/GB virus C RNA sequences in the semen of infected men. *J Infect Dis.*
1998;177(4):848-854.
113. Viazov S, Riffelmann M, Sarr S, Ballauff A, Meisel H, Roggendorf M. Transmission of
GBV-C/HGV from drug-addicted mothers to their babies. *J Hepatol.* 1997;27(1):85-90.
114. Zanetti AR, Tanzi E, Romano L, et al. Multicenter trial on mother-to-infant transmission
of GBV-C virus. The Lombardy Study Group on Vertical/Perinatal Hepatitis Viruses
Transmission. *J Med Virol.* 1998;54(2):107-112.
115. Hino K, Moriya T, Ohno N, et al. Mother-to-infant transmission occurs more frequently
with GB virus C than hepatitis C virus. *Arch Virol.* 1998;143(1):65-72.
116. Lin HH, Kao JH, Yeh KY, et al. Mother-to-infant transmission of GB virus C/hepatitis G
virus: the role of high-titered maternal viremia and mode of delivery. *J Infect Dis.*
1998;177(5):1202-1206.
117. Santos LM, Lobato RC, Barral MFM, Goncalves CV, da Hora VP, Martinez AMB.
Prevalence and vertical transmission of human pegivirus among pregnant women
infected with HIV. *Int J Gynaecol Obstet.* 2017;138(1):113-118.

118.   Wang JT, Tsai FC, Lee CZ, et al. A prospective study of transfusion-transmitted GB
701   virus C infection: similar frequency but different clinical presentation compared with
702   hepatitis C virus. *Blood.* 1996;88(5):1881-1886.

703   119.   Prati D, Zanella A, Bosoni P, et al. The incidence and natural course of transfusion-
704   associated GB virus C/hepatitis G virus infection in a cohort of thalassemic patients. The
705   Cooleycare Cooperative Group. *Blood.* 1998;91(3):774-777.

706   120.   Nakao H, Okamoto H, Fukuda M, et al. Mutation rate of GB virus C/hepatitis G virus over
707   the entire genome and in subgenomic regions. *Virology.* 1997;233(1):43-50.

708   121.   Suzuki Y, Katayama K, Fukushi S, et al. Slow evolutionary rate of GB virus C/hepatitis G
709   virus. *J Mol Evol.* 1999;48(4):383-389.

710   122.   Chivero ET, Stapleton JT. Tropism of human pegivirus (formerly known as GB virus
711   C/hepatitis G virus) and host immunomodulation: insights into a highly successful viral
712   infection. *J Gen Virol.* 2015;96(Pt 7):1521-1532.

713   123.   Lauck M, Sibley SD, Hyeroba D, et al. Exceptional simian hemorrhagic fever virus
714   diversity in a wild African primate community. *J Virol.* 2013;87(1):688-691.

715   124.   Lauck M, Switzer WM, Sibley SD, et al. Discovery and full genome characterization of
716   two highly divergent simian immunodeficiency viruses infecting black-and-white colobus
717   monkeys (Colobus guereza) in Kibale National Park, Uganda. *Retrovirology.*
718   2013;10:107.

719   125.   Gartner LP, Hiatt JL. *Color atlas of histology.* 5th ed. Philadelphia: Wolters Kluwer
720   Health/Lippincott William & Wilkins; 2009.