Acibenzolar-S-methyl activates stomatal-based defense systemically in Japanese radish by inducing peroxidase-dependent reactive oxygen species production

Nanami Sakata[1], Takako Ishiga[1], Shizuku Taniguchi[2] and Yasuhiro Ishiga[1*]

[1] Faculty of Life and Environmental Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba,
Ibaraki, Japan.

[2]Syngenta Japan, 780 Kuno-cho, Ushiku, Ibaraki, Japan.

*Correspondence:

Yasuhiro Ishiga

ishiga.yasuhiro.km@u.tsukuba.ac.jp

Keywords: Acibenzolar-S-methyl, *Pseudomonas cannabina* pv. *alisalensis*, Japanese radish,
Stomatal-based defense, Systemic acquired resistance, Plant defense activator, Peroxidase,
Reactive oxygen species

Abstract

Acibenzolar-S-methyl (ASM) is a well-known plant activator, which is a synthetic analog of salicylic
acid (SA). Recently, copper fungicides and antibiotics are major strategies for controlling bacterial
diseases. However, resistant strains have already been found. Therefore, there is an increasing
demand for sustainable new disease control strategies. We investigated the ASM disease control
effect against *Pseudomonas cannabina* pv. *alisalensis* (*Pcal*), which causes bacterial blight on
Japanese radish. In this study, we demonstrated that ASM effectively suppressed *Pcal* disease
symptom development associated with reduced bacterial populations on Japanese radish leaves.
Interestingly, we also demonstrated that ASM activated systemic acquired resistance (SAR),
including stomatal-based defense, not only on ASM treated leaves, but also on untreated upper and
lower leaves. Reactive oxidative species (ROS) are essential second messengers in stomatal-based
defense. We found that ASM induced stomatal closure by inducing ROS production through
peroxidase. These results indicate that stomatal closure induced by ASM treatment is effective for
preventing *Pcal* pathogen invasion into plants, and in turn reduction of disease development.

Introduction

28 Recently, several bacterial pathogens were identified as causal agents of severe worldwide
29 epidemics. *Pseudomonas cannabina* pv. *alisalensis* (*Pcal*) causes bacterial blight of crucifer plants
30 (Brassicaceae plants) (Takikawa and Takahashi, 2014) and inflicts great damage on crucifer
31 production including cabbage, Chinese cabbage, and Japanese radish. On Japanese radish, disease
32 symptoms caused by a newly isolated *Pcal* were observed on leaves and roots (Takikawa and

33 Takahashi, 2014). Discoloration of roots and necrotic leaf lesions have serious implications for the
34 commercial value of Japanese radish. Chemical treatments such as copper fungicides and antibiotics
35 are popular strategies for bacterial disease control. However, *Pcal* strains resistant against these
36 chemicals were found (Takahashi et al., 2013). Therefore, the demand for developing sustainable
37 alternative strategies has been increasing.

38 Acibenzolar-S-methyl (ASM), a synthetic analog of salicylic acid (SA), has been used to protect
39 crops from several diseases by activating plant defense (Kunz et al., 1997; Oostendorp et al., 2001).
40 We demonstrated that ASM soil drench application induced systemic acquired resistance (SAR)
41 against *Pcal* (Ishiga et al., 2020). SAR processes can be divided into three steps: local immune
42 activation, information relay from local to systemic tissues by mobile signals, and defense activation
43 and priming in systemic tissues (Jung et al., 2009; Shah and Zeier, 2013; Wang et al., 2018). Thus,
44 our previous results implied that mobile signals generated by ASM triggered SAR in untreated
45 leaves. Pre-treating cucumber plant first leaves with ASM protected whole plants from fungal
46 infection with *Colletotrichum orbiculare* (Cools and Ishii, 2002). In cucumber plants, *SAR* genes
47 were rapidly and highly expressed in upper leaves after first leaf treatment with ASM (Cools and
48 Ishii, 2002; Narusaka et al., 1999; Narusaka et al., 2001). However, the ASM-induced SAR
49 mechanism is still unclear. We also demonstrated that ASM protects cabbage plants from *Pcal* by
50 activating stomatal-based defense (Ishiga et al., 2020). Furthermore, ASM induced stomatal-based
51 defense resulted in reduced *Pcal* bacterial populations as well as disease symptoms on cabbage
52 within 4 h after soil drench, and this positive effect is prolonged for at least three weeks (Ishiga et al.,
53 unpublished). Additionally, many foliar bacterial pathogens target stomata as an entry site. Therefore,
54 ASM is expected to have a powerful disease control effect. However, whether ASM is effective at
55 suppressing disease development against *Pcal* on other crops (in addition to cabbage) has not been
56 evaluated.

57 ROS production at the cell surface, which is called oxidative burst, is one of the earliest defenses
58 detectable during pathogen-associated molecular pattern (PAMP)-triggered immunity (PTI) and
59 effector-triggered immunity (ETI). Doke (1983) first demonstrated that ROS members possibly
60 function as the chemical signals required for hypersensitive response (HR) induction. The oxidative
61 burst induced by pathogen-derived signals is essential to defense mechanisms throughout the plant
62 (Fobert and Després, 2005). Apoplastic ROS are generated from plasma membrane-localized
63 NADPH oxidases and cell wall-localized peroxidases. ROS are known as antimicrobial molecules
64 involved in cell wall reinforcement and callose deposition. ROS also act as local and systemic
65 secondary messengers triggering additional immune responses, including stomatal movement.
66 Stomatal closure in response to various stress is brought about by loss of guard cell turgor, which is
67 caused by ion channel modulation. A complex signaling network involving ROS production
68 regulates these channels (Kim et al., 2010; Kollist et al., 2014). ASM treatment of the first leaves via
69 fungal inoculation caused rapid H_2O_2 accumulation below the penetration site on ASM-untreated
70 third leaves (Lin and Ishii, 2009; Park et al., 2019), which might have contributed to the triggered
71 defense responses against fungal pathogen infection. Endogenous SA is essential for both apoplastic
72 ROS production and stomatal closure induced by chitosan as microbe-associated molecular patterns
73 (MAMPs) (Prodhan et al., 2017). Since ASM is a synthetic analog of SA, it is tempting to speculate
74 that ASM induced stomatal closure is closely related to ROS production.

75 We demonstrate that ASM successfully suppresses disease development and bacterial multiplication
76 in Japanese radish by activating stomatal defense. Furthermore, to gain knowledge into the ASM
77 action mechanism on stomata, we investigated its effect on ROS production. ASM activated ROS

78   production mediated by peroxidases, which in turn activated stomatal-based defense. Moreover,
79   ASM also induces *PR* gene expression on Japanese radish. Importantly, ASM controlled bacterial
80   blight disease not only on the treated leaves, but also on Japanese radish upper and lower leaves.
81   Thus, our results highly support that ASM can be an additional disease management tool to prevent
82   crop disease losses against bacterial pathogens.

Materials and Methods

Plant materials and chemical treatment

Japanese radish (*Raphanus sativus* var. *longipinnatus*) cv. Natsutsukasa plants were used for all
experiments. Seedlings were grown in 9 cm pots at 24°C with a light intensity of 200 μmol/(m^2 sec)
and 16 h light/8 h dark in a growth chamber or were maintained in the greenhouse under a natural
photoperiod at 21.2±2.4°C. Acibenzolar-S-methyl (ASM, marketed as ACTIGARD®) was supplied
courtesy of Syngenta. To evaluate the effect of ASM on plant defense, Japanese radish fourth leaves
were treated with ASM (100 ppm) by dip-treatment at 3 weeks after sowing or when fifth true leaves
were unfolded. Salicylhydroxamic acid (SHAM) and diphenyleneiodonium chloride (DPI) were
obtained from Sigma-Aldrich (Sigma-Aldrich, St Louis, MO, USA). Plants were treated with ASM 4
h, 1 d, and 1 w before *Pcal* inoculation (Supplementary Figure 1). Plants were dip-treated with water
or mock-inoculated with water as controls.

Bacterial strains and growth conditions

97   Pathogenic *Pseudomonas cannabina* pv. *alisalensis* strain KB211 (*Pcal*) (Sakata et al., 2019) was
98   kindly provided by the Nagano Vegetable and Ornamental Crops Experiment Station, Nagano, Japan.
99   *Pcal* was grown at 28°C for 24 h on Kings B (KB) (King et al., 1954) agar medium. For inoculum,
100   bacteria were suspended in sterile distilled H$_2$O, and cell densities measured at 600 nm (OD$_{600}$) using
101   a Biowave CO8000 Cell Density Meter (biochrom, Cambridge, UK).

Bacterial inoculation

Plants were spray-inoculated with a bacterial suspension in sterile distilled water containing 0.025%
Silwet L-77 (bio medical science, Tokyo, Japan). For *Pcal* lesion area and disease symptoms
measurement, Japanese radish plants were spray-inoculated with a bacterial suspension (5 x 10^7
colony forming units [CFU]/ml) to runoff. Dip-inoculation method was used for stomatal assay. For
bacterial growth assay, Japanese radish plants were spray-inoculated with a bacterial suspension (5 x
10^7 CFU/ml). The plants were then incubated in growth chambers at approximately 100% RH for the
first 24 h, then at approximately 70% RH for the rest of the experiment. Lesion area and bacterial
population were assessed 1-week post-inoculation (wpi).

Real-time quantitative RT-PCR

For expression profiles of Japanese radish defense genes in response to ASM, plants were dip-treated
with water as a control or ASM on fourth leaves. After 4 h total RNA was extracted using RNAiso
Plus (Takara Bio, Shiga, Japan) according to the manufacturer's protocol and used for real-time
quantitative RT-PCR (qRT-PCR) as described (Ishiga and Ichinose 2016). Two micrograms of total
RNA were treated with gDNA Remover (TOYOBO, Osaka, Japan) to eliminate genomic DNA, and

the DNase-treated RNA was reverse transcribed using the ReverTra Ace® qPCR RT Kit (TOYOBO).
The cDNA (1:50) was then used for RT-qPCR with the primers shown below with
THUNDERBIRD® qPCR Mix (TOYOBO) on a Thermal Cycler Dice Real Time System (Takara
Bio). Radish *Glyceraldehyde 3-phosphate dehydrogenase* (*GAPDH*) gene was used as an internal
control. Average CT values, calculated using the second derivative maximum method from triplicate
samples, were used to determine the fold expression relative to the controls (0 time). Primers used in
gene-specific PCR amplification for *PR1* were 5′- AAAGCTACGCCGACCGACTACGAG -3′ and
5′- CCAGAAAAGTCGGCGCTACTCCA -3′; for *PR2*, 5′- GTACGCTCTGTTCAAACCGACCC -
3′ and 5′- TTTCCAACGATCCTCCGCCTGA -3′; for *PR3*, 5′-
TCTTTGGTCAGACTTCCCACGAG -3′ and 5′- GATGGCTCTTCCACACTGTCCGTA -3′; and
for *GAPDH*, 5′- CGCTTCCTTCAACATCATTCCCA -3′ and 5′-
TCAGATTCCTCCTTGATAGCCTT -3′.

Stomatal assay

A modified method was used to assess stomatal response as described previously (Chitrakar and
Melotto, 2010). Briefly, Japanese radish plants were grown for 3 weeks after sowing as described
previously. *Pcal* was grown at 28°C for 24 h on KB agar medium, then suspended in sterile distilled
water to 1 x 10^8 CFU/ml. Dip-inoculated or water mock-inoculated radish leaves were directly
imaged at 4 h post-inoculation (hpi) using a Nikon optical microscope (Eclipse 80i). Additionally, 1
week after ASM treatment, leaves were treated with SHAM (1 mM) and DPI (10 µM) with and
without *Pcal*. The aperture width of at least 100 stomata was measured 4 h post inoculation and
treatment (Supplementary Figure 1). The average and standard error for the stomatal aperture width
were calculated. The stomatal apertures were evaluated in three independent experiments.

Results

ASM suppresses *Pcal* disease development in radish and triggers SAR

To investigate if ASM effectively suppresses *Pcal* disease development in Japanese radish plants,
plants were inoculated with *Pcal*, 4 h, 1 d, and 1 w after ASM dip-treatment on only the fourth leaf.
Then disease symptoms were observed 7 dpi (Supplementary Figure 1). ASM-treated Japanese radish
plants showed less disease symptoms than control water-treated plants inoculated with *Pcal* in all
treatments (Supplementary Figure 2). We also evaluated the disease development by measuring
lesion areas. *Pcal* lesion area on ASM-treated Japanese radish plants was significantly smaller than
that of control water-treated plants within 4 h post-treatment (hpt) (Figures 1A, B, C). Furthermore,
lesion area was reduced not only on the treated Japanese radish leaves, but also on the untreated
upper and lower leaves (Figures 1A, B, C). To investigate whether ASM suppresses bacterial growth
as well as lesion area, we also measured bacterial population 7 dpi. In ASM-treated Japanese radish
fourth leaves, *Pcal* populations were around ten times less compared to those of the water-treated
control 4 hpt (Figure 2A). In addition, at 1 d post-treatment (dpt) and 1w post-treatment (wpt) with
ASM, *Pcal* populations were around 100 times less compared to those of the water-treated control
(Figures 2B, C). Moreover, in third and fifth leaves, *Pcal* populations were also less compared to
those of the water-treated control (Figures 2A, B, C). These results indicate that ASM is able to
suppress symptom development and bacterial multiplication in Japanese radish, and this effect is
acquired systemically.

ASM activates stomatal-based defense against *Pcal*

We demonstrated that ASM activated stomatal-based defense against *Pcal*, resulting in reduced
disease development and bacterial populations on cabbage (Ishiga et al., 2020; Ishiga et al.,
unpublished). To examine whether ASM activates stomatal-based defense on Japanese radish, we
observed stomatal response after *Pcal* inoculation after ASM dip-treatment on fourth leaves. After
Pcal inoculation, ASM treated Japanese radish leaves exhibited stomatal closure (Figures 3B, E, H).
Interestingly, we also observed stomatal closure of the untreated upper and lower leaves (Figures 3A,
C, D, F, G, I). These results indicate ASM activated stomatal-based defense against *Pcal* on radish
systemically. Furthermore, stomatal closure was observed on ASM mock-inoculated Japanese radish
leaves at 1 dpt and 1 wpt (Figures 3D, E, F, G, H, I), but not 4 hpt (Figures 3A, B, C). These results
indicate that ASM suppresses stomatal opening regardless of *Pcal* inoculation 1 dpt.

ASM induced stomatal closure by inducing peroxidase-dependent ROS production

Reactive oxygen species (ROS) are essential second messengers in stomatal immunity (Joshi-Saha et
al., 2011). Stomatal closure by the phytohormone salicylic acid (SA) requires ROS production by cell
wall peroxidases (Khokon et al., 2011; Mori et al., 2001). Since ASM is a synthetic analog of SA, we
investigated whether ASM induced stomatal closure by producing ROS through peroxidase by using
a peroxidase inhibitor, salicylhydroxamic acid (SHAM), and an NADPH oxidase inhibitor,
diphenylene iodonium (DPI). ASM-induced stomatal closure was suppressed by SHAM (Figures 4A,
B, C), indicating that cell wall peroxidases are involved in mediating ROS production during ASM-
induced stomatal closure. Conversely, exogenous DPI application did not inhibit the ASM-induced
stomatal closure (Figures 4A, B, C). We obtained the same results after *Pcal* inoculation
(Supplementary Figures 3A, B, C). These results indicate that ASM induces stomatal closure via
ROS production mediated by peroxidases.

ASM induces defense-related gene expression in Japanese radish

ASM also induced defense-related gene expression, including PR proteins (Ishiga et al., 2020;
Kouzai, et al., 2018; Kouzai, et al., 2018; Nakashita et al., 2002; Yasuda et al., 2008). To investigate
the effect of ASM treatment on Japanese radish defense gene expression, we investigated the
expression profiles of *PR1*, *PR2*, and *PR3* in response to ASM. The expression of *PR1*, *PR2*, and
PR3 was significantly induced in fourth leaves 4 h after ASM treatment (Figures 5A, B, C).
Furthermore, in both untreated upper and lower leaves, all gene expression was induced or tended to
be induced (Figures 5A, B, C). These results indicate that ASM induced defense-related gene
expression triggered systemic acquired resistance in Japanese radish.

Discussion

There is an increased demand for developing sustainable disease control strategies. ASM is predicted
as an additional disease management tool to protect numerous crops from pathogens without the
occurrence of chemically resistant strains. Our results clearly showed that an ASM dip-treatment
effectively suppressed *Pcal* lesion formation and bacterial multiplication on Japanese radish plants
(Figure 1; Figure 2; Supplementary Figure 2). Previously, we also demonstrated that an ASM soil
drench suppressed *Pcal* disease development associated with reduced bacterial population in cabbage

(Ishiga et al., 2020). These results indicate that ASM is a powerful tool against *Pcal*, which is the
causal agent of bacterial blight of cruciferous plants.

The most important finding was that ASM greatly enhanced the systemic activation of defense
response. The reduction of disease development and bacterial growth was observed not only on
treated fourth leaves, but also on untreated third and fifth leaves (Figure 1; Figure 2; Supplementary
Figure 2). Additionally, ASM triggered stomatal-based defense and SAR-related gene expression
induction on both treated and untreated leaves (Figure 3; Figure 5). Cools and Ishii (2002)
demonstrated that ASM pre-treatment of cucumber first leaves protected whole plants from infection
with the virulent fungal pathogen *Colletotrichum orbiculare*. Three hours post-ASM treatment on
first leaves, ASM-induced systemic priming of *PAL1* expression in cucumber plants occurred
rapidly, with enhanced expression in the third leaves (Cools and Ishii, 2002). Interestingly, our
results showed ASM-induced systemic defense was acquired not only on untreated upper leaves, but
also on untreated lower leaves (Figure 5). Kubota and Abiko (2000) demonstrated that 7 days after a
first inoculation with *Colletotrichum lagenarium* and *Pythium ultimum* on cucumber cotyledons and
primary leaves, induced resistance was observed against the challenge inoculation with those
pathogenic fungi in hypocotyl. These results implied that foliar leaf treatment can prevent disease of
lower plant parts, including hypocotyls and roots (Kubota and Abiko, 2000). Importantly, *Pcal*
causes discoloration of Japanese radish hypocotyl and root (Takikawa and Takahashi, 2014). It is
expected that ASM-induced systemic defense could be a powerful disease management tool against
bacterial blight on Japanese radish.

The gene expression profiles of *PR1*, *PR2*, and *PR3* were induced within 4 h in both treated and
untreated leaves (Figure 5). Benzo (1,2,3) thiadiazole-7-carboxylic acid (acibenzolar), but not ASM
itself was detected in untreated third leaves 3 dpt on first leaves (Ishii et al., 2002). SABP2 catalyzes
ASM conversion into acibenzolar to induce SAR (Tripathi et al., 2010). Therefore, acibenzolar may
function similar to SA to activate NPR1 and SAR. Although the ASM control effect was obvious
within 4 hpt (Figure 1; Figure 2) and SAR-related gene expression was also induced within 4 hpt
(Figure 5), acibenzolar was not detected in untreated leaves 1 dpt (Ishii et al., 2002). Therefore, it is
tempting to speculate that putative signal molecules are rapidly translocated from treated to untreated
leaves. Several candidates for this long-distance signal have been identified, including methyl
salicylate (MeSA), an SFD1/GLY1-derived G3P-dependent signal, the lipid-transfer protein DIR1,
the dicarboxylic acid azelaic acid (AzA), the abietane diterpenoid dehydroabirtinal (DA), jasmonic
acid (JA), and the amino acid derivative pipecolic acid (Pip) (Dempsey and Klessig, 2012). Among
them, Pip acts as primary regulator of SAR (Návarová et al., 2012; Zeier, 2013). In *Arabidopsis*
thaliana, these responses include camalexin (phytotoxin) accumulation and expression of defense-
related genes, including *ALD1*, *FMO1*, and *PR1* (Bernsdorff et al., 2016). Further analysis will be
needed to elucidate the mobile signals which induce SAR in Japanese radish.

ASM induced stomatal-based defense against *Pcal* within 4 hpt in both cabbage and Japanese radish
(Figure 3; Ishiga et al., 2020). We also showed that ASM-triggered stomatal closure was suppressed
by a peroxidase inhibitor, SHAM but not by an NADPH oxidase inhibitor, DPI (Figure 4;
Supplementary Figure 3). Khokon et al. (2011) reported that SA-induced stomatal closure
accompanied ROS production mediated by peroxidase in *A. thaliana*. Therefore, these results
indicated that ASM-induced stomatal closure is closely related to ROS production through
peroxidase. Our results also showed that ASM induced stomatal-based defense was observed not
only on treated leaves, but also on untreated leaves (Figure 3). Lin and Ishii (2009) demonstrated that
rapid H_2O_2 accumulation occurred in cucumber stem xylem fluids during ASM-induced SAR.

Takahashi and Shinozaki (2019) reported that light stress-mediated ABA accumulation induced the
ROS/Ca^{2+} wave propagates from local leaves to systemic leaves with a rapid transmission speed,
suggesting that local ROS/Ca^{2+} wave functions as long-distance signal to activate stomatal closure
during light stress conditions. Therefore, it is possible that ROS induced by ASM treatment activates
stomatal closure in a systemic response during pathogen infection.

Based on our results, we propose a stomatal closure model induced by ASM treatment (Figure 6).
ASM induces ROS production mediated by SHAM-sensitive peroxidase, leading to stomatal closure.
Then, local ROS functions as a long-distance signal to induce stomatal closure in a systemic response
after ASM treatment. In addition to ROS, it is expected that some mobile signals are rapidly
transported and induce ROS production, and in turn stomatal closure. Further analysis will be needed
to understand the ASM-induced SAR mechanism.

We clearly demonstrated that ASM dip-treatment protects Japanese radish plants against *Pcal,* a
causal agent of bacterial blight disease, by activating SAR, such as stomatal-based defense.
Furthermore, we observed that ASM induced defense response was acquired not only on treated
leaves, but also untreated upper and lower leaves. These results highlight the role of ASM as an
efficient sustainable disease control strategy.

Acknowledgments

262　We thank Dr. Christina Baker for editing the manuscript. *Pcal* was kindly provided by the Nagano
263　Vegetable and Ornamental Crops Experiment Station, Nagano, Japan. This work was supported in
264　part by the JST ERATO NOMURA Microbial Community Control Project, JST, Japan.

References

Bernsdorff, F., Döring, A.C., Gruner, K., Schuck, S., Bräutigam, A., and Zeier, J. (2016). Pipecolic
acid orchestrates plant systemic acquired resistance and defense priming via salicylic acid-dependent
and-independent pathways. *Plant Cell* 28, 102–129. doi: 10.1105/tpc.15.00496

Chitrakar, R., and Melotto, M. (2010) Assessing stomatal response to live bacterial cells using whole
leaf imaging. *J. Vis. Exp*. 44:2185. doi: 10.3791/2185

Cools, H.J., and Ishii, H. (2002). Pre-treatment of cucumber plants with acibenzolar-S-methyl
systemically primes a *phenylalanine ammonia lyase* gene (*PAL1*) for enhanced expression upon
attack with a pathogenic fungus. *Physiol. Mol. Plant Pathol*. 61, 273–280. doi:
10.1006/pmpp.2003.0439

Dempsey, D.A., and Klessig, D.F. (2012). SOS - too many signals for systemic acquired resistance?
Trends Plant Sci. 17, 538–545. doi: 10.1016/j.tplants.2012.05.011

Doke, N. (1983). Generation of superoxide anion by potato tuber protoplasts during the
hypersensitive response to hyphal wall components of *Phytophthora infestans* and specific inhibition

of the reaction by suppressors of hypersensitivity. *Physiol. Plant Pathol.* 23, 359–367. doi:
10.1016/0048-4059(83)90020-6

Fobert, P.R., and Després, C. (2005). Redox control of systemic acquired resistance. *Curr. Opin.*
Plant Biol. 8, 378–382. doi: 10.1016/j.pbi.2005.05.003

Ishiga, T., Iida, Y., Sakata, N., Ugajin, T., Hirata, T., Taniguchi, S., et al. (2020). Acibenzolar-S-
methyl activates stomatal-based defense against *Pseudomonas cannabina* pv. *alisalensis* in cabbage.
J. Gen. Plant Pathol. 48–54. doi: 10.1007/s10327-019-00883-5

Ishiga, Y., and Ichinose, Y. (2016). *Pseudomonas syringae* pv. *tomato* OxyR is required for virulence
in tomato and *Arabidopsis*. *Mol. Plant Microbe Interact.* 29, 119–131. doi: 10.1094/MPMI-09-15-
0204-R

Ishii, H., Kato, T., Yamanaka, M., Cools, H., and Sugiyama, T. (2002) Induction activity of systemic
disease resistance and metabolism of acibenzolar-S-methyl in cucumber (abstract in Japanese). Abstr.
27th Ann. Meet. Pesticide Sci. Soc. Jpn.

Joshi-Saha, A., Valon, C., and Leung, J. (2011). A brand new START: abscisic acid perception and
transduction in the guard cell. *Sci. Signal.* 4:re4. doi: 10.1126/scisignal.2002164

Jung, H.W., Tschaplinski, T.J., Wang, L., Glazebrook, J., and Greenberg, J.T. (2009). Priming in
systemic plant immunity. Science. 324:89–91. doi: 10.1126/science.1170025

Khokon, A.R., Okuma, E., Hossain, M.A., Munemasa, S., Uraji, M., Nakamura, Y. et al. (2011).
Involvement of extracellular oxidative burst in salicylic acid-induced stomatal closure in
Arabidopsis. *Plant Cell Environ.* 34, 434–443. doi: 10.1111/j.1365-3040.2010.02253.x

Kim, T.H., Böhmer, M., Hu, H., Nishimura, N., and Schroeder, J.I. (2010). Guard cell signal
transduction network: advances in understanding abscisic acid, CO_2, and Ca^{2+} signaling. *Annu. Rev.*
Plant Biol. 61, 561–591. doi: 10.1146/annurev-arplant-042809-112226

King, E.O., Ward, M.K., and Raney, D.E. (1954). Two simple media for the demonstration of
pyocyanin and fluorescin. *J. Lab. Clin. Med.* 44, 301–307.

Kollist, H., Nuhkat, M., and Roelfsema, M.R. (2014). Closing gaps: linking elements that control
stomatal movement. *New Phytol.* 203, 44–62. doi: 10.1111/nph.12832

Kouzai, Y., Kimura, M., Watanabe, M., Kusunoki, K., Osaka, D., Suzuki, T. et al. (2018a). Salicylic
acid-dependent immunity contributes to resistance against *Rhizoctonia solani*, a necrotrophic fungal
agent of sheath blight, in rice and *Brachypodium distachyon*. *New Phytol.* 217, 771–783. doi:
10.1111/nph.14849

Kouzai, Y., Noutoshi, Y., Inoue, K., Shimizu, M., Onda, Y., and Mochida, K. (2018b).
Benzothiadiazole, a plant defense inducer, negatively regulates sheath blight resistance in
Brachypodium distachyon. *Sci. Rep.* 8, 17358. doi: 10.1038/s41598-018-35790-w

Kubota, M., and Abiko, K. (2000). Induced resistance in hypocotyl of cucumber by infection with
Colletotrichum lagenarium in leaves. *J. Gen. Plant Pathol.* 66, 128–131. doi: 10.1007/PL00012933

Kunz, W., Schurter, R., and Maetzke, T. (1997). The chemistry of benzothiadiazole plant activators.
Pest. Sci. 50, 275–282. doi: 10.1002/(SICI)1096-9063(199708)50:4<275::AID-PS593>3.0.CO;2-7

Lin, T., and Ishii, H. (2009). Accumulation of H_2O_2 in xylem fluids of cucumber stems during ASM-
induced systemic acquired resistance (SAR) involves increased LOX activity and transient
accumulation of shikimic acid. *Eur. J. Plant Pathol.* 125, 119–130. doi: 10.1007/s10658-009-9464-9

Mori, I.C., Pinontoan, R., Kawano, T., and Muto, S. (2001). Involvement of superoxide generation in
salicylic acid-induced stomatal closure in *Vicia faba*. *Plant Cell Physiol.* 42, 1383–1388. doi:
10.1093/pcp/pce176

Nakashita, H., Yasuda, M., Nishioka, M., Hasegawa, S., Arai, Y., Uramoto, M., et al. (2002).
Chloroisonicotinamide derivative induces a broad range of disease resistance in rice and tobacco.
Plant Cell Physiol. 43, 823–831. doi: 10.1093/pcp/pcf097

Narusaka, Y., Narusaka, M., Horio, T., and Ishii, H. (1999). Induction of disease resistance in
cucumber by acibenzolar-S-methyl and expression of resistance-related genes. *Ann. Phytopath. Soc.*
Japan. 65, 119–122.

Narusaka, Y., Narusaka, M., and Ishii, H. (2001). Searching by differential display for new genes
implicated in induced disease resistance of cucumber treated with acibenzolar-S-methyl. *J. Gen.*
Plant Pathol. 67, 207–211. doi: 10.1007/PL00013013

Návarová, H., Bernsdorff, F., Döring, A.C., and Zeier, J. (2012). Pipecolic acid, an endogenous
mediator of defense amplification and priming, is a critical regulator of inducible plant immunity.
Plant Cell 24, 5123–5141. doi: 10.1105/tpc.112.103564

Oostendorp, M., Kunz, W., Dietrich, B., and Staub, T. (2001). Induced disease resistance in plants by
chemicals. *Eur. J. Plant Pathol.* 107, 19–28. doi: 10.1023/A:1008760518772

Park, P., Kurihara, T., Nita, M., Ikeda, K., Inoue, K., and Ishii, H. (2019). Application of energy-
filtering electron microscopy (EFEM) for analysis of hydrogen peroxide and lignin in epidermal
walls of cucumber leaves triggered by acibenzolar-S-methyl treatment prior to inoculation with
Colletotrichum orbiculare. *Plant Path.* 68, 1624–1634. doi: 10.1111/ppa.13095

Prodhan, M.Y., Issak, M., Nakamura, T., Munemasa, S., Nakamura, Y., and Murata, Y. (2017).
Chitosan signaling in guard cells requires endogenous salicylic acid. *Biosci. Biotechnol. Biochem.* 81,
1536–1541. doi: 10.1080/09168451.2017.1332979

Sakata, N., Ishiga, T., Saito, H., Nguyen, V.T., and Ishiga, Y. (2019). Transposon mutagenesis
reveals *Pseudomonas cannabina* pv. *alisalensis* optimizes its virulence factors for pathogenicity on
different hosts. *PeerJ*. 7:e7698. doi: 10.7717/peerj.7698. eCollection 2019

Shah, J., and Zeier, J. (2013). Long-distance communication and signal amplification in systemic
acquired resistance. *Front. Plant Sci.* 4, 30. doi: 10.3389/fpls.2013.00030. eCollection 2013

Takahashi, F., Ochiai, M., Ikeda, K., and Takikawa, Y. (2013). Streptomycin and copper resistance in
Pseudomonas cannabina pv. *alisalensis* (abstract in Japanese). Jpn. J. Phytopathol. 35.

Takahashi, F., and Shinozaki, K. (2019). Long-distance signaling in plant stress response. *Curr.*
Opin. Plant Biol. 47, 106–111. doi: 10.1016/j.pbi.2018.10.006

Takikawa, Y., and Takahashi, F. (2014). Bacterial leaf spot and blight of crucifer plants
(Brassicaceae) caused by *Pseudomonas syringae* pv. *maculicola* and *P. cannabina* pv. *alisalensis*. *J.*
Gen. Plant Pathol. 80, 466–474. doi: 10.1007/s10327-014-0540-4

Tripathi, D., Jiang, Y.L., and Kumar, D. (2010). SABP2, a methyl salicylate esterase is required for
the systemic acquired resistance induced by acibenzolar-S-methyl in plants. *FEBS Lett.* 584, 3458–
3463. doi: 10.1016/j.febslet.2010.06.046

Wang, Y., Schuck, S., Wu, J., Yang, P., Döring, A.C., Zeier, J., et al. (2018). A MPK3/6-WRKY33-
ALD1-pipecolic acid regulatory loop contributes to systemic acquired resistance. *Plant Cell* 30,
2480–2494. doi: 0.1105/tpc.18.00547

Yasuda, M., Ishikawa, A., Jikumaru, Y., Seki, M., Umezawa, T., Asami, T., et al. (2008).
Antagonistic interaction between systemic acquired resistance and the abscisic acid-mediated abiotic
stress response in *Arabidopsis*. *Plant Cell* 20, 1678–1692. doi: 10.1105/tpc.107.054296

Zeier, J. (2013). New insights into the regulation of plant immunity by amino acid metabolic
pathways. *Plant Cell Environ.* 36, 2085–2103. doi: 10.1111/pce.12122

Figure legends

Figure 1. Lesion area (%) of Japanese radish inoculated with *Pcal* after dip-treatment with
ASM on fourth leaves. Greenhouse grown Japanese radish plants were spray-inoculated with *Pcal*
(5×10^7 CFU/ml) 4 h (**A**), 1 d (**B**), and 1 w (**C**) after ASM dip-treatment (100 ppm) on fourth leaves.
Disease symptoms were monitored by measuring lesion areas at 1-week post inoculation (wpi).
Vertical bars indicate the standard error for four independent replications. Asterisks indicate a
significant difference from the water treatment control in a *t*-test (*$p < 0.05$; ** $p < 0.01$).

Figure 2. Bacterial populations of *Pcal* inoculated plants after ASM dip-treatment. Three-week-
old Japanese radish plants were spray-inoculated with *Pcal* suspensions (5×10^7 CFU/ml) 4 h (**A**), 1
d (**B**), and 1 w (**C**) after ASM dip-treatment (100 ppm) on fourth leaves. Bacterial populations were
determined by dilution plating on selective medium as described in the methods section at 1-week
post inoculation (wpi). Vertical bars indicate the standard error for three independent experiments.
Asterisks indicate a significant difference from the water treatment control in a *t*-test (*$p < 0.05$; ** p
< 0.01).

Figure 3. Stomatal aperture width (μm) in Japanese radish plants dip inoculated with *Pcal*
suspensions (1×10^8 CFU/ml) after ASM treatment. Japanese radish leaves were inoculated with
Pcal 4 h (**A–C**), 1 d (**D–F**), and 1 w (**G–I**) after ASM dip-treatment (100 ppm) on fourth leaves.
Stomatal aperture width (μm) was measured in third (**A, D, G**), fourth (**B, E, H**), and fifth (**C, F, I**)
leaves. Each time point after ASM treatment, radish leaves were dip-inoculated, then imaged using a
Nikon optical microscope (Eclipse 80i). In all bar graphs, vertical bars indicate the standard error for
three biological replicates. Significant differences ($p < 0.05$) are indicated by different letters based
on a Tukey's honestly significant difference (HSD) test.

Figure 4. Stomatal aperture width (µm) in Japanese radish plants treated with SHAM and DPI
after ASM treatment. Japanese radish leaves were treated with SHAM (1 mM) and DPI (10 µM) 1
w after ASM dip-treatment (100 ppm) on fourth leaves. Stomatal aperture width (µm) was measured
on third **(A)**, fourth **(B)**, and fifth **(C)** leaves using a Nikon optical microscope (Eclipse 80i). In all
bar graphs, vertical bars indicate the standard error for three biological replicates. Significant
differences ($p < 0.05$) are indicated by different letters based on a Tukey's honestly significant
difference (HSD) test.

Figure 5. Gene expression profiles involved in Japanese radish plant defense after ASM
treatment. Three-week-old Japanese radish plants were treated with water as a control and ASM
(100 ppm). After 4 h expression of *PR1* **(A)**, *PR2* **(B)**, and *PR3* **(C)** was determined using real-time
quantitative reverse transcription-polymerase chain reaction (RT-qPCR) with gene-specific primer
sets. Expression was normalized using *GAPDH*. Vertical bars indicate the standard error for three
biological replicates. Asterisks indicate a significant difference from the water control in a *t*-test (*p
< 0.05; ** $p < 0.01$).

Figure 6. Proposed model of stomatal closure induced by ASM treatment by inducing
peroxidase-dependent ROS production. ASM induces stomatal closure by inducing ROS
production through peroxidase. Then, local ROS and some mobile signals function as long-distance
signals to induce stomatal closure in a systemic response after ASM treatment. X indicates mobile
signals.

A

B

C

A

B

C

A

B

C

Figure 5

Figure 6

Systemic
responses

Mobile signals

Peroxidase

SHAM

ROS

Stomatal
closure

3rd & 5th leaves

Local
responses

ASM

Peroxidase

SHAM

ROS

Stomatal
closure

4th leaf