1 **Single cell resolution regulatory landscape of the mouse kidney highlights cellular**
2 **differentiation programs and renal disease targets**
3
4

5 Zhen Miao[1,2,3,*], Michael S. Balzer[1,2,*], Ziyuan Ma[1,2], Hongbo Liu[1,2], Junnan Wu[1,2], Rojesh
6 Shrestha[1,2], Tamas Aranyi[1,2], Amy Kwan[4], Ayano Kondo[4], Marco Pontoglio[5], Junhyong Kim[6],
7 Mingyao Li[7], Klaus H. Kaestner[2,4] and Katalin Susztak[1,2,4]
8
9
[1]Renal, Electrolyte, and Hypertension Division, Department of Medicine, University of
Pennsylvania, Perelman School of Medicine, Philadelphia, PA 19104, USA
[2]Institute for Diabetes, Obesity, and Metabolism, University of Pennsylvania, Perelman School of
Medicine, Philadelphia, PA 19104, USA
[3]Graduate Group in Genomics and Computational Biology, University of Pennsylvania, Perelman
School of Medicine, Philadelphia, PA 19104, USA
[4]Department of Genetics, University of Pennsylvania, Perelman School of Medicine,
Philadelphia, PA 19104, USA
[5]Epigenetics and Development Laboratory, Université de Paris Inserm U1151/CNRS UMR
8253, Institut Necker Enfants Malades, Paris, France
[6]Department of Biology, University of Pennsylvania, Philadelphia, PA 19104, USA
[7]Department of Epidemiology and Biostatistics, University of Pennsylvania, Perelman School of
Medicine, Philadelphia, PA 19104, USA
[#]These two authors contributed equally to this work
24
Correspondence:
Katalin Susztak, MD, PhD
Professor of Medicine
University of Pennsylvania, Perelman School of Medicine
3400 Civic Center Blvd,
Smilow Translational building 12-123,
Philadelphia, PA 19104
Phone: (215) 898-2009
ksusztak@pennmedicine.upenn.edu

Grant support:
Work in the Susztak lab is supported by the NIH DK076077, DK087635, and DK105821. MSB is
supported by a German Research Foundation grant (BA 6205/2-1).

Abstract

The kidney has a very limited capacity to repair. Defining cellular differentiation during development therefore could aid the understanding of homeostatic and maladaptive regeneration.

Here, we profiled open chromatin and gene expression in developing and adult mouse kidneys at single cell resolution. We show critical reliance of gene expression on distal regulatory elements (enhancers). We define key cell type-specific transcription factors and major gene-regulatory circuits for kidney cells. Dynamic chromatin and expression changes during nephron progenitor differentiation demonstrated that podocyte commitment occurs early and is associated with sustained *Foxl1* expression. Renal tubule cells followed a more complex differentiation, where *Hfn4a* was associated with proximal and *Tfap2b* with distal fate. Mapping single nucleotide variants associated with human kidney disease identified critical cell types, developmental stages, genes, and regulatory mechanisms.

We provide a global single cell resolution view of chromatin accessibility of kidney development. The dataset is available via an interactive public website.

Keywords

Kidney development; differentiation; single cell; chromatin accessibility; transcription factor; gene regulatory network; cis-regulatory elements; enhancer; kidney disease.

Introduction

The mammalian kidney maintains fluid, electrolyte, and metabolite balance of the body and plays an essential role in blood pressure regulation and red blood cell homeostasis. The human kidney makes roughly 180 liters of primary filtrate each day that is then reabsorbed and modified by a long tubule segment. To perform this highly choreographed and sophisticated function, the kidney contains close to 20 highly specialized epithelial cells. The renal glomerulus acts as a 60 kD size-selective filter. The proximal part of the tubules is responsible for reclaiming more than 70% of the primary filtrate, which is done via unregulated active and passive paracellular transport [1], while the loop of Henle plays an important role in concentrating the urine. The distal convoluted tubule is critical for regulated electrogenic sodium reabsorption and potassium secretion. The last segment of kidney tubules is the collecting duct where the final concentration of the urine is determined via regulation of water channels as well as acid or base secretion. Understanding the development of these diverse cell types in the kidney is essential to understand kidney homeostasis, disease, and regeneration.

The mammalian kidney develops from the intermediate mesoderm via a complex interaction between the ureteric bud and the metanephric mesenchyme [2]. In the mouse kidney, Six2 marks the self-renewing nephron progenitor population [3]. The nephron progenitors commit and undergo a mesenchymal-to-epithelial transformation giving rise to the renal vesicle [3]. The renal vesicle then undergoes segmentation and elongation, giving rise to epithelia from the podocytes to the distal convoluted tubules, while the ureteric bud becomes the collecting duct. Unbiased and hypothesis-driven studies have highlighted critical stages and drivers of early kidney development [4], that have been essential for the development of *in vitro* kidney organoid differentiation protocols [5-7]. However, cells in organoids are still poorly differentiated, improving cellular differentiation and maturation of these structures remains a major challenge [8]. Thus, the understanding of late kidney development, especially the cell type-specific driver transcription factors (TFs) is of great importance [9-11]. Alteration in Wnt, Notch, Bmp, and Egf signaling significantly impacts cellular differentiation, but only a handful of TFs that directly drive the differentiation of distinct segments have been identified, such as *Pou3f3*, *Lhx1*, *Irx2*, *Foxc2* and *Mafb* [12]. Further understanding of the terminal differentiation program could aid the understanding of kidney disease development.

While single cell RNA sequencing (scRNA-seq) has improved our understanding of kidney

development in mice and humans [9,10,13,14], it provides limited information of TFs, which are usually

lowly expressed. Equally difficult is to understand how genes are regulated from scRNA-seq data

alone. Chromatin state profiles, on the other hand, provide valuable insight to gene regulation

mechanisms during cell differentiation, since they show not only the accessibility of the gene

transcription start site (TSS), but also of distal regulatory regions such as enhancers. It is believed

that enhancers are critical for establishing the cell type-specific gene expression pattern, but it has

not been shown conclusively on a single cell level. Together with gene expression, open chromatin

profiles can define the gene regulatory logic, which is the fundamental element of cell identity.

However, there is a scarcity of open chromatin information by Assay for Transposase-Accessible

Chromatin using sequencing (ATAC-seq) or chromatin immunoprecipitation (ChIP) data by ChIP-

seq related to kidney development. In addition, epigenetic changes observed in bulk analyses

mostly represent changes in cell composition, rather than cell type-specific changes [15], making it

challenging to interpret bulk ATAC-seq data.

To this end, here we generated a single cell open chromatin and corresponding expression survey

for the developing and adult mouse kidney, which will be available for the community via a

searchable website (**http://susztaklab.com/VisCello_snATAC/** and **http://susztaklab.com/igv/**).

Using this atlas, we have produced a new epigenome-based classification of developing and

mature cells and defined cell type-specific regulatory networks. We also investigated key TFs and

cell-cell interactions associated with developmental cellular transitions. Finally, we used the single

cell open chromatin information to pinpoint putative target genes and cell types of several chronic

kidney disease noncoding genome-wide association study (GWAS) loci.

Results

Single cell accessible chromatin landscape of the developing and adult mouse kidneys

To characterize the accessible chromatin landscape of the developing and adult mouse kidneys at

single cell resolution, we performed single nuclei ATAC-seq (snATAC-seq) on kidneys of mice

on postnatal day 0 (P0) at 3 and 8 weeks of age (**Figure 1a, Methods**). In parallel, we also

performed bulk (whole kidney) ATAC-seq analysis at matched developmental stages. Following

sequencing, we aggregated all high-quality mapped reads in each sample irrespective of barcode.

The combined snATAC-seq dataset from all samples showed the expected insert size periodicity

(**Figure S1a**) with a strong enrichment of signal at TSSs (**Figure S1b**), indicating high data quality.

The snATAC-seq data showed high concordance with the bulk ATAC data (Spearman correlation

coefficient >0.84, **Methods**, **Figure S1c**).

We next revealed cell type annotations from the open chromatin information. After conducting

stringent filtering of the number of barcodes, promoter ratio and mitochondria ratio (**Methods,**

Figure S1d), we kept 28,316 cells across the samples (**Figures 1b, S1f-g**). Cells were then

clustered using snapATAC [16], which binned the whole genome into 5 kb regions and used diffusion

map and principal component analysis for dimension reduction (**Methods**). Prior to clustering, we

used Harmony [17], an iterative batch correction method, to correct for variability across samples.

Using batch-corrected low dimensional embeddings, we clustered all cells together and retained

13 clusters, all of which had consistent representation across the number of peaks, samples and

read depth profiles (**Figures 1b, S1e-g**). As expected, some clusters such as nephron progenitors

and stromal cells were enriched in the developing kidney (P0).

In order to identify the cell type-specific open chromatin regions, we conducted peak calling using

MACS2 [18] on each cell type separately. The peaks were then merged to obtain a comprehensive

open chromatin set. We found that the single nuclei open chromatin set showed good concordance

with bulk ATAC-seq samples, with most of the peaks in bulk ATAC-seq data captured by the

single nuclei data. On the other hand, single nuclei chromatin accessibility data showed roughly

50% more accessible chromatin peaks (total of 300,693 peaks) than the bulk ATAC-seq data

(**Figure 1e, Methods**), indicating that the snATAC-seq data was particularly powerful in
identifying open chromatin areas that are accessible in single cell types.

To determine the cell types represented by each cluster, we examined chromatin accessibility
around the TSS and gene body regions of the cognate known cell type-specific marker genes [19].
Based on the accessibility of the known marker genes, we identified clusters representing nephron
progenitors, endothelial cells, podocytes, proximal tubule segment 1 and segment 3 cells, loop of
Henle, distal convoluted tubule, connecting tubule, collecting duct principal cells, collecting duct
intercalated cells, stromal and immune cells (**Figure 1b**). **Figures 1d and S1h** show chromatin
accessibility information for key cell type marker genes, such as *Uncx* and *Cited1* for nephron
progenitors, *Nphs1* and *Nphs2* for podocytes, *Akr1c21* for both segments of proximal tubules,
Slc34a1 and *Slc5a2* for segment 1 of proximal tubules, *Kap* for segment 3 of proximal tubules,
Slc12a1 and *Umod* for loop of Henle, *Scl12a3* and *Pvalb* for distal convoluted tubule, *Trpv5* for
connecting tubule, *Aqp2* and *Fxyd4* for principal cells, *Atp6v1g3* and *Atp6v0d2* for intercalated
cells, *Egfl7* for endothelial cells, *C1qb* for immune cells and *Col3a1* for different types of stromal
cells, respectively [19].

To understand cell type-specific gene expression changes, we also generated a single cell RNA
sequencing (scRNA-seq) atlas for mouse kidney samples at the same developmental stages. The
single cell transcriptome profiles of P0 and adult mouse kidneys were derived and processed as
described in **Methods**. Rigorous quality control yielded a set of 43,636 single cells (**Figures 1b,**
S1i). Quality control metrics such as gene counts, UMI counts and mitochondrial gene percentage
along with batch correction results are shown in **Figures S1j-m**. By unbiased clustering [20] we
obtained 17 distinct cell populations in the combined P0 and adult mouse datasets (**Figure S1i**).
On the basis of marker gene expression, we identified kidney epithelial, immune and endothelial
cells (**Figures 1f, S1n-o**), closely resembling the clustering obtained from snATAC-seq analysis.
We then conducted differential expression analysis on the clusters and identified key marker genes
for each cell type (**Supplemental Table 1**).

To compare the consistency between cluster assignment in the snATAC-seq data and the scRNA-
seq data, we next derived a gene activity score for the top 3,000 highly variable genes in each

snATAC-seq cluster and computed the Pearson's correlation coefficient between each snATAC

cluster and scRNA cluster (**Methods**). This analysis indicated good concordance between the two

datasets (**Figures 1g, S1p**). While the correlation between gene expression and inferred gene

activity score was high, we noted some differences in cell proportions, which was mostly related

to the sample preparation-induced cell drop-out (**Figures S1g, i**). Consistent with previous

observations that single cell preparations better capture immune cells than single nuclear

preparations [21], we noted that the immune cell repertoire was limited in the snATAC-seq dataset;

on the other hand, stromal cells were better captured by the nuclear preparation.

Finally, to allow the interactive use of this dataset by the community, we not only made the raw

data   available   but   also   the   processed   dataset   via   our   searchable   website

(**http://susztaklab.com/VisCello_snATAC/** and **http://susztaklab.com/igv/**). For example, here

we show the chromatin accessibility landscape of *Ace2*, which is of major interest currently due to

the COVID19 epidemic. We can observe an open chromatin region around the transcription start

site of *Ace2* only in proximal tubules, which is consistent with its expression in proximal tubules

(**Figure S1q**).

Characterization of the cell type-specific regulatory landscape

To characterize different genomic elements captured by snATAC-seq data, we first stratified the

genome into promoters, exons, 5' and 3' untranslated regions, introns, and distal regions using the

GENCODE annotation [22] (**Methods**). We noticed that consistent with bulk ATAC-seq data, most

peaks in snATAC-seq data were in regions characterized as distal elements or introns, (**Figure**

S2a) and relatively small portions (<10%) were in promoter or 5' untranslated regions. The

genomic elements proportion was stable across developmental stages. In addition, almost half of

the open chromatin peaks overlapped with P0 or adult H3K27Ac ChIP-seq signals (**Figure S2b**),

indicating the contribution of enhancer regions to accessible chromatins.

To study the open chromatin heterogeneity in different cell types, we derived a cell type-specific

accessible chromatin landscape by conducting pairwise Fisher's exact test for each peak between

every cluster (Benjamini-Hochberg adjusted q value ≤ 0.05, **Methods**). In total, we identified

60,684 differentially accessible open chromatin peaks (DAPs) across the 13 cell types

(**Supplemental Table 2, Figure 2a**). Among these peaks, most showed high specificity for a single

cluster. However, we noticed overlaps between the S1 and S3 proximal tubule segments-specific

peaks, as well as between the loop of Henle and distal convoluted tubule segments, which is

consistent with their biological similarities. In addition to the cell type-specific peaks, we also

found some cell-type independent open chromatin areas (present across nephron progenitors,

podocytes, proximal tubule and loop of Henle cells), likely consistent with basal housekeeping

genes and regulatory elements. (**Figures 2a, S2c**).

We noticed that many genes had strong cell type-specific DAPs at their TSS. Other genes, however,

had accessible chromatin at their TSS in multiple cell types. For example, *Umod*, the loop of

Henle-specific marker gene, showed accessible chromatin at its TSS at multiple tubule cell types

(**Figure S2d, S1h**). Rather than with its TSS, cell type-specific chromatin accessibility of *Umod*

strongly correlated with an upstream open chromatin peak, which is likely an enhancer region,

indicated by the H3K27Ac ChIP-seq signal (**Figure S2d**). Consistently, we noticed the enrichment

of intronic regions and distal elements (**Figures S2e-f**) in cell type-specific DAPs, indicating their

role in cell type-specific gene regulation.

These observations motivated us to study cis-regulatory elements using the snATAC-seq data and

scRNA-seq data. We reasoned that a subset of the cell type-specific cis-regulatory elements should

regulate cell type-specific gene expression in cis. Inspired by [23], we aligned DAPs and

differentially expressed genes from our snATAC-seq and scRNA-seq datasets, and inferred the

putative regulatory peaks by their proximity (**Methods**). Such cis-regulatory elements predictions

were confirmed by comparing with cis-regulatory elements inferred previously [24], as we

recapitulated roughly 20% of elements from their analysis. In addition, our analysis was able to

identify several known enhancers such as for *Six2* and *Slc6a18* [24,25] (**Figure S2g**). To quantify the

contribution of cis-regulatory elements, we analyzed peak co-accessibility patterns using Cicero

[26]. By using a heuristic co-accessible score 0.4 as a cutoff, we identified 1,214,638 and 926,288

cis-regulatory element links in the P0 and adult data, respectively. Some of these are likely

promoter-enhancer regulatory units. Among these co-accessible elements, only 149,389 were

common in P0 and adult kidneys, while most were developmental stage-dependent. While this

observation needs further experimental validation, it highlights dynamic changes in gene
regulation during development, which includes both chromatin opening and looping.

Given the large contribution of distal regulatory elements to chromatin accessibility, we next
looked into identifying key TFs that occupy the cell type-specific open chromatin regions. Until
now, information on cell type-specific TFs in the kidney has been scarce. Therefore, we performed
motif enrichment analysis on the cell type-specific open chromatin regions using HOMER [27].
HOMER was designed as a differential motif discovery algorithm that scores motifs by computing
enrichment of motif sequences in target compared to a reference set. To reduce false discovery,
we focused on the known motifs. The full list of cell type-specific TF binding motifs is shown in
Supplemental Table 3. Since several TFs have identical or similar binding sequences, we next
correlated motif enrichment with scRNA-seq TF expression. Using this combined motif
enrichment and gene expression approach, we have defined the mouse kidney cell type-specific
TF landscape. Examples include *Six2* and *Hoxc9* in nephron progenitors, *Wt1* and *Mafb* in
podocytes, *Hnf4a*, *Ppara*, and *Bhle41* in proximal tubules, *Esrrb* and *Foxa1* in loop of Henle, *Vdr*
in distal convoluted tubule, *Elf5* in principal cells, *Tcfcp2l1* in intercalated cells, *Erg* and *Sox17* in
endothelial cells, *Spi1* and *Batf* in immune cells, and *Twist1* and *Nr2f2* in stromal cells (**Figures
2a, S2h**).

In order to study the putative target genes of TFs, we examined TF regulon activity using Single-
Cell rEgulatory Network Inference and Clustering (SCENIC) [28]. SCENIC was designed to reveal
TF-centered gene co-expression networks. By inferring a gene correlation network followed by
motif-based filtration, SCENIC keeps only potential direct targets of each TF as modules
(regulons). The activity of each regulon in each cell was quantified and then binarized to "on" or
"off" based on activity distribution across cells (**Methods**). SCENIC was also able to conduct
clustering based on the regulon states of each cell. SCENIC results (**Figure 2b**) indicated strong
enrichment in *Trps1*, *Hnf1b*, *Maf*, *Hnf1a*, and *Hnf4a* regulon activity in proximal tubules, *Hmga2*,
Hoxc6, *Hoxd11*, *Meox1*, *Six2*, *Tcf4*, and *Uncx* in nephron progenitors, *Esrrg*, and *Ppargc1a* in loop
of Henle, *Hmgb3* in proliferating cells and *Foxc1*, *Foxc2*, *Foxd1*, *Lef1*, and *Mafb* in podocytes,
respectively. While the expression of several of these TFs was relatively low and was further
exacerbated by transcript drop-outs, many TFs did not show strong cell type enrichment. The

regulon-based analysis, however, showed a very clear enrichment. SCENIC also successfully
reported multiple downstream target genes. The full list of regulons and their respective target
genes can be found in **Supplemental Table 4,** scaled and binarized regulon activity is also
available in **Supplemental Table 5.** Examples of regulon activity, corresponding TF expression,
and target gene expression are depicted in **Figures 2c, S2i**. For example, TFs such as *Eya1*, *Hoxc8*,
Hoxc9, *Pax2*, *Spock2* and *Wnt4* are important downstream targets within the regulon of nephron
progenitor-specific TF *Uncx*, indicating an important transcriptional hierarchy of nephron
development [29].

In summary, we generated a comprehensive atlas for the cell type-specific regulatory elements and
TF-centered regulatory network.

The regulatory trajectory of nephron progenitor differentiation

All cells in the body differentiate from the same genetic template. Cell type-specific chromatin
opening and closing events associated with TF binding changes set up the cell type-specific
regulatory landscape resulting in cell type specification and development. We found that closing
of open chromatin regions was the predominant event during the nephron progenitor
differentiation (**Figure S2a**). We then evaluated the cellular differentiation trajectory in the
snATAC-seq and scRNA-seq datasets (**Methods**). We identified multiple nephron progenitor sub-
groups (**Figures 3a-b**), which will need to be carefully mapped to prior gene expression- and
anatomical location-driven nephron progenitor sub-classification. Consistently, across both data
modalities, we identified that the podocyte precursors differentiated early from the nephron
progenitor pool (**Figures 3a-b**). The tubule cell trajectory was more complex with a shared
intermediate stage and later differentiation into proximal tubules and distal tubules/loop of Henle
(**Figures 3a-b, S3a-c**). We also integrated snATAC-seq and scRNA-seq data to obtain a single
trajectory (**Methods**). The cell types in this dataset were correctly mapped and the trajectory
resembled the path observed in individual analyses of the scRNA and snATAC datasets (**Figures
S3e-g**). The robustness of developmental trajectories was further supported by obtaining similar
results when performing RNA velocity analysis using Velocyto [30] (**Figure S3d**) and by comparing
with previous human and mouse kidney developmental studies [9,13,14].

Building on both the SCENIC-generated gene regulatory network and the robust differentiation trajectories of the snATAC-seq and scRNA-seq datasets, we next aimed to understand chromatin dynamics, identify TFs and driver pathways for cell type specification and differentiation. To this end, we first determined variation in chromatin accessibility along the 3 differentiation trajectories using ChromVAR [31]. ChromVAR estimates the accessibility dynamics of motifs in snATAC-seq data (**Methods**). We observed three different patterns when analyzing genes of interest (**Figure 3c**): *1) Decrease of TF motif accessibility in all lineages.* For example, *Sox11* motif enrichment score was high in nephron progenitor cells at the beginning of all 3 trajectories. It then decreased in all 3 lineages in parallel, underlining the role of *Sox11* in early kidney development. Several other TFs followed this pattern such as *Six2* and *Sox9. 2) Cell type-specific maintenance of chromatin accessibility with advancing differentiation.* We observed that chromatin accessibility for the *Wt1* motif was high initially, but declined in proximal tubule and loop of Henle lineages, while its expression increased in the podocyte lineage. This is consistent with the important role of *Wt1* in nephron progenitors and podocytes [32,33]. Other TFs that followed this pattern include *Foxc2* and *Foxl1. 3) A de novo increase in chromatin accessibility with cell type commitment and advancing differentiation.* For example, the chromatin accessibility of *Hnf4a* and *Pou3f3* motif increased in proximal tubule and loop of Henle trajectories, respectively, coinciding with the cellular differentiation program [34]. A large number of TFs followed this pattern such as *Mafb* (in podocytes), *Hnf4a* and *Hnf1a* (in proximal tubule), *Hnf1b* (in both proximal tubule and loop of Henle) as well as *Esrrb* and *Tfap2b* (in loop of Henle).

Next, we correlated changes in chromatin accessibility-based TF motif enrichment with TF expression and their respective target genes along Monocle-generated trajectories. To this end, we used the scRNA-seq differentiation trajectories to find TFs and target genes differentially expressed over pseudotime (**Supplemental Table 6**). We also noticed a good concordance of time-dependent changes of TF and target gene expression along with TF motif enrichment, including the lineages for podocytes (e.g., *Foxc2*, *Foxl1*, *Mafb*, *Magi2*, *Nphs1*, *Nphs2*, *Plat*, *Synpo*, *Thsd7a*, *Wt1*, and *Zbtb7c*), proximal tubule (e.g., *Ace2*, *Atp1a1*, *Dab2*, *Hnf1a*, Hnf4a, *Hsd17b2*, *Lrp2*, *Maf*, *Slc12a3*, *Slc22a12*, *Slc34a1*, and *Wnt9b*), loop of Henle (e.g.,*Cyfip2*, *Cytip*, *Esrrb*, *Esrrg*, *Irx1*, *Irx2*, *Mecom*, *Pla2g4a*, *Pou3f3*, *Ppargc1a*, *Stat3*, *Sytl2*, *Tfap2b*, *Thsd4*, and *Umod*), as well as for

both proximal tubule and loop of Henle (e.g., *Bhlhe40*, *Hnf1b*, and *Tmprss2*), respectively (**Figures**

3c, S3i). Most interestingly, we noticed two distinct patterns of how gene expression was related

to chromatin accessibility. While gene expression of TFs increased over pseudotime, its

corresponding motif accessibility either increased in parallel (such as *Hnf4a* and *Pou3f3*) or

maintained in a lineage-specific manner (such as *Wt1*). This might indicate different regulatory

mechanisms during differentiation.

We next aimed to interrogate the stage-dependent chromatin dynamics along the identified

differentiation trajectory. The differentiation trajectory was binned into 15 developmental steps

based on the lineage specification (**Figures S3b-c**). These stages were labeled as NP (nephron

progenitor), IM (intermediate cells), Podo (podocytes), PT (proximal tubule), LOH (loop of Henle),

and DCT (distal convoluted tubule), however, this designation will need to be matched with prior

cell marker-based annotations. To study the chromatin opening and closing, we conducted

differential chromatin accessibility analysis between subsequent stages. To understand the

biological processes controlled by the epigenetic changes, we examined the nearest genes and

performed functional annotation (**Methods**). We found that open chromatin profiles were

relatively stable in the early precursor stages such as NP1 to NP3, with fewer than 70 DAPs

identified (**Supplemental Table 7, Figure S4a**). The podocyte differentiation branch was

associated with marked increase in the number of DAPs, (796 DAPs between NP3 and Podo1).

This mainly represented the closing of chromatin areas around nephron progenitor-specific genes

such as *Osr1*, *Gdnf*, *Sall1*, *Pax2* and opening of areas around podocyte-specific genes and key TFs

such as *Foxc2* and *Efnb2*, both of which are validated to be important for early podocyte

differentiation [35,36]. At later stages, there was a strong increase in expression of actin filament-

based processes and a significant decrease in *Notch* and *Ctnnb1* in the podocyte lineages

(**Supplemental Table 8**). Fewer chromatin closing events were observed (234 DAPs) between

NP3 and intermediate cells 1 (IM1), mainly associated with closing of the chromatin around *Osr1*

and opening around tubule cell-specific TFs such as *Lhx1* and *Pax3* (**Figure 4**). The decrease in

Six2 expression only occurred at the IM2 stage, at which we also observed an increase in tubule

specification genes such as *Hnf1a*. Gene ontology results from the 820 up-regulated peaks between

PT1 and IM2 showed enrichment associated with typical proximal tubule functions including

sodium-dependent phosphate transport, maintenance of osmotic response in the loop of Henle and

active sodium transport in the distal convoluted tubule (**Figure S4a**, the full list can be found in

Supplemental Tables 7, 8 and 9).

In addition to analyzing changes along the trajectory, we also specifically examined cell-fate

decision events. We studied the chromatin opening and closing during the first cell commitment

event. We found that podocyte specification from nephron progenitors was associated with

differential opening of *Foxl1*, *Zbt7c,* and *Smad2* in the podocyte lineage and *Lhx1*, *Sall1*, *Dll1*,

Jag1, *Cxcr3* and *Pax3* in the other lineage, respectively. While the role of several TFs has been

established for podocyte specification, the expression of *Foxl1* has not been described in the kidney

until now (**Figure 4**). Our analysis pinpointed that four peaks in the vicinity of *Foxl1* were

accessible only in podocyte lineage, which locate in +53,381 bp, +152,832 bp, +237,019 bp, and

+268,550 bp of the *Foxl1* TSS, respectively. To confirm the expression of *Foxl1* in nephron

progenitors and podocytes, we performed immunofluorescence studies on developing kidneys

(E13.5, P0 and P6). Consistent with the computational analysis, we found strong expression of

FOXL1 in nephron progenitors (E13.5). At later stages, it was present in comma and S shape body

and finally in the glomerular podocytes (**Figure S4b**). Consistently, while there was no expression

within cells destined to become proximal tubule or loop of Henle cells, gene expression of *Foxl1*

increased in cells along the podocyte trajectory (**Figure S4c**). While further experimental

validation will be important, our study has illustrated the critical role of open chromatin state

information and dynamics in cellular differentiation.

The intermediate cells (IM) gave rise to proximal and distal branches, representing the proximal

tubules and the loop of Henle as well as distal convoluted tubule segments. The proximal tubule

region was characterized by chromatin opening around *Hnf4a, Maf*, *Tprkb*, and *Gpat2*. The loop

of Henle and distal convoluted tubule segments were remarkable for multiple DAPs in the vicinity

of *Tfap2a*, *Tfap2b*, *Cited4*, *Ephb2*, *Ephb3*, *Hoxd8*, *Mecom*, and *Prmd16*, indicating a critical novel

role for these TFs in distal tubule differentiation (**Figures 3c, 4, S4c**). Consistently, we saw a

reduction in chromatin accessibility of *Six2* promoter and enhancers along all three trajectories

(podocyte, proximal tubule and loop of Henle) (**Figure S4d**). There was also a decrease in

expression of *Jag1* and *Heyl* in the distal loop of Henle segment, consistent with the putative role

of Notch driving the proximal tubule fate [37] (**Supplemental Table 11)**. Another striking

observation was that tubule segmentation and specification occurred early by an increase in

chromatin accessibility around *Lhx1*, *Hnf1a* and *Hnf4a* and *Maf* for proximal tubule and *Tfap2b*

for loop of Henle. Terminal differentiation of proximal tubule and loop of Henle cells was strongly

linked to nuclear receptors that regulate metabolism, such as *Esrra* and *Ppara* in proximal tubules

and *Esrra* and *Ppargc1a* in the loop of Henle segment, once more indicating the critical role of

metabolism of driving gene expression and differentiation [38].

In summary, we reconstructed the developmental and differentiation trajectories of podocytes,

proximal tubule and loop of Henle cells. We defined chromatin and gene expression dynamics and

identified numerous putative TFs for kidney cell specification and differentiation.

Stromal-to-epithelial communication is critical in the developing and adult kidneys

Previous studies indicated that the survival, renewal, and differentiation of nephron progenitors is

largely regulated through its cross-talk with the adjacent ureteric bud [39]. To investigate the complex

cellular communication network, we used CellPhoneDB [40] to systematically infer potential cell-

cell communication in the developing and adult kidney. CellPhoneDB provides a comprehensive

database and a statistical method for the identification of ligand-receptor interactions in scRNA-

seq data. Analysis of our scRNA-seq dataset indicated that the number of cell-cell interaction pairs

was larger in developing kidney compared to the adult kidney (**Figure 5a**). In the developing

kidney, the stroma showed the greatest number of interactions among all cell types. This is

consistent with the well-known role of epithelial-stromal interactions in driving kidney

development. Of the identified interactions, many were related to stroma-secreted molecules such

as collagen 1, 3, 4, 6, and 14 (**Figure 5b**). Furthermore, the stroma seemed to interact with most

cell types, such as podocytes and different tubule cells. Interestingly, the nephron progenitor

cluster showed important ligand-receptor interaction between *Fgf1*, *Fgf8* as well as *Fgf9* and the

corresponding receptor *Fgfr1*, which is consistent with the well-known role of FGF signaling in

kidney development [41]. Of the manifold identified interactions in the fetal kidney, stromal

interaction and the VEGF-involving interaction remained significant in the adult data set,

underscoring the importance of endothelial-to-epithelial communication.

We next individually examined the expression of several key pathways known to play important
roles in kidney development, such as Gdnf-Ret, sonic hedgehog, FGF, Bmp, Wnt and others [9].
Expression of these key ligand-receptor pairs showed strong cell type specificity (**Figure 5c**). For
example, *Robo2* of the Gdnf-Ret pathway was expressed in nephron progenitors and in podocytes
of P0 and adult kidney. Gdnf signaling through the Ret receptor is required for normal growth of
the ureteric bud during kidney development [42] and the Slit2/Robo2 pathway is implicated with
congenital kidney anomalies (Hwang et al., Hum Genet 2015) and important for maintenance of
podocyte foot process integrity [43]. *Eya1*, however, is genetically upstream of *Gdnf* and acts as a
positive regulator for its activation [44]. Consistently, we noted distinct cell type specificity of *Eya1*
expression only in nephron progenitors, which was also true for other important signaling
molecules such as *Ptch1, Smo* and *Gli3* of the sonic hedgehog pathway. *Fgfr1* showed the highest
expression in nephron progenitors as well as in fetal and adult stroma, underscoring the importance
of FGF signaling for cell-cell interactions in both the developing and developed kidney. Most
interestingly, some cell-cell interactions between specific cell types that were observed in fetal
kidney were abrogated in adult kidney because of the loss of expression of either ligand or receptor,
such as *Pdgfc* in nephron progenitor signaling to its receptor *Pdgfra* in stroma, *Npnt* from several
epithelial cells signaling to *Itga8* in nephron progenitors, *Tnc-Itga9* signaling from nephron
progenitors to stroma and *Rspo3* in stroma signaling to *Sdc4* in several epithelial cells. Because
not much is known about some of these markers, the significance of these putative interactions
requires further investigation. For example, *Rspo3* has been implicated in nephron progenitor-
associated interactions during nephrogenesis [10]. Mutations in the *Itga8* gene are known to cause
isolated congenital anomalies of kidney and urinary tract in humans [45] and *Pdgfra* has been
regarded as a commitment marker in kidney differentiation [10].
446
447   In summary, we inferred cell-cell interactions in the developing and adult kidneys and found the
448   critical role of stromal-epithelial interactions in the developing kidney.
449

450 **Single cell chromatin accessibility identified human kidney GWAS target regulatory regions,**
451 **genes and cell types**

452

453 Finally, we examined whether single cell level chromatin accessibility data can help identify cell
454 and gene targets for human kidney disease development. GWAS have been exceedingly successful
455 in identifying nucleotide variations associated with specific diseases or traits. However, more than
456 90% of the identified genetic variants are in the non-coding region of the genome. Initial
457 epigenome annotation studies indicated that GWAS hits are enriched in tissue-specific enhancer
458 regions. As there are many different cell types in the kidney with differing function, understanding
459 the true cell type specificity of these enhancers is critically important. Here, we reasoned that
460 single cell accessible chromatin information could be extremely useful to identify the cell type-
461 specific enhancer regions and thereby the target cell type for the GWAS hits, however, such maps
462 have not been generated for the human kidney. We combined three recent kidney disease GWAS
463 [46-48], and obtained 26,637 single nucleotide polymorphisms (SNPs) that passed genome-wide
464 significance level of which we retained 7,923 after lift-over from human to mouse.

465

466 Specifically, we examined loci where functional validation studies reported conflicting results on
467 target cell types and target genes (**Figure 6).** The *SHROOM3* locus has shown a reproducible
468 association with kidney function in multiple GWAS [48]. However, previous functional follow-up
469 studies have reported confusing and somewhat contradictory results. While one study indicated
470 that the genetic variants were associated with an increase in SHROOM3 levels in tubule cells
471 inducing kidney fibrosis [49], the other suggested that the variant was associated with lower
472 SHROOM3 levels in podocytes resulting in chronic kidney disease development [50]. We found an
473 open chromatin (likely promoter) area in multiple cell types such as nephron progenitors,
474 podocytes, loop of Henle, distal convoluted tubule, principal cells and intercalated cells (**Figure**
475 **6a**). We also identified intronic open chromatin areas only in nephron progenitors and podocytes
476 that overlapped with the GWAS significant variants (**Figure 6a**). Consistent with the cis-
477 regulatory open chromatin, the strongest expression of *Shroom3* was observed in podocytes and
478 nephron progenitor cells. Expression of *Shroom3* in the adult bulk kidney was below our detection
479 limit. To further understand the regulatory dynamics of this locus in the developing mouse kidneys,
480 we examined gene expression and epigenome annotation data generated from bulk mouse kidney

samples at different stages of development for H3K27ac and H3K4me1 in adult and fetal samples

(**Figure 6b**). Interesting to note that the GWAS-significant SNP that showed strong nephron

progenitor-specific enrichment also coincided with the *Six2* binding area. Finally, the Cicero-based

co-accessible analysis connected the GWAS top variants, which located in an intronic enhancer

region of *Shroom3*, with *Shroom3* exons, indicating that *Shroom3* is the likely target gene of the

variant (**Figure 6a**).

Next, we analyzed the chromosome 15 GWAS region, where we identified some open chromatin

regions that were uniformly open in all examined cell types. *Dab2* expression, on the other hand,

strongly correlated with open distal enhancer regions in proximal tubule cells (**Figure 6c**). This is

consistent with earlier publications indicating the role of proximal tubule-specific DAB2 playing

a role in kidney disease development [51]. Interestingly, while single cell analysis indicated an

additional distal enhancer in intercalated cells, the GWAS-significant region coincided with the

proximal tubule-specific enhancer region and showed strong coregulation (**Figure 6d**). Regulatory

annotation of the developing kidney indicated strong enhancer marks in the adult but not in the

fetal kidney.

Lastly, we examined the region around *Uncx*, for which reproducible association with kidney

function was shown in multiple GWAS [46,47]. Interestingly, the GWAS locus demonstrated a strong

open chromatin region in nephron progenitors but not in any other differentiated cell types (**Figure

6e**). Consistently, in bulk chromatin accessibility data we only observed regulatory activity such

as H3K27ac, H3Kme1 and we show *Six2*-binding at this locus in fetal kidneys. The locus did not

show H3K27ac enrichment in the adult kidney, while H3K4me1 remained positive (**Figure 6f**).

Uncx expression was strong in the fetal kidney samples, but we could not detect its expression in

the adult kidney (**Figure 6e**). A closer view of these loci is shown in **Figure S6**.

These results indicate that variants associated with kidney disease development are located in

regions with cell type- and developmental stage-specific regulatory activity and illustrate the

critical role of snATAC-seq in defining target genes and target cell types for GWAS variants.

Discussion

In summary, here we present the first cellular resolution open chromatin map for the developing and adult mouse kidney. Using this dataset, we identified key cell type-specific regulatory networks for kidney cells, defined the cellular differentiation trajectory, characterized regulatory dynamics and identified key driving TFs for nephron development, especially for the terminal differentiation of epithelial cells. Furthermore, our results shed light on the cell types and target genes for genetic variants associated with kidney disease development.

By performing massively parallel single cell profiling of chromatin state, we were able to define the key regulatory logic for each kidney cell type by investigating cis-regulatory elements and TF-target gene interaction. We found that most cell type-specific open chromatin regions are within distal regulatory elements and intronic regions. Our studies identified a massive amount of highly dynamic co-regulated peaks indicating the important correlation between distal regulatory elements and gene expression. Future studies will examine the relative contribution of promoters and enhancer openness in gene expression regulation. However, these studies highlight that both chromatin opening and looping are critical for gene regulation.

We also observed that the single cell open chromatin atlas was able to define more distinct cell types even in the developing kidney compared to scRNA-seq analysis. Given the continuous nature of RNA expression, it has been exceedingly difficult to dissect specific cell types in the developing kidney [9,10,13]. In addition, it has been difficult to resolve the cell type origin of lowly expressed transcripts in scRNA-seq data. However, this is not the case for snATAC-seq data, which were able to capture the chromatin state irrespective of gene expression magnitude. There were several examples where accessible peaks were identified in specific cell types even for lowly expressed genes such as Shroom3.

We identified critical cell type-specific TFs by integrating multiple computational analyses. TF identification is challenging in scRNA-seq data since the expression of several cell type-specific TFs is low and some of them do not show a high degree of cell type-specificity. By extracting motif information, snATAC-seq data provides additional information for TF identification.

Together with regulon analysis, as implemented in SCENIC, we have identified several TFs as well as their target genes that are important for kidney development. Leveraging this newly identified cell type-specific regulatory network will be essential for future studies of cellular reprogramming of precursors into specific kidney cell types and for better understanding homeostatic and maladaptive regeneration.

Our studies revealed dynamic chromatin accessibility that tracks with renal cell differentiation. These states may reveal mechanisms governing the establishment of cell fate during development, in particular those underlying the emergence of specific cell types. We found a consistent and coherent pattern between gene expression and open chromatin information, where the nephron progenitors differentiated into two branches representing podocytes and tubule cells [52]. We found that podocytes commitment occurred earlier, while tubule differentiation and segmentation appeared to be more complex. This podocyte specification correlated with the maintenance of expression of *Foxc2* and *Foxl1* expression in podocytes. While *Foxc2* has been known to play a role in nephron progenitors and podocytes, this is the first description of *Foxl1* in kidney and podocyte development. Our studies are consistent with recent observations from organoid models that recapitulated podocyte differentiation better than tubule cell differentiation [53]. Our study also sheds light on tubule differentiation and segmentation. We confirmed the key role of *Hnf4a* in proximal tubules. We have identified a large number of new transcriptional regulators such as *Tfap2a* that seem to be critical for the distal portion of the nephron. Our data indicate that distal tubule differentiation is linked to the loop of Henle, a critically important observation needing further confirmation. Furthermore, the terminal differentiation of proximal tubule cells correlated with the increase in *Ppara* and *Esrra* expression, both of which are known regulators of oxidative phosphorylation and fatty acid oxidation [38]. Loop of Henle differentiation strongly correlated with *Essrb* and *Ppargc1a* expression. These studies potentially indicate that cell specification events occur early and metabolism controls terminal differentiation of tubule cells [54]. Impaired metabolic fitness of proximal tubules has been a key contributor to kidney dysfunction, explaining the critical association with tubule metabolism and function.

Furthermore, we show that single cell and stage level epigenome annotation is critical for the annotation of human GWAS. Most identified GWAS signals are in the non-coding region of the

genome. Due to the linkage disequilibrium structure of the human genome, each GWAS locus

contains a large number of variants, each passing genome-wide significance level [55]. Furthermore,

as these signals are often non-coding, the target gene and the target cell type remain unknown.

While molecular quantitative trait locus studies and bulk epigenome annotation experiments have

been important to define the molecular pathways leading to disease development from the

identified signals, these methods have limited resolution, as cell type-specific enhancer regions

cannot be identified by bulk analysis [56]. Additionally, bulk molecular quantitative trait locus

studies suffer from the same linkage disequilibrium problems as GWAS analyses [57]. Our results

indicate that multiple GWAS regions are conserved between mice and humans. Single cell open

chromatin information enables not only the identification of affected cell types, but also the

understanding of co-regulation of the open chromatin area. It is also able to highlight critical target

genes. Performing single cell open chromatin analysis on human kidney tissue samples will be

essential to further understand molecular pathways altered by genetic variants. Here we showed

three important examples. We confirmed the role of *Dab2* and its specific expression in the

proximal tubule during kidney disease development, as its implication therein has been shown in

previous expression quantitative trait locus and bulk epigenome analysis experiments [51].

Furthermore, we showed that the GWAS variants map only to those regions where chromatin is

open exclusively in nephron progenitors, whereas chromatin becomes inaccessible as

differentiation progresses during later stages, such as *Shroom3* and *Uncx*. This is an interesting

and important novel mechanism, indicating that the altered expression of this gene might play a

role in the development rewiring of the kidney. This mechanism is similar to genes associated with

autism that are known to be expressed in the fetal but not in the adult stages [58] and highlights the

critical role of understanding chromatin accessibility at multiple stages of differentiation.

While we have generated a large amount of high-quality data, this information will need further

experimental validation, which is beyond the scope of the current manuscript. In addition, one

needs to be aware of the limitations when interpreting different computational analyses, for

example, the motif enrichment analyses such as implemented by HOMER, SCENIC, and

chromVAR, are not able to distinguish between TFs with similar binding sites. Future high-

throughput studies that analyze open chromatin and gene expression information from the same

cells will be exceedingly helpful to correlate open chromatin and gene expression information

along the differentiation trajectory [24,59,60].

In summary, our dataset provides critical novel insight into the cell type-specific gene regulatory

network, cell differentiation program, and disease development.

Materials and Methods

Single cell RNA sequencing of P0 mice

1-day-old mouse neonate was decapitated with surgical scissors, 2 kidneys were harvested and minced into 1 mm^3 pieces and incubated with digestion solution containing Enzyme D, Enzyme R and Enzyme A from Multi Tissue Dissociation Kit (Miltenyi, 130-110-201) at 37 °C for 15 min with agitation. Reaction was deactivated by adding 10% FBS, then solution was passed through a 40 μm cell strainer. After centrifugation at 1,000 RPM for 5 min, cell pellet was incubated with 500 μL of RBC lysis buffer on ice for 3 min. We centrifuged the cells at 1,000 RPM for 5 min at 4 °C and resuspended the cells in the buffer for further steps. Cell number and viability were analyzed using Countess AutoCounter (Invitrogen, C10227). The cell concentration was 2.2 million cells/mL with 92% viability. 10,000 cells were loaded into the Chromium Controller (10X Genomics, PN-120223) on a Chromium Single Cell B Chip (10X Genomics, PN-120262) and processed to generate single cell gel beads in the emulsion (GEM) according to the manufacturer's protocol (10X Genomics, CG000183). The library was generated using the Chromium Single Cell 3' Reagent Kits v3 (10X Genomics, PN-1000092) and Chromium i7 Multiplex Kit (10X Genomics, PN-120262) according to the manufacturer's manual. Quality control for constructed library was performed by Agilent Bioanalyzer High Sensitivity DNA kit (Agilent Technologies, 5067-4626) for qualitative analysis. Quantification analysis was performed by Illumina Library Quantification Kit (KAPA Biosystems, KK4824). The library was sequenced on an Illumina HiSeq or NextSeq 2x150 paired-end kits using the following read length: 28 bp Read1 for cell barcode and UMI, 8 bp I7 index for sample index and 91 bp Read2 for transcript.

Single cell ATAC sequencing

3-week-old and 8-week-old mice were euthanized and perfused with chilled 1x PBS via left ventricle. Kidneys (0.25 g) were harvested, minced and lysed in 5 mL lysis buffer for 15 min. 1-day-old mice were decapitated with surgical scissors, and both kidneys were harvested. Kidneys were minced and lysed in 2 mL lysis buffer for 15 min. Tissue lysis reaction was then blocked by adding 10 mL 1x PBS into each tube, and solution was passed through a 40 μm cell strainer. Cell debris and cytoplasmic contaminants were removed by Nuclei PURE Prep Nuclei Isolation Kit (Sigma, NUC-201) after centrifugation at 13,000 RPM for 45 min. Nuclei concentration was

calculated with Countess AutoCounter (Invitrogen, C10227). Diluted nuclei suspension was

loaded and incubated in transposition mix from Chromium Single Cell ATAC Library & Gel Bead

Kit (10X Genomics, PN-1000110) by targeting 10,000 nuclei recovery. GEMs were then captured

on the Chromium Chip E (10x Genomics, PN-1000082) in the Chromium Controller according to

the manufacturer's protocol (10X Genomics, CG000168). Libraries were generated using the

Chromium Single Cell ATAC Library & Gel Bead Kit and Chromium i7 Multiplex Kit N (10X

Genomics, PN-1000084) according to the manufacturer's manual. Quality control for constructed

library was perform by Agilent Bioanalyzer High Sensitivity DNA kit. The library was sequenced

on an Illumina HiSeq 2x50 paired-end kits using the following read length: 50 bp Read1 for DNA

fragments, 8 bp i7 index for sample index, 16 bp i5 index for cell barcodes and 50 bp Read2 for

DNA fragments.

Bulk ATAC sequencing

Bulk ATAC-seq was performed as described earlier [61,62]. Briefly, 50,000 nuclei/sample were

tagmented with Tn5 transposase (Illumina) in 50 μl reaction volume including Tween-20 (0.1%)

(Sigma) and digitonin (0.01%) (Promega). The reaction was carried out at 37 °C for 30 min in a

thermomixer at 1,000 RPM. After purification of DNA with Qiagen Minelute Reaction Cleanup

kit (Qiagen), samples were subjected to library amplification (8-10 cycles). Libraries were purified

with AmpureXP beads (Beckman Coulter) and their quality was assessed by Agilent High

sensitivity DNA Bioanalysis chip (Agilent). Libraries were submitted to 150 bp PE sequencing.

snATAC-seq data analysis

Data processing and quality control

Raw fastq files were aligned to the mm10 (GRCm38) reference genome and quantified using Cell

Ranger ATAC (v. 1.1.0). We only kept valid barcodes with number of fragments ranging from

1,000 to 40,000 and mitochondria ratio less than 10%. One of the important indicators for ATAC-

seq data quality is the fraction of peaks in promoter regions, so we did further filtration based on

promoter ratio. We noticed the promoter ratio seemed to follow a binary distribution, with most of

cells either having a promoter ratio around 5% (background) or more than 20% (valid cells)

(**Figure S1d**). We therefore filtered out cells with a promoter ratio <20%. After this stringent

quality control, we obtained 11,429 P0 single cells (5,993 in P0_batch_1 and 5,436 in P0_batch_2)

and 16,887 adult single cells (7,129 in P56_batch_3, 6,397 in P56_batch_4, and 3,361 in
P21_batch_5).

Preprocessing

Since snATAC-seq data are very sparse, previous methods either conducted peak calling or
binarization before clustering. Here, we chose to do binarization instead of peak calling for two
reasons: 1) Peak calling is time consuming; 2) Many peaks are cell type-specific, open chromatin
regions in rare populations are more likely to be treated as background. After binarizing fragments
into 5 kb bins and removing the fragments not matched to chromosomes or aligned to the
mitochondria, we binarized the cell-bin matrix. In order to only keep bins that were informative
for clustering, we removed the top 5% most accessible bins and bins overlapping with ENCODE
blacklist. The 484,606 remaining bins were used as input for clustering.

Dimension reduction, batch effect correction and clustering

Clustering was conducted using snapATAC [16], a single-cell ATAC-seq algorithm scalable to large
dataset. Previous benchmarking evaluation has shown that snapATAC was one of the best-
performing methods for snATAC-seq clustering [63]. Diffusion map was applied as a dimension
reduction method using function *runDiffusionMaps*. To remove batch effect, we used Harmony [17],
in which the low dimensional embeddings obtained from the diffusion map were used as input.
Harmony iteratively pulled batch-specific centroid to cluster centroid until convergence to remove
the variability across batches. After batch correction, a graph was constructed using k Nearest
Neighbor (kNN) algorithm with k=15, which was then used as input for Louvain clustering. We
used the first 20 dimensions for the Louvain algorithm. The number of dimensions was chosen
using a method recommended by snapATAC, although we noticed that the clustering results were
similar among a series of dimensions from 18 to 30.

Cell type annotation

We used a published list of marker genes [9,19] to annotate kidney cell types. In order to infer gene
expression of each cell type, we built a cell-gene activity score matrix by integrating all fragments
that overlapped with gene transcript. We used GENCODE Mouse release VM16 [22] as reference
annotation.

Peak calling and visualization

Peak calling was conducted for each cell type separately using MACS2 [18]. We aggregated all fragments obtained from the same cell types to build a pseudo-bulk ATAC data and conducted peak calling with parameters "--nomodel --keep-dup all --shift 100 --ext 200 --qval 1e-2 -B --SPMR --call-summits". By specifying "--SPMR", MACS2 generated "fragment pileup per million reads" pileup files, which were converted to bigwig format for visualization using UCSC bedGraphToBigWig tool.

We also visualized public chromatin ChIP-seq data and RNA-seq data obtained from ENCODE Encyclopedia (https://www.encodeproject.org/) with the following identifiers: ENCFF338WZP, ENCFF872MVE, ENCFF455HPY, ENCFF049LRQ, ENCFF179NTO, ENCFF071PID, ENCFF746MFH, ENCFF563LOO, ENCFF184AYF, ENCFF107NQP, ENCFF465THI, ENCFF769XWI, ENCFF591DAX. The Six2 ChIP-seq data were obtained from [64] and the WGBS data were obtained from [65].

Genomic elements stratification

Mouse mm10 genome annotation files were download from UCSC Table Browser (https://genome.ucsc.edu/cgi-bin/hgTables) using GENCODE VM23. TSS upstream 5 kb regions were included as promoter regions, but the results were similar when using 2 kb upstream regions as promoters. We then studied the number of overlapped regions between open chromatin regions identified from the snATAC-seq and bulk ATAC-seq dataset and genome annotations. Since one open chromatin region could overlap with multiple genomic elements, we defined an order of genomic elements as exon > 5'-UTR > 3'-UTR > intron > promoter > distal elements. To be more specific, if one peak overlapped with both exon and 5'-UTR, the algorithm would count it as an exon-region peak.

Identification of differentially accessible regions

Peaks identified in each cell type were combined to build a union peak set. Overlapping peaks were then merged to one peak using *reduce* function from the GenomicRanges package. This resulted in 300,755 peaks, which were used to build binarized cell-by-peak matrix. Differentially

accessible peaks (DAPs) for each cell type were identified by pairwise peak comparison.

Specifically, for each peak, we conducted a Fisher's exact test between a cell type and each of the

other cell types. To address multiple testing problem, we used the Benjamini-Hochberg approach

(BH correction) to correct p values. Peaks with corrected p values below significance level (0.05)

in all pairwise tests were defined as DAPs. In total, we obtained 60,683 DAPs, which were used

for motif enrichment analysis.

Motif enrichment analysis

Motif enrichment analysis was conducted using DAPs by HOMER v4.10.4 [27] with parameters

background="automatic" and scan.size=300. We noticed that *de novo* motif identification only

generated few significant results, so we focused on known motifs for our following study. We used

the significance level of 0.05 for BH corrected p value to determine the enriched results. The motif

enrichment results are provided in **Supplemental Table 3**.

Peak-peak correlation analysis

Peak-peak correlation analysis was conducted using Cicero [26]. In order to find developmental

stage-specific peak-peak correlations, the analysis was conducted for P0 and adult separately.

Cicero uses Graphic Lasso with distance penalty to assess the co-accessibility between different

peaks. Cicero analysis was conducted using the *run_cicero* function with default parameters. A

heuristic cutoff of 0.25 score of co-accessibility was used to determine the connections between

two peaks.

snATAC-seq trajectory analysis

snATAC-seq trajectory was conducted using Cicero, which extended Monocle3 to the snATAC-

seq analysis. We obtained the preprocessed P0 snATAC-seq cell-peak matrix from snapATAC as

input for Cicero and conducted dimension reduction using Latent Semantic Indexing (LSI) and

visualized using UMAP. Trajectory graph was built using the function *learn_graph*. Batch effect

was not observed between the two P0 batches, and the trajectory graph was consistent with cell

type assignment with clustering analysis (**Figures S3 a-b**).

In order to study how open chromatin changes are associated with the cell fate decision, we first
binned the cells into 15 groups based on their pseudotime and cell type assignment. Next, we
studied the DAPs between each group and its ancestral group using the same methods described
above. The number of newly open and closed chromatins were reported using pie charts. The exact
peak locations are provided in the **Supplemental Table 7**.

Genes and gene ontology terms associated with snATAC-seq trajectory
Based on the binned trajectory graphs and DAPs between each group and its ancestral group, we
next used GREAT tool [66] to study the enrichment of associated genes and gene ontology (GO)
terms along the trajectory. We used the newly open or closed peaks as test regions and all the peaks
from peak-calling output as the background regions for the analysis. The output can be found in
the **Supplemental Table 8 and 9**.

Predict cis-regulatory elements
We implemented two methods to study cis-regulatory elements in the snATAC-seq data. The first
method was inspired by [23], which was based on the observation that there was co-enrichment in
the genome between the snATAC-seq cell type-specific peaks and scRNA-seq cell type-specific
genes. This method links a gene with a peak if 1) they were both specific in the same cell type, 2)
they were in *cis*, meaning that the peak is in ±100 kb region of the TSS of the corresponding gene,
and 3) the peak did not directly overlap with the TSS of the gene. This method successfully inferred
several known distal elements such as for *Six2* and *Slc6a18* (**Figure S2g**).

Alternatively, we assessed the co-accessibility of two peaks. We implemented Cicero [26], which
aggregates similar cells to obtain a set of "meta-cells" and address the issue with sparsity in the
snATAC-seq data. We used *run_cicero* function with default parameters to predict cis-regulatory
elements (CREs). Although it is recommended to use 0.25 as a cutoff for co-accessibility score,
we noticed that this resulted in a great amount of CREs, which could contain many false positives.
Thus, we used a more stringent score of 0.4 for the cutoff and retained 1,214,638 and 926,288
CRE links in the P0 and adult data, respectively.

Bulk ATAC sequencing analysis

Bulk ATAC-seq raw fastq files were processed using the end-to-end tool ENCODE ATAC-seq pipeline (**Software and Algorithms**). This tool provided a standard workflow for ATAC-seq data quality control, adaptor removal, alignment, and peak calling. To obtain high quality ATAC-seq peaks, peak calling results from two biological replicates were compared and only those peaks that were present in both replicates were kept, which were further used to compare with snATAC-seq peaks.

Correlation of bulk and single nuclei ATAC sequencing data

snATAC-seq reads were aggregated to a pseudo-bulk data for the comparison purpose. To prevent the effect of sex chromosome and mitochondria chromosome, reads from chromosome X, Y and M were excluded from our analysis. We used multiBigwigSummary tool from deeptools [67] to study the correlation between different samples. Specifically, the whole genome was binned into equally sized (10 kb) windows, and the reads in each bin were aggregated, generating a bin-read count vector for each of the sample. The correlation of these vectors was computed as a measure of pairwise similarity between samples.

To compare the number of peaks in these two datasets, we used as input the narrowpeak files from the snATAC-seq and bulk ATAC-seq analysis. We filtered out bulk ATAC-seq peaks with q value > 0.01 to be consistent with the snATAC-seq setting. Since the snATAC peaks were called after merging different time points, we also took the union set of bulk ATAC-seq peaks from different time points. We then used *findoverlap* function in GenomicRanges package [68] to find and report overlapped peaks.

Comparison between single nuclei ATAC sequencing data and single cell RNA sequencing data

In order to compare the cluster assignment between snATAC-seq data and scRNA-seq data, we obtained the average gene expression values and peak accessibility in each cluster for P0 and adult samples separately. We next transformed snATAC-seq data by summing up the reads within gene body and 2 kb upstream regions to build gene activity score matrix, as suggested in Seurat [20]. Then, we normalized the data and computed the mean expression and mean gene activity scores in each

cell type, and calculated z scores of each gene. Pearson's correlation coefficient was then

calculated among top 3,000 highly variable genes between snATAC-seq data and scRNA-seq data.

We found high concordance between these two datasets in terms of cell type assignment (**Figure**

S4).

Single cell RNA sequencing data analysis

Alignment and quality control

Raw fastq files were aligned to the mm10 (Ensembl GRCm38.93) reference genome and

quantified using CellRanger v3.1.0 (http://10xgenomics.com). Seurat R package v3.0 [69] was used

for data quality control, preprocessing and dimensional reduction analysis. After gene-cell data

matrix generation of both P0 and adult datasets, matrices were merged and poor-quality cells with

<200 or >3,000 expressed genes and mitochondrial gene percentages >50 were excluded, leaving

25,138 P0 and 18,498 adult cells for further analytical processing, respectively (**Figures S1j-k**).

Pre-processing, batch effect correction and dimension reduction

Data were normalized by RPM following log transformation and 3,000 highly variable genes were

selected for scaling and principal component analysis (PCA). Harmony R package v1.0 [17] was

used to correct batch effects. The top 20 dimensions of Harmony embeddings were used for

downstream uniform manifold approximation and projection (UMAP) visualization and clustering

(**Figures S1l-m**).

Cell clustering, identification of marker genes and differentially expressed genes

Louvain algorithm with resolution 0.4 was used to cluster cells, which resulted in 18 distinct cell

clusters. A gene was considered to be differentially expressed if it was detected in at least 25% of

one group and with at least 0.25 log fold change between two groups and the significant level of

BH-adjusted p value <0.05 in Wilcoxon rank sum test was used. We used a list of marker genes

[9,19] to manually annotate cell types. 2 distal convoluted tubule clusters were merged based on the

marker gene expression, resulting in a total of 17 clusters (**Figures S1i, n, o**).

scRNA-seq trajectory analysis

<u>Monocle3</u>

To construct single cell pseudotime trajectory and to identify genes whose expression changed as

the cells underwent transition, Monocle3 v0.1.3 [70] was applied to P0 cells of the following Seurat

cell clusters: nephron progenitors (NP), proliferating cells, stroma-like cells, podocytes, loop of

Henle (LOH), early proximal tubule (PT), proximal tubule S1, proximal tubule S3 cells.

To show cell trajectories from both small (nephron progenitors) and large cell populations

(proximal tubule), an equal number of 450 cells per cluster was randomly subsampled. Cells were

re-clustered by Monocle3 using a resolution of 0.0005 with k-nearest neighbor (kNN) k=29.

Highly variable genes along pseudotime were identified using differential *GeneTest* function and

cells were ordered along pseudotime trajectory. NP cluster was defined as earliest principal node.

In order to find genes differentially expressed along pseudotime, trajectories for podocytes, loop

of Henle, and proximal tubule clusters were analyzed separately with the *fit_models* function of

Monocle3. Genes with a q value <0.05 in the differential *GeneTest* analysis were kept. In an

alternate approach, *graph_test* function of Monocle3 was used and trajectory-variable genes were

collected into modules at a resolution of 0.01.

<u>RNA velocity</u>

To calculate RNA velocity, Python-based Velocyto command-line tool as well as Velocyto.R

package were used as instructed [30]. We used Velocyto to calculate the single-cell

trajectory/directionality using spliced and unspliced reads. From loom files produced by the

command-line tool, we subset the exact same cells that were previously selected randomly for

Monocle trajectory analysis. This subset was loaded into R using the SeuratWrappers v0.1.0

package. RNA velocity was estimated using gene-relative model with k-nearest neighbor (kNN)

cell pooling (k = 25). The parameter n was set at 200, when visualizing RNA velocity on the

UMAP embedding.

Gene regulatory network inference

In order to identify TFs and characterize cell states, we employed *cis*-regulatory analysis using the

R package SCENIC v1.1.2.2 [71], which infers the gene regulatory network based on co-expression

and DNA motif analysis. The network activity is then analyzed in each cell to identify recurrent

cellular states. In short, TFs were identified using GENIE3 and compiled into modules (regulons),

which were subsequently subjected to *cis*-regulatory motif analysis using RcisTarget with two

gene-motif rankings: 10 kb around the TSS and 500 bp upstream. Regulon activity in every cell

was then scored using AUCell. Finally, binarized regulon activity was projected onto Monocle3-

created UMAP trajectories.

Ligand-receptor interactions

To assess cellular crosstalk between different cell types, we used the CellPhoneDB repository to

infer cell-cell communication networks from single cell transcriptome data [40]. We used the Python

package CellPhoneDB v2.1.2 together with the database v2.0.0 to predict cell type-specific ligand-

receptor complexes as per the authors' instructions. Only receptors and ligands expressed in more

than 5% of the cells in the specific cluster were considered. 1,000 iterations were used for pairwise

comparison between cell types and considered for further statistical analysis.

Immunofluorescence staining

Mouse kidneys were fixed with 4% paraformaldehyde overnight, rinsed in PBS, and dehydrated

for paraffin embedding. Antigen retrieval was performed using Tris-EDTA buffer pH 9.0 with a

pressure cooker (PickCell Laboratories, Agoura Hills, CA) and antibody staining performed as

described [72]. Antibodies used were as follows: guinea pig FOXL1 (1:1,500) [73], mouse E-cadherin

(1:250; BD Transducton 610182, Franklin Lakes, NJ). Cy2-, Cy3-, and Cy5-conjugated donkey

secondary antibodies were purchased from Jackson ImmunoResearch Laboratories, Inc.

Fluoresecence images were collected on a Keyence microscope.

Material Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, Antibodies, Peptides, and Recombinant Proteins		
Guinea pig anti-Foxl1	own production	(ref. 73)
Mouse anti-E-cadherin	BD Transducton	Cat#610182

Cy2-conjugated donkey secondary antibody	Jackson ImmunoResearch Laboratories	Cat#715-225-150
Cy3-conjugated donkey secondary antibody	Jackson ImmunoResearch Laboratories	Cat#715-165-150
Cy5-conjugated donkey secondary antibody	Jackson ImmunoResearch Laboratories	Cat#715-175-150
DPBS	Corning	Cat# 21-031-CV
Tet System Approved FBS	Clontech	Cat# 631106
Nonidet™ P40 Substitute	Sigma	Cat# 74385
Magnesium Chloride Solution	Sigma	Cat# M1028
Ultrapure BSA (50 mg/ml)	Thermo Fisher	Cat# AM2616
RNAse inhibitor	Applied Biosystems	Cat# 100021540
Critical Commercial Assays		
Bioanalyzer High Sensitivity DNA kit	Agilent Technologies	5067-4626
Chromium Cell B Chip	10X Genomics	PN-120262
Chromium Chip E	10X Genomics	PN-1000082
Chromium Controller	10X Genomics	PN-120223
Chromium i7 Multiplex Kit	10X Genomics	PN-120262
Chromium Single Cell 3' Reagent Kits v3	10X Genomics	PN-1000092
Chromium Single Cell ATAC Library & Gel Bead Kit	10X Genomics	PN-1000110
Chromium Single Cell ATAC Library & Gel Bead Kit and Chromium i7 Multiplex Kit N	10X Genomics	PN-1000084
Countess AutoCounter	Invitrogen	C10227
Illumina Library Quantification Kit	KAPA Biosystems	KK4824
Multi Tissue dissociation kit	Miltenyi	130-110-201
Nuclei PURE Prep Nuclei Isolation Kit	Sigma	NUC-201
Deposited Data		
sci-CAR seq data	(ref. 24)	GSE117089
Mouse adult kidney WGBS data	(ref. 65)	GSM1051156
Mouse kidney H3K27ac and H3K4me1 CHIP-seq, WGBS, and RNA-seq	ENCODE project	ENCFF338WZP, ENCFF872MVE, ENCFF455HPY, ENCFF049LRQ, ENCFF179NTO, ENCFF071PID, ENCFF746MFH, ENCFF563LOO, ENCFF184AYF, ENCFF107NQP,

		ENCFF465THI, ENCFF769XWI, ENCFF591DAX
Six2 ChIP-seq data in nephron progenitor cells	(ref. 64)	GUDMAP database (RID:Q-Y4CY)
Software and Algorithms		
bedtools v. 2.29.2	open source	https://bedtools.readthedocs.io/en/latest/
Cell Ranger ATAC v. 1.1.0	10X Genomics	https://support.10xgenomics.com/single-cell-atac/software/downloads/latest
Cell Ranger v. 3.1.0	10X Genomics	https://support.10xgenomics.com/single-cell-gene-expression/software/downloads/latest
CellPhoneDB v. 2.1.2	open source	https://www.cellphonedb.org
ChromVAR v. 3.1.0	open source	http://bioconductor.org/packages/release/bioc/html/chromVAR.html
Cicero v. 1.5.5	open source	https://github.com/cole-trapnell-lab/cicero-release
deeptools v. 2.0	open source	https://deeptools.readthedocs.io/en/develop/
ENCODE ATAC-seq pipeline	open source	https://github.com/kundajelab/atac_dnase_pipelines
HOMER v. 4.10.4	open source	http://homer.ucsd.edu/homer/motif/
IGV	open source	http://software.broadinstitute.org/software/igv/
MACS2 v. 2.2.6	open source	https://github.com/taoliu/MACS
Monocle3 v. 0.1.3	open source	http://cole-trapnell-lab.github.io/monocle-release/
Harmony	open source	https://github.com/immunogenomics/harmony
SCENIC v. 1.1.2.2	open source	https://aertslab.org/#scenic
Seurat R package v. 3.0	open source	https://satijalab.org/seurat/
snapATAC	open source	https://github.com/r3fang/SnapATAC
UCSC bedgraphtobigwig	open source	http://hgdownload.soe.ucsc.edu/admin/exe/linux.x86_64.v385/
UCSC liftOver	open source	https://genome.ucsc.edu/cgi-bin/hgLiftOver
VelocytoR	open source	https://github.com/velocyto-team/velocyto.R
VisCello	open source	https://github.com/qinzhu/VisCello

Acknowledgements

Work in the Susztak lab is supported by the NIH DK076077, DK087635, and DK105821. MSB is supported by a German Research Foundation grant (BA 6205/2-1).

Author Contributions

KS and ZM designed and conceived the experiment. ZYM, JW, RS, and TA conducted the experiment. ZM conducted snATAC-seq bioinformatics analysis with advice from KS, HL, ML, and JK. MSB and ZM conducted scRNA-seq bioinformatics analysis with advice from KS. AMK and AYK conducted immunofluorescence staining with supervision from KHK. KS, ZM, and MSB wrote the manuscript and all authors edited and approved of the final manuscript.

Conflict of Interest

Authors declare no competing interests.

Lead Contact and Materials Availability

Raw data files and data matrix are being uploaded onto GEO and an accession number will be provided when it becomes available. The annotated and analyzed data can be viewed at **http://susztaklab.com/VisCello_snATAC/** and **http://susztaklab.com/igv/**. Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact: Katalin Susztak. Email: ksusztak@pennmedicine.upenn.edu.

References

Reidy, K., Kang, H. M., Hostetter, T. & Susztak, K. Molecular mechanisms of diabetic kidney disease. *J Clin Invest* **124**, 2333-2340, doi:10.1172/JCI72271 (2014).

Costantini, F. & Kopan, R. Patterning a complex organ: branching morphogenesis and nephron segmentation in kidney development. *Dev Cell* **18**, 698-712, doi:10.1016/j.devcel.2010.04.008 (2010).

Park, J. S. *et al.* Six2 and Wnt regulate self-renewal and commitment of nephron progenitors through shared gene regulatory networks. *Dev Cell* **23**, 637-651, doi:10.1016/j.devcel.2012.07.008 (2012).

Harding, S. D. *et al.* The GUDMAP database--an online resource for genitourinary research. *Development (Cambridge, England)* **138**, 2845-2853, doi:10.1242/dev.063594 (2011).

Wu, H. *et al.* Comparative Analysis and Refinement of Human PSC-Derived Kidney Organoid Differentiation with Single-Cell Transcriptomics. *Cell Stem Cell* **23**, 869-881 e868, doi:10.1016/j.stem.2018.10.010 (2018).

Takasato, M. *et al.* Kidney organoids from human iPS cells contain multiple lineages and model human nephrogenesis. *Nature* **526**, 564-568, doi:10.1038/nature15695 (2015).

Morizane, R. & Bonventre, J. V. Generation of nephron progenitor cells and kidney organoids from human pluripotent stem cells. *Nat Protoc* **12**, 195-207, doi:10.1038/nprot.2016.170 (2017).

Nishinakamura, R. Human kidney organoids: progress and remaining challenges. *Nature Reviews Nephrology* **15**, 613-624, doi:10.1038/s41581-019-0176-x (2019).

Combes, A. N. *et al.* Correction: Single cell analysis of the developing mouse kidney provides deeper insight into marker gene expression and ligand-receptor crosstalk (doi:10.1242/dev.178673). *Development* **146**, doi:10.1242/dev.182162 (2019).

Adam, M., Potter, A. S. & Potter, S. S. Psychrophilic proteases dramatically reduce single-cell RNA-seq artifacts: a molecular atlas of kidney development. *Development* **144**, 3625-3632, doi:10.1242/dev.151142 (2017).

Schmidt-Ott, K. M. How to grow a kidney: patient-specific kidney organoids come of age. *Nephrol Dial Transplant* **32**, 17-23, doi:10.1093/ndt/gfw256 (2017).

Menon, R. *et al.* Single-cell analysis of progenitor cell dynamics and lineage specification in the human fetal kidney. *Development (Cambridge, England)* **145**, doi:10.1242/dev.164038 (2018).

13 Lindstrom, N. O. *et al.* Progressive Recruitment of Mesenchymal Progenitors Reveals a
Time-Dependent Process of Cell Fate Acquisition in Mouse and Human Nephrogenesis.
Dev Cell **45**, 651-660 e654, doi:10.1016/j.devcel.2018.05.010 (2018).

14 Hochane, M. *et al.* Single-cell transcriptomics reveals gene expression dynamics of
human fetal kidney development. *PLOS Biology* **17**, e3000152,
doi:10.1371/journal.pbio.3000152 (2019).

15 Park, J., Liu, C. L., Kim, J. & Susztak, K. Understanding the kidney one cell at a time.
Kidney Int **96**, 862-870, doi:10.1016/j.kint.2019.03.035 (2019).

16 Fang, R. *et al.* Fast and Accurate Clustering of Single Cell Epigenomes Reveals
Cis-Regulatory Elements in Rare Cell Types. *bioRxiv*, 615179,
doi:10.1101/615179 (2019).

17 Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nat Methods* **16**, 1289-1296, doi:10.1038/s41592-019-0619-0 (2019).

18 Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137,
doi:10.1186/gb-2008-9-9-r137 (2008).

19 Park, J. *et al.* Single-cell transcriptomics of the mouse kidney reveals potential cellular
targets of kidney disease. *Science* **360**, 758-763, doi:10.1126/science.aar2131 (2018).

20 Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-
1902.e1821, doi:10.1016/j.cell.2019.05.031 (2019).

21 Wu, H., Kirita, Y., Donnelly, E. L. & Humphreys, B. D. Advantages of Single-Nucleus
over Single-Cell RNA Sequencing of Adult Kidney: Rare Cell Types and Novel Cell
States Revealed in Fibrosis. *J Am Soc Nephrol* **30**, 23-32, doi:10.1681/ASN.2018090912
(2019).

22 Frankish, A. *et al.* GENCODE reference annotation for the human and mouse genomes.
Nucleic Acids Res **47**, D766-D773, doi:10.1093/nar/gky955 (2019).

23 Zhu, Q. *et al.*, doi:10.1101/848846 (2019).

24 Cao, J. *et al.* Joint profiling of chromatin accessibility and gene expression in thousands
of single cells. *Science* **361**, 1380-1385, doi:10.1126/science.aau0730 (2018).

25 O'Brien, L. L. *et al.* Transcriptional regulatory control of mammalian nephron
progenitors revealed by multi-factor cistromic analysis and genetic studies. *PLoS Genet*
14, e1007181, doi:10.1371/journal.pgen.1007181 (2018).

26 Pliner, H. A. *et al.* Cicero Predicts cis-Regulatory DNA Interactions from Single-Cell
Chromatin Accessibility Data. *Mol Cell* **71**, 858-871 e858,
doi:10.1016/j.molcel.2018.06.044 (2018).

27   Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime
999   cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-
589, doi:10.1016/j.molcel.2010.05.004 (2010).

28   Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nature*
Methods **14**, 1083-1086, doi:10.1038/nmeth.4463 (2017).

29   Nittoli, V. *et al.* Characterization of paralogous uncx transcription factor encoding genes
in zebrafish. *Gene X* **2**, 100011, doi:10.1016/j.gene.2019.100011 (2019).

30   La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494-498,
doi:10.1038/s41586-018-0414-6 (2018).

31   Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. *Nat*
Methods **14**, 975-978, doi:10.1038/nmeth.4401 (2017).

32   Guo, J. K. *et al.* WT1 is a key regulator of podocyte function: reduced expression levels
cause crescentic glomerulonephritis and mesangial sclerosis. *Hum Mol Genet* **11**, 651-
659, doi:10.1093/hmg/11.6.651 (2002).

33   Kann, M. *et al.* WT1 targets Gas1 to maintain nephron progenitor cells by modulating
FGF signals. *Development* **142**, 1254-1266, doi:10.1242/dev.119735 (2015).

34   Nakai, S. *et al.* Crucial roles of Brn1 in distal tubule formation and function in mouse
kidney. *Development* **130**, 4751-4759, doi:10.1242/dev.00666 (2003).

35   Nilsson, D., Heglind, M., Arani, Z. & Enerbäck, S. Foxc2 is essential for podocyte
function. *Physiol Rep* **7**, e14083-e14083, doi:10.14814/phy2.14083 (2019).

36   Takahashi, T. *et al.* Temporally compartmentalized expression of ephrin-B2 during renal
glomerular development. *J Am Soc Nephrol* **12**, 2673-2682 (2001).

37   Cheng, H. T. & Kopan, R. The role of Notch signaling in specification of podocyte and
proximal tubules within the developing mouse kidney. *Kidney Int* **68**, 1951-1952,
doi:10.1111/j.1523-1755.2005.00627.x (2005).

38   Kang, H. M. *et al.* Defective fatty acid oxidation in renal tubular epithelial cells has a key
role in kidney fibrosis development. *Nat Med* **21**, 37-46, doi:10.1038/nm.3762 (2015).

39   Das, A. *et al.* Stromal–epithelial crosstalk regulates kidney progenitor cell differentiation.
Nature Cell Biology **15**, 1035-1044, doi:10.1038/ncb2828 (2013).

40   Vento-Tormo, R. *et al.* Single-cell reconstruction of the early maternal-fetal interface in
humans. *Nature* **563**, 347-353, doi:10.1038/s41586-018-0698-6 (2018).

41 Trueb, B., Amann, R. & Gerber, S. D. Role of FGFRL1 and other FGF signaling proteins
in early kidney development. *Cell Mol Life Sci* **70**, 2505-2518, doi:10.1007/s00018-012-
1189-9 (2013).

42 Costantini, F. & Shakya, R. GDNF/Ret signaling and the development of the kidney.
Bioessays **28**, 117-127, doi:10.1002/bies.20357 (2006).

43 Fan, X. *et al.* SLIT2/ROBO2 signaling pathway inhibits nonmuscle myosin IIA activity
and destabilizes kidney podocyte adhesion. *JCI Insight* **1**, e86934,
doi:10.1172/jci.insight.86934 (2016).

44 Sajithlal, G., Zou, D., Silvius, D. & Xu, P. X. Eya 1 acts as a critical regulator for
specifying the metanephric mesenchyme. *Dev Biol* **284**, 323-336,
doi:10.1016/j.ydbio.2005.05.029 (2005).

45 Hwang, D. Y. *et al.* Mutations of the SLIT2-ROBO2 pathway genes SLIT2 and SRGAP1
confer risk for congenital anomalies of the kidney and urinary tract. *Hum Genet* **134**, 905-
916, doi:10.1007/s00439-015-1570-5 (2015).

46 Wuttke, M. *et al.* A catalog of genetic loci associated with kidney function from analyses
of a million individuals. *Nat Genet* **51**, 957-972, doi:10.1038/s41588-019-0407-x (2019).

47 Hellwege, J. N. *et al.* Mapping eGFR loci to the renal transcriptome and phenome in the
VA Million Veteran Program. *Nat Commun* **10**, 3842, doi:10.1038/s41467-019-11704-w
(2019).

48 Pattaro, C. *et al.* Genetic associations at 53 loci highlight cell types and biological
pathways relevant for kidney function. *Nature communications* **7**, 10023,
doi:10.1038/ncomms10023 (2016).

49 Menon, M. C. *et al.* Intronic locus determines SHROOM3 expression and potentiates
renal allograft fibrosis. *J Clin Invest* **125**, 208-221, doi:10.1172/JCI76902 (2015).

50 Khalili, H. *et al.* Developmental Origins for Kidney Disease Due to Shroom3 Deficiency.
J Am Soc Nephrol **27**, 2965-2973, doi:10.1681/ASN.2015060621 (2016).

51 Qiu, C. *et al.* Renal compartment-specific genetic variation analyses identify new
pathways in chronic kidney disease. *Nat Med* **24**, 1721-1731, doi:10.1038/s41591-018-
0194-4 (2018).

52 Kobayashi, A. *et al.* Six2 defines and regulates a multipotent self-renewing nephron
progenitor population throughout mammalian kidney development. *Cell Stem Cell* **3**,
169-181, doi:10.1016/j.stem.2008.05.020 (2008).

53 Subramanian, A. *et al.* Single cell census of human kidney organoids shows
reproducibility and diminished off-target cells after transplantation. *Nat Commun* **10**,
5462, doi:10.1038/s41467-019-13382-0 (2019).

54   Han, S. H. *et al.* PGC-1alpha Protects from Notch-Induced Kidney Fibrosis
1066   Development. *J Am Soc Nephrol* **28**, 3312-3322, doi:10.1681/ASN.2017020130 (2017).

1067   55   Schaid, D. J., Chen, W. & Larson, N. B. From genome-wide associations to candidate
1068   causal variants by statistical fine-mapping. *Nat Rev Genet* **19**, 491-504,
1069   doi:10.1038/s41576-018-0016-z (2018).

1070   56   Cusanovich, D. A. *et al.* The cis-regulatory dynamics of embryonic development at
1071   single-cell resolution. *Nature* **555**, 538-542, doi:10.1038/nature25981 (2018).

1072   57   Calderon, D. *et al.* Inferring Relevant Cell Types for Complex Traits by Using Single-
1073   Cell Gene Expression. *Am J Hum Genet* **101**, 686-699, doi:10.1016/j.ajhg.2017.09.009
1074   (2017).

1075   58   Parikshak, N. N. *et al.* Integrative functional genomic analyses implicate specific
1076   molecular pathways and circuits in autism. *Cell* **155**, 1008-1021,
1077   doi:10.1016/j.cell.2013.10.031 (2013).

1078   59   Zhu, C. *et al.* An ultra high-throughput method for single-cell joint analysis of open
1079   chromatin and transcriptome. *Nature Structural & Molecular Biology* **26**, 1063-1070,
1080   doi:10.1038/s41594-019-0323-x (2019).

1081   60   Chen, S., Lake, B. B. & Zhang, K. High-throughput sequencing of the transcriptome and
1082   chromatin accessibility in the same cell. *Nature Biotechnology* **37**, 1452-1457,
1083   doi:10.1038/s41587-019-0290-0 (2019).

1084   61   Buenrostro, J. D., Wu, B., Chang, H. Y. & Greenleaf, W. J. ATAC-seq: A Method for
1085   Assaying Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biol* **109**, 21 29 21-
1086   21 29 29, doi:10.1002/0471142727.mb2129s109 (2015).

1087   62   Corces, M. R. *et al.* An improved ATAC-seq protocol reduces background and enables
1088   interrogation of frozen tissues. *Nat Methods* **14**, 959-962, doi:10.1038/nmeth.4396
1089   (2017).

1090   63   Chen, H. *et al.* Assessment of computational methods for the analysis of single-cell
1091   ATAC-seq data. *Genome Biol* **20**, 241, doi:10.1186/s13059-019-1854-5 (2019).

1092   64   O'Brien, L. L. *et al.* Differential regulation of mouse and human nephron progenitors by
1093   the Six family of transcriptional regulators. *Development* **143**, 595-608,
1094   doi:10.1242/dev.127175 (2016).

1095   65   Hon, G. C. *et al.* Epigenetic memory at embryonic enhancers identified in DNA
1096   methylation maps from adult mouse tissues. *Nature Genetics* **45**, 1198-1206,
1097   doi:10.1038/ng.2746 (2013).

1098   66   McLean, C. Y. *et al.* GREAT improves functional interpretation of cis-regulatory
1099   regions. *Nat Biotechnol* **28**, 495-501, doi:10.1038/nbt.1630 (2010).

1100   67   Ramirez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data
1101   analysis. *Nucleic Acids Res* **44**, W160-165, doi:10.1093/nar/gkw257 (2016).

1102   68   Lawrence, M. *et al.* Software for Computing and Annotating Genomic Ranges. *PLOS*
1103   *Computational Biology* **9**, e1003118, doi:10.1371/journal.pcbi.1003118 (2013).

1104   69   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902
1105   e1821, doi:10.1016/j.cell.2019.05.031 (2019).

1106   70   Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis.
1107   *Nature* **566**, 496-502, doi:10.1038/s41586-019-0969-x (2019).

1108   71   Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nat*
1109   *Methods* **14**, 1083-1086, doi:10.1038/nmeth.4463 (2017).

1110   72   Zhao, J., Hu, Z. Z., Zheng, X. G. & Ng, S. W. [2-(Tetra-zol-1-yl)acetato-kappaO]tris-(tri-
1111   phenyl-phosphine-kappaP)silver(I) mono-hydrate. *Acta Crystallogr Sect E Struct Rep*
1112   *Online* **65**, m1601, doi:10.1107/S1600536809048144 (2009).

1113   73   Aoki, R. *et al.* Foxl1-expressing mesenchymal cells constitute the intestinal stem cell
1114   niche. *Cell Mol Gastroenterol Hepatol* **2**, 175-188, doi:10.1016/j.jcmgh.2015.12.004
1115   (2016).

1116   74   Uhlen, M. *et al.* Proteomics. Tissue-based map of the human proteome. *Science* **347**,
1117   1260419, doi:10.1126/science.1260419 (2015).
1118

Figure Legends

Figure 1. snATAC-seq and scRNA-seq identified major cell types in developing and adult mouse kidney.

(A) Schematics of the study design. Kidneys from P0 and adult mice were processed for snATAC-seq and scRNA-seq followed by data processing and analysis including cell type identification and peak calling.

(B) UMAP embeddings of snATAC-seq data and scRNA-seq data. Using marker genes, cells were annotated into nephron progenitors (NP), collecting duct intercalated cells (IC), collecting duct principal cells (PC), proximal tubule segment 1 and 3 (PT S1 and PT S3), loop of Henle (LOH), distal convoluted tubules (DCT), stromal cells (Stroma), podocytes (Podo), endothelial cells (Endo) and immune cells (Immune). In scRNA-seq data, the same cell types were identified, with an additional proliferative population and immune cells were clustered into neutrophils and macrophages.

(C) UMAP embeddings of snATAC-seq and scRNA-seq data colored by P0 and adult batches.

(D) Genome browser view of read density in each snATAC-seq cluster at cell type marker gene transcription start sites. We used Uncx for nephron progenitors, *Nphs2* for podocytes, *Akr1c21* for proximal tubules, *Slc12a1* for loop of Henle, *Slc12a3* for distal convoluted tubule, *Trpv5* for connecting tubule, *Aqp2* for collecting duct principal cells, *Atp6v1g3* for intercalated cells, *Egfl7* for endothelial cells, *C1qb* for immune cells and *Col3a1* for stroma. Additional marker gene examples are shown in **Figure S1h**.

(E) Comparison of peaks identified from snATAC-seq data and bulk ATAC-seq data. Peaks that are identified in both datasets are colored blue, and peaks that are dataset-specific are grey.

(F) Violin plots showing cell type-specific gene expression in scRNA-seq data. With the exception of proximal tubule, the same marker genes as in snATAC-seq data were used (*Slc5a2* and *Slc22a30* for proximal tubule S1 and S3, respectively).

(G) Correlation between snATAC-seq gene activity scores and gene expression values in P0 data. The correlation of the adult dataset is shown in **Figure S1p**.

Figure 2. Cell type-specific gene regulatory landscape of the mouse kidney.

(A) Left panel: Heatmap showing examples of the cell type-specific differentially accessible peaks

(DAPs) (yellow: open chromatin, blue: closed chromatin) (full results are shown in **Supplemental**

Table 2). Middle panel: Examples of cell type-specific motif enrichment analysis using Homer.

(full results are shown in **Supplemental Table 3**). Right panel: TF expression z score heatmap

that corresponds to the motif enrichment in each cell type.

(B) Regulon activity heatmap. Each column represents a single cell, colored by cluster assignment

and ordered by hierarchical clustering; each row represents binarized regulon activities ("on-

black", "off-white") and ordered by hierarchical clustering.

(C) UMAP depiction of regulon activity ("on-blue", "off-grey") and TF gene expression (red scale)

of exemplary regulons for proximal tubule (*Hnf1a*), nephron progenitors (*Uncx*), loop of Henle

(*Ppargc1a*), proliferating cells (*Hmgb3*) and podocytes (*Mafb*). Examples of target gene

expression of the *Uncx* regulon (*Eye1*, *Hoxc8*, *Pax2*, *Spock2* and *Wnt4*) are shown in purple scale.

Expression of target genes of *Hnf1a*, *Ppargc1a*, *Hmgb3* and *Mafb* is shown in **Figure S3d**.

(D) tSNE representation of regulon density as a surrogate for stability of regulon states, as inferred

by SCENIC algorithm.

1164

Figure 3. The cellular trajectory of nephron progenitor differentiation.

(A) UMAP representation of snATAC-seq nephron progenitor differentiation trajectory towards

podocytes, proximal tubule, loop of Henle and distal convoluted tubule, respectively, as inferred

by Cicero. Cells are colored by pseudotime.

(B) UMAP representation of scRNA-seq nephron progenitor differentiation trajectory towards

podocytes, proximal tubule and loop of Henle, respectively, as inferred by Monocle3. Cells are

colored by pseudotime.

(C) Pseudotime-dependent chromatin accessibility and gene expression changes along the

proximal tubule (red), podocyte (green) and loop of Henle (blue) cell lineages. The first column

shows the dynamics of chromVAR TF enrichment score, the second column shows the dynamics

of TF gene expression values and the third and fourth column represent the dynamics of SCENIC-

reported target gene expression values of corresponding TFs, respectively. Additional examples

are given in **Figure S3e**.

1178

Figure 4. Chromatin dynamics of nephron progenitor differentiation.

Di-graph representing cell type and lineage divergence, as derived from Cicero trajectory inference.

Nephron progenitors (NP), podocytes (Podo), intermediate stage (IM), proximal tubule (PT), loop

of Henle (LOH) and distal convoluted tubule (DCT) are connected with their developmental

precursor stages and represented by ascending numbering. Arrows represent cell differentiation

along respective trajectories. Genes listed next to the trajectories were derived from analyzing gene

enrichment of differentially assessible peaks (DAPs) between two stages. Genes colored red were

derived from the opening DAPs between two stages, genes colored blue were derived from the

closing DAPs between two stages, and genes colored green were derived from opening DAPs

between two branches. Three important genes, *Foxl1*, *Hnf4a* and *Tfap2b* are shown along with

their cell type-specific accessibility peaks and immunostaining results. Peaks that were open

during the development of specific cell types are shown in red boxes. Immunofluorescence

staining of fetal mouse kidney shows FOXL1 in red along cellular differentiation (from right to

left) from early progenitor stage (asterisk) over comma-shaped (+) and S shaped bodies (cross)

towards podocytes within primitive glomeruli (#). HNF4A and TFAP2B in human adult kidney

samples (taken from the Human Protein Atlas, http://www.proteinatlas.org [74]) are visualized by

immunohistochemistry in brown.

1196

Figure 5. Cell-cell communication analysis in the developing and adult mice highlighted the

critical role of stroma in driving cell differentiation.

(A) Heatmaps showing the number of cell-cell interactions in the scRNA-seq dataset of P0 (top)

and adult (bottom) kidneys, as inferred by CellPhoneDB. Dark blue and dark red colors denote

low and high numbers of cell-cell interactions, respectively.

(B) CellPhoneDB-derived measures of cell-cell interaction scores and p values. Each row shows

a ligand-receptor pair, and each column shows the 2 interacting cell types, which is binned by cell

type. Columns are sub-ordered by first interacting cell type into stroma, podocytes, endothelial

cells, proximal tubule, loop of Henle and nephron progenitors. Color scale denotes the mean values

for all the interacting partners, where mean value refers to the total mean of the individual partner

average expression values in the interacting cell type pairs. Orange scale denotes P0, blue scale

denotes adult. Dot size denotes corresponding p values of the permutation test.

(C) Dot plots of RNA expression of important cell-cell communication candidates within the Gdnf-

Ret, Sonic hedgehog, Fgf, Bmp, Wnt and other pathways in both P0 (top) and adult (bottom)

kidney. Dot size denotes percentage of cells expressing the marker. Color scale represents average

gene expression values, orange denotes P0, blue denotes adult. Arrows indicate ligand-receptor

pairs.

1214

Figure 6. Single cell level chromatin accessibility highlighted human kidney GWAS target

genes and cell types.

(A, C, E) From top to bottom: Cicero-inferred co-accessibility of open chromatin regions in mouse

orthologues of human *Shroom3*, *Dab2* and *Uncx* loci; Gene browser view of the single nucleotide

polymorphisms within the regions; gene browser view of chromatin accessibility for nephron

progenitors (NP), collecting duct intercalated cells (IC), collecting duct principal cell types (PC),

proximal tubules segment 1 and 3 (PT S1 and PT S3), loop of Henle (LOH), distal convoluted

tubule (DCT), stromal cells (stroma), podocytes (Podo), endothelial cells (Endo) and immune cells

(Immune). Right subpanel shows violin plots of scRNA-seq gene expression in P0 (orange) and

adult (blue) kidneys.

(B, D, F) Whole kidney H3K27ac, H3K4me1 and Six2 ChIP-seq, whole genome bisulfate

sequencing (WGBS) and RNA-seq data in E15.5, P0 and adult kidney samples.

Supplemental Information

1228

Supplemental Tables

Supplemental Table 1. Cell type marker genes derived from scRNA-seq analysis.

Supplemental Table 2. Cell type-specific open chromatin derived from snATAC-seq analysis.

Supplemental Table 3. Cell type-specific motif enrichment.

Supplemental Table 4. Regulons and respective target genes inferred by SCENIC.

Supplemental Table 5. Binarized regulon activities in each cell type inferred by SCENIC.

Supplemental Table 6. Differentially expressed genes along pseudotime in distinct lineages in

scRNA-seq data.

Supplemental Table 7. Differentially accessible peaks along pseudotime in distinct lineages in

snATAC-seq data.

Supplemental Table 8. Nearest genes of differentially accessible peaks along pseudotime in

distinct lineages in snATAC-seq data.

Supplemental Table 9. GO enrichment of differentially accessible peaks along pseudotime in

distinct lineages inferred by GREAT analysis.

Supplemental Table 10. ChromVAR cell-TF enrichment score matrix.

Supplemental Table 11. Nearest genes of differentially accessible peaks at bifurcation events

along pseudotime in distinct lineages inferred by GREAT analysis.

1246

1247

Supplemental Figures

Figure S1. Quality control and data processing methods for snATAC-seq and scRNA-seq

data analysis.

(A) Insert size distribution of the 5 snATAC-seq samples showing periodic patterns.

(B) Transcription start sites (TSS) signal enrichment of the 5 snATAC-seq samples.

(C) Spearman correlation between snATAC-seq datasets and bulk ATAC-seq of binned genomic

regions.

(D) Distribution of number of unique molecular identifiers (UMIs, x axis) and promoter ratio (y

axis) in 5 samples shown by dot plot.

(E) Violin plots representing the number of accessible peaks across different clusters in the snATAC-seq dataset indicating similar distributions.

(F) UMAP representation of the snATAC-seq dataset colored by batches.

(G) Stacked bar graphs representing absolute numbers and percentages of identified cell types across snATAC-seq batches.

(H) Genome browser view of cell type-specific peaks at the TSS of marker genes for 13 cell types in the snATAC-seq dataset.

(I) From left to right: Stacked bar graphs showing the percentage of different cell types in the P0 and adult scRNA-seq datasets, tables showing the number of cells in each cell type (nCells) and corresponding percentage. NP, nephron progenitor; Podo, podocyte; PT, proximal tubule; S1, segment 1; S3, segment 3; LOH, loop of Henle; DCT, distal convoluted tubule cells; PC, collecting duct principal cells; IC, collecting duct intercalated cells; Endo, endothelial cells; Macro, macrophages; Neutro, neutrophils.

(J) UMAP representation of scRNA-seq data colored by the mitochondrial gene ratio (Mt %).

(K) Violin plots showing number of informative genes per single cell and unique molecular identifiers (UMIs) per single cell. Blue denotes adult kidney, orange denotes P0 kidney.

(L, M) Principal component (PC) representation of combined adult and P0 scRNA-seq dataset (left panel) and violin plots of corresponding embeddings values (right panel) before (L) and after (M) batch correction using Harmony.

(N) Dot plot of cell type-specific marker genes. Dot size denotes percentage of cells expressing the marker. Color scale represents average expression, orange denotes P0, blue denotes adult kidney.

(O) Feature plots of representative marker genes projected on UMAP dimension.

(P) Correlation between snATAC-seq gene activity scores and gene expression values in adult data, which is complementary to **Figure 1g**.

(Q) We provide the processed chromatin accessibility dataset via a searchable, interactive website (http://susztaklab.com/igv/). Ace2 was used as an example, and we show proximal tubule-specific enrichment of peaks at transcription start sites of the *Ace2* (Angiotensin-converting enzyme 2) gene (red boxes).

Figure S2. Characterization of the cell type-specific regulatory landscape

(A) Bar graph representing the number of accessible peaks in distal elements, promoters, introns,

5'-UTR, 3'UTR and exons, as distributed across samples of snATAC-seq data and bulk ATAC-

seq data.

(B) Overlap of scATAC-seq differentially accessible peaks among cell types with H3K27Ac ChIP-

seq data.

(C) Number of shared and unique peaks among snATAC-seq cell types. Cell types include nephron

progenitors and cells differentiated from nephron progenitors.

(D) Genome browser view of Umod as an example for distal open chromatin region and its target

promoter region.

(E) Distribution of different open chromatin elements in snATAC-seq cell types.

(F) Distribution of different open chromatin elements among differentially accessible peaks (DAPs)

in snATAC-seq cell types.

(G) Genome browser representations of single cell open chromatin data for individual cell types

at chromosomal loci around Six2 and Slc6a18, along with their known distal elements (red boxes).

Corresponding chromosomal interaction of open chromatin regions, as inferred by Cicero

(**Methods**), is depicted at the top.

(H) Genome browser views of representative marker genes demonstrating cell type-specific

chromatin accessibility for proximal tubule (*Hnf4a* and *Hmgb3*), several tubular segments (*Hnf1b*),

loop of Henle and distal convoluted tubule (*Esrrb* and *Ppargc1a*) as well as nephron progenitors

and podocytes (Wt1).

(I) UMAP depiction of regulon activity ("on-blue", "off-grey") and RNA expression (red scale) of

exemplary regulons of proximal tubule (*Hnf1a*), nephron progenitors (*Six2*), loop of Henle

(*Ppargc1a*), proliferating cells (*Hmgb3*) and podocytes (*Mafb*), respectively. Exemplary target

gene expression for the respective TF is shown in purple scale.

1312

Figure S3. snATAC-seq and scRNA-seq cell differentiation trajectories.

(A) UMAP representation of snATAC-seq trajectory lineages of podocytes, proximal tubule and

loop of Henle cells from nephron progenitors colored by 2 P0 batches.

(B) UMAP representation of snATAC-seq trajectory lineages of podocytes, proximal tubule and

loop of Henle cells from nephron progenitors colored by original cell type assignment as in **Figure**

1b.

(C) UMAP representation of scRNA-seq trajectory lineages of podocytes, proximal tubule and

loop of Henle cells from nephron progenitors colored by original cell type assignment as **in Figure**

1b.

(D) UMAP representation of RNA velocity of scRNA-seq trajectory inferred by VelocytoR,

colored by original cell type assignment. Each dot is one cell and each arrow represents the time

derivative of the gene expression state.

(E) UMAP representation of snATAC-scRNA integration results colored by cell type assignment.

(F) UMAP representation of snATAC-scRNA integration results colored by technologies

(snATAC=red, scRNA=grey). Podo: podocytes, PT: proximal tubule, LOH: loop of Henle, DCT:

distal convoluted tubule, NP: nephron progenitors, IM: intermediate stage cells.

(G) Dot plot showing snATAC-scRNA integration cell type assignment confusion matrix. Each

column represents the original cell type assignment of snATAC-seq data, and each row represents

the predicted cell type assignment by the integration analysis scRNA-seq data. Each dot represents

the number of cells that were matched in the integrated data.

(H) Heatmap of chromVAR enrichment results. The original data matrix is given in **Supplemental**

Table 10.

(I) Pseudotime-dependent chromatin accessibility and gene expression changes along the proximal

tubule (red), podocytes (green) and loop of Henle (blue) cell lineages. The first column represents

the dynamics of chromVAR TF enrichment score, the second column represents the dynamics of

TF gene expression values, and the third and fourth column represent the dynamics of SCENIC-

reported target gene expression values.

1340

Figure S4. Chromatin dynamics of nephron progenitor differentiation.

(A) Di-graph representing cell type and lineage divergence, as derived from Cicero trajectory

inference. Nephron progenitors (NP), podocytes (Podo), intermediate stage (IM), proximal tubule

(PT), loop of Henle (LOH) and distal convoluted tubule (DCT) are connected with their

developmental precursor stages and ordered by ascending numbering. Pie charts represent

differentially assessible peaks (DAPs) between two stages, where the size of pie charts is

proportional to the number of DAPs, orange color represents the number of open peaks, grey color

the number of closed peaks. Bar graphs depict gene ontology (GO) term analysis of genes nearby

DAPs derived from GREAT analysis (full list in **Supplemental Table 9**).

(B) Immunofluorescence staining of fetal mouse kidney. Upper panel and insert denote E13.5 stage,

lower panel denotes P6 mouse. Blue staining represents nuclei (DAPI), green staining represents

tubular epithelium (E-Cadherin) and red staining represents progenitor cells (FOXL1) along a

developmental trajectory from early progenitor stage (asterisk) over comma-shaped (+) and S

shaped bodies (cross) towards podocytes within primitive glomeruli (#).

(C) Pseudotime-dependent chromatin accessibility and gene expression changes along the

proximal tubule (red), podocytes (green) and loop of Henle (LOH, blue) cell lineages for important

bifurcation TFs in the podocyte (*Foxl1*) and distal tubule (*Tfap2b*) lineage.

(D) Bar graphs denote the percentage of cells with accessible chromatin of several *Six2* promoters

and enhancers (gene loci numbered 1-3) as well as putative *Foxl1* enhancers (gene loci numbered

4-7) along pseudotime. Exact gene loci of enhancers and promoters are given above each

respective graph. Changes along pseudotime are depicted for 3 lineages from nephron progenitors

(NP) to podocytes, proximal tubule (PT) and loop of Henle (LOH) cells, respectively. The right

upper subpanel depicts the genome browser overview of chromatin accessibility for the NP and

therefore corresponds to the first bar in graphs on the left. The right lower subpanel depicts zoom-

in versions of the 7 loci for all 3 lineages.

1366

Figure S5. Single cell level chromatin accessibility highlighted human kidney GWAS target

genes and cell types.

Open chromatin and co-accessibility view at alternative scales to those shown in **Figure 6**.

(A, C, E) From top to bottom: Cicero-inferred co-accessibility of open chromatin regions in mouse

orthologues of human *Shroom3*, *Dab2* and *Uncx* loci; Gene browser view of the single nucleotide

polymorphisms within the regions; gene browser view of chromatin accessibility for nephron
progenitors (NP), collecting duct intercalated cells (IC), collecting duct principal cell types (PC),
proximal tubules segment 1 and 3 (PT S1 and PT S3), loop of Henle (LOH), distal convoluted
tubule (DCT), stromal cells (stroma), podocytes (Podo), endothelial cells (Endo) and immune cells
(Immune). Right subpanel shows violin plots of scRNA-seq gene expression in P0 (orange) and
adult (blue) kidneys.
(B, D, F) Whole kidney H3K27ac, H3K4me1 and Six2 ChIP-seq, whole genome bisulfate
sequencing (WGBS) and RNA-seq data in E15.5, P0 and adult kidney samples.

Figure 1

Figure 2

A

N = 22,248 cell type-specific peaks (subset)

Normalized accessibility signal
0   3

Normalized gene expression
-1.5   1.5

Homer enrichment

Motif	% target seq	P-val
Six2	35.73	1e-270
Hoxc9	18.33	1e-151
Wt1	26.07	1e-125
Mafb	12.79	1e-20
Hnf4a	29.78	1e-491
Ppara	12.14	1e-267
Bhlhe41	18.71	1e-31
Esrrb	24.56	1e-20
Foxa1	20.89	1e-2
Vdr	4.74	1e-5
Elf5	18.49	1e-34
Tcfcp2l1	8.34	1e-74
Erg	64.77	1e-1009
Sox17	22.37	1e-189
PU.1	34.78	1e-410
Batf	13.82	1e-25
Twist1	3.05	1e-10
Nr2f2	29.11	1e-24

NP, Podo, PTS1, PTS3, LOH, DCT, PC, IC, Endo, immune, stroma1, stroma2

Cell type-specific TF gene expression

Six2, Hoxc9, Wt1, Mafb, Hnf4a, Ppara, Bhlhe41, Esrrb, Foxa1, Vdr, Elf5, Tfcp2l1, Erg, Sox17, Spi1, Batf, Twist1, Nr2f2

NP Podo PT LOH DCT PC IC Endo immune stroma

B

Cells

Regulons

NP, Podo, LOH, PT S1, PT S3, Stroma-like, Prolif, Early PT

Trps1, Hnf1b, Maf, Hnf4a, Hnf1a, Hoxd11, Ors1, Hoxc8, Hoxc9, Six2, Meox1, Hoxc6, Hoxa11, Uncx, Tcf4, Hoxc10, Hmga2, Nfia, Nr2f2, Mef2c, Fli1, Hic1, Pbx1, Irf2, Hoxb8, Hoxb4, Hoxd8, Foxq1, Hoxd9, Esrrg, Ppargc1a, Klf4, Nfat5, E2f3, Hmgb3, Sox11, Junb, Jun, Fosb, Tfdp1, Bhlhe40, Zmiz1, Jund, Irf3, Brf1, Lef1, Tead1, Foxc1, Foxc2, Wt1, Mafb, Foxd1

Cell cluster
- LOH
- PT S1
- PT S3
- Early PT
- Stroma-like
- Podo
- Prolif
- NP

Regulon activity
- ■ "On" □ "Off"

D Regulon density

high

low

LOH, PT S3, PTS1, earlyPT, Podo, Stroma-like, Prolif, NP

C

Regulon activity	TF gene expression	Target gene expression
off ☐ on ■	low→high (red)	low→high (purple)

Hnf1a — *Eya1*

Uncx — *Hoxc8*

Ppargc1a — *Pax2*

Hmgb3 — *Spock2*

Mafb — *Wnt4*

Figure 3

Figure 4

Figure 5

A *Shroom3*
C *Dab2*
E *Uncx*

chr5:92,675,198-92,787,886

chr15:6,346,126-6,415,942

chr5:139,525,148-139,576,090