The island rule explains consistent patterns of body size evolution in terrestrial vertebrates

Ana Benítez-López[1,2]*, Luca Santini[1,3], Juan Gallego-Zamorano[1], Borja Milá[4], Patrick Walkden[5], Mark A.J. Huijbregts[1,†], Joseph A. Tobias[5,†]

[1]Department of Environmental Science, Institute for Wetland and Water Research, Radboud University, P.O. Box 9010, NL-6500 GL, Nijmegen, the Netherlands.

[2]Integrative Ecology Group, Estación Biológica de Doñana, CSIC, 41092, Sevilla, Spain

[3]National Research Council, Institute of Research on Terrestrial Ecosystems (CNR-IRET), Via Salaria km 29.300, 00015, Monterotondo (Rome), Italy

[4]National Museum of Natural Sciences, Spanish National Research Council (CSIC), Madrid 28006, Spain

[5]Department of Life Sciences, Imperial College London, Silwood Park, Buckhurst Road, Ascot, Berkshire SL5 7PY, United Kingdom

*Correspondence to: ana.benitez@ebd.csic.es; abenitez81@gmail.com

†These two authors contributed equally

Abstract

Island faunas can be characterized by gigantism in small animals and dwarfism in large animals, but the extent to which this so-called 'island rule' provides a general explanation for evolutionary trajectories on islands remains contentious. Here we use a phylogenetic meta-analysis to assess patterns and drivers of body size evolution across a global sample of paired island-mainland populations of terrestrial vertebrates. We show that 'island rule' effects are widespread in mammals, birds and reptiles, but less evident in amphibians, which mostly tend towards gigantism. We also found that the magnitude of insular dwarfism and gigantism is mediated by climate as well as island size and isolation, with more pronounced effects in smaller, more remote islands for mammals and reptiles. We conclude that the island rule is pervasive across vertebrates, but that the implications for body size evolution are nuanced and depend on an array of context-dependent ecological pressures and environmental conditions.

Introduction

From giant pigeons to dwarf elephants, islands have long been known to generate evolutionary oddities[1]. Understanding the processes by which island lineages evolve remains a prominent theme in evolutionary biology, not least because they include many of the world's most bizarre and highly threatened organisms[2]. The classic insular pattern of both small-animal gigantism and large-animal dwarfism in relation to mainland relatives has been described as a macro-evolutionary or biogeographical rule – the 'island rule'[3-5] (Fig. 1). However, previous research has cast doubt on the generality of this pattern[6], suggesting that body size shifts are asymmetrical, with reduced size in some clades (e.g. carnivores, heteromyid rodents, and artiodactyls) or increased size in others (e.g. murid rodents)[7,8]. Even in these cases, the underlying mechanisms driving patterns of insular gigantism and dwarfism remain unclear.

Several mechanisms have been proposed to explain the island rule, including reduced predation, relaxed competition and resource limitation in island environments[9]. In theory, each of these factors may be accentuated in smaller, more isolated islands, where lower levels of interspecific competition and predation could lead to 'ecological release', allowing small-bodied species to increase in body size[5,9]. Conversely, among large-bodied species, limited resource availability could select for smaller body sizes with reduced energy requirements, leading to insular dwarfism. Climatic conditions may also influence body size evolution on islands since primary productivity and associated resource availability are strongly influenced by climate[9,10]. Although previous studies of body size evolution on islands have tested the effects of these different mechanisms, many have focused on relatively restricted geographic and taxonomic scales and did not directly address the island rule in its broad sense across multiple species within a taxon[10-13], with notable exceptions[9,14-16].

Most work on the island rule has been restricted to mammals (e.g.[4,7,14,17]), although the hypothesis has also been tested in amphibians[18], reptiles[19-21], birds[15,22], fish[23], insects[24], molluscs[25], and plants[26]. The highly inconsistent results of these studies (e.g.[5,6,27]) are perhaps unsurprising because they typically deal with single species or pool together data on different traits from numerous sources without controlling for variation in study design or accounting for sampling variance. Accordingly, a recent systematic review based on a simplified scoring system[27] concluded that empirical support for the island rule is not only potentially biased but

also generally low, particularly for non-mammalian taxa. However, scoring approaches provide

only limited information as they do not account for heterogeneity between studies, taxonomic

representativeness, sample size, or precision in the estimates.

These limitations are best addressed with formal meta-analyses[28,29], hence we tested the island

rule hypothesis by applying phylogenetic meta-regressions to a global dataset of 2,479 island-

mainland comparisons for 1,166 insular and 886 mainland species of terrestrial vertebrates

(Supplementary Dataset 1, Fig. 2). Our analytical framework allows us to control for multiple

types of variation, including data source, sample size imbalance, intraspecific and intra-

population variability, and phylogenetic relatedness (see Methods). For each island-mainland

comparison, we calculated the log response ratio ($lnRR$) as the natural logarithm of the ratio

between the mean body size of individuals from an insular population M_i and that of mainland

relatives M_m ($lnRR = \log[M_i/Mm]$)[30]. Then, we regressed $lnRR$ against the body mass of the

mainland population (M_m)(Fig.1).

This framework provides a clear set of predictions in the context of evolutionary trajectories on

islands[4,6,14]. Specifically, since negative values of $lnRR$ indicate dwarfism and positive values

indicate gigantism, a positive intercept and negative slope of the $lnRR$-mainland mass

relationship supports the island rule (Fig. 1). Given the contentiousness of the generality in the

island rule, we assessed the robustness of our results against potential biases derived of

regressing ratios[31,32], using small samples, imputing missing data, or the influence of using data

from the island rule literature or derived from other studies focused on unrelated questions (i.e.

publication bias; see Methods). Finally, we use our framework to assess how body size shifts are

related to island size, island isolation, island productivity and climate, as well as species diet.

The extent to which these different factors explain insular body size shifts allows us to re-

evaluate a range of hypotheses for the mechanisms underlying "island rule" effects on body size,

including ecological release[9], immigrant selection[9], resource limitation[9,33,34],

thermoregulation[9,15,35], water availability[36,37] and starvation resistance[9,33] (Supplementary Table

1, Extended Data Fig. 1).

Results

The generality of the island rule

We found that *lnRR* and mainland body mass were negatively related for mammals, birds and reptiles, with small species tending to gigantism and large species to dwarfism (Fig. 3). The relationship was weakly negative but statistically non-significant for amphibians, with a tendency towards gigantism across all body sizes (Fig. 3, Table 1). We obtained similar results using size ratios corrected for small sample size (*lnRR*$^{\Delta}$), or by regressing island mass against mainland mass, with support for the island rule across all groups except for amphibians (Supplementary Table 3-4). This indicates that our analyses are robust to small sample size bias[38] or any potential spurious correlation associated to ratio regression models[31,32] (Extended Data Fig. 2). Further, neither imputation nor publication bias influenced our results (Supplementary Table 5-6), with no apparent differences between island-mainland comparisons sampled from studies formally testing the island rule or compiled from unrelated data sets.

Mainland body mass explained 11.4, 7.0 and 17.6% of the variance in mammals, birds and reptiles, respectively. The amount of further variance accounted for by phylogeny (0.0–29.8%), data source (1.8–25.1%), and species (25.9–53.2%) fluctuated widely among taxa (Extended Data Fig. 3). Phylogeny accounted for a relatively large amount of variance in mammals (20.1%) and reptiles (29.8%), but even in these cases the overall patterns were not driven by large effects in particular clades. Some groups tended towards gigantism and others towards dwarfism, while others contained both dwarfs or giants depending on body size (e.g., Primata, Rodentia, and Carnivora in mammals, and Viperidae, Scincidae and Iguanidae in reptiles; Extended Data Fig. 4).

Ecological mechanisms underlying body size evolution on islands

The pattern of body size evolution in our island-mainland comparisons provides some insight into the likely mechanisms driving "island rule" effects (Extended Data Fig. 5-8, Supplementary Table 7, Supplementary Dataset 2). Overall, insular size shifts arise through some combination of ecological release from predation and competition, resource limitation, biased colonization (i.e. immigrant selection), and starvation resistance. The fact that no single factor explained island effects on body size is not surprising because some hypotheses shared overlapping predictions, making them difficult to disentangle.

Shifts in body mass of mammals were mostly explained by island size and spatial isolation ($Q_m =$

12.20, P = 0.002, Fig. 4a), resulting in more pronounced gigantism or dwarfism in small and

remote islands. Birds showed similar size shifts in relation to spatial isolation and island area, but

these were not statistically significant (Supplementary Table 7). In both mammals and birds,

temperature had similar effects across the size range, with body size consistently larger in cool

islands and smaller in warm islands (Extended Data Figs. 5e, 6e). Hence, in these groups, even

large species that had undergone dwarfism were significantly larger in cool insular environments

than in warm ones. Contrary to the starvation resistance hypothesis, small-sized birds did not

become larger in highly seasonal islands, but large-sized birds had reduced dwarfism on islands

with high seasonality in temperatures ($Q_m =$ 12.33, P < 0.001, Extended Data Fig. 6).

In reptiles, the combination of island area and spatial isolation were the most important factors

explaining variation in body size (Fig. 4c), with productivity and seasonality also supported but

with weaker effects (Extended Data Fig. 7). Similar to mammals, the tendency towards dwarfism

or gigantism in large-bodied or small-bodied reptiles was more apparent in isolated small-sized

islands, with stronger effects of area than isolation (Supplementary Table 7). The effects of

productivity and seasonality were only partially in line with predictions, as small-sized species

were larger on islands with high seasonality, but smaller on islands with high productivity

(Extended Data Fig. 7). In turn, large-bodied reptiles were smaller on islands with low

productivity and high seasonality.

Finally, the relationship between size ratio and mainland mass in amphibians was slightly steeper

in small and remote islands (Fig. 4d), with island area being marginally more important than

spatial isolation (Extended Data Fig. 8). The effect of seasonality was clearer, with amphibian

species inhabiting islands with high seasonality (unpredictable environments) tending toward

gigantism, whereas those from islands with low seasonality (predictable environments) being

similar in size to mainland counterparts (Extended Data Fig. 8). We found no effects of diet for

any of the four taxa, or precipitation for amphibians, contrary to the water availability

hypothesis.

Discussion

Based on comprehensive morphometric data from a worldwide sample of island fauna, we show consistent patterns of body size evolution across terrestrial vertebrates in accordance with predictions of the island rule. This finding was robust to alternative modelling approaches (island mass vs mainland mass regressions), small sample bias, data imputation, and publication bias. Moreover, we have demonstrated that insular size shifts are contextual and depend not only on the body size of mainland relatives (island rule *sensu stricto*) but also on the physiographic and climatic characteristics of particular island environments[9].

Repeated evolutionary trajectories on islands

We found a clear negative relationship between insular body size variation and the body mass of mainland individuals in mammals, birds, and reptiles. Mainland body mass explains between 7.0 and 17.6% of the variation in insular size divergence in these three taxonomic groups, which is similar to that reported in smaller-scale studies of bats (15%), birds (13%), snakes (42%), non-volant and terrestrial mammals (11–21%), and turtles (8%)[5,14,15,39,40]. Contrary to these earlier studies, our analyses are corrected not only for phylogenetic relatedness, but also for variability between species and intrapopulation variability, thereby strengthening the evidence for predictable evolutionary trajectories on islands. Nevertheless, the island rule provides only a partial explanation for these trajectories because substantial variation around the trend line remains unexplained. We also conducted the first multispecies test of island rule effects in amphibians, showing that the relationship goes in the expected direction but with a weak effect (1.4 %), possibly because the body mass range in amphibians is narrower and limited to small sizes (~ 0.5-50 g) and thus most amphibians tend to gigantism on islands with reduced predation risk.

Our findings are in contrast with a number of studies rejecting the island rule, including a recent review of evidence from across mammals, birds and reptiles[27], as well as other taxon-specific studies focused on lizards[20,41] and turtles[21]. On the other hand, the patterns we detect are consistent with analyses supporting the island rule in snakes[19], mammals[4,9] and birds[5,15] We conclude that the contradictory results of previous studies may have been related to sampling bias, heterogeneity between sources and species, and phylogenetic relatedness (i.e. statistical non-independence). By accounting for these effects in our global models we are able to

demonstrate that vertebrate animals evolve in largely consistent ways on islands. Further, we

have shown that the island rule is not clade-specific and instead applies to numerous clades

within major taxonomic groups, particularly in mammals and birds.

A corollary that emerges from the island rule is that body size converges on islands. Specifically,

if insular environments select for intermediate body sizes, closer to the optimal size of the focal

clade, then the size spectrum of organisms found on islands should be narrower compared to the

mainland[42,43]. Theoretically, the optimal body size towards which small and large species may

converge in low-diversity systems such as islands should correspond to the point where the trend

intersects the horizontal dashed line in the relationship between size ratio and mainland mass, at

which point fitness is maximized[42] (but see[44]). Interestingly, the shift between dwarfism and

gigantism in our models occurred at approximately 100-250 g in endotherms, slightly larger than

the 100 g adult body mass proposed for mammals[42] (but see[43]), and the mode of the global body

size distribution of birds that separate between small- and large-bodied species (60 g)[22,45,46].

Additionally, our analyses suggest that the optimal body size for island reptiles should be ca. 20

g, which is marginally higher than the modal body size of Lepidosaurs (14.1 g)[47]. Whether there

is an optimal body size in island biotas has been the subject of much debate[44], but overall we

expect that phenotypic variability in morphometric traits will be substantially narrowed if

directional selection is operating in island assemblages, a feature that warrants further

investigation. Additionally, optimal phenotypes should vary with the environmental

characteristics of islands, in particular their area and isolation, climate, productivity and

seasonality. For example, in mammals, our results suggest that the optimal body size would be

ca. 100 g and ca. 900 g in warm and cold islands, respectively.

Ecological mechanisms influencing body size variation

Because body size is intimately linked to many physiological and ecological characteristics of

vertebrates, it may be associated with a variety of environmental factors. As a consequence, the

body size of colonizing species may predictably evolve as the result of selective pressures

associated with insular environments (e.g., low food resources, few competitors, no predators)

and others that act across larger geographic scales (e.g., climate). For mammals and reptiles, our

results suggest that insular body size shifts are indeed governed by spatial isolation and island

size, with individuals becoming dwarfs or giants in remote islands of limited size. Furthermore,

the slope of the relationship between size ratio and mainland mass was slightly steeper for birds

and amphibians in small remote islands than in large islands near continental land masses (Fig.

4). This points to a combination of resource limitation (with small islands having fewer resources

to maintain large-sized organisms[48,49]) along with release from interspecific competition and

predation pressure in small, species-poor islands. The pattern is also consistent with biased

colonization favoring larger individuals with higher dispersal abilities (immigration selection[50]).

Conversely, our results showed that body size divergence on islands close to the mainland was

minimal, reflecting two non-mutually exclusive processes. First, many of these islands were

connected to the continent by land bridges so recently that phenotypic differences have not had

time to accumulate. Second, regular dispersal between mainland and island populations promotes

gene flow, with introgression counteracting divergent selection[51,52].

Besides island physiography (area and isolation), other relevant factors were temperature

conditions in endotherms and resource availability and seasonality in ectothermic organisms.

Mammals and birds both responded to island temperature in line with the heat conservation

hypothesis, with small- and large-sized species exhibiting exacerbated gigantism and diminished

dwarfism, presumably to conserve heat in colder, harsher insular environments. Additionally,

temperature seasonality was an important determinant of the size of large-bodied birds, with

populations on highly seasonal islands being similar in size to mainland populations. One

possibility is that larger size in these cases may help maintain energy reserves during periods of

low food availability, allowing them to thrive in otherwise hostile environments. Another

possibility is that bird populations on highly seasonal islands – which tend to be situated at

relatively high latitudes – are more often seasonally mobile or even migratory, potentially

increasing gene flow with mainland populations or weakening adaptation to the local

environment[53]. These findings add new insights to previous results regarding the role of thermal

and feeding ecology on morphological divergence in island birds[54,55]. Traditionally, changes in

feeding ecology were thought to be the prime force in driving morphological divergence in

island birds[54,55]. Yet, our results imply that physiological mechanisms related to heat

conservation ('thermoregulation hypothesis') and energy constraints ('starvation resistance

hypothesis') may also shape body size evolution in island birds.

In reptiles, we find some evidence that resource availability and seasonality are important factors

explaining body size evolution, with some deviations from the patterns predicted. As

hypothesized, large species are much smaller on islands with low resource availability, and small

species are larger on islands with high seasonality. Yet, unexpectedly, small species are larger on

islands with low productivity, perhaps because increased intraspecific competition favors large

individuals under the high population densities that reptiles often attain on islands[56,57].

Overall, most amphibians tended to gigantism, presumably as a result of increased growth rate or

lower mortality due to reduced predation pressure on islands[58]. Additionally, we found that body

size of amphibians consistently increased on islands where resources were highly seasonal and

unpredictable, perhaps to maximize energy reserves and withstand long periods without food, for

example during aestivation or hibernation[59]. We did not find a clear relationship between

precipitation and body size, suggesting that water availability is not a key factor. It appears that

gigantism in island amphibians is mostly driven by physiological mechanisms that maximize

growth rate, particularly in smaller, more isolated islands. These findings should be further

explored when more data on island-mainland pairwise populations of amphibians become

available.

Body size evolution in extinct species

Our analyses focused on extant species for which we could gather information on the variation

around the morphometric estimates, along with sample size (essential for meta-analyses). The

widespread extinction of large species on islands, including dwarf morphotypes of large species

such as insular elephants in Sicily and the Aegean islands[60,61], may have masked the historical

pattern of phenotypic variation on islands[62]. Giant insular birds[54,63], primates[64,65], and lizards[66],

along with large insular turtle species, went extinct during the Holocene and late Pleistocene[67],

most likely because of overhunting and the introduction of invasive species[68,69]. Overall, it is

estimated that human colonization of oceanic islands was followed by the extinction of 27% of

insular endemic mammals[70], as well as over 2000 bird species in the Pacific region alone[71], with

these losses biased towards large-bodied, flightless, ground☐nesting species[68]. Extinct species

may shed new light on size evolution in insular vertebrates because species extinctions have

substantially altered the biogeography of body size in island faunas, potentially leading to

downsized insular communities[72,73]. For example, the predominance in our dataset of smaller-

bodied organisms could reflect the extinction of large species on islands[68], or simply the fact that

few islands support large species. Either way, further studies should include data from extinct

species as this may alter or strengthen the signal that we report for extant species[39].

We foresee that, under global change, the extinction of insular species and the introduction of

novel (invasive) species may trigger new equilibria, with concomitant shifts in the composition

of insular communities and the opening of novel niches to which species may respond via

genetic adaptations and phenotypic plasticity. Recent evidence indicates that even introduced

species on islands, which were not included in our analysis, predictably evolve towards dwarfism

or gigantism[74-76]. In theory, as the Anthropocene gathers pace, further extinctions will drive a

decline in mean body size of the overall island community, pushing optimal body sizes towards

the lower end of body size ranges in the different vertebrate groups.

Conclusions

Of the many evolutionary implications of living on islands – together known as the 'island

syndrome'[2] – the effects on body size are the most widely known and controversial. We have

shown that these 'island rule' effects are widespread in vertebrate animals, although the evidence

for amphibians is inconclusive. Morphological changes were directional for species at the

extremes of the body size range in mammals, birds and reptiles, following the predicted pattern

of convergence towards intermediate "optimum" body sizes, in line with optimal body size

theory[42,43,45]. Although this convergence towards morphological optima may result from natural

selection or phenotypic plasticity, the exact mechanism producing these changes on islands is

still not well understood. Nonetheless, we found that consistent transitions towards intermediate

body sizes were associated with a combination of factors, indicating a range of different

ecological mechanisms. Our results highlight the contextual nature of insular size shifts, where

island physiographic, climatic and ecological characteristics play a fundamental role in shaping

body size evolution, reinforcing the idea that large-scale macroevolutionary patterns do not arise

from single mechanisms but are often the result of multiple processes acting together[77,78].

Methods

Data collection

We collected baseline morphometric data from articles included in a recent assessment of the island rule[27], as well as other compilations assembled to test the hypothesis in reptiles[20], mammals[6], and birds[15]. To expand this sample, we then performed a literature search (February 2020) in Web Of Science Core Collection (WOS) using the following search string: ("island rule" OR "island effect" OR "island syndrome" OR island*) AND (gigantism OR dwarfism OR "body size" OR weight OR SVL OR snout-vent length OR length OR size) AND (mammal* OR bird* OR avian OR amphibia* OR reptile*) (Appendix 1). Because this search was complementary to the data we have gathered from previous compilations[6,15,20,27], we only downloaded the first 500 hits out of a total of 33,431 hits ordered by relevance, and removed duplicates already included in our dataset. We reviewed every island-mainland comparison reported in published studies and traced primary source data when possible to extract original measurements. We also extracted data from all studies containing morphometric measurements for insular populations when these could be matched with equivalent data published elsewhere for relevant mainland taxa. We excluded problematic data, such as comparisons that were not supported by taxonomic or phylogenetic evidence, or which reported morphometric data restricted to single specimens or without sample size. In addition, we excluded comparisons based on extinct taxa since they are often known from very few or incomplete specimens (Supplementary Dataset 3).

It has been argued that research on the island rule might be prone to ascertainment bias, where researchers are more likely to notice and measure animals of extreme body size when conducting research on islands[41]. To help overcome this problem, we collected body size data not only from studies testing the island rule, or reporting dwarfism and gigantism in island fauna, but also from studies that did not specifically test hypotheses related to the island rule. We matched unpaired insular populations with independent data from mainland populations by performing species-specific searches in WOS and Google Scholar. We also compiled morphometric data for 442 insular and 407 mainland bird species from an independent global dataset of avian functional traits[79].

Large islands may be more 'mainland like' in relation to factors that are thought to affect body

size (i.e. competition, resource availability and predation[5]). Thus, when major islands were at

least 10 times larger than a nearby island, we treated the large island as the mainland

comparison, following previous studies testing the island rule[4,5,20]. Consequently, a single mid-

sized island can simultaneously be treated as the continent in comparisons with smaller islands,

and the island in comparisons with larger continents. When authors reported data referring to an

entire archipelago instead of a specific island (3.2% of cases), we used the size of the largest

island as island area. Removing these cases from our analyses did not qualitatively affect our

results (Supplementary Table 8).

Our final dataset contained 529 data sources and 2,479 island-mainland comparisons[7,10,36,58,79-

602]. In total, we collated morphometric measurements for 63,561 insular and 154,875 mainland

specimens representing mammals (1,058 island-mainland comparisons), birds (695

comparisons), reptiles (547 comparisons) and amphibians (179 comparisons) from across the

globe (Fig. 2). 2,068 island-mainland comparisons (83.4%) were within species (e.g. subspecies)

comparisons, and 411 (16.6%) were between-species comparisons. Insular populations were

sampled from an array of islands varying widely in size (0.0009–785,753 km^2), climate, and

level of spatial isolation (0.03–3,835 km from mainland). To explore the drivers of body-size

shifts in insular populations, we also sampled species with a wide range of average body masses

(0.18–234,335 g). We collated data on body size indices (body mass, body length, cranial and

dental measurements) of different taxa in island and mainland populations following strict

morphological, phylogenetic and biogeographic criteria. Specifically, we always compared the

same body size index for island and mainland populations. For within-species comparisons, we

compared island and mainland populations based on the information given by the authors of the

relevant study (e.g. taking note of which mainland source populations are likely to inhabit a

particular island because of colonization history or isolation via rising sea levels[89,101,240,386,548]).

When we matched comparisons independently, we used information published in the study

reporting the insular form, selecting the geographically closest mainland population whenever

possible. In addition, we prioritized latitudinal alignment of mainland and island populations to

avoid confounding effects of latitudinal variation in body size. In the case of island endemics, we

compared island populations to their closest mainland relative whenever these were identifiable

by phylogenetic data or other information reported in each particular study. This usually meant

selecting their sister species or the geographically closest representative of a sister clade or

polytomy (Supplementary Dataset 1). If we could not reliably establish the closest mainland

relative, we discarded the data (see Supplementary Dataset 3).

When more than one body size index was reported in published studies, we prioritized those

indices related most closely to body mass (Supplementary Table 2). For mammals, we selected

indices in this order of preference: body mass, body length, cranial length (greatest skull length

or condylobasal length), and dentition (e.g. canine length)[5]. For birds, preferred indices were

body mass, wing length, tarsus length and bill length. Finally, for amphibians and reptiles, size

was reported as body mass, snout-vent-length (SVL), carapace length (CL, for turtles) and total

length (TL, including SVL and tail length). In all cases, we included measurements for adults

only. To avoid size biases attributable to sexual size dimorphism, we calculated the pooled mean

for both sexes and the combined SD using standard formulae for combining groups[603]. When

information was only available for one sex (male or female), we restricted our size comparisons

to the sex for which we had morphometric data in both mainland and island populations. Data

from zoos or studies that could not be georeferenced were discarded.

To overcome the problem that different authors report size using different indices, we used

allometric relationships to convert island and mainland size to body mass equivalents, thereby

enabling cross-taxa and cross-study comparisons. Although this conversion is imprecise,

morphological indices and body mass are nonetheless highly correlated across the global scale

and wide range of body sizes within our samples (providing more accurate predictions than

simply assuming an exponent ~3, as in previous studies testing the island rule[5,9]). We used

published allometric relationships where available (see Supplementary Table 2), or derived them

based on published datasets[47,179,604-609] and other data sources (Supplementary Dataset 4). To

calculate allometric relationships, we used OLS (Ordinary Least Square) models of the $\log^{10}$

transformed body mass against the $\log^{10}$ transformed body size index (e.g. condylobasal length,

Supplementary Table 2, Supplementary Dataset 4).

For birds, we complemented published data with standardized morphometric measurements from

3,618 museum specimens and live individuals of 436 insular and 404 mainland bird species

(see[79]). We used wing length in the main analyses instead of tarsus length because the former is a

better predictor of body mass in our dataset ($R^2_{wing} = 0.89$ vs $R^2_{tarsus} = 0.69$, Supplementary Table

2) (see also[610]). Although wing length may change during moult or thereafter because of wear,

these effects are negligible in relation to interspecific differences[79] and minimized by calculating

averages across multiple individuals. Further, interobserver differences between measurements

may explain some variation in wing length estimates, but again this bias was shown to have

negligible effects in our dataset by comparing repeated measures from different observers

(see[79]). To assess the consistency in our results, we repeated analyses using tarsus length, another

popular proxy of overall body size in birds[611]. Our results were unchanged (Fig. S2).

To select suitable comparisons for museum specimens, we first classified species as either

insular or continental by overlapping IUCN range polygons with a GIS land layer including

continental land masses. For each insular species we then identified continental sister species

from avian phylogenies[612], using the method described above. We excluded bird species that are

highly pelagic or aerial (e.g. swifts) and fully migratory species because in these groups it is

unclear whether insular and mainland forms experience different environments[15]. Further, we

also excluded flightless bird species, because morphological changes may be due to

flightlessness rather than island dwelling per se[15].

We calculated the response ratio (*lnRR,* eq. 1) as effect size in our meta-regressions, where we

divided the mean body mass of individuals from an insular population $\bar{M}_i$ by that of the nearest

mainland relative, $\bar{M}_m$, and then applied the natural logarithm. Unlike unlogged ratios, the

sampling distribution of *lnRR* is normal, particularly for small samples[30], and thus less prone to

statistical artefacts associated with ratio-based regressions.

$$lnRR = \ln\left(\frac{\bar{M}_i}{\bar{M}_m}\right)$$ (Eq. 1)

Response ratios greater than zero indicate a shift towards larger sizes (gigantism) whereas ratios

less than zero indicate shifts towards smaller sizes (dwarfism). Besides mean measurements, we

recorded measures of variation, i.e. standard deviation (SD), standard error (SE) or coefficient of

variation (CV), and sample sizes of the body size indices in island and mainland organisms. SD

and sample sizes were used to calculate sampling variances (Eq. 2), which were then used to

weight each response ratio (coupled with the amount of heterogeneity, i.e. the variance in the

underlying effects)[30].

$$\hat{\sigma}^2(lnRR) = \frac{SD_i^2}{N_i \bar{X}_i^2} + \frac{SD_m^2}{N_m \bar{X}_m^2} \; ; \qquad\qquad\qquad\qquad (\text{Eq. 2})$$

SDs were extracted from raw data when possible. If ranges were provided instead of SD (or SE

or CV), we calculated SD following[613]. If neither ranges nor measures of variation were reported,

but the reported sample size was > 1, we imputed SD based on the coefficient of variation from

all complete cases ("Bracken approach"[614]). Imputation was done for 22% of all cases in

mammals, 1.1% in birds, 11% in reptiles and 7.3% in amphibians, all within the upper limit of

imputations (<30% of all cases per group) advised in previous studies[90].

For each study and island-mainland comparison, we compiled the mainland and island names,

the study reference, the body size index used, the geographic coordinates, the distance to the

closest mainland (spatial isolation, km) and the island area (km^2). We completed missing data on

island characteristics using the UNEP island database (http://islands.unep.ch/) and the

Threatened Island Biodiversity Database (TIB, http://tib.islandconservation.org/). Missing

information was calculated using Google Earth. Additionally, we extracted the Normalized

Difference Vegetation Index (NDVI) as a proxy for resource availability on islands[615]. We also

calculated the standard deviation of NDVI to assess seasonality in leaf or vegetation cover, as an

index of seasonality in available resources. NDVI was downloaded from NASA Ames

Ecological Forecasting Lab (https://lpdaacsvc.cr.usgs.gov/appeears/task/area).

Because climate influences both resource requirements and primary productivity, body size

evolution should also be influenced by climatic conditions on islands. We thus extracted island

climatic conditions from WorldClim v. 2.0 (http://worldclim.org[616]). Specifically, we used

variables that are more closely associated with the proposed underlying mechanisms of

Bergmann's rule (i.e. thermoregulation and starvation resistance): mean annual temperature,

annual precipitation, and seasonality of temperature and precipitation[617]. We assumed that the

time period for these bioclimatic variables (1970–2000), although not necessarily matching the

actual time period of body size evolution in the insular populations, roughly represents the

climatic conditions in the Holocene, a period relatively climatically stable where most of our

populations became isolated (i.e., after the last glacial maximum; see also[9]). Because climatic

variability across cells substantially exceeds variation within cells in the Holocene, current layers

are considered adequate for geographic comparisons. All spatial variables were downloaded at

0.1-degree resolution, and we averaged all cells per island to obtain a mean value of each

environmental variable (e.g., temperature, NDVI, precipitation, etc.). Finally, for each species

included in our dataset, we collated diet information from EltonTraits for birds and mammals[618],

and from other sources for reptiles[608,619], and classified species as carnivores (> 50% diet

consisting of vertebrates) or non-carnivores (< 50% diet consisting of vertebrates), following

previous studies[79,620]. As all amphibians in our dataset are carnivores[621], we did not record their

diet.

Data analyses

To test the island rule hypothesis, we used phylogenetic meta-regressions between *lnRR* and

body mass of mainland relatives, following most previous studies of the island rule (e.g. [4,5,7]

[6,622,623]). A negative slope for this relationship would support the island rule (Fig. 1).

The use of multiple populations of the same species can overestimate the actual number of

degrees of freedom, generating type-1 errors. We controlled for this by adding 'Species' as a

random effect intercept in our analyses. Additionally, body size evolution in insular vertebrates

is heavily influenced by phylogenetic effects, with species within entire clades seemingly

showing either dwarfism or gigantism[6]. Thus, we accounted for phylogeny by including the

phylogenetic relatedness correlation matrix as a random effect. The species term captures the

similarities of effect sizes within the same species, while the phylogenetic term represents the

similarity due to relatedness[624]. We also added 'Source' as a random effect intercept to account

for between-source variability and the fact that we had multiple response ratios per study. In

some cases, 'Source' represented the combination of two sources of data, one for the island size

and one for the mainland size. Finally, we included an observation level random effect, which

represents the residual variance that needs to be explicitly modelled in a meta-analysis[29]. Total

heterogeneity, and heterogeneity due to phylogeny, source and species identity were computed

following Nakagawa & Santos (2012)[29].

We tested the robustness of our results against several potential limitations. Because multiple

island populations were often compared with a single mainland population, we accounted for

these repeated measures in a variance-covariance matrix where the diagonal includes the

sampling variances, and the off-diagonals of the matrix represent the shared variance

(covariance) among the response ratios due to the common mainland population[625]. Further, we

compared our main results to models fitted with *lnRR* and sampling variances corrected for small

sample size[38]. Another potential problem is that regressions using ratios may lead to spurious

correlations[31,32]. Thus, we conducted an additional analysis testing the statistical significance of

body size trends by regressing island mass against mainland mass, following previous

studies[4,5,20,41]. Phylogenetic meta-regressions were run using island mass as the response

variable, and mainland mass as the predictor (both transformed with natural-log), with random

effects as specified above, and sampling variance $sd_i^2/mass_i^2*N_i$. This approach has some

limitations in being harder to visualize and less effective in considering the sampling variance of

measurements (representing intrapopulation variability), yet nonetheless provides an alternative

approach for assessing the robustness of our results, in line with previous studies[4,5,20,41]. Finally,

we assessed publication bias by testing the influence of data source on the relationship between

size ratio and mainland mass. This involved comparing whether patterns differed in island-

mainland pairs extracted from studies testing the island rule (38.6% of cases) versus pairs

extracted from studies not testing the island rule (61.4% of cases).

Testing ecological hypotheses explaining insular size shifts

To evaluate the relative role of key mechanisms proposed to influence body size evolution in

island fauna, we compiled a further range of variables (Supplementary Table 1, Extended Data

Fig. 1). These included island area (linked to both resource limitation and to ecological release

from both predation and competition) and spatial isolation (linked to reduced colonization from

mainland populations for smaller taxa, i.e. immigration selection[50]). In addition, we included

climatic and resource seasonality, which are linked to the starvation resistance hypothesis, and

productivity and species diet, each of which are linked to resource limitation. Because body size

evolution may be influenced by climate (e.g. Bergmann's rule)[9,626], we also included mean

temperature, which is linked to body size adaptations for enhancing heat conservation or

dissipation (thermoregulation hypothesis). For amphibians, we included precipitation as a proxy

for water supply linked to aquatic habitats, moisture and humidity (water availability

hypothesis).

We modelled interactions between body size and each of the explanatory variables because we

expected these factors to differentially affect species of different sizes, thus producing different

effects in small, medium-sized and large species. In line with the ecological release and resource

limitation hypotheses, we expected the slope of the *lnRR*-mainland mass relationship to be

steeper in smaller islands, isolated from the mainland and with fewer or no predators (Fig. 1).

Further, if resource availability is a key factor, we also expected large species to undergo

dwarfism on islands with low productivity[48,49], and for dwarfism to be accentuated in dietary

niches with high energy requirements, including carnivory[9]. In addition, high seasonality in

resources and in temperature was expected to result in increased gigantism in small-sized

species, because energy reserves increase faster than energy depletion as body size increases

(starvation resistance hypothesis)[9,627]. We hypothesized that smaller species would benefit

comparatively more by increasing in size than larger species. Because amphibians are generally

small-sized, we also fitted a model for this group with only additive terms (mainland mass +

sdNDVI) where seasonality in resources would result in larger body sizes for all species. Finally,

mechanisms driven by thermoregulation and water availability predict that body size shifts are

associated with temperature and rainfall, respectively. Mean temperature was expected to

predominantly affect endotherms and small ectotherms with good thermoregulating abilities

(reptiles and anurans) living on cold islands which, compared to similar-sized species on islands

with a mild climate, would exhibit more pronounced gigantism to enhance heat conservation. We

fitted the effect of temperature as an interactive (mainland mass × Tmean) or additive term

(mainland mass + Tmean) to assess whether only small species or all species would increase in

size in low temperature islands (see details in Supplementary Table 1, Extended Data Fig 1,

Supplementary Table 7).

Prior to modeling, all the moderators (explanatory variables) were inspected and $\log_{10}$-

transformed if necessary to meet normality assumptions in model errors. We considered a result

to be significant when the 95% confidence interval (CI) did not cross zero. We assessed the

explained heterogeneity using Omnibus test for moderators (Q_m) and the percentage of variance

explained by the moderators using R^2 marginal[628]. All figures show the relationship between size

response ratio and body mass, and how this might be altered by the mechanisms explained

above.

All analyses were performed in R 3.5.3[629] using the packages *metafor v2.0*[630] and *metagear*

v0.4[631] for the meta-regression models and data imputation, *metaDigitise v1.0*[632] for data

extraction from plots, *ape v5.2*[633] for estimating branch lengths and resolving polytomies, *rotl*

v3.0.4[634] for building the phylogenies for our species by searching the Open Tree Taxonomy[635]

and retrieving the phylogenetic relationships from the Open Tree of Life[636], *sf v0.7-3* [637] and

raster v2.7-15[638] for spatial analyses, *dplyr v0.8.0.1*[639] and *reshape2 v1.4.3*[640] for data

manipulation and *ggplot2 v 3.3.0.9000*[641] and *ggpubr v0.1.8* [642] for data visualization. ArcMap

10.5 was used for Figure 2. Silhouettes in figures were extracted from 'phylopic'

(https.phylopic.org). The PRISMA Checklist for systematic reviews is available in Appendix 3.

Data availability

542   All data are available at https://github.com/anabenlop/Island_Rule and

543   https://figshare.com/projects/Body_size_evolution_in_insular_vertebrates/89102.

Code availability

545   The code to conduct the analyses is available at https://github.com/anabenlop/Island_Rule.

References

1   Foster, J. B. Evolution of mammals on islands. *Nature* **202**, 234-235 (1964).

2   Baeckens, S. & Van Damme, R. The island syndrome. *Curr. Biol.* **30**, R338-R339 (2020).

3   Whittaker, R. J. & Fernández-Palacios, J. M. *Island biogeography: ecology, evolution,*
and conservation. (Oxford University Press, 2007).

4   Lomolino, M. V. Body size of mammals on islands: the island rule reexamined. *Am. Nat.*
125, 310-316 (1985).

5   Lomolino, M. V. Body size evolution in insular vertebrates: generality of the island rule.
J. Biogeogr. **32**, 1683-1699 (2005).

6   Meiri, S., Cooper, N. & Purvis, A. The island rule: made to be broken? *Proc. R. Soc.*
Lond., Ser. B: Biol. Sci. **275**, 141-148 (2008).

7   Meiri, S., Dayan, T. & Simberloff, D. The generality of the island rule reexamined. *J.*
Biogeogr. **33**, 1571-1577 (2006).

8   Meiri, S., Dayan, T. & Simberloff, D. Body size of insular carnivores: little support for
the island rule. *Am. Nat.* **163**, 469-479 (2004).

9   Lomolino, M. V., Sax, D. F., Palombo, M. R. & van der Geer, A. A. Of mice and
mammoths: evaluations of causal explanations for body size evolution in insular
mammals. *J. Biogeogr.* **39**, 842-854 (2012).

10   Millien, V. Relative effects of climate change, isolation and competition on body-size
evolution in the Japanese field mouse, *Apodemus argenteus*. *J. Biogeogr.* **31**, 1267-1276
(2004).

11   Angerbjorn, A. Gigantism in island populations of wood mice (Apodemus) in Europe.
Oikos **47**, 47-56 (1986).

12   Schillaci, M. A., Meijaard, E. & Clark, T. The effect of island area on body size in a
primate species from the Sunda Shelf Islands. *J. Biogeogr.* **36**, 362-371 (2009).

13 Radtkey, R. R., Fallon, S. M. & Case, T. J. Character displacement in some
Cnemidophorus lizards revisited: a phylogenetic analysis. *Proceedings of the National*
Academy of Sciences **94**, 9740-9745 (1997).
14 McClain, C. R., Durst, P. A., Boyer, A. G. & Francis, C. D. Unravelling the determinants
of insular body size shifts. *Biol. Lett.* **9**, 20120989 (2013).
15 Clegg, S. M. & Owens, P. The 'island rule'in birds: medium body size and its ecological
explanation. *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* **269**, 1359-1365 (2002).
16 Raia, P. & Meiri, S. The island rule in large mammals: Paleontology meets ecology.
Evolution **60**, 1731-1742 (2006).
17 Bromham, L. & Cardillo, M. Primates follow the 'island rule': implications for
interpreting *Homo floresiensis. Biol. Lett.* **3**, 398-400 (2007).
18 Montesinos, R., da Silva, H. R. & de Carvalho, A. L. G. The 'island rule'acting on anuran
populations (Bufonidae: *Rhinella ornata*) of the Southern Hemisphere. *Biotropica* **44**,
506-511 (2012).
19 Boback, S. M. Body size evolution in snakes: evidence from island populations. *Copeia*
2003, 81-94 (2003).
20 Meiri, S. Size evolution in island lizards. *Global Ecol. Biogeogr.* **16**, 702-708 (2007).
21 Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C. & Meiri, S. Is the island rule
general? Turtles disagree. *Global Ecol. Biogeogr.* **23**, 689-700 (2014).
22 Boyer, A. G. & Jetz, W. Biogeography of body size in Pacific island birds. *Ecography*
33, 369-379 (2010).
23 Herczeg, G., Gonda, A. & Merilä, J. Evolution of gigantism in nine☐spined sticklebacks.
Evolution **63**, 3190-3200 (2009).
24 Palmer, M. Testing the 'island rule'for a tenebrionid beetle (Coleoptera, Tenebrionidae).
Acta Oecol. **23**, 103-107 (2002).
25 McClain, C. R., Boyer, A. G. & Rosenberg, G. The island rule and the evolution of body
size in the deep sea. *J. Biogeogr.* **33**, 1578-1584 (2006).
26 Biddick, M., Hendriks, A. & Burns, K. Plants obey (and disobey) the island rule. *Proc.*
Natl. Acad. Sci. USA **116**, 17632-17634 (2019).
27 Lokatis, S. & Jeschke, J. M. The island rule: An assessment of biases and research trends.
Journal of Biogeography **45**, 289-303 (2018).
28 Gurevitch, J., Koricheva, J., Nakagawa, S. & Stewart, G. Meta-analysis and the science
of research synthesis. *Nature* **555**, 175 (2018).
29 Nakagawa, S. & Santos, E. S. Methodological issues and advances in biological meta-
analysis. *Evol. Ecol.* **26**, 1253-1274 (2012).
30 Hedges, L. V., Gurevitch, J. & Curtis, P. S. The meta☐analysis of response ratios in
experimental ecology. *Ecology* **80**, 1150-1156 (1999).
31 Atchley, W. R., Gaskins, C. T. & Anderson, D. Statistical properties of ratios. I.
Empirical results. *Syst. Zool.* **25**, 137-148 (1976).
32 Prairie, Y. T. & Bird, D. F. Some misconceptions about the spurious correlation problem
in the ecological literature. *Oecologia* **81**, 285-288 (1989).
33 Phillips, L. B., Hansen, A. J. & Flather, C. H. Evaluating the species energy relationship
with the newest measures of ecosystem energy: NDVI versus MODIS primary
production. *Remote Sens. Environ.* **112**, 3538-3549 (2008).
34 Olesen, J. M. & Valido, A. Lizards as pollinators and seed dispersers: an island
phenomenon. *Trends Ecol. Evol.* **18**, 177-181 (2003).

617	35	Zamora☐Camacho, F., Reguera, S. & Moreno☐Rueda, G. Bergmann's Rule rules body
618		size in an ectotherm: heat conservation in a lizard along a 2200☐metre elevational
619		gradient. *J. Evol. Biol.* **27**, 2820-2828 (2014).
620	36	Valenzuela-Sánchez, A., Cunningham, A. A. & Soto-Azat, C. Geographic body size
621		variation in ectotherms: effects of seasonality on an anuran from the southern temperate
622		forest. *Front. Zool.* **12**, 37 (2015).
623	37	Ashton, K. G. Do amphibians follow Bergmann's rule? *Can. J. Zool./Rev. Can. Zool.* **80**,
624		708-716 (2002).
625	38	Lajeunesse, M. J. Bias and correction for the log response ratio in ecological
626		meta☐analysis. *Ecology* **96**, 2056-2063 (2015).
627	39	Lomolino, M. V. *et al.* Of mice and mammoths: generality and antiquity of the island
628		rule. *J. Biogeogr.* **40**, 1427-1439 (2013).
629	40	Boback, S. M. & Guyer, C. Empirical evidence for an optimal body size in snakes.
630		*Evolution* **57**, 345-451 (2003).
631	41	Meiri, S., Raia, P. & Phillimore, A. B. Slaying dragons: limited evidence for unusual
632		body size evolution on islands. *J. Biogeogr.* **38**, 89-100 (2011).
633	42	Brown, J. H., Marquet, P. A. & Taper, M. L. Evolution of body size: consequences of an
634		energetic definition of fitness. *Am. Nat.* **142**, 573-584 (1993).
635	43	Damuth, J. Cope's rule, the island rule and the scaling of mammalian population density.
636		*Nature* **365**, 748-750 (1993).
637	44	Raia, P., Carotenuto, F. & Meiri, S. One size does not fit all: no evidence for an optimal
638		body size on islands. *Global Ecol. Biogeogr.* **19**, 475-484 (2010).
639	45	Maurer, B. The evolution of body size in birds. II. The role of reproductive power. *Evol.*
640		*Ecol.* **12**, 935-944 (1998).
641	46	Blackburn, T. M. & Gaston, K. J. The distribution of body sizes of the world's bird
642		species. *Oikos*, 127-130 (1994).
643	47	Feldman, A., Sabath, N., Pyron, R. A., Mayrose, I. & Meiri, S. Body sizes and
644		diversification rates of lizards, snakes, amphisbaenians and the tuatara. *Global Ecol.*
645		*Biogeogr.* **25**, 187-197 (2016).
646	48	McNab, B. K. Minimizing energy expenditure facilitates vertebrate persistence on
647		oceanic islands. *Ecol. Lett.* **5**, 693-704 (2002).
648	49	McNab, B. K. Geographic and temporal correlations of mammalian size reconsidered: a
649		resource rule. *Oecologia* **164**, 13-23 (2010).
650	50	Durst, P. A. & Roth, V. L. Mainland size variation informs predictive models of
651		exceptional insular body size change in rodents. *Proceedings of the Royal Society B:*
652		*Biological Sciences* **282**, 20150239 (2015).
653	51	Tobias, J. A., Ottenburghs, J. & Pigot, A. L. Avian Diversity: Speciation,
654		Macroevolution, and Ecological Function. *Annu. Rev. Ecol., Evol. Syst.* **51**, 533-560
655		(2020).
656	52	Li, J.-W. *et al.* Rejecting strictly allopatric speciation on a continental island: prolonged
657		postdivergence gene flow between Taiwan (*Leucodioptron taewanus*, Passeriformes
658		Timaliidae) and Chinese (*L. canorum canorum)* hwameis. *Mol. Ecol.* **19**, 494-507 (2010).
659	53	Somveille, M., Rodrigues, A. S. & Manica, A. Why do birds migrate? A macroecological
660		perspective. *Global Ecol. Biogeogr.* **24**, 664-674 (2015).
661	54	Blondel, J. Evolution and ecology of birds on islands: trends and prospects. *Vie et milieu*
662		*(1980)* **50**, 205-220 (2000).

55 Grant, P. R. Patterns on islands and microevolution. *Evolution on islands* (1998).

56 Novosolov, M. *et al.* Power in numbers. Drivers of high population density in insular
lizards. *Global Ecol. Biogeogr.* **25**, 87-95 (2016).

57 Santini, L. *et al.* Global drivers of population density in terrestrial vertebrates. *Global*
Ecol. Biogeogr. **27**, 968-979 (2018).

58 Castellano, S. & Giacoma, C. Morphological variation of the green toad, *Bufo viridis*, in
Italy: a test of causation. *J. Herpetol.*, 540-550 (1998).

59 Ashton, K. G. Body size variation among mainland populations of the western rattlesnake
(*Crotalus viridis*). *Evolution* **55**, 2523-2533 (2001).

60 Athanassiou, A., van der Geer, A. A. & Lyras, G. A. Pleistocene insular Proboscidea of
the Eastern Mediterranean: a review and update. *Quat. Sci. Rev.* **218**, 306-321 (2019).

61 Herridge, V. L. & Lister, A. M. Extreme insular dwarfism evolved in a mammoth. *Proc.*
R. Soc. Lond., Ser. B: Biol. Sci. **279**, 3193-3200 (2012).

62 Faurby, S. & Svenning, J.-C. Resurrection of the island rule: human-driven extinctions
have obscured a basic evolutionary pattern. *Am. Nat.* **187**, 812-820 (2016).

63 Steadman, D. W. *Extinction and biogeography of tropical Pacific birds*. (University of
Chicago Press, 2006).

64 MacPhee, R. D. & Horovitz, I. New craniodental remains of the Quaternary Jamaican
monkey *Xenothrix mcgregori* (Xenotrichini, Callicebinae, Pitheciidae), with a
reconsideration of the Aotus hypothesis. *Am. Mus. Novit.* **2004**, 1-51 (2004).

65 MacPhee, R. & Fleagle, J. Postcranial remains of Xenothrix mcgregori (Primates,
Xenotrichidae) and other late Quaternary mammals from long Mile Cave, Jamaica. *Bull.*
Am. Mus. Nat. Hist. N. Y. **206**, 287-321 (1991).

66 Pregill, G. Body size of insular lizards: a pattern of Holocene dwarfism. *Evolution* **40**,
997-1008 (1986).

67 Hansen, D. M., Donlan, C. J., Griffiths, C. J. & Campbell, K. J. Ecological history and
latent conservation potential: large and giant tortoises as a model for taxon substitutions.
Ecography **33**, 272-284 (2010).

68 Boyer, A. G. Extinction patterns in the avifauna of the Hawaiian islands. *Divers. Distrib.*
14, 509-517 (2008).

69 White, A. W., Worthy, T. H., Hawkins, S., Bedford, S. & Spriggs, M. Megafaunal
meiolaniid horned turtles survived until early human settlement in Vanuatu, Southwest
Pacific. *Proc. Natl. Acad. Sci. USA* **107**, 15512-15516 (2010).

70 Alcover, J. A., Sans, A. & Palmer, M. The extent of extinctions of mammals on islands.
J. Biogeogr. **25**, 913-918 (1998).

71 Steadman, D. W. Prehistoric extinctions of Pacific island birds: biodiversity meets
zooarchaeology. *Science* **267**, 1123-1131 (1995).

72 Pérez-Méndez, N., Jordano, P., García, C. & Valido, A. The signatures of Anthropocene
defaunation: cascading effects of the seed dispersal collapse. *Sci. Rep.* **6**, 24820 (2016).

73 Heinen, J. H., van Loon, E. E., Hansen, D. M. & Kissling, W. D. Extinction‐driven
changes in frugivore communities on oceanic islands. *Ecography* **41**, 1245-1255 (2018).

74 van der Geer, A. A. E. Changing invaders: trends of gigantism in insular introduced rats.
Environ. Conserv. **45**, 203-211 (2018).

75 van der Geer, A. A. E., Lomolino, M. V. & Lyras, G. 'On being the right size' - Do aliens
follow the rules? *J. Biogeogr.* **45**, 515-529 (2018).

76 Mathys, B. A. & Lockwood, J. L. Rapid evolution of great kiskadees on Bermuda: an
assessment of the ability of the island rule to predict the direction of contemporary
evolution in exotic vertebrates. *J. Biogeogr.* **36**, 2204-2211 (2009).

77 Quinn, J. F. & Dunham, A. E. On hypothesis testing in ecology and evolution. *Am. Nat.*
122, 602-617 (1983).

78 McGill, B. J. & Nekola, J. C. Mechanisms in macroecology: AWOL or purloined letter?
Towards a pragmatic view of mechanism. *Oikos* **119**, 591-603 (2010).

79 Pigot, A. L. *et al.* Macroevolutionary convergence connects morphological form to
ecological function in birds. *Nat. Ecol. Evol.* **4**, 230-239 (2020).

80 Krause, M. A., Burghardt, G. M. & Gillingham, J. C. Body size plasticity and local
variation of relative head and body size sexual dimorphism in garter snakes (*Thamnophis*
sirtalis). *J. Zool.* **261**, 399-407 (2003).

81 Krystufek, B., Tvrtkovic, N., Paunovic, M. & Ozkan, B. Size variation in the Northern
white-breasted hedgehog Erinaceus roumanicus: latitudinal cline and the island rule.
Mammalia **73**, 299-306 (2009).

82 Kubo, M. O. & Takatsuki, S. Geographical Body Size Clines in Sika Deer: Path Analysis
to Discern Amongst Environmental Influences. *Evol. Biol.* **42**, 115-127 (2015).

83 Kuchling, G., Rhodin, A. G., Ibarrondo, B. R. & Trainor, C. R. A new subspecies of the
snakeneck turtle *Chelodina mccordi* from Timor-Leste (East Timor)(Testudines:
Chelidae). *Chelonian Conserv. Biol.* **6**, 213-222 (2007).

84 Kuo, C.-Y., Lin, Y.-T. & Lin, Y.-S. Sexual size and shape dimorphism in an agamid
lizard, *Japalura swinhonis* (Squamata: Lacertilia: Agamidae). *Zool. Stud.* **48**, 351-361
(2009).

85 Kurta, A. & Ferkin, M. The correlation between demography and metabolic rate: a test
using the beach vole (Microtus breweri) and the meadow vole (Microtus pennsylvanicus).
Oecologia **87**, 102-105 (1991).

86 Kutrup, B., Cakir, E., Colak, Z., Bulbul, U. & Karaoglu, H. Age and growth of the green
toad, *Bufo viridis* (Laurenti, 1768) from an island and a mainland population in Giresun,
Turkey. *J. Anim. Vet. Adv.* **10**, 1469-1472 (2011).

87 Kwet, A., Steiner, J. & Zillikens, A. A new species of Adenomera (Amphibia: Anura:
Leptodactylidae) from the Atlantic rain forest in Santa Catarina, southern Brazil. *Stud.*
Neotrop. Fauna Environ. **44**, 93-107 (2009).

88 Lampert, K. P., Bernal, X. E., Rand, A. S., Mueller, U. G. & Ryan, M. J. Island
populations of *Physalaemus pustulosus*: History influences genetic diversity and
morphology. *Herpetologica* **63**, 311-319 (2007).

89 Lawlor, T. E. The evolution of body size in mammals: evidence from insular populations
in Mexico. *Am. Nat.* **119**, 54-72 (1982).

90 Lee, D. E. *et al.* Growth, Age at Maturity, and Age-Specific Survival of the Arboreal
Salamander (Aneides lugubris) on Southeast Farallon Island, California. *J. Herpetol.* **46**,
64-71 (2012).

91 Li, Y. *et al.* Reduced predator species richness drives the body gigantism of a frog
species on the Zhoushan Archipelago in China. *J. Anim. Ecol.* **80**, 171-182 (2011).

92 Libois, R. M. & Fons, R. Le mulot des Iles d'Hyeres: un cas de gigantisme insulaire. *Vie*
Milieu **40**, 217-222 (1990).

93 Lichtenbelt, W. D. V. & Albers, K. B. Reproductive adaptations of the green iguana on a
semiarid island. *Copeia*, 790-798 (1993).

94　Lim, B. The distribution, food habits and parasite patterns of the leopard cat
(*Prionailurus bengalensis*) in Peninsular Malaysia. *The Journal Wildlife and Parks* **17**
(1999).

95　Lin, L.-H., Mao, F., Chen, C. & Ji, X. Reproductive traits of the gray ratsnake *Ptyas*
korros from three geographically distinct populations. *Current Zoology* **58**, 820-827
(2012).

96　Lindell, L. E., Forsman, A. & Merila, J. Variation in number of ventral scales in snakes:
effects on body size, growth rate and survival in the adder, *Vipera berus*. *J. Zool.* **230**,
101-115 (1993).

97　Ljubisavljević, K., Džukić, G., Vukov, T. & Kalezić, M. Morphological variability of the
Hermann's tortoise (*Testudo hermanni*) in the Central Balkans. *Acta Herpetologica* **7**,
253-262 (2012).

98　Lo Cascio, P. & Corti, C. Indagini sull'ecologia dei rettili sauri della RNO e del SIC
"Isola di Lampedusa". *Naturalista sicil* **32**, 319-354 (2008).

99　Lo Cascio, P. & Pasta, S. Preliminary data on the biometry and the diet of a micro-insular
population of *Podarcis wagleriana* (Reptilia: Lacertidae). *Acta Herpetologica* **1**, 147-152
(2006).

100　Lo Valvo, M. & Giacalone, G. Biometrical analyses of a Sicilian green toad, *Bufo siculus*
(Stõck et al. 2008), population living in sicily *Int. J. Morphol.* **31**, 681-686 (2013).

101　Long, E. S., Courtney, K. L., Lippert, J. C. & Wall-Scheffler, C. M. Reduced body size of
insular black-tailed deer is caused by slowed development. *Oecologia* **189**, 675-685
(2019).

102　López-Martín, J., Ruiz-Olmo, J. & Padró, I. Comparison of skull measurements and
sexual dimorphism between the Minorcan pine marten (*Martes martes minoricensis*) and
the Iberian pine marten (*M. m. martes*): A case of insularity. *Mamm. Biol.* **71**, 13-24
(2006).

103　Lötters, S. *et al.* Bio-sketches and partitioning of sympatric reed frogs, genus Hyperolius
(Amphibia; Hyperoliidae), in two humid tropical African forest regions. *J. Nat. Hist.* **38**,
1969-1997 (2004).

104　Luiselli, L., Filippi, E. & Capula, M. Geographic variation in diet composition of the
grass snake (*Natrix natrix*) along the mainland and an island of Italy: the effects of
habitat type and interference with potential competitors. *Herpetological Journal* **15**, 221-
230 (2005).

105　Luiselli, L., Petrozzi, F., Mebert, K., Zuffi, M. A. L. & Amori, G. Resource partitioning
and dwarfism patterns between sympatric snakes in a micro-insular Mediterranean
environment. *Ecol. Res.* **30**, 527-535 (2015).

106　Luz, A. C., Vicente, L. & Monasterio, C. in *Mainland and insular lizards. A*
Mediterranean perspective (eds C. Corti, P. Lo Cascio, & Biaggini M) 111-123
(Firenze University Press, 2006).

107　Lymberakis, P., Poulakakis, N., Kaliontzopoulou, A., Valakos, E. & Mylonas, M. Two
new species of Podarcis (Squamata; Lacertidae) from Greece. *Syst. Biodivers.* **6**, 307-318
(2008).

108　Lynch, J., Conroy, J., Kitchener, A., Jefferies, D. & Hayden, T. Variation in cranial form
and sexual dimorphism among five European populations of the otter *Lutra lutra*. *J. Zool.*
238, 81-96 (1996).

| 799 | 109 | Lyon Jr, M. W. Mammals collected in eastern Sumatra by Dr. WL Abbott during 1903, 1906, and 1907, with descriptions of new species and subspecies. *Proc. U. S. Natl. Mus.* (1908). |

109 Lyon Jr, M. W. Mammals collected in eastern Sumatra by Dr. WL Abbott during 1903,
1906, and 1907, with descriptions of new species and subspecies. *Proc. U. S. Natl. Mus.*
(1908).
110 Lyon, M. W. Mammals of Banka, Mendanau, and Billiton Islands, between Sumatra and
Borneo. *Proc. U. S. Natl. Mus.* **31**, 575–612 (1906).
111 Lyon, M. W. Mammals collected by Dr. WL Abbott on Borneo and some of the small
adjacent islands. *Proc. U. S. Natl. Mus.* **40**, 53-146 (1911).
112 Lyon, M. W. Tree shrews: an account of the mammalian family Tupaiidae. *Proceedings*
of the U. S. National Museum **45**, 1-188 (1913).
113 Lyon, M. W. Mammals collected by Dr. WL Abbott on the chain of islands lying off the
western coast of Sumatra, with descriptions of twenty-eight new species and subspecies.
Proc. U. S. Natl. Mus. **52**, 437–462 (1916).
114 Maddock, S. *Systematics and phylogeography of Seychelles amphibians.* PhD thesis,
UCL (University College London), (2016).
115 Madeira, B. d. C. M. A. *Sexual dimorphism and reproductive phenology of common birds*
in São Tomé island: conservation implications PhD thesis, Universidade de Lisboa,
(2018).
116 Madsen, T. & Shine, R. Phenotypic plasticity in body sizes and sexual size dimorphism
in European grass snakes. *Evolution* **47**, 321-325 (1993).
117 Mageski, M. *et al.* The island rule in the Brazilian frog *Phyllodytes luteolus* (Anura:
Hylidae): incipient gigantism? *Zoologia* **32**, 329-333 (2015).
118 Magnanou, E., Fons, R., Blondel, J. & Morand, S. Energy expenditure in Crocidurinae
shrews (Insectivora): Is metabolism a key component of the insular syndrome? *Comp.*
Biochem. Physiol., A: Mol. Integr. Physiol. **142**, 276-285 (2005).
119 Maharadatunkamsi, H. S., Kitchener, D. & Schmitt, L. Relationships between
morphology, genetics and geography in the cave fruit bat *Eonycteris spelaea* (Dobson,
1871) from Indonesia. *Biol. J. Linn. Soc.* **79**, 511-522 (2003).
120 Major, R. E. Latitudinal and insular variation in morphology of a small Australian
passerine: consequences for dispersal distance and conservation. *Aust. J. Zool.* **60**, 210-
218 (2012).
121 Mallick, S., Driessen, M. & Hocking, G. Biology of the southern brown bandicoot
(*Isoodon obesulus*) in south-eastern Tasmania. II. Demography. *Aust. Mammal.* **20**, 339-
347 (1998).
122 Malmquist, M. G. Character displacement and biogeography of the pygmy shrew in
northern Europe. *Ecology* **66**, 372-377 (1985).
123 Manier, M. K. Geographic variation in the long-nosed snake, *Rhinocheilus lecontei*
(Colubridae): beyond the subspecies debate. *Biol. J. Linn. Soc.* **83**, 65-85 (2004).
124 Manríquez-Morán, N. L., Cruz, M. V.-S. & Mendez-De La Cruz, F. R. Reproductive
biology of the parthenogenetic lizard, *Aspidoscelis cozumela. Herpetologica* **61**, 435-439
(2005).
125 Marinelli, L. & Millar, J. S. The ecology of beach-dwelling *Peromyscus maniculatus* on
the Pacific Coast. *Canadian Journal of Zoology* **67**, 412-417 (1989).
126 Martins, M., Araujo, M. S., Sawaya, R. J. & Nunes, R. Diversity and evolution of
macrohabitat use, body size and morphology in a monophyletic group of Neotropical
pitvipers (Bothrops). *J. Zool.* **254**, 529-538 (2001).

127 Martins, M., Arnaud, G. & Avila-Villegas, H. Juvenile recruitment, early growth, and
morphological variation in the endangered Santa Catalina island rattlesnake, *Crotalus*
catalinensis. *Herpetol. Conserv. Biol.* **7**, 376-382 (2012).
128 Marunouchi, J., Ueda, H. & Ochi, O. Variation in age and size among breeding
populations at different altitudes in the Japanese newts, *Cynops pyrrhogaster*. *Amphibia-*
Reptilia **21**, 381-396 (2000).
129 Matsui, M. & Ota, H. Parameters of fecundity in *Microhyla ornata* from the Yaeyama
Group of the Ryukyu Archipelago. *Japanese Journal of Herpetology* **10**, 73-79 (1984).
130 Matsui, M., Shimada, T. & Sudin, A. First record of the tree frog genus Chiromantis from
Borneo with the description of a new species (Amphibia: Rhacophoridae). *Zool. Sci.* **31**,
45-51 (2014).
131 Matsui, M., Toda, M. & Ota, H. A new species of frog allied to Fejervarya limnocharis
from the Southern Ryukyus, Japan (Amphibia: Ranidae). *Curr. Herp.* **26**, 65-79 (2007).
132 Mazák, J. H. & Groves, C. P. A taxonomic revision of the tigers (Panthera tigris) of
Southeast Asia. *Mamm. Biol.* **71**, 268-287 (2006).
133 McCord, W. P. & Iverson, J. B. A new species of Ocadia (Testudines: Batagurinae) from
southwestern China. *Proc. Biol. Soc. Wash.* **107**, 52-59 (1994).
134 McCord, W. P. & Thomson, S. A. A new species of Chelodina (Testudines: Pleurodira:
Chelidae) from northern Australia. *J. Herpetol.* **36**, 255-267 (2002).
135 McFadden, K. W. & Meiri, S. Dwarfism in insular carnivores: a case study of the pygmy
raccoon. *J. Zool.* **289**, 213-221 (2013).
136 McLaughlin, J. F. & Roughgarden, J. Avian predation on Anolis lizards in the
northeastern caribbean: Inter□island contrast. *Ecology* **70**, 617-628 (1989).
137 Medway, L. Observations on the fauna of Pulau Tioman and Pulau Tulai. 2. The
mammals *Natl. Mus. Singap. Bull.* **34**, 9-32 (1966).
138 Mees, G. F. *A systematic review of the Indo-Australian Zosteropidae (Part III).* (EJ Brill,
1969).
139 Meijaard, E. & Groves, C. A taxonomic revision of the Tragulus mouse-deer
(Artiodactyla). *Zool. J. Linn. Soc.* **140**, 63-102 (2004).
140 Meijaard, E. & Groves, C. P. Morphometrical relationships between South-east Asian
deer (Cervidae, tribe Cervini): evolutionary and biogeographic implications. *J. Zool.* **263**,
179-196 (2004).
141 Meik, J. M., Lawing, A. M. & Pires-daSilva, A. Body size evolution in insular speckled
rattlesnakes (Viperidae: *Crotalus mitchellii*). *Plos One* **5** (2010).
142 Melo, M. *et al.* Multiple lines of evidence support the recognition of a very rare bird
species: the Príncipe thrush. *J. Zool.* **282**, 120-129 (2010).
143 Melo, M. & O'Ryan, C. Genetic differentiation between Príncipe Island and mainland
populations of the grey parrot (*Psittacus erithacus*), and implications for conservation.
Mol. Ecol. **16**, 1673-1685 (2007).
144 Melton, R. Body size and island Peromyscus: a pattern and a hypothesis. *Evolutionary*
Theory **6**, 113-126 (1982).
145 Menzies, J. & Tyler, M. The systematics and adaptations of some Papuan microhylid
frogs which live underground. *J. Zool.* **183**, 431-464 (1977).
146 Merriam, C. H. in *North American Fauna Nº 10*   5-34 (Government Printing Office,
1895).

147 Merriam, C. H. Synopsis of the American shrews of the genus Sorex. *North American*
Fauna **10**, 57-125 (1895).
148 Merriam, C. H. Six new mammals from Cozumel Island, Yucatan. *Proc. Biol. Soc. Wash.*
14, 99-104 (1901).
149 Merriam, C. H. Descriptions of ten new kangaroo rats. *Proc. Biol. Soc. Wash.* **20**, 75-80
(1907).
150 Miller, C. & Miller, T. Population dynamics and diet of rodents on Rangitoto Island, New
Zealand, including the effect of a 1080 poison operation. *N. Z. J. Ecol.*, 19-27 (1995).
151 Miller, G. S. Mammals collected by Dr. WL Abbott in the region of the Indragiri River,
Sumatra. *Proc. Acad. Nat. Sci. Phila.* **54**, 143–159 (1902).
152 Miller, G. S. *Catalogue of the Mammals of Western Europe (Europe Exclusive of Russia)*
in the Collection of the British Museum. (Order of the Trustees of the British Museum,
1912).
153 Millien, V. & Damuth, J. Climate change and size evolution in an island rodent species:
New perspectives on the island rule. *Evolution* **58**, 1353-1360 (2004).
154 Molina-Borja, M. & Rodriguez-Dominguez, M. A. Evolution of biometric and life-
history traits in lizards (Gallotia) from the Canary Islands. *J. Zool. Syst. Evol. Res.* **42**, 44-
53 (2004).
155 Monadjem, A., McCleery, R. A. & Collier, B. A. Activity and movement patterns of the
tortoise *Stigmochelys pardalis* in a subtropical savanna. *J. Herpetol.* **47**, 237-242 (2013).
156 Montesinos, R., da Silva, H. R. & Gomes de Carvalho, A. L. The 'island rule'acting on
anuran populations (Bufonidae: *Rhinella ornata*) of the Southern Hemisphere. *Biotropica*
44, 506-511 (2012).
157 Montgomery, C. E., Boback, S. M., Green, S. E., Paulissen, M. A. & Walker, J. M.
Cnemidophorus lemniscatus (Squamata: Teiidae) on Cayo Cochino Pequeño, Honduras:
extent of island occupancy, natural history, and conservation status. *Herpetol. Conserv.*
Biol. **6**, 10-24 (2011).
158 Moratelli, R., Wilson, D. E., Novaes, R. L., Helgen, K. M. & Gutiérrez, E. E. Caribbean
Myotis (Chiroptera, Vespertilionidae), with description of a new species from Trinidad
and Tobago. *J. Mammal.* **98**, 994-1008 (2017).
159 Mori, A. & Hasegawa, M. Early growth of *Elaphe quadrivirgata* from an insular gigantic
population. *Curr. Herp.* **21**, 43-50 (2002).
160 Mori, A., Ikeuchi, I. & Hasegawa, M. Calling activity of an anuran assemblage in a
temporary pond in a dry forest of Madagascar. *Curr. Herp.* **34**, 140-148 (2015).
161 Mori, E. *et al.* Skull shape and Bergmann's rule in mammals: hints from Old World
porcupines. *J. Zool.* **308**, 47-55 (2019).
162 Motokawa, M. Geographic variation in the Japanese white-toothed shrew *Crocidura*
dsinezumi. *Acta Theriologica* **48**, 145-156 (2003).
163 Motokawa, M. & Lin, L.-K. Geographic variation in the mole-shrew *Anourosorex*
squamipes. *Mammal Study* **27**, 113-120 (2002).
164 Motokawa, M., Lin, L.-K., Harada, M. & Hattori, S. Morphometric geographic variation
in the Asian lesser white-toothed shrew *Crocidura shantungensis* (Mammalia,
Insectivora) in East Asia. *Zool. Sci.* **20**, 789-795 (2003).
165 Munks, S. The breeding biology of *Pseudocheirus peregrinus viverrinus* on Flinders
Island, Bass Strait. *Wildl. Res.* **22**, 521-533 (1995).
166 MVZ. (University of California, Berkeley).

167   Myers, S., Brown, G. & Kleindorfer, S. Divergence in New Holland honeyeaters
936   (*Phylidonyris novaehollandiae*): evidence from morphology and feeding behavior. *J.*
937   *Ornithol.* **151**, 287-296 (2010).
938   168   Nagorsen, D. Body weight variation among insular and mainland American martens.
939   *Martens, sables, and fishers biology and conservation. Comstock, Ithaca, NY*, 85-97
940   (1994).
941   169   Nagy, Z. T., Glaw, F. & Vences, M. Systematics of the snake genera Stenophis and
942   Lycodryas from Madagascar and the Comoros. *Zool. Scr.* **39**, 426-435 (2010).
943   170   Nanova, O. & Prôa, M. Cranial features of mainland and Commander Islands (Russia)
944   Arctic foxes (Vulpes lagopus) reflect their diverging foraging strategies. *Polar Res.* **36**, 7
945   (2017).
946   171   Napier, P. H. *Catalogue of Primates in the British Museum (Natural History) and*
947   *Elsewhere in the British Isles. Part III: Family Cercopithecidae, subfamily Colobinae.*,
948   Vol. 3 (British Museum (Natural History), 1985).
949   172   Nelson, E. Descriptions of new birds from the Tres Marias Islands, western Mexico.
950   *Proc. Biol. Soc. Wash.* **12**, 5-11 (1898).
951   173   Nelson, E. W. in *North American Fauna* 1-314 (US Government Printing Office, 1909).
952   174   Newman, D. G. Activity, dispersion, and population densities of Hamilton's frog
953   (*Leiopelma hamiltoni*) on Maud and Stephens Islands, Nez Zealands. *Herpetologica* **46**,
954   319-330 (1990).
955   175   Nguyen, S. T., Oshida, T., Dang, P. H., Bui, H. T. & Motokawa, M. A new species of
956   squirrel (Sciuridae: Callosciurus) from an isolated island off the Indochina Peninsula in
957   southern Vietnam. *J. Mammal.* **99**, 813-825 (2018).
958   176   Nijman, V. Group composition and monandry in grizzled langurs, *Presbytis comata*, on
959   Java. *Folia Primatol.* **88**, 237-254 (2017).
960   177   Nor, S. M. *The mammalian fauna on the islands at the northern tip of Sabah, Borneo*.
961   (Field Museum of Natural History., 1996).
962   178   Norman, F. & Hurley, V. Gonad measurements and other parameters from Chestnut Teal
963   Anas castanea collected in the Gippsland Lakes region, Victoria. *Emu-Austral*
964   *Ornithology* **84**, 52-55 (1984).
965   179   Nowak, R. M. & Walker, E. P. *Walker's Mammals of the World*. Vol. 1 (JHU Press,
966   1999).
967   180   O'Connell, D. P. *et al.* A sympatric pair of undescribed white-eye species (Aves:
968   Zosteropidae: Zosterops) with different origins. *Zool. J. Linn. Soc.* **186**, 701-724 (2019).
969   181   Oates, J. F., Davies, A. G. & Delson, E. in *Colobine Monkeys: Their Ecology, Behaviour*
970   *and Evolution*   (eds A. Glyn Davies & John F. Oates) (Cambridge University Press,
971   1994).
972   182   Oh, H.-S., Yoshinaga, Y., Kaneko, T., Iida, H. & Mori, T. Taxomic re-examination of the
973   *Apodemus agrarius chejuensis*, comparing external and cranial morphological characters
974   among four Asian Apodemus species. *J. Fac. Agr. Kyushu Univ* **47**, 373-386 (2003).
975   183   Ohdachi, S., Abe, H., Oh, H. & Han, S. Morphological relationships among populations
976   in the *Sorex caecutiens/shinto* group (Eulipotyphla, Soricidae) in East Asia, with a
977   description of a new subspecies from Cheju Island, Korea. *Mamm. Biol.* **70**, 345-358
978   (2005).

979	184	Okada, S., Izawa, M. & Ota, H. Growth and reproduction of *Gekko hokouensis* (Reptilia :
980		Squamata) on Okinawajima Island of the Ryukyu Archipelago, Japan. *J. Herpetol.* **36**,
981		473-479 (2002).
982	185	O'Keeffe, J., O'Boyle, I. & Fogarty, U. Observations on the pathology of tuberculosis in
983		badgers. Report No. 1898473838, (University College Dublin. Centre for Veterinary
984		Epidemiology and Risk Analysis, Dublin, 1997).
985	186	Olsen, P., Debus, S., Czechura, G. & Mooney, N. Comparative feeding ecology of the
986		grey goshawk *Accipiter novaehollandiae* and brown goshawk *Accipiter fasciatus*. *Aust.*
987		*Bird Watch.* **13**, 178-192 (1990).
988	187	Oneto, F., Ottonello, D. & Salvidio, S. Primi dati sulla biometria di *Euleptes euopaea*
989		(Genè, 1839) dell'isola del Tino (La Spezia, Liguria). *Doriana, Ann. Mus. Civ. Stor. nat*
990		*G. Doria, Genova* **8**, 1-8 (2008).
991	188	Oromi, N. *et al.* Geographical variations in adult body size and reproductive life history
992		traits in an invasive anuran, *Discoglossus pictus*. *Zoology* **119**, 216-223 (2016).
993	189	Osgood, W. H. Natural history of the Queen Charlotte islands. *North American Fauna*, 1-
994		87 (1901).
995	190	Osgood, W. H. Revision of the mice of the American genus *Peromyscus*. *North American*
996		*Fauna* **28**, 1-285 (1909).
997	191	Ota, H. Taxonomic redefinition of *Japalura swinhonis* Günther (Agamidae: Squamata),
998		with a description of a new subspecies of *J. polygonata* from Taiwan. *Herpetologica*,
999		280-294 (1991).
1000	192	Ota, H., Lau, M. W., Weidenhofer, T., Yasukawa, Y. & Bogadek, A. Taxonomic review
1001		of the geckos allied to *Gekko chinensis* Gray 1842 (Gekkonidae Reptilia) from China and
1002		Vietnam. *Trop. Zool.* **8**, 181-196 (1995).
1003	193	Owens, J. R. *Ecology and Behavior of the Bioko Island Drill (Mandrillus leucophaeus*
1004		*poensis).* PhD thesis, Drexel University, (2013).
1005	194	Pabijan, M., Gehring, P.-S., Koehler, J., Glaw, F. & Vences, M. A new microendemic
1006		frog species of the genus Blommersia (Anura: Mantellidae) from the east coast of
1007		Madagascar. *Zootaxa* **2978**, 34-50 (2011).
1008	195	Pafilis, P. *et al.* Reproductive biology of insular reptiles: marine subsidies modulate
1009		expression of the "island syndrome". *Copeia*, 545-552 (2011).
1010	196	Pafilis, P., Kapsalas, G., Lymberakis, P., Protopappas, D. & Sotiropoulos, K. Diet
1011		composition of the Karpathos marsh frog (*Pelophylax cerigensis*): what does the most
1012		endangered frog in Europe eat? *Anim. Biodivers. Conserv.* **42**, 1-8 (2019).
1013	197	Pafilis, P., Meiri, S., Foufopoulos, J. & Valakos, E. Intraspecific competition and high
1014		food availability are associated with insular gigantism in a lizard. *Naturwissenschaften*
1015		**96**, 1107-1113 (2009).
1016	198	Pafilis, P. *et al.* Body size affects digestive performance in a Mediterranean lizard.
1017		*Herpetological Journal* **26**, 199-205 (2016).
1018	199	Pagh, S. *et al.* Methods for the identification of farm escapees in feral mink (*Neovison*
1019		*vison*) populations. *PloS One* **14** (2019).
1020	200	Pahl, L. & Lee, A. Reproductive traits of 2 populations of the common ringtail possum,
1021		*Pseudocheirus peregrinus*, in Victoria. *Aust. J. Zool.* **36**, 83-97 (1988).
1022	201	Palacios, F. & Fernández, J. A new subspecies of hare from Majorca (Balearic Islands).
1023		*Mammalia* **56**, 71-86 (1992).

202 Papakosta, M. A. Biometric variation in *Martes foina* from mainland Greece and the
Aegean Islands. *Turkish Journal of Zoology* **41**, 654-663 (2017).
203 Parker, W. S. & Pianka, E. R. Comparative ecology of populations of the lizard *Uta*
stansburiana. *Copeia*, 615-632 (1975).
204 Parrish, G. & Gill, B. Natural history of the lizards of the Three Kings Islands, New
Zealand. *N. Z. J. Zool.* **30**, 205-220 (2003).
205 Pasachnik, S. A. *et al.* Body size, demography, and body condition in Utila spiny-tailed
iguanas, *Ctenosaura bakeri*. *Herpetol. Conserv. Biol.* **7**, 391-398 (2012).
206 Pearson, D. *et al.* Ecological notes on crowned snakes *Elapognathus coronatus* from the
Archipelago of the Recherche in southwestern Australia. *Aust. Zool.* **31**, 610-617 (2001).
207 Pedrono, M. & Markwell, T. Maximum size and mass of the ploughshare tortoise,
Geochelone yniphora. *Chelonian Conserv. Biol.* **4**, 190-190 (2001).
208 Perez-Ramos, E. *et al.* *Aspidoscelis sexlineata* (Sauria: Teiidae) in Mexico: distribution,
habitat, morphology, and taxonomy. *Southwest. Nat.* **55**, 419-425 (2010).
209 Persson, S. *et al.* Influence of age, season, body condition and geographical area on
concentrations of chlorinated and brominated contaminants in wild mink (*Neovison*
vison) in Sweden. *Chemosphere* **90**, 1664-1671 (2013).
210 Peters, J. A. The snakes of the subfamily Dipsadinae. *Miscellaneous Publications of the*
Museum of Zoology, University of Michigan **114** (1960).
211 Pocock, R. The civet cats of Asia. Part II. . *J. Bombay Nat. Hist. Soc.* **36**, 629-656 (1933).
212 Pocock, R. in *Proceedings of the Zoological Society of London.* 613-684 (Wiley Online
Library).
213 Pocock, R. I. *The Fauna Of British India Including Ceylon And Burma.* (Taylor and
Francis Ltd, 1939).
214 Pons, J. M., Kirwan, G. M., Porter, R. F. & Fuchs, J. A reappraisal of the systematic
affinities of Socotran, Arabian and East African scops owls (Otus, Strigidae) using a
combination of molecular, biometric and acoustic data. *Ibis* **155**, 518-533 (2013).
215 Abe, H. Habitat factors affecting the geographic size variation of Japanese moles.
Mammal Study **21**, 71-87 (1996).
216 Abramov, A. V., Jenkins, P. D., Rozhnov, V. V. & Kalinin, A. A. Description of a new
species of Crocidura (Soricomorpha: Soricidae) from the island of Phu Quoc, Vietnam.
Mammalia **72**, 269-272 (2008).
217 Abramov, A. V. & Puzachenko, A. Y. Sexual dimorphism of craniological characters in
Eurasian badgers, *Meles spp.*(Carnivora, Mustelidae). *Zoologischer Anzeiger-A Journal*
of Comparative Zoology **244**, 11-29 (2005).
218 Adams, N. E., Dean, M. D. & Pauly, G. B. Morphological divergence among populations
of *Xantusia riversiana*, a night lizard endemic to the Channel Islands of California.
Copeia **106**, 550-562 (2018).
219 Aguilar-Moreno, M. *et al.* Dimorfismo sexual de *Aspidoscelis costata costata* (Squamata:
Teiidae) en el sur del Estado de México, México. *Rev. Chil. Hist. Nat.* **83**, 585-592
(2010).
220 Ajtić, R. Morphological, biogeographical and ecological characteristics of Kotschy's
gecko (*Cyrtodactylus kotschyi* Steindachner, 1870 Gekkonidae) from the mainland
portion of its distribution range. *Fauna Balkana* **3**, 1-70 (2014).
221 Albrecht, G. H., Jenkins, P. D. & Godfrey, L. R. Ecogeographic size variation among the
living and subfossil prosimians of Madagascar. *Am. J. Primatol.* **22**, 1-50 (1990).

1070	222	Allen, G. M. *The Mammals of China and Mongolia*. (American Museum of Natural
1071		History, 1938).
1072	223	Allen, J. A. A preliminary study of the South American opossums of the genus Didelphis.
1073		*Bulletin of the American Museum of Natural History* **16**, 249-279 (1902).
1074	224	Allen, J. A. Mammals collected in Alaska and northern British Columbia by the Andrew
1075		J. Stone expedition of 1902. *Bulletin of the American Museum of Natural History* **19**,
1076		521-567 (1903).
1077	225	Allen, J. A. Notes on American deer of the genus Mazama. *Bulletin of the American*
1078		*Museum of Natural History* **34**, 521-553 (1915).
1079	226	Allen, J. A. Review of the South American Sciuridae. *Bulletin of the American Museum*
1080		*of Natural History* **34**, 147-309 (1915).
1081	227	Allen, J. A. & Chapman, F. M. On a collection of mammals from the island of Trinidad,
1082		with descriptions of new species. *Bulletin of the American Museum of Natural History* **5**,
1083		203-234 (1898).
1084	228	Allen, J. A., Kermode, F. & Andrews, R. C. The white bear of southwestern British
1085		Columbia. *Bulletin of the American Museum of Natural History* **26**, 233-238 (1909).
1086	229	Altunisik, A., Kalayci, T. E., Gul, C., Ozdemir, N. & Tosunoglu, M. A
1087		skeletochronological study of the smooth newt *Lissotriton vulgaris* (Amphibia: Urodela)
1088		from an island and a mainland population in Turkey. *Ital. J. Zool.* **81**, 381-388 (2014).
1089	230	Amor, N. & Farjallah, S. Morphological variation of the African green toad, *Bufo*
1090		*boulengeri* (Amphibia: Anura) in Tunisia. *Pak. J. Zool.* **43** (2011).
1091	231	Anderson, R. P. & Handley, C. O. Dwarfism in insular sloths: Biogeography, selection,
1092		and evolutionary rate. *Evolution* **56**, 1045-1058 (2002).
1093	232	Andrade, P. *et al.* Ecomorphological patterns in the Blackcap *Sylvia atricapilla*: insular
1094		versus mainland populations. *Bird Study* **62**, 498-507 (2015).
1095	233	Andrews, R. M. Evolution of life histories: a comparison of Anolis lizards from matched
1096		island and mainland habitats. *Breviora* **454**, 1-51 (1979).
1097	234	Angarita-Sierra, T. & Lynch, J. D. A new species of Ninia (Serpentes: Dipsadidae) from
1098		Chocó-Magdalena biogeographical province, western Colombia. *Zootaxa* **4244**, 478-492
1099		(2017).
1100	235	Angelici, F., Capizzi, D., Amori, G. & Luiselli, L. Morphometric variation in the skulls
1101		of the crested porcupine *Hystrix cristata* from mainland Italy, Sicily, and northern Africa.
1102		*Mamm. Biol.* **68**, 165-173 (2003).
1103	236	Aowphol, A., Rujirawan, A., Taksintum, W., Arsirapot, S. & Mcleod, D. S. Re-
1104		evaluating the taxonomic status of Chiromantis in Thailand using multiple lines of
1105		evidence (Amphibia: Anura: Rhacophoridae). *Zootaxa* **3702**, 101-123 (2013).
1106	237	Araújo, M. & Martins, M. The defensive strike of five species of lanceheads of the genus
1107		Bothrops (Viperidae). *Braz. J. Biol.* **67**, 327-332 (2007).
1108	238	Arendt, W. J. & Arendt, A. L. Bill deformity in a Pearly-eyed Thrasher from Montserrat,
1109		West Indies. *Auk* **54**, 324-332 (1986).
1110	239	Arntzen, J. & García-París, M. Morphological and allozyme studies of midwife toads
1111		(genus Alytes), including the description of two new taxa from Spain. *Contributions to*
1112		*Zoology* **65**, 5-34 (1995).
1113	240	Aubret, F. Island colonisation and the evolutionary rates of body size in insular neonate
1114		snakes. *Heredity* **115**, 349-356 (2015).

1115   241   Auer, M. & Taskavak, E. Population structure of syntopic *Emys orbicularis* and
1116   *Mauremys rivulata* in western Turkey. *Biologia* **59**, 81-84 (2004).
1117   242   Aumann, T. *Aspects of the biology of the Brown Goshawk Accipiter fasciatus in south-*
1118   *eastem Australia.* MSc thesis, Monash University, (1986).
1119   243   Baier, F. & Hoekstra, H. E. The genetics of morphological and behavioural island traits
1120   in deer mice. *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* **286** (2019).
1121   244   Baker-Gabb, D. J. *Comparative ecology and behaviour of swamp harriers Circus*
1122   *approximans, spotted harriers C. assimilis and other raptors in Australia and New*
1123   *Zealand*, Royal Australasian Ornithological Union, (1982).
1124   245   Bakhuis, W. L. Size and sexual differentiation in the lizard *Iguana iguana* on a semi-arid
1125   island. *J. Herpetol.* **16**, 322-325 (1982).
1126   246   Balham, R. W. Grey and mallard ducks in the Manawatu district, New Zealand. *Emu-*
1127   *Austral Ornithology* **52**, 163-191 (1952).
1128   247   Bangma, J. T. *et al.* Perfluorinated alkyl acids in plasma of American alligators (*Alligator*
1129   *mississippiensis*) from Florida and South Carolina. *Environ. Toxicol. Chem.* **36**, 917-925
1130   (2017).
1131   248   Bangma, J. T. *et al.* Variation in perfluoroalkyl acids in the American alligator (*Alligator*
1132   *mississippiensis*) at Merritt Island National Wildlife Refuge. *Chemosphere* **166**, 72-79
1133   (2017).
1134   249   Banks, R. A new insular subspecies of spiny pocket mouse (Mammalia: Rodentia). *Proc.*
1135   *Biol. Soc. Wash.* **80**, 101-104 (1967).
1136   250   Barbo, F. E. *et al.* Another new and threatened species of lancehead genus Bothrops
1137   (Serpentes, Viperidae) from Ilha dos Franceses, Southeastern Brazil. *Zootaxa* **4097**, 511-
1138   529 (2016).
1139   251   Barbour, T. & Allen, G. M. The white-tailed deer of eastern United States. *J. Mammal.* **3**,
1140   65-78 (1922).
1141   252   Bartle, J. & Sagar, P. Intraspecific variation in the New Zealand bellbird *Anthornis*
1142   *melanura*. *Notornis* **34**, 253-306 (1987).
1143   253   Barwick, R. E. *Studies on the scincid lizard Egernia cunninghami (Gray, 1832).* PhD
1144   thesis, The Australian National University, (1965).
1145   254   Başkale, E., Ulubeli, S. A. & Kaska, Y. Age structures and growth parameters of the
1146   Levantine frog, *Pelophylax bedriagae*, at different localities in Denizli, Turkey. *Acta*
1147   *Herpetologica* **13**, 147-154 (2018).
1148   255   Bee, J. W. & Hall, E. R. *Mammals of northern Alaska on the Arctic Slope*. (Museum of
1149   Natural History, University of Kansas, 1956).
1150   256   Bejakovic, D., Kalezic, M. L., Aleksic, I., Dzukic, G. & CrnobrnjaIsailovic, J. Female
1151   reproductive cycle and clutch traits in the Dalmatian wall lizard (*Podarcis melisellensis*).
1152   *Folia Zoologica* **44**, 371-380 (1995).
1153   257   Bell, B. D. Observations on the ecology and reproduction of the New Zealand
1154   Leiopelmid frogs. *Herpetologica* **34**, 340-354 (1978).
1155   258   Bell, R. C. & Irian, C. G. Phenotypic and genetic divergence in reed frogs across a
1156   mosaic hybrid zone on São Tomé Island. *Biol. J. Linn. Soc.* **128**, 672-680 (2019).
1157   259   Bennett, D. & Hampson, K. Further Observations of Varanus olivaceus on the Polillo
1158   Islands. *Hampson et al, Wildlife and Conservation in the Pollilo Islands. Pollilo project*
1159   *Final Report, Multimedia CD. Viper Press, Glossop* (2003).

260 Benson, S. B. Two new Pocket Mice: genus Perognathus, from the Californias.
University of California Publications in Zoology **32**, 449-454 (1930).
261 Bentz, E. J., Rodríguez, M. R., John, R. R., Henderson, R. W. & Powell, R. Population
densities, activity, microhabitats, and thermal biology of a unique crevice-and litter-
dwelling assemblage of reptiles on Union Island, St. Vincent and the Grenadines.
Herpetol. Conserv. Biol. **6**, 40-50 (2011).
262 Beovides-Casas, K. & Mancina, C. A. Natural history and morphometry of the Cuban
iguana (*Cyclura nubila* Gray, 1831) in Cayo Siju, Cuba. *Anim. Biodivers. Conserv.* **29**, 1-
8 (2006).
263 Berardo, F. *Habitat preferenziali e dinamica di popolazione di Testudo hermanni nelle*
aree costiere del Molise. PhD thesis, Universitá degli studi del Molise, (2015).
264 Berry, R. The evolution of an island population of the house mouse. *Evolution* **18**, 468-
483 (1964).
265 Berry, R., Jakobson, M. & Peters, J. The house mice of the Faroe Islands: a study in
microdifferentiation. *J. Zool.* **185**, 73-92 (1978).
266 Bertolero, A. in *Enciclopedia Virtual de los Vertebrados Españoles.* (ed A. Salvador,
Marco, A.) (Museo Nacional de Ciencias Naturales, 2015).
267 Bertolero, A., Cheylan, M. & Nougarede, J.-P. Accroissement de la fécondité chez la
tortue d'Hermann *Testudo hermanni hermanni* en condition insulaire: Un contre-exemple
du syndrome insulaire? *Revue d'écologie* **62**, 93-98 (2007).
268 Bischoff, W. Bemerkungen zur innerartlichen Variabilität von *Gallotia atlantica* (Peters
& Doria, 1882) (Lacertidae). *Bonn. Zool. Beitr.* **36**, 489-506 (1985).
269 Bishop, C. A. & Rouse, J. D. Polychlorinated biphenyls and organochlorine pesticides in
plasma and the embryonic development in Lake Erie water snakes (*Nerodia sipedon*
insularum) from Pelee Island, Ontario, Canada (1999). *Arch. Environ. Contam. Toxicol.*
51, 452-457 (2006).
270 Blackburn, D. C. A new squeaker frog (Arthroleptidae: Arthroleptis) from Bioko island,
Equatorial Guinea. *Herpetologica* **66**, 320-334 (2010).
271 Blondel, J., Perret, P., Anstett, M. C. & Thebaud, C. Evolution of sexual size dimorphism
in birds: test of hypotheses using blue tits in contrasted Mediterranean habitats. *J. Evol.*
Biol. **15**, 440-450 (2002).
272 Boback, S. M. A morphometric comparison of island and mainland boas (*Boa*
constrictor) in Belize. *Copeia* **2006**, 261-267 (2006).
273 Bosc, V. Inventaire du Phyllodactyle d'Europe *Euleptes europaea* sur l'îlot de Roscana
(Corse du Sud). 1-18 (Conservatoire des Espaces Naturels de Corse, 2009).
274 Boye, P. Notes on the morphology, ecology and geographic origin of the Cyprus long-
eared hedgehog (*Hemiechinus auritus dorotheae*). *Bonn. Zool. Beitr.* **42**, 115-123 (1991).
275 Braithwaite, L. & Miller, B. The mallard, Anas platyrhynchos, and mallard-black duck,
Anas superciliosa rogersi, hybridization. *Wildl. Res.* **2**, 47-61 (1975).
276 Brandon-Jones, D. The taxonomic affinities of the Mentawai Islands Sureli, *Presbytis*
potenziani (Bonaparte, 1856)(Mammalia: Primata: Cercopithecidae). *Raffles Bull. Zool.*
41, 331-357 (1993).
277 Brasileiro, C. A., Haddad, C. F., Sawaya, R. J. & Sazima, I. A new and threatened island-
dwelling species of Cycloramphus (Anura: Cycloramphidae) from southeastern Brazil.
Herpetologica **63**, 501-510 (2007).

278   Brecko, J. *et al.* Functional and ecological relevance of intraspecific variation in body
1206   size and shape in the lizard Podarcis melisellensis (Lacertidae). *Biol. J. Linn. Soc.* **94**,
1207   251-264 (2008).
1208   279   Brisbin Jr, I. L. & Lenarz, M. S. Morphological comparisons of insular and mainland
1209   populations of southeastern white-tailed deer. *J. Mammal.* **65**, 44-50 (1984).
1210   280   Bromham, L. & Cardillo, M. Primates follow the 'island rule': implications for
1211   interpreting *Homo floresiensis*. *Biol. Lett.* **3**, 398-400 (2007).
1212   281   Brown, J. L., Maan, M. E., Cummings, M. E. & Summers, K. Evidence for selection on
1213   coloration in a Panamanian poison frog: a coalescent□based approach. *J. Biogeogr.* **37**,
1214   891-901 (2010).
1215   282   Bruschi, S. *et al.* Comments on the status of the Sardinian-Corsican lacertid lizard
1216   *Podarcis tiliguerta. Proc. Calif. Acad. Sci.* **57**, 225-245 (2006).
1217   283   Buden, D. W. Morphological variation and distributional ecology of the giant
1218   Micronesian gecko (*Perochirus scutellatus*) of Kapingamarangi Atoll. *Pac. Sci.* **52**, 250-
1219   258 (1998).
1220   284   Burt, W. H. Descriptions of heretofore unknown mammals from islands in the Gulf of
1221   California, Mexico. *Transactions of the San Diego Society of Natural History* **16**, 161-
1222   185 (1932).
1223   285   Butynski, T. M., de Jong, Y. A. & Hearn, G. W. Body measurements for the monkeys of
1224   Bioko Island, Equatorial Guinea. *Primate Conservation* **24**, 99-105 (2009).
1225   286   Cabot, J. & Urdiales, C. The subspecific status of Sardinian warblers *Sylvia
1226   melanocephala* in the Canary Islands with the description of a new subspecies from
1227   Western Sahara. *Bull. Br. Ornithol. Club* **125**, 230-240 (2005).
1228   287   Camerano, L. Ricerche intorno alla variazione del *Phyllodactylus europaeux* Gené. *Bull.
1229   Mus Zool. Anat. comp. R. Univ Torino* **19** 1-28 (1904).
1230   288   Camps, D. in *Enciclopedia Virtual de los Vertebrados Españoles*   (ed A. Salvador,
1231   Barja, I.) (Museo Nacional de Ciencias Naturales, 2017).
1232   289   Capula, M. *et al.* Thermal ecology of the aeolian wall lizard, Podarcis raffonei. what
1233   about body temperatures in microinsular lizards. *Scripta Herpetologica. Studies on
1234   Amphibians and Reptiles in honour of Benedetto Lanza. Monografie della Societas
1235   Herpetologica Italica* **3**, 39-47 (2014).
1236   290   Carbone, M. *Caratteristiche della popolazione di Testudo hermanni Gmelin del Parco
1237   Naturale della Maremma.* MSc thesis, Università degli studi di Genova, (1988).
1238   291   Carrascal, L. M., Moreno, E. & Valido, A. Morphological evolution and changes in
1239   foraging behaviour of island and mainland populations of Blue Tit (*Parus caeruleus*)—a
1240   test of convergence and ecomorphological hypotheses. *Evol. Ecol.* **8**, 25-35 (1994).
1241   292   Carretero, M. & Llorente, G. Morphometry in a community of Mediterranean lacertid
1242   lizards, and its ecological relationships. *Hist. Anim.* **2**, 77-99 (1993).
1243   293   Case, T. J., Cody, M. L. & Ezcurra, E. *A new island biogeography of the Sea of Cortés.*
1244   (Oxford University Press on Demand, 2002).
1245   294   Case, T. J. & Schwaner, T. D. Island/mainland body size differences in Australian
1246   varanid lizards. *Oecologia* **94**, 102-109 (1993).
1247   295   Castellano, S., Rosso, A., Doglio, S. & Giacoma, C. Body size and calling variation in
1248   the green toad (*Bufo viridis*). *J. Zool.* **248**, 83-90 (1999).
1249   296   Castilla, A. M. & Bauwens, D. Reproductive characteristics of the lacertid lizard
1250   *Podarcis atrata. Copeia* **2000**, 748-756 (2000).

297 Catzeflis, F., Maddalena, T., Hellwing, S. & Vogel, P. Unexpected findings on the
taxonomic status of East Mediterranean *Crocidura russula* auct. (Mammalia,
Insectivora). *Zeitschrift für Säugetierkunde* **50**, 185-201 (1985).
298 Celis-Diez, J. L. *et al.* Population abundance, natural history, and habitat use by the
arboreal marsupial *Dromiciops gliroides* in rural Chiloé Island, Chile. *J. Mammal.* **93**,
134-148 (2012).
299 Ceríaco, L. M. *et al.* Description of a new endemic species of shrew (Mammalia,
Soricomorpha) from PrÍncipe Island (Gulf of Guinea). *Mammalia* **79**, 325-341 (2015).
300 Chamberlain, J. The Block Island meadow mouse, *Microtus provectus*. *J. Mammal.* **35**,
587-589 (1954).
301 Charles☐Dominique, P. Urine marking and territoriality in *Galago alleni* (Waterhouse,
1837—Lorisoidea, Primates)—a field study by radio☐telemetry. *Z. Tierpsychol.* **43**, 113-
138 (1977).
302 Chasen, F. & Kloss, C. B. On a collection of mammals from the lowlands and islands of
North Borneo. *Bull. Raffles Mus.* **6**, 1-82 (1931).
303 Chen, S.-L. *et al.* Taxonomic status of the Korean populations of the genus Scincella
(Squamata: Scincidae). *J. Herpetol.* **35**, 122-129 (2001).
304 Cherem, J. J., Olimpio, J. & Ximenez, A. Descrição de uma nova espécie do gênero
Cavia Pallas, 1766 (Mammalia-Caviidae) das Ilhas dos Moleques do Sul, Santa Catarina,
Sul do Brasil. *Biotemas* **12**, 95-117 (1999).
305 Chondropoulos, B. P. & Lykakis, J. J. Ecology of the Balkan wall lizard, *Podarcis
taurica ionica* (Sauria: Lacertidae) from Greece. *Copeia*, 991-1001 (1983).
306 Clark, R. J. Herpetofauna of the islands of the Argo-Saronic Gulf, Greece. *Proc. Calif.
Acad. Sci.* **35**, 23-36 (1967).
307 Clegg, S. M. *et al.* Microevolution in island forms: the roles of drift and directional
selection in morphological divergence of a passerine bird. *Evolution* **56**, 2090-2099
(2002).
308 Clover, R. C. Phenetic relationships among populations of *Podarcis sicula* and *P.
melisellensis* (Sauria: Lacertidae) from islands in the Adriatic Sea. *Syst. Biol.* **28**, 284-298
(1979).
309 Cole, C. J., Dessauer, H. C., Townsend, C. R. & Arnold, M. G. Unisexual lizards of the
genus Gymnophthalmus (Reptilia, Teiidae) in the Neotropics: genetics, origin, and
systematics. American Museum novitates; no. 2994. (1990).
310 Colyn, M. Données pondérales sur les primates Cercopithecidae d'Afrique Centrale
(Bassin du Zaïre/Congo). *Mammalia* **58**, 483-488 (1994).
311 Copley, P., Read, V., Robinson, A. & Watts, C. Preliminary studies of the Nuyts
Archipelago bandicoot *Isoodon obesulus nauticus* on the Franklin Islands, South
Australia. *Bandicoots and Bilbies'.(Eds JH Seebeck, PR Brown, RL Wallis and CM
Kemper.) pp*, 345-356 (1990).
312 Cordero Rivera, A., Velo-Antón, G. & Galán, P. Ecology of amphibians in small coastal
Holocene islands: local adaptations and the effect of exotic tree plantations. *Munibe* **25**,
94-103 (2007).
313 Corti, C. & Zuffi, M. Aspects of population ecology of *Testudo hermanni hermanni* from
Asinara Island, NW Sardinia (Italy, Western Mediterranean Sea): preliminary data.
Amphibia-Reptilia **24**, 441-447 (2003).

314   Cowan, I. M. Insularity in the genus Sorex on the north coast of British Columbia. *Proc.*
1297   *Biol. Soc. Wash.* **54**, 95-107 (1941).
1298   315   Cox, D. T. *et al.* Patterns of seasonal and yearly mass variation in West African tropical
1299   savannah birds. *Ibis* **153**, 672-683 (2011).
1300   316   Creer, S., Chou, W. H., Malhotra, A. & Thorpe, R. S. Offshore insular variation in the
1301   diet of the Taiwanese bamboo viper *Trimeresurus stejnegeri* (Schmidt). *Zool. Sci.* **19**,
1302   907-913 (2002).
1303   317   Cruz-Elizalde, R. *et al.* Sexual dimorphism and natural history of the Western Mexico
1304   Whiptail, *Aspidoscelis costata* (Squamata: Teiidae), from Isla Isabel, Nayarit, Mexico.
1305   *North-Western Journal of Zoology* **10**, 374-381 (2014).
1306   318   Cumbo, V., Licata, F., Mercurio, E., Anz, S. & Lo Valvo, M. in *VIII Congresso*
1307   *Nazionale Societas Herpetologica Italica.* 401-404 (Ianieri Edizioni).
1308   319   D'Angelo, S., Galia, F. & Lo Valvo, M. Biometric characterization of two Sicilian pond
1309   turtle (*Emys trinacris*) populations of south-western Sicily. *Revista Española de*
1310   *Herpetología* **22**, 15-22 (2008).
1311   320   Dagosto, M., Gebo, D. L. & Dolino, C. Positional behavior and social organization of the
1312   Philippine tarsier (*Tarsius syrichta*). *Primates* **42**, 233-243 (2001).
1313   321   Darevsky, I. Two new species of the worm-like lizard Dibamus (Sauria, Dibamidae),
1314   with remarks on the distribution and ecology of Dibamus in Vietnam. *Asiatic*
1315   *Herpetological Research* **4**, 1-12 (1992).
1316   322   David, P., Vidal, N. & Pauwels, O. S. A morphological study of Stejneger's pitviper
1317   *Trimeresurus stejnegeri* (Serpentes, Viperidae, Crotalinae), with the description of a new
1318   species from Thailand. *Russian Journal of Herpetology* **8**, 205-222 (2001).
1319   323   De la Cruz, J. O. & Casas Jr, E. V. Captive Observations and Comparative Morphology
1320   of Philippine Tarsier (*Carlito syrichta*) in Brgy. Hugpa, Biliran, Biliran: A Preliminary
1321   Study. *Philippine Journal of Natural Sciences* **20**, 46-54 (2015).
1322   324   Decker, D. Systematics of the coatis, genus Nasua (Mammalia: Procyonidae). *Proc. Biol.*
1323   *Soc. Wash.* **104**, 370-386 (1991).
1324   325   Delany, M. Variation in the long-tailed field-mouse (*Apodemus sylvaticus* (L.)) in north-
1325   west Scotland I. Comparisons of individual characters. *Proceedings of the Royal Society*
1326   *of London. Series B. Biological Sciences* **161**, 191-199 (1964).
1327   326   Delany, M. & Healy, M. Variation in the white-toothed shrews (*Crocidura spp.*) in the
1328   British Isles. *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* **164**, 63-74 (1966).
1329   327   Delany, M. & Healy, M. Variation in the long-tailed field-mouse (*Apodemus sylvaticus*
1330   (L.)) in the Channel Islands. *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* **166**, 408-421 (1967).
1331   328   Delaugerre, M. La variation géographique chez *Phyllodactylus europaeus* Gené (Reptilia,
1332   Sauria, Gekkonidae). Etude de la population de l'îlot Sperduto Grande (Sud de la Corse,
1333   réserve naturelle des Iles Lavezzi). *Publications de la Société Linnéenne de Lyon* **54**,
1334   262-269 (1985).
1335   329   Delaugerre, M. Le Phyllodactyle d'Europe sur l'île de Port-Cros: mise en place d'un suivi
1336   géographique; la population de la Gabinière. 38 (Parc national de Port-Cros, Hyères,
1337   2003).
1338   330   Delaugerre, M. Archipel de la Galite «Notes herpétologiques 2008» Petites îles de
1339   Méditerranée 08. (2009).
1340   331   Delaugerre, M. & Dubois, A. La variation geographique et la variabilite
1341   intrapopulationnelle chez *Phyllodactylus europaeus* (Reptilia, Sauria, Gekkonidae).

Bulletin du Museum National d'Histoire Naturelle Section A Zoologie Biologie et
Ecologie Animales **7**, 709-736 (1985).

332   Delibes de Castro, M. Sobre las Ginetas de la Isla de Ibiza (*Genetta genetta isabelae* n.
ssp.). *Doñana. Acta Vertebrata* **4**, 139-160 (1977).

333   Delibes, M. & Amores, F. The Stone Marten M*artes foina* (Erxleben, 1777)(Mammalia,
Carnivora) from Ibiza (Pitiusic, Balearic Islands). *Miscellania Zoologica (Spain)* (1986).

334   Delson, E., Terranova, C. J., Jungers, W. L., Sargis, E. J. & Jablonski, N. G. Body mass
in Cercopithecidae (Primates, Mammalia): estimation and scaling in extinct and extant
taxa. Anthropological papers of the AMNH; no. 83. (2000).

335   Dietzen, C., Garcia-del-Rey, E., Castro, G. D. & Wink, M. Phylogeography of the blue tit
(*Parus teneriffae*-group) on the Canary Islands based on mitochondrial DNA sequence
data and morphometrics. *J. Ornithol.* **149**, 1-12 (2008).

336   Djong, H. T., Matsui, M., Kuramoto, M., Nishioka, M. & Sumida, M. A new species of
the *Fejervarya limnocharis* complex from Japan (Anura, Dicroglossidae). *Zool. Sci.* **28**,
922-929 (2011).

337   Dodd Jr, C. K. Population structure and the evolution of sexual size dimorphism and sex
ratios in an insular population of Florida box turtles (*Terrapene carolina bauri*).
Canadian Journal of Zoology **75**, 1495-1507 (1997).

338   Donihue, C. M., Brock, K. M., Foufopoulos, J. & Herrel, A. Feed or fight: testing the
impact of food availability and intraspecific aggression on the functional ecology of an
island lizard. *Funct. Ecol.* **30**, 566-575 (2016).

339   Driessen, M. M. & Rose, R. K. *Isoodon obesulus* (Peramelemorphia: Peramelidae).
Mammalian Species **47**, 112-123 (2015).

340   Du, W.-G., Ji, X. & Zhang, Y.-P. Inter-population variation in life-history traits of a
Chinese lizard (*Takydromus septentrionalis*, Lacertidae). *The Herpetological Journal* **16**,
233-237 (2006).

341   Du, W.-G., Ji, X., Zhang, Y.-P., Xu, X.-F. & Shine, R. Identifying sources of variation in
reproductive and life-history traits among five populations of a Chinese lizard
(*Takydromus septentrionalis*, Lacertidae). *Biol. J. Linn. Soc.* **85**, 443-453 (2005).

342   Dumbell, G. S. *The ecology, behaviour and management of New Zealand Brown Teal, or*
Pateke (Anas aucklandica chlorotis) PhD thesis, University of Auckland, (1987).

343   Dunham, A. E., Tinkle, D. W. & Gibbons, J. W. Body size in island lizards: a cautionary
tale. *Ecology* **59**, 1230-1238 (1978).

344   Dunn, E. R. & Saxe Jr, L. Results of the Catherwood-Chaplin West Indies Expedition,
1948. Part V. Amphibians and reptiles of San Andrés and Providencia. *Proc. Acad. Nat.*
Sci. Phila., 141-165 (1950).

345   Dunstone, N. *et al.* Uso del hábitat, actividad y dieta de la güiña (*Oncifelis guigna*) en el
Parque Nacional Laguna San Rafael, XI Región, Chile. *Boletín del Museo Nacional de*
Historia Natural (Chile) **51**, 147-158 (2002).

346   Elton, S. & Morgan, B. J. Muzzle size, paranasal swelling size and body mass in
Mandrillus leucophaeus. *Primates* **47**, 151-157 (2006).

347   Esteves, R. G. *Comportamento alimentar e aspectos reprodutivos de Bothrops insularis*
(Amaral, 1921)(Serpentes: Viperidae), na criação ex-situ no Instituto Vital Brazil MSc
thesis, Universidade Federal do Estado do Rio de Janeiro, (2016).

348   Faaborg, J. & Winters, J. E. Winter resident returns and longevity and weights of Puerto
Rican birds. *Bird-Banding*, 216-223 (1979).

Fang, Y.-P. & Lee, L.-L. Re-evaluation of the Taiwanese white-toothed shrew, *Crocidura tadae* Tokuda and Kano, 1936 (Insectivora: Soricidae) from Taiwan and two offshore islands. *J. Zool.* **257**, 145-154 (2002).

Faraone, F. P. *Indagini sulla variazione fenotipica in ambienti insulari e microinsulari delle specie di Podarcis Wagler, 1820 (Reptilia, Lacertidae) presenti in Sicilia e in alcune isole circumsiciliane.* PhD thesis, Universitá degli studi di Palermo, (2011).

Faraone, F. P., Giacalone, G. & Lo Valvo, M. in *Atti VIII Congresso Nazionale Societas Herpetologica Italica.* 247-252.

Fitch, H. S. Variation in clutch and litter size in New World reptiles. *Univ Kans Mus Nat Hist Misc Publ* **76**, 1-76 (1985).

Fooden, J. Systematic review of the rhesus macaques, *Macaca mulatta* (Zimmermann, 1780). *Fieldiana Zool* **96**, 180 (2000).

Fooden, J. & Albrecht, G. H. Latitudinal and insular variation of skull size in crab☐eating macaques (Primates, Cercopithecidae: *Macaca fascicularis*). *Am. J. Phys. Anthropol.* **92**, 521-538 (1993).

Ford, H. The honeyeaters of Kangaroo Island. *South Australian Ornithologist* **27**, 134–138 (1976).

Fornasiero, S., Corti, C., Luiselli, L. & Zuffi, M. A. Sexual size dimorphism, morphometry and phenotypic variation in the Whip snake *Hierophis viridiflavus* from a central Mediterranean area. *Revue d'écologie* (2007).

Forsman, A. Variation in sexual size dimorphism and maximum body size among adder populations: effects of prey size. *J. Anim. Ecol.*, 253-267 (1991).

Forsman, A., Merila, J. & Ebenhard, T. Phenotypic evolution of dispersal-enhancing traits in insular voles. *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* **278**, 225-232 (2011).

Foster, J. *The evolution of the native land mammals of the Queen Charlotte Islands and the problem of insularity.* PhD thesis, University of British Columbia, (1963).

Fox, N. C. Some morphological data on the Australasian Harrier (Circus approximans gouldi) in New Zealand. *Notornis* **24**, 9-19 (1977).

Franco, M., Quijano, A. & Soto-Gamboa, M. Communal nesting, activity patterns, and population characteristics in the near-threatened monito del monte, *Dromiciops gliroides*. *J. Mammal.* **92**, 994-1004 (2011).

Fukada, H. Growth and maturity of the Japanese rat snake, *Elaphe climacophora* (Reptilia, Serpentes, Colubridae). *J. Herpetol.*, 269-274 (1978).

Galán, P. Reproductive characteristics of an insular population of the lizard *Podarcis hispanica* from Northwest Spain (Cies Islands, Galicia). *Copeia* **2003**, 657-665 (2003).

Ganem, G., Granjon, L., Ba, K. & Duplantier, J. M. Body size variability and water balance - A comparison between mainland and island populations of *Mastomys huberti* (Rodentia, Muridae) in Senegal. *Experientia* **51**, 402-410 (1995).

Garcés-Restrepo, M. F., Giraldo, A. & Carr, J. L. Population ecology and morphometric variation of the chocoan river turtle (*Rhinoclemmys nasuta*) from two localities on the Colombian Pacific coast. *Boletín Científico. Centro de Museos. Museo de Historia Natural* **17**, 160-171 (2013).

Garcia-Porta, J., Smid, J., Sol, D., Fasola, M. & Carranza, S. Testing the island effect on phenotypic diversification: insights from the Hemidactylus geckos of the Socotra Archipelago. *Sci. Rep.* **6** (2016).

1433   367   Gaulke, M. Overview on the present Knowledge on *Varanus mabitang* Gaulke and Curio,
1434   2001, Including new Morphological and Meristic Data. *Biawak* **4**, 50-58 (2010).
1435   368   Geissmann, T. *Evolution of communication in gibbons (Hylobatidae).* PhD thesis,
1436   Universität Zürich, (1993).
1437   369   Giacalone, G., Abbate, M., Fritz, U. & Lo Valvo, M. Preliminary data on distribution,
1438   morphometric and genetic characterization of Hermann's tortoise in Sicily. *Herpetologia*
1439   *Sardiniae*, 282-286 (2008).
1440   370   Giagia, E. V. *Cytotaxonomical study of eastern European hedgehog Erinaceus concolor*
1441   *M. in Greece* PhD thesis, University of Patras, (1977).
1442   371   Gil Escobedo, L. J. *Ámbitos de Hogar de la Iguana de Órgano Ctenosaura palearis*
1443   *(Sauria: Iguanidae) en el Bosque Tropical Estacionalmente Seco de Cabañas, Zacapa,*
1444   *Guatemala* BSc thesis, Universidade de San Carlos de Guatemala, (2016).
1445   372   Gill, B. & McLean, I. G. Morphometrics of the whitehead *Mohoua albicilla* on Little
1446   Barrier Island, New Zealand. *N. Z. J. Zool.* **13**, 267-271 (1986).
1447   373   Glaw, F., Hawlitschek, O., Glaw, K. & Vences, M. Integrative evidence confirms new
1448   endemic island frogs and transmarine dispersal of amphibians between Madagascar and
1449   Mayotte (Comoros archipelago). *Science of Nature* **106** (2019).
1450   374   Glaw, F., Koehler, J., Townsend, T. M. & Vences, M. Rivaling the World's Smallest
1451   Reptiles: Discovery of Miniaturized and Microendemic New Species of Leaf Chameleons
1452   (Brookesia) from Northern Madagascar. *Plos One* **7** (2012).
1453   375   Glaw, F., Nagy, Z. T., Franzen, M. & Vences, M. Molecular phylogeny and systematics
1454   of the pseudoxyrhophiine snake genus Liopholidophis (Reptilia, Colubridae): evolution
1455   of its exceptional sexual dimorphism and descriptions of new taxa. *Zool. Scr.* **36**, 291-300
1456   (2007).
1457   376   Glaw, F., Vences, M., Andreone, F. & Vallan, D. Revision of the *Boophis majori* group
1458   (Amphibia : Mantellidae) from Madagascar, with descriptions of five new species. *Zool.*
1459   *J. Linn. Soc.* **133**, 495-529 (2001).
1460   377   Goldberg, S. R., Bursey, C. R. & Arreola, J. Gastrointestinal helminths from eight species
1461   of Aspidoscelis (Squamata: Teiidae) from Mexico. *West. N. Am. Nat.*, 223-227 (2014).
1462   378   Goldman, E. A. & Jackson, H. H. Raccoons of North and Middle America. *North*
1463   *American Fauna*, 1-153 (1950).
1464   379   Goltsman, M., Kruchenkova, E. P., Sergeev, S., Volodin, I. & Macdonald, D. W. 'Island
1465   syndrome' in a population of Arctic foxes (*Alopex lagopus*) from Mednyi Island. *J. Zool.*
1466   **267**, 405-418 (2005).
1467   380   González Quintero, E. P. *Análisis taxonómico del coyote (Canis latrans) de la Península*
1468   *de Baja Calfornia, México.* MSc thesis, Centro de Investigaciones Biológicas del
1469   Noroeste, S.C, (2004).
1470   381   González Rossell, A. *Ecología y conservación de la iguana (Cyclura nubila nubila) en*
1471   *Cuba.* PhD thesis, Universidad de Alicante, (2018).
1472   382   Granjon, L. & Cheylan, G. Biometric differentiation of black rat (*Rattus rattus*)
1473   populations in the West Mediterranean islands. *Mammalia* **54**, 213-231 (1990).
1474   383   Grant, P. *A systematic study of the terrestrial birds of the Tres Marias Islands, Mexico.*
1475   (Peabody Museum of Natural History, Yale University, 1965).
1476   384   Grant, P. The coexistence of two wren species of the genus Thryothorus. *The Wilson*
1477   *Bulletin*, 266-278 (1966).

385 Grant, P. *Further information on the relative length of the tarsus in land birds.* (Yale
Peabody Museum of Natural History, 1966).
386 Grant, P. Ecological and morphological variation of Canary Island blue tits, *Parus*
caeruleus (Aves: Paridae). *Biol. J. Linn. Soc.* **11**, 103-129 (1979).
387 Grinnell, J. The species of the mammalian genus Sorex of West-Central California.
University of California Publications in Zoology **10**, 179-195 (1913).
388 Grinnell, J., Dixon, J. S. & Linsdale, J. M. *The fur bearing mammals of California.*,
(University of California Press, 1937).
389 Guarino, F. M., Garcia, G. & Andreone, F. Huge but moderately long-lived: age structure
in the mountain chicken, *Leptodactylus fallax*, from Montserrat, West Indies.
Herpetological Journal **24**, 167-173 (2014).
390 Gursky, S. Effects of radio transmitter weight on a small nocturnal primate. *Am. J.*
Primatol. **46**, 145-155 (1998).
391 Gutsche, A. & Streich, W. J. Demography and Endangerment of the Utila Island Spiny-
Tailed Iguana, *Ctenosaura bakeri*. *J. Herpetol.* **43**, 105-113 (2009).
392 Hadi, S., Ziegler, T. & Hodges, J. K. Group structure and physical characteristics of
Simakobu monkeys (*Simias concolor*) on the Mentawai Island of Siberut, Indonesia.
Folia Primatol. **80**, 74-82 (2009).
393 Hagen, B. Zur Kleinsäugerfauna Siziliens. *Bonn. Zool. Beitr.* **5**, 1-15 (1954).
394 Hai, B. T., Tu, L. N., Duong, V. T. & Son, N. T. in *HỘI NGHỊ KHOA HỌC TOÀN*
QUỐC VỀ SINH THÁI VÀ TÀI NGUYÊN SINH VẬT LẦN THỨ 7 - National Scientific
Conference on Ecological and Biological Resources. 670-676.
395 Harding, L. E. *Trachypithecus cristatus* (Primates: Cercopithecidae). *Mammalian Species*
42, 149-165 (2010).
396 Hasegawa, M. Insular radiation in life-history of the lizard *Eumeces okadae* in the Izu
islands, Japan. *Copeia*, 732-747 (1994).
397 Hasegawa, M. & Moriguchi, H. Geographic variation in food habits, body size and life
history traits of the snakes on the Izu Islands, p. 414–432. *Current herpetology in East*
Asia. M. Matui, T. Hikida, and RC Goris (eds.). Herpetological Society of Japan,
Tokyo.[Google Scholar] (1989).
398 Hawlitschek, O., Nagy, Z. T. & Glaw, F. Island evolution and systematic revision of
Comoran snakes: why and when subspecies still make sense. *PLoS One* **7** (2012).
399 Heaney, L. R. Island area and body size of insular mammals: evidence from the
tri□colored squirrel (*Callosciurus prevosti*) of Southeast Asia. *Evolution* **32**, 29-44
(1978).
400 Heaney, L. R. & Timm, R. M. Systematics and distribution of shrews of the genus
Crocidura (Mammalia: Insectivora) in Vietnam. (1983).
401 Hedges, S. B. & Thomas, R. At the lower size limit in amniote vertebrates: a new
diminutive lizard from the West Indies. *Caribb. J. Sci.* **37**, 168-173 (2001).
402 Heinsohn, G. E. Ecology and reproduction of the Tasmanian bandicoots (*Perameles*
gunni and *Isoodon obesulus*). (1966).
403 Hemelaar, A. Age, growth and other population characteristics of *Bufo bufo* from
different latitudes and altitudes. *J. Herpetol.*, 369-388 (1988).
404 Heo, J.-H. *et al.* Can an invasive prey species induce morphological and behavioral
changes in an endemic predator? Evidence from a South Korean snake (*Oocatochus*
rufodorsatus). *Asian Herpetological Research* **5**, 245-254 (2014).

405 Hernández Gallegos, O., López Moreno, A. E., Méndez de la Cruz, F. R. & Walker, J.
Home range of the parthenogenetic lizard *Aspidoscelis maslini* (Fritts, 1969), on a beach
strand. (2018).

406 Hernandez-Gallegos, O., Esthela Lopez-Moreno, A., Fernando Mendez-Sanchez, J.,
Lloyd Rheubert, J. & Roberto Mendez-de la Cruz, F. Home range of *Aspidoscelis*
cozumela (Squamata: Teiidae): a parthenogenetic lizard microendemic to Cozumel
Island, Mexico. *Rev. Biol. Trop.* **63**, 771-781 (2015).

407 Hernandez-Salinas, U., Ramirez-Bautista, A., Pavon, N. P. & Rosas Pacheco, L. F.
Morphometric variation in island and mainland populations of two lizard species from the
Pacific Coast of Mexico. *Rev. Chil. Hist. Nat.* **87**, 1-9 (2014).

408 Herrel, A., Cottam, M. D., Godbeer, K., Sanger, T. & Losos, J. B. An ecomorphological
analysis of native and introduced populations of the endemic lizard *Anolis maynardi* of
the Cayman Islands. *Breviora* **522**, 1-10 (2011).

409 Herrel, A., Spithoven, L., Van Damme, R. & De Vree, F. Sexual dimorphism of head size
in *Gallotia galloti*: testing the niche divergence hypothesis by functional analyses. *Funct.*
Ecol. **13**, 289-297 (1999).

410 Hershkovitz, P. Mammals of Northern Colombia, Preliminary Report No. 6: Rabbits
(Leporidae), with Notes on the Classification and Distribution of the South American
Forms. *Proc. U. S. Natl. Mus.* (1950).

411 Hervias-Parejo, S. *et al.* Small size does not restrain frugivory and seed dispersal across
the evolutionary radiation of Galapagos lava lizards. *Current Zoology* **65**, 353-361
(2019).

412 Heyer, M. M., Heyer, W. R. & de Sá, R. O. *Leptodactylus pentadactylus*. *Catalogue of*
American Amphibians and Reptiles (CAAR) (2012).

413 Heyer, W., Rand, A., Cruz, C., Peixoto, O. L. & Nelson, C. E. Frogs of Boracéia.
Arquivos de Zoologia (Museu de Zoologia da Universidade Sao Paulo) **31**, 231-410
(1990).

414 Higgins, P. J. *Handbook of Australian, New Zealand and Antarctic birds. Volume 4:*
Parrots to Dollarbird., Vol. 4 (Oxford University Press, 1999).

415 Higgins, P. J. & Davies, S. J. J. F. *Handbook of Australian, New Zealand and Antarctic*
Birds. Volume 3: Snipe to Pigeons. Vol. 3 (Oxford University Press, 1996).

416 Higgins, P. J. & Peter, J. M. *Handbook of Australian, New Zealand and Antarctic Birds.*
Volume 6: Pardalotes to Shrike-thrushes. Vol. 6 (Oxford University Press, 2002).

417 Higgins, P. J., Peter, J. M. & Cowling, S. J. *Handbook of Australian, New Zealand and*
Antarctic Birds. Volume 7: Boatbill to Starlings. Vol. 7 (Oxford University Press, 2006).

418 Higgins, P. J., Peter, J. M. & Steele, W. K. *Handbook of Australian, New Zealand and*
Antarctic Birds. Volume 5: Tyrant-flycatchers to Chats. Vol. 5 (Oxford University Press,
2001).

419 Hill, J. The Robinson collection of Malaysian mammals. *Bull. Raffles Mus.* **29**, 6-22
(1960).

420 Hitchmough, R. A. & McCallum, J. The mammals, birds, reptiles and freshwater fish of
the eastern island group of the Bay of Islands. *Tane* **26**, 127-134 (1980).

421 Hofmeyr, M. D. Egg production in *Chersina angulata*: an unusual pattern in a
Mediterranean climate. *J. Herpetol.* **38**, 172-179 (2004).

422 Hollingsworth, B. D. The systematics of chuckwallas (Sauromalus) with a phylogenetic
analysis of other iguanid lizards. *Herpetol. Monogr.*, 38-191 (1998).

423   How, R. A., Cowan, M. A., Teale, R. J. & Schmitt, L. H. Environmental correlates of
reptile variation on the Houtman Abrolhos archipelago, eastern Indian Ocean. *J.*
Biogeogr. (2020).

424   Huang, W.-S. Ecology and reproductive patterns of the agamid lizard *Japalura swinhonis*
on an east Asian island, with comments on the small clutch sizes of island lizards. *Zool.*
Sci. **24**, 181-188 (2007).

425   Hummelinck, P. W. *Studies on the fauna of Curaçao, Aruba, Bonaire, and the*
Venezuelan Islands: No. 6. (1940).

426   Hutterer, R. Variation and evolution of the Sicilian shrew: taxonomic conclusions and
description of a possibly related species from the Pleistocene of Morocco (Mammalia:
Soricidae). *Bonn. Zool. Beitr.* **42**, 241-251 (1991).

427   Huyghe, K., Vanhooydonck, B., Herrel, A., Tadić, Z. & Van Damme, R. Morphology,
performance, behavior and ecology of three color morphs in males of the lizard *Podarcis*
melisellensis. Integr. Comp. Biol. **47**, 211-220 (2007).

428   Iliopoulou-Georgudaki, J. The relationship between climatic factors and forearm length
of bats: evidence from the chiropterofauna of Lesvos island (Greece–East Aegean).
Mammalia **50**, 475-482 (1986).

429   Inger, R. F., Stuart, B. L. & Iskandar, D. T. Systematics of a widespread southeast Asian
frog, *Rana chalconota* (Amphibia: Anura: Ranidae). *Zool. J. Linn. Soc.* **155**, 123-147
(2009).

430   Iverson, J. B., Hines, K. N. & Valiulis, J. M. The nesting ecology of the Allen Cays rock
iguana, *Cyclura cychlura inornata* in the Bahamas. *Herpetol. Monogr.* **18**, 1-36 (2004).

431   Jacobs, D. Morphological divergence in an insular bat, *Lasiurus cinereus semotus. Funct.*
Ecol., 622-630 (1996).

432   Jenkins, P. & Veitch, C. Sexual dimorphism and age determination in the North Island
saddleback (*Philesturnus carunculatus rufaster*). *N. Z. J. Zool.* **18**, 445-450 (1991).

433   Jenkins, P. D., Abramov, A. V., Bannikova, A. A. & Rozhnov, V. V. Bones and genes:
resolution problems in three Vietnamese species of Crocidura (Mammalia,
Soricomorpha, Soricidae) and the description of an additional new species. *ZooKeys*, 61
(2013).

434   Jenssen, T. A. *et al.* Morphological characteristics of the lizard *Anolis carolinensis* from
South Carolina. *Herpetologica*, 401-411 (1995).

435   Jessop, R. Biometrics and moult of red-capped plovers in Victoria, Tasmania, South
Australia and North West Australia. *Stilt* **17** 29-35 (1990).

436   Jessop, T. S. *et al.* Maximum body size among insular Komodo dragon populations
covaries with large prey density. *Oikos* **112**, 422-429 (2006).

437   Ji, X. & Wang, Z. W. Geographic variation in reproductive traits and trade-offs between
size and number of eggs of the Chinese cobra (*Naja atra*). *Biol. J. Linn. Soc.* **85**, 27-40
(2005).

438   Ji, X., Xie, Y. Y., Sun, P. Y. & Zheng, X. Z. Sexual dimorphism and female reproduction
in a viviparous snake, *Elaphe rufodorsata. J. Herpetol.* **31**, 420-422 (1997).

439   Jimenez, J. E. & McMahon, E. in *Canids: foxes, wolves, jackals, and dogs: status survey*
and conservation action plan (eds C. Sillero-Zubiri, M. Hoffmann, & D.W Macdonald)
(IUCN, 2004).

440 Joubert, L. & Cheylan, M. La tortue d'Hermann de Corse: résultat des recherches menées
en 1985 et 1986. *Travaux scientifiques—Parc naturel régional et réserves naturelles de*
Corse, 1-54 (1989).
441 Judd, F. W. & Ross, R. K. Year-to-year variation in clutch size of island and mainland
populations of Holbrookia propinqua (Reptilia, Lacertilia, Iguanidae). . *J. Herpetol.* **12**,
203-207 (1978).
442 Kang, C. *et al.* Differential predation drives the geographical divergence in multiple traits
in aposematic frogs. *Behav. Ecol.* **28**, 1122-1130 (2017).
443 Keast, A. Competitive interactions and the evolution of ecological niches as illustrated by
the Australian honeyeater genus Melithreptus (Meliphagidae). *Evolution*, 762-784
(1968).
444 Keehn, J. E., Nieto, N. C., Tracy, C. R., Gienger, C. M. & Feldman, C. R. Evolution on a
desert island: body size divergence between the reptiles of Nevada's Anaho Island and the
mainland around Pyramid Lake. *J. Zool.* **291**, 269-278 (2013).
445 Kim, D.-I. *et al.* Patterns of morphological variation in the Schlegel's Japanese gecko
(*Gekko japonicus*) across populations in China, Japan, and Korea. *Journal of Ecology*
and Environment **43**, 34 (2019).
446 Kim, S.-I., Oshida, T., Lee, H., Min, M.-S. & Kimura, J. Evolutionary and
biogeographical implications of variation in skull morphology of raccoon dogs
(*Nyctereutes procyonoides*, Mammalia: Carnivora). *Biol. J. Linn. Soc.* **116**, 856-872
(2015).
447 King, R. B. Body size variation among island and mainland snake populations.
Herpetologica **45**, 84-88 (1989).
448 King, R. B. Variation in brown snake (*Storeria dekayi*) morphology and scalation: Sex,
family, and microgeographic differences. *J. Herpetol.* **31**, 335-346 (1997).
449 King, T. The birds of the Lesio-Louna and Lefini Reserves, Batéké Plateau, Republic of
Congo. *Malimbus* **33**, 1-41 (2011).
450 Kitchener, D., Hisheh, S., Schmitt, L. & Maryanto, I. Morphological and genetic
variation in *Aethalops alecto* (Chiroptera, Pteropodidae) from Java, Bali and Lombok Is,
Indonesia. *Mammalia* **57**, 255-272 (1993).
451 Kitchener, D. & Schmitt, L. Morphological and genetic variation in *Suncus murinus*
(Soricidae: Crocidurinae) from Java, Lesser Sunda islands, Maluku and Sulawesi,
Indonesia. *Mammalia* **58**, 433-452 (1994).
452 Kitchener, D., Schmitt, L., Hisheh, S., How, R. & Cooper, N. Morphological and genetic
variation in the bearded tomb bats (Taphozous: Emballonuridae) of Nusa Tenggara,
Indonesia. *Mammalia* **57**, 63-84 (1993).
453 Klauber, L. M. The gopher snakes of Baja California: with descriptions of a new
subspecies of *Pituophis catenifer*. *Transactions of the San Diego Society of Natural*
History **11**, 1-40 (1946).
454 Klauber, L. M. A new gopher snake (Pituophis) from Santa Cruz Island, California
Transactions of the San Diego Society of Natural History **11**, 41-48 (1946).
455 Klauber, L. M. Some new and revived subspecies of rattlesnakes. *Transactions of the San*
Diego Society of Natural History **11**, 61-116 (1949).
456 Klauber, L. M. *Rattlesnakes: their habits, life histories, and influence on mankind*.
(University of California Press, 1997).
457 Kloss, C. B. On mammals collected in Siam. *J Nat Hist Soc Siam* **4**, 333-407 (1919).

458 Klütsch, C., Misof, B., Grosse, W. R. & Moritz, R. Genetic and morphometric
differentiation among island populations of two Norops lizards (Reptilia: Sauria:
Polychrotidae) on independently colonized islands of the Islas de Bahia (Honduras). *J.*
Biogeogr. **34**, 1124-1135 (2007).
459 Knapp, C. R., Iverson, J. B. & Owens, A. K. Geographic variation in nesting behavior
and reproductive biology of an insular iguana (*Cyclura cychlura*). *Can. J. Zool./Rev. Can.*
Zool. **84**, 1566-1575 (2006).
460 Kohno, H. Reptiles in a seabird colony: herpetofauna of Nakanokamishima Island of the
Yaeyama group, Ryukyu Archipelago. *Island Studies in Okinawa* **9**, 73-89 (1991).
461 Koopman, K. F. & Diamond, J. Zoogeography of mammals from islands off the
northeastern coast of New Guinea. *Am. Mus. Novit.* **2690** (1979).
462 Kraus, C., Trillmich, F. & Künkele, J. Reproduction and growth in a precocial small
mammal, *Cavia magna*. *J. Mammal.* **86**, 763-772 (2005).
463 Proetzel, D., Ruthensteiner, B., Scherz, M. D. & Glaw, F. Systematic revision of the
Malagasy chameleons *Calumma boettgeri* and *C. linotum* (Squamata: Chamaeleonidae).
Zootaxa **4048**, 211-231 (2015).
464 Qualls, C. P., Shine, R., Donnellan, S. & Hutchinsonm, M. The evolution of viviparity
within the Australian scincid lizard *Lerista bougainvillii*. *J. Zool.* **237**, 13-26 (1995).
465 Quin, D. G., Smith, A. P. & Norton, T. W. Eco-geographic variation in size and sexual
dimorphism in sugar gliders and squirrel gliders (Marsupialia: Petauridae). *Aust. J. Zool.*
44, 19-45 (1996).
466 Rader, J. A., Dillon, M. E., Chesser, R. T., Sabat, P. & del Rio, C. M. Morphological
divergence in a continental adaptive radiation: South American ovenbirds of the genus
Cinclodes. *Auk* **132**, 180-190 (2015).
467 Radočaj, M., Jelić, D., Karaica, D. & Kapelj, S. Morphological and reproductive traits of
the insular population of *Podarcis siculus* (Reptilia: Lacertidae) from Krk Island
(Croatia) *Hyla* **2**, 5-22 (2011).
468 Raia, P. *et al.* The blue lizard spandrel and the island syndrome. *BMC Evol. Biol.* **10**, 289
(2010).
469 Rajaratnam, R. *Ecology of leopard cat (Prionailurus bengalensis) in Tabin Wildlife*
Reserve,Sabah, Malaysia. PhD thesis, Universiti Kebangsaan Malaysia (2000).
470 Rausch, R. L. Geographic variation in size in North American brown bears, *Ursus arctos*
L., as indicated by condylobasal length. *Can. J. Zool./Rev. Can. Zool.* **41**, 33-45 (1963).
471 Reboucas, R., da Silva, H. R., Sanuy, D. & Sole, M. Sexual maturity and growth of male
toads (*Rhinella ornata*): A comparison between insular and mainland populations. *Zool.*
Anz. **283**, 12-19 (2019).
472 Reboucas, R., da Silva, H. R. & Sole, M. Frog size on continental islands of the coast of
Rio de Janeiro and the generality of the island rule. *PloS One* **13** (2018).
473 Redfield, J. Distribution, abundance, size, and genetic variation of *Peromyscus*
maniculatus on the Gulf Islands of British Columbia. *Can. J. Zool./Rev. Can. Zool.* **54**,
463-474 (1976).
474 Renaud, S. *et al.* Morphometrics and genetics highlight the complex history of Eastern
Mediterranean spiny mice. *Biol. J. Linn. Soc.* (2020).
475 Reynolds, R. G., Niemiller, M. L. & Fitzpatrick, B. M. Genetic analysis of an endemic
archipelagic lizard reveals sympatric cryptic lineages and taxonomic discordance.
Conserv. Genet. **13**, 953-963 (2012).

476 Rhodin, A. G. Chelid turtles of the Australasian archipelago: a new species of Chelodina
from Roti island, Indonesia. *Breviora* **498**, 1-31 (1994).

477 Rickart, E. A., Heaney, L. R., Goodman, S. M. & Jansa, S. Review of the Philippine
genera Chrotomys and Celaenomys (Murinae) and description of a new species. *J.*
Mammal. **86**, 415-428 (2005).

478 Ridgway, R. *The birds of North and Middle America: A descriptive catalogue of the*
higher groups, genera, species, and subspecies of birds known to occur in North
America, from the Arctic lands to the Isthmus of Panama, the West Indies and other
islands of the Caribbean sea, and the Galapagos Archipelago. (US Government Printing
Office, 1904).

479 Rivas, G. A. *et al.* A distinctive new species of Gonatodes (Squamata:
Sphaerodactylidae) from Isla La Blanquilla, Venezuela, with remarks on the distribution
of some other Caribbean sphaerodactylid lizards. *South Am. J. Herpetol.* **8**, 5-18 (2013).

480 Robertson, H., Whitaker, A. & Fitzgerald, B. Morphometrics of forest birds in the
Orongorongo Valley, Wellington, New Zealand. *N. Z. J. Zool.* **10**, 87-97 (1983).

481 Rocha, C. F. & Vrcibradic, D. Reproductive traits of two sympatric viviparous skinks
(*Mabuya macrorhyncha* and *Mabuya agilis*) in a Brazilian restinga habitat.
Herpetological Journal **9**, 43-53 (1999).

482 Rodríguez-Cabal, M. A., Amico, G. C., Novaro, A. J. & Aizen, M. A. Population
characteristics of *Dromiciops gliroides* (Philippi, 1893), an endemic marsupial of the
temperate forest of Patagonia. *Mamm. Biol.* **73**, 74-76 (2008).

483 Rog, S., Ryan, M. J., Mueller, U. & Lampert, K. P. Evidence for morphological and
genetic diversification of túngara frog populations on islands. *Herpetol. Conserv. Biol.* **8**,
228-239 (2013).

484 Rogers, K., Rogers, A. & Rogers, D. Bander's Aid. *Supplement Number One. Royal*
Australasian Ornithologists' Union Report (1990).

485 Ross, R. K. & Judd, F. W. Comparison of lipid cycles of *Holbrookia propinqua* from
Padre island and mainland Texas. *J. Herpetol.* **16**, 53-60 (1982).

486 Rowe-Rowe, D. & Crafford, J. Density, body size, and reproduction of feral house mice
on Gough Island. *S. Afr. J. Zool.* **27**, 1-5 (1992).

487 Runemark, A., Sagonas, K. & Svensson, E. I. Ecological explanations to island
gigantism: dietary niche divergence, predation, and size in an endemic lizard. *Ecology* **96**,
2077-2092 (2015).

488 Sabater Pi, J. Contribution to the ecology of *Colobus polykomos satanas* (Waterhouse,
1838) of Rio Muni, Republic of Equatorial Guinea. *Folia Primatol.* **19**, 193-207 (1973).

489 Sagonas, K., Pafilis, P. & Valakos, E. D. Effects of insularity on digestion: living on
islands induces shifts in physiological and morphological traits in island reptiles. *The*
Science of Nature **102**, 55 (2015).

490 Sahimi, H. N. M., Chubo, J. K., Mohd, M., Saripuddin, N. B. & Ab Rahim, S. S. The
distribution and population density of Bornean tarsier, *Tarsius bancanus borneanus*
(Elliot) in secondary and rehabilitated forests of Universiti Putra Malaysia, Bintulu
Sarawak Campus, Sarawak, Malaysia. *Tropical Life Sciences Research* **29**, 139 (2018).

491 Salvidio, S., Cresta, P. & Dolmen, D. The common toad *Bufo bufo* population of Hitra
island, central Norway. *Fauna Norvegica Series A* **14**, 51-55 (1993).

492 Sanches, V. Q. A. & Grings, D. R. Daily movement and habitat use of *Iguana iguana*
(Linnaeus, 1758) in an urban second growth Amazonian forest fragment in Brazil.
Herpetology Notes **11**, 93-96 (2018).
493 Sanderson, J., Sunquist, M. E. & Iriarte, A. W. Natural history and landscape-use of
guignas (Oncifelis guigna) on Isla Grande de Chiloé, Chile. *J. Mammal.* **83**, 608-613
(2002).
494 Sargis, E. J., Millien, V., Woodman, N. & Olson, L. E. Rule reversal: Ecogeographical
patterns of body size variation in the common treeshrew (Mammalia, Scandentia). *Ecol.*
Evol. **8**, 1634-1645 (2018).
495 Sá☐Sousa, P., Almeida, A., Rosa, H., Vicente, L. & Crespo, E. Genetic and
morphological relationships of the Berlenga wall lizard (*Podarcis bocagei berlengensis*:
Lacertidae). *J. Zool. Syst. Evol. Res.* **38**, 95-102 (2000).
496 Sá-Sousa, P. & Harris, D. J. *Podarcis carbonelli* Pérez-Mellado, 1981 is a distinct
species. *Amphibia-Reptilia* **23**, 459-468 (2002).
497 Sá-Sousa, P., Vicente, L. & Crespo, E. Morphological variability of *Podarcis hispanica*
(Sauria: lacertidae) in Portugal. *Amphibia-Reptilia* **23**, 55-69 (2002).
498 Scali, S. *et al.* in *11th Ordinary General Meeting Societas Europaea Herpetologica*
(SEH). 161-166.
499 Scheffer, V. B. & Dalquest, W. W. A new shrew from Destruction Island, Washington. *J.*
Mammal. **23**, 333-335 (1942).
500 Schifter, H., Cunningham-van Someren, G. & van Someren, G. C. The Avifauna of the
North Nandi Forest, Kenya. *Annalen des Naturhistorischen Museums in Wien. Serie B für*
Botanik und Zoologie **100**, 425-479 (1998).
501 Schlotfeldt, B. E. & Kleindorfer, S. Adaptive divergence in the Superb Fairy-wren
(*Malurus cyaneus*): a mainland versus island comparison of morphology and foraging
behaviour. *Emu* **106**, 309-319 (2006).
502 Schultz, A. H. Observations on the growth, classification and evolutionary specialization
of gibbons and siamangs. *Human Biology* **5**, 212 (1933).
503 Schultz, A. H. The relative weight of the testes in primates. *The Anatomical Record* **72**,
387-394 (1938).
504 Schwaner, T. D. & Sarre, S. D. Body size of tiger snakes in Southern Australia, with
particular reference to *Notechis ater serventyi* (Elapidae) on Chappell Island. *J. Herpetol.*
22, 24-33 (1988).
505 Schwaner, T. D. & Sarre, S. D. Body size and sexual dimorphism in mainland and island
tiger snakes. *J. Herpetol.* **24**, 320-322 (1990).
506 Schwartz, A. Three new mammals from southern Florida. *J. Mammal.* **33**, 381-385
(1952).
507 Schwartz, A. Snakes of the genus Alsophis in Puerto Rico and the Virgin Islands. *Studies*
on the Fauna of Curaçao and other Caribbean Islands **23**, 177-227 (1966).
508 Scott, D. Notes on the Eastern Hogsnose Snake, *Heterodon platyrhinos* Latreille
(Squamata, Colubridae), on a Virginia Barrier Island. *Brimleyana*, 51-55 (1986).
509 Selcer, K. W. & Judd, F. W. Variation in the reproductive ecology of *Holbrookia*
propinqua (Sauria, Iguanidae). *Texas Journal of Science* **34**, 125-135 (1982).
510 Senczuk, G., García, A., Colangelo, P., Annesi, F. & Castiglia, R. Morphometric and
genetic divergence in island and mainland populations of *Anolis nebulosus* (Squamata:

1795   Polychrotidae) from Jalisco (Mexico): an instance of insular gigantism. *Ital. J. Zool.* **81**,
1796   204-214 (2014).

1797   511   Seock, M., Nam, K.-B. & Yoo, J.-C. Distribution and movement tendencies of short-
1798   tailed viper snakes (*Gloydius saxatilis*) by altitude. *Asian Herpetological Research* **8**, 39-
1799   47 (2017).

1800   512   Serrano-Cardozo, V. H., Ramírez-Pinilla, M. P., Ortega, J. E. & Cortes, L. A. Annual
1801   reproductive activity of *Gonatodes albogularis* (Squamata: Gekkonidae) living in an
1802   anthropicarea in Santander, Colombia. *South Am. J. Herpetol.* **2**, 31-38 (2007).

1803   513   Shaidani, N.-I. *The biogeographic origins and trophic ecology of Maine's island red-*
1804   *backed salamanders (Plethodon cinereus)* MSc thesis, University of Maine, (2017).

1805   514   Sharples, C. M., Fa, J. E. & BELL, D. J. Geographical variation in size in the European
1806   rabbit *Oryctolagus cuniculus* (Lagomorpha: Leporidae) in western Europe and North
1807   Africa. *Zool. J. Linn. Soc.* **117**, 141-158 (1996).

1808   515   Shekelle, M. Observations of wild Sangihe island tarsiers *Tarsius sangirensis*. *Asian*
1809   *Primates Journal* **3**, 18-23 (2013).

1810   516   Shimada, T., Maeda, S. & Sakakibara, M. A morphological study of *Cynops*
1811   *pyrrhogaster* from the Chita Peninsula: rediscovery of the "extinct" Atsumi race endemic
1812   to peninsular regions of Aichi prefecture, Central Japan. *Curr. Herp.* **35**, 38-52 (2016).

1813   517   Shine, R. Venomous snakes in cold climates: ecology of the Australian genus Drysdalia
1814   (Serpentes: Elapidae). *Copeia*, 14-25 (1981).

1815   518   Shine, R., Sun, L.-x., Zhao, E. & Bonnet, X. A review of 30 years of ecological research
1816   on the Shedao pitviper, *Gloydius shedaoensis*. *Herpetological Natural History* **9**, 1-14
1817   (2003).

1818   519   Sicuro, F. L. & Oliveira, L. F. B. Variations in leopard cat (*Prionailurus bengalensis*)
1819   skull morphology and body size: sexual and geographic influences. *PeerJ* **3**, e1309
1820   (2015).

1821   520   Siler, C. D., McVay, J. D., Diesmos, A. C. & Brown, R. M. A new species of fanged
1822   frog, genus Limnonectes (Amphibia: Anura: Dicroglossidae) from southeast Mindanao
1823   Island, Philippines. *Herpetologica* **65**, 105-114 (2009).

1824   521   Siliceo-Cantero, H. H., García, A., Reynolds, R. G., Pacheco, G. & Lister, B. C.
1825   Dimorphism and divergence in island and mainland Anoles. *Biol. J. Linn. Soc.* **118**, 852-
1826   872 (2016).

1827   522   Simberloff, D., Dayan, T., Jones, C. & Ogura, G. Character displacement and release in
1828   the small Indian mongoose, *Herpestes javanicus*. *Ecology* **81**, 2086-2099 (2000).

1829   523   Sinclair, E. Morphological variation among populations of the quokka, Setonix
1830   brachyurus (Macropodidae: Marsupialia), in Western Australia. *Aust. J. Zool.* **46**, 439-
1831   449 (1998).

1832   524   Siqueira, C. C., Van Sluys, M., Ariani, C. V. & Rocha, C. F. Feeding ecology of *Thoropa*
1833   *miliaris* (Anura, Cycloramphidae) in four areas of Atlantic rain forest, southeastern
1834   Brazil. *J. Herpetol.* **40**, 520-525 (2006).

1835   525   Slábová, M. & Frynta, D. Morphometric variation in nearly unstudied populations of the
1836   most studied mammal: The non-commensal house mouse (*Mus musculus domesticus*) in
1837   the Near East and Northern Africa. *Zoologischer Anzeiger-A Journal of Comparative*
1838   *Zoology* **246**, 91-101 (2007).

1839   526   Sleeman, D., Cussen, R., Southey, A. & O'Leary, D. The Badgers *Meles meles* (L.) of
1840   Coney Island, Co Sligo. *The Irish Naturalists' Journal*, 10-18 (2002).

527 Sleeman, D. P., Davenport, J., Cussen, R. E. & Hammond, R. F. The small-bodied
badgers (*Meles meles* (L.)) of Rutland Island, Co. Donegal. *The Irish Naturalists'*
Journal, 1-6 (2009).
528 Sleeman, P., Cussen, R., O'Donoughue, T. & Costello, E. Badgers (*Meles meles*) on
Fenit island, and their presence or absence on other Islands in co. Kerry, Ireland. *Small*
carnivore conservation **10** (2001).
529 Smith, F. A. Evolution of body size among woodrats from Baja California, Mexico.
Funct. Ecol., 265-273 (1992).
530 Smith, R. J. & Jungers, W. L. Body mass in comparative primatology. *J. Hum. Evol.* **32**,
523-559 (1997).
531 Smith, T. B. & Temple, S. A. Feeding habits and bill polymorphism in hook-billed kites.
The Auk **99**, 197-207 (1982).
532 Society, F. o. M.-L. I.   (2008).
533 Sody, H. Notes on some primates, carnivora, and the babirusa from the Indo-Malayan
and Indo-Australian regions. *Treubia* **20**, 121-190 (1949).
534 Song, L., Fa-Hong, Y. & Xue-Fei, L. Cranial morphometric study of four giant flying
squirrels (Petaurista)(Rodentia: Sciuridae) from China. *Zool. Res.* **33**, 119-126 (2012).
535 Sparkman, A. M. *et al.* Convergence in reduced body size, head size, and blood glucose
in three island reptiles. *Ecol. Evol.* **8**, 6169-6182 (2018).
536 Stager, K. E. The avifauna of the Tres Marias islands, Mexico. *The Auk* **74**, 413-432
(1957).
537 Stamenković, S. & Matić, R. Morphological correlates of prey consumed by *Podarcis*
melisellensis (Braun, 1877) and *P. siculus* (Rafinesque, 1810)(Sauria, Lacertidae) from
two mainland regions in the eastern Adriatic area. *Archives of Biological Sciences* **65**,
1015-1025 (2013).
538 Steven, D. M. Recent evolution in the genus Clethrionomys. *Symp. Soc. Exp. Biol.* **7**,
310-319 (1953).
539 Stoddart, D. M. & Braithwaite, R. W. A strategy for utilization of regenerating heathland
habitat by the brown bandicoot (*Isoodon obesulus*; Marsupialia, Peramelidae). *J. Anim.*
Ecol., 165-179 (1979).
540 Storer, R. W. Variation in the red-tailed hawks of southern Mexico and Central America.
The Condor **64**, 77-78 (1962).
541 Strickland, D. & Norris, D. R. An example of phenotypic adherence to the island rule? -
Anticosti gray jays are heavier but not structurally larger than mainland conspecifics.
Ecol. Evol. **5**, 3687-3694 (2015).
542 Struhsaker, T. T. *The red colobus monkeys: variation in demography, behavior, and*
ecology of endangered species. (Oxford University Press, 2010).
543 Stuart-Smith, J. F., Stuart-Smith, R. D., Swain, R. & Wapstra, E. Size dimorphism in
Rankinia Tympanocryptis diemensis (Family Agamidae): sex-specific patterns and
geographic variation. *Biol. J. Linn. Soc.* **94**, 699-709 (2008).
544 Stubbs, D. & Swingland, I. R. The ecology of a Mediterranean tortoise (*Testudo*
hermanni): a declining population. *Can. J. Zool./Rev. Can. Zool.* **63**, 169-180 (1985).
545 Swarth, H. The lemming of Nunivak Island, Alaska. *Proc. Biol. Soc. Wash* **44**, 101-104
(1931).
546 Swarth, H. S. *Birds and mammals of the 1909 Alexander Alaska expedition*. Vol. 7 (The
University Press, 1911).

547   Takada, Y., Sakai, E., Uematsu, Y. & Tateishi, T. Morphometric variation of house mice
(*Mus musculus*) on the Izu Islands. *Mammal Study* **24**, 51-65 (1999).
548   Takada, Y., Sakai, E., Uematsu, Y. & Tateishi, T. Morphological variation of large
Japanese field mice, *Apodemus speciosus* on the Izu and Oki Islands. *Mammal Study* **31**,
29-40 (2006).
549   Takada, Y., Yamada, H. & Tateishi, T. Morphometric variation of Japanese wild mice on
islands. *Journal of the Mammalogical Society of Japan* **19**, 113-128 (1994).
550   Takenaka, T. & Hasegawa, M. Female-biased mortality and its consequence on adult sex
ratio in the freshwater turtle *Chinemys reevesii* on an island. *Curr. Herp.* **20**, 11-17
(2001).
551   Tamarin, R. H. Dispersal, population regulation, and K-selection in field mice. *The*
American Naturalist **112**, 545-555 (1978).
552   Tanaka, K. Phenotypic plasticity of body size in an insular population of a snake.
Herpetologica **67**, 46-57 (2011).
553   Tanaka, K. & Ota, H. Natural history of two colubrid snakes, *Elaphe quadrivirgata* and
Rhabdophis tigrinus, on Yakushima Island, southwestern Japan. *Amphibia-Reptilia* **23**,
323-331 (2002).
554   Taverne, M. *et al.* Diet variability among insular populations of Podarcis lizards reveals
diverse strategies to face resource limited environments. *Ecol. Evol.* **9**, 12408-12420
(2019).
555   Taylor, E. H. The lizards of Thailand. *University of Kansas Science Bulletin* **44**, 687-
1077 (1963).
556   Taylor, H. L. & Cooley, C. R. A multivariate analysis of morphological variation among
parthenogenetic teiid lizards of the *Cnemidophorus cozumela* complex. *Herpetologica*,
67-76 (1995).
557   Terada, C., Tatsuzawa, S. & Saitoh, T. Ecological correlates and determinants in the
geographical variation of deer morphology. *Oecologia* **169**, 981-994 (2012).
558   Terborgh, J., Faaborg, J. & Brockmann, H. J. Island colonization by Lesser Antillean
birds. *The Auk* **95**, 59-72 (1978).
559   Tessa, G., Crottini, A., Giacoma, C., Guarino, F. M. & Randrianirina, J. E. Comparative
longevity and age at sexual maturity in twelve rainforest frogs of the genera Boophis,
Gephyromantis, and Mantidactylus (Anura: Mantellidae) from Madagascar.
Phyllomedusa: Journal of Herpetology **16**, 13-21 (2017).
560   Thomas, D. & Broughton, E. *Status of three Canadian caribou populations north of 70 in*
winter 1977. (Canadian Wildlife Service, 1978).
561   Thouless, C. & Bassri, K. A. Taxonomic status of the Farasan Island gazelle. *J. Zool.*
223, 151-159 (1991).
562   Tsai, T. Sexual dimorphism of Chinese green tree viper, *Trimeresurus stejnegeri*
stejnegeri. *Biol Bull Nat Taiwan Normal Univ* **33**, 13-22 (1998).
563   Uller, T. *et al.* Genetic differentiation predicts body size divergence between island and
mainland populations of common wall lizards (*Podarcis muralis*). *Biol. J. Linn. Soc.* **127**,
771-786 (2019).
564   Ursin, E. *Geographical variation in Apodemus sylvaticus and A. flavicollis (Rodentia,*
Muridae) in Europe, with special reference to Danish and Latvian populations.
(Munksgaard, 1956).

565 Valakos, E. D. & Polymeni, R. M. The food of *Cyrtodactylus kotschyi* (Steindachner,
1870) (Sauria - Gekkonidae) during the wet season in the Mediterranean insular
ecosystems of the Aegean. *Herpetological Journal* **1**, 474-477 (1990).
566 Van Heezik, Y. M., Cooper, J. & Seddon, P. J. Population characteristics and
morphometrics of angulate tortoises on Dassen Island, South Africa. *J. Herpetol.*, 447-
453 (1994).
567 Van Weenen, J. *Aspects of the ecology of Cunningham's skink, Egernia cunninghami, on*
West Island, South Australia. BSc (Honours) thesis, University of Adelaide, (1995).
568 Vanek, J. P. & Burke, R. L. Insular dwarfism in female eastern hog-nosed snakes
(*Heterodon platirhinos*; Dipsadidae) on a barrier island. *Can. J. Zool./Rev. Can. Zool.* **98**,
157-164 (2020).
569 Vanzolini, P. E. & Reboucas-Spieker, R. Distribution and differentiation of animals along
the coast and in continental islands of the state of São Paulo, Brazil. 3. Reproductive
differences between *Mabuya caissara* and *Mabuya macrorhyncha* (Sauria, Scincidae).
Papéis Avulsos de Zoologia, São Paulo **29**, 95-109 (1976).
570 Vasconcelos, R. & Carranza, S. Systematics and biogeography of *Hemidactylus*
homoeolepis Blanford, 1881 (Squamata: Gekkonidae), with the description of a new
species from Arabia. *Zootaxa* **3835**, 501-527 (2014).
571 Velo-Anton, G. & Cordero-Rivera, A. Ethological and phenotypic divergence in insular
fire salamanders: diurnal activity mediated by predation? *Acta Ethologica* **20**, 243-253
(2017).
572 Vences, M. Erste untersuchungen über die fortpflanzungsbiologie des Iberisichen
Scheibenzünglers *Discoglossus galganoi* (Amphibia: Anura: Discoglossidae). *Acta*
Biologica Benrodis **6**, 89-98 (1994).
573 Vences, M., Glaw, F., Mercurio, V. & Andreone, F. Review of the Malagasy tree snakes
of the genus Stenophis (Colubridae). *Salamandra* **40**, 161-179 (2004).
574 Vences, M., Köhler, J., Pabijan, M. & Glaw, F. Two syntopic and microendemic new
frogs of the genus Blommersia from the east coast of Madagascar. *African Journal of*
Herpetology **59**, 133-156 (2010).
575 Ventura, J. & Fuster, M. J. L. Morphometric analysis of the black rat, *Rattus rattus*, from
Congreso Island (Chafarinas Archipielago, Spain). *Orsis: organismes i sistemes* **15**, 91-
102 (2000).
576 Verdon, E. & Donnelly, M. A. Population structure of Florida box turtles (*Terrapene*
carolina bauri) at the southernmost limit of their range. *J. Herpetol.* **39**, 572-577 (2005).
577 Vershinin, A. The biology and trapping of the ermine in Kamchatka. *Byull. Mosk. o-va*
Ispyt. Prir. Otd. Biol **77**, 16-26 (1972).
578 Vervust, B., Grbac, I. & Van Damme, R. Differences in morphology, performance and
behaviour between recently diverged populations of *Podarcis sicula* mirror differences in
predation pressure. *Oikos* **116**, 1343-1352 (2007).
579 Villeneuve, A. R. Habitat selection and population density of the world's smallest
chameleon, *Brookesia micra,* on Nosy Hara,. Madagascar. *Herpetol. Conserv. Biol.* **12**,
334-341 (2017).
580 Vogel, P. & Sofianidou, T. The shrews of the genus Crocidura on Lesbos, an eastern
Mediterranean island. *Bonn. Zool. Beitr.* **46**, 339-347 (1996).
581 Vogrin, M. Sexual Dimorphism in *Podarcis sicula campestris*. *Turkish Journal of*
Zoology **29**, 189-191 (2005).

Vrcibradic, D. & Rocha, C. F. Observations on the natural history of the lizard *Mabuya macrorhyncha* Hoge (Scincidae) in Queimada Grande island, Sao Paulo, Brazil. *Rev. Bras. Zool.* **22**, 1185-1190 (2005).

Walker, R., Woods-Ballard, A. J. & Rix, C. E. Population density and seasonal activity of the threatened Madagascar spider tortoise (*Pyxis arachnoides arachnoides*) of the southern dry forests; South West Madagascar. *Afr. J. Ecol.* **46**, 67 (2008).

Wang, Y., Li, Y., Wu, Z. & Murray, B. Insular shifts and trade□offs in life□history traits in pond frogs in the Zhoushan Archipelago, China. *J. Zool.* **278**, 65-73 (2009).

Watkins, G. G. Proximate causes of sexual size dimorphism in the iguanian lizard Microlophus occipitalis. *Ecology* **77**, 1473-1482 (1996).

Wayne, R. K. *et al.* A morphologic and genetic study of the island fox, *Urocyon littoralis*. *Evolution* **45**, 1849-1868 (1991).

West, A. G. *Variation in mandible shape and body size of house mice Mus musculus across the New Zealand archipelago: A Trans-Tasman comparison using geometric morphometrics* MSc thesis, University of Waikato, (2017).

Whitehead, V. *Populationd dynamics of the eastern box turtle (Terrapene carolina carolina) in the Maryville College Woods, Maryville, Tennessee: a report of a senior study*, Maryville College, (2017).

Woodman, N. *et al.* A new southern distributional limit for the Central American rodent *Peromyscus stirtoni*. *Caribb. J. Sci.* **38**, 281-284 (2002).

Wu, Z., Li, Y. & Murray, B. R. Insular shifts in body size of rice frogs in the Zhoushan Archipelago, China. *J. Anim. Ecol.* **75**, 1071-1080 (2006).

Xu, F., Adler, G. H. & Li, Y. Covariation in insular life-history traits of the rice frog (*Fejervarya limnocharis*) in Eastern China. *Asian Herpetological Research* **4**, 28-35 (2013).

Yabe, T. Population structure and male melanism in the Reeves' turtle, Chinemys reevesii. *Japanese Journal of Herpetology* **15**, 131-137 (1994).

Yasukawa, Y., Ota, H. & Iverson, J. B. Geographic variation and sexual size dimorphism in Mauremys mutica (Cantor, 1842)(Reptilia: Bataguridae), with description of a new subspecies from the southern Ryukyus, Japan. *Zool. Sci.* **13**, 303-317 (1996).

Yurgenson, P. B. Ermines of the Far East territory. *Byull. Mosk. o-va Ispyt. Prir. Otd. Biol* **45**, 239-243 (1936).

Yusefi, G. H., Kiabi, B. H., Khalatbari, L., Faizolahi, K. & Monteiro, N. M. Morphological analysis of Brandt's hedgehog (Paraechinus hypomelas) reflects the isolation history of Persian Gulf islands and has implications for taxonomy. *Biol. J. Linn. Soc.* **119**, 497-510 (2016).

Yustian, I. *Ecology and conservation status of Tarsius bancanus saltator on Belitung Island, Indonesia*. (Cuvillier Verlag, 2007).

Ziegler, T., Vences, M., Glaw, F. & Böhme, W. Remarks on the genital morphology of the Malagasy snake genus Liophidium (Reptilia, Serpentes, Colubridae). *Acta biologica Benrodis* **8** (1996).

Zihlman, A. L., Mootnick, A. R. & Underwood, C. E. Anatomical contributions to hylobatid taxonomy and adaptation. *Int. J. Primatol.* **32**, 865-877 (2011).

Zuffi, M. & Gariboldi, A. Sexual dimorphism of the European pond terrapin, Emys orbicularis (L., 1758) from Italy. *Scientia herpetologica* **1995**, 124-129 (1995).

600 Zuffi, M. A., Sacchi, R., Pupin, F. & Cencetti, T. Sexual size and shape dimorphism in
the Moorish gecko (*Tarentola mauritanica,* Gekkota, Phyllodactylidae). *North-Western*
Journal of Zoology **7**, 189-197 (2011).

601 Zuffi, M. A. L., Odetti, F., Batistoni, R. & Mancino, G. Geographic variation of sexual
size dimorphism and genetics in the European pond turtle, *Emys orbicularis* and *Emys*
trinacris, of Italy. *Ital. J. Zool.* **73**, 363-372 (2006).

602 Zug, G. R., Hedges, S. B. & Sunkel, S. Variation in reproductive parameters of three
neotropical snakes, *Coniophanes fissidens*, *Dipsas catesbyi*, and *Imantodes cenchoa*.
Smithson. Contrib. Zool. **300** (1979).

603 Higgins, J. P. *et al. Cochrane handbook for systematic reviews of interventions.* (John
Wiley & Sons, 2019).

604 Santini, L., Benítez☐López, A., Ficetola, G. F. & Huijbregts, M. A. Length–mass
allometries in amphibians. *Integr. Zool.* **13**, 36-45 (2018).

605 Lislevand, T., Figuerola, J. & Székely, T. Avian body sizes in relation to fecundity,
mating system, display behavior, and resource sharing. *Ecology* **88**, 1605-1605 (2007).

606 Van Valkenburgh, B. in *Body size in mammalian paleobiology: estimation and biological*
implications (eds J. Damuth & Bruce J. MacFadden) Ch. 10, 181-206 (Cambridge
University Press, 1990).

607 Jones, K. E. *et al.* PanTHERIA: a species☐level database of life history, ecology, and
geography of extant and recently extinct mammals. *Ecology* **90**, 2648-2648 (2009).

608 Scharf, I. *et al.* Late bloomers and baby boomers: ecological drivers of longevity in
squamates and the tuatara. *Global Ecol. Biogeogr.* **24**, 396-405 (2015).

609 Regis, K. W. & Meik, J. M. Allometry of sexual size dimorphism in turtles: a comparison
of mass and length data. *PeerJ* **5**, e2914 (2017).

610 Gosler, A., Greenwood, J., Baker, J. & Davidson, N. The field determination of body size
and condition in passerines: a report to the British Ringing Committee. *Bird Study* **45**, 92-
103 (1998).

611 Rising, J. D. & Somers, K. M. The measurement of overall body size in birds. *The Auk*
106, 666-674 (1989).

612 Jetz, W., Thomas, G., Joy, J., Hartmann, K. & Mooers, A. The global diversity of birds in
space and time. *Nature* **491**, 444-448 (2012).

613 Wan, X., Wang, W., Liu, J. & Tong, T. Estimating the sample mean and standard
deviation from the sample size, median, range and/or interquartile range. *BMC Med. Res.*
Methodol. **14**, 135 (2014).

614 Bracken, M. in *Effective care of the newborn infant* (ed J.C. Sinclair and M.B. Bracken)
13-20 (Oxford University Press, 1992).

615 Pettorelli, N. *et al.* The Normalized Difference Vegetation Index (NDVI): unforeseen
successes in animal ecology. *Clim. Res.* **46**, 15-27 (2011).

616 Fick, S. E. & Hijmans, R. J. WorldClim 2: new 1☐km spatial resolution climate surfaces
for global land areas. *Int. J. Climatol.* **37**, 4302-4315 (2017).

617 Blackburn, T. M., Gaston, K. J. & Loder, N. Geographic gradients in body size: a
clarification of Bergmann's rule. *Divers. Distrib.* **5**, 165-174 (1999).

618 Wilman, H. *et al.* EltonTraits 1.0: Species☐level foraging attributes of the world's birds
and mammals: Ecological Archives E095☐178. *Ecology* **95**, 2027-2027 (2014).

619 Meiri, S. Traits of lizards of the world: Variation around a successful evolutionary
design. *Global Ecol. Biogeogr.* **27**, 1168-1172 (2018).

620 Benítez-López, A. *et al.* The impact of hunting on tropical mammal and bird populations.
Science **356**, 180-183 (2017).
621 Solé, M. & Rödder, D. in *Amphibian Ecology and Conservation: a handbook of*
techniques. (ed C. Kenneth Dodd Jr) 167-184 (Oxford University Press, 2010).
622 McClain, C. R., Durst, P. A. P., Boyer, A. G. & Francis, C. D. Unravelling the
determinants of insular body size shifts. *Biol. Lett.* **9**, 5pp.-5pp. (2013).
623 Rozzi, R. Space-time patterns of body size variation in island bovids: The key role of
predatory release. *J. Biogeogr.* **45**, 1196-1207 (2018).
624 Hadfield, J. & Nakagawa, S. General quantitative genetic methods for comparative
biology: phylogenies, taxonomies and multi◻trait models for continuous and categorical
characters. *J. Evol. Biol.* **23**, 494-508 (2010).
625 Lajeunesse, M. J. On the meta◻analysis of response ratios for studies with correlated and
multi◻group designs. *Ecology* **92**, 2049-2055 (2011).
626 McClain, C. R., Durst, P. A. P., Boyer, A. G. & Francis, C. D. Unravelling the
determinants of insular body size shifts. *Biol. Lett.* **9** (2013).
627 Lindstedt, S. L. & Boyce, M. S. Seasonality, fasting endurance, and body size in
mammals. *The American Naturalist* **125**, 873-878 (1985).
628 Nakagawa, S. & Schielzeth, H. A general and simple method for obtaining R2 from
generalized linear mixed◻effects models. *Methods Ecol. Evol.* **4**, 133-142 (2013).
629 R-Core-Team. *R: A language and environment for statistical computing.*, (R Foundation
for Statistical Computing, 2018).
630 Viechtbauer, W. Conducting meta-analyses in R with the metafor package. *J. Stat. Softw.*
36, 1-48 (2010).
631 Lajeunesse, M. J. Facilitating systematic reviews, data extraction and meta◻analysis with
the metagear package for R. *Methods Ecol. Evol.* **7**, 323-330 (2016).
632 Pick, J. L., Nakagawa, S. & Noble, D. W. Reproducible, flexible and high◻throughput
data extraction from primary literature: The metaDigitise r package. *Methods Ecol. Evol.*
10, 426-431 (2019).
633 Paradis, E., Claude, J. & Strimmer, K. APE: analyses of phylogenetics and evolution in R
language. *Bioinformatics* **20**, 289-290 (2004).
634 Michonneau, F., Brown, J. W. & Winter, D. J. rotl: an R package to interact with the
Open Tree of Life data. *Methods Ecol. Evol.* **7**, 1476-1481 (2016).
635 Rees, J. A. & Cranston, K. Automated assembly of a reference taxonomy for
phylogenetic data synthesis. *Biodivers. Data J.* (2017).
636 Hinchliff, C. E. *et al.* Synthesis of phylogeny and taxonomy into a comprehensive tree of
life. *Proc. Natl. Acad. Sci. USA* **112**, 12764-12769 (2015).
637 Pebesma, E. Simple features for R: standardized support for spatial vector data. *The R*
Journal **10**, 439-446 (2018).
638 Hijmans, R. J. *et al.* Package 'raster'. *R package* (2015).
639 Wickham, H., Francois, R., Henry, L. & Müller, K. dplyr: A Grammar of Data
Manipulation. R package version 0.4. 3. *R Found. Stat. Comput., Vienna. https://CRAN.*
R-project. org/package= dplyr (2015).
640 Wickham, H. reshape2: Flexibly reshape data: a reboot of the reshape package. *R*
package version **1** (2012).
641 Wickham, H. *ggplot2: elegant graphics for data analysis*. (Springer, 2016).

642   Kassambara, A. ggpubr:"ggplot2" based publication ready plots. R package version 0.1.
6. (2017).

2116

Acknowledgements

We are grateful to K. B. Aubry, J. E. Keehn, S. Michaelides and D. Strickland for sharing their

data with us, S. Meiri for pointing out valuable sources of measurement data, and to A. Sánchez-

Tojar, P. Peres-Neto and S. Nakagawa for useful discussion on the analytical framework. ABL

was supported by a Juan de la Cierva-Incorporación grant (IJCI-2017-31419) from the Spanish

Ministry of Science, Innovation and Universities. LS and MAJH were supported by the ERC

project (62002139 ERC – CoG SIZE 647224). We thank numerous biological collections, in

particular the Natural History Museum, Tring, for providing access to specimens. Bird trait data

collection was supported by Natural Environment Research Council grant nos. NE/I028068/1

and NE/P004512/1 (to JAT).

Author Contributions

2128 AB-L conceived and coordinated the research, led the analyses and wrote the first draft; AB-L,

2129 LS, JG-Z, MAJH and JAT helped to develop the conceptual framework; LS compiled the

2130 environmental rasters; JAT, PW and BM provided morphometric data. All authors contributed to

2131 the data collection from the literature and to the writing of the final manuscript.

Competing interests

2133 The authors declare no competing interests.

2134

Figures and Tables

Figure 1. Conceptual figure showing body size evolution in island populations. According to the island rule, changes in body size of island populations are dependent on the body mass of mainland relatives, with small species tending to increase in size on islands (gigantism) and large species tending to decrease in size (dwarfism). By plotting the log response ratio (*lnRR*) between insular mass and mainland mass, against mainland mass, we can test if insular populations adhere to the rule (intercept > 0 and slope < 0) (blue line). Mechanisms proposed to drive 'island rule' effects are mainly based on reduced predation, inter- and intra-specific competition, and food availability, suggesting that the relationship will steepen in small, remote islands (red line).

Figure 2. Location of island populations included in our analyses for mammals (N = 1058, blue), birds (N = 695, orange), reptiles (N = 547, yellow), and amphibians (N = 179, green). The size of each point indicates the number of species sampled on each island; some points overlap. See Fig. S1 for a 4-panel figure with the location of insular populations separated for each taxonomic group.

Figure 3. Relationship between *lnRR* (log response ratio between island mass and mainland body mass) and body mass in the mainland for (a) mammals (N = 1058), (b) birds (N = 695), (c) reptiles (N = 547) and (d) amphibians (N = 179). Models were fitted using phylogenetic multi☐level meta-regression models with mainland body mass as moderator, and observation-level ID, source ID, species ID and phylogeny as random effects. *lnRR* > 0 indicates gigantism; *lnRR* < 0 indicates dwarfism; and *lnRR* = 0 indicates stasis (no shift in body size from mainland to island populations). The size of points represents the inverse of the sampling variance for each paired island-mainland response ratio in the model. Shaded areas represent 95% confidence intervals. Note that y-axes have different scales.

Figure 4. The effect of island area and spatial isolation on insular size shifts in terrestrial vertebrates for (a) mammals (N = 1058), (b) birds (N = 695), (c) reptiles (N = 547) and (d) amphibians (N = 179). Continuous variables are represented at the 10% and 90% quantile for each extreme (close vs remote islands; small vs large islands). *lnRR* > 0 indicates gigantism; *lnRR* < 0 indicates dwarfism; and *lnRR* = 0 indicates stasis (no shift in body size from mainland to island populations). Shaded areas represent 95% confidence intervals.

2165 **Table 1.** Parameter estimates for the phylogenetic meta-regression models testing the generality
2166 of the island rule in terrestrial vertebrates. k: number of island-mainland comparisons ($lnRR$),
2167 Q_m: test of moderators ($\log_{10}$(mainland mass). R^2_m: marginal R^2, estimated percentage of
2168 heterogeneity explained by the moderator (fixed effects). R^2_c: conditional R^2, percentage of
2169 heterogeneity attributable to fixed and random effects.

2170

Class	k	Intercept (CI)	Slope (CI)	Q_m (p-value)	R^2_m	R^2_c
Mammals	1058	0.208 (0.052 – 0.365)	-0.088 (-0.122 – -0.055)	27.30 (p < 0.001)	11.4	56.4
Birds	695	0.216 (0.117 – 0.315)	-0.104 (-0.145 – -0.064)	25.40 (p < 0.001)	7.0	43.4
Reptiles	547	0.410 (0.006 – 0.814)	-0.305 (-0.419 – -0.190)	27.21 (p < 0.001)	17.6	66.6
Amphibians	179	0.195 (0.012 – 0.377)	-0.107 (-0.320 – 0.107)	0.96 (p = 0.328)	1.4	67.1

2171

2172

2173

2174

2175

2176

$$RR = \log\left(\frac{\text{Insular species mass}}{\text{Mainland species mass}}\right)$$

Low resource availability
No competitors
No predators

No predators
No competitors