Visual gene expression reveals a cone to rod developmental progression in deep-sea fishes

Nik Lupše[1*], Fabio Cortesi[2], Marko Freese[3], Lasse Marohn[3], Jan-Dag Pohlman[3], Klaus Wysujack[3], Reinhold Hanel[3], Zuzana Musilova[1*]

1) Department of Zoology, Faculty of Science, Charles University, Vinicna 7, 12844 Prague, Czech Republic

2) Queensland Brain Institute, University of Queensland, Brisbane 4072 QLD, Australia

3) Thünen Institute of Fisheries Ecology, Herwigstraße 31, 27572, Bremerhaven, Germany

* corresponding authors; lupsen@natur.cuni.cz, zuzana.musilova@natur.cuni.cz

Keywords: opsin, evolution, mesopelagic, adaptation, convergence, phototransduction, vision, gene expression, rhodopsin

Abstract

Vertebrates use cone cells in the retina for colour vision and rod cells to see in dim light. Many deep-sea fishes have adapted to their environment to have only rod cells in the retina, while both rod and cone genes are still preserved in their genomes. As deep-sea fish larvae start their lives in the shallow, and only later submerge to the depth, they have to cope with diverse environmental conditions during ontogeny. Using a comparative transcriptomic approach in 20 deep-sea fish species from eight teleost orders, we report on a developmental cone-to-rod switch. While adults mostly rely on rod opsin (*RH1*) for vision in dim light, larvae almost exclusively express middle-wavelength-sensitive ("green") cone opsins (*RH2*) in their retinas. The phototransduction cascade genes follow a similar ontogenetic pattern of cone- followed by rod-specific gene expression in most species, except for the pearleye and sabretooth (Aulopiformes), in which the cone cascade remains dominant throughout development. By inspecting the whole genomes of five deep-sea species (four of them sequenced within this study: *Idiacanthus fasciola, Chauliodus sloani*; Stomiiformes; *Coccorella atlantica,* and *Scopelarchus michaelsarsi*; Aulopiformes), we found that deep-sea fish possess one or two copies of the rod *RH1* opsin gene, and up to seven copies of the cone *RH2* opsin genes in their genomes, while other cone opsin classes have been mostly lost. Our findings hence provide

34   molecular evidence for a limited opsin gene repertoire and a conserved vertebrate pattern

35   whereby cone photoreceptors develop first and rod photoreceptors are added only at later

36   developmental stages.

INTRODUCTION

Vision is a primary sense used by most vertebrates for navigation, predator avoidance, communication and to find food and shelter. At its initiation, vertebrate vision is enabled by cone (photopic, colour vision) and rod (scotopic) photoreceptors in the retina containing a light absorbing pigment that consists of an opsin protein covalently bound to a vitamin-A-derived chromophore (Lamb 2013). The absorbance of photons by the chromophore leads to a conformational change of the opsin protein, which initiates a photoreceptor-specific G-protein-coupled phototransduction cascade, propagating the signal to the brain (Downes and Gautam 1999, Larhammar et al. 2009, Lamb et al. 2019). It is thought that the development of the visual system follows a conserved molecular pattern whereby cone specific genes are activated first before the rod molecular pathway is initiated later during ontogeny (Mears et al. 2001, Shen and Raymond 2004, Sernagor et al. 2006). However, whether this is the case for all vertebrates and especially for those that have retinas that contain only rods as adults, remains unclear.

Changes in the light environment, ecology, and phylogenetic inertia are thought to be primary drivers for visual system diversity in vertebrates (Hunt et al. 2014). For example, most mesopelagic deep-sea fishes (200 – 1,000 m depth), either living strictly at depth or migrating to the shallows at night, have evolved visual systems that are sensitive to the dominant blue light (~ 470 – 490 nm) of their environment (Turner et al. 2009). Moreover, as the daylight and the bioluminescent light emitted by deep-sea critters are quickly dimmed with depth and distance, deep-sea fish visual systems have evolved peculiar morphologies to maximise photon capture including barrel-eyes, reflective tapeta and the use of rod-dominated and in many cases rod-only retinas that might be stacked into multiple banks (reviewed in de Busserolles et al. 2020). However, most mesopelagic fishes start their lives in the shallow well-lit epipelagic zone (0 – 200 m depth) (Moser and Smith 1993, Sassa and Hirota 2013). Consequently, their visual systems must cope with a variety of light intensities and spectra throughout development.

Studies investigating the gene expression in the retina of deep-sea fishes are scarce and usually focus on a selected few species (Zhang et al. 2000, Douglas et al. 2016, de Busserolles et al. 2017, Musilova et al. 2019a, Byun et al. 2020). In adults, species with pure rod retinas tend to only express rod opsin(s) (Douglas et al. 2016, Musilova et al. 2019a), albeit two species of pearlsides (*Maurolicus* spp.) have been found to express cone-specific genes (i.e., cone transduction pathway and opsin genes) inside rod-looking cells (de Busserolles et al. 2017). It remains unknown whether deep-sea fishes that have a low proportion of cone photoreceptors as adults (e.g., Munk 1990, Collin et al. 1998, Bozanno et al. 2007, Pointer et al. 2007, Biagioni

et al. 2016) also express cone-specific genes at any stages of their lives or whether these fishes
rely on the rod machinery alone. To investigate whether the retinal development in deep-sea
fishes follows a similar cone-to-rod molecular pathway as found in other vertebrates or whether
some species start their lives with the rod pathway activated, we set out to sequence the retinal
transcriptomes of 20 deep-sea fish species, including the larval stages in ten species, belonging
to eight different teleost orders (Argentiniformes, Aulopiformes, Beryciformes,
Myctophiformes, Pempheriformes, Scombriformes, Stomiiformes and Trachichthyiformes).
We have further investigated the genomic repertoire in five selected species.
79
RESULTS AND DISCUSSION
Opsin gene repertoire in the genome. In teleost fishes, gene duplications and deletions
followed by functional diversification have resulted in extant species having between 1-40
visual opsin genes within their genomes (Musilova et al. 2019a, Musilova et al. 2021). These
genes are defined by their photoreceptor specificity, their phylogeny, and their spectrum of
maximal sensitivity (λ_{max}) and fall within five major classes, four cone opsins ('ultraviolet or
UV sensitive' *SWS1*: 347–383 nm, 'blue' *SWS2*: 397–482 nm, 'green' *RH2*: 452–537 nm and
'red' *LWS*: 501–573 nm) and one rod opsin ('blue-green' rhodopsin, *RH1* or Rho: 447–525
nm) (Carleton et al. 2020). We analysed the whole genomes of five deep-sea species (sawtail
fish *Idiacanthus fasciola*, viperfish *Chauliodus sloani*; both Stomiiformes; sabretooth
Coccorella atlantica, pearleye *Scopelarchus michaelsarsi*; both Aulopiformes; and fangtooth
Anoplogaster cornuta; Trachichthyiformes), four of them sequenced for the purpose of this
study. All species possess one or two copies of the rod opsin *RH1* gene, and one to seven copies
of the *RH2* cone opsin (Fig. 1). All other cone opsin classes, i.e., the *SWS1*, *SWS2* (except for
the fangtooth) and *LWS* are missing and have been putatively lost during evolution in these
five species. This is in accordance with the observation that the *LWS* gene abundance decreases
with the habitat depth (Musilova et al. 2019a). Such limited genomic repertoire most likely
represents an evolutionary response to the deep-sea scotopic environment where the shortest
(UV-violet) and longest (red) wavelengths of light get filtered out first in the water column, as
opposed to middle-range wavelengths that can penetrate greater depths (reviewed in Musilova
et al. 2021, De Busserolles et al. 2020, and Carleton et al. 2020). The increased *RH2* diversity
observed in the two aulopiform species, on the other hand, illustrates the versatility of this cone
opsin class and confirms its dominance in various dimmer-light habitats (Musilova and Cortesi
2021). Here we confirm that *RH2* is undoubtedly the most important (and often the only) cone
opsin gene present in deep-sea fish genomes.

Visual opsin gene expression. Transcriptomic sequencing of 20 deep-sea teleost species revealed that deep-sea fishes mainly express rod opsins and/or green-sensitive cone opsins (*RH2*s) in their retinas (Fig. 1, Table 1). While larvae mostly expressed *RH2*, adults and juveniles mostly expressed *RH1* and in a few cases a combination of both. We found none or very low expression of any of the other cone opsin genes: the red sensitive *LWS* was not expressed at all, the UV sensitive *SWS1* was only found in the larva of the whalefish, *Gyrinomimus* sp. (Beryciformes), and the blue/violet sensitive *SWS2* only in the larvae of the whalefish, and the fangtooth, *Anoplogaster cornuta* (Trachichthyiformes)*,* (Fig. 1, Table 1). Differences in gene expression patterns are likely to be driven by ontogenetic transitions in light habitat from bright to dim environments and by changes in ecological demands, as discussed in more detail below.

Similar to the opsin genes, we also detected ontogenetic differences in the expression of phototransduction cascade genes (Fig. 2). Here we focused on the comparison of five species from three teleost orders for which we had both larval and adult specimens available and found that the cone-specific genes were mostly expressed in the larval stages (e.g., cone transducin, *GNAT2*), while adults from three species mostly expressed rod-specific genes (e.g., rod transducin, *GNAT1*; Fig. 2b). Hence, at the molecular level, the visual systems of deep-sea fishes start out with a cone-based expression pattern. Moreover, in the fangtooth, where samples from various sized specimens were available, we found that the cone-specific expression was gradually replaced with the rod profile as the fish grew (Fig. 2c, Table S1). This sequence is similar to the visual development in shallower living fishes (e.g., Atlantic cod (Valen et al. 2016), zebrafish (Sernagor et al. 2006)) and terrestrial vertebrates (e.g., mice (Mears et al. 2001), rhesus monkey (La Vail et al. 1991)), where cone photoreceptors are first to develop, followed by temporally and spatially distinct rods (Raymond 1995, Shen and Raymond 2004). The cone-to-rod developmental sequence is therefore likely to be shared across vertebrates, even in species that have pure rod retinas as adults.

Ontogenetic shift in expression profiles and the transition phase. The observed developmental changes in the visual system are best explained by the different habitats larval and adult deep-sea fishes inhabit. In general, deep-sea fish larvae live in the shallow epipelagic zone (Moser and Smith 1993) where ambient light levels are sufficiently high to warrant a cone-based visual system. After metamorphosis, deep-sea fishes start to submerge deeper and take up a life at different depths in the mesopelagic or even bathypelagic (below 1,000 m depth) zone, where the sun- and moonlight is gradually replaced by bioluminescence as the main

source of light (Denton 1990). In this extremely dim environment, rods work at their best and

cone photoreceptors would be obsolete for the most part at least. Rod-based vision is also

favoured in those deep-sea species that exhibit diel vertical migrations to feed in the plankton

rich surface layers at night (de Busserolles et al. 2020). In addition, we discovered that in some

species there was a switch in the expressed type of cone *RH2* opsin (Fig. 1). For example, in

Aulopiformes, the larvae expressed an alternative *RH2* copy that is presumably sensitive to

longer wavelengths of light compared to the *RH2* that was found in adults (Table 2). This

clearly shows that larval and adult deep-sea fishes rely on different opsin expression profiles,

which is similar to ontogenetic changes in opsin gene expression in diurnal shallow-water

fishes such as freshwater cichlids (Carleton et al. 2016) and coral reef dottybacks (Cortesi et

al. 2015, Cortesi et al. 2016) or between the freshwater and deep-sea maturation stages in eels

(Zhang et al. 2000).

Our data furthermore suggest that the ontogenetic change in visual gene expression

precedes morphological changes such as metamorphosis from larva to juvenile and also habitat

transitions. For example, in the fangtooth, the larvae which were collected from the shallows

(0 – 300 m) showed increasing amounts of *RH1* expression with growth, despite displaying

larval phenotypes throughout (horns and small teeth; Fig. 1). A similar pattern of changing

opsin gene expression ahead of metamorphosis has also been reported from shallow-water

fishes such as European eels (Bowmaker et al. 2008), dottybacks (Cortesi et al. 2016) and

surgeonfishes (Tettamanti et al. 2019). Interestingly, all our fangtooth larvae (including the

smallest individual with a total length of 4 mm) already expressed a small amount of *RH1* (Fig.

2c). Whether fangtooth start their lives with a pure cone retina or low-levels of rod opsin

expression are normal even in pre-flexation larvae remains therefore unclear. In addition to the

green-sensitive cone opsin *RH2*, the smallest fangtooth larvae also expressed low levels of the

blue-sensitive *SWS2*, potentially conferring dichromatic colour vision to the early life stages of

this species (Fig. 1).

Photoreceptor cell identities. Interestingly, two aulopiform species; the Atlantic sabretooth,

Coccorella atlantica, and the Bigfin pearleye, *Scopelarchus michaelsarsi*, despite expressing

mostly *RH1* as adults, retained a cone-dominated phototransduction cascade expression profile

akin to the one found in the larval stages (Fig. 2, Table S1). This begs the question whether the

photoreceptors they are using are cones or rods in nature. Initially described in snakes and

geckos (Simoes et al. 2016, Schott et al. 2019) and recently also in a deep-sea fish (de

Busserolles et al. 2017), it appears that the dichotomy of rods and cones is not always as clear

cut as one might think. For example, adult deep-sea pearlsides, *Maurolicus* spp. have a retina
that expresses ~ 99% *RH2* and ~ 1% *RH1* with corresponding cone and rod phototransduction
gene expressions. Their photoreceptors, however, are all rod-shaped and careful histological
examination has shown that these consist of a tiny proportion of true rods and a majority of
transmuted rod-like cones (de Busserolles et al. 2017). In the case of pearlsides, and also in
geckos and snakes, the opsin and phototransduction genes correspond to each other making it
possible to distinguish photoreceptor types at the molecular level. However, in the aulopiforms,
high expression of rod opsin is seemingly mismatched with high levels of cone
phototransduction gene expression (Fig. 2). In salamanders, the opposite pattern can be found
whereby a cone opsin is combined with the rod phototransduction cascade inside a rod looking
cone photoreceptor (Mariani 1986). Anatomically, the retina of *S. michaelsarsi* is composed of
mostly rods with low numbers of cone cells (Collin et al. 1998), while the adult retina of
Evermanella balbo, an evermannellid species related to *C. atlantica,* appears to consist of two
differently looking rod populations (Wagner et al. 2019). It is therefore likely, as found in
pearlsides (de Busserolles et al. 2017), that these fishes have a high proportion of transmuted
rod-like cone photoreceptors, but that they use *RH1* instead of a cone opsin as the visual
pigment. Alternatively, a proportion of true rods might make use of the cone phototransduction
cascade. Either way, combining more stable rod opsin in a rod-shaped cell with the cone-
specific cascade is likely to increase sensitivity while also maintaining high transduction and
recovery speeds of cells (Baylor 1987, Kawamura and Tachibanaki 2012, Luo et al. 2020).
Histology, fluorescent in-situ hybridisation and ideally physiological recordings are needed to
ultimately disentangle the identity of photoreceptor cells in aulopiforms.
195
Evolutionary history of deep-sea fish opsins. While the majority of adult fishes relied on a
single *RH1* copy, we found three species that expressed multiple *RH1* copies: The Warming's
lanternfish, *Ceratoscopelus warmingii* (Myctophiformes), expressed three different *RH1*
genes, and *S. michaelsarsi* and the basslet, *Howella brodiei* (Pempheriformes), expressed two
copies each. Larvae and a few adult deep-sea fishes mostly expressed a single *RH2* copy, except
for the pearleyes, *Scopelarchus* spp., and the Reinhardt's lanternfish, *Hygophum reinhardtii*,
which expressed three larval copies each, and the whalefish (*Gyrinomimus* sp.) which
expressed five larval copies (Fig. 1).
204   The *RH1* and *RH2* phylogenies revealed that most deep-sea fish visual opsins cluster
together by species or order (Fig. 3). For example, in the whalefish all *RH2s* are clustered
together suggesting that these genes are lineage or species-specific duplicates (Fig. 3b).

However, there were a few exceptions, suggesting more ancient duplication events. In
Scopelarchus the two *RH1* copies are not in a sister relationship and in fact result in different
clusters, suggesting that these copies originated in the common ancestor of aulopiforms or
perhaps even earlier (Fig. 3a). Also, the *RH2s* in aulopiforms (*Scopelarchus*, *Coccorella*)
cluster by ontogenetic stage, making it likely that the developmental switch in gene expression
was already present in the aulopiform ancestor (Fig. 3b).

Molecular complexity of deep-sea fish visual systems. The complexity of the deep-sea fish
visual systems at the molecular level varied quite substantially. For example, the three
Stomiiformes species: The Ribbon sawtail fish, *Idiacanthus fasciola*, and two species of
viperfish, *Chauliodus sloani* and *Ch. danae*, appeared to have a very basic visual set up; these
fishes were found to express a single *RH2* cone opsin as larvae and a single *RH1* rod opsin as
adults (Fig. 1). On the contrary, several deep-sea fish orders examined here expressed more
than one opsin gene. Adult lanternfishes and basslets have rod-only retinas but expressed
multiple *RH1* copies that have functionally diversified (Fig. 3). Other species expressed both
cone and rod opsins as adults (the aulopiform species and *Scombrolabrax*), which is somewhat
similar to the opsin gene expression profiles found in shallow-living nocturnal reef fishes
(Cortesi et al. 2020).
The most complex visual system in this study was found in *S. michaelsarsi*. In general,
this species is known for its numerous morphological and anatomical adaptations to vision in
the depth, including having barrel eyes with a main and an accessory retina, rods that are
organised in bundles, large ganglion cells and corneal lens pads (Collin et al., 1998). The two
copies of *RH1* (*RH1a* and *RH1b*) it expressed showed high sequence diversity differing in 79
out of 354 amino acids, eight of which are known key tuning sites likely to change the spectral
sensitivity of the pigments via a shift in λ_{max} (Fig. 3, Table 3) (Yokoyama 2008, Musilova et
al. 2019a, Yokoyama and Yia 2020). This supports the findings by Pointer et al. (2007) who
found by using in vitro expression of pigments in another pearleye species, *S. analis*, two rod
photoreceptors with different absorption maxima at 479 and 486 nm. Interestingly, Pointer et
al. (2007) also speculate that another short-shifted opsin (previously measured in *Sc. analis* to
have λ_{max} at 444 nm by Partridge et al., 1992) possibly belongs to the SWS2 class. Our data
however does not support this expectation as no SWS2 gene is found in the genome of
Scopelarchus michaelsarsi. The existence and identity of such short-sensitive opsins in
pearleyes remains therefore elusive. The situation is less clear for the green-sensitive *RH2*
opsin. While in *S. analis* cones have been found in the accessory and main retinas, in *S.*

michaelsarsi cone photoreceptors appear restricted to the accessory retina alone (Collin et al.
1998). This is intriguing as it suggests substantial differences in visual systems even between
closely related species from the same genus.
244
The visual ecology of deep-sea fishes. We found molecular support for deep-sea visual
adaptations on multiple levels:
1) Opsin gene diversity in the genome. Data from the genomes of five deep-sea species revealed
the diversity of the opsin genes (Fig. 1C). Retinal transcriptomes in the Stomiiformes pointed
towards a simple visual system that is based on a single expressed opsin gene at different
developmental stages (*RH2* in larvae, *RH1* in adults) (Fig. 1). A search for visual opsins in the
stomiiform genomes sequenced for the purpose of this study (*I. fasciola* and *Ch. sloani*), as
well as in several published genomes (Musilova et al., 2019a) revealed that Stomiiformes are
likely to have lost all cone opsin gene families except for *RH2* (Fig. 1). In the case of *RH2*,
they seem to only have a single or at most two gene copies, which is substantially less than
other teleosts (Musilova et al. 2019a; Musilova and Cortesi, 2021). The stomiiform example,
therefore, shows that a decrease in light intensity and the spectrum of light in the deep-sea may
not only restrict gene expression at adult stages, but also lead to the loss of opsin genes
altogether. Similarly, a loss in opsin and other vision-related genes (e.g., otx5b, crx) has
previously been reported from shallow living fishes that are either nocturnal (Luehrmann et al.
2019), live in murky waters (Liu et al. 2019), or inhabit caves or similarly dim environments
(Huang et al. 2019, Musilova et al. 2019a). Contrarily, the genomes of two aulopiform species
(*C. atlantica* and *S. michaelsarsi*) revealed expanded cone opsin gene repertoires achieved
mostly through *RH2* duplications (*C. atlantica* with three *RH2*, and *S. michaelsarsi* with seven
RH2 genes; Fig 1). Both species also possess two copies of the rod opsin gene. These species
inhabit relatively shallower depths (300-400m) compared to other deep-sea fishes such as the
Stomiiformes (Table 1). It is likely that having extra copies of *RH2* cone opsins may benefit
their vision at these photon-richer depths. It is also possible that an evolutionary stochasticity
and the gene content in the ancestor have contributed to the observed pattern. To be able to
clearly state this, future research should be done on multiple aulopiform species.
270
2) Visual gene expression. Previous work has found that the expression of the longest- (*LWS* -
red) and shortest- (*SWS1* - UV) sensitive opsins is reduced or absent in deeper living coral reef
fishes (Cortesi et al. 2020) and in fishes inhabiting deep freshwater lakes (e.g., Hunt et al. 1997,
Sugawara et al. 2005, Musilova et al. 2019b), which is correlated with a loss of short- and long-

wavelengths with depth. Also, many deep-sea fish lineages have lost *LWS* from their genomes

(Musilova et al. 2019a, Cortesi et al. 2021). Supporting these findings, we show here that deep-

sea fishes lack any *LWS* expression even in the shallow-living larval stages (Fig. 1). Similarly,

SWS1 is not expressed in any of the species studied, except for in the larval whalefish, and is

also absent from many deep-sea fish genomes (Fig. 1) (Musilova et al. 2019a). However,

shallow larval stages are likely to explain why all deep-sea fishes studied to date maintain at

least some cone opsins in their genomes (Musilova et al. 2019a).

282   Most deep-sea fish larvae expressed a single *RH2* gene, but the larvae of some species

(fangtooth*,* whalefish, and lanternfish) expressed multiple cone opsin genes (Fig. 1). This is

likely to provide them with similar visual systems to the larvae of shallow-living marine (Britt

et al. 2001) and freshwater species (Carleton et al. 2016), possibly aiding in detecting residual

light and discriminating brightness and/or colours. Juvenile deep-sea fishes, on the other hand,

showed rod-based expression profiles also found in the adult stages (Fig. 1). This shift in opsin

gene expression correlates with developmental changes in ecology. As opposed to the adults

which are exposed to a narrow and dim light environment where food is scarce, larvae typically

live in well-lit, broad spectrum shallow waters where food and predators are abundant (Moser

and Smith 1993).

3) Functional adaptation in key spectral tuning sites. When multiple *RH1* copies were

expressed, they often showed distinct differences in key amino acid sites that are likely to shift

the spectral sensitivities of the pigments (Fig. 3 and Table 3) (Yokoyama 2008, Musilova et al.

2019a). Estimating spectral absorbance from these sites resulted in very similar λ_{max} values to

those that were measured in vivo from the same or closely related species using

microspectrophotometry (MSP) or similar techniques (Partridge et al. 1989, Collin and

Marshall 2003, Pointer et al. 2007) (Fig. 3, Table 3). We therefore integrated the available

functional evidence (MSP values) with our molecular data (protein sequence and gene

expression levels) to better understand the function of the visual system in the three species

that expressed multiple rod opsins (Fig. 3, Table 3).

303   The three different *RH1*s in *C. warmingii* differed in 15 key-tuning sites. Our data

revealed a dominant rod opsin copy (*RH1-1*), and a shorter- (*RH1-2*) and a longer-shifted (*RH1-*

3) copies with lower transcript levels (Figs. 1 and 3). The MSP in *C. warmingii* revealed two

distinct rod types with λ_{max} values of 488 and 468 nm in past (Collin and Marshall 2003)

corresponding most likely to the *RH1a/RH1-1* and *RH1b/RH1-3* genes, respectively. It is

possible that multiple *RH1* copies are coexpressed within the same photoreceptor, something

that has previously been reported for cone opsins in shallow-water marine (Savelli et al. 2018,

Stieb et al. 2019) and freshwater fishes (Dalton et al. 2014, Torres-Dowdall et al. 2017).

Coexpression could produce visual pigment mixtures that shift photoreceptor sensitivity and

enhance visual contrast, aiding in predator-prey interactions or mate detection (Dalton et al.

2014). Alternatively, we predict that a third rod photoreceptor type with longer spectral

sensitivity (*RH1-2*; Fig 3c) exists, possibly overlooked during MSP, which can happen

especially if the cell type is rare. While the function of having multiple rod opsins in *C.*

warmingii remains to be investigated, several possible benefits for a multi-rod visual system

have recently been proposed including that it might enable conventional or unusual colour

vision in dim light, it might be used to increase visual sensitivity, or enhance an object's

contrast against a certain background (Musilova et al. 2019a).

In *H. brodiei*, the second *RH1* copy (*RH1-2*) differed in two key-tuning sites, E122Q (-

15 nm) and G124S (-11 nm), known to cause major short-wavelength shifts in other fishes

(Yokoyama 2008). This is in accord with the MSP measurements in its sister species, *Howella*

sherborni, which found two different rod types with spectral sensitivities of 463 and 492 nm

(Fig. 3) (Partridge et al. 1989). Having multiple differently tuned rod photoreceptors, one

centred on the prevailing light (bioluminescence and/or ambient light ~ 480 – 490 nm) and a

second one that is offset from it (i.e., the offset pigment hypothesis; Lythgoe 1966), may be

beneficial to break counter illumination of prey - a way of active camouflage in mesopelagic

organisms where ventral photophores emit bioluminescent light that matches the residual

down-welling light (Denton et all. 1985). Hence, revealing an individual's silhouette could help

to distinguish prey and predators from the background lighting, or visually finding mates.

However, apart from lanternfishes with three (or more) and basslets with two rod opsins, or

exceptional cases of tube-eye (six) and spinyfin (38), the majority of the deep-sea fishes seem

to have only one rod opsin (Musilova et al., 2019a).

Differences in spectral sensitivity between photoreceptors can also be due to the

chromophore type that binds to the opsin protein; visual pigments with a vitamin A1-based

chromophore (typical in marine fishes) confer shorter-shifted λ_{max} values compared to those

with a vitamin A2-based chromophore (typical in freshwater fishes) (Carleton et al. 2016).

Cyp27c1 is the enzyme responsible for converting A1- to A2-based chromophores (Enright et

al. 2015), with high expression levels suggesting the presence of longer-shifted visual

pigments. However, *Cyp27c1* was not expressed in our dataset suggesting that the visual

pigments of these deep-sea fishes are based on A1-retinal alone (Table S2).

Conclusions

So far, the development of deep-sea fish vision at the molecular level had not been studied in detail and only limited morphological information is available. In this study we compared opsin and visual gene expression between 20 deep-sea fish species revealing a major change in expression between larval and adult stages. While deep-sea fish genomes contain both cone and rod opsin genes, larvae rely on the cone pathway and most adult fishes switch to a rod-dominated or rod-only visual system. The cone- versus rod-specific phototransduction cascade genes follow the opsins in some lineages, however, not in aulopiforms. We detected reduced opsin gene repertoires in the genomes of five deep-sea fish species composed only of one rod (*RH1*) and one or two cone (*RH2*, or *RH2* and *SWS2*) opsin gene classes. Interestingly, we have discovered lineage-specific opsin gene duplications, possibly allowing for increased visual sensitivity and certain kind of colour vision in the depth in some species. Overall, our molecular results support a conserved developmental progression in vertebrates whereby cones appear first in the retina and rod photoreceptors are added later during development.

MATERIALS AND METHODS

Specimens used in this study were collected in the Sargasso Sea during three multipurpose fishery surveys conducted by the German Thünen Institute of Fisheries Ecology onboard the research vessels *Maria S. Merian* in March to April in 2015, and *Walther Herwig III* in 2017 and in 2020. The sampling of adults occurred during both day and night at depths of 600 – 1'000 m using a mid-water pelagic trawl net (Engel Netze, Bremerhaven, Germany) with an opening of 30 m x 20 m, a length of 145 m, and mesh sizes (knot to knot) from 90 cm decreasing stepwise to 40, 20, 10, 5, 4, 3, 2 cm, with a 1.5-cm mesh in the 27-m-long codend. The larvae were mostly collected using an Isaacs-Kidd Midwater Trawl net (IKMT; 6.2 m^2 mouth-opening, 0.5 mm mesh size; Hydro-Bios Apparatebau GmbH) at depths of 0 - 300 m by double-oblique transect tows. Adult fish were flash-frozen at -80 ^{0}C upon arrival on board and their fin clip was stored in 96% ethanol. Larval samples were fixed in RNAlater^{TM} (ThermoFisher) and stored at -80 ^{0}C until further use.

To sequence the whole genome of *Idiacanthus fasciola, Chauliodus sloani, Coccorella atlantica,* and *Scopelarchus michaelsarsi,* the genomic DNA was extracted from the fin clip using the DNeasy Blood and Tissue kit (Qiagen) following the enclosed protocol. The library preparation and genome sequencing on Illumina NovaSeq platform (150 bp PE and the yield over 20 Gb per genome) has been outsourced to the sequencing centre Novogene, Singapore (https://en.novogene.com/). To analyse the opsin gene repertoire, the raw genomic reads were mapped in Geneious software version 11.0.3 (Kearse et al. 2012) against the opsin references (single exons of all five opsin classes from the reference species: Nile tilapia, Round goby, Blind cavefish, Spotted gar), as well as against the genes found in the transcriptomes of each species. The parameters were set to the Medium Sensitivity to capture all reads that matched any visual opsin gene. The captured reads mapping to all exons were then remapped against one reference per exon and the species-specific consensus sequence was generated. If present, multiple paralogous genes were disentangled manually, and the consensus sequence was exported for each variant (see below more details for the transcriptomic analysis). The obtained consensus sequences served as references for the second round of mapping, whereby all genomic reads were again mapped with the Low Sensitivity settings, and each reference was then elongated by the overlapping sequence. This step was repeated until the full gene region was covered. In case of *Scopelarchus michaelsarsi*, we were not able to cover the full length of five out of seven *RH2* genes due to the repetitions and these genes were reported in two parts, always one covering the exons 1 and 2, and one covering exons 3, 4 and 5. The genomic

raw reads are available from GenBank (BioProject PRJNA754116) and the opsin gene
sequences are provided in Supplementary file 1.

Total RNA was extracted from the whole eyes using either the RNeasy micro or mini
kit (Qiagen) and the extracted RNA concentration and integrity were subsequently verified on
a 2100 Bioanalyzer (Agilent). RNAseq libraries for 31 samples were constructed in-house from
unfragmented total RNA using Illumina's NEBNext Ultra II Directional RNA library
preparation kit, NEBNext Multiplex Oligos and the NEBNext Poly(A) mRNA Magnetic
Isolation Module (New England Biolabs). Multiplexed libraries were sequenced on the
Illumina HiSeq 2500 platform as 150 bp paired-end (PE) reads. Library construction and
sequencing (150 bp PE) for an additional 10 samples was outsourced to Novogene, Singapore
(https://en.novogene.com/). We additionally re-analysed 11 retinal transcriptomes previously
published in Musilova et al. (2019a). Together, then, our dataset comprised 53 samples of
which, based on morphology, 26 were classified as larvae, 6 as juveniles and 21 as adults.
Sample IDs, number of raw reads, individual accession numbers for BioProject PRJNA754116
and further parameters are listed in Table 1.

The sequence data was quality-checked using FastQC (Andrews 2010). Opsin gene
expression was then quantified using Geneious software version 11.0.3 (Kearse et al. 2012).
For each sample we first mapped the reads against a general fish reference dataset comprising
all visual opsin genes from the Nile tilapia, *Oreochromis niloticus* and the zebrafish, *Danio*
rerio, with the Medium-sensitivity settings in Geneious. This enabled us to identify cone and
rod opsin specific reads. If present, paralogous genes were subsequently disentangled following
the methods in Musilova et al. 2019a and de Busserolles et al. 2017. Briefly, we created species-
specific references of the expressed opsin genes and their several copies (Musilova et al. 2019a)
and re-mapped the transcriptome reads with Medium-Low sensitivity to obtain copy-specific
expression levels. If multiple opsin genes were found to be expressed, we report their
proportional expression in relation to the total opsin gene expression (Fig. 1). We used the same
pipeline to quantify expression of phototransduction cascade genes in five focal deep-sea
species (Fig. 2, Table S1), and to search for the expression of the *cyp27c1* gene (Table S2).

To analyse key amino-acid substitutions in *RH1* and *RH2* and potential shifts in their
absorbance, we first translated the opsin coding sequences into amino acid sequences, and then
aligned them with the bovine *RH1* (GenBank Acc.No: M12689). We have specifically focused
on the positions identified as key-tuning sites in Yokoyama (2008) and Musilova et al. (2019a).
For details, see Tables 2 and 3. Unfortunately, we were not able to estimate the sensitivity shift
of rod opsin copies in *C. warmingii* as only four of the amino acids that were substituted at the

15 key-tuning amino acid sites corresponded with previously tested cases (Yokoyama, 2008,

Musilova et al. 2019a). Out of the three copies, *RH1-2* has three out of four longer-shifting

amino acid variants in these four sites and we assume it is therefore red-shifted. *RH1-1* is most

likely sensitive to 488 nm, and *RH1-3*, being the shortest, to 468nm (Fig. 3c).

A dataset containing *RH1* opsin gene sequences mined from our dataset and additional

*RH1*s obtained from GenBank (GenBank accession numbers listed in Figure 3), were aligned

using the MAFFT (Katoh et al. 2009) plugin as implemented in Geneious, and a phylogenetic

tree was subsequently reconstructed using MrBayes v3.2.1 (Ronquist and Huelsenbeck 2003)

(Fig. 3a). Trees were produced using the Markov chain Monte Carlo analysis which ran for 1

million generations. Trees were sampled every 100 generations, and the printing frequency was

1000, discarding the first 25% of trees as burn-in. The evolutionary model chosen was GTR

model with gamma-distributed rate variation across sites and a proportion of invariable sites.

Posterior probabilities (PP) were calculated to evaluate statistical confidence at each node. We

used the same approach with an *RH2*-specific reference dataset to reconstruct the phylogenetic

relationship between the transcriptome-derived deep-sea *RH2* genes (Fig. 3b).

Acknowledgements:

We would like to express our thanks to both scientific and technical crew of the Maria S.

Merian and Walther Herwig III research cruises in 2015, 2017 and 2020. In addition, we thank

Tina Blancke for help with the sample management, and Veronika Truhlářová for technical

support and lab management. We would also like to thank three anonymous reviewers for their

comments improving the final version of the manuscript. NL and ZM were supported by the

Swiss National Science Foundation (PROMYS - 166550), ZM by the PRIMUS Research

Programme (Charles University), the Czech Science Foundation (21-31712S) and the Basler

Stiftung fuer Experimentelle Zoologie, and FC by an Australian Research Council (ARC)

DECRA Fellowship (DE200100620).

References:

Andrews S. 2017. FastQC: a quality control tool for high throughput sequence data. 2010.

Baylor DA. 1987. Photoreceptor signals and vision. Proctor lecture. Investigative
ophthalmology & visual science 28.1

Betancur-R R., et al. 2017. Phylogenetic classification of bony fishes. BMC evolutionary
biology, 17(1), 162.

Biagioni LM, Hunt DM, Collin SP. 2016. Morphological characterization and topographic
analysis of multiple photoreceptor types in the retinae of mesopelagic hatchetfishes with
tubular eyes. Frontiers in Ecology and Evolution, 4, 25.

Britt LL, Loew ER, McFarland N. 2001. Visual pigments in the early life stages of Pacific
northwest marine fishes. Journal of Experimental Biology 204.14.

Bowmaker JK, Hunt DM, Jeffery G. 2008. Eel visual pigments revisited: The fate of retinal
cones during metamorphosis. Visual neuroscience, 25(3), 249.

Bozzano A, Pankhurst PM, Sabatés A. 2007. Early development of eye and retina in
lanternfish larvae. Visual neuroscience, 24(3), 423-436.

Byun JH, et al. 2020. Gene expression patterns of novel visual and non-visual opsin families
in immature and mature Japanese eel males. PeerJ, 8, e8326.

Carleton KL, Kocher TD. 2001. Cone opsin genes of African cichlid fishes: tuning spectral
sensitivity by differential gene expression. Molecular biology and evolution 18.8.

Carleton KL, Dalton BE, Escobar-Camacho D, Nandamuri SP. 2016. Proximate and
ultimate causes of variable visual sensitivities: insights from cichlid fish radiations. Genesis,
54(6), 299-325.

Carleton KL, Escobar-Camacho D, Stieb SM, Cortesi F, Marshall NJ. 2020. Seeing the
rainbow: mechanisms underlying spectral sensitivity in teleost fishes. Journal of Experimental
Biology, 223(8).

Collin SP, Hoskins RV, Partridge JC. 1998. Seven Retinal Specializations in the Tubular
Eye of the Scopelarchus michaelsarsi: A Case Study in Visual Optimization. Brain, Behavior
and Evolution, 1998(09), 291–314.

Collin SP, Marshall NJ. 2003. Sensory Processing in Aquatic Environments. Springer-Verlag
New York.

Cortesi F, et al. 2015. Ancestral duplications and highly dynamic opsin gene evolution in
percomorph fishes. Proceedings of the National Academy of Sciences 112.5.

Cortesi F, et al. 2016. From crypsis to mimicry: changes in colour and the configuration of
the visual system during ontogenetic habitat transitions in a coral reef fish. Journal of
Experimental Biology, 219(16), 2545-2558.

Cortesi F, et al. 2020. Visual system diversity in coral reef fishes. Seminars in Cell &
Developmental Biology. Academic Press.

Cortesi F, **et al.** 2021. Multiple ancestral duplications of the red-sensitive opsin gene (LWS)
in teleost fishes and convergent spectral shifts to green vision in gobies. bioRxiv:
https://doi.org/10.1101/2021.05.08.443214.

Dalton BE, Lu J, Leips J, Cronin TW, Carleton KL. 2015. Variable light environments
induce plastic spectral tuning by regional opsin coexpression in the African cichlid fish,
Metriaclima zebra. Molecular ecology 24,16 (2015).

de Busserolles F, et al. 2017. Pushing the limits of photoreception in twilight conditions: The
rod-like cone retina of the deep-sea pearlsides. Science advances, 3(11), eaao4709.

de Busserolles F, Fogg L, Cortesi F, Marshall J. 2020. The exceptional diversity of visual
adaptations in deep-sea teleost fishes. Seminars in Cell & Developmental Biology. Academic
Press.

Dalton BE, Loew ER, Cronin TW, Carleton KL. 2014. Spectral tuning by opsin
coexpression in retinal regions that view different parts of the visual field. Proceedings of the
Royal Society B: Biological Sciences 281,1797.

Denton EJ. 1990. Light and vision at depths greater than 200 metres. Light and life in the sea
127-148.

Denton EJ, Herring PJ, Widder EA, Latz MF, Case JF. 1985 The roles of filters in the
photophores of oceanic animals and their relation to vision in the oceanic environment.
Proceedings of the Royal Society of London. Series B. Biological Sciences 225, 1238.

Douglas RH, Hunt DM, Bowmaker JK. 2003. Spectral sensitivity tuning in the deep-sea. In
Sensory Processing in Aquatic Environments (pp. 323-342). Springer, New York, NY.

Douglas RH, Partridge JC, Marshall NJ. 1998. The visual systems of deep-sea fish. I.
Optics, tapeta, visual and lenticular pigmentation. Prog. Ret. Eye Res, 17(4), 597-636.

Douglas RH, Genner MJ, Hudson AG, Partridge JC, Wagner HJ. 2016. Localisation and
origin of the bacteriochlorophyll-derived photosensitizer in the retina of the deep-sea dragon
fish *Malacosteus niger*. Scientific reports, 6, 39395.

Downes GB, Gautam N. 1999. The G protein subunit gene families. Genomics, 62(3), pp.544-
552.

Enright JM, et al. 2015. *Cyp27c1* red-shifts the spectral sensitivity of photoreceptors by
converting vitamin A1 into A2. Current Biology, 25(23), 3048-3057.

Hope AJ, Partridge JC, Dulai KS, Hunt DM. 1997. Mechanisms of wavelength tuning in
the rod opsins of deep-sea fishes. Proceedings of the Royal Society of London. Series B:
Biological Sciences, 264(1379), 155-163.

Huang Z, Titus T, Postlethwait JH, Meng F. 2019. Eye Degeneration and Loss of otx5b
Expression in the Cavefish Sinocyclocheilus tileihornes. Journal of molecular evolution 87.7.

Hunt DM, Dulai KS, Partridge JC, Cottrill P, Bowmaker JK. 2001. The molecular basis
for spectral tuning of rod visual pigments in deep-sea fish. Journal of Experimental Biology,
204(19), 3333-3344.

Hunt DM, Fitzgibbon J, Slobodyanyuk SJ, Bowmaker JK., Dulai KS. 1997. Molecular
evolution of the cottoid fish endemic to Lake Baikal deduced from nuclear DNA evidence.
Molecular Phylogenetics and Evolution, 8(3), 415-422.

Hunt DM, Hankins MW, Collin SP, Marshall NJ. 2014. Evolution of visual and non-visual
pigments (Vol. 4). Boston, MA: Springer.

Katoh K, Asimenos G, Toh H. 2009. Multiple alignment of DNA sequences with MAFFT.
In Bioinformatics for DNA sequence analysis (pp. 39-64). Humana Press.

Kawamura S, Tachibanaki S. 2012. Explaining the functional differences of rods versus
cones. Wiley Interdisciplinary Reviews: Membrane Transport and Signaling 1.5.

Kearse M, et al. 2012. Geneious Basic: an integrated and extendable desktop software
platform for the organization and analysis of sequence data. Bioinformatics, 28(12), 1647-
1649.

La Vail MM, Rapaport DH, Rakic P. 1991. Cytogenesis in the monkey retina. J. Comp.
Neurol. 309, 86-114. doi:10.1002/cne.903090107.

Lamb TD. 2013. Evolution of phototransduction, vertebrate photoreceptors and retina.
Progress in retinal and eye research, 36, 52-119.

Lamb TD. 2019. Evolution of the genes mediating phototransduction in rod and cone
photoreceptors. Progress in Retinal and Eye Research, p.100823.

Larhammar D, Nordström K, Larsson TA 2009.**.** Evolution of vertebrate rod and cone
phototransduction genes. Philosophical Transactions of the Royal Society B: Biological
Sciences, 364(1531), pp.2867-2880.

Liu D, et al. 2019. The cone opsin repertoire of osteoglossomorph fishes: gene loss in
mormyrid electric fish and a long wavelength-sensitive cone opsin that survived 3R. Molecular
biology and evolution 36.3.

Lythgoe JN. 1966. Visual pigments and under- water vision. In: Light as an Ecological Factor
(Rackham, O., ed.). Oxford: Blackwell.

Luehrmann M, et al. 2019. Cardinalfishes (Apogonidae) show visual system adaptations
typical of nocturnally and diurnally active fish. Molecular ecology 28.12.

Luo DG, et al. 2020. Apo-Opsin and Its Dark Constitutive Activity across Retinal Cone
Subtypes." Current Biology 30.24.

Ma JX, et al. 2001. A visual pigment expressed in both rod and cone photoreceptors. Neuron,
32(3), 451-461.

Mariani AP. 1986. Photoreceptors of the larval tiger salamander retina. Proceedings of the
Royal society of London. Series B. Biological sciences 227.1249.

Mears AJ, et al. 2001. *Nrl* is required for rod photoreceptor development. Nature genetics,
29(4), 447-452.

Moser HG, Smith PE. 1993. Larval fish assemblages and oceanic boundaries. Bull. Mar. Sci,
53(2), 283-289.

Munk O. 1990. Changes in the visual cell layer of the duplex retina during growth of the eye
of a deep-sea teleost, Gempylus serpens Cuvier, 1829. Acta Zoologica, 71(2), 89-95.

Musilova Z, et al. 2019a. Vision using multiple distinct rod opsins in deep-sea fishes. Science,
364(6440), 588-592.

Musilova Z, et al. 2019b. Evolution of the visual sensory system in cichlid fishes from crater
lake Barombi Mbo in Cameroon. Molecular Ecology, (August), 5010–5031.

Musilova Z, Salzburger W, Cortesi F. 2021. The Visual Opsin Gene Repertoires of Teleost
Fishes: Evolution, Ecology and Function. Annual Review of Cell and Developmental Biology,
37, https://doi.org/10.1146/annurev-cellbio-120219-024915.

Musilova Z, Cortesi F. 2021. Multiple ancestral and a plethora of recent gene duplications
during the evolution of the green sensitive opsin genes (RH2) in teleost fishes. bioRxiv,
https://doi.org/10.1101/2021.05.11.443711.

Partridge JC, Shand J, Archer SN, Lythgoe JN, van Groningen-Luyben WA. 1989.
Interspecific variation in the visual pigments of deep-sea fishes. Journal of Comparative
Physiology A, 164(4), 513-529.

Partridge JC, Archer SN, van Oostrum J. 1992. Single and multiple visual pigments in deep-
sea fishes. J. Mar. Biol. Assoc. U. K. 72, 113-130

Pointer MA, Carvalho LS, Cowing JA, Bowmaker JK, Hunt DM. 2007. The visual
pigments of a deep-sea teleost, the pearl eye Scopelarchus analis. Journal of Experimental
Biology, 210(16), 2829-2835.

Raymond PA. 1995. Development and morphological organization of photoreceptors. In
Neurobiology and Clinical Aspects of the Outer Retina (pp. 1-23). Springer, Dordrecht.

Reif WE. 1985. Functions of scales and photophores in mesopelagic luminescent sharks. Acta
Zoologica, 66(2), 111-118.

Ronquist F, et al. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model
choice across a large model space. Systematic biology, 61(3), 539-542.

Sassa C, Hirota Y. 2013. Seasonal occurrence of mesopelagic fish larvae on the onshore side
of the Kuroshio off southern Japan. Deep Sea Research Part I: Oceanographic Research Papers,
81, 49-61.

Savelli I, Flamarique IN, Iwanicki T, Taylor JS. 2018. Parallel opsin switches in multiple
cone types of the starry flounder retina: tuning visual pigment composition for a demersal life
style. Scientific reports 8:1.

Schott RK, Bhattacharyya N, Chang BS. 2019. Evolutionary signatures of photoreceptor
transmutation in geckos reveal potential adaptation and convergence with snakes. Evolution,
73(9), 1958-1971.

Schott RK, et al. 2016. Evolutionary transformation of rod photoreceptors in the all-cone
retina of a diurnal garter snake. Proceedings of the National Academy of Sciences, 113(2), 356-
361.

Sernagor E, Eglen S, Harris B, Wong R. 2006. Retinal development. Cambridge University
Press.

Shen YC, Raymond PA. 2004. Zebrafish cone-rod (*crx*) homeobox gene promotes
retinogenesis. Developmental biology, 269(1), 237-251.

Simoes BF, et al. 2016. Multiple rod–cone and cone–rod photoreceptor transmutations in
snakes: evidence from visual opsin gene expression. Proceedings of the Royal Society B:
Biological Sciences, 283(1823), 20152624.

Stieb SM, et al. 2019. A detailed investigation of the visual system and visual ecology of the
Barrier Reef anemonefish, *Amphiprion akindynos*. Scientific reports 9.1.

Sugawara T, et al. 2005. Parallelism of amino acid changes at the RH1 affecting spectral
sensitivity among deep-water cichlids from Lakes Tanganyika and Malawi. Proceedings of the
National Academy of Sciences, 102(15), 5448-5453.

Tettamanti V, de Busserolles F, Lecchini D, Marshall NJ, Cortesi F. 2019. Visual system
development of the spotted unicornfish, *Naso brevirostris* (Acanthuridae). Journal of
Experimental Biology, 222(24).

Torres-Dowdall J, et al. 2017. Rapid and parallel adaptive evolution of the visual system of
Neotropical Midas cichlid fishes. Molecular biology and evolution 34.10.

Turner JR, White EM, Collins MA, Partridge JC, Douglas RH. 2009. Vision in lanternfish
(Myctophidae): adaptations for viewing bioluminescence in the deep-sea. Deep Sea Research
Part I: Oceanographic Research Papers 56, 6.

Underwood G. 1968. Some suggestions concerning vertebrate visual cells. Vision research,
8(4), 483-488.

Valen R, et al. 2016. The two-step development of a duplex retina involves distinct events of
cone and rod neurogenesis and differentiation. Developmental biology, 416(2), 389-401.

Wagner HJ, Partridge JC, Douglas RH. 2019. Observations on the retina and 'optical fold'of
a mesopelagic sabretooth fish, *Evermanella balbo*. Cell and Tissue Research, 378(3), 411-425.

Yokoyama S. 2008. Evolution of dim-light and color vision pigments. Annu. Rev. Genomics
Hum. Genet., 9, 259-282.

Yokoyama S, Jia H. 2020. Origin and adaptation of green-sensitive (RH2) pigments in
vertebrates. FEBS Open Bio, 10(5), 873-882.

Zhang H, et al. 2020. Molecular cloning of fresh water and deep-sea rod opsin genes from
Japanese eel *Anguilla japonica* and expressional analyses during sexual maturation. Febs
Letters, 469(1), 39-43.

*Fig. 1: Cone and rod opsin gene expression in larval and adult deep-sea fishes. A) Transcriptomes were used to characterise the opsin gene expression in the retinas of 20 deep-sea fish species, belonging to eight different orders and mapped onto a simplified teleost phylogeny (topology after Betancur et al. 2017). Boxes highlight the five species for which both larval and adult samples were available. B) Proportional opsin gene expression for each individual (horizontal bar) at different developmental stages. Different colours correspond to cone (colours) or rod (shades of grey) opsin genes, depicted as the proportional expression over the total sum of visual opsins expressed. Different shades of the same colour represent multiple copies of the same gene family. Based on the opsin gene expression, the larvae (left column) show a pure-cone or cone-dominated retina, while the adults (right column) have a pure-rod or rod-dominated visual system. Juvenile specimens in two species had an adult expression profile. Note that some species expressed multiple RH1 copies (*Scopelarchus, Howella brodiei *and*

Ceratoscopelus warmingii *adults) or multiple RH2 copies (*Gyrinomimus *sp. larva,* Hygophum reinhardtii *larva).*

Notably, adults and larvae of Scopelarchus *sp. and* Coccorella atlantica *expressed different copies of RH2 (more*

details in Fig. 2). Details about the samples and expression levels are listed in Table 1. C) The genomic repertoire

of the visual opsins is shown for five species Idiacanthus fasciola, Chauliodus sloani, Coccorella atlantica,

Scopelarchus michaelsarsi *(all this study), and* Anoplogaster cornuta *(Musilova et al. 2019a). The rod RH1 opsin*

and the cone RH2 opsin genes are present in all studied species in one or multiple (up to seven) copies. The SWS2

opsin gene was found only in the fangtooth, and the SWS1 and LWS are missing from all five studied genomes.

Fig. 2: Phototransduction cascade gene expression in the retinas of five deep-sea fish species. *A) Heat map of the expression of individual phototransduction cascade genes for each sample, based on normalised numbers of reads (FPKM). B) Pie charts comparing mean values of relative expression of the opsin genes (rod RH1 and cone*

RH2), photoreceptor-specific cascade transducin genes (rod-type GNAT1 and cone-type GNAT2), and all cascade

genes (photoreceptor-specific transducins, arrestins and phosphodiesterases) summarized. The black square

highlights the two aulopiform species with the discordance between the opsin type (rod-specific) and

*phototransduction cascade genes (cone-specific) in adults. C) Focus on the common fangtooth (*Anoplogaster

cornuta*) transitional phase shown as a sequence for seven larval and one adult sample. Size given as standard*

length (SL). Note that all fangtooth larvae expressed both RH1 and RH2, with an increasing proportion of RH1

to RH2 as the larvae increased in size (with the exception of the largest larva where RH1:RH2 was 28:72). Smaller

larvae also expressed the SWS2 gene. These individuals all had traits of larval phenotypes (dorsal and ventral

horns and small teeth; Fig. 1) and were collected relatively shallow between 0-300 m using the plankton trawls.

Fig. 3: Gene trees of the A) RH1 and B) RH2 opsin genes found in the retinal transcriptomes of deep-sea fishes.
Species with multiple copies are highlighted in colour. Additional gene sequences from public databases are listed
with their GenBank accession numbers. Note the topology within Aulopiformes; the adult RH2s of Cocorrella
atlantica and Scopelarchus cluster together as do the major larval RH2s. Yellow circles mark lineage-specific
gene duplication events, while red circles pinpoint the ancestral duplication of RH1 impacting the Scopelarchus
genus, and the duplication of RH2 in the aulopiform ancestor (or at least the common ancestor of Coccorella and
Scopelarchus). C) Key tuning spectral site mutations in species with multiple rod opsins and indicative wavelength
shifts based on the previous in vitro experiments. The known shifts are listed in nanometers. Blue and green letters
in the tables stand for the shorter- and longer-shifting amino acid variants, respectively. Multiple different rod
opsins have been found in three species, Ceratoscopelus warmingii (Myctophiformes), Howella brodiei
(Pempheriformes) and Scopelarchus michaelsarsi (Aulopiformes). Note that the RH1 copies in Scopelarchus seem
to show a mixed pattern - the longer-wavelength sensitive copy (RH1a; confirmed by in vitro measurements by
Pointer et al. 2007) carries also several shorter-shifting amino-acid sites as compared to RH1b). For functional

interpretation we considered microspectrophotometry measurements from # = Pointer et al. 2007, † = Collin and

Marshall 2003 and ‡ = Partridge et al. 1989.

Table 1: *Samples used in the study and results of the opsin gene expression in the eyes or retina. * https://obis.org/*

Species	Order	Median species depth (m)*	Stage	Code	Size (mm)	Date of collection	Raw reads	Reads after bacteria filtering	Gene expression				Accession no.	Source
									RH1	RH2	SWS1	SWS2		
Anoplogaster cornuta	Trachichthyiformes	1000-2000	Larva	39016	4	April 2015	36,373,372	36,125,630	<0.01	0.99		<0.01	SAMN10473242	GenBank
			Larva	39017	4	April 2015	34,712,538	34,604,364	<0.01	0.99		<0.01	SAMN10473243	GenBank
			Larva	4_23	9	April 2015	41,592,840	41,519,070	0.02	0.98		<0.01	SAMN10473246	GenBank
			Larva	4_22	11	April 2015	75,109,270	74,990,214	0.11	0.89		<0.01	SAMN10473245	GenBank
			Larva	4_21	11-12	April 2015	28,945,804	28,869,552	0.22	0.78		<0.01	SAMN10473244	GenBank
			Larva	261s03	12	30.3.2017	48,458,044	48,457,472	0.47	0.53		<0.01	SAMN20741930	this study
			Larva	126A1	20	1.3.2020	79,371,594	79,347,702	0.28	0.72			SAMN20746021	this study
			Adult	56H6	102	6.-7.4.2017	11,658,040	11,635,610	1				SAMN20748547	this study
Ceratoscopelus warmingii	Myctophiformes	400-500	Adult	300s03	63	7.-8.4.2017	49,078,974	44,114,639	0.66 [RH1-1] 0.01 [RH1-2] 0.33 [RH1-3]				SAMN20752508	this study
			Adult	S1	n/a	April 2015	8,796,786	8,789,961	0.94 [RH1-1] 0.02 [RH1-2] 0.04 [RH1-3]				SAMN10473255	GenBank
			Adult	S2	n/a	April 2015	8,655,248	8,649,342	0.93 [RH1-1] 0.02 [RH1-2] 0.05 [RH1-3]				SAMN10473256	GenBank
			Adult	S3	n/a	April 2015	7,904,714	7,899,617	0.94 [RH1-1] 0.01 [RH1-2] 0.05 [RH1-3]				SAMN10473257	GenBank
Chauliodus sloani	Stomiiformes	800-900	Larva	67I2_2	11	3.4.2017	34,445,252	34,371,394	0.01	0.99			SAMN20769443	this study
			Larva	67I1	17	3.4.2017	20,701,686	20,683,764	<0.01	0.99			SAMN20769824	this study
			Juvenile	109B6	20	5.4.2017	24,854,866	24,835,646	1				SAMN20771653	this study
Chauliodus danae	Stomiiformes	800-900	Juvenile	109C7	24	30.3.2017	32,865,334	32,834,300	1				SAMN20771814	this study
			Juvenile	109A2	20	5.4.2017	25,389,318	25,344,182	1				SAMN20777471	this study
			Juvenile	109D7	96	30.3.2017	18,540,248	18,533,484	1				SAMN20777490	this study
Coccorella atlantica	Aulopiformes	300-400	Larva	109G8	21	30.3.2017	26,376,620	26,359,060	0.01	0.99 [RH2-1]			SAMN20793157	this study
			Adult	56C7	77	25.3.2017	36,794,070	36,790,854	0.69	0.31 [RH2-2]			SAMN20793158	this study
			Adult	297_7	68	6.-7.4.2017	43,948,534	40,511,279	0.96	0.04 [RH2-3]			SAMN20793260	this study
Grammatostomias flagellibarba	Stomiiformes	500-600	Adult	56H8	138	6.-7.4.2017	9,775,520	9,742,864	1				SAMN20794729	this study
Gyrinomimus myersi	Beryciformes	1000-2000	Larva	S25	n/a	April 2015	19,270,096	19,122,986	<0.01	0.45 [RH2-1] 0.13 [RH2-2] 0.08 [RH2-3] 0.11 [RH2-4] 0.15 [RH2-5]	0.07	<0.01	SAMN10473271	GenBank
Howella brodiei	Pempheriformes	500-600	Adult	56D8	76	25.3.2017	56,905,264	43,936,178	0.17 [RH1-1] 0.83 [RH1-2]				SAMN20797093	this study
			Adult	56D9	68	25.3.2017	61,909,674	61,590,270	0.07 [RH1-1] 0.93 [RH1-2]				SAMN20799523	this study
Hygophum reinhardtii	Myctophiformes	100-200	Larva	67I2_1	5	3.4.2017	17,222,626	17,122,880		0.26 [RH2-1] 0.68 [RH2-2] 0.06 [RH2-3]			SAMN20799641	this study
Idiacanthus fasciola	Stomiiformes	600-700	Larva	71C2	22	22.3.2017	23,382,468	23,379,360		1			SAMN20800764	this study
			Larva	67D2	22	31.3.2017	6,647,384	6,623,496		1			SAMN20800770	this study
			Larva	71C1	18	22.3.2017	23,632,774	23,497,715		1			SAMN20800774	this study
			Larva	67I7	25	5.4.2017	25,713,960	25,701,768		1			SAMN20800775	this study
			Larva	71B9	14	22.3.2017	43,695,620	43,633,424		1			SAMN20801851	this study
			Larva	67B8	17	29.3.2017	28,333,642	28,313,024		1			SAMN20804813	this study
			Larva	109A1	41	5.4.2017	27,834,274	27,818,750		1			SAMN20804946	this study
			Adult	228s01	210	22.-23.3.2017	21,704,528	21,622,274	1				SAMN20805116	this study
			Adult	67F8	178	1.4.2017	15,744,604	15,739,004	1				SAMN20805515	this study
			Adult	67B6	75	29.3.2017	15,439,954	11,368,044	1				SAMN20808344	this study
Melamphaes sp.	Beryciformes	600-700	Juvenile	4_26	n/a	April 2015	38,583,370	38,534,507	1				SAMN10473273	GenBank
			Juvenile	4_28	n/a	April 2015	35,489,970	35,431,421	1				SAMN10473274	GenBank
Melanolagus bericoides	Argentiniformes	800-900	Larva	71H3	7	26.3.2017	26,219,646	26,210,402		1			SAMN20834672	this study
Pteraclis aesticola	Scombriformes	0-300	Larva	109H4	8	30.3.2017	19,918,922	19,870,484	0.22	0.78			SAMN20844147	this study
Pollichthys mauli	Stomiiformes	300-400	Adult	109I2	34	30.3.2017	55,767,498	54,875,747	1				SAMN20845044	this study
			Adult	109I3	36	30.3.2017	68,655,686	67,082,709	1				SAMN20848441	this study
Scombrolabrax heterolepis	Scombriformes	800-900	Adult	,56E3	91	25.3.2017	65,770,014	65,527,135	0.98	0.02			SAMN20857047	this study
			Adult	,56E4	77	25.3.2017	60,643,652	60,472,479	0.93	0.07			SAMN20857932	this study
Scopelarchus guentheri	Aulopiformes	200-300	Larva	109H1	21	30.3.2017	37,542,802	37,416,486		0.96 [RH2_larval] 0.04 [RH2-2_larval]			SAMN20859427	this study
Scopelarchus michaelsarsi	Aulopiformes	300-400	Larva	109B9	21	5.4.2017	9,971,250	9,919,148		1 [RH2_larval]			SAMN20860823	this study
			Larva	71B7	20	22.3.2017	43,695,620	43,682,534	<0.01 [RH1a] <0.01 [RH1b]	0.96 [RH2_larval] 0.03 [RH2-2_larval]			SAMN20865446	this study
			Adult	272_10	66	2.4.2017	45,565,202	43,729,338	0.80 [RH1a] 0.10 [RH1b]	0.10 [RH2_adult]			SAMN20867027	this study
Scopelogadus mizolepis	Beryciformes	800-900	Adult	,57E2	43	5.4.2017	5,160,876	5,160,672	1				SAMN20867114	this study
			Adult	300s01	62	7.-8.4.2017	15,448,240	15,420,650	1				SAMN20867573	this study
Scopelosaurus hoedti	Aulopiformes	500-600	Adult	297_9	69	6.-7.4.2017	43,948,534	42,248,053	0.98	0.02			SAMN20872996	this study
Vinciguerria poweriae	Stomiiformes	100-300	Larva	109H2	18	30.3.2017	25,863,628	25,755,100		1			SAMN20878005	this study
			Larva	109C8	16	30.3.2017	22,914,004	22,858,398		1			SAMN20882114	this study

691   ***Table 2:** Key-tuning amino acid sites in the cone opsin RH2 gene*

Species	Order	83	122	207	255	292	λmax (nm)	Reference
Bovine RH1		D	E	M	I	A	500	Yokoyama 2008
Ancestral teleost		D	Q	M	I	A	488	Yokoyama & Yia 2020
Melanolagus bericoides	Argentiniformes	G	Q	.	V	.		
Coccorella atlantica adult	Aulopiformes	G	Q	.	.	.		
Coccorella atlantica larval	Aulopiformes	G	Q	.	V	.		
*Scopelarchus michaelsarsi_* RH2_adult	Aulopiformes	?	Q	I	C	T		
Scopelarchus guentheri _RH2_larval	Aulopiformes	G	Q	.	V	.		
*Scopelarchus michaelsarsi_* RH2_ larval	Aulopiformes	G	Q	.	V	.	505	Pointer et al. 2007
*Scopelarchus guentheri_* RH2-2_ larval	Aulopiformes	G	Q	.	C	.		
*Scopelarchus michaelsarsi_* RH2-2_ larval	Aulopiformes	G	Q	.	C	.		
Scopelosaurus hoedti	Aulopiformes	G	Q	.	.	.		
Gyrinomimus myersi RH2-1	Beryciformes	G	Q	.	F	.		
Gyrinomimus myersi RH2-2	Beryciformes	G	Q	L	F	.		
Gyrinomimus myersi RH2-3	Beryciformes	G	Q	L	F	.		
Gyrinomimus myersi RH2-4	Beryciformes	G	Q	L	F	.		
Gyrinomimus myersi RH2-5	Beryciformes	G	Q	L	F	.		
Lepisosteus platyrhincus (shallow outgroup)	Lepisosteiformes	G	.	.	.	.		
Hygophum reinhardtii RH2-1	Myctophiformes	G	Q	.	V	.		
Hygophum reinhardtii RH2-2	Myctophiformes	G	Q	.	V	.		
Hygophum reinhardtii RH2-3	Myctophiformes	G	Q	.	V	.		
Lepidopus fitchi RH2-1	Scombriformes	G	.	.	V	.	496	Yokoyama & Yia 2020
Lepidopus fitchi RH2-2	Scombriformes	G	Q	.	V	.		
Lepidopus fitchi RH2-3	Scombriformes	G	Q	.	V	.	506	Yokoyama & Yia 2020
Lepidopus fitchi RH2-4	Scombriformes	G	Q	.	V	.		
Pteraclis aesticola	Scombriformes	G	Q	L	.	.		
Scombrolabrax heterolepis	Scombriformes	G	Q	.	.	.		
Aristostomias scintillans	Stomiiformes	G	Q	L	V	.	468	Yokoyama & Yia 2020
Chauliodus sp.	Stomiiformes	G	Q	.	F	.		
Grammatostomias flagellibarba	Stomiiformes	G	Q	.	.	.		
Idiacanthus fasciola	Stomiiformes	G	Q	.	V	.		
Vinciguerria poweriae	Stomiiformes	G	Q	.	F	.		
Anoplogaster cornuta	Trachichthyiformes	G	Q	.	.	.		

Table 3: *Key-tuning amino acid sites in the rhodopsin RH1 gene. * = value for* Scopelarchus analis*; † = two pigments reported without assignment to the gene; see also Figure 3; ‡ = value for closely related species,* Evermannella balbo*; § = value for* Scopelogadus beani*.*

Species	Order	83	90	96	102	111	113	118	122	124	132	164	183	188	194	195	207	208	211	214	253	261	265	269	289	292	295	299	300	317	λmax (nm)	Reference
bovine RH1		D	G	Y	Y	N	E	T	E	A	A	A	M	G	P	H	M	F	H	I	M	F	W	A	T	A	A	A	V	M	500	Yokoyama 2008
ancestral teleost RH1		D	G	Y	Y	.	E	T	E	A	A	A	M	.	L	N	M	F	H	I	M	F	W	A	T	A	A	A	L	M		Musilova et al. 2019
Bathysaurus ferox	Aulopiformes	N	.	.	.	.	.	.	S	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	S	.	.	T	L	.	481	Collin & Marshall 2003
Bathysaurus mollis	Aulopiformes	N	.	.	.	.	.	.	S	.	.	.	.	.	R	A	.	.	.	.	.	.	.	T	S	.	.	T	L	.	479	Collin & Marshall 2003
Coccorella atlantica	Aulopiformes	N	.	.	.	S	.	.	M	.	.	.	.	.	R	A	.	.	.	C	.	.	.	.	.	S	.	I	.	.	480‡	Douglas et al. 1998
Scopelarchus michaelsarsi RH1a	Aulopiformes	N	.	.	.	.	.	.	S	.	.	.	.	.	R	A	.	.	.	C	.	.	.	.	S	S	S	I	.	.	486*	Pointer et al. 2007
Scopelarchus michaelsarsi RH1b	Aulopiformes	N	.	.	.	.	.	.	S	.	.	.	.	.	L	K	.	.	.	V	.	.	.	.	.	.	.	.	L	.	479*	Pointer et al. 2007
Scopelosaurus hoedti	Aulopiformes	.	.	.	.	.	.	.	.	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	S	S	S	I	.	.		
Gyrinomimus myersi	Beryciformes	N	.	.	.	.	.	.	.	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	.	S	.	I	.	.		
Melamphaes sp.	Beryciformes	.	.	.	.	.	.	.	Q	.	S	.	.	.	R	V	.	.	.	.	G	.	.	.	S	.	.	S	I	.		
Scopelogadus mizolepis	Beryciformes	.	.	.	.	.	.	.	Q	G	S	.	.	.	R	V	.	.	.	.	G	.	.	T	.	.	.	S	.	.	488§	Douglas et al. 1998
Lepisosteus oculatus (outgroup)	Lepisosteiformes	.	.	.	.	.	.	.	Q	S	.	.	.	.	L	K	.	.	.	L	.	Y	.	G	.	.	.	L	.	.		
Ceratoscopelus warmingii RH1-1	Myctophiformes	.	.	.	.	T	.	.	Q	.	S	.	.	.	R	A	.	.	.	.	G	.	.	.	A	.	.	S	I	.	488 †	Collin & Marshall 2003
Ceratoscopelus warmingii RH1-2	Myctophiformes	.	.	W	.	I	.	.	I	T	.	.	.	.	R	A	.	.	.	.	L	Y	.	.	.	.	.	I	.	.	undetected	Collin & Marshall 2003
Ceratoscopelus warmingii RH1-3	Myctophiformes	N	.	.	.	T	.	.	Q	.	S	S	.	.	N	V	.	Y	.	G	.	.	.	.	S	.	.	I	.	.	468†	Collin & Marshall 2003
Hygophum reinhardtii	Myctophiformes	.	.	.	A	.	.	.	Q	.	.	.	.	.	R	A	.	.	.	.	G	.	.	.	.	.	.	I	.	.		
Howella brodiei RH1-1	Pempheriformes	N	.	.	.	.	.	.	G	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	G	S	S	S	I	.		
Howella brodiei RH1-2	Pempheriformes	N	.	.	.	.	.	.	Q	S	.	.	.	.	R	A	.	.	.	A	.	.	.	.	S	S	S	I	.	.		
Pteraclis aesticola	Scombriformes	.	.	.	.	.	.	.	Q	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	.	.	.	S	I	.		
Scombrolabrax heterolepis	Scombriformes	.	.	.	.	.	.	.	Q	.	.	.	.	.	R	A	.	.	.	.	.	.	S	.	S	.	.	S	I	.		
Chauliodus spp.	Stomiiformes	N	.	.	.	.	.	.	.	G	.	.	.	.	R	A	.	.	.	V	.	.	.	.	A	S	.	.	L	V	484	Collin & Marshall 2003
Grammatostomias flagellibarba	Stomiiformes	N	.	.	.	.	.	.	.	.	.	.	.	.	R	A	.	.	.	V	.	.	.	.	S	.	.	.	L	.	480 - 487	Kenaley et al. 2014
Idiacanthus fasciola	Stomiiformes	N	.	.	.	.	.	.	.	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	S	.	.	.	L	.	485	Collin & Marshall 2003
Pollichthys mauli	Stomiiformes	N	.	.	.	.	.	.	.	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	S	.	.	.	L	.		
Anoplogaster cornuta	Trachichthyiformes	N	.	.	.	.	.	.	.	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	T	S	.	S	I	.	485	Collin & Marshall 2003