The expression of deep-sea fish visual genes supports a conserved cone-to-

rod vertebrate retinal development

Nik Lupše[1*], Fabio Cortesi[2], Marko Freese[3], Lasse Marohn[3], Jan-Dag Pohlman[3], Klaus

Wysujack[3], Reinhold Hanel[3], Zuzana Musilova[1*]

1) Department of Zoology, Faculty of Science, Charles University, Vinicna 7, 12844 Prague,

Czech Republic

2) Queensland Brain Institute, University of Queensland, Brisbane 4072 QLD, Australia

3) Thünen Institute of Fisheries Ecology, Herwigstraße 31, 27572, Bremerhaven, Germany

* corresponding authors; lupsen@natur.cuni.cz, zuzana.musilova@natur.cuni.cz

Keywords: opsin, evolution, mesopelagic, adaptation, convergence, phototransduction, vision,

gene expression, rhodopsin

Abstract

The ontogeny of the vertebrate retina is thought to follow a conserved pattern whereby cone

photoreceptors develop first and rod photoreceptors are added only at later developmental

stages. Whether this is also the case for species that use only rods as adults, such as many deep-

sea fishes, remained unclear. Since many deep-sea fishes start their lives in the shallow, well-

lit epipelagic zone cones might still be favoured during early developmental stages. Using a

comparative transcriptomic approach in 20 deep-sea species from eight teleost orders, we show

that while adults mostly rely on rod opsin (*RH1*) for vision in dim light, larvae almost

exclusively express mid-wavelength-sensitive cone opsins (*RH2*) in their retinas. The

phototransduction cascade genes follow a similar ontogenetic pattern of cone- followed by rod-

specific gene expression in most orders, except for the pearleye and sabretooth (Aulopiformes),

in which the cone cascade remains dominant throughout development. Hence, our findings

provide strong molecular support for a conserved cone-to-rod retinal development in

vertebrates.

INTRODUCTION

Vision is a primary sense used by most vertebrates for navigation, predator avoidance, communication and to find food and shelter. At its initiation, vertebrate vision is enabled by cone (photopic, colour vision) and rod (scotopic, mostly colour blind) photoreceptors in the retina containing a light absorbing pigment that consists of an opsin protein covalently bound to a vitamin-A-derived chromophore[1]. The absorbance of photons by the chromophore leads to a conformational change of the opsin protein, which initiates a photoreceptor-specific G-protein-coupled phototransduction cascade, propagating the signal to the brain[2,3,4]. It is thought that the development of the visual system follows a conserved molecular pattern whereby cone specific genes are activated first before the rod molecular pathway is initiated later during ontogeny[5,6,7]. However, whether this is the case for all vertebrates and especially for those that have retinas that contain only rods as adults, remains unclear.

Changes in the light environment, ecology, and phylogenetic inertia are thought to be primary drivers for visual system diversity in vertebrates[8]. For example, most mesopelagic deep-sea fishes (200 – 1,000 m depth), either living strictly at depth or migrating to the shallows at night, have evolved visual systems that are sensitive to the dominant blue light (~ 470 – 490 nm) of their environment[9]. Moreover, as the daylight and the bioluminescent light emitted by deep-sea critters are quickly dimmed with depth, deep-sea fish visual systems have evolved peculiar morphologies to maximise photon capture including barrel-eyes, reflective tapeta and the use of rod-dominated and in many cases rod-only retinas that might be stacked into multiple banks (reviewed in de Busserolles et al.[10]). However, most mesopelagic fishes start their lives in the shallow well-lit epipelagic zone (0 – 200 m depth)[11,12]. Consequently, their visual systems must cope with a variety of light intensities and spectra throughout development.

Studies investigating the gene expression in the retina of deep-sea fishes are scarce and usually focus on a selected few species[13,14,15,16,17]. In adults, species with pure rod retinas tend to only express rod opsin(s)[14,16], albeit two species of pearlsides (*Maurolicus* spp.) have been found to express cone-specific genes (i.e., cone transduction pathway and opsin genes) inside rod-looking cells[15]. It remains unknown whether deep-sea fishes that have a low proportion of cone photoreceptors as adults[18,19,20,21,22] also express cone-specific genes at any stages of their lives or whether molecularly these fishes rely on the rod machinery alone. To investigate whether the retinal development in deep-sea fishes follows a similar cone-to-rod molecular pathway as found in other vertebrates or whether some species start their lives with rod pathway activated, we set out to sequence the retinal transcriptomes of 20 deep-sea fish species,

including the larval stages in ten species, belonging to eight different teleost orders

(Argentiniformes, Aulopiformes, Beryciformes, Myctophiformes, Pempheriformes,

Scombriformes, Stomiiformes and Trachichthyiformes).

RESULTS AND DISCUSSION

Visual opsin gene expression. In teleost fishes, gene duplications and deletions followed by

functional diversification have resulted in extant species having between 1-40 visual opsin

genes within their genomes. These genes are defined by their phylogeny and their spectrum of

maximal sensitivity (λ_{max}) and fall within five major classes, four cone opsins ('ultraviolet or

UV sensitive' *SWS1*: 347–383 nm, 'blue' *SWS2*: 397–482 nm, 'green' *RH2*: 452–537 nm and

'red' *LWS*: 501–573 nm) and one rod opsin ('blue-green' rhodopsin, *RH1* or Rho: 447–525

nm)[23]. Transcriptomic sequencing of 20 deep-sea teleost species revealed that deep-sea fishes

mainly express rod opsins and/or green-sensitive cone opsins (*RH2*s) in their retinas (Fig. 1,

Table 1). While larvae mostly expressed *RH2*, adults and juveniles mostly expressed *RH1* and

in a few cases a combination of both. We found none or very low expression of any of the other

cone opsin genes: the red sensitive *LWS* was not expressed at all, the UV sensitive *SWS1* was

only found in the larva of the whalefish, *Gyrinomimus* sp. (Beryciformes), and the blue/violet

sensitive *SWS2* only in the larvae of the whalefish, and the fangtooth, *Anoplogaster cornuta*

(Trachichthyiformes)*,* (Fig. 1, Table 1). Differences in gene expression patterns are likely to

be driven by ontogenetic transitions in light habitat from bright to dim environments and also

by changes in ecological demands, as discussed in more detail below. Accordingly, the

genomic opsin gene repertoire of the fangtooth (*A. cornuta*; Trachichthyiformes) and

ribbonfish (*I. fasciola*; Stomiiformes) is just composed of *SWS2* and *RH2*, or only *RH2*

opsin(s), respectively. Both species have lost the *LWS* and *SWS1* opsin genes also in their

genome (Fig. 1).

Similar to the opsin genes, we also detected ontogenetic differences in the expression

of phototransduction cascade genes (Fig. 2). Here we focused on the comparison of five species

from three teleost orders for which we had both larval and adult specimens available and found

that the cone-specific genes were mostly expressed in the larval stages (e.g., cone transducin,

GNAT2), while adults from three species mostly expressed rod-specific genes (e.g., rod

transducin, *GNAT1*; Fig. 2b). Hence, at the molecular level, the visual systems of deep-sea

fishes start out with a cone-based expression pattern. Moreover, in the fangtooth, where

samples from various sized specimens were available, we found that the cone-specific

expression was gradually replaced with the rod profile as the fish grew (Fig. 2c, Table S1).

This is similar to the visual development in shallower living fishes (e.g., Atlantic cod[24],

zebrafish[7]) and terrestrial vertebrates (e.g., mice[5], rhesus monkey[25]), where cone

photoreceptors are first to develop, followed by temporally and spatially distinct rods[6,26]. The

cone-to-rod developmental sequence is therefore likely to be shared across vertebrates, even in

species that have pure rod retinas as adults.

Ontogenetic shift in expression profiles and the transition phase. The developmental

changes in the visual system we uncovered are best explained by the different habitats larval

and adult deep-sea fishes inhabit. In general, deep-sea fish larvae live in the shallow epipelagic

zone[11] where ambient light levels are sufficiently high to warrant a cone-based visual system.

After metamorphosis, deep-sea fishes start to submerge deeper and take up a life at different

depths in the mesopelagic or even bathypelagic (below 1,000 m depth) zone, where the sun-

and moonlight is gradually replaced by bioluminescence as the main source of light[27]. In this

extremely dim environment, rods work at their best and cone photoreceptors would be obsolete

for the most part at least. Rod-based vision is also favoured in those deep-sea species that

exhibit diel vertical migrations to feed in the plankton rich surface layers at night[10]. In addition,

we discovered that in some species there was a switch in the type of cone-based *RH2* opsin that

was expressed (Fig. 1). For example, in Aulopiformes, the larvae expressed an *RH2* that is

presumably sensitive to longer wavelengths of light compared to the *RH2* that was found in

adults (Table 2). This clearly shows that larval and adult deep-sea fishes rely on different opsin

expression profiles, which is similar to ontogenetic changes in opsin gene expression in diurnal

shallow-water fishes such as freshwater cichlids[28] and coral reef dottybacks[29,30] or between the

freshwater and deep-sea maturation stages in eels[13].

Our data furthermore suggests that the ontogenetic change in visual gene expression

precedes morphological changes such as metamorphosis from larva to juvenile and also habitat

transitions. For example, in the fangtooth, the larvae which were collected from the shallows

(0 – 300 m) showed increasing amounts of *RH1* expression with growth, despite displaying

larval phenotypes throughout (horns and small teeth; Fig. 1). A similar pattern of changing

opsin gene expression ahead of metamorphosis has also been reported from shallow-water

fishes such as European eels[31], dottybacks[30] and surgeonfishes[32]. Interestingly, all our

fangtooth larvae (incl. the smallest individual with a total length of 4 mm) already expressed a

small amount of *RH1* (Fig. 2c). Whether fangtooth start their lives with a pure-cone retina or

low-levels of rod opsin expression are normal even in pre-flexation larvae remains therefore

unclear. In addition to the green-sensitive cone opsin *RH2*, the fangtooth larvae also expressed

low levels of the blue-sensitive *SWS2*, potentially conferring dichromatic colour vision to the
early life stages of this species (Fig. 1).

Photoreceptor cell identities. Interestingly, two aulopiform species; the Atlantic sabretooth,
Coccorella atlantica, and the Bigfin pearleye, *Scopelarchus michaelsarsi*, despite expressing
mostly *RH1* as adults, retained a cone-dominated phototransduction cascade expression profile
akin to the one found in the larval stages (Fig. 2, Table S1). This begs the question whether the
photoreceptors they are using are of cone or rod nature. Initially described in snakes and
geckos[33,34] and recently also in a deep-sea fish[15], it appears that the dichotomy of rods and
cones is not always as clear cut as one might think. For example, adult deep-sea pearlsides,
Maurolicus spp. have a retina that expresses ~ 99% *RH2* and ~ 1% *RH1* with corresponding
cone and rod phototransduction gene expressions. Their photoreceptors, however, are all rod-
shaped and careful histological examination has shown that these consist of a tiny proportion
of true rods and a majority of transmuted rod-like cones[15]. In the case of the pearlside, and also
in geckos and snakes, the opsin and phototransduction genes correspond to each other making
it possible to distinguish photoreceptor types at the molecular level. However, in the
aulopiforms, high expression of rod opsin is seemingly mismatched with high levels of cone
phototransduction gene expression (Fig. 2). In salamanders, the opposite pattern can be found
whereby a cone opsin is combined with the rod phototransduction cascade inside a rod looking
cone photoreceptor[35]. Anatomically, the retina of *S. michaelsarsi* is composed of mostly rods
with low numbers of cone cells[19], while the adult retina of *Evermanella balbo,* an
evermannellid species related to *C. atlantica,* appears to consist of two differently looking rod
populations[36]. It is therefore likely, that similar to what was found in pearlsides[15], these fishes
have a high proportion of transmuted rod-like cone photoreceptors that use *RH1* instead of a
cone opsin as the visual pigment. Alternatively, a proportion of true rods might make use of
the cone phototransduction cascade. Either way, combining more stable rod opsin in a rod-
shaped cell with the cone-specific cascade is likely to increase sensitivity while also
maintaining high transduction and recovery speeds of cells[37,38,39]. Histology, fluorescent in-situ
hybridisation and ideally physiological recordings are needed to ultimately disentangle the
identity of photoreceptor cells in aulopiform.

Evolutionary history of deep-sea fish opsins. While the majority of adult fishes relied on a
single *RH1* copy, three species were found that expressed multiple *RH1* copies: The Warming's
lanternfish, *Ceratoscopelus warmingii* (Myctophiformes), expressed three different *RH1*

genes, and *S. michaelsarsi* and the basslet, *Howella brodiei* (Pempheriformes), expressed two

copies each. Larvae and a few adult deep-sea fishes mostly expressed a single *RH2* copy, except

for the pearleyes, *Scopelarchus* spp., and the Reinhardt's lanternfish, *Hygophum reinhardtii*,

which expressed three larval copies each, and the whalefish which expressed five larval copies

(Fig. 1).

The *RH1* and *RH2* phylogenies revealed that most deep-sea fish visual opsins cluster

together by species or order (Fig. 3). For example, in the whalefish all *RH2s* are clustered

together suggesting that these genes are lineage or species-specific (Fig. 3b). However, there

were a few exceptions, suggesting more ancient duplication events. In *Scopelarchus* the two

RH1 copies are not in a sister relationship and result in different clusters, suggesting that these

copies originated in the common ancestor of aulopiforms or perhaps even earlier (Fig. 3a).

Also, the *RH2s* in äulopiforms (*Scopelarchus*, *Coccorella*) cluster by ontogenetic stage,

making it likely that the developmental switch in gene expression was already present in the

aulopiform ancestor (Fig. 3b).

Molecular complexity of deep-sea fish visual systems. The complexity of deep-sea fish

visual systems at the molecular level varied quite substantially. For example, the three

Stomiiformes species: The Ribbon sawtail fish, *Idiacanthus fasciola*, and two species of

viperfish, *Chauliodus sloani* and *Ch. danae*, appeared to have a very basic visual set up; these

fishes were found to express a single *RH2* cone opsin and a single *RH1* rod opsin as larvae and

adults, respectively (Fig. 1). On the contrary, a number of the deep-sea fish orders examined

here expressed more than two opsin genes. Adult lanternfishes and basslets have rod-only

retinas but expressed multiple *RH1* copies that have functionally diversified (Fig. 3). Other

species expressed both cone and rod opsins as adults (some aulopiform species and

Scombrolabrax), which is similar to the opsin gene expression profiles found in shallow-living

nocturnal reef fishes[40].

The most complex visual system described here was found in *S. michaelsarsi*. In

general, this species is known for its numerous morphological and anatomical adaptations to

vision in the deep-sea, including having barrel eyes with a main and an accessory retina, rods

that are organised in bundles, large ganglion cells and corneal lens pads[19]. The two copies of

RH1 (*RH1-1* and *RH1-2,* corresponding to the *RH1a* and *RH1b* in *Sc. analis* in Pointer et al.[21])

it expressed showed high sequence diversity differing in 66 amino acids, three of which are

key tuning sites likely to change the spectral sensitivity of the pigments via a shift in λ_{max} (Fig.

3, Tables 2 and 3)[16,41,42]. This supports the findings by Pointer et al.[21] which using

microscpetrophotometry (MSP) in another pearleye species, *S. analis*, found two rod

photoreceptors with different absorption maxima at 479 and 486 nm. The situation is less clear

for the green-sensitive *RH2* opsin. While in *S. analis* cones have been found in the accessory

and main retinas, in *S. michaelsarsi* cone photoreceptors appear restricted to the accessory

retina alone[19]. This is intriguing as it suggests substantial differences in visual systems even

between closely related species from the same genus.

The visual ecology of deep-sea fishes. We found molecular support for deep-sea visual

adaptations on multiple levels:

1) Opsin gene diversity in the genome. Retinal transcriptomes in the Stomiiformes pointed

towards a simple visual system that expresses a single opsin gene at different developmental

stages (*RH2* in larvae, *RH1* in adults) (Fig. 1). Besides *RH2*, searching for visual opsins in a

newly sequenced (*I. fasciola*) and several published genomes[16], revealed that Stomiiformes are

likely to have lost all other cone opsin gene families (Fig. 1). For *RH2*, they seem to only have

a single or at most two gene copies, which is substantially less compared to other teleosts[16].

The Stomiiform example, therefore, shows that a decrease in light intensity and the spectrum

of light in the deep-sea may not only restrict gene expression at adult stages, but also lead to

the loss of opsin genes altogether. Similarly, a loss in opsin and other vision-related genes (e.g.

otx5b, *crx*) has previously been reported from shallow living fishes that are either nocturnal[43],

live in murky waters[44], or inhabit caves or similarly dim environments[16,45].

2) Visual gene expression. Previous work has found that the expression of the longest- (*LWS* -

red) and shortest- (*SWS1* - UV) sensitive opsins is reduced or absent in deeper living coral reef

fishes[40] and also in fishes inhabiting deep freshwater lakes[46,47,48] which is correlated with a loss

of short- and long-wavelengths with depth. Accordingly, we found here that deep-sea fishes

lack any *LWS* expression even in the shallow-living larval stages (Fig. 1), which is

accompanied by the loss of *LWS* cone opsin in many deep-sea fish lineages[16]. Similarly, *SWS1*

is not expressed in any of the species studied, except for in the larval whalefish, and is also

absent from many deep-sea fish genomes (Fig. 1)[16]. However, shallow larval stages are likely

to explain why all deep-sea fishes studied to date maintain at least some cone opsins in their

genomes[16].

Most deep-sea fish larvae expressed a single *RH2* gene, but the larvae of some species

(fangtooth, whalefish and lanternfish) expressed multiple cone opsin genes, likely providing

them with similar visual systems to the larvae of shallow-living marine[49] and freshwater

species[28], possibly aiding them in detecting residual light, discriminating brightness and/or

colours. Juvenile deep-sea fishes, on the other hand, showed rod-based expression profiles also

found in the adult stages (Fig. 1). This shift in opsin gene expression correlates with

developmental changes in ecology. As opposed to the adults which are exposed to a narrow

and dim light environment where food is scarce, larvae typically live in well-lit, broad spectrum

shallow waters where food and predators are abundant[11].

3) Functional adaptation in key spectral tuning sites. When multiple *RH1* copies were

expressed, they often showed distinct differences in key amino acid sites that are likely to shift

the spectral sensitivities of the pigments (Fig. 3 and Tables 2 and 3)[16,41]. As such, the three

different *RH1*s in *C. warmingii* differed in 15 key-tuning sites. Opsin gene expression together

with spectral-sensitivity estimations revealed a dominant rod opsin copy (*RH1-1*), and a

shorter- (*RH1-3*) and a longer-shifted (*RH1-2*) copy with lower transcript levels (Figs. 1 and

3). The MSP in *C. warmingii* found two distinct rod types with λ_{max} values of 488 and 468 nm[50]

corresponding most likely to the *RH1-1* and *RH1-3*, respectively. It is possible that multiple

RH1 copies are coexpressed within the same photoreceptor, something that has previously been

reported for cone opsins in shallow-water marine[51,52] and freshwater fishes[53,54]. Coexpression

could produce visual pigment mixtures that shift photoreceptor sensitivity and enhance visual

contrast, aiding in predator-prey interactions or mate finding[53]. Alternatively, a third rod

photoreceptor type might have been overlooked during MSP, which can occur especially if the

cell-type is rare. While the function of having multiple rod opsins in *C. warmingii* remains to

be investigated, several possible benefits for a multi-rod visual system have recently been

proposed including that it might enable conventional or unusual colour vision in dim light, it

might be used to increase visual sensitivity, or enhance object's contrast against a certain

background[16].

In *H. brodiei*, the second *RH1* copy (*RH1-2*) differed in two key tuning sites, E122Q (-15 nm)

and G124S (-11 nm), known to cause major short-wavelength shifts in other fishes[41]. This

supports the MSP measurements in its sister species, *Howella sherborni*, which found two

different rod types with spectral sensitivities of 463 and 492 nm (Fig. 2)[55].

Having multiple differently tuned rod photoreceptors, one centred on the prevailing light

(bioluminescence and/or ambient light ~ 480 – 490 nm) and a second one that is offset from it

(i.e. the offset pigment hypothesis[56]), may be beneficial to break counter illumination - a way

of active camouflage in mesopelagic organisms where ventral photophores emit

bioluminescent light that matches the residual down-welling light[57]. Hence, revealing an

individual's silhouette could help to distinguish prey and predators from the background
lighting, or visually finding mates. Apart of lanternfishes with three (or more) and basslets with
two rod opsins, or exceptional cases of tube-eye (six) and spinyfin (38), the majority of the
deep-sea fishes however seems to have only one rod opsin[16].

273
Conclusions
So far, the development of deep-sea fish vision at the molecular level had not been studied and
only limited morphological information was available. In this study we compared opsin and
visual gene expression between 20 deep-sea fish species revealing a major change in expression
between larval and adult stages. While deep-sea fish genomes contain both cone and rod opsin
genes, larvae rely on the cone pathway and most adult fishes switch to a rod-dominated or rod-
only visual system. The cone vs. rod-specific phototransduction cascade genes follow the
opsins in some lineages, however, not in aulopiforms. Interestingly, we have discovered
lineage-specific opsin gene duplications, possibly allowing for increased visual sensitivity and
certain kind of colour vision in the depth in some species. Our molecular results therefore
support a conserved developmental progression in vertebrates whereby cones appear first in
the retina and rod photoreceptors are added later during development.

Fig. 1: Cone and rod opsin gene expression in larval and adult deep-sea fishes. *A) Transcriptomes were used to characterise the opsin gene expression in the retinas of 20 deep-sea fish species, belonging to eight different orders and mapped onto a simplified teleost phylogeny (topology after Betancur et al.[63]). Boxes highlight the five species for which both larval and adult samples were available. The genomic repertoire of the visual opsins is shown for* Idiacanthus fasciola *(this study), and* Anoplogaster cornuta[16]. *B) Proportional opsin gene expression for each individual (horizontal bar) at different developmental stages. Different colours correspond to cone (colours) or rod (shades of grey) opsin genes, depicted as the proportional expression over the total sum of visual opsins expressed. Different shades of the same colour represent multiple copies of the same gene family. Based on the opsin gene expression, the larvae (left column) show a pure-cone or cone-dominated retina, while the adults (right column) have a pure-rod or rod-dominated visual system. Juveniles specimens in two species had an*

297 *adult expression profile. Note that some species expressed multiple RH1 copies (*Scopelarchus, Howella brodiei

298 *and* Ceratoscopelus warmingii *adults) or multiple RH2 copies (*Gyrinomimus *sp. larva,* Hygophum reinhardtii

299 *larva). Notably, adults and larvae of* Scopelarchus *sp. and* Coccorella atlantica *expressed different copies of RH2*

300 *(more details in Fig. 3). Details about the samples and expression levels are listed in Table 1.*

Fig. 2: Phototransduction cascade gene expression in the retina of five deep-sea fish species. A) Heat map of the expression of individual phototransduction cascade genes for each sample, based on normalised numbers of reads (FPKM). B) Pie charts comparing mean values of relative expression of the opsin genes (rod RH1 and cone RH2), photoreceptor-specific cascade transducin genes (rod-type GNAT1 and cone-type GNAT2), and all cascade genes (photoreceptor-specific transducins, arrestins and phosphodiesterases) summarized. The black square highlights

the two aulopiform species with the discordance between the opsin type (rod-specific) and phototransduction

*cascade genes (cone-specific) in adults. C) Focus on the common fangtooth (*Anoplogaster cornuta*) transitional*

phase shown as a sequence for six larval samples; size given as standard length (SL). Note that all fangtooth

larvae expressed both RH1 and RH2, with an increasing proportion of RH1 to RH2 as the larvae increased in

size. These individuals all had traits of larval phenotypes (dorsal and ventral horns and small teeth; Fig. 1) and

were collected relatively shallow between 0-300 m using the plankton trawls.

Fig. 3*: Gene trees of the A) RH1 and B) RH2 opsin genes mined from the retinal transcriptomes of deep-sea*

fishes. Species with multiple copies are highlighted in colour. Additional gene sequences from public databases

are listed with their GenBank accession numbers. Note the topology within Aulopiformes; the adult RH2s of

Cocorrella atlantica *and* Scopelarchus *cluster together as do the major larval RH2s. Stars mark gene*

duplication events with red stars pinpointing (A) the ancestral duplication of RH1 impacting the Scopelarchus

genus, and (B) the duplication of RH2 in the aulopiform ancestor (or at least the common ancestor of

Coccorella and Scopelarchus). *C) Predicted rod photoreceptor spectral sensitivities based on key spectral*

tuning site mutations in species with multiple rod opsins. Multiple rod opsins differed in three species,

Ceratoscopelus warmingii *(Myctophiformes),* Howella brodiei *(Pempheriformes) and* Scopelarchus michaelsarsi

(Aulopiformes). For functional interpretation we considered also direct microspectrophotometry measurements

from # = Pointer et al.[21], † = Collin & Marshall[50] and ‡ = Partridge et al.[55].

MATERIALS AND METHODS

Specimens used in this study were collected in the Sargasso Sea during two multipurpose fishery surveys conducted by the German Thünen Institute of Fisheries Ecology onboard the research vessel *Walther Herwig III* in March to April in 2014 and in 2017. The sampling of adults occurred during both day and night at depths of 600 – 1'000 m using a mid-water pelagic trawl net (Engel Netze, Bremerhaven, Germany) with an opening of 30 m x 20 m, a length of 145 m, and mesh sizes (knot to knot) from 90 cm decreasing stepwise to 40, 20, 10, 5, 4, 3, 2 cm, with a 1.5-cm mesh in the 27-m-long codend. The larvae were mostly collected using an Isaacs-Kidd Midwater Trawl net (IKMT; 6.2 m^2 mouth-opening, 0.5 mm mesh size; Hydro-Bios Apparatebau GmbH) at depths of 0 - 300 m by double-oblique transect tows. Adult fish were flash-frozen at -80 ^{0}C upon arrival on board. Larval samples were fixed in RNAlater^{TM} (ThermoFisher) and stored at -80 ^{0}C until further use.

Total RNA was extracted from the whole eyes using either the RNeasy micro or mini kit (Qiagen) and the extracted RNA concentration and integrity were subsequently verified on a 2100 Bioanalyzer (Agilent). RNAseq libraries for 31 samples were constructed in-house from unfragmented total RNA using Illumina's NEBNext Ultra II Directional RNA library preparation kit, NEBNext Multiplex Oligos and the NEBNext Poly(A) mRNA Magnetic Isolation Module (New England Biolabs). Multiplexed libraries were sequenced on the Illumina HiSeq 2500 platform as 150 bp paired-end (PE) reads. Library construction and sequencing (150 bp PE) for an additional 10 samples was outsourced to Novogene, Singapore (https://en.novogene.com/). We additionally re-analyzed 11 retinal transcriptomes previously published in Musilová et al.[16]. Together, then, our dataset comprised 52 samples of which, based on morphology, 25 were classified as larvae, 5 as juveniles and 22 as adults. Sample IDs, number of raw reads and further parameters are listed in Table 1.

The sequence data was quality-checked using FastQC[58]. Opsin gene expression was then quantified using Geneious software version 11.0.3[59]. For each sample we first mapped the reads against a general fish reference dataset comprising all visual opsin genes from the Nile tilapia, *Oreochromis niloticus* and the zebrafish, *Danio rerio*, with the Medium-sensitivity settings in Geneious. This enabled us to identify cone and rod opsin specific reads. If present, paralogous genes were subsequently disentangled following the methods in Musilova et al.[16] and de Busserolles et al.[15]. Briefly, we created species-specific references of the expressed opsin genes and their several copies[16] and re-mapped the transcriptome reads with Medium-Low sensitivity

to obtain copy-specific expression levels. If multiple opsin genes were found to be expressed,

we report their proportional expression in relation to the total opsin gene expression (Fig. 1).

We used the same pipeline to quantify expression of phototransduction cascade genes in five

focal deep-sea species (Fig. 2, Tables S1).

To check for key amino-acid substitutions in *RH1* and *RH2* and potential shifts in its

absorbance, we first translated the opsin coding sequences into amino acid sequences, and then

aligned them with the bovine *RH1* (GenBank Acc.No: M12689). We have specifically focused

on the positions identified as key-tuning sites in Yokoyama[41] and Musilova et al.[16]. For details,

see Tables 2 and 3. Unfortunately, we were not able to estimate the exact sensitivity shift in *C.*

warmingii as only four of the amino acids that were substituted at the 15 key-tuning amino acid

sites corresponded with previously tested cases[16,41]. Out of the three copies, *RH1-2* has three

out of four longer-shifting amino acid variants in these four sites and we assume is therefore

red-shifted. *RH1-1* is most likely sensitive to 488 nm, and *RH1-3*, being the shortest, to 468nm.

In order to properly estimate the λ_{max} , we have further checked the samples for expression of

the *Cyp27c1* gene, the factor assuming conversion of the A1 retinal to A2. Expression of

Cyp27c1 would assume that opsin protein bounds to the A2 instead of A1, which would cause

a maximum shift of ca 25-30 nm[60]. The expression of this gene has not been detected

suggesting only A1 retinal in the studied species of the deep-sea fishes (Table S2).

A dataset containing *RH1* opsin gene sequences from our transcriptomes, and additional *RH1*s

obtained from GenBank (link to NCBI) were aligned using the MAFFT[61] plugin as

implemented in Geneious and a phylogenetic tree was subsequently reconstructed using

MrBayes v3.2.1[62] (Fig. 2). Trees were produced using the Markov chain Monte Carlo analysis

which ran for 1 million generations. Trees were sampled every 100 generations, and the

printing frequency was 1000, discarding the first 25% of trees as burn-in. The evolutionary

model chosen was GTR model with gamma-distributed rate variation across sites and a

proportion of invariable sites. Posterior probabilities (PP) were calculated to evaluate statistical

confidence at each node. We used the same approach with an *RH2*-specific reference dataset

to reconstruct the phylogenetic relationship between the transcriptome-derived deep-sea *RH2*

genes (Fig. 3).

To sequence the whole genome of *Idiacanthus fasciola*, we extracted the DNA from a fin clip of one adult specimen using the DNA Tissue kit (QiaGen). Library preparation and genome sequencing (150bp PE) was outsourced to Novogene and we obtained 20 million reads. To search for the opsin genes, genome reads were then mapped against opsin references, as described above in the case of transcriptomes.

Acknowledgements:

We would like to express our thanks to both scientific and technical crew of the Walther Herwig III research cruises in 2014 and 2017. In addition, we thank Tina Blancke for help with the sample management, and Veronika Truhlářová for technical support and lab management. NL and ZM were supported by the Swiss National Science Foundation (PROMYS - 166550) , ZM by the PRIMUS Research Programme (Charles University), the Czech Science Foundation (21-31712S) and the Basler Stiftung fuer Experimentelle Zoologie, and FC by an Australian Research Council (ARC) DECRA Fellowship (DE200100620).

References:

1. Lamb, T. D. Evolution of phototransduction, vertebrate photoreceptors and retina. *Progress in retinal and eye research*, 36, 52-119 (2013).

2. Downes, G. B. & Gautam, N. The G protein subunit gene families. *Genomics*, 62(3), pp.544-552 (1999).

3. Larhammar, D, Nordström K & Larsson, T. A. Evolution of vertebrate rod and cone phototransduction genes. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1531), pp.2867-2880 (2009).

4. Lamb, T. D. Evolution of the genes mediating phototransduction in rod and cone photoreceptors. *Progress in Retinal and Eye Research*, p.100823 (2019).

5. Mears, A. J., et al. Nrl is required for rod photoreceptor development. *Nature genetics*, 29(4), 447-452 (2001).

6. Shen, Y. C. & Raymond, P. A. Zebrafish cone-rod (crx) homeobox gene promotes retinogenesis. *Developmental biology*, 269(1), 237-251 (2004).

7. Sernagor, E., Eglen, S., Harris, B. & Wong, R. (Eds.). Retinal development. Cambridge University Press (2006).

8. Hunt, D. M., Hankins, M.W., Collin, S.P. & Marshall, N.J. Evolution of visual and non-visual pigments (Vol. 4). Boston, MA: Springer (2014).

9. Turner, J. R., White, E. M., Collins, M. A., Partridge, J. C. & Douglas, R. H. Vision in
lanternfish (Myctophidae): adaptations for viewing bioluminescence in the deep-sea." *Deep*
Sea Research Part I: Oceanographic Research Papers 56, 6 (2009).

10. de Busserolles, F., Fogg, L., Cortesi, F. & Marshall, J. The exceptional diversity of visual
adaptations in deep-sea teleost fishes. *Seminars in Cell & Developmental Biology*. Academic
Press (2020).

11. Moser, H. G. & Smith, P. E. Larval fish assemblages and oceanic boundaries. *Bull. Mar.*
Sci, 53(2), 283-289 (1993).

12. Sassa, C. & Hirota, Y. Seasonal occurrence of mesopelagic fish larvae on the onshore
side of the Kuroshio off southern Japan. *Deep Sea Research Part I: Oceanographic Research*
Papers, 81, 49-61 (2013).

13. Zhang, H., et al. Molecular cloning of fresh water and deep-sea rod opsin genes from
Japanese eel *Anguilla japonica* and expressional analyses during sexual maturation. *Febs*
Letters, 469(1), 39-43 (2020).

14. Douglas, R. H., Genner, M. J., Hudson, A. G., Partridge, J. C. & Wagner, H. J.
Localisation and origin of the bacteriochlorophyll-derived photosensitizer in the retina of the
deep-sea dragon fish *Malacosteus niger*. *Scientific reports*, 6, 39395 (2016).

15. de Busserolles, F., et al. Pushing the limits of photoreception in twilight conditions: The
rod-like cone retina of the deep-sea pearlsides. *Science advances*, 3(11), eaao4709 (2017).

16. Musilova, Z., et al. Vision using multiple distinct rod opsins in deep-sea fishes. *Science*,
364(6440), 588-592 (2019).

17. Byun, J. H., et al. Gene expression patterns of novel visual and non-visual opsin families
in immature and mature Japanese eel males. *PeerJ*, 8, e8326 (2020).

18. Munk, O. Changes in the visual cell layer of the duplex retina during growth of the eye of
a deep-sea teleost, *Gempylus serpens* Cuvier, 1829. *Acta Zoologica*, 71(2), 89-95 (1990).

19. Collin, S. P., Hoskins, R. V. & Partridge, J. C. Seven Retinal Specializations in the
Tubular Eye of the Scopelarchus michaelsarsi: A Case Study in Visual Optimization. *Brain,*
Behavior and Evolution, 1998(09), 291–314 (1998).

20. Bozzano, A., Pankhurst, P. M. & Sabatés, A. Early development of eye and retina in
lanternfish larvae. *Visual neuroscience*, 24(3), 423-436 (2007).

21. Pointer, M. A., Carvalho, L. S., Cowing, J. A., Bowmaker, J. K. & Hunt, D. M. The
visual pigments of a deep-sea teleost, the pearl eye *Scopelarchus analis*. *Journal of*
Experimental Biology, 210(16), 2829-2835 (2007).

22. Biagioni, L. M., Hunt, D. M. & Collin, S. P. Morphological characterization and
topographic analysis of multiple photoreceptor types in the retinae of mesopelagic
hatchetfishes with tubular eyes. *Frontiers in Ecology and Evolution*, 4, 25 (2016).

23. Carleton, K. L., Escobar-Camacho, D., Stieb, S. M., Cortesi, F. & Marshall, N. J.
Seeing the rainbow: mechanisms underlying spectral sensitivity in teleost fishes. *Journal of*
Experimental Biology, 223(8) (2020).

24. Valen, R., et al. The two-step development of a duplex retina involves distinct events of
cone and rod neurogenesis and differentiation. *Developmental biology*, 416(2), 389-401
(2016).

25. La Vail, M. M., Rapaport, D. H. & Rakic, P. Cytogenesis in the monkey retina. *J. Comp.*
Neurol. 309, 86-114. doi:10.1002/cne.903090107 (1991).

26. Raymond, P. A. Development and morphological organization of photoreceptors. In
Neurobiology and Clinical Aspects of the Outer Retina (pp. 1-23). Springer, Dordrecht (1995).

27. Denton, E. J. Light and vision at depths greater than 200 metres. Light and life in the sea
(1990): 127-148.

28. Carleton, K. L., Dalton, B. E., Escobar-Camacho, D. & Nandamuri, S. P. Proximate
and ultimate causes of variable visual sensitivities: insights from cichlid fish radiations.
Genesis, 54(6), 299-325 (2016).

29. Cortesi, F., et al. Ancestral duplications and highly dynamic opsin gene evolution in
percomorph fishes. *Proceedings of the National Academy of Sciences* 112.5 (2015).

30. Cortesi, F., et al. From crypsis to mimicry: changes in colour and the configuration of the
visual system during ontogenetic habitat transitions in a coral reef fish. *Journal of Experimental*
Biology, 219(16), 2545-2558 (2016).

31. Bowmaker, J. K., Hunt, D. M., & Jeffery, G. Eel visual pigments revisited: The fate of
retinal cones during metamorphosis. *Visual neuroscience*, 25(3), 249 (2008).

32. Tettamanti, V., de Busserolles, F., Lecchini, D., Marshall, N.J. & Cortesi, F. Visual
system development of the spotted unicornfish, *Naso brevirostris* (Acanthuridae). Journal of
Experimental Biology, 222(24) (2019).

33. Simoes, B. F., et al. Multiple rod–cone and cone–rod photoreceptor transmutations in
snakes: evidence from visual opsin gene expression. *Proceedings of the Royal Society B:*
Biological Sciences, 283(1823), 20152624 (2016).

34. Schott, R. K., Bhattacharyya, N. & Chang, B. S. Evolutionary signatures of
photoreceptor transmutation in geckos reveal potential adaptation and convergence with
snakes. *Evolution*, 73(9), 1958-1971 (2019).

35. Mariani, A. P. Photoreceptors of the larval tiger salamander retina. *Proceedings of the*
Royal society of London. Series B. Biological sciences 227.1249 (1986)

36. Wagner, H. J., Partridge, J. C. & Douglas, R. H. Observations on the retina and 'optical
fold' of a mesopelagic sabretooth fish, *Evermanella balbo*. *Cell and Tissue Research*, 378(3),
411-425 (2019).

37. Baylor, D. A. Photoreceptor signals and vision. Proctor lecture. *Investigative*
ophthalmology & visual science 28.1 (1987)

38. Kawamura, S. & Tachibanaki, S. Explaining the functional differences of rods versus
cones. *Wiley Interdisciplinary Reviews: Membrane Transport and Signaling* 1.5 (2012)

39. Luo, DG., et al. "Apo-Opsin and Its Dark Constitutive Activity across Retinal Cone
Subtypes." *Current Biology* 30.24 (2020)

40. Cortesi, F., et al. Visual system diversity in coral reef fishes. *Seminars in Cell &*
Developmental Biology. Academic Press (2020).

41. Yokoyama, S. Evolution of dim-light and color vision pigments. *Annu. Rev. Genomics*
Hum. Genet., 9, 259-282 (2008).

42. Yokoyama, S. & Jia, H. Origin and adaptation of green-sensitive (RH2) pigments in
vertebrates. *FEBS Open Bio*, 10(5), 873-882 (2020).

43. Luehrmann, M. et al. Cardinalfishes (Apogonidae) show visual system adaptations typical
of nocturnally and diurnally active fish. *Molecular ecology* 28.12 (2019).

44. Liu, D.. et al. The cone opsin repertoire of osteoglossomorph fishes: gene loss in mormyrid
electric fish and a long wavelength-sensitive cone opsin that survived 3R. *Molecular biology*
and evolution 36.3 (2019).

45. Huang, Z., Titus, T., Postlethwait, J. H., & Meng, F. Eye Degeneration and Loss of
otx5b Expression in the Cavefish *Sinocyclocheilus tileihornes*. *Journal of molecular evolution*
87.7 (2019).

46. Hunt, D. M., Fitzgibbon, J., Slobodyanyuk, S. J., Bowmaker, J. K. & Dulai, K. S.
Molecular evolution of the cottoid fish endemic to Lake Baikal deduced from nuclear DNA
evidence. *Molecular Phylogenetics and Evolution*, 8(3), 415-422 (1997).

47. Sugawara, T., et al. Parallelism of amino acid changes at the RH1 affecting spectral
sensitivity among deep-water cichlids from Lakes Tanganyika and Malawi. *Proceedings of the*
National Academy of Sciences, 102(15), 5448-5453 (2005).

48. Musilova, Z., et al. Evolution of the visual sensory system in cichlid fishes from crater
lake Barombi Mbo in Cameroon. *Molecular Ecology*, (August), 5010–5031 (2019).

49. Britt, L. L., Loew, E. R. & McFarland, W. N. Visual pigments in the early life stages of
Pacific northwest marine fishes. *Journal of Experimental Biology* 204.14 (2001).

50. Collin, S. P. & Marshall, N. J. Sensory Processing in Aquatic Environments. Springer-
Verlag New York (2003).

51. Savelli, I., Flamarique, I. N., Iwanicki, T. & Taylor, J. S. Parallel opsin switches in
multiple cone types of the starry flounder retina: tuning visual pigment composition for a
demersal life style. *Scientific reports* 8, no. 1 (2018).

52. Stieb, S. M., et al. A detailed investigation of the visual system and visual ecology of the
Barrier Reef anemonefish, *Amphiprion akindynos*. *Scientific reports* 9.1 (2019).

53. Dalton, B. E., Loew, E. R., Cronin, T. W. & Carleton, K. L. Spectral tuning by opsin
coexpression in retinal regions that view different parts of the visual field. *Proceedings of the
Royal Society B: Biological Sciences* 281,1797 (2014).

54. Torres-Dowdall, J., et al. Rapid and parallel adaptive evolution of the visual system of
Neotropical Midas cichlid fishes. *Molecular biology and evolution* 34.10 (2017).

**55. Partridge, J. C., Shand, J., Archer, S. N., Lythgoe, J. N., & van Groningen-Luyben,
W. A. H. M.** Interspecific variation in the visual pigments of deep-sea fishes. *Journal of
Comparative Physiology A*, 164(4), 513-529 (1989).

56. Lythgoe, J.N. Visual pigments and under- water vision. In: Light as an Ecological Factor
(Rackham, O., ed.). Oxford: Blackwell (1966).

57. Denton, E. J., Herring, P. J., Widder, E. A. Latz, M. F. & Case, J. F. The roles of filters
in the photophores of oceanic animals and their relation to vision in the oceanic environment.
Proceedings of the Royal Society of London. Series B. Biological Sciences 225, 1238 (1985).

58. Andrews, S. FastQC: a quality control tool for high throughput sequence data. 2010 (2017).

59. Kearse, M., et al. Geneious Basic: an integrated and extendable desktop software platform
for the organization and analysis of sequence data. Bioinformatics, 28(12), 1647-1649 (2012).

60. Enright, J. M., et al. Cyp27c1 red-shifts the spectral sensitivity of photoreceptors by
converting vitamin A1 into A2. *Current Biology*, 25(23), 3048-3057 (2015).

61. Katoh, K., Asimenos, G. & Toh, H. Multiple alignment of DNA sequences with MAFFT.
In Bioinformatics for DNA sequence analysis (pp. 39-64). Humana Press (2009).

62. Ronquist, F., et al. MrBayes 3.2: efficient Bayesian phylogenetic inference and model
choice across a large model space. *Systematic biology*, 61(3), 539-542 (2012).

63. Betancur-R, R., et al. Phylogenetic classification of bony fishes. *BMC evolutionary
biology*, 17(1), 162 (2017).

64. Douglas, R. H., Partridge, J. C. & Marshall, N. J. The visual systems of deep-sea fish.

I. Optics, tapeta, visual and lenticular pigmentation. *Prog. Ret. Eye Res*, 17(4), 597-636 (1998).

65. Kenaley, C. P., DeVaney, S. C., & Fjeran, T. T. The complex evolutionary history of

seeing red: Molecular phylogeny and the evolution of an adaptive visual system in deep-sea

dragonfishes (Stomiiformes: Stomiidae). *Evolution*, 68(4), 996-1013 (2014).

Table 1: Samples used in the study and results of the opsin gene expression in the eyes or retina

species	order	stage	code	size	date of collection	raw reads	reads after bacteria filtering	RH1	RH2	SWS1	SWS2	accession number
Anoplogaster cornuta	Trachichthyiformes	Larva	39016	4mm		36,373,372	36,125,630	<0.01	0.99		<0.01	
		Larva	39017	4mm		34,712,538	34,604,364	<0.01	0.99		<0.01	
		Larva	4_23	9mm		41,592,840	41,519,070	0.02	0.98		<0.01	
		Larva	4_22	11mm		75,109,270	74,990,214	0.11	0.89		<0.01	
		Larva	4_21	11-12 mm		28,945,804	28,869,552	0.22	0.78		<0.01	
		Larva	261s03	12mm		48,458,044	48,457,472	0.47	0.53		<0.01	
		Adult	56H6			11,658,040	11,635,610	1				
Ceratoscopelus warmingii	Myctophiformes	Adult	300s03	63mm		49,078,974	44,114,639	$0.66^{RH1\text{-}1}$ $0.01^{RH1\text{-}2}$ $0.33^{RH1\text{-}3}$				
		Adult	S1			8,796,786	8,789,961	$0.94^{RH1\text{-}1}$ $0.02^{RH1\text{-}2}$ $0.04^{RH1\text{-}3}$				
		Adult	S2			8,655,248	8,649,342	$0.93^{RH1\text{-}1}$ $0.02^{RH1\text{-}2}$ $0.05^{RH1\text{-}3}$				
		Adult	S3			7,904,714	7,899,617	$0.94^{RH1\text{-}1}$ $0.01^{RH1\text{-}2}$ $0.05^{RH1\text{-}3}$				
Chauliodus sloani	Stomiiformes	Larva	67I2_2	11mm		34,445,252	34,371,394	0.01	0.99			
		Larva	67I1	17mm		20,701,686	20,683,764	<0.01	0.99			
		Adult	109B6	20mm		24,854,866	24,835,646	1				
Chauliodus danae	Stomiiformes	Juvenile	109C7	24mm		32,865,334	32,834,300	1				
		Juvenile	109A2	20mm		25,389,318	25,344,182	1				
		Juvenile	109D7			18,540,248	18,533,484	1				
Coccorella atlantica	Aulopiformes	Larva	109G8			26,376,620	26,359,060	0.01	$0.99^{RH2\text{-}1}$			
		Adult	56C7			36,794,070	36,790,854	0.69	$0.31^{RH2\text{-}2}$			
		Adult	297_7	68mm		43,948,534	40,511,279	0.96	$0.04^{RH2\text{-}3}$			
Grammatostomias flagellibarba	Stomiiformes	Adult	56H8			9,775,520	9,742,864	1				
Gyrinomimus myersi	Beryciformes	Larva	S25			19,270,096	19,122,986	<0.01	$0.45^{RH2\text{-}1}$ $0.13^{RH2\text{-}2}$ $0.08^{RH2\text{-}3}$ $0.11^{RH2\text{-}4}$ $0.15^{RH2\text{-}5}$	0.07	<0.01	
Howella brodiei	Pempheriformes	Adult	56D8			56,905,264	43,936,178	$0.17^{RH1\text{-}1}$ $0.83^{RH1\text{-}2}$				
		Adult	56D9			61,909,674	61,590,270	$0.07^{RH1\text{-}1}$ $0.93^{RH1\text{-}2}$				
Hygophum reinhardtii	Myctophiformes	Larva	67I2_1	5mm		17,222,626	17,122,880		$0.26^{RH2\text{-}1}$ $0.68^{RH2\text{-}2}$ $0.06^{RH2\text{-}3}$			
Idiacanthus fasciola	Stomiiformes	Larva	71C2			23,382,468	23,379,360		1			
		Larva	67D2			6,647,384	6,623,496		1			
		Larva	71C1			23,632,774	23,497,715		1			
		Larva	67I7	25mm		25,713,960	25,701,768		1			
		Larva	71B9			43,695,620	43,633,424		1			
		Larva	67B8			28,333,642	28,313,024		1			
		Larva	109A1	41mm		27,834,274	27,818,750		1			
		Adult	228s01			21,704,528	21,622,274	1				
		Adult	67F8			15,744,604	15,739,004	1				
		Adult	67B6			11,368,044	15,439,954	1				
Melamphaes sp.	Beryciformes	Juvenile	4_26			38,583,370	38,534,507	1				
		Juvenile	4_28			35,489,970	35,431,421	1				
Melanolagus bericoides	Argentiniformes	Larva	71H3			26,219,646	26,210,402		1			
Pteraclis aesticola	Scombriformes	Larva	109H4			19,918,922	19,870,484	0.22	0.78			
Pollichthys mauli	Stomiiformes	Adult	109I2			55,767,498	54,875,747	1				
		Adult	109I3			68,655,686	67,082,709	1				
Scombrolabrax heterolepis	Scombriformes	Adult	,56E3			65,770,014	5,527,135	0.98	0.02			
		Adult	,56E4			60,643,652	60,472,479	0.93	0.07			
Scopelarchus guentheri	Aulopifoformes	Larva	109H1			37,542,802	37,416,486		0.96^{RH2_larval} $0.04^{RH2\_2nd\_larval}$			
Scopelarchus michaelsarsi	Aulopifoformes	Larva	109B9	21mm		9,971,250	9,919,148		1^{RH2_larval}			
		Larva	71B7			43,695,620	43,682,534	$<0.01^{RH1\text{-}1}$ $<0.01^{RH1\text{-}2}$	0.96^{RH2_larval} $0.03^{RH2\_2nd\_larval}$			
		Adult	272_10	66mm		45,565,202	43,729,338	$0.80^{RH1\text{-}1}$ $0.10^{RH1\text{-}2}$	0.10^{RH2_adult}			
Scopelogadus mizolepis	Beryciformes	Adult	,57E2			5,160,876	5,160,876	1				
		Adult	300s01			15,448,240	15,420,650	1				
Scopelosaurus hoedti	Aulopiformes	Adult	297_9	69mm		43,948,534	42,248,053	0.98	0.02			
Vinciguerria poweriae	Stomiiformes	Larva	109H2			25,863,628	25,755,100		1			
		Larva	109C8	16mm		22,914,004	22,858,398		1			

Table 2: key-tuning amino acid sites in the cone opsin RH2 gene

species	order	83	122	207	255	292	lmax	ref.
Bovine RH1		D	E	M	I	A	500 nm	41
Ancestral teleost		D	Q	M	I	A	488 nm	42
Melanolagus bericoides	Argentiniformes	G	Q	.	V	.		
Coccorella atlantica adult	Aulopiformes	G	Q	.	.	.		
Coccorella atlantica larval	Aulopiformes	G	Q	.	V	.		
Scopelarchus michaelsarsi adult	Aulopiformes	?	Q	I	C	T		
Scopelarchus guentheri larval	Aulopiformes	G	Q	.	V	.		
Scopelarchus michaelsarsi larval	Aulopiformes	G	Q	.	V	.	505 nm	21
Scopelarchus guentheri 2nd larval	Aulopiformes	G	Q	.	C	.		
Scopelarchus michaelsarsi 2nd larval	Aulopiformes	G	Q	.	C	.		
Scopelosaurus hoedti	Aulopiformes	G	Q	.	.	.		
Gyrinomimus myersi RH2-1	Beryciformes	G	Q	.	F	.		
Gyrinomimus myersi RH2-2	Beryciformes	G	Q	L	F	.		
Gyrinomimus myersi RH2-3	Beryciformes	G	Q	L	F	.		
Gyrinomimus myersi RH2-4	Beryciformes	G	Q	L	F	.		
Gyrinomimus myersi RH2-5	Beryciformes	G	Q	L	F	.		
Lepisosteus platyrhincus	Lepisosteiformes	G	.	.	.	.		
Hygophum reinhardtii RH2-1	Myctophiformes	G	Q	.	V	.		
Hygophum reinhardtii RH2-2	Myctophiformes	G	Q	.	V	.		
Hygophum reinhardtii RH2-3	Myctophiformes	G	Q	.	V	.		
Lepidopus fitchi RH2-1	Scombriformes	G	.	.	V	.	496 nm	42
Lepidopus fitchi RH2-2	Scombriformes	G	Q	.	V	.		
Lepidopus fitchi RH2-3	Scombriformes	G	Q	.	V	.	506 nm	42
Lepidopus fitchi RH2-4	Scombriformes	G	Q	.	V	.		
Pteraclis aesticola	Scombriformes	G	Q	L	.	.		
Scombrolabrax heterolepis	Scombriformes	G	Q	.	.	.		
Aristostomias scintillans	Stomiiformes	G	Q	L	V	.	468 nm	42
Chauliodus sp.	Stomiiformes	G	Q	.	F	.		
Grammastomias flagellibarba	Stomiiformes	G	Q	.	.	.		
Idiacanthus fasciola	Stomiiformes	G	Q	.	V	.		
Vinciguerria poweriae	Stomiiformes	G	Q	.	F	.		
Anoplogaster cornuta	Trachichthyiformes	G	Q	.	.	.		

Table 3: key-tuning amino acid sites in the rhodopsin RH1 gene

Species	Order	83	90	96	102	111	113	118	122	124	132	164	183	188	194	195	207	208	211	214	253	261	265	269	289	292	295	299	300	317	lmax	ref.
bovine RH1		D	G	Y	Y	N	E	T	E	A	A	A	M	G	P	H	M	F	H	I	M	F	W	A	T	A	A	A	V	M	500 nm	41
ancestral teleost RH1		D	G	Y	Y		E	T	E	A	A	A	M		L	N	M	F	H	I	M	F	W	A	T	A	A	A	L	M		16
Bathysaurus ferox	Aulopiformes	N	.	.	.	.	.	.	.	S	.	.	.	.	R	A	.	.	.	.	.	.	.	.	.	S	.	T	L	.	481 nm	50
Bathysaurus mollis	Aulopiformes	N	.	.	.	.	.	.	.	S	.	.	.	.	R	A	.	.	.	.	.	.	.	.	T	S	.	T	L	.	479 nm	50
Coccorella atlantica	Aulopiformes	N	.	.	S	.	.	M	.	.	.	.	.	.	R	A	.	.	.	C	.	.	.	.	.	S	.	.	I	.	480 nm‡	64
Scopelarchus spp. RH1-1	Aulopiformes	N	.	.	.	.	.	.	.	.	.	.	.	.	R	A	.	.	.	C	.	.	.	.	.	S	S	S	I	.	486 nm	21
Scopelarchus spp. RH1-2	Aulopiformes	N	.	.	.	.	.	.	.	S	.	.	.	.	L	K	.	.	.	V	.	.	.	.	.	S	.	.	L	.	479 nm	21
Scopelosaurus hoedti	Aulopiformes	.	.	.	.	.	.	.	.	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	.	S	S	S	.	.		
Gyrinomimus myersi	Beryciformes	N	.	.	.	.	.	.	.	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	.	S	.	.	I	.		
Melamphaes sp.	Beryciformes	.	.	.	.	.	.	.	Q	.	S	.	.	.	R	V	.	.	.	.	.	.	.	G	.	S	.	S	I	.		
Scopelogadus mizolepis	Beryciformes	.	.	.	.	.	.	.	Q	G	S	.	.	.	R	V	.	.	.	.	.	.	.	G	T	.	.	S	.	.	488 nm§	64
Lepisosteus oculatus	Lepisosteiformes	.	.	.	.	.	.	.	Q	S	.	.	.	.	L	K	.	.	.	L	.	Y	.	G	.	.	.	.	L	.		
Ceratoscopelus warmingii RH1-1	Myctophiformes	.	.	.	.	T	.	.	Q	.	S	.	.	.	R	A	.	.	.	.	.	.	.	G	.	.	.	S	I	.	?468 nm†	50
Ceratoscopelus warmingii RH1-2	Myctophiformes	.	.	W	.	I	.	.	I	T	.	.	.	.	R	A	.	.	.	V	.	Y	.	.	.	.	.	.	I	.	prob. 488 nm†	50
Ceratoscopelus warmingii RH1-3	Myctophiformes	.	.	.	.	A	.	.	Q	.	S	S	.	.	N	V	.	.	Y	.	.	.	.	G	.	S	.	.	I	.	?468 nm†	50
Hygophum reinhardtii	Myctophiformes	.	.	.	.	A	.	.	Q	.	.	.	.	.	R	A	.	.	.	.	.	.	.	G	.	.	.	.	I	.		
Howella brodiei RH1-1	Pempheriformes	N	.	.	.	.	.	.	.	.	.	.	.	.	R	A	.	.	.	.	.	.	.	G	.	S	S	S	I	.		
Howella brodiei RH1-2	Pempheriformes	N	.	.	.	.	.	.	Q	S	.	.	.	.	R	A	.	.	.	A	.	.	.	.	.	S	S	S	I	.		
Pteraclis aesticola	Scombriformes	.	.	.	.	.	.	.	Q	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	.	.	.	S	I	.		
Scombrolabrax heterolepis	Scombriformes	.	.	.	.	.	.	.	Q	.	.	.	.	.	R	A	.	.	.	.	.	.	.	S	.	S	.	S	I	.		
Chauliodus spp.	Stomiiformes	N	.	.	.	.	.	.	G	.	.	.	.	.	R	A	.	.	.	V	.	.	.	.	A	S	.	.	L	V	484 nm	50
Grammatostomias flagellibarba	Stomiiformes	N	.	.	.	.	.	.	G	.	.	.	.	.	R	A	.	.	.	V	.	.	.	.	.	S	.	.	L	.	480 - 487 nm	65
Idiacanthus fasciola	Stomiiformes	N	.	.	.	.	.	.	.	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	.	S	.	.	L	.	485 nm	50
Pollichthys mauli	Stomiiformes	N	.	.	.	.	.	.	.	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	.	S	.	.	L	.	485 nm	50
Anoplogaster cornuta	Trachichthyiformes	N	.	.	.	.	.	.	.	.	.	.	.	.	R	A	.	.	.	.	.	.	.	.	T	S	.	S	I	.	485 nm	50

† = two pigments reported without assignment to the gene; see also Figure 3

‡ = for *Evermannella balbo* ; sequence not available

§ = for *Scopelogadus beani* ; sequence not available