A network linking scene perception and spatial memory systems in posterior cerebral cortex.

Adam Steel[1], Madeleine M. Billings[1], Edward H. Silson[2], Caroline E. Robertson[1]

[1]Department of Psychology and Brain Sciences, Dartmouth College, Hanover, NH, 03755

[2]Psychology, School of Philosophy, Psychology, and Language Sciences, University of Edinburgh,
Edinburgh, UK EH8 9JZ

Corresponding author: Adam Steel, Department of Psychology and Brain Sciences, Dartmouth
College, 3 Maynard Street, Hanover, NH, 03755; email: adam.steel@dartmouth.edu; tel: (202)
640 9340

Abstract word count: 101

Main text word count: 4750

Figures: 5

Extended data figures: 15

Extended data files: 5

Acknowledgements: The authors thank Ratan Murty, Iris Groen, Martin Hebart, and Brad
Duchaine for comments on drafts of this manuscript, and Chris Baker, Nancy Kanwisher, Laura
Lewis, and Michael Cohen for helpful discussion. We also thank Tommy Botch, Yeo Bi Choi, and
Anna Mynick for assistance with data collection. This work was supported by a grant from NVIDIA
to CER. AS is supported by the Neukom Institute for Computational Science.

Author contribution statement: AS and CER conceived the idea and designed the research. AS
collected the data. AS and MMB analyzed the data. AS wrote the initial manuscript draft. AS, EHS,
and CER wrote the paper.

Conflict of interest statement: The authors declare no conflict of interest

ABSTRACT

Here, we report a network of brain areas bridging the spatial-memory and scene-perception
systems of the human brain. Using fine-grained individual-subject fMRI, we reveal three cortical
areas of the human brain, each lying immediately anterior to a region of the scene perception
network in posterior cerebral cortex, that selectively activate when recalling familiar real-world
locations. Despite their close proximity to the scene-perception areas, network analyses show
that these regions constitute a distinct functional network that interfaces with memory systems
during naturalistic scene understanding. These "place-memory areas" offer a new framework for
understanding how the brain implements memory-guided visual behaviors, including navigation.

MAIN TEXT

Introduction

As we navigate through our world, the visual scene in front of us is seamlessly integrated with
our memory of the broader spatial environment. The neural systems supporting visual scene
processing in posterior cerebral cortex[1–9] and spatial memory in the hippocampus and medial
temporal lobe[10–18] are well-described. But how do visual and spatio-mnemonic systems interface
in the brain to give rise to memory-guided visual experience?

Two disparate lines of inquiry yield different hypotheses. On the one hand, mechanistic accounts
of memory often posit that explicit recall of visual stimuli reinstates perceptual representations
in visual areas[19–23], including the three areas of the scene-perception network (parahippocampal
place area (PPA[2]), occipital place area (OPA; also referred to as the transverse occipital sulcus[3,24–]
[27]), and medial place area (MPA; also referred to as retrosplenial complex[8,9,19,28–31]). However,
recent studies question the co-localization of perceptual and memory-based representations and
instead suggest a perception to memory transition moving anteriorly from areas of the scene-
perception netwok[1,32–36]. These later findings are consistent with neuropsychological
observations, where patients with intact perception but disrupted mental imagery, and *vice*
versa, have been described[37–39]. Resolving this discrepancy is critical to understanding how
contextual information from memory is brought to bear on visual representations in the brain.
In short, do scene perception and memory share common neural substrates?

Here, we sought to describe the neural basis of perceptually- and mnemonically-driven
representations of real-world scenes in the human brain. To do this, we conducted a series of
experiments using fine-grained, individual-subject fMRI to assess whether these responses are
co-localized in the brain (Experiment 1). Surprisingly, this experiment revealed strong evidence

to the contrary: in all subjects, visually recalling real-world locations evoked activity in three
previously undescribed areas in posterior cerebral cortex, each lying immediately anterior to one
of the three regions of the scene-perception network. We next characterized the functional and
network properties of these "place-memory areas" (Experiments 2-3) and explicitly tested the
wide-spread view that the neural substrates of perception and memory recall (i.e., mental
imagery) activate shared regions of the human brain (Experiment 4). Together, our results reveal
a network of brain areas that collectively bridge the scene perception and spatial memory
systems of the human brain and may facilitate integration of the local visual environment with
spatial memory representations.

Results

Distinct topography of scene-perception and place-memory activity
In Experiment 1, we mapped the topography of perceptual (henceforth "scene-perception") and
mnemonic (henceforth "place-memory") activity related to real-world scene processing in 14
adult participants. We first independently localized the three regions of the scene-perception
network[1] by comparing activation when participants viewed images of scenes, faces, and objects
(Methods). Then, in the same individuals, we localized areas that showed preferential BOLD-
activation when participants recalled personally familiar real-world locations (e.g. their house)
versus faces (e.g. their mother)[33]. We subsequently compared the topography (i.e. anatomical
location and spatial extent) of the scene-perception and place-memory preferring areas.

Comparison of scene-perception and place-memory activation revealed a striking topographical
pattern: in all participants, we found three clusters of place-memory activity in posterior cerebral
cortex, each paired with one of the three scene-perception regions (Fig. 1). We henceforth refer
to these clusters as 'place-memory areas' for brevity. On the lateral, ventral, and medial cortical
surfaces, the pairs of place-memory and scene-perception areas exhibited a systematic
topographical relationship: the center-of-mass of each place-memory area was located
consistently anterior to the center-of-mass of its corresponding scene-perception area in every
individual participant (Lateral pairs: $t(13)=16.41$, $p<0.0001$, $d=4.39$; Ventral pairs: $t(13)=12.115$,
$p<0.0001$, $d=3.24$; Medial pairs: $t(13)=5.99$, $p<0.0001$, $d=1.6$; Extended data Fig. 1a;
Supplementary File 1). Importantly, control analyses using 1) a more conservative contrast
(scenes > faces + objects) to define scene-perception areas[40,41] and 2) a probabilistic ROI-based
definition of PPA[41] confirmed that the anterior shift of memory relative to perception was not
due to our definition of the scene-perception areas (Extended data Figs. 2 & 3).

Across the cortical surfaces, the degree of overlap between the scene-perception and place-
memory areas varied systematically (Fig. 1, inset). On the ventral surface, place-memory

activation emerged rapidly at the anterior 60-70% of the scene-perception area (PPA) and
continued anteriorly beyond the scene-perception area's anterior extent (Extended data Fig. 4).
In contrast, on the medial surface, the scene-perception area (MPA) was encompassed within
the most posterior portion of the place-memory responsive region. On the lateral surface, the
place-memory and scene-perception area (OPA) partially overlapped, but to a lesser extent than
on the other cortical surfaces. This systematic variation across the cortical surfaces was present
in all participants (Extended data figure 1a, Extended data file 1).

Fig. 1. Distinct topography of place-memory and scene-perception activity in posterior cerebral cortex. In all
participants, three place-memory areas were observed, each located significantly anterior to one region of the scene-
perception network. One example participant in Experiment 1 is shown (See Extended data Fig. 1 and Extended data
file 1 for thresholded and unthresholded activation maps for all participants (n=14)). The participant's scene
perception ROIs are outlined in white, and place memory activity is shown in warm colors. The scene-perception
network (parahippocampal place area [PPA], occipital place area [OPA], and medial place area [MPA]) was localized
by comparing the BOLD response when participants viewed images of scenes versus with faces (outlined in white,
thresholded at vertex-wise p < 0.001). Place-memory areas on each surface were localized in separate fMRI runs by
comparing the BOLD response when participants recalled personally familiar places versus people (warm colors,
thresholded at vertex-wise p < 0.001). Polar plots: for each cortical surface, the center of mass of place-memory
activation was significantly anterior to the center of mass of scene-perception activation in all participants (all ts > 5,
p < 0.001). In contrast, face memory activation was spatially co-localized with the face-selective fusiform face area
(FFA) on the ventral surface, and no anterior shift was observed (cool colors, $t_9 = 0.1211$, p = 0.906). Statistical
analyses revealed no difference between the hemispheres, so, for clarity, only right hemisphere is shown. Inset: While
the activation during place memory was systematically anterior to activation during scene perception, the spatial
overlap between perception and memory activation varied across the cortical surfaces. Note that the axes (posterior-
anterior) of each polar plot is aligned to its associated cortical surface. The data in the polar plot reflects distance in
millimeters.

We performed two additional analyses to characterize the cortical locations of the place memory
areas. First, we asked: do the locations of the place-memory areas fall in retinotopic, or non-
retinotopic cortex? We hypothesized that, unlike scene perception areas, which process visual
input, place memory areas would lie in non-retinotopic cortex. To test this, we performed a group
analysis of the place memory localizer and compared the peak activity for each area to
retinotopic parcels based on a probabilistic atlas[42]. Strikingly, on all cortical surfaces, place-
memory areas were consistently located anterior to the cortical retinotopic maps (Figure 2a) with
minimal overlap (Extended data Fig. 5). This location in non-retinotopic cortex is potentially
consistent with their role as preferentially mnemonic, rather than visual, areas. Second, we
asked: how do the locations of the place-memory areas relate to known anatomical landmarks
in the brain? To address this question, we compared the results of the group place-memory
localizer to anatomical parcels from the Glasser atlas[43]. This revealed a remarkable pattern: the
peaks of the place-memory preferring activity fell at the intersection of areas known to be
important for visual and spatial processing. Medially, the peak fell within the parietal occipital
sulcus area 1 near the retrosplenial cortex; ventrally, the peak fell between parahippocampal
area 1 and the presubiculum, and laterally the peak fell within area PGp and PGi (Figure 2b). Thus,
the place-memory areas straddle visual and spatio-mnemonic functional areas, and, based on
this localization, these areas are poised to play a role in bridging these processes. In contrast, the
scene-perception areas fell in retinotopic cortex, and areas associated with perception (Extended
data Fig. 6).

Retintopic atlas (Wang et al. 2014)

Memory for places vs. people
t > 7.3, p = 1e-6, q = 1.5e-4

Anatomical
Parcellation (Glasser et al. 2016)

Fig. 2. Location of place-memory activity in relation to known functional and anatomical landmarks. a. The place-memory areas are anterior to and have minimal overlap with retinotopic cortex. We conducted a group analysis of the place-memory localizer (place > people memory recall; thresholded at vertex-wise t > 7.3, p = 1e-6) and compared the resulting activation to the most probable location of the cortical retinotopic maps using the atlas (overlaid) from Wang et al. 2014. The peak of place memory activity was anterior to the retinotopic maps on the lateral and ventral surfaces and there was very little overlap between place-memory activity and retinotopic maps. b. The place-memory areas fall at the intersection between anatomical parcels known to be involved in visual processing and those associated with spatial memory. Comparing the peaks of place-memory activity with parcels from Glasser et al. 2016[43] (overlaid) revealed that the place-memory areas fell at the intersection of parcels associated with visual and spatial processing. Activation maps are replotted in panels a and b to allow comparison between parcellations.

In contrast to the striking distinction between scene-perception and place-memory, memory-related activation for faces was centered on the perceptually-driven FFA on the ventral surface (Difference of centers-of-mass: t(9)=0.1211, p=0.906, d=0.17), and we did not observe consistent memory-driven activation near the occipital face area on the lateral surface. The anterior shift for places was greater than that for faces on all surfaces (Main effect ROI: F(3,85)=14.842, p<0.001, t vs faces: all ps<0.001, all ds > 2.26). Importantly, face and place memories are complex and multifaceted, and factors such as vividness and memory age can influence the activity elicited by memory recall[44–46]. However, the difference between categories was not due to subjective

differences in recall ability, as participants reported equal subjective vividness for place and face
memory recall, although vividness of recollection was not related to activation magnitude in
place-memory or scene-perception areas (Extended data Fig. 7). We reasoned that one potential
explanation for the anterior distinction of memory versus perception for scenes/places but not
faces might be the BOLD signal-dropout artifact affecting lateral ventral temporal cortex.
However, we ruled out this explanation by replicating Experiment 1 using an advanced multi-
echo fMRI sequence that improved signal from the lateral and anterior temporal lobe and
obtaining similar results: an anterior shift for place-memory relative to scene-perception, but no
distinction between the location of face-memory relative to face-perception activity (Extended
data Fig. 1b). All in all, the consistent topographical relationship between perception and memory
observed for scenes was not observed for faces.

These findings show that place memory recall engages three distinct brain areas that are
systematically paired with the scene-perception areas. We next sought to characterize the
functional and network properties of these areas, specifically to test whether scene-perception
and place-memory areas were functionally, as well as anatomically, dissociable. To ensure that
we evaluated functionally homogenous regions and to control for differences in ROI sizes across
regions, we constrained the scene-perception and place-memory areas to the unique members
of the top 300 most scene-perception/place-memory preferring vertices for all subsequent
experiments. Importantly, on all cortical surfaces, the constrained place-memory area was
anterior to the scene-perception area (Extended data Fig. 8).

Place-memory areas respond preferentially to familiar places
In Experiment 1, we identified three place memory areas that showed category-selectivity for
remembered places as compared with perceived scenes. But are these areas only driven by top-
down, constructive memory recall tasks, or more broadly driven by memory tasks, including tasks
that rely less on top-down signals, such as recognition memory? We examined this question in
Experiment 2. In this experiment, participants provided a list of real-world locations that were
familiar to them. Then, in the scanner, participants performed a covert recognition task, wherein
they were shown panning videos of their personally familiar locations or unfamiliar locations
taken from another participant's familiar locations (created using Google StreetView; Extended
data Video 1-4; see Materials and Methods). We then compared activity of the place-memory
areas with the scene-perception areas across the familiarity conditions. We hypothesized that if
the place-memory areas were preferentially driven by mnemonic tasks as compared with scene
areas, the place-memory network would respond more strongly than the scene-perception
network when viewing videos of familiar versus unfamiliar locations.

As predicted, we found that the place-memory areas were preferentially driven by videos of
familiar compared to unfamiliar locations, and, critically, this difference was significantly greater

than observed in the scene-perception areas (Region x Familiarity interaction – Lateral:
F(1,91)=20.98, p<0.001, t(13)=6.40, p<0.001, *d*=2.26; Ventral: F(1,91)=7.00, p=0.01, t(13)=4.443,
p<0.001, *d*=1.55; Medial: F(1,91)=7.321, p=0.008, t(13)=7.19, *d*=1.05; Fig. 3a,b). The scene-
perception areas were also driven more by familiar compared to unfamiliar videos (OPA: t(13) =
4.38, p = 0.0001, *d* = 1.171; PPA: t(13) = 4.46, p = 0.0001, *d* = 1.192; MPA: t(13) = 7.231, p < 0.0001,
d = 1.932) but to a significantly lower extent than the place-memory areas, as noted above.
Importantly, the familiarity effect was not observed in control regions, the amygdala (Extended
data Fig. 9) or early visual cortex (Extended data Fig. 10), arguing against a purely attention-
related account of this effect. However, consistent with its role in recognition memory, the
hippocampus activated more when participants saw familiar compared to unfamiliar locations
(F(1,39)=8.14, p=0.0069, t(13)=2.54, p=0.004, *d*=0.75; Extended data Fig. 11). These data suggest
that the place-memory areas are broadly driven by memory-related task demands, including
both recall and recognition memory tasks, to greater extent than the scene-perception areas.

Fig. 3. The place-memory areas respond preferentially to familiar stimuli and form a distinct functional network
during scene-understanding. a,b. The place-memory areas preferentially activate to familiar stimuli. a. Experiment
3. Participants viewed viewing panning movies of personally familiar places versus unfamiliar places, tailored to each
participant using Google StreetView (see Extended data Videos 1-4). The cortical surface depicts the contrast of BOLD
activity for a single participant, thresholded at vertex-wise p < 0.001. Only significant vertices within the scene
perception (white) and place memory (burgundy) areas are shown. b. Average t-statistic of vertices in the scene-
perception (open bars) and place-memory areas (filled bars) when viewing videos of personally familiar places
compared to unfamiliar places. On each cortical surface, the place-memory areas showed an enhanced response to
familiar stimuli compared to the scene-perception areas (all ts > 2.6, ps < 0.01). Connected points depict individual
participants. The hippocampus also showed a preferential response to familiar compared to unfamiliar place movies
(Extended data Fig. 9a). The amygdala (Extended data Fig. 9b) and early visual cortex (Extended data Fig. 10) did not
show a preferential response to familiar place movies, arguing against a purely attentional account of this effect.
Abbreviations: OPA – Occipital place area, LPMA – Lateral place memory area, PPA – Parahippocampal place area,
VPMA – Ventral place memory area, MPA – Medial place area, MPMA – Medial place memory area.

Place-memory and scene-perception areas form different functional networks
While we observed a strong dissociation between the scene-perception and place-memory areas
under controlled experimental conditions, real-world behavior requires a dynamic interaction
between perceptual and memory processes. How do the scene-perception and place-memory
areas interact under naturalistic conditions? In Experiment 3 we investigated this question, and
we hypothesized that, if place-memory areas and scene-perception areas were functionally
distinct, these groups of areas would form separable functional networks during a naturalistic
scene viewing task. Further we hypothesized that, given their functional role in place memory
tasks, place-memory areas would affiliate more with memory structures in the brain than the
scene-perception areas.

Participants watched an 11-minute video that featured a concatenated series of architectural
tours, college admissions videos (including Dartmouth College), and real estate advertisements
(see Methods). This video was designed to concurrently engage perceptual and mnemonic
representations of complex real-world environments by engaging short-term memory built up
over the course of the video, and long-term memory processes engaged during video segments
that included familiar locations (i.e. Dartmouth College). We analyzed the co-fluctuation of
activity among the scene-perception and place-memory areas during the movie by calculating
the correlation between the activity timeseries for each region pair, whilst partialling out the
activation of all other regions (Fig. 4a). Accounting for the activation of all other regions allowed
us to assess the co-fluctuation of these areas above-and-beyond their shared response to the
video stimulus, which causes all regions to exhibit overwhelming positively correlated activity.

This analysis revealed that the place-memory areas constitute a distinct functional network that
is dissociated from the scene-perception network: the correlation among the scene-perception
and place-memory areas (i.e., within network) was significantly greater than the correlation
between these areas (i.e., between network) (Main effect Network: $F_{(60,2)}=47.99$, $p<0.0001$;
Perception:Perception v Perception:Memory, $t_{(12)}=7.31$, $p<0.0001$, *d*=4.63; Memory:Memory v
Perception:Memory $t_{(12)}=5.31$, $p=0.0002$, *d*=3.794; Memory:Memory v Perception:Perception
$t_{(12)}=1.18$, $p=0.26$, *d*=0.48; Fig. 4b). The difference between within- versus between-network
correlations remained significant even after removing the two lowest pairwise correlation values
(MPA with the ventral place memory area and PPA with the lateral place memory area;
Perception:Perception v Perception:Memory, $t_{(12)} = 4.88$, $p = 0.0003$, d = 3.11; Memory:Memory
v Perception:Memory, SPN: $t_{(12)} = 3.27$, $p = 0.001$, d = 2.39), which confirmed that the observed
network-differentiation was not driven by outlying pairwise associations.

Next, we assessed whether the scene-perception and place-memory areas associated more with
the visual and spatio-mnemonic systems, respectively, by assessinng their co-fluctuations with
early visual cortex and the hippocampus. Here, we observed a double dissociation (Main effect

of Network: F(3,84)=55.5, p<0.001), whereby the scene-perception areas were more correlated
to early visual cortex compared to the place-memory areas (t(12)=6.05, p<0.001, *d*=4.26), while
the place-memory areas were more strongly correlated with the hippocampus (t(12)=10.64,
p<0.001, *d*=6.32) (Fig. 4c). Together, these data demonstrate that the place-memory areas
constitute a distinct functional network, which affiliates with the hippocampus during naturalistic
scene understanding.

Fig 4. The place-memory areas constitute a distinct function network and associate closely with the hippocampus. a.
Experiment 2. To assess whether the place-memory areas and scene-perception areas form distinct functional
networks, participants watched an 11-minute video designed to elicit naturalistic scene understanding, comprised of
several college admissions videos, real-estate listings, and architectural tours. For each participant (n=13), the
average timeseries from the scene-perception areas (pink, parahippocampal place area [PPA], occipital place area
[OPA], and medial place area [MPA], the place-memory areas (burgundy, medial, ventral, and lateral place memory
areas [MPMA, LPMA, VPMA]), and the pairwise partial correlation was calculated. The correlation matrix depicts the
average partial correlation of each area from all participants (ordered by Ward similarity). b. The average pairwise
partial correlation of within-network activity (Scene-perception network x Scene-perception network [SPN x SPN] and
place-memory network x place memory network [PMN x PMN]) was significantly higher than the correlation of
between network activity (SPN x PMN) (F_{(2,60)}=50.915, p<0.001). c. The scene-perception and place-memory areas
differentially associate with the brain's visual and memory systems (F_{(3,84)}=55.5, p<0.001). The scene-perception areas
were more correlated with early visual cortex (t_{12}=6.05, p<0.001), while the place-memory areas were more
correlated with the hippocampus (t_{12}=10.64, p<0.001), which is further evidence for their roles in perception and
*memory, respectively. In all plots, n.s., p > 0.05; *p < 0.05; **p < 0.01; ***p < 0.001*

Place-memory areas, not scene-perception areas, activate during visual mental imagery
The observed distinction between the functional networks supporting scene perception and
place memory in posterior cerebral cortex is inconsistent with a classic theory in cognitive
neuroscience: that perception and recall (i.e. mental imagery) of high-level visual stimuli engage
the same neural circuitry[19,29–31,44,47]. In Experiment 4, we explored this discrepancy by
characterizing the relative roles of the scene-perception and place-memory areas in mental

imagery and perception. Specifically, we compared activity when participants performed explicit
place-memory recall in a manner identical to Experiment 1 (i.e. mental imagery) to activity
evoked by viewing panning videos of unfamiliar places (Fig. 5a).

Consistent with our results in Experiment 1, we found that mental imagery and perception
differentially engage the place-memory and scene-perception areas, respectively (Network x
Task interaction – Lateral: $F_{(1,91)}=237.37$; $p<0.001$; Ventral: $F_{(1,91)}=78.19$; $p<0.001$; Medial:
$F_{(1,91)}=28.96$; $p<0.001$). Specifically, we observed a double dissociation on all cortical surfaces.
Compared to the place-memory areas, the scene-perception areas showed greater activation
during perception than mental imagery (Lateral: $t(13)=7.33$; $p<0.001$, $d=2.80$; Ventral: $t(13)=6.21$,
$p<0.001$, $d=1.80$; Medial: $t(13)=4.15$, $p<0.001$, $d=0.97$; Fig. 5b,c). In contrast, compared to the
scene-perception areas, the place-memory areas were more active during mental imagery
(Lateral: $t(13)=8.35$, $p<0.001$, $d=3.33$; Ventral: $t(13)=6.63$, $p<0.001$, $d=2.44$; Medial: $t(13)=5.44$,
$p<0.001$, $d=1.25$; Fig. 5b, c). Remarkably, PPA, a region that prior studies found to be active during
mental imagery[29], showed below-baseline activation during mental imagery ($t(13)=-3.132$,
$p=0.008$), as did OPA ($t(13)=-2.407$, $p=0.03$). Within the scene-perception network, only MPA
showed above-baseline activation during both mental imagery ($t(13)=2.94$, $p=0.01$) and
perception ($t(13)=4.89$, $p=0.003$). Early visual cortex (Extended data Fig. 11), hippocampus
(Extended data Fig. 12a), and amygdala (Extended data Fig. 12b) all showed greater activation
during perception compared to imagery. These findings contradict the classic understanding that
mental imagery recruits the same neural substrates as perception[19,29–31,44,47], and suggest that
the place-memory network, not the scene-perception network, supports explicit visual recall (i.e.
mental imagery) of places.

Fig. 5. Perception and explicit memory recall (i.e. mental imagery) differentially engage the scene-perception and place-memory areas. a. In Experiment 4, participants (n = 14) saw panning movies of unfamiliar places (perception trials) or performed explicit memory recall (i.e. mental imagery) of personally familiar places (mental imagery trials). b. BOLD activation during mental imagery of places compared to baseline for a representative subject. Below baseline activation within the perceptual areas PPA (ventral) and OPA (lateral) are highlighted. Only vertices within the scene-perception network (SPN; white) and place-memory network (PMN; burgundy) are shown. c. The scene-perception areas and place-memory areas are differentially engaged during scene perception and mental imagery. Activation versus baseline of the scene-perception (open bars) and place-memory areas (filled bars) during perception of places (pink) or mental imagery of places (red). A linear mixed effects model analysis revealed that on each cortical surface, there was a significant dissociation in activation during perception and mental imagery, where the scene perception

areas were significantly more active during perception, while the place memory areas were significantly more active
during mental imagery of places (ROI x Task interaction – Lateral: $F_{(1,91)}$=237.37; p<0.001; Ventral: $F_{(1,91)}$=78.19;
p<0.001; Medial: $F_{(1,91)}$=28.96; p<0.001). Early visual cortex (Extended data Fig. 11) and amygdala (Extended data
Fig. 12) also showed below baseline responses during mental imagery; hippocampus responded more to perception
compared to imagery, but positively in both conditions (Extended data Fig. 12). In all plots, n.s., p > 0.05; *p < 0.05;
**p < 0.01; ***p < 0.001

Discussion

To summarize, we found that scene perception and place memory are subserved by two distinct,
but overlapping, functional networks in posterior cerebral cortex. The three regions of the place-
memory network fall immediately anterior to regions of the scene-perception network on all
three cortical surfaces – PPA (ventral), MPA (medial), and OPA (lateral). Despite their topographic
association with the scene-perception areas, the place-memory areas were functionally
dissociable on the basis of their relative preference for mnemonic tasks: the place memory areas
responded more when participants viewed videos of familiar compared to unfamiliar places. The
place-memory areas also formed a dissociable functional network during extended naturalistic
scene understanding, and this network was more strongly correlated with the hippocampus
compared with primary visual cortex. Taken together, our data suggest that the place-memory
areas might be a key interface between the scene perception and spatial memory systems in the
human brain.

Multiple recent studies have suggested that the anterior aspects of ventral temporal cortex,
including PPA, harbor relatively more abstract[48] and context-related[35,49] information than
posterior aspects, consistent with their functional connectivity with memory structures[1,32–34,50].
Our data situate these prior findings within a broader topographical organization of posterior
cerebral cortex: three place-memory selective areas lie immediately anterior to the three scene-
selective visual areas on all three cortical surfaces. These findings reveal a new mechanistic step
between the regions of the brain that support spatial memory and those that support visual
scene analysis, closing a critical gap in our understanding of how integrates perceptual and
memory representations.

Topography of scene-perception and place-memory areas
The topographical relationship between the scene-perception and place-memory areas was
strikingly consistent across individuals: on all cortical surfaces, place-memory selective activity
was located immediately anterior to scene-perception activity. However, intriguingly, the degree
of overlap varied systematically by cortical surface. Specifically, on the ventral surface, the scene-
selective area PPA was activated by scene perception and located immediately posterior to an
area that was activated by both perception and memory (the ventral place memory area, VPMA).

The underlying organizational principles of ventral temporal cortex have been hotly debated.
Along the medial-lateral axis, multiple organizational gradients have been identified, including
eccentricity[24,26,40], animacy[51–54], and real-world size[55]. On the other hand, the anterior-posterior
axis has been conceptualized as a gradient from concrete/perceptual representations to
abstract/conceptual representations[40,46,56–61]. How do our results relate to these gradients?
Theories of medial-lateral organization can account for the presence of scene-preferring visually
responsive regions in medial ventral temporal cortex, because scene-selective areas occur in
peripherally-biased ventral temporal cortex and prefer large objects. However, theories of
anterior-posterior organization do not explain how visual scene representations in PPA are
transformed into allocentric spatial representations in anterior medial temporal lobe[11,12,16,62].
The present results address this knowledge gap, suggesting that a distinct area might subserve
this process: the place-memory responsive region is well-positioned to integrate visual
information represented in PPA with the allocentric spatial representations thought to reside in
hippocampus and entorhinal cortex[11,62]. Thus, by introducing areas that may interface between
visual cortex and memory structures, the present work fills this important gap in our
understanding of how the brain may implement visually guided behaviors.

In contrast to the organization of the ventral surface, the scene-selective visual area on the
medial surface, MPA, was activated by both scene perception and place memory and was
contained within a larger place-memory responsive area (the medial place memory area, MPMA)
that extended anteriorly into medial parietal cortex and retrosplenial cortex proper (Brodmann's
area 29/30). Medial parietal cortex has traditionally been thought of as a homogenous area
engaged during episodic memory processes[33,63–65]. However, many studies have focused on
understanding the presence of a unique scene-preferring visual area in this region[1,5,34,66], and we
and others have demonstrated that specific subdivisions of medial parietal cortex show category-
preferences during memory recall, including the place-memory responsive area described
here[33,63,64]. But few studies have produced an integrative account of the medial parietal cortex's
mnemonic and perceptual functions because they are often investigated separately[50]. Thus, the
present work unifies prior work by describing the topographical and functional distinction
between MPA and MPMA: MPA may be best described as a visually responsive subregion of a
larger, memory responsive area. This raises the question, what function could this organization
subserve? Previous studies have found that medial parietal cortex contains signals that reflect
the local spatial environment from an egocentric perspective, including portions of the visual
field outside of the current field-of-view[67]. Separately, studies in humans and animal models have
shown that retrosplenial cortex proper (Brodmann area 29/30) represents the current heading
direction[68–72]. Taking these results together, we suggest that that MPMA, which overlaps with
retrosplenial cortex, might represent the local spatial environment in an egocentric reference
frame, with MPA reflecting a visual signal underlying this computation.

On the lateral surface, scene-selective OPA was activated by perception and located immediately
posterior to an area that was activated only during memory (the lateral place memory area,
LPMA). Intersecting vertices between the two regions showed joint responsivity to both
perception and memory. Among all of the surfaces, the place memory area on the lateral surface
is perhaps most topographically striking and theoretically surprising. Unlike for MPA and PPA, the
relationship of OPA to memory-related processes has not been widely discussed. One study
found that panoramic scene content associated with a visually-presented, discrete field of view
could be decoded on the lateral surface[67], suggesting a role for OPA beyond perceptual
processing. However, OPA is more broadly characterized as involved in visual analysis: it has been
shown to code three-dimensional information about spaces[73] as well as navigational
affordances[74]. Using resting state fMRI analyses, two studies have shown an area anterior to OPA
where the BOLD response is more strongly correlated to anterior PPA compared to posterior PPA,
leading to speculation that this area (sometimes referred to as caudal IPL) might be involved in
scene memory, but this had not been experimentally tested [32,34]. Thus, our finding of a
functionally distinct place-memory area on the lateral surface significantly advances our
understanding of scene-perception and place-memory processes on the lateral surface of the
human brain. Future work should focus on identifying the specific contribution of LPMA to
memory-guided visual behavior.

Content-specificity in the default mode network
While we focused on the intersection between the place-memory and scene-perception areas, it
is interesting that the place memory network resides within the caudal inferior parietal lobule,
medial temporal lobe, and posterior parietal cortex, which are also hubs of the default mode
network on the lateral, ventral, and medial surfaces, respectively[75]. Recent conceptualizations of
human cortical organization consider these hubs as the apex of a macroscale gradient, spanning
from unimodal (i.e. sensory or motor) areas to transmodal, high-level areas[76,77]. These
transmodal areas are thought to facilitate cognitive processes, such as autobiographical memory,
in a content-agnostic fashion[18,46,75,76,78,79]. Intriguingly, however, our data show that place-
memory selective subregions exist within each of these default mode network hubs, broadly
mirroring the category-selective subdivisions of the visual system[33,40,80]. Based on our
observations here and prior work[18,33,63,64,80], we hypothesize that content-specificity may be a
defining feature of cortical organization at each level of processing, from visual to transmodal
cortex. Neuropsychological patients support this assertion: damage to specific areas within the
caudal inferior parietal lobule, posterior parietal cortex, or medial temporal lobe cause severe
navigational impairments, while leaving other memory domains, such as face memory, intact[81,82].
This content-specific organization likely persists in prefrontal areas[83–85], although traditional
functional imaging analysis approaches might fail to detect tightly interdigitated features due to
topographic heterogeneity across individuals[83–85]. Revealing the relationship between
transmodal cortex and content-specific networks within individual participants will be

paramount to understanding the neuroanatomical substrates underlying complex cognitive
processes, and future studies directly comparing these systems could help to elucidate this issue.

A bridge between perceptual and memory structures
In addition to residing at the intersection of transmodal and perceptual cortex, the place-memory
areas lie at the intersection between perceptual areas and memory structures in the brain,
including the hippocampus and enthorinal cortex. Although traditionally the hippocampus has
been associated with episodic memory[45,86], recently the hippocampus has been implicated in
both perceptual and mnemonic tasks, including scene construction[87–89]. Consistent with this
view, we found that the hippocampus, like the place-memory areas, responded preferentially to
perceiving familiar stimuli compared to unfamiliar stimuli, suggesting that it has a role in visual
recognition-related processes. Interestingly, unlike the place-memory areas, when we directly
compared perception and recall in Experiment 4, the hippocampus responded more during
perception compared to recall, which is qualitatively more similar to the scene-perception areas
than the place-memory areas. Together, these experiments suggest that the hippocampus is
involved in both perceptual and memory processes, but we cannot delineate what the
hippocampus specifically contributes in each case. For example, during perception of unfamiliar
stimuli, the hippocampus may be encoding relevant perceptual features, or comparing the
present visual stimuli versus other internal representations, or both. Future work is necessary to
disentangle the role of the hippocampus in scene perception and situate it in the context of the
place-memory and scene-perception networks.
461
Different neural substrates for perception and recall (i.e. mental imagery)
Based on previous fMRI studies, it has been widely assumed that perception and recall (i.e.
mental imagery) of high-level stimuli (such as scenes) recruit the same neural substrates,
including category-selective areas in ventral temporal cortex[31,90]. For example, prior studies
found that both perceiving and recalling scenes evoked activity in overlapping, scene-selective
cortical areas and concluded that perception and memory shared neural substrates[22,29,47].
However, in contrast, neuropsychological studies have suggested that mental imagery and visual
recognition may have dissociable neural substrates[37,39,91], given observations of patients with
preserved visual recognition but impoverished mental imagery and *vice versa*. Our findings
reconcile this discrepancy, by showing that distinct, but neighboring regions underlie perception
and recall (i.e. mental imagery) of places. These neighboring regions, which are evident at the
individual subject level, likely account for recent group-level findings of an anterior shift in
category-level decoding of scenes during recall vs. perception[36]. In sum, we observe minimal
overlap between the neural substrates of scene perception and place memory: the (anterior)
place-memory network, but not the (posterior) scene-perception network, preferentially
activates during recall (i.e. mental imagery).

So, why might previous fMRI studies have reported that the substrates of perception and imagery
in ventral temporal cortex are shared? First, the initial fMRI investigation into mental imagery
and perception of high-level visual stimuli[29] used familiar exemplars for perception, which could
recruit the VPMA (similar to Experiment 2 here). Second, this initial study, which used a
combination of individual subject mapping and ROI analyses, reported that only a portion of PPA
was active during imagery[29], but later studies did not investigate this distinction. So, the mental
imagery-related activation and decoding observed in later studies (e.g. [30]) may also have arisen
from nearby memory areas. Consistent with this hypothesis, we note that if we analyze PPA and
VPMA combined as a single area, we find positive activation during mental imagery in the
majority of our participants (Extended data Fig. 13). Third, typical whole-brain group-analyses
could obscure the topographical relationship between perception- and memory-related activity
by averaging across individuals[19,31,44]. As such, our individualized analysis approach is uniquely
suited to detect this fine-grained distinction. Importantly, as in some previous studies of mental
imagery[47], participants in our study recalled *familiar* locations in the recall condition but viewed
unfamiliar locations in the perception condition. Future work should consider mapping the
evolution of place-memory area response longitudinally as stimulus familiarity develops to better
understand how familiarity shapes the activity of the place-memory areas.

Are scenes special? Distinct perceptual-mnemonic topography for scenes, not faces
Finally, the distinct topography of the scene-perception and place-memory networks in posterior
cerebral cortex was unique to the spatial domain – we found no separable networks for face-
perception and memory (although, consistent with prior work[33], we did observe a people-
memory preferring area in medial parietal cortex in most participants). These findings challenge
the hypothesis that a concrete/abstract posterior-anterior gradient may broadly characterize
high-level visual cortex[36,58,61]. Why might a 'place-memory network', anterior to the scene-
perception network, exist, while an analogous 'face-memory network' does not? We hypothesize
that this distinction might relate to the 'goal' of scene processing: integrating the current field-
of-view with the larger spatial context[35,67]. In contrast, the 'goal' of visual face processing is face
recognition, which requires nuanced discrimination of a discrete stimulus, but not broader
visuospatial context[6,92,93]. Neuropsychological patients support this differentiation. For example,
lesions of the scene-perception network can cause impairments of spatial processing while
leaving scene recognition and certain navigational abilities intact[82]. On the other hand, lesions of
critical face-perception areas including the occipital face area or FFA tend produce significant and
selective impairments in conscious face recognition *and* imagery[94,95]. In sum, we suggest that
visual scene processing regions may have unique processing requirements relative to other visual
categories (e.g., object or face processing), and generalizing computational principles from other
categories may be limited. To test this hypothesis, future studies should characterize the
topography of mnemonic and perceptual signals in high-level visual cortex using multiple
categories beyond those used here (e.g. objects, bodies, words).

Conclusion

To conclude, we have described the place memory network: a set of brain areas in posterior

cerebral cortex that support recall of familiar locations. The discovery of these functionally

distinct areas, residing between the brain's perception and memory systems, provides a new

framework for considering the neural substrates of memory-guided visual behaviors.

MATERIALS AND METHODS

Participants

Fourteen adults (9 females; age=25.7±3.6 STD years old) participated in the Experiments 1, 2, and

4. Of these, 13 participants took part in Experiment 3 (one participant requested to abort the

scan; 8 females, age=25.9±3.6 STD years old). In addition, a subset of the original participants

(N=6; 2 females, age=26.3±4.4 STD years old) completed the multi-echo replication of Experiment

1. Participants had normal or correct-to-normal vision, were not colorblind, and were free from

neurological or psychiatric conditions. Written consent was obtained from all participants in

accordance with the Declaration of Helsinki and with a protocol approved by the Dartmouth

College Institutional Review Board (Protocol #31288). We confirmed that participants were able

to perform mental imagery by assessing performance Vividness of Visual Imagery Questionnaire

[96]. All participants scored above 9 with eyes open, indicating satisfactory performance

(mean=15.2, range=9.75-19.25).

Procedure

All participants took part in Experiments 1-4. In Experiment 1, we compared i) the topography of

activation during a scene/face perceptual localizer with ii) the topography of place/people

memory-recall related activation collected in separate fMRI runs. In Experiment 2, we

characterized the differential functional responses of the place-memory and scene-perception

areas during a recognition memory task, by comparing activation when participants viewed

panning movies of personally familiar as compared with unfamiliar places. In Experiment 3, we

tested whether the scene-perception and place-memory areas dissociated into distinct

functional networks under naturalistic conditions by evaluating the correlated activity of these

areas during a movie-watching task. Finally, in Experiment 4, we evaluated the relative

involvement of the scene-perception and place-memory areas in perception and visual recall (i.e.

mental imagery) by comparing their activation when participants performed mental imagery of

personally familiar places as compared with viewing panning videos of unfamiliar places.

Stimuli – Experiments 1-4

Personally familiar place- and face-memory stimulus lists
For Experiments 1 and 4, we asked participants to generate a list of 36 familiar people and places
(72 stimuli total). These stimuli were generated prior to scanning based on the following
instructions:

For your scan, you will be asked to visualize people and places that are personally familiar to you.
So, we need you to provide these lists for us. For personally familiar people, please choose people
that you know personally (no celebrities) that you can visualize in great detail. You do not need
to be in contact with these people now – just as long as you knew them personally and remember
what they look like. So, you could choose your childhood friend even if you are no longer in touch
with this person. Likewise, for personally familiar places, please list places that you have been to
and can imagine in great detail. You should choose places that are personally relevant to you, so
you should avoid choosing places that you have only been one time, and you should not choose
famous places where you have never been. You can choose places that span your whole life, so
you could do your current kitchen as well as the kitchen from your childhood home.

Familiar panning movie stimuli
For Experiment 2, we generated panning movies depicting places that were personally familiar
to our participants. To do this, participants used the GoogleMaps plug-in iStreetView
(istreetview.com) to locate and download 24 photospheres of personally familiar places based
on the following instructions:

In another part of your scan, you will see videos of places that will be familiar or unfamiliar to
you. You will use iStreetView to locate places with which you are familiar. iStreetView has access
to all photospheres that Google has access to, as well as all of the photospheres from Google
Street View. So, you should do your best to find photospheres taken from perspectives where you
have actually stood in your life. Like before, please find places that are personally familiar to you
(such as in front of your childhood school), rather than famous places (like the Eiffel tower), even
if you have been to that famous place in your life.

Participants were also asked to specify whether a particular viewpoint would be most
recognizable to them so that this field-of-view could be included in the panning movie. Once the
photospheres were chosen, panning movies were generated by concatenating a series of 90°
fields-of-view images taken from the photospheres that panned 120° to the right (in steps of
0.4°), then back to the left. The videos always began and ended at the same point of view, and
the duration of the video was 10 seconds. Extended data Videos 1-4 show examples of these
videos.

Naturalistic movie stimulus

In Experiment 3, participants watched an 11-minute movie depicting various, real-world scenes.
The movie consisted of an architectural tour, three college admissions videos, one tourism
informational video, and one real-estate listing downloaded from YouTube. Specifically, the
segments were: 1) an architectural tour of Moosilauke Ravine Lodge in Warren, NH (0:00-2:00),
2) an admission video of Princeton University in Princeton, NJ (2:00-3:00), 3) an admission video
of Dartmouth College in Hanover, NH (3:00-3:51), 4) an admission video of Cornell University in
Ithaca, NY (3:52-6:40), 5) a tourism informational video of the University of Oxford in Oxford,
United Kingdom (6:40-8:53), 6) and a real-estate listing of a lake front property in Enfield, NH
(8:53-11:00). Each segment included footage of people and places, ground-level and aerial
footage, and indoor and outdoor footage. Videos were chosen to match closely for content,
footage type, and video quality, while varying in the degree to which participants would be
familiar with the places. No audio was included in the video. Participants were not asked whether
places were familiar to them, although all participants were Dartmouth students/staff and
therefore familiar with the Dartmouth Campus. The full video can be found here:
https://bit.ly/2Arl3I0.

Questionnaire - Vividness ratings

To ensure that participants could reliably visualize the familiar stimuli used in Experiments 1 and
4, and to confirm that there were no differences in visualization between stimulus categories, we
had participants rate their ability to richly visualize each stimulus. Specifically, after the scanning
session we gave the participants the following instructions:

Please rate each stimulus on how well you are able to visualize each stimulus from 1-5 (1 – I
cannot visualize this person/place, 5 – as though I am seeing the person/place). Please rate the
stimuli base upon how well you can visualize the stimulus, and not try to remember how well you
visualized the stimulus during the scan.

Procedure - Experiment 1: Perceptual category localizer

Face- and scene-selective perceptual areas were localized using a block-design functional
localizer. Participants were shown images of faces, scenes, and objects (image presentation
duration: 500 ms; ISI: 500 ms; block duration: 24 s; number of blocks per condition: 6). Blocks
were presented continuously with no baseline periods between blocks. Participants were
instructed to passively view the images and performed no task during the perceptual localizer
experiment. Two runs of the localizer were performed.

Procedure - Experiment 1: Memory activation localizer

People and place selective memory activation was localized using a procedure adapted
from[33]. On each trial, participants were shown the written name of one of 36 personally

familiar people and 36 places that they provided to us prior to the MRI scan (see *Personally*
familiar place- and face-memory stimulus lists above). Each stimulus was presented one time
during the memory localizer task. Participants were instructed to visualize this stimulus from
memory as vividly as possible for the duration of the trial (10 s) following the instructions
detailed below. Trials were separated by a variable inter-trial interval (4-8 s). The task was
performed over 4 runs, with 9 people and place stimuli presented during each run in a
pseudo-randomized order. No more than 3 instances of a stimulus category (i.e. person or
place) could appear consecutively

Localizer task instructions were as follows:

In this task, you will see the name of a person or place from the list you provided us. When the
name appears, I want you to perform memory recall of that stimulus. The recall should be as vivid
as possible, and you should try to do this the whole time that the name is on the screen.

If the name you see is a person, I want you to imagine that person's face in as much detail as
possible. You should picture their head in front of you as the person turns their head from left to
right and up and down, so that you can see all of the sides of their face. They can make different
expressions, as well.

The most important thing is that you are picturing their face; when some people perform memory
recall, they will imagine interacting with that person or touching and hugging them. We do not
want you to do this, just imagine the face. Also, do not picture their bodies, arms, or legs. Try to
focus just on their face as it moves in your mind's eye.

If the name you see is a place, I want you to imagine that you are in the place. You should direct
your attention around the place, looking to the left and right of you, and looking up and down,
and behind you. We want you to have the experience of being in this place as much as possible.

When you are imagining places, you should imagine yourself standing in one single spot. Don't
walk around the place or imagine yourself from above. Also, we want it to be as though you are
in the place, rather than thinking about the objects in the place. So, as you "look" around, don't
think about the individual objects or list them in your mind. Instead, we want you to image that
the place has materialized all around you, and all of the things that would be within your field of
view are "visible" at the same time. For example, if you are imagining your kitchen, do not imagine
the chairs, table, and windows sequentially. Instead, try to imagine all of these features at once,
and that you're looking around and experiencing them as you would if you were in the room.

Between each trial, participants were instructed to clear their mind, relax, and wait for the next
trial to begin.

Procedure - Experiment 2: Familiar/unfamiliar places experiment

The goal of Experiment 2 was to determine whether the place memory areas respond
preferentially to familiar stimuli (i.e., stimuli that could be recognized) presented visually. To this
end, we compared brain activation when participants saw panning videos of personally familiar
places with activation when viewing panning videos of unfamiliar places. Videos were taken from
Google Maps as described above (see section Familiar panning movie stimuli). To control for low-
level variation in stimuli, unfamiliar videos were randomly sampled from other participants'
familiar videos. Videos of Dartmouth College and Hanover, NH were excluded from the possible
unfamiliar places.

The experiment was performed across 6 runs, with each run featuring 8 familiar and 8 unfamiliar
place videos. There were 24 videos in each condition, and each video was seen twice. Before the
experiment, participants were told that they would be seeing videos of familiar and unfamiliar
places panning from left to right, then back again and that they should imagine that they were
standing in that place, turning their head like the video. The instructions provided to the
participants are reproduced below.

You will see panning movies of places that you are familiar with, as well as places that will not be
familiar to you. Regardless of which you see, we want you to imagine that you are in that location
and the video is your perspective as you turn your head from left to right, and back again. You
should not actually turn your head.

Be sure that for all scenes you are pretending you are in that location, whether or not they are
familiar to you. Of course, if it is a familiar location, you might be able to predict what is coming
into view. That is fine, just do your best to imagine you're in that place for both conditions.

Each trial lasted 10 seconds. Trials were separated by a variable inter-trial interval (4-8 s). All
videos were repeated two times in the experiment: once in the first three runs, and once in the
last three runs.

Procedure – Experiment 3: Naturalistic movie watching experiment

The goal of Experiment 3 was to characterize the network properties of the scene-perception and
place-memory areas during naturalistic conditions. 13 participants took part in a single run of an
11-minute movie watching experiment (see *Naturalistic movie stimulus* above for stimulus
details). We chose a natural movie watching task, rather than a simple resting-state scan, to avoid
the possibility that mind-wandering or mental imagery might cause a decoupling of the
perception and memory networks artificially[78,97].

Procedure – Experiment 4: Mental imagery versus perception experiment

Experiment 4 explicitly tested whether the place memory network was activated to a greater
extent than the scene perception network during mental imagery. Each imaging run featured two
types of trials, perception and mental imagery trials. On perception trials, participants saw a
panning movie of an unfamiliar place, and were instructed to imagine that they were in that place
turning their head (as in Experiment 3). On mental imagery trials, participants saw the name of a
personally familiar place, and performed mental imagery of that place following the instructions
given in Experiment 1 (instructed to imagine that they were in that place turning their head). This
task was presented over 6 runs. Each run featured 8 familiar place word stimuli for mental
imagery and 8 unfamiliar place videos. As in Experiment 3, 24 unique stimuli were used in each
condition (24 place word stimuli taken from the list generated in Experiment 1, and 24 panning
videos of unfamiliar places). Trials lasted 10 s and were separated by a variable inter-trial interval
(4-8 s). All stimuli were repeated two times in the experiment, once in the first three runs, and
once in the last three runs. Familiar stimuli for mental imagery were randomly sampled from
each participant's stimulus list used during Experiment 1. Unfamiliar places videos for the
perception condition were not repeated from Experiment 2.

MRI acquisition and preprocessing

699
All data was collected on a Siemens Prisma 3T MRI scanner equipped with a 32-channel head coil
at Dartmouth College. Images were transformed from dicom to nifti files using dcm2niix [98], which
applies slice time correction by default.

T1 image
For registration purposes, a high-resolution T1-weighted magnetization-prepared rapid
acquisition gradient echo (MPRAGE) imaging sequence was acquired (TR=2300 ms, TE=2.32 ms,
inversion time=933 ms, Flip angle=8°, FOV=256 x 256 mm, slices=255, voxel size=1 x 1 x 1 mm).
T1 images segmented and surfaces were generated using Freesurfer[99–101].

Single-echo fMRI

Acquisition
In Experiments 1-4, single-echo T2*-weighted echo-planar images covering the temporal,
parietal, and frontal cortices were acquired using the following parameters: TR=2000 ms, TE=32
ms, GRAPPA=2, Flip angle=75°, FOV=240 x 240 mm, Matrix size=80 x 80, slices=34, voxel size=3 x
3 x 3 mm. To minimize dropout caused by the ear canals, slices were oriented parallel to temporal
lobe[102]. The initial two frames were discarded by the scanner to achieve steady state.

Preprocessing

Task fMRI (Experiments 1, 3, and 4)

Task fMRI data was preprocessed using AFNI[103]. In addition to the frames discarded by the fMRI
scanner during acquisition, the initial two frames were discarded to allow T1 signal to achieve
steady state. Signal outliers were attenuated (3dDespike). Motion correction was applied, and
parameters were stored for use as nuisance regressors (3dVolreg). Data were then iteratively
smoothed to achieve a uniform smoothness of 5mm FWHM (3dBlurToFWHM).

Naturalistic movie watching (Experiment 2)
Naturalistic movie watching data were preprocessed using a combination of AFNI and FSL tools.
Signal outliers were attenuated (3dDespike). Motion correction was applied. Data were then
iteratively smoothed to achieve a uniform smoothness of 5mm FWHM (3dBlurToFWHM).

For denoising, independent component analysis (ICA) was applied to decompose the data into
signals and sources using FSL's melodic[104–106]. These were classified as signal or noise by one
investigator (AS) using the approach described in Griffanti et al. (2014)[107]. Components classified
as noise were then regressed out of the data (fsl_regfilt). Motion from volume registration was
not included in the regression, as motion is generally well captured by the ICA decomposition[107].
Time series were then transferred to the high-density SUMA standard mesh (std.141) using
@SUMA_Make_Spec_FS and @Suma_AlignToExperiment.

Multi-echo fMRI

Acquisition
To better characterize activation during mental imagery of personally familiar people, we
acquired a replication dataset of Experiment 1 using a multi-echo T2*-weighted sequence. Multi-
echo imaging afforded the benefit of mitigating dropout with short echo times, while maintaining
a high level of BOLD contrast with long echo times[108–110]. The sequence parameters were:
TR=2000 ms, TEs=[11.00, 25.33, 39.66, 53.99 ms], GRAPPA=3, Flip angle=75, FOV=240 x 240 mm,
Matrix size=80 x 80, slices=40, Multi-band factor=2, voxel size=3 x 3 x 3 mm. As with single-echo
acquisition, the slices were oriented parallel to the temporal lobe. The initial two frames were
discarded by the scanner.

Preprocessing
Multi-echo data preprocessing was implemented based on the multi-echo preprocessing pipeline
from afni_proc.py. Initial preprocessing steps were carried out on each echo separately. Signal
outliers were attenuated (3dDespike). Motion correction parameters were estimated from the
second echo (3dVolreg); these alignment parameters were then applied to all echoes.

Data were then denoised using multi-echo ICA (tedana.py[109–111]). The optimal combination of the
four echoes was calculated, and the echoes were combined to form a single, optimally weighted

time series (T2smap.py). Data were then subjected to PCA, and thermal noise was removed using

the Kundu decision tree, which selectively eliminates components that explain a small amount of

variance and do not have a TE-dependent signal decay across echoes. Subsequently, ICA was

performed to separate the time series into independent spatial components and their associated

signals. These components were classified as signal and noise based on known properties of the

$T2^*$ signal decay of BOLD versus noise. The retained components were then recombined to

construct the optimally combined, denoised time series. Following denoising, images were

blurred with a 5mm gaussian kernel (3dBlurInMask) and normalized to percent signal change.

fMRI analysis

Region of interest definitions (Scene-perception and Place-memory areas, Experiment 1)

To define category selective perceptual areas, the scene perception localizer was modeled by

fitting gamma function of the trial duration with a square wave for each condition (Scenes, Faces,

and Objects) using 3dDeconvolve. Estimated motion parameters were included as additional

regressors of no-interest. To compensate for slow signal drift, 4^{th} order polynomials were

included for single-echo data. Polynomial regressors were not included multiecho data analysis.

Scene and face areas were drawn based on a general linear test comparing the coefficients of the

GLM during scene versus face blocks as well as scene versus face + objects blocks. These contrast

maps were then transferred to the SUMA standard mesh (std.141) using @SUMA_Make_Spec_FS

and @Suma_AlignToExperiment. A vertex-wise significance of $p<0.001$ along with expected

anatomical locations was used to define the regions of interest – in particular, visually responsive

PPA was not permitted to extend posteriorly beyond the fusiform gyrus.

To define category-selective memory areas, the familiar people/places memory data was

modeled by fitting a gamma function of the trial duration for trials of each condition (people and

places) using 3dDeconvolve. Estimated motion parameters were included as additional

regressors of no-interest. To compensate for slow signal drift, 4^{th} order polynomials were

included for single-echo data. Polynomial regressors were not included multiecho data analysis.

Activation maps were then transferred to the suma standard mesh (std.141) using

@SUMA_Make_Spec_FS and @Suma_AlignToExperiment. People and place memory areas were

drawn based on a general linear test comparing coefficients of the GLM for people and place

memory. A vertex-wise significance threshold of $p<0.001$ was used to draw ROIs.

To control for possible signal-to-noise differences introduced by variably sized ROIs, for analysis

of correlated activity during movie watching (Experiment 3) and activation in Experiments 2 and

4, scene-perception and place-memory ROIs were constrained to unique members of the top 300

vertices from the perception and memory areas. These ROIs are referred to as "constrained ROIs"

in subsequent sections.

Importantly, our findings in Experiments 1-4 remained significant when two additional
approaches to defining scene-selective areas were used: 1) the contrast scenes > faces + objects,
and 2) a disk of 300 nodes drawn around the center of each ROI (See Extended Data Figs. 14-15).

Analysis of topography of perception and memory areas (Experiment 1)
The topography of the perception and memory areas was compared in two ways: by identifying
whether there was a significant anterior shift from perception to memory and quantifying the
overlap between category selective perception and memory vertices at the significance threshold
p<0.001.

To quantify the anterior displacement of perception and memory areas, we calculated the
weighted center of mass for the scene-perception/place-memory selective area on each surface,
where the contribution of each vertex to the center of mass was weighted by its selectivity (t-
statistic). The distance between the center of mass for perception and memory in the y-
dimension (posterior-anterior) was then compared using a linear mixed effects model with ROI
(perception/memory) and Hemisphere (lh/rh) as factors separately for each surface. Because
there was no significant effect of hemisphere, the data are presented collapsed across
hemisphere (all ps>0.10). In addition, to determine whether the anterior displacement of
memory compared to perception was specific to scenes, we compared the distance in the y
direction between the PPA and ventral place memory area (VPMA) with the FFA and the face-
memory selectivity are on the ventral surface using a linear mixed effects model with Category
(scene/face), ROI (perception/memory) and Hemisphere (lh/rh) as factors.

To further examine the relationship between perception and memory on each surface, we then
qualitatively compared the overlap between the category-selective perception and memory
areas on each surface.

Group place-memory localizer analysis
Group place-memory activity was defined by comparing beta-values during place- versus people-
memory recall activation for each subject using 3dttest++ in AFNI. Data were thresholded at
vertex-wise q < 0.00015. The peak of the group localizer results were compared with 1) a
probabilistic retinotopic atlas[42] and 2) an anatomical parcellation based on multi-modal data
from the Human Connectome Project[43] aligned to the SUMA standard mesh (std.141).

Anatomical region of interest definitions (Hippocampus and Primary Visual Cortex)
Hippocampus and primary visual cortex (occipital pole) were defined for each participant using
FreeSurfer's automated segmentation (aparc+aseg).

Task fMRI analyses (Experiments 2 & 4)
For Experiments 2 and 4, a gamma function of duration 10s was used to model responses for
trials in each condition (Experiment 2: familiar/unfamiliar places videos; Experiment 4: mental

imagery/perception). These regressors, along with motion parameters and 4th order polynomials
were fit using 3dDeconvolve. Activation maps were then transferred to the SUMA standard mesh
(std.141) using @SUMA_Make_Spec_FS and @SUMA_AlignToExperiment.

For analysis, the average t-statistic (compared to baseline) for each condition (Experiment 3:
familiar versus unfamiliar videos; Experiment 4: mental imagery versus perception tasks) from
the constrained ROIs was calculated. The average t-statistics were compared using a linear mixed
effects model [112] in R [113] for each surface separately (i.e. PPA v VPMA, MPA v MPMA, OPA v LPMA
were separately compared). Trial condition (Experiment 3 – Familiarity: familiar/unfamiliar;
Experiment 4 – Task: perception/imagery) and Hemisphere (lh/rh) were included as factors.
There was no significant effect of hemisphere in any test, and thus data are presented collapsed
across hemisphere (all ps>0.10). Post-hoc tests were implemented using the emmeans package
[114]. In addition, to determine whether a region was significantly active above or below baseline,
the t-static from each ROI was compared versus zero. T-statistics were chosen to aid comparison
across areas, which is typical in these studies[6].

Analysis of correlated activity timeseries during movie watching (Experiment 3)
In Experiment 3, we analyzed the co-fluctuation of activity patterns in each ROI during a
naturalistic movie-watching task. For each participant, we first extracted the average time course
of each constrained ROI, as well as the hippocampus and early visual cortex, which were
anatomically defined based on each participant's FreeSurfer segmentation/parcellation. We then
calculated the correlation of the time series from each region pair while partialing out the time
series from all other region pairs. The correlation matrices were calculated for each hemisphere
separately. The average pairwise correlation within- and between- networks (perception: PPA,
MPA, OPA; memory: VPMA, medial place memory area (MPMA), and lateral place memory area
(LPMA)) were then compared using a linear mixed effects model with Connection (PxP, MxM,
and PxM) and Hemisphere (lh/rh) as factors. The average correlation of each network with
hippocampus and early visual cortex was compared using a linear mixed effects model [112] with
Network (Perception/Memory) and Hemisphere (lh/rh) as factors. Significant model terms were
determined using an analysis of variance implemented in R[113]. There was no significant effect of
hemisphere in any test, and thus data are presented collapsed across hemisphere (all ps>0.10).

DATA AVAILABILITY STATEMENT

851   The data that support the claims of this study are available upon reasonable request to the
852   corresponding author.

CODE AVAILABILITY STATEMENT

854   The code used in this study are available upon reasonable request to the corresponding author.

REFERENCES

856
857

1.   Epstein, R. A. & Baker, C. I. Scene Perception in the Human Brain. *Annu. Rev. Vis. Sci.* **5**,
annurev-vision-091718-014809 (2019).

2.   Epstein, R. & Kanwisher, N. A cortical representation the local visual environment. *Nature*
392, 598–601 (1998).

3.   Dilks, D. D., Julian, J. B., Paunov, A. M. & Kanwisher, N. The occipital place area is causally
and selectively involved in scene perception. *J. Neurosci.* **33**, 1331–1336 (2013).

4.   Marchette, S. A., Vass, L. K., Ryan, J. & Epstein, R. A. Outside looking in: Landmark
generalization in the human navigational system. *J. Neurosci.* **35**, 14896–14908 (2015).

5.   Marchette, S. A., Vass, L. K., Ryan, J. & Epstein, R. A. Anchoring the neural compass: Coding
of local spatial reference frames in human medial parietal lobe. *Nat. Neurosci.* **17**, 1598–
1606 (2014).

6.   Silson, E. H., Chan, A. W. Y., Reynolds, R. C., Kravitz, D. J. & Baker, C. I. A retinotopic basis
for the division of high-level scene processing between lateral and ventral human
occipitotemporal cortex. *J. Neurosci.* **35**, 11921–11935 (2015).

7.   Kornblith, S., Cheng, X., Ohayon, S. & Tsao, D. Y. A network for scene processing in the
macaque temporal lobe. *Neuron* **79**, 766–781 (2013).

8.   Maguire, E. A. *et al.* Knowing where and getting there: A human navigation network.
Science (80-.). **280**, 921–924 (1998).

9.   Epstein, R. A., Higgins, J. S., Jablonski, K. & Feiler, A. M. Visual Scene Processing in Familiar
and Unfamiliar Environments. *J. Neurophysiol.* **97**, 3670–3683 (2007).

10.   O'Keefe, J. & Nadel, L. *The Hippocampus as a Cognitive Map*. (Oxford University Press,
1978).

11.   Moser, E. I., Kropff, E. & Moser, M.-B. Place Cells, Grid Cells, and the Brain's Spatial
Representation System. *Annu. Rev. Neurosci.* **31**, 69–89 (2008).

12.   Doeller, C. F., Barry, C. & Burgess, N. Evidence for grid cells in a human memory network.
Nature **463**, 657–661 (2010).

13.   Ekstrom, A. D. *et al.* Cellular networks underlying human spatial navigation. *Nature* **425**,
184–187 (2003).

14.   Maguire, E. A., Woollett, K. & Spiers, H. J. London taxi drivers and bus drivers: A structural
MRI and neuropsychological analysis. *Hippocampus* **16**, 1091–1101 (2006).

15.   Howard, L. R. *et al.* The hippocampus and entorhinal cortex encode the path and euclidean

distances to goals during navigation. *Curr. Biol.* **24**, 1331–1340 (2014).

16.   Julian, J. B., Keinath, A. T., Frazzetta, G. & Epstein, R. A. Human entorhinal cortex
represents visual space using a boundary-anchored grid. *Nat. Neurosci.* **21**, 191–194
(2018).

17.   Tsitsiklis, M. *et al.* Single-Neuron Representations of Spatial Targets in Humans. *Curr. Biol.*
30, 245-253.e4 (2020).

18.   Woolnough, O. *et al.* Category Selectivity for Face and Scene Recognition in Human Medial
Parietal Cortex. *Curr. Biol.* (2020). doi:10.1016/j.cub.2020.05.018

19.   Pearson, J. The human imagination: the cognitive neuroscience of visual mental imagery.
Nat. Rev. Neurosci. 1–11 (2019). doi:10.1038/s41583-019-0202-9

20.   Wheeler, M. E., Petersen, S. E. & Buckner, R. L. Memory's echo: Vivid remembering
reactivates sensory-specific cortex. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 11125–11129 (2000).

21.   Buchsbaum, B. R., Lemire-Rodger, S., Fang, C. & Abdi, H. The neural basis of vivid memory
is patterned on perception. *J. Cogn. Neurosci.* **24**, 1867–1883 (2012).

22.   Johnson, M. R. & Johnson, M. K. Decoding individual natural scene representations during
perception and imagery. *Front. Hum. Neurosci.* **8**, 59 (2014).

23.   Schultz, H. *et al.* Content tuning in the medial temporal lobe cortex: Voxels that perceive,
retrieve. *eNeuro* **6**, (2019).

24.   Hasson, U., Levy, I., Behrmann, M., Hendler, T. & Malach, R. Eccentricity bias as an
organizing principle for human high-order object areas. *Neuron* **34**, 479–490 (2002).

25.   Hasson, U., Harel, M., Levy, I. & Malach, R. Large-scale mirror-symmetry organization of
human occipito-temporal object areas. *Neuron* **37**, 1027–1041 (2003).

26.   Levy, I., Hasson, U., Avidan, G., Hendler, T. & Malach, R. Center-periphery organization of
human object areas. *Nat. Neurosci.* **4**, 533–539 (2001).

27.   Kamps, F. S., Pincus, J. E., Radwan, S. F., Wahab, S. & Dilks, D. D. Late Development of
Navigationally Relevant Motion Processing in the Occipital Place Area. *Curr. Biol.* (2020).
doi:10.1016/j.cub.2019.12.008

28.   Ino, T. *et al.* Mental navigation in humans is processed in the anterior bank of the parieto-
occipital sulcus. *Neurosci. Lett.* (2002). doi:10.1016/S0304-3940(02)00019-8

29.   O'Craven, K. M. & Kanwisher, N. Mental imagery of faces and places activates
corresponding stimulus-specific brain regions. *J. Cogn. Neurosci.* **12**, 1013–1023 (2000).

30.   Boccia, M. *et al.* The dynamic contribution of the high-level visual cortex to imagery and
perception. *Hum. Brain Mapp.* **40**, 2449–2463 (2019).

31.   Dijkstra, N., Bosch, S. E. & van Gerven, M. A. J. Shared Neural Mechanisms of Visual
923   Perception and Imagery. *Trends in Cognitive Sciences* **23**, 423–434 (2019).

924   32.   Baldassano, C., Esteva, A., Fei-Fei, L. & Beck, D. M. Two Distinct Scene-Processing Networks
925   Connecting Vision and Memory. *eNeuro* **3**, (2016).

926   33.   Silson, E. H., Steel, A., Kidder, A., Gilmore, A. W. & Baker, C. I. Distinct subdivisions of
927   human medial parietal cortex support recollection of people and places. *Elife* **8**, (2019).

928   34.   Silson, E. H., Steel, A. D. & Baker, C. I. Scene-Selectivity and Retinotopy in Medial Parietal
929   Cortex. *Front. Hum. Neurosci.* **10**, 412 (2016).

930   35.   Peer, M., Ron, Y., Monsa, R. & Arzy, S. Processing of different spatial scales in the human
931   brain. *Elife* **8**, (2019).

932   36.   Bainbridge, W. A., Hall, E. H. & Baker, C. I. Distinct Representational Structure and
933   Localization for Visual Encoding and Recall during Visual Imagery. *Cereb. Cortex* **00**, 1–16
934   (2020).

935   37.   Bartolomeo, P. The relationship between visual perception and visual mental imagery: A
936   reappraisal of the neuropsychological evidence. *Cortex* **38**, 357–378 (2002).

937   38.   Bartolomeo, P. *et al.* Multiple-domain dissociation between impaired visual perception
938   and preserved mental imagery in a patient with bilateral extrastriate lesions.
939   *Neuropsychologia* **36**, 239–249 (1998).

940   39.   Behrmann, M., Moscovitch, M. & Winocur, G. Intact Visual Imagery and Impaired Visual
941   Perception in a Patient With Visual Agnosia. *J. Exp. Psychol. Hum. Percept. Perform.* **20**,
942   1068–1087 (1994).

943   40.   Grill-Spector, K. & Weiner, K. S. The functional architecture of the ventral temporal cortex
944   and its role in categorization. *Nature Reviews Neuroscience* **15**, 536–548 (2014).

945   41.   Weiner, K. S. *et al.* Defining the most probable location of the parahippocampal place area
946   using cortex-based alignment and cross-validation. *Neuroimage* **170**, 373–384 (2018).

947   42.   Wang, L., Mruczek, R. E. B., Arcaro, M. J. & Kastner, S. Probabilistic maps of visual
948   topography in human cortex. *Cereb. Cortex* **25**, 3911–3931 (2015).

949   43.   Glasser, M. F. *et al.* A multi-modal parcellation of human cerebral cortex. *Nature* **536**, 171–
950   178 (2016).

951   44.   Dijkstra, N., Bosch, S. E., Marcel, X., Van Gerven, A. J. & van Gerven, M. A. J. Vividness of
952   Visual Imagery Depends on the Neural Overlap with Perception in Visual Areas. *J. Neurosci.*
953   **37**, 1367–1373 (2017).

954   45.   Squire, L. R. Memory systems of the brain: A brief history and current perspective. in
955   *Neurobiology of Learning and Memory* **82**, 171–177 (Academic Press, 2004).

46. Renoult, L., Irish, M., Moscovitch, M. & Rugg, M. D. From Knowing to Remembering: The
Semantic–Episodic Distinction. *Trends in Cognitive Sciences* **23**, 1041–1057 (2019).

47. Ishai, A., Ungerleider, L. G. & Haxby, J. V. Distributed Neural Systems for the Generation of
Visual Images. *Neuron* **28**, 979–990 (2000).

48. Baumann, O. & Mattingley, J. B. Functional organization of the parahippocampal cortex:
Dissociable roles for context representations and the perception of visual scenes. *J.*
Neurosci. **36**, 2536–2542 (2016).

49. Silson, E. H., Steel, A., Kidder, A., Gilmore, A. W. & Baker, C. I. Distinct subdivisions of
human medial parietal cortex are recruited differentially for memory recall of places and
people. doi:10.1101/554915

50. Silson, E. H. *et al.* A Posterior-Anterior Distinction between Scene Perception and Scene
Construction in Human Medial Parietal Cortex. *J. Neurosci.* **39**, 705–717 (2019).

51. Wiggett, A. J., Pritchard, I. C. & Downing, P. E. Animate and inanimate objects in human
visual cortex: Evidence for task-independent category effects. *Neuropsychologia* **47**, 3111–
3117 (2009).

52. Thorat, S., Proklova, D. & Peelen, M. V. The nature of the animacy organization in human
ventral temporal cortex. *Elife* **8**, (2019).

53. Sha, L. *et al.* The animacy continuum in the human ventral vision pathway. *J. Cogn.*
Neurosci. **27**, 665–678 (2015).

54. Martin, A., Wiggs, C. L., Ungerleider, L. G. & Haxby, J. V. Neural correlates of category-
specific knowledge. *Nature* **379**, 649–652 (1996).

55. Konkle, T. & Caramazza, A. Tripartite organization of the ventral stream by animacy and
object size. *J. Neurosci.* **33**, 10235–10242 (2013).

56. Yamins, D. L. K. *et al.* Performance-optimized hierarchical models predict neural responses
in higher visual cortex. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 8619–8624 (2014).

57. Hong, H., Yamins, D. L. K., Majaj, N. J. & Dicarlo, J. J. Explicit information for category-
orthogonal object properties increases along the ventral stream. *Nat. Neurosci.* **19**, 613–
622 (2016).

58. Peelen, M. V. & Caramazza, A. Conceptual object representations in human anterior
temporal cortex. *J. Neurosci.* **32**, 15728–15736 (2012).

59. Hoffman, P. & Lambon Ralph, M. A. From percept to concept in the ventral temporal lobes:
Graded hemispheric specialisation based on stimulus and task. *Cortex* **101**, 107–118
(2018).

60. Güçlü, U. & van Gerven, M. A. J. Deep neural networks reveal a gradient in the complexity

of neural representations across the ventral stream. *J. Neurosci.* **35**, 10005–10014 (2015).

61.   Martin, A. GRAPES—Grounding representations in action, perception, and emotion
systems: How object properties and categories are represented in the human brain.
Psychon. Bull. Rev. **23**, 979–990 (2016).

62.   Peer, M., Brunec, I. K., Newcombe, N. S. & Epstein, R. A. Structuring Knowledge with
Cognitive   Maps   and   Cognitive   Graphs. *Trends   Cogn.   Sci.* (2020).
doi:10.1016/j.tics.2020.10.004

63.   Chrastil, E. R., Tobyne, S. M., Nauer, R. K., Chang, A. E. & Stern, C. E. Converging meta-
analytic   and   connectomic   evidence   for   functional   subregions   within   the   human
retrosplenial region. *Behav. Neurosci.* **132**, 339–355 (2018).

64.   Chrastil, E. R. Heterogeneity in human retrosplenial cortex: A review of function and
connectivity. *Behavioral Neuroscience* **132**, 317–338 (2018).

65.   Andrews-Hanna, J. R., Reidler, J. S., Sepulcre, J., Poulin, R. & Buckner, R. L. Functional-
Anatomic Fractionation of the Brain's Default Network. *Neuron* **65**, 550–562 (2010).

66.   Vass, L. K. & Epstein, R. A. Common Neural Representations for Visually Guided
Reorientation and Spatial Imagery. (2016). doi:10.1093/cercor/bhv343

67.   Robertson, C. E., Hermann, K. L., Mynick, A., Kravitz, D. J. & Kanwisher, N. Neural
Representations Integrate the Current Field of View with the Remembered 360° Panorama
in Scene-Selective Cortex. *Curr. Biol.* **26**, 2463–2468 (2016).

68.   Shine, J. P., Valdés-Herrera, J. P., Hegarty, M. & Wolbers, T. The Human Retrosplenial
Cortex   and   Thalamus   Code   Head   Direction   in   a   Global   Reference   Frame. (2016).
doi:10.1523/JNEUROSCI.1268-15.2016

69.   Nau, M., Navarro Schröder, T., Frey, M. & Doeller, C. F. Behavior-dependent directional
tuning in the human visual-navigation network. *Nat. Commun.* **11**, 3247 (2020).

70.   Cho, J. & Sharp, P. E. Head direction, place, and movement correlates for cells in the rat
retrosplenial cortex. *Behav. Neurosci.* **115**, 3–25 (2001).

71.   Jacob, P.-Y. *et al.* An independent, landmark-dominated head-direction signal in
dysgranular retrosplenial cortex. *Nat. Neurosci.* **20**, 173–175 (2017).

72.   Chen, L. L., Lin, L. H., Green, E. J., Barnes, C. A. & McNaughton, B. L. Head-direction cells in
the rat posterior cortex. I. Anatomical distribution and behavioral modulation. *Exp. brain*
Res. **101**, 8–23 (1994).

73.   Lescroart, M. D. & Gallant, J. L. Human Scene-Selective Areas Represent 3D Configurations
of Surfaces. *Neuron* **101**, 178-192.e7 (2019).

74.   Bonner, M. F. & Epstein, R. A. Coding of navigational affordances in the human visual

system. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 4793–4798 (2017).

75.   Andrews-Hanna, J. R., Smallwood, J. & Spreng, R. N. The default network and self-
1026   generated thought: Component processes, dynamic control, and clinical relevance. *Ann.*
1027   *N. Y. Acad. Sci.* **1316**, 29–52 (2014).

1028   76.   Margulies, D. S. *et al.* Situating the default-mode network along a principal gradient of
1029   macroscale cortical organization. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 12574–12579 (2016).

1030   77.   Murphy, C. *et al.* Distant from input: Evidence of regions within the default mode network
1031   supporting perceptually-decoupled and conceptually-guided cognition. *Neuroimage* **171**,
1032   393–401 (2018).

1033   78.   Ranganath, C. & Ritchey, M. Two cortical systems for memory-guided behaviour. *Nature*
1034   *Reviews Neuroscience* **13**, 713–726 (2012).

1035   79.   Ritchey, M. & Cooper, R. A. Deconstructing the Posterior Medial Episodic Network. *Trends*
1036   *Cogn. Sci.* **xx**, (2020).

1037   80.   Peer, M., Salomon, R., Goldberg, I., Blanke, O. & Arzy, S. Brain system for mental
1038   orientation in space, time, and person. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 11072–7 (2015).

1039   81.   Takahashi, N., Kawamura, M., Shiota, J., Kasahata, N. & Hirayama, K. Pure topographic
1040   disorientation due to right retrosplenial lesion. *Neurology* **49**, 464–9 (1997).

1041   82.   Aguirre, G. K. & D'Esposito, M. Topographical disorientation: a synthesis and taxonomy.
1042   *Brain* **122**, 1613–1628 (1999).

1043   83.   DiNicola, L. M., Braga, R. M. & Buckner, R. L. Parallel distributed networks dissociate
1044   episodic and social functions within the individual. *J. Neurophysiol.* **123**, 1144–1179 (2020).

1045   84.   Braga, R. M., Van Dijk, K. R. A., Polimeni, J. R., Eldaief, M. C. & Buckner, R. L. Parallel
1046   distributed networks resolved at high resolution reveal close juxtaposition of distinct
1047   regions. *J. Neurophysiol.* **121**, 1513–1534 (2019).

1048   85.   Braga, R. M. & Buckner, R. L. Parallel Interdigitated Distributed Networks within the
1049   Individual Estimated by Intrinsic Functional Connectivity. *Neuron* **95**, 457-471.e5 (2017).

1050   86.   Squire, L. R. & Zola-Morgan, S. The medial temporal lobe memory system. *Science* **253**,
1051   1380–1386 (1991).

1052   87.   Maguire, E. A. & Mullally, S. L. The hippocampus: A manifesto for change. *J. Exp. Psychol.*
1053   *Gen.* **142**, 1180–1189 (2013).

1054   88.   Maguire, E. A., Intraub, H. & Mullally, S. L. Scenes, Spaces, and Memory Traces: What Does
1055   the Hippocampus Do? *Neuroscientist* **22**, 432–439 (2016).

89.   Lee, A. C. H., Yeung, L. K. & Barense, M. D. The hippocampus and visual perception.

Frontiers in Human Neuroscience **6**, 91 (2012).

90.   Pearson, J., Rademaker, R. L. & Tong, F. Evaluating the Mind's Eye. *Psychol. Sci.* **22**, 1535–
1542 (2011).

91.   Bartolomeo, P., Hajhajate, D., Liu, J. & Spagna, A. Assessing the causal role of early visual
areas in visual mental imagery. *Nat. Rev. Neurosci.* **21**, 517–517 (2020).

92.   Grill-Spector, K., Weiner, K. S., Kay, K. & Gomez, J. The Functional Neuroanatomy of Human
Face Perception. *Annu. Rev. Vis. Sci.* **3**, 167–196 (2017).

93.   Freiwald, W., Duchaine, B. & Yovel, G. Face Processing Systems: From Neurons to Real-
World Social Perception. *Annu. Rev. Neurosci.* **39**, 325–346 (2016).

94.   Steeves, J. K. E. *et al.* The fusiform face area is not sufficient for face recognition: Evidence
from a patient with dense prosopagnosia and no occipital face area. *Neuropsychologia* **44**,
594–609 (2006).

95.   Duchaine, B. & Yovel, G. A Revised Neural Framework for Face Processing. *Annu. Rev. Vis.*
Sci. **1**, 393–416 (2015).

96.   MARKS, D. F. VISUAL IMAGERY DIFFERENCES IN THE RECALL OF PICTURES. *Br. J. Psychol.*
64, 17–24 (1973).

97.   Raichle, M. E. The Brain's Default Mode Network. *Annu. Rev. Neurosci.* **38**, 433–447 (2015).

98.   Li, X., Morgan, P. S., Ashburner, J., Smith, J. & Rorden, C. The first step for neuroimaging
data analysis: DICOM to NIfTI conversion. *J. Neurosci. Methods* **264**, 47–56 (2016).

99.   Fischl, B. *et al.* Whole brain segmentation: Automated labeling of neuroanatomical
structures in the human brain. *Neuron* **33**, 341–355 (2002).

100.   Dale, A. M., Fischl, B. & Sereno, M. I. Cortical surface-based analysis: I. Segmentation and
surface reconstruction. *Neuroimage* **9**, 179–194 (1999).

101.   Fischl, B. FreeSurfer. *NeuroImage* **62**, 774–781 (2012).

102.   Weiskopf, N., Hutton, C., Josephs, O. & Deichmann, R. Optimal EPI parameters for
reduction of susceptibility-induced BOLD sensitivity losses: A whole-brain analysis at 3 T
and 1.5 T. (2006). doi:10.1016/j.neuroimage.2006.07.029

103.   Cox, R. W. AFNI: Software for analysis and visualization of functional magnetic resonance
neuroimages. *Comput. Biomed. Res.* **29**, 162–173 (1996).

104.   Smith, S. M. *et al.* Advances in functional and structural MR image analysis and
implementation as FSL. in *NeuroImage* **23**, S208–S219 (Academic Press, 2004).

105.   Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W. & Smith, S. M. FSL.
Neuroimage **62**, 782–790 (2012).

106.   Woolrich, M. W. *et al.* Bayesian analysis of neuroimaging data in FSL. *Neuroimage* **45**,
(2009).

107.   Griffanti, L. *et al.* ICA-based artefact removal and accelerated fMRI acquisition for
improved resting state network imaging. *Neuroimage* **95**, 232–247 (2014).

108.   Poser, B. A., Versluis, M. J., Hoogduin, J. M. & Norris, D. G. BOLD contrast sensitivity
enhancement and artifact reduction with multiecho EPI: Parallel-acquired inhomogeneity-
desensitized fMRI. *Magn. Reson. Med.* **55**, 1227–1235 (2006).

109.   Evans, J. W., Kundu, P., Horovitz, S. G. & Bandettini, P. A. Separating slow BOLD from non-
BOLD baseline drifts using multi-echo fMRI. *Neuroimage* **105**, 189–197 (2015).

110.   Kundu, P., Inati, S. J., Evans, J. W., Luh, W. M. & Bandettini, P. A. Differentiating BOLD and
non-BOLD signals in fMRI time series using multi-echo EPI. *Neuroimage* **60**, 1759–1770
(2012).

111.   DuPre, E. *et al.* ME-ICA/tedana: 0.0.6. (2019). doi:10.5281/ZENODO.2558498

112.   Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. & Team, R. C. nlme: Linear and Nonlinear
Mixed Effects Models. (2019).

113.   R Core Team. R: A language and environment for statistical computing. R Foundation for
Statistical Computingitle. (2013).

114.   Lenth, R., Singmann, H., Love, J., Buerkner, P. & Herve, M. emmeans. (2020).

1108

1109

1110   **A network linking perception and memory systems in posterior cerebral cortex.**

1111   Extended data

1112

1113

1114   Adam Steel[1], Madeleine M. Billings[1], Edward H. Silson[2], Caroline E. Robertson[1]

1115

[1]Department of Psychology and Brain Sciences, Dartmouth College, Hanover, NH, 03755

[2]Psychology, School of Philosophy, Psychology, and Language Sciences, University of Edinburgh,
Edinburgh, UK EH8 9JZ

Corresponding author: Adam Steel, Department of Psychology and Brain Sciences, Dartmouth
College, 3 Maynard Street, Hanover, NH, 03753; email: adam.steel@dartmouth.edu; tel: (202)
640 9340

1122

a

subj001   subj002   subj003   subj004

subj005   subj006   subj007   subj008

subj009   subj010   subj011   subj012

subj013   subj014

-15 > t-statistic < 15

Memory for places vs. people
t > 3.3, p < 0.001

b

subj002   subj005   subj006   subj008   subj011   subj013

-30 > t-statistic < 30

Memory for places vs. people
t > 5.0, p < 0.0005

Multi-echo ventral surface

Ant

Places
People

Lat   15   30   Med

Post

1123

Extended data Fig. 1. Place-memory selective activation is anterior to scene-selective perceptual activation in all
participants. *a. In each participant, scene-selective perceptual areas (parahippocampal place area [PPA], occipital*
place area [OPA], and medial place area [MPA]) were localized by comparing BOLD activation when viewing images
of places compared to people); the white outlined region shows each scene-selective area (PPA, OPA, MPA)
thresholded at vertex-wise p < 0.001. Place-memory selective areas were localized by comparing BOLD activation
when participants recalled personally familiar places versus personally familiar faces. These maps were thresholded
at vertex-wise p < 0.001. In all participants and each cortical surface, the memory activation falls significantly anterior
to scene perception activation. Face-selective perception (grey outline) and face-selective memory (cool colors) areas
on the ventral surface are shown for comparison thresholded at (vertex-wise p < 0.001). At this threshold, people-
memory activation on the ventral surface was only visible in 12/14 participants on the right hemisphere and 13/14
participants in the left hemisphere (not shown). In contrast to scene areas, the center of mass of people-memory was
not anterior to face-perception. b. Signal dropout from the air-tissue interface of the ear canal is known to obscure
activity in the lateral and anterior temporal lobe, and therefore could prevent us from observing an anterior bias
during face memory recall activity relative to perception of faces. To ensure this was not the case, we replicated the
comparison of perception and memory localizers using multi-echo fMRI in a subset of participants, with a short-TE
(11 ms) to mitigate the dropout artifact [108–110]. We were able to replicate the anterior bias in place-memory selective
activation (relative to activation during scene-perception) in the replication cohort. In addition, we did not observe
any consistent anterior bias for people memory activation relative to activation during face perception. All
participants unthresholded activation maps for both perception and memory localizers can be found in Extended data
file 1.

1144

1145

Subject 3 **Subject 2** **Subject 12** **Subject 14**

PPA (Experiment 1)

Perception of Scenes > (Faces + Objects)
t > 3.3, p < 0.001

1146

Extended data Fig. 2. Contrasting scenes versus faces and objects does not impact the location of scene-perceptual
areas. *The anterior bias of the place-memory areas relative to scene-perception areas persists when using a more*
conservative contrast to define scene areas (scenes > faces + objects)[41], instead of the contrast used in Experiment 1
(scenes > faces). Left. Four example participants demonstrate the minimal impact of including visually objects when
defining PPA. PPA defined by comparing blocks of scenes versus faces (as in Experiment 1) is outlined in black, and
the PPA defined by comparing blocks scenes versus faces and objects is shown in hot colors. There was not significant
shift in the weighted center-of-mass of PPA between the two ROI definitions (Left hemisphere: t(13) = 0.99, p = 0.34,
Right hemisphere: t(13) = 0.96, p = 0.35), nor was there any difference in the center-of-mass of the two definitions of
MPA (Left hemisphere: t(13) = 0.37, p = 0.71, Right hemisphere: t(13) = 1.65, p = 0.12). The weighted center of mass
of OPA was shifted significantly posterior when objects were included in the contrast (Left hemisphere: t(13) = 2.87,
p = 0.013, Right hemisphere: t(13) = 2.22, p = 0.045), and thus the original contrast would have been more difficult
to observe an anterior shift of memory relative to perception. Right. Polar plots comparing scene-perception and
place-memory areas. The center of the polar plot refers to the weighted center-of-mass of the scene-selective regions
defined by contrasting scenes versus faces and objects in each participant, and each data point refers to the weighted
center-of-mass of the place-memory area. The center-of-mass of the place-memory area was significantly anterior
to the weighted center-of-mass of the scene-perception area on all surfaces (Medial surface -- Left hemisphere: t(13)
= 7.00, p < 0.0001, Right hemisphere: t(13) = 5.24, p = 0.00016; Ventral surface -- Left hemisphere: t(13) = 10.27, p <
0.0001, Right hemisphere: t(13) = 6.97, p < 0.0001; Lateral surface -- Left hemisphere: t(13) = 16.67, p < 0.0001, Right
hemisphere: t(13) = 9.89, p < 0.0001). The polar plots are ordered with ventral (top), lateral (middle), and medial
(bottom). Anterior is towards the top of the figure for all surfaces, and the distance is given in millimeters.

1167

1168

1169

1170

1171 ***Extended data Fig 3. The ventral place-memory area falls anterior to probabilistically-defined PPA.*** *Left. Data from*
1172 *an example participant. The ventral place-memory area (hot colors) is anterior to the most probable location of PPA[41].*
1173 *Right. Polar plot showing the center of mass of the most probable location of PPA (Polar plot center) compared to*
1174 *the weighted center of mass of the ventral place memory area (data points). Consistent with the individually localized*
1175 *PPA (Figure 1), the center of mass of the ventral place memory area is anterior to the probabilistically defined PPA.*

1176

1177

Extended data Fig 4. Normalized place memory response along the posterior-anterior axis of the ventral scene-
perception area (PPA). *In each participant (light grey lines), PPA was divided into 20 evenly spaced bins, and we*
calculated the mean place-memory activity (place-memory versus people-memory) from each bin. The place-memory
activity from each participant was then demeaned and normalized, and the group average was calculated (black
line). ***Note that the y-axis shows demeaned and normalized place-memory responses, and therefore the direction***
of response (positive/negative) should not be interpreted. *At each bin, the red heatmap shows the proportion of*
participants with a place-memory area vertex within a given bin. At the group level, in both the left and right
hemisphere, there was an apparent transition from perceptual to memory-related processing from approximately
50%-70% of the distance along the anterior-posterior axis of PPA. However, at the individual participant level, the
transition was often sharper. It is impossible to estimate the true steepness of the transition between scene-
perception and place-memory activity, however, due to the inherent smoothness of fMRI data. As such, these results
should be interpreted with caution.

1190

1191

1192 ***Extended data Fig 5. Minimal overlap between cortical retinotopic maps and the place memory areas on the***
1193 ***ventral and lateral surfaces.*** *The overlap between the maximum probability project of cortical retinotopic maps from*
1194 *Wang et al. 2014 overlaid on one individual participant's ventral and lateral place-memory activity. The place-*
1195 *memory areas on the ventral surface fall largely anterior to the retinotopic regions PHC1 and PHC2. On the lateral*
1196 *surface, the place-memory area primarily fell outside of retinotopic cortex. Inset. Group-average pie charts showing*
1197 *the proportion of surface vertices of the place memory areas contained within each retinotopic region at the*
1198 *individual participant-level. Across all participants, the majority of vertices fell outside of the cortical retinotopic*
1199 *maps. The medial surface is not shown due to the lack of retinotopic maps in medial parietal cortex.*

Extended data fig. 6. Location of scene perception activity in relation to known functional and anatomical landmarks. left. Unlike the place-memory areas, the scene-perception fall largely within retinotopic cortex. We conducted a group analysis of the scene-perception localizer (place > people memory recall; thresholded at vertex-wise t > 7.3, p = 1e-6) and compared the resulting activation to the most probable location of the cortical retinotopic maps using the atlas (overlaid) from Wang et al. 2014. Right. Unlike the place-memory areas, the scene-perception areas fall in areas of cortex associated with perception based on their location relative to a widely used anatomical parcellation (Ventral: PHC 1-3, Medial: dorsal transitional visual area, Lateral: LO1/LO2, V3; Glasser et al. 2016).

1211

Extended data Fig.7. Participants reported no difference in vividness between imagery for people and places.
Outside of the scanner, participants rated each personally familiar stimulus for vividness of visual imagery on a scale
of 1-5, where 1 indicates no imagery possible and 5 indicates "as if you were seeing the <stimulus>." There was no
significant difference in vividness between people and place stimuli (t(13) = 0.92, p = 0.373), suggesting that the
topographical difference between place and people imagery activation was not due to differences of vividness
between the stimulus categories. Additionally, we sought to rule out that the richness of recollection might impact
the activity of the scene-perception and place-memory areas. So, we tested whether the "vividness" of a memory
covaried with the magnitude of the place-memory network response on a given trial during Experiment 1 using a
linear mixed effects model with Network (scene-perception/place-memory), stimulus type (person/place),
hemisphere (left/right), and vividness as factors. We found no overall effect of vividness in any region (Main effect of
Vividness -- Lateral: F(1,3716) = 0.7647, p = 0.3819; Ventral: F(1,3716) = 0.1657, p = 0.684; medial -- F(1,3716) =
0.0001, p = 0.99), nor did we find any interaction between vividness and stimulus type (Vividness x Stimulus type
interaction -- lateral: F(1,3716) = 0.4437, p = 0.5054; ventral: F(1,3716) = 1.0179 p = 0.3131; medial: F(1,3716) =
1.5674, p = 0.2107) or vividness and network (Vividness x Network interaction -- lateral: F(1,3716) = 0.001, p = 0.97;
ventral: F(1,3716) = 0.525, p = 0.46; medial: F(1,3716) = 0.016, p = 0.90).

1227

1228

1229

Extended data Figure 7. **Anterior shift of place-memory relative to scene perception remains consistent after constraining to maximally selective surface vertices.** (Left). Top 300 in each participant most scene-selective surface vertices were used for as regions of interest for Experiments 2-4. (Right). Across the lateral, medial and ventral surfaces, we compared the weighted center-of-mass of both the scene-perception and place-memory selective areas after constraining the regions to the top 300 most scene-perception/place-memory preferring surface vertices. Using the constrained ROI definitions, we still observed a significant anterior shift from scene-perception to place-memory on all cortical surfaces in both the left and right hemisphere (Lateral - left hemisphere: t(13) = 9.35, p < 0.0001, right hemisphere: t(13) = 8.62, p < 0.0001; Ventral - left hemisphere: t(13) = 4.84, p = 0.0003, right hemisphere: t(13) = 7.45, p < 0.0001; Medial - left hemisphere: t(13) = 2.86, p = 0.013, right hemisphere: t(13) = 2.36, p = 0.034).

1241

1242

1243

1244

1245

Extended data Fig. 9. Hippocampus responds preferentially to familiar place movies, while amygdala does not. *As*
an exploratory analysis, we tested how the hippocampus, which we expected would preferentially respond to familiar
images, responded in Experiment 2, where participants viewed familiar and unfamiliar panning movies. The average
T-statistic from all voxels within the hippocampus and amygdala (defined by Freesurfer segmentation) was extracted
for each participant. We then compared the activation for each region separately using a linear mixed effects model
with Hemisphere (left/right) and Familiarity (familiar/unfamiliar) as factors. a. As predicted, the hippocampus
preferentially activated to familiar compared to unfamiliar stimuli (Main effect of Familiarity: F(1,39)=8.14, p =
0.0069; t(13)=2.54, p = 0.004). Activation was stronger in the right hemisphere compared to the left hemisphere
(Main effect of Hemisphere: F(1,39)=8.61, p = 0.0056; t(13)=2.38, p=0.0055); however, there was no interaction
between Hemisphere and Familiarity (Hemisphere x Familiarity interaction: F(1,39)=0.33, p = 0.56). b. In contrast, in
the amygdala, there was no effect of Familiarity (F(1,39)=0.29, p = 0.59), Hemisphere (F(1,39)=0.07, p = 0.78), or
interaction between Hemisphere and Familiarity (F(1,39)=0.003, p = 0.95), arguing against a simple account of
attention modulating the response to familiar compared to unfamiliar stimuli.

1259

1260

Extended data Fig. 10. Early visual cortex responses to familiar and unfamiliar place movies do not differ. *One*
explanation for the increased activation to familiar compared to unfamiliar stimuli is that greater attention is being
paid to familiar stimuli, causing a global increase in activation of visually-responsive areas. To confirm that this was
not the case, we compared activation of the occipital pole (defined in each participant from their Freesurfer
parcellation) when participants viewed familiar versus unfamiliar panning movies. We compared the responses using
a mixed effects model with Hemisphere (lh/rh) and Familiarity (familiar/unfamiliar) as factors. We found that
responses in the right hemisphere were stronger than the left hemisphere (Main effect of Hemisphere: $F(1,39) =$
13.29, $p =0.0008$; $t39=3.65$, $p = 0.0008$). However, we found no effect of Familiarity ($F(1,39)=3.19$, $p = 0.08$) or
interaction between Hemisphere and Familiarity ($F(1,39)=0.01$, $p = 0.91$), confirming that a simple attentional
account cannot explain the familiarity effect observed in the place-memory areas.

1271

1272

1273

***Extended data Fig. 11. Early visual cortex responds more strongly to perception than mental imagery**. As an*
exploratory analysis, we investigated the response in early visual cortex (occipital pole defined in individual
participants from their Freesurfer segmentation) during perception of unfamiliar places and mental imagery of
familiar places. We compared the responses using a mixed effects model with Hemisphere (lh/rh) and Task
(imagery/perception) as factors. As expected, we found that early visual cortex responded significantly more to
perception compared to mental imagery (Main effect of Task: F(1,39) = 541.38, p < 0.0001; t(13) = 16.83, p < 0.0001).
There was no effect of Hemisphere (F(1,39) = 1.22, p = 0.22) or interaction between Hemisphere and Task (F(1,39) =
1.05, p = 0.31).

1282

1283

1284

Extended data Fig. 12. Both hippocampus and amygdala respond more strongly during perception compared to
mental imagery. *As an exploratory analysis, we investigated whether the hippocampus and amygdala — subcortical*
areas implicated in memory processes — responded differently when viewing panning movies of unfamiliar places
and mental imagery of familiar places. The average T-statistic from all voxels within the hippocampus and amygdala
(defined by Freesurfer segmentation) was extracted for each participant. We then separately compared the
activation of each region using a linear mixed effects model with Hemisphere (left/right) and Task
(Imagery/Perception) as factors. Both the hippocampus and amygdala responded more strongly during perception
compared to mental imagery (Hippocampus: F(1,39)=28.67, p < 0.0001 t(13)=3.31, p < 0.0001; Amygdala:
F(1,39)=37.22, p < 0.0001; t(13)=4.01, p < 0.0001). Responses did not differ by hemisphere in either region
(Hippocampus: F(1,39)=0.57, p = 0.45; Amygdala: F(1,39)=2.86, p = 0.09), and there was no interaction between
Hemisphere and Task in either region (Hippocampus: F(1,39)=1.65, p = 0.20; Amygdala: F(1,39)=0.02, p = 0.89).

1296
1297

1298

Combined perception and memory ROI

1299

Extended data Fig. 13. All cortical surfaces (lateral, ventral, medial) show positive activation during perception
and imagery when scene-perception and place-memory areas are considered together. *Previous studies of mental*
imagery (e.g. [29]), have found that mental imagery and perception activate shared neural substrates. However, these
studies have not considered the scene-perception and place-memory areas separately. Therefore, in order to relate
our results to previous studies of mental imagery, the average t-statistic of the scene-perception and place-memory
areas from each cortical surface were extracted and combined into a single "combined ROI". Consistent with prior
work, we found that mental imagery positively activated the combined ROI in the majority of participants (lateral:
10/14, mean t-stat = 1.96; ventral: 9/14, mean t-stat = 0.56; medial: 10/14, mean t-stat = 3.12). Group activation of
the lateral (t(13) = 2.99, p = 0.011) and medial (t(13) = 4.08, p = 0.0013) surfaces were significantly above zero during
mental imagery. Despite the majority of participants showing positive activation, the ventral surface was not
significantly above zero at the group level (t(13) = 1.03, p = 0.32).

1311

1312   a

1313

1314   b

1315

1316   c

1317

Extended data Fig. 14. Results from all experiments remain unchanged when scene-perception areas are defined
using the contrast scenes > faces + objects. *a. Experiment 2. Place memory areas exhibit an increased response to*
viewing panning movies of familiar places that is greater than the scene perception areas defined using the contrast
Scenes > Faces + Objects. Lateral (LPMA v OPA): t(13) = 2.65, p = 0.02, D = 0.7; Medial (MPMA v MPA): t(13) = 5.15,

p = 0.00019, D = 1.13; Ventral (VPMA v PPA): t(13) = 3.18, p = 0.007, D = 0.85). b. Experiment 3. Experiment 3. (left)
The place memory areas and scene perception areas exhibit greater within network connectivity compared to
between network connectivity when the scene perception areas defined using the contrast Scenes > Faces + Objects.
P v PxM: t(12) = 2.97, p = 0.01, D = 2.11; M v PxM: t(12) = 2.20, p = 0.048, D = 1.59; P v M: T(12) =3.4 , p = 0.005 , D =
0.93. (right) The scene perception network is more connected to primary visual cortex than the place memory
network, while the place memory network is more connected to hippocampus than the scene perception network
when the scene perception areas are defined using the contrast Scenes > Faces + Objects. PxV1 - PxHipp v MxV1 -
MxHipp: t(12) = 6.3, p < 0.0001, D = 4.52. c. Experiment 4. Place memory areas respond more when recalling familiar
places compared to the scene perception areas when defined using the contrast Scenes > Faces + Objects. Lateral:
Perception - t(13) = 7.092; p < 0.0001 , D = 2.3, Memory - t(13) = 8.509; p < 0.0001, D = 3.94; Ventral: Perception -
t(13) = 4.803; p = 0.0003, D = 1.19, Memory - T(13) = 6.50; p < 0.0001, D = 2.77; Medial: Perception - t(13) = 0.83; p
= 0.41, D = 0.25, Memory - T(13) = 4.82; p = 0.0003 , D = 1.17.

1334

1335   a

1336

1337   b

1338

1339   c

1340

Extended data Fig. 15. Results from all experiments remain unchanged when scene-perception areas are defined
using the 100 vertices closest to the scene-perception and place-memory localizer activation peaks. *a. Experiment*
2. Place memory areas exhibit an increased response to viewing panning movies of familiar places that is greater
than the scene perception areas are defined using the 100 vertices closest to the localizer activation peak. Medial:

t(13) = 5.98, p < 0.0001, D = 1.60; Ventral: t(13) = 4.80, p = 0.0003, D = 1.28; Lateral: t(13) = 4.32, p = 0.0008 , D =
1.28. b. Experiment 3. (Left) The place memory areas and scene perception areas exhibit greater within network
connectivity compared to between network connectivity when the scene perception areas defined using the 100
vertices closest to the localizer activation peak. P v PxM: t(12) = 4.65, p = 0.0005, D = 2.84; M v PxM: t(12) = 3.45, p
= 0.004, D = 2.14; P v M: t(12) =0.23, p = 0.79, D = 0.1. (Right) The scene perception network is more connected to
primary visual cortex than the place memory network, while the place memory network is more connected to
hippocampus than the scene perception network when the scene perception areas are defined using the contrast
Scenes > Faces + Objects. PxV1 - PxHipp v MxV1 - MxHipp: t(12) = 5.8, p < 0.0001, D = 4.14. c. Experiment 4. Place
memory areas respond more when recalling familiar places compared to the scene perception areas when the areas
are defined using the 100 vertices closest to the localizer activation peak. Lateral: Perception – t(13) = 6.91, p < 0.0001
, D = 2.43, Memory – t(13) = 7.53, p < 0.0001, D = 2.7; Ventral: Perception – t(13) = 3.83, p = 0.002, D = 1.15, Memory
– t(13) =6.34 , p < 0.0001, D = 1.85; Medial: Perception – t(13) =5.7 , p < 0.0001, D = 1.1; Memory – t(13) = 3.97, p =
0.0015, D = 1.12.

1358

Extended data file 1. All participants unthresholded activation maps. This file is a gif image
displaying, alternatingly, the unthresholded activation maps for the scene perception (perception
of scene versus faces) and place memory (memory recall of places versus people) localizer tasks.
The anterior shift of memory relative to perception can be seen in all participants. Note that, for
clarity, only the right hemisphere is shown because no significant interaction with hemisphere
was found.

Extended data Video 1. Example panning video of a familiar outdoor place used in Experiments
2 and 4 (Oxford, United Kingdom).

Extended data Video 2. Example panning video of a familiar indoor place used in Experiments 2
and 4 (Hanover, NH, United States).

Extended data Video 3. Example panning video of a familiar city street used in Experiments 2 and
4 (Yongkang, Zhejiang, China).

Extended data Video 4. Example panning video of a familiar place used in Experiments 2 and 4
(Zandvoort, Netherlands).

1373

1374

1375

1376