1

3 **HSATII RNA is induced *via* a non-canonical ATM-regulated DNA-damage response**

4 **pathway and facilitates tumor cell proliferation and movement**

9 Maciej T. Nogalski[a,1] and Thomas Shenk[a]

15 [a] Department of Molecular Biology, Princeton University, Princeton, NJ 08544-1014, USA

18 [1] To whom correspondence may be addressed: nogalski@princeton.edu

26 **Keywords:** HSATII, DNA damage response, human cytomegalovirus, breast cancer cells

ABSTRACT

Pericentromeric human satellite II (HSATII) repeats are normally silent, but can be actively transcribed in tumor cells, where increased HSATII copy number is associated with a poor prognosis in colon cancer, and in human cytomegalovirus (HCMV)-infected cells, where the RNA facilitates viral replication. Here, we report that HCMV infection or treatment of ARPE-19 diploid epithelial cells with the DNA-damaging agents, etoposide and zeocin, induced HSATII RNA expression, and a kinase-independent function of ATM was required for the induction. Additionally, various breast cancer cell lines growing in adherent, 2-dimensional cell culture expressed HSATII RNA at different levels, and levels were markedly increased when cells were either infected with HCMV or treated with zeocin. High levels of HSATII RNA expression correlated with enhanced migration of breast cancer cells, and knockdown of HSATII RNA reduced cell migration and the rate of cell proliferation. Our investigation links high expression of HSATII RNA to the DNA damage response, centered on a non-canonical function of ATM, and demonstrates a role for the satellite RNA in tumor cell proliferation and movement.

SIGNIFICANCE

HSATII RNA is associated with cancer progression, immunostimulation and, as we recently reported, it plays an important role in herpesvirus infections. However, the understanding of cellular processes responsible for the expression of HSATII RNA has been limited. Our current investigation identified a non-canonical, ATM kinase-independent DNA-damage response pathway as a common cellular mechanism regulating HSATII RNA induction in virus-infected cells or cells treated with DNA-damaging agents. Additionally, our study provides a link between expression of HSATII RNA and the cellular growth and migration phenotypes of cancer cells, establishing a new paradigm to study the biological consequences of HSATII RNA expression, i.e., treatment of normal diploid and tumor cells with DNA-damaging agents.

INTRODUCTION

Repetitive DNA sequences comprise more than 50% of the human genome (1) and include short (SINE) and long (LINE) interspersed nuclear elements, DNA transposons, long terminal repeat (LTR) transposons and satellite repeats. Even though repetitive DNA elements are ubiquitous in the human genome, there is a relatively limited understanding of their functions and the molecular mechanisms regulating their expression. Satellite DNAs (satDNAs), which account for ~3% of the genome (1), are constituents of centromeric and pericentromeric heterochromatin, and have been implicated in chromosome organization and segregation, kinetochore formation as well as heterochromatin regulation (2). Next-generation sequencing showed these genomic sites, originally thought to be largely transcriptionally inert, can produce non-coding RNAs (ncRNAs), contributing to satDNA functions (3-6).

Altered patterns of transcription, including the induction of ncRNA accumulation, occur in tumors (7-9). Several ncRNAs originating from satDNA regions of the genome are expressed in cancer cells, such as human alpha-satellite repeat (Alpha/ALR) RNA, human satellite II (HSATII) RNA and its mouse counterpart GSAT RNA (10-13). While some satDNA transcription is stress-dependent (14) or triggered during apoptotic, differentiation or senescence programs in cells (15, 16), HSATII RNA accumulation was found to be refractory to these generalized stressors. Rather, it was induced in a series of colorectal cancer cells when they were grown under non-adherent conditions or as xenografts in mice (17). High expression of Alpha/ALR and HSATII RNAs can lead to their reverse transcription and stable reintegration into the human genome, expanding their genomic copy numbers (17). Elevated copies of genomic HSATII sequences were found in primary human colon tumors and correlated with lower survival rates of colon cancer patients (17). Additionally, HSATII and GSAT RNAs have a nucleotide motif usage that is distinct from that commonly seen in most non-coding transcripts, resulting in their immunostimulatory potential (18).

Viruses de-regulate many biological processes, often in a similar manner to aberrations seen in cancer cells. Those biological changes caused by infections not only resemble cancer, but in some cases contribute to oncogenesis (19). Indeed, it is estimated that 12-20% of all cancers have a viral etiology and often are linked to persistent or chronic viral infections (20-22).

Human cytomegalovirus (HCMV) is a β-herpesvirus that infects a large percentage of the adult population worldwide. Infection in immunocompetent people is typically asymptomatic. In contrast, HCMV is a life-threatening opportunistic pathogen in immunosuppressed individuals (23-25), and a major infectious cause of birth defects (26). Many of the biological changes seen in HCMV-infected cells resemble those common to cancers, including activation of pro-oncogenic pathways, changes in cellular metabolism and increased cell survival (27-32). Given similarities in the molecular symptoms of HCMV infections and tumorigenesis, as well as its reliable presence in glioblastoma tumors, HCMV has been suggested to play a role, perhaps an oncomodulatory role, in the etiology of several human

cancers (29, 33-39). However, its high prevalence has made it difficult to establish causality in the

disease (40).

Recently, we determined that HCMV infection induces HSATII RNA in fibroblasts, among other satDNA

transcripts, to a similar extent as reported for tumor cells (41). Elevated HSATII RNA was also detected

in biopsies of CMV colitis, showing that the RNA is elevated in human infections and raising the

possibility that elevated HSATII RNA may influence the pathogenesis of HCMV disease. When the

amount of HSATII RNA in HCMV-infected cells was reduced by knockdown (KD) with locked nucleic

acid (LNA) oligonucleotides, viral gene expression, DNA accumulation and yield were reduced.

Moreover, KD experiments revealed that HSATII RNA affects several infected-cell processes, including

protein stability, posttranslational modifications and motility of infected cells.

The HCMV IE1 and IE2 proteins induce HSATII RNA in infected cells (41); and are known to regulate

the Rb and E2F protein families, which contribute to efficient viral replication in part through induction of

a DNA damage response (DDR) (42, 43). Here, we have tested the possibility that these two activities

of the viral IE1 and IE2 proteins - induction of a DDR and HSATII RNA - are related. We report that the

DNA-damaging agents, zeocin and etoposide, which induce DNA double-strand breaks (DSBs) and a

DDR (44, 45), markedly induce HSATII RNA in diploid epithelial cells growing in adherent, 2-

dimensional culture. Consistent with this observation, ATM, a key regulator of DDRs, is required for

induction of the satellite RNA by infection or DNA-damaging agents. HSATII RNA induction requires a

non-canonical, kinase-independent ATM function. Finally, we demonstrate that HSATII RNA is

expressed in a variety of breast cancer cell lines grown in adherent culture, the RNA is further induced

in these cell lines by HCMV infection or zeocin treatment, and HSATII RNA levels correlate with their

migration and proliferation rates. In sum, our data argues that the DDR leads to elevated HSATII RNA

levels, potentially enhancing the fitness of both HCMV and tumor cells.

RESULTS

The DDR regulates HSATII RNA expression. Ectopic expression of the HCMV IE1 and IE2 proteins

can induce HSATII RNA expression in the absence of infection (41). As IE1 and IE2 induce a DDR that

supports efficient viral replication (42, 43), we speculated that the DDR could play a role in the

expression of human satDNAs.

To test this hypothesis, ARPE-19 cells were stressed by three regimens that induce a DDR: UV-C

irradiation, H_2O_2 treatment or serum withdrawal (46-48). Exposure of these diploid epithelial cells, which

normally express HSATII RNAs at very low basal levels, to these stressing agents induced HSATII

RNA to a very limited extent, 2-3 fold (Fig. 1*A*), in contrast to the 100-1000-fold induction observed in

response to virus infection (41). As a proxy to monitor successful DDR activation, the induction of p53

activity was monitored by assaying the levels of p53-dependent cyclin-dependent kinase inhibitor 1

(p21; encoded by the CDKN1A gene) (46) and p53-regulated Damage Induced Noncoding (DINO)

lncRNA (49). As expected, both were elevated in cells exposed to all three treatments (Fig. 1*A*).

We reasoned that the failure of these treatments to induce HSATII RNA might be a consequence of the

specific type of DNA damage and the resulting response that they induce. For example, while UV-C

irradiation can lead to DNA double-strand brakes (DSBs), the specific role of UV-C in DNA damage

depends on the replicative state of treated cells, as those breaks usually arise from the replication of

unrepaired UV-induced DNA lesions (50, 51). It was possible that UV-C and the other treatments were

not efficient in inducing HSATII RNA. Therefore, we tested the effects of zeocin and etoposide.

Etoposide, similarly to doxorubicin, is an anticancer agent that poisons topoisomerase II, stabilizing

DNA-protein complexes, which ultimately leads to stalled replication forks (52). Zeocin belongs to the

bleomycin/phleomycin-family of antibiotics and, as a radiomimetic, has similar effects on DNA as

ionizing radiation (53). Both drugs induce DNA DSBs (44, 45), and etoposide has been shown to

stimulate the expression and retrotransposition of another repeated element, mouse B2 SINE RNA (54,

55). ARPE-19 cells were exposed to increasing concentrations of zeocin or etoposide and HSATII,

CDKN1A and DINO RNA expression were monitored. Zeocin caused a rapid, dose-dependent

induction of HSATII RNA, reaching an ~500-fold increase relative to untreated cells, at the maximum

dose tested (400 μg/mL) (Fig. 1*B*). Etoposide also elevated HSATII RNA levels, reaching an ~700-fold

increase at the highest dose tested (500 μM) (Fig. 1*C*). CDKN1A and DINO RNA expression mirrored

HSATII RNA accumulation, reaching a 6-7-fold induction in cells exposed to 400 μg/mL of zeocin (Fig.

1*B*) and ~25-fold induction in cells treated with 500 μM of etoposide (Fig. 1*C*). No ARPE-19 cell toxicity

was detected at any drug concentration after a 24 h treatment (Figs. 1*B* and *C*).

The kinetics of HSATII, CDKN1A and DINO RNA expression were monitored by RT-qPCR from 0-24 h

post treatment (hpt) with zeocin (200 μg/mL) or etoposide (200 μM). HSATII RNA expression started to

increase between 7-12 hpt with either zeocin or etoposide, reaching a 300-fold induction at 24 hpt

compared to solvent control-treated cells (Figs. 1*D* and *E*). The dynamics of CDKN1A and DINO RNA

expression were accelerated compared to HSATII RNA expression, as both transcripts reached their

maximum levels within the first 2-4 hpt, suggesting a possible decoupling of DDR/p53-dependent

pathways from the mechanism regulating HSATII RNA induction.

Zeocin treatment substantially mimics the transcriptional effect of HCMV infection on HSATII

RNA expression. To compare the induction HSATII RNA by virus versus DNA-damaging agent,

ARPE-19 cells were infected with HCMV or treated with zeocin. The infection used an epithelial cell-

grown HCMV strain (TB40/E-GFP-epi) that efficiently infects ARPE-19 cells (56). Mock-infection and

solvent treatment served as controls. After 24 h, RNA was isolated and analyzed by using RNA-seq. By

analyzing only uniquely mapped HSATII reads, RNA-seq analysis confirmed that zeocin effectively

induces HSATII expression in ARPE-19 cells (Fig. 2*A*). Moreover, as in HCMV-infected cells (41),

HSATII RNA in zeocin-treated cells is produced preferentially from chromosome 1, 2, 10, and 16. We

detected a higher prevalence of HSATII-specific reads from chromosome 1 in cells exposed to zeocin

(~15%) compared to only ~5% of these reads in samples from HCMV-infected cells (Fig. 2B). However,

in both HCMV- or zeocin-treated cells the majority (~86% or 76%, respectively) of HSATII specific

reads mapped to chromosome 16 (Fig. 2*B*).

Read coverage tracks at the HSATII-rich locus on chromosome 16 were mapped with the Integrative

Genomics Viewer (IGV) using STAR aligner (57)-created BAM files (Fig. 2*C*). The data from mock-

infected and solvent-treated cells showed very few, if any, reads mapped to this HSATII locus (Fig. 2*C*).

The read coverage tracks based on RNA-seq data from HCMV-infected or zeocin-treated ARPE-19

cells display a similar distribution in this HSATII–rich locus spanning approximately 17 kb of DNA.

Therefore, these data strengthened our view that common mechanisms might be responsible for the

induction of HSATII RNA in both HCMV-infected and DNA-damaging drug-treated cells.

Zeocin treatment partially mimics the HCMV-induced transcriptome. To assess common biological

processes regulated by infection and drug, ARPE-19 cells were HCMV-infected or zeocin-treated, and,

again, mock-infection or solvent treatment served as controls. The RNA-seq analysis determined that

654 and 914 cellular RNAs were significantly ($q < 0.05$) modulated in HCMV-infected and zeocin-

treated cells, respectively, and 87 RNAs were modulated by both treatments (Fig. 3*A*). The 87 co-

regulated genes were further analyzed by Gene Set Enrichment Analysis (GSEA) (58) using the

hallmark gene set of the Molecular Signatures Database (MSigDB) (59) and found to be strongly

associated with multiple gene subsets related to DNA damage/repair and other processes, which, if

deregulated, associate with enhanced cancer progression (60-63) (Fig. 3*B*). As we were interested in

determining the common molecular mechanisms that govern HSATII RNA induction in infected and

zeocin-treated cells, we focused on the role of HCMV- and drug-induced DDR pathways in HSATII

RNA expression and its possible effect on cancer cell biology.

ATM is required for HSATII RNA expression. Different types of DNA damage govern the activation of

DDR pathways through key regulators: ATM, ATR and DNA-PKcs (64-66), and each of these PI3-

kinases, when activated, phosphorylates the histone H2A family member, H2AX, generating γ-H2AX

(67). HCMV infection induces accumulation of γ-H2AX (42, 43) as does zeocin (68). To confirm that

both agents induce this marker in ARPE-19 epithelial cells, untreated, HCMV-infected, or zeocin-

treated (100 μg/mL) cells were fixed after 24 h, stained with antibody recognizing γ-H2AX protein and

representative images were captured. Both HCMV-infected and zeocin-treated cells had elevated γ-

H2AX signals (Fig. 4*A*), however, cells exposed to zeocin were characterized by a higher number of γ-

H2AX-positive cells (65%) than HCMV-infected cells (14%) at 24 h after infection or drug treatment

(Fig. 4*B*).

We next scrutinized possible contributions of ATM, ATR and/or DNA-PKcs to HSATII RNA

accumulation. ARPE-19 cells were transfected with NT siRNA or siRNAs targeting ATM, ATR or

PRKDC (encodes DNA-PKcs) transcripts. 48 h later, the cells were infected with HCMV or treated with

zeocin (100 μg/mL) for 24 h. All siRNAs efficiently reduced their targeted transcripts (75-90% decrease)

(Fig. 5*A*). However, only ATM-specific siRNA treatment led to a significant reduction of HSATII RNA

levels (~70% decrease) in HCMV-infected (Fig. 5*A*, left panel) or zeocin-treated cells (Fig. 5A, right

panel). To directly assess the importance of ATM kinase activity (66, 69, 70) in HSATII RNA induction

the effect of Ku-55933 or AZ31, ATM kinase inhibitors (71, 72), were tested. Neither drug compromised

the viability of ARPE-19 cells when tested at concentrations up to 50 μM (Fig. 5B, left panel). However,

because we noticed morphological changes in treated cells at the highest dose (50 μM), we used Ku-

55933 and AZ31 at 20 μM, which is in the range of concentrations at which these drugs are typically

used (71-73). Cells were pretreated with Ku-55933 or AZ31 for 2 h before being treated with zeocin

(100 μg/mL). Interestingly, at 24 hpt neither ATM kinase inhibitor affected HSATII RNA levels (Fig. 5*B*,

right panel). However, both kinase inhibitors negatively (~70% decrease) affected expression of

CDKN1A and DINO transcripts when compared to DMSO-treated control cells, demonstrating that the

drugs were active.

To further establish a role for ATM in HSATII RNA expression, we used ataxia-telangiectasia

fibroblasts, A-T(-) cells, that carry a missense mutation in the carboxy-terminal PI3-kinase-like domain

of ATM, and contain very low levels of mutant protein (74, 75). A derivative of this cell line, A-T(+) cells,

with a stably expressed, functional ATM gene (75) was used as a control. Cells were HCMV-infected or

treated with zeocin (100 μg/mL). Mock infection and solvent treatment served as controls. After 24 h,

infected and drug-treated A-T(+) cells were characterized by a 71x or 137x induction of HSATII RNA,

respectively; and zeocin-treated, but not HCMV-infected, cells contained higher levels of CDKN1A and

DINO RNA. HSATII RNA was also induced in A-T(-) cells following HCMV infection or zeocin treatment,

but to an ~6-fold lower level when compared A-T(+) cells (Fig. 5*C*). Both treatments resulted in

significantly lower expression of CDKN1A and DINO RNAs (~15x and ~5x lower, respectively) in A-T(-)

cells when compared to A-T(+) cells. Together these data further implicate ATM in HSATII RNA

expression following infection or treatment with DNA-damaging drugs.

Next, to confirm a kinase-independent role for ATM in HSATII RNA expression, we ectopically

expressed wild-type ATM (ATM*wt*), kinase-dead ATM (ATM*mut*) or GFP, as a control, in the A-T(-)

fibroblasts and followed with zeocin treatment. As originally demonstrated by Ziv *et al.* A-T(-) fibroblasts

are characterized by a very low level of ATM protein (75) and transfection of the cells with ATM-

encoding plasmids resulted in the efficient expression of ATM*wt* or ATM*mut* (Fig. 5D, left panel).

ATM*wt*- but not ATM*mut*-expressing cells responded to zeocin treatment with enhanced

phosphorylation of Chk2 at threonine 68, as predicted. HSATII RNA levels were monitored in these

cells. Similarly to the data presented in figure 5*C*, zeocin treatment increased HSATII expression in

untransfected or GFP A-T(-) cells (Fig. 5*D*, right panel). Importantly, however, both ATM*wt*- and

ATM*mut*-expressing A-T(-) cells responded to zeocin treatment with similarly increased HSATII RNA

expression, when compared to untransfected or GFP-expressing cells. This experiment confirms that

ATM regulates HSATII expression, but independently from its kinase activity.

HSATII RNA expression is induced by ATM *via* a Chk1/2- and p53-independent pathway. ATM

and ATR phosphorylate two downstream kinases, checkpoint 1 and 2 (Chk1 and Chk2 encoded by the

CHEK1 and CHEK2 genes, respectively), which ultimately regulate p53 activity in a context-dependent

manner (66, 69, 70). Chk2 is normally activated by ATM-mediated phosphorylation, so we tested

whether it is required for accumulation of HSATII RNA. As a control, we also tested Chk1. 48 h prior to

HCMV infection or treatment with zeocin (100 μg/mL), ARPE-19 cells were transfected with NT siRNA

or siRNA specifically targeting CHEK1 or CHEK2 transcripts (encoding Chk1 or Chk2, respectively).

The treatment resulted in a robust decrease in these transcripts at 24 hpi or 24 hpt (Fig. 6*A*). However,

we did not detect a statistically significant decrease of HSATII RNA levels in CHEK1 or CHEK2 KD

cells upon HCMV infection or zeocin treatment, consistent with the lack of a requirement for ATM

kinase activity (Fig. 5), and further arguing that the canonical ATM-Chk2 signaling cascade does not

play an important role in the expression of HSATII RNA.

Since ATM (76) and Chk2 (77) both phosphorylate and stabilize/activate p53, we also investigated the

possible involvement of p53 in HSATII RNA induction. Control and p53 KD ARPE-19 cells were HCMV-

infected or treated with zeocin (100 μg/mL), and HSATII, CDKN1A and DINO RNA expression were

assessed at 24 (Fig. 6*B*) and 120 hpi/hpt (Fig. 6*C*). TP53-specific siRNA strongly decreased the

targeted p53 transcript at both times tested (~83% and 87%), when compared to NT siRNA-treated

cells. p53 KD had a modest effect (10% decrease) on HSATII RNA expression at 120 hpi when

compared to control cells, but zeocin treatment either had no effect (24 hpt) or slightly increased

HSATII RNA expression (120 hpt). In control experiments, p53 KD had little effect on CDKN1A and

DINO expression at 24 h post HCMV infection, but significantly decreased (~65%) both transcripts at

120 hpi. In comparison, p53 KD strongly (80% decreased) affected CDKN1A and DINO expression at

both 24 and 120 h post zeocin treatment.

In sum, ATM protein is required for HSATII RNA accumulation in response to infection or zeocin

treatment. However, ATM kinase activity and two downstream elements of the DDR response that are

normally activated by ATM-mediated phosphorylation, Chk2 and p53, are not required for HSATII RNA

expression.

Breast cancer cells differ in HSATII RNA levels and sensitivity to HCMV infection and zeocin treatments. HSATII RNA expression has been detected in several cancers, including breast (10, 12), and HSATII copy number gains are a common and negative prognostic feature of colorectal tumors (17). Given the induction of HSATII RNA by HCMV infection (41) and DNA-damaging agents (Fig. 1), and knowing that deregulated DDRs allow cancer cells to achieve high proliferation rates while making them more susceptible to DNA-damaging agents (53, 78), we explored the consequences of induced HSATII RNA expression in breast cancer cells. Initially, we assessed HSATII RNA expression levels in breast cancer cell lines that differed by p53 status, hormonal dependency and invasiveness (Fig. 7*A*). We tested less invasive tumor cell lines: MDA-MB-175VII (ER+, PR-, HER2-, p53 wt), MDA-MB-361 (ER+, PR+/-, HER2-, p53 wt), MCF7 (ER+, PR+, HER2-, p53 wt); more invasive triple negative breast cancer (TNBC) cell lines with mutated p53: SUM1315M02, MDA-MB-231, BT-549; and "normal" MCF-10A breast epithelial cells as well as "normal" diploid ARPE-19 cells (79). Cells were cultured in an adherent, 2-dimensional format using identical media conditions across all lines to prevent variations in results potentially caused by different culture conditions. Upon reaching 80-90% confluency, RNA samples were collected. The basal levels of HSATII RNA in the three less invasive cell lines (MCF-7, MDA-MB-175VII and MDA-MB-361) were 5-6-fold higher when compared to HSATII expression in ARPE-19 cells; and the three more invasive cell lines (SUM1315M02, MDA-MB-231, BT-549) were characterized with 14- to 53-fold higher levels of HSATII RNA compared to less invasive breast cancer cells or ARPE-19 cells. The "normal" MCF-10A breast epithelial cells had HSATII RNA levels similar to the less invasive group of tumor cells, suggesting that the basal level of HSATII RNA might be influenced by the tissue origin of tested cells.

Next, we tested the induction of HSATII RNA in a selection of the breast cancer cell lines in response to HCMV infection (Fig. 7*B*). At 48 hpi, all tested cells, except BT-549 cells, demonstrated significantly elevated but varied levels of HSATII RNA (from ~5-fold increase in MCF-7 and MDA-MB-231 to ~100-fold increase in MDA-MB-361 and SUM1315MO$_2$) compared to its levels in uninfected cells. As we previously established that HCMV IE1 and IE2 proteins collaborate to induce HSATII RNA (41) and knowing that HCMV replication is limited or abortive in cancer cells (56, 80, 81), we monitored expression of UL123 and UL122 viral transcripts, encoding the IE1 and IE2 proteins. UL123 and UL122 RNA expression was detected in the breast cancer cell lines upon HCMV infection (Fig. 7B). However, our data did not detect any correlation between levels of HSATII RNA and viral transcripts.

We also tested the effect of zeocin treatment on HSATII RNA induction in these breast cancer cell lines (Fig. 7*C*). At 24 and 96 hpt, all cell lines responded to zeocin with significantly elevated levels of HSATII RNA compared to its levels in solvent control-treated cells. Additionally, a longer exposure (96 h) of cells to the DNA-damaging drug correlated with even higher expression of HSATII RNA. At 96 hpt

HSATII RNA levels increased by as much as 240-fold in the case of MCF-10A cells.

DNA damage-induced HSATII RNA enhances motility and proliferation of breast cancer cells.

We previously reported that HSATII RNA KD reduced the motility of HCMV-infected ARPE-19 cells

(41). Since HCMV infection and zeocin treatment elevated HSATII RNA expression in breast cancer

cell lines (Fig. 7*B* and *C*), we tested whether induced HSATII RNA influences migration of these cells.

Infection of breast cancer cells at a high multiplicity with HCMV leads to viral protein expression in only

a portion of the cells (56), and this would confound interpretation of the migration assay. Therefore, we

treated ARPE-19, MCF-7 and SUM1315MO2 cells with zeocin (200 μg/mL) or H_2O as a solvent control

to guarantee a uniform effect across the cell population (Fig. 8*A*). At 24 hpt, transwell migration assays

(82) were performed using equal numbers of cells and cell motility was assessed after an additional 24

h. ARPE-19 and MCF-7 cells showed an ~ 70% and 100% increase in migration, respectively, upon

zeocin treatment when compared to solvent control-treated cells. Interestingly, this phenomenon was

not observed with SUM1315MO2 cells. Indeed, zeocin caused a slight decrease in the motility of these

cells. Although additional cells must be assayed to reach a firm conclusion, this result raises the

possibility that cells with high basal HSATII RNA levels might be less sensitive to zeocin treatment.

As MDA-MB-231, BT-549 and SUM1315MO2 cells are highly metastatic (83, 84), we tested effects of

HSATII RNA KD on the migratory abilities of these three TNBC cells using transwell migration and

wound healing (85) assays. For the transwell migration assay, MDA-MB-231, BT-549 and

SUM1315MO2 cells were transfected with either NT-LNA or HSATII-LNAs and motility was assessed

24 h later (Fig. 8*B*). HSATII-deficient MDA-MB-231, BT-549 and SUM1315 cells exhibited significantly

reduced migration relative to control-treated cells. MDA-MB-231 and BT-549 cells were the most

sensitive to the treatment with HSATII-LNAs, exhibiting 20- and 5-fold decreases in mean migration,

respectively. The wound healing assay provided consistent results (Fig. 8*C*). HSATII-deficient MDA-

MB-231, BT-549 and SUM1315 cells were significantly slower in closing wounds compared to control

treated cells.

As the aggressiveness of TNBC cells is also related to their high proliferation rates, we tested the

effects of HSATII KD on the proliferation of MDA-MB-231, BT-549 and SUM1315MO2 cells (Fig. 8*D*).

MDA-MB-231, BT-549 and SUM1315MO2 cells were transfected with either NT-LNA or HSATII-LNAs

and 24 h later equal numbers of cells were seeded in 2-dimensional cultures and cell proliferation

monitored for 5 consecutive days. Although the cell lines proliferated at different rates, the growth rate

was significantly reduced for each cell line following HSATII KD.

Our data suggest that HSATII RNA is an important regulator linking a DDR with processes involved in

cancer cell migration and proliferation. Further, at least some DNA-damaging cancer therapeutics,

through the induction of HSATII RNA, might enhance the migration and proliferation of tumor cells.

DISCUSSION

Given the ability of HCMV IE1 and 2 proteins to induce a DDR (42, 43) and HSATII RNA (41), and knowing that DNA damage-inducing treatments stimulate the expression of SINEs (54, 55), we tested the possibility that a chemically-induced DDR can also induce HSATII RNA. Zeocin and etoposide robustly induced HSATII RNA in diploid ARPE-19 epithelial cells (Figs. 1*B*) with kinetics (Figs. 1*C*) similar to those described previously for the induction of the transcript by HCMV infection (41). In contrast, UV-C did not induce HSATII (Fig. 1*A*). UV-C causes helix-distorting lesions in DNA, such as pyrimidine(6-4)pyrimidone photoproducts and cyclobutane pyrimidine dimers (50), that can subsequently be converted into DSBs during DNA replication; whereas zeocin (86), etoposide (87) and HCMV infection (88) directly induce DSBs. Our UV-C assay employed confluent monolayers of cells, conditions that did not favor cellular DNA replication and induction of DSBs, so it is possible that extensive DNA DSBs were not produced during the course of the experiment.

Zeocin induced expression from a similar set of HSATII loci as did HCMV-infection (Fig. 2*B* and *C*), and a subset of genes induced by zeocin and virus infection overlapped and included key elements of DDR signaling cascades. Infection and zeocin treatment induced γ-H2AX foci in diploid epithelial cells, a marker for DSBs (Fig. 4). The ATM kinase is the primary mediator of the response to DSBs (89), and KD of ATM in diploid epithelial cells (Fig. 5*A* and *B*) blocked HSATII RNA induction by the DNA-damaging drugs and infection. Zeocin, etoposide and virus infection each induce a DDR and HSATII RNA, and these responses depend on ATM. However, although ATM protein is clearly required, two different active ATM kinase inhibitors (Ku-55933 and AZ31) failed to block HSATII RNA induction by zeocin (Fig. 5*B*). Moreover, ectopic expression of wild-type ATM or kinase dead ATM restored the ability of ATM-deficient ataxia-telangiectasia A-T(-) cells to efficiently induce HSATII RNA in response to zeocin treatment (Fig. 5*D*). Further, KD of Chk2, which is activated by ATM-mediated phosphorylation, and KD of p53, which is phosphorylated and activated by both ATM and Chk2, had no effect on HSATII RNA induction by zeocin (Fig. 6*A* and *B*). Thus, our data indicate that ATM kinase activity and the canonical ATM-Chk2-p53 signaling cascade are not required for HSATII RNA accumulation. Different signaling events fit with the observation that the ATM kinase-dependent induction of CDKN1A and DINO RNAs occurs much more rapidly than the ATM kinase-independent induction of HSATII in response to zeocin (Fig. 1*D*) or etoposide (Fig. 1*E*).

Even though its kinase activity is not essential, ATM protein (or mRNA) is clearly required for HSATII RNA induction. There is precedent for ATM kinase-independent function. ATM controls mitophagy through its interaction with the Parkin E3 ubiquitin ligase; ATM acts via the protein-protein interaction, independently of its kinase activity (90). We anticipate that the kinase-independent role of ATM in the induction of HSATII RNA might be driven by a similar mechanism.

Overexpression of HSATII RNA has been reported for lung, ovarian, prostate and osteosarcoma

tumors as well as tumor-derived cell lines (10, 12, 13). Given our current data revealing the importance

of the DDR in HSATII RNA expression and with the known impact of deregulated DDRs and cell cycle

checkpoints on oncogenesis (91, 92), we probed the physiological consequences of HSATII RNA

expression in cultured tumor cells. We tested several breast cancer cell lines growing in adherent, 2-

dimensional culture, and higher endogenous HSATII RNA levels correlated with a highly metastatic

phenotype (Fig. 7*A*), and HSATII levels were substantially increased in most cell lines by HCMV

infection or zeocin treatment (Fig. 7*B* and *C*). Cells characterized by low basal HSATII RNA expression,

i.e. ARPE-19 and MCF-7 cells, showed an enhanced migratory phenotype when exposed to zeocin,

while SUM1315MO2 cells with higher uninduced levels of HSATII RNA did not (Fig. 8*A*). Perhaps the

non-responsive cells had a sufficiently high level of the satellite RNA before induction. HSATII RNA KD

markedly reduced migration (Fig. 8*B* and *C*) and proliferation (Fig. 8*D*) in the three TNBC cell lines

tested, suggesting a direct relationship between high levels of the RNA and cell phenotype. These

observations argue that higher HSAII RNA levels correlate with enhanced proliferation and migration,

characteristics of aggressive, metastatic tumors; and they are consistent with an earlier report showing

that HSATII copy number gains are a negative prognostic feature of colorectal cancers (17).

DNA-damaging cancer therapeutics can lead to drug resistance (53, 78) and raise concerns about the

development of *de novo* primary tumors (93). The induction of HSATII RNA accumulation by DNA-

damaging drugs in cultured cells (Fig 1 and 7) and the correlation between elevated HSATII RNA and

enhanced tumor cell proliferation and migration (Fig. 8) prompts us to speculate that DNA-damaging

chemotherapeutics have potential to enhance malignancy of tumor cells that survive treatment. This

effect could be of long or short duration, depending on how long HSATII RNA remains elevated after

drug treatment. Negative consequences could derive from immediate effects of HSATII RNA

overexpression, as we have measured here, and perhaps overexpression of other non-coding RNAs. In

addition, longer-term effects might result from reverse transcription of the overexpressed RNAs

followed by integration and copy number expansion in tumors, as described for HSATII (17) and

another repeated element, LINE-1 (94).

HCMV RNAs and proteins are found in a variety of human malignancies, and HCMV-infected cells have

many properties suggesting the virus might influence tumor progression (95). The ability of HCMV

infection, like radiomimetic drugs, to sponsor a DDR (43, 96) with induction of HSATII RNA in healthy

cells (41) (Figs. 1 and 2) and breast cancer cells (Fig. 7*B*) leads to a second speculation. Perhaps

HSATII RNA induction contributes to the apparent oncomodulatory activity of the virus. Indeed,

numerous viruses, many of which are considered tumor viruses, are known to induce a DDR (42); and,

therefore, might also induce HSATII RNA contributing to malignancy. For HCMV, even a limited or

abortive viral replication documented in cancer cells (56, 80, 81) might lead, *via* IE1/IE2 protein-

mediated HSAII RNA induction, to enhanced migratory and proliferative characteristics of cancer cells.

In sum, this study demonstrates that DNA DSBs induce HSATII RNA via a DDR that requires ATM

protein but not its kinase activity, demonstrates a link between expression of the RNA and cellular

growth and migration phenotype, and establishes a new paradigm to study the biological consequences

of HSATII RNA expression - treatment of normal diploid and tumor cells with DNA-damaging agents.

METHODS AND MATERIALS

Cells, viruses and drugs. Human retinal pigment epithelial (ARPE-19) cells, human foreskin

fibroblasts (HFFs) (30) and breast epithelial (MCF-10A) cells as well as breast cancer cell lines: MCF-7,

MDA-MB-175VII, MDA-MB-361, and MDA-MB-231 were from the American Type Culture Collection

(ATCC). SUM1315MO2 breast cancer cells (97) were generously provided by Stephen Ethier (Medical

University of South Carolina). A-T(-) fibroblasts, originally named AT22IJE-T, and A-T(+) fibroblasts

with a stably expressed functional ATM were created by the lab of Yosef Shiloh (Sackler School of

Medicine) (74, 75) and generously provided by Matthew Weitzman (Perelman School of Medicine

University of Pennsylvania). ARPE-19, MCF-7, MDA-MB-175VII, MDA-MB-361, MDA-MB-231 and

SUM1315MO2 cells were cultured in 10% FBS/DMEM with added Ham's F-12 nutrient mixture (Sigma-

Aldrich) and supplemented with 1x MEM Non-Essential Amino Acids Solution (ThermoFisher

Scientific), 1 mM sodium pyruvate, 1x GlutaMAX and 10 mM Hepes, pH 7.4. HFFs, A-T(-) and A-T(+)

cells were cultured in 10% FBS/DMEM. Media were supplemented with penicillin G sodium salt (100

units/ml) and streptomycin sulfate (95 units/ml). IE1 and IE2-expressing ARPE-19 cells were described

previously (41).

An infectious BAC clone of HCMV strain TB40/E-GFP (TB40/Ewt-GFP) (98) was electroporated into

ARPE-19 cells or HFFs to generate viral progeny that were passaged once more in ARPE-19 cells or

HFFs, respectively. TB40/E-GFP-epi designates TB40/E-GFP virus grown in ARPE-19 cells (99). All

viral stocks were partially purified by centrifugation through a 20% D-sorbitol cushion in buffer

containing 50 mM Tris HCl, 1 mM $MgCl_2$, pH 7.2, resuspended in DMEM and stored in aliquots at -

80^oC. Infections were performed by treating cells with viral inoculum for 2 h, followed by removal of the

inoculum and washing with phosphate-buffered saline (PBS; Sigma-Aldrich) before applying fresh

medium. To quantify extracellular virions, media from infected cells was collected and spun for 10 min.

at 2000 x g to remove debris. To quantify intracellular virions, infected cells were washed with PBS,

scrapped and intracellular virus released by 3 rounds of freezing and thawing and then spun for 10 min.

at 2000 x g to remove debris. Viral stocks and sample supernatant were titered by plaque assay.

Zeocin (InvivoGen) was dissolved in water and stored at -20°C. Etoposide (Millipore Sigma), Ku-55933

(Tocris) and AZ31 (Selleckchem) were dissolved in DMSO and stored at -20°C.

RNA sequence analysis (RNA-seq). RNA was collected in QIAzol Lysis Reagent (Qiagen) and
isolated using the miRNeasy Mini Kit (Qiagen). DNA was removed from samples using Turbo DNase
(Thermo Fisher Scientific) and RNA quality was assessed using the Bioanalyzer 2100 (Agilent
Technologies). Sequencing libraries were prepared using the TruSeq Stranded Total RNA with Ribo-
Zero kit (Illumina), and sequenced on Illumina HiSeq2500 sequencer instrument in paired-end, rapid
mode (2x 150bp).

For RNA-seq analysis, the Galaxy instance of Princeton University was used (100). RNA-Seq data
were de-multiplexed based on indexes. Phred quality scores were checked by using the FastQC toolkit
and were greater than 26 for more than 90% of the read length. Human and HCMV fasta and
annotation (.gtf) files were created for mapping by combining sequences and annotations from Ensembl
annotation, build 38 (GRCh38), Repbase elements (release 19) and TB40 (EF999921) when
appropriate. Quality filtered reads were mapped to the concatenated human-virus genomes using RNA
STAR (57) (Galaxy version 2.6.0b-2). The featureCounts program (101) (Galaxy version 1.6.4) was
used to measure gene expression. The resulting files were used as input to determine differential
expression for each gene utilizing DESeq2 (102) (Galaxy version 2.11.40.6). Fold changes in gene
expression were considered significant when the adjusted *P* value (q value) for multiple testing with the
Benjamini–Hochberg procedure, which controls for the False Discovery Rate (FDR), was <0.05.
To analyze HSATII expression, aligned reads were assigned using the featureCounts function of
Rsubread package (103) with the external HSATII annotation obtained from a RepeatMasker (104)
using GRCh38 and Repbase consensus sequences. This produced the raw read counts. HSATII
expression in terms of CPM (counts per million reads) was computed and normalized across samples
using the trimmed mean of M-values method (TMM) (105). To calculate the percent of HSATII reads
originating from each chromosome, we identified uniquely mapped reads that exclusively overlapped
with HSATII repeat. The number of normalized counts of HSATII reads mapped to each chromosome
was computed and the percentage of these reads mapping to each chromosome was calculated as
described previously (41). To visualize read coverage tracks for alignment files, RNA STAR-generated
BAM files were imported into the Integrative Genomics Viewer (106) (IGV; version 2.6.3) and aligned
with human genome assembly GRCh38 focused on the HSATII locus on chromosome 16. The scale of
read coverage peaks was adjusted for experimentally paired samples. To create Venn diagrams, based
on data from the DESeq2 analysis the list of significantly expressed genes in HCMV-infected or zeocin-
treated cells compared to control cells were imported into a web-based tool InteractiVenn (107).
Gene Set Enrichment Analysis (GSEA) (58) was performed on the list of common genes differentially
expressed in both HCMV-infected and zeocin-treated cells using the hallmark gene set of the Molecular
Signature Database (MSigDB) (59). A matrix of differentially expressed genes from the data set
significantly matching the hallmark gene set of MSigDB was composed and a list of the hallmark gene

subsets was ordered based on a number of overlapping genes, *p* value determining the probability of

association with a given gene set and a FDR q-value.

Quantification of nucleic acids. For analysis by qRT-PCR, RNA was extracted from samples

collected in QIAzol lysis reagent using the miRNeasy kit (Qiagen). cDNA was made from 1 μg of total

RNA with random hexamer primers and MultiScribe reverse transcriptase (Applied Biosystems),

according to the manufacturer's protocol. Transcript-specific primer sequences used in the study were

either previously reported (41) or newly designed: CDKN1A Forward 5'-TTTAACAGCCTGCTCCCTTG-

3', Reverse 5'-AGTTTGCAACCATGCACTTG-3'; DINO Forward 5'-GGAGGCAAAAGTCCTGTGTT-3',

Reverse 5'-GGGCTCAGAGAAGTCTGGTG-3' (49); E2F1 Forward 5'-CCGTGGACTCTTCGGAGAAC-

3', Reverse 5'-ATCCCACCTACGGTCTCCTC-3'; E2F2 Forward 5'-TGGGTAGGCAGGGGAATGTT-3',

Reverse 5'-GCCTTGTCCTCAGTCAGGTG-3'; E2F3a Forward 5'-

TTTAAACCATCTGAGAGGTACTGATGA-3', Reverse 5'-CGGCCCTCCGGCAA-3' (108); E2F3b

Forward 5'-TTTAAACCATCTGAGAGGTACTGATGA-3', Reverse 5'- CCCTTACAGCAGCAGGCAA-3'

(108); E2F4 Forward 5'-TGGAAGGTATCGGGCTAAT-3', Reverse 5'-

CAATCAGTTTGTCAGCAATCTC-3'; E2F5 Forward 5'-CGGCAGATGACTACAACTTTA-3', Reverse 5'-

GATAACAGTCCCAAGTTTCCA-3'; ATM Forward 5'-GCACTGAAAGAGGATCGTAAA-3', Reverse 5'-

GAGGGAACAAAGTCGGAATAC-3'; ATR Forward 5'-ATATCACCCAAAAGGCGTCGT-3', Reverse 5'-

TGCTCTTTTGGTTCATGTCCAC-3'; PRKDC Forward 5'-AAATGGGCCAGAAGATCGCA-3', Reverse

5'-AGGTCCAGGGCTGGAATTTT-3'; CHEK1 Forward 5'-TTTGGACTTCTCTCCAGTAAAC-3', Reverse

5'-GCTGGTATCCCATAAGGAAAG-3'; CHEK2 Forward 5'-CGCGGTCGTGATGTCTCGGG-3',

Reverse 5'-CGCTGCCATGGGGCTGTGAA-3'; TP53 Forward 5'-AGGGATGTTTGGGAGATGTAAG-3',

Reverse 5'-CCTGGTTAGTACGGTGAAGTG-3' (109). qPCR was performed using SYBR Green master

mix (Applied Biosystems) on a QuantStudio™ 6 Flex Real-Time PCR System (Applied Biosystems).

Transcript levels were analyzed using the delta delta Ct method and GAPDH was used as an internal

control (110). Error ranges are reported as standard deviation of the mean (SD).

Analysis of proteins. For Western blot analysis, transfected A-T(-) were harvested using lysis buffer

(50 mM Tris-HCl, pH 7.5, 5 mM EDTA, 100 mM NaCl, 1% Triton X-100, 0.1% SDS, and 10% glycerol).

Samples were mixed with 6xSDS sample buffer (325 mM Tris pH 6.8, 6% SDS, 48% glycerol, 0.03%

bromophenol blue containing 9% 2-mercaptoethanol). Proteins were separated by electrophoresis

(SDS-PAGE) and transferred to ImmunoBlot polyvinylidene difluoride (PVDF) membranes (BioRad

Laboratories). Western blot analyses were performed using primary antibodies recognizing ATM

(D2E2; Cell Signaling Technology), GFP (D5.1; Cell Signaling Technology), p-Chk2 (T68) (2661; Cell

Signaling), Chk2 (2662; Cell Signaling) and β-actin-HRP (Abcam). Donkey anti-rabbit (GE Healthcare

Biosciences) conjugated with horseradish peroxidase was used as a secondary antibody. Western

blots were developed using WesternSure ECL Detection Reagents (Licor).

Immunofluorescence. For immunofluorescence assays, ARPE-19 cells grown in 96-well plates were infected with TB40/E-GFP-epi (3 IU/cell), treated with zeocin (200 μg/mL) or left untreated. After 24 h, cells were fixed in 100% methanol (Sigma) and incubated in blocking solution [PBS with 3% bovine serum albumin prior to the addition of a primary mouse monoclonal antibody specific for phospho-Histone H2AX (Ser13; clone JBW301; Millipore Sigma). Following incubation, cells were washed with PBS, and a donkey Alexa488-conjugated anti-mouse secondary antibody (Santa Cruz Biotechnology) was applied. Nuclei were counterstained with the Hoechst stain. Cells were visualized and % of γ-H2AX-positive cells was calculated using the Operetta high-content imaging and analysis system (Perkin Elmer) with 20X objective.

Plasmid transfection. Plasmids used in the study: pCMV-Neo-Bam was a gift from Bert Vogelstein (Addgene plasmid #16440). pCMV-Neo-Bam E2F3a and pCMV-Neo-Bam E2F3b were gifts from Jacqueline Lees (Addgene plasmid #37970 and #37970, respectively). pCMVHA E2F1 and pCMVHA E2F2 were gifts from Kristian Helin (Addgene plasmids #24225 and #24226, respectively). ARPE-19 cells at 70-80% confluency were transfected with 1 μg of each plasmid using Lipofectamine 3000 (Thermo Fischer Scientific) according to the manufacturer's instructions. 72 h later, plasmid-transfected cells were harvested using TRIzol and samples were stored at -80°C.

Cell stress induction. For the induction of stress by serum deprivation (111), cells were washed with 1xPBS, fed with medium lacking FBS and cultured for 24 h. For the oxidative stress (112), cells were washed with 1xPBS, fed with medium lacking FBS, but containing 100 μM of H_2O_2 and cultured for 24 h. For UV-irradiation-based stress (113), cells were washed with 1xPBS, exposed to the ultra violet light (50 J/m^2) using a Stratalinker (Stratagene), washed with 1xPBS, supplemented with 10%FBS/DMEM and cultured for 24 h.

Locked nucleic acid and siRNA KD. Locked nucleic acid (LNA™; Exiqon, Skelstedt, Denmark) oligonucleotides targeting HSATII RNA were described previously (41). Control NT-LNA or HSATII-LNAs (100-200 nM) were transfected into cells using Lipofectamine RNAiMAX® Reagent (Thermo Fisher Scientific) according to the manufacturer's instructions. LNA-treated ARPE-19 cells were incubated for 24-48 h before being subjected to subsequent experimental procedures.

siRNAs targeting E2F1 (E2F1 siRNA), E2F2 (E2F2 siRNA), E2F3 (E2F3 siRNA), E2F4 (E2F4 siRNA), E2F5 (E2F5 siRNA), CHEK1 (CHEK1 siRNA) and CHEK2 (CHEK2 siRNA) were siGENOME SMARTpools (Dharmacon). siRNAs targeting ATM (ATM siRNA), ATR (ATR siRNA), PRKDC (PRKDC siRNA) and TP53 (TP53 siRNA) were ON-TARGETplus SMARTpools (Dharmacon). All siRNAs and appropriate NT siRNAs (siGENOME Non-targeting siRNA Pool #1 and ON-TARGETplus Non-targeting Control Pool; Dharmacon) were purchased from Life Technologies. ARPE-19 cells were grown to ~80% confluence and then transfected with siRNA using Lipofectamine®

RNAiMAX Reagent (Life Technologies). Cells transfected with non-specific, scrambled siRNA (NT

siRNA) (Life Technologies) served as a negative control. Following 48 h incubation with siRNA, cells

were washed with 1xPBS and subjected to subsequent experimental procedures.

Cell migration, proliferation and toxicity assays. Wound healing and transwell migration assays

were performed as described previously (41). For wound healing assays, confluent monolayers of NT-

LNA- or HSATII-LNA-transfected MDA-MB-231, BT-549 and SUM1315MO2 cells were scratched with a

1-ml pipet to create wounds. The process of wound closure was monitored in time using a Nikon

Eclipse TE2000-U inverted microscope. The average wound width (in arbitrary units) of ARPE-19 cells

was calculated from 6 measurements for each experimental arm from the captured images using

ImageJ software (114). Results are plotted as a mean percent of remaining wound width (SD). For

transwell assays, NT-LNA- or HSATII-LNA -transfected MDA-MB-231, BT-549 and SUM1315MO2 or

zeocin-treated ARPE-19, MCF-7 and SUM1315MO2 cells were trypsinized and 50,000 cells were

seeded onto each filter in FBS-free medium containing ITS Liquid Media Supplement (Sigma-Aldrich).

After 24 h, filters were washed, cells were fixed and migrated cells on the bottom surface of the filter

were stained with 0.2% crystal violet solution. Migrated cells were imaged and quantified using ImageJ

software (114).

To monitor proliferation, cells were transfected with NT-LNA or HSATII-LNAs. After 48 hpt, equal

number of cells (2500 cells/well) were seeded into 96-well plates and cultured in 10% FBS/DMEM with

added Ham's F-12 nutrient mixture and supplemented with 1x MEM Non-Essential Amino Acids

Solution, 1 mM sodium pyruvate, 1x GlutaMAX and 10 mM Hepes. Cell numbers were quantified using

a colorimetric 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-

tetrazolium (MTS)-based CellTiter 96® AQ$_{ueous}$ One Solution Cell Proliferation Assay (Promega,

Madison, WI) according to the manufacturer's protocol. Absorbance was measured at 490 nm using the

SpectraMax Plus 384 Microplate reader (Molecular Devices).

To assay toxicity, ARPE-19 cells were treated with zeocin, etoposide, Ku-55933 or AZ31 at specified

concentrations for 24 h or with water or DMSO as solvent controls. Each tested concentration of a

specific drug contained the same concentration of solvent. At indicated times, the CellTiter 96® AQ$_{ueous}$

One Solution Cell Proliferation Assay (Promega) was performed according to the manufacturer's

instructions. Absorbance was measured at 490 nm using the SpectraMax Plus 384 Microplate reader

(Molecular Devices).

Statistical analysis. To determine statistical significance between two conditions in experiments,

unpaired, two-tailed *t*-tests with Welch`s correction were performed, otherwise one-way ANOVA was

performed between the arrays of data from distinct samples to determine *P* values. *P* value <0.05 was

considered significant. Significance is shown by the presence of asterisks above data points with one,

two, three or four asterisks representing *P*<0.05, *P*<0.01, *P*<0.001 or P<0.0001, respectively. Only

575 significant P values are reported.

576 **Data availability.** Raw RNA-seq data will be available from the NCBI Gene Expression Omnibus

577 (GEO). All relevant experimental data are available from the authors.

ACKNOWLEDGMENTS

We thank Stephen Ethier (Medical University of South Carolina) for generously providing
SUM1315MO2 cells; Matthew Weitzman (Perelman School of Medicine, University of Pennsylvania) for
generously providing A-T(-) and A-T(+) fibroblasts, Adam Oberstein (University of Illinois) for creating
IE1/IE2-expressing ARPE-19 cells, Alexander Solovyov (Memorial Sloan Kettering Cancer Center) for
advice and assistance with RNA-seq analyses and members of the T.S. laboratory for scientific
discussions.

This work was supported by National Institutes of Health Grants AI112951 and AI142520. M.T.N. was
partially supported by American Cancer Society Fellowship PF-14-116-01-MPC.

AUTHOR CONTRIBUTIONS

M.T.N. designed and performed experiments, analyzed the data and wrote the manuscript. T.S.
supervised the study, analyzed the data and edited the manuscript.

DECLARATION OF INTEREST

Icahn School of Medicine at Mt. Sinai and Princeton University have submitted a provisional patent
application based on the use of HSATII-specific LNAs for HSATII knockdown, and M.T.N. and T.S. are
co-inventors.

REFERENCES

1. A. J. Levine, D. T. Ting, B. D. Greenbaum, P53 and the defenses against genome instability caused by transposons and repetitive elements. *Bioessays* **38**, 508-513 (2016).
2. Z. Pezer, J. Brajkovic, I. Feliciello, D. Ugarkovc, Satellite DNA-mediated effects on genome regulation. *Genome Dyn.* **7**, 153-169 (2012).
3. A. Eymery *et al.*, A transcriptomic analysis of human centromeric and pericentric sequences in normal and tumor cells. *Nucleic Acids Res.* **37**, 6340-6354 (2009).
4. L. E. Hall, S. E. Mitchell, R. J. O'Neill, Pericentric and centromeric transcription: a perfect balance required. *Chromosome Res.* **20**, 535-546 (2012).
5. M. A. Garrido-Ramos, Satellite DNA: An Evolving Topic. *Genes (Basel)* **8** (2017).
6. S. Louzada *et al.*, Decoding the Role of Satellite DNA in Genome Architecture and Plasticity-An Evolutionary and Clinical Affair. *Genes (Basel)* **11** (2020).
7. K. Gronbaek, C. Hother, P. A. Jones, Epigenetic changes in cancer. *APMIS* **115**, 1039-1059 (2007).
8. R. Kanwal, S. Gupta, Epigenetic modifications in cancer. *Clin. Genet.* **81**, 303-311 (2012).
9. A. Sanchez Calle, Y. Kawamura, Y. Yamamoto, F. Takeshita, T. Ochiya, Emerging roles of long non-coding RNA in cancer. *Cancer Sci.* **109**, 2093-2100 (2018).
10. D. T. Ting *et al.*, Aberrant overexpression of satellite repeats in pancreatic and other epithelial cancers. *Science* **331**, 593-596 (2011).
11. K. I. Leonova *et al.*, p53 cooperates with DNA methylation and a suicidal interferon response to maintain epigenetic silencing of repeats and noncoding RNAs. *Proc. Natl. Acad. Sci. U.S.A.* **110**, E89-98 (2013).
12. L. L. Hall *et al.*, Demethylated HSATII DNA and HSATII RNA Foci Sequester PRC1 and MeCP2 into Cancer-Specific Nuclear Bodies. *Cell Rep.* **18**, 2943-2956 (2017).
13. X. D. Ho *et al.*, Analysis of the Expression of Repetitive DNA Elements in Osteosarcoma. *Front. Genet.* **8**, 193 (2017).
14. R. Valgardsdottir *et al.*, Transcription of Satellite III non-coding RNAs is a general stress response in human cells. *Nucleic Acids Res.* **36**, 423-434 (2008).
15. H. Bouzinba-Segard, A. Guais, C. Francastel, Accumulation of small murine minor satellite transcripts leads to impaired centromeric architecture and function. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 8709-8714 (2006).
16. M. De Cecco *et al.*, Genomes of replicatively senescent cells undergo global epigenetic changes leading to gene silencing and activation of transposable elements. *Aging Cell* **12**, 247-256 (2013).
17. F. Bersani *et al.*, Pericentromeric satellite repeat expansions through RNA-derived DNA intermediates in cancer. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 15148-15153 (2015).
18. A. Tanne *et al.*, Distinguishing the immunostimulatory properties of noncoding RNAs expressed in cancer cells. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 15154-15159 (2015).
19. N. A. Krump, J. You, Molecular mechanisms of viral oncogenesis in humans. *Nat. Rev. Microbiol.* **16**, 684-698 (2018).
20. E. A. Mesri, M. A. Feitelson, K. Munger, Human viral oncogenesis: a cancer hallmarks analysis. *Cell Host Microbe* **15**, 266-282 (2014).
21. M. E. McLaughlin-Drubin, K. Munger, Viruses associated with human cancer. *Biochim. Biophys. Acta* **1782**, 127-150 (2008).
22. M. K. White, J. S. Pagano, K. Khalili, Viruses and human cancers: a long road of discovery of molecular paradigms. *Clin. Microbiol. Rev.* **27**, 463-481 (2014).
23. I. G. Sia, R. Patel, New strategies for prevention and therapy of cytomegalovirus infection and disease in solid-organ transplant recipients. *Clin. Microbiol. Rev.* **13**, 83-121, table of contents (2000).
24. S. S. Kanj, A. I. Sharara, P. A. Clavien, J. D. Hamilton, Cytomegalovirus infection following liver transplantation: review of the literature. *Clin. Infect. Dis.* **22**, 537-549 (1996).

25. G. Gerna *et al.*, Human cytomegalovirus end-organ disease is associated with high or low
systemic viral load in preemptively treated solid-organ transplant recipients. *New Microbiol.* **35**,
279-287 (2012).
26. T. M. Lanzieri, S. C. Dollard, S. R. Bialek, S. D. Grosse, Systematic review of the birth
prevalence of congenital cytomegalovirus infection in developing countries. *Int. J. Infect. Dis.* **22**,
44-48 (2014).
27. M. Michaelis, H. W. Doerr, J. Cinatl, The story of human cytomegalovirus and cancer:
increasing evidence and open questions. *Neoplasia* **11**, 1-9 (2009).
28. G. Herbein, The Human Cytomegalovirus, from Oncomodulation to Oncogenesis. *Viruses* **10**
(2018).
29. Q. Lepiller, K. Aziz Khan, V. Di Martino, G. Herbein, Cytomegalovirus and tumors: two players
for one goal-immune escape. *Open Virol. J* **5**, 60-69 (2011).
30. L. Vastag, E. Koyuncu, S. L. Grady, T. E. Shenk, J. D. Rabinowitz, Divergent effects of human
cytomegalovirus and herpes simplex virus-1 on cellular metabolism. *PLoS Pathog.* **7**, e1002124
(2011).
31. P. Kumari *et al.*, Essential role of HCMV deubiquitinase in promoting oncogenesis by targeting
anti-viral innate immune signaling pathways. *Cell Death Dis.* **8**, e3078 (2017).
32. D. Collins-McMillen *et al.*, Human Cytomegalovirus Promotes Survival of Infected Monocytes via
a Distinct Temporal Regulation of Cellular Bcl-2 Family Proteins. *J. Virol.* **90**, 2356-2371 (2015).
33. L. Soroceanu, C. S. Cobbs, Is HCMV a tumor promoter? *Virus Res.* **157**, 193-203 (2011).
34. B. Bhattacharjee, N. Renzette, T. F. Kowalik, Genetic analysis of cytomegalovirus in malignant
gliomas. *J. Virol.* **86**, 6815-6824 (2012).
35. M. El-Shinawi *et al.*, Human cytomegalovirus infection enhances NF-kappaB/p65 signaling in
inflammatory breast cancer patients. *PLoS One* **8**, e55755 (2013).
36. M. Michaelis *et al.*, Selection of a highly invasive neuroblastoma cell population through long-
term human cytomegalovirus infection. *Oncogenesis* **1**, e10 (2012).
37. P. Ranganathan, P. A. Clark, J. S. Kuo, M. S. Salamat, R. F. Kalejta, Significant association of
multiple human cytomegalovirus genomic Loci with glioblastoma multiforme samples. *J. Virol.*
86, 854-864 (2012).
38. L. Harkins *et al.*, Specific localisation of human cytomegalovirus nucleic acids and proteins in
human colorectal cancer. *Lancet* **360**, 1557-1563 (2002).
39. L. E. Harkins *et al.*, Detection of human cytomegalovirus in normal and neoplastic breast
epithelium. *Herpesviridae* **1**, 8 (2010).
40. J. Cinatl, M. Scholz, R. Kotchetkov, J. U. Vogel, H. W. Doerr, Molecular mechanisms of the
modulatory effects of HCMV infection in tumor cell biology. *Trends Mol. Med.* **10**, 19-23 (2004).
41. M. T. Nogalski *et al.*, A tumor-specific endogenous repetitive element is induced by
herpesviruses. *Nat. Commun.* **10**, 90 (2019).
42. E. Xiaofei, T. F. Kowalik, The DNA damage response induced by infection with human
cytomegalovirus and other viruses. *Viruses* **6**, 2155-2185 (2014).
43. X. E *et al.*, An E2F1-mediated DNA damage response contributes to the replication of human
cytomegalovirus. *PLoS Pathog.* **7**, e1001342 (2011).
44. D. J. Smart *et al.*, Assessment of DNA double-strand breaks and gammaH2AX induced by the
topoisomerase II poisons etoposide and mitoxantrone. *Mutat. Res.* **641**, 43-47 (2008).
45. K. Shimada *et al.*, TORC2 signaling pathway guarantees genome stability in the face of DNA
strand breaks. *Mol. Cell.* **51**, 829-839 (2013).
46. T. Waldman, K. W. Kinzler, B. Vogelstein, p21 is necessary for the p53-mediated G1 arrest in
human cancer cells. *Cancer Res.* **55**, 5187-5190 (1995).
47. M. Valverde *et al.*, Hydrogen Peroxide-Induced DNA Damage and Repair through the
Differentiation of Human Adipose-Derived Mesenchymal Stem Cells. *Stem Cells Int.* **2018**,
1615497 (2018).
48. B. Benedict *et al.*, Loss of p53 suppresses replication-stress-induced DNA breakage in G1/S
checkpoint deficient cells. *Elife* **7** (2018).
49. A. M. Schmitt *et al.*, An inducible long noncoding RNA amplifies DNA damage signaling. *Nat.*
Genet. **48**, 1370-1376 (2016).

50. L. F. Batista, B. Kaina, R. Meneghini, C. F. Menck, How DNA lesions are turned into powerful
killing structures: insights from UV-induced apoptosis. *Mutat. Res.* **681**, 197-208 (2009).
51. W. P. Roos, B. Kaina, DNA damage-induced cell death: from specific DNA lesions to the DNA
damage response and apoptosis. *Cancer Lett.* **332**, 237-248 (2013).
52. S. Liu *et al.*, Distinct roles for DNA-PK, ATM and ATR in RPA phosphorylation and checkpoint
activation in response to replication stress. *Nucleic Acids Res.* **40**, 10780-10794 (2012).
53. D. Woods, J. J. Turchi, Chemotherapy induced DNA damage response: convergence of drugs
and pathways. *Cancer Biol. Ther.* **14**, 379-389 (2013).
54. C. R. Hagan, R. F. Sheffield, C. M. Rudin, Human Alu element retrotransposition induced by
genotoxic stress. *Nat. Genet.* **35**, 219-220 (2003).
55. C. M. Rudin, C. B. Thompson, Transcriptional activation of short interspersed elements by DNA-
damaging agents. *Genes Chromosomes Cancer* **30**, 64-71 (2001).
56. A. Oberstein, T. Shenk, Cellular responses to human cytomegalovirus infection: Induction of a
mesenchymal-to-epithelial transition (MET) phenotype. *Proc. Natl. Acad. Sci. U.S.A.*
10.1073/pnas.1710799114 (2017).
57. A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).
58. A. Subramanian *et al.*, Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 15545-15550
(2005).
59. A. Liberzon *et al.*, The Molecular Signatures Database (MSigDB) hallmark gene set collection.
Cell Syst. **1**, 417-425 (2015).
60. J. M. Ford, M. B. Kastan, "11 - DNA Damage Response Pathways and Cancer" in Abeloff's
Clinical Oncology (Sixth Edition)*,* J. E. Niederhuber, J. O. Armitage, M. B. Kastan, J. H.
Doroshow, J. E. Tepper, Eds. (Content Repository Only!, Philadelphia, 2020), pp. 154-
164.e154.
61. W. Lu, Y. Kang, Epithelial-Mesenchymal Plasticity in Cancer Progression and Metastasis. *Dev.*
Cell **49**, 361-374 (2019).
62. L. C. Kim, R. S. Cook, J. Chen, mTORC1 and mTORC2 in cancer and the tumor
microenvironment. *Oncogene* **36**, 2191-2201 (2017).
63. A. Hanna, L. A. Shevde, Hedgehog signaling: modulation of cancer properies and tumor
mircroenvironment. *Mol. Cancer* **15**, 24 (2016).
64. J. Falck, J. Coates, S. P. Jackson, Conserved modes of recruitment of ATM, ATR and DNA-
PKcs to sites of DNA damage. *Nature* **434**, 605-611 (2005).
65. G. Giglia-Mari, A. Zotter, W. Vermeulen, DNA damage response. *Cold Spring Harb. Perspect.*
Biol. **3**, a000745 (2011).
66. A. N. Blackford, S. P. Jackson, ATM, ATR, and DNA-PK: The Trinity at the Heart of the DNA
Damage Response. *Mol. Cell* **66**, 801-817 (2017).
67. J. Yuan, R. Adamski, J. Chen, Focus on histone variant H2AX: to be or not to be. *FEBS Lett.*
584, 3717-3724 (2010).
68. F. Delacote *et al.*, Chronic exposure to sublethal doses of radiation mimetic Zeocin selects for
clones deficient in homologous recombination. *Mutat. Res.* **615**, 125-133 (2007).
69. J. Smith, L. M. Tho, N. Xu, D. A. Gillespie, The ATM-Chk2 and ATR-Chk1 pathways in DNA
damage signaling and cancer. *Adv. Cancer Res.* **108**, 73-112 (2010).
70. Y. H. Ou, P. H. Chung, T. P. Sun, S. Y. Shieh, p53 C-terminal phosphorylation by CHK1 and
CHK2 participates in the regulation of DNA-damage-induced C-terminal acetylation. *Mol. Biol.*
Cell **16**, 1684-1695 (2005).
71. I. Hickson *et al.*, Identification and characterization of a novel and specific inhibitor of the ataxia-
telangiectasia mutated kinase ATM. *Cancer Res.* **64**, 9152-9159 (2004).
72. J. Greene *et al.*, The novel ATM inhibitor (AZ31) enhances antitumor activity in patient derived
xenografts that are resistant to irinotecan monotherapy. *Oncotarget* **8**, 110904-110913 (2017).
73. Y. Li, D. Q. Yang, The ATM inhibitor KU-55933 suppresses cell proliferation and induces
apoptosis by blocking Akt in cancer cells with overactivated Akt. *Mol. Cancer. Ther.* **9**, 113-125
(2010).

74. Y. Ziv *et al.*, Cellular and molecular characteristics of an immortalized ataxia-telangiectasia
(group AB) cell line. *Cancer Res.* **49**, 2495-2501 (1989).

75. Y. Ziv *et al.*, Recombinant ATM protein complements the cellular A-T phenotype. *Oncogene* **15**,
159-167 (1997).

76. C. E. Canman *et al.*, Activation of the ATM kinase by ionizing radiation and phosphorylation of
p53. *Science* **281**, 1677-1679 (1998).

77. A. Hirao *et al.*, DNA damage-induced activation of p53 by the checkpoint kinase Chk2. *Science*
287, 1824-1827 (2000).

78. K. Cheung-Ong, G. Giaever, C. Nislow, DNA-damaging agents in cancer chemotherapy:
serendipity and chemical biology. *Chem. Biol.* **20**, 648-659 (2013).

79. X. Dai, H. Cheng, Z. Bai, J. Li, Breast Cancer Cell Line Classification and Its Relevance with
Breast Tumor Subtyping. *J. Cancer* **8**, 3131-3141 (2017).

80. S. Xu, X. Schafer, J. Munger, Expression of Oncogenic Alleles Induces Multiple Blocks to
Human Cytomegalovirus Infection. *J. Virol.* **90**, 4346-4356 (2016).

81. Q. Lepiller, W. Abbas, A. Kumar, M. K. Tripathy, G. Herbein, HCMV activates the IL-6-JAK-
STAT3 axis in HepG2 cells and primary human hepatocytes. *PLoS One* **8**, e59591 (2013).

82. C. R. Justus, N. Leffler, M. Ruiz-Echevarria, L. V. Yang, In vitro cell migration and invasion
assays. *J. Vis. Exp.* 10.3791/51046 (2014).

83. Y. L. Liu *et al.*, Assessing metastatic potential of breast cancer cells based on EGFR dynamics.
Sci. Rep. **9**, 3395 (2019).

84. A. V. Taubenberger, V. M. Quent, L. Thibaudeau, J. A. Clements, D. W. Hutmacher, Delineating
breast cancer cell interactions with engineered bone microenvironments. *J. Bone Miner. Res.*
28, 1399-1411 (2013).

85. C. C. Liang, A. Y. Park, J. L. Guan, In vitro scratch assay: a convenient and inexpensive method
for analysis of cell migration in vitro. *Nat. Protoc.* **2**, 329-333 (2007).

86. G. M. Ehrenfeld *et al.*, Copper-dependent cleavage of DNA by bleomycin. *Biochemistry* **26**, 931-
942 (1987).

87. Y. Pommier, E. Leo, H. Zhang, C. Marchand, DNA topoisomerases and their poisoning by
anticancer and antibacterial drugs. *Chem. Biol.* **17**, 421-433 (2010).

88. M. T. Pickering, T. F. Kowalik, Rb inactivation leads to E2F1-mediated DNA double-strand
break accumulation. *Oncogene* **25**, 746-755 (2006).

89. C. J. Bakkenist, M. B. Kastan, DNA damage activates ATM through intermolecular
autophosphorylation and dimer dissociation. *Nature* **421**, 499-506 (2003).

90. A. Sarkar *et al.*, Ataxia telangiectasia mutated interacts with Parkin and induces mitophagy
independent of kinase activity. Evidence from mantle cell lymphoma. *Haematologica*
10.3324/haematol.2019.234385 (2020).

91. P. Bouwman, J. Jonkers, The effects of deregulated DNA damage signalling on cancer
chemotherapy response and resistance. *Nat. Rev. Cancer* **12**, 587-598 (2012).

92. F. G. Giancotti, Deregulation of cell signaling in cancer. *FEBS Lett.* **588**, 2558-2570 (2014).

93. L. Livraghi, J. E. Garber, PARP inhibitors in the management of breast cancer: current data and
future prospects. *BMC Med.* **13**, 188 (2015).

94. B. Rodriguez-Martin *et al.*, Pan-cancer analysis of whole genomes identifies driver
rearrangements promoted by LINE-1 retrotransposition. *Nat. Genet.* **52**, 306-319 (2020).

95. C. Cobbs, Cytomegalovirus is a tumor-associated virus: armed and dangerous. *Curr. Opin.*
Virol. **39**, 49-59 (2019).

96. M. H. Luo, K. Rosenke, K. Czornak, E. A. Fortunato, Human cytomegalovirus disrupts both
ataxia telangiectasia mutated protein (ATM)- and ATM-Rad3-related kinase-mediated DNA
damage responses during lytic infection. *J. Virol.* **81**, 1934-1950 (2007).

97. R. M. Neve *et al.*, A collection of breast cancer cell lines for the study of functionally distinct
cancer subtypes. *Cancer Cell* **10**, 515-527 (2006).

98. E. Murphy *et al.*, Coding potential of laboratory and clinical strains of human cytomegalovirus.
Proc. Natl. Acad. Sci. U.S.A. **100**, 14976-14981 (2003).

99. K. Wu, A. Oberstein, W. Wang, T. Shenk, Role of PDGF receptor-alpha during human
cytomegalovirus entry into fibroblasts. *Proc. Natl. Acad. Sci. U.S.A.* **115**, E9889-E9898 (2018).

100. E. Afgan *et al.*, The Galaxy platform for accessible, reproducible and collaborative biomedical
813   analyses: 2018 update. *Nucleic Acids Res.* **46**, W537-W544 (2018).
814   101. Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose program for assigning
815   sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).
816   102. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-
817   seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).
818   103. Y. Liao, G. K. Smyth, W. Shi, The Subread aligner: fast, accurate and scalable read mapping by
819   seed-and-vote. *Nucleic Acids Res.* **41**, e108 (2013).
820   104. A. F. A. Smit, R. Hubley, P. Green (2013-2015) RepeatMasker Open-4.0.
821   105. M. D. Robinson, A. Oshlack, A scaling normalization method for differential expression analysis
822   of RNA-seq data. *Genome Biol.* **11**, R25 (2010).
823   106. H. Thorvaldsdottir, J. T. Robinson, J. P. Mesirov, Integrative Genomics Viewer (IGV): high-
824   performance genomics data visualization and exploration. *Brief. Bioinform.* **14**, 178-192 (2013).
825   107. H. Heberle, G. V. Meirelles, F. R. da Silva, G. P. Telles, R. Minghim, InteractiVenn: a web-
826   based tool for the analysis of sets through Venn diagrams. *BMC Bioinformatics* **16**, 169 (2015).
827   108. Y. Gao *et al.*, A double-negative feedback loop between E2F3b and miR- 200b regulates
828   docetaxel chemosensitivity of human lung adenocarcinoma cells. *Oncotarget* **7**, 27613-27626
829   (2016).
830   109. H. H. Gari, G. D. DeGala, M. S. Lucia, J. R. Lambert, Loss of the oncogenic phosphatase PRL-3
831   promotes a TNF-R1 feedback loop that mediates triple-negative breast cancer growth.
832   *Oncogenesis* **5**, e255 (2016).
833   110. K. J. Livak, T. D. Schmittgen, Analysis of relative gene expression data using real-time
834   quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402-408 (2001).
835   111. D. Rajapakse, K. Peterson, S. Mishra, G. Wistow, Serum starvation of ARPE-19 changes the
836   cellular distribution of cholesterol and Fibulin3 in patterns reminiscent of age-related macular
837   degeneration. *Exp. Cell. Res.* **361**, 333-341 (2017).
838   112. S. Upadhyay, S. Vaish, M. Dhiman, Hydrogen peroxide-induced oxidative stress and its impact
839   on innate immune responses in lung carcinoma A549 cells. *Mol. Cell. Biochem.* **450**, 135-147
840   (2019).
841   113. S. Nakajima *et al.*, UV light-induced DNA damage and tolerance for the survival of nucleotide
842   excision repair-deficient human cells. *J. Biol. Chem.* **279**, 46674-46677 (2004).
843   114. W. S. Rasband (1997-2010) ImageJ. (U.S. National Institutes of Health, Bethesda, Maryland,
844   USA).
845

FIGURES

Figure 1.

Fig. 1. DNA damage response regulates HSATII RNA expression. (*A*) Limited induction of HSATII RNA by UV, H_2O_2, or serum withdrawal. ARPE-19 cells were exposed to UV-irradiation (50 J/m^2), H_2O_2 (100 nM) or serum withdrawal (no FBS). RNA samples were collected at 24 hpt. RT-qPCR was performed to quantify HSATII, CDKN1A or DINO transcripts, using GAPDH as an internal control. Data are presented as a fold change (mean ± SD; n=3). *$P<0.05$, **$P<0.01$, ***$P<0.001$. (*B* and *C*) Induction of HSATII RNA by treatment with DNA-damaging drugs. ARPE-19 cells were treated with increasing concentrations of zeocin (*B*) or etoposide (*C*) with H_2O or DMSO, respectively, as solvent controls. RNA samples were collected at 24 hpt, and RT-qPCR was performed to quantify HSATII, CDKN1A and DINO transcripts, using GAPDH as an internal control. Data are presented as a fold change (mean ± SD; n=3). Insert: At 24 hpt, cell viability was assessed at each indicated drug concentrations. Data is presented as % viable cells, (mean ± SD; n = 3). (*D* and *E*) Drug-induced HSATII RNA accumulation occurs after a delay. ARPE-19 cells were treated with zeocin (200 μg/mL) (*D*), etoposide (200 μM) (*E*) and H_2O or DMSO, respectively, as solvent controls. RNA samples were collected at indicated times. RT-qPCR was performed to quantify HSATII, CDKN1A and DINO transcripts, using GAPDH as an internal control. Data are presented as a fold change (mean ± SD; n=3).

Figure 2.

Fig. 2. Zeocin treatment partially mimics the transcriptional effect of HCMV infection on HSATII RNA expression. ARPE-19 cells were infected with HCMV (1 TCID50/cell), or treated with zeocin (100 µg/mL) or H_2O as a solvent control, and RNA samples were collected at either 24 hpi or 24 hpt, respectively. RNA was isolated and analyzed using RNA-seq. (*A*) Quantification of HSATII RNA induction by RNA-seq analysis. HSATII RNA expression in terms of counts per million reads mapped (CPM) was computed and normalized across samples (n=2). (*B*) Chromosomal distribution of HSATII RNA expression. HSATII chromosomal origin in HCMV-infected or zeocin-treated cells was depicted based on the number of unique HSATII reads mapped to specific chromosomal loci. Data are presented as a percentage of total HSATII reads (mean ± SD. n=2). (*C*) Example loci of HSATII RNA expression on chromosome 16. HSATII read coverage for HCMV-infection versus zeocin treatment was based on reads mapped to human genome and read coverage track at the HSATII locus on chromosome 16 was visualized using IGV.

937 **Figure 3.**

A

HCMV   Zeocin

567   87   827

B

Gene Set Name	p-value	FDR q-value
HALLMARK_E2F_TARGETS	$2.64 \ e^{-17}$	$1.32 \ e^{-15}$
HALLMARK_G2M_CHECKPOINT	$3.31 \ e^{-11}$	$8.26 \ e^{-10}$
HALLMARK_EPITHELIAL_MESENCHYMAL_TRANSITION	$9.52 \ e^{-5}$	$1.59 \ e^{-3}$
HALLMARK_INTERFERON_GAMMA_RESPONSE	$1.13 \ e^{-3}$	$1.13 \ e^{-2}$
HALLMARK_MTORC1_SIGNALING	$1.13 \ e^{-3}$	$1.13 \ e^{-2}$
HALLMARK_HEDGEHOG_SIGNALING	$3.04 \ e^{-3}$	$2.53 \ e^{-2}$
HALLMARK_DNA_REPAIR	$4.87 \ e^{-3}$	$3.48 \ e^{-2}$
HALLMARK_APOPTOSIS	$5.92 \ e^{-3}$	$3.7 \ e^{-2}$

Fig. 3. Zeocin treatment partially mimics the HCMV-induced transcriptome. ARPE-19 cells were mock- or TB40/E-GFP-epi-infected (1 $TCID_{50}$/cell), or treated with zeocin (100 μg/mL) or H_2O as a solvent control. After 24 h, RNA samples were collected, RNA was isolated and analyzed using RNA-seq. (*A*) Cellular genes modulated by both HCMV and zeocin. Venn diagram showing significantly expressed genes (q<0.05) either in HCMV-infected or zeocin-treated cells vs. respective controls. (*B*) Categories of gene sets modulated by both HCMV and zeocin. GSEA was performed on the genes differentially expressed in both HCMV-infected and zeocin-treated cells using the hallmark gene set of MSigDB. Identified specific gene set names are categorized based on increasing *p*-value and FDR q-value.

Figure 4.

Fig. 4. HCMV and zeocin induce γ-H2AX in ARPE-19 cells. Cells were mock- or TB40/E-GFP-epi-infected (3 TCID50 /cell) or treated with zeocin (200 μg/mL). After 24 h, cells were stained for γ-H2AX, nuclei were counterstained with the Hoechst stain, cells were visualized (*A*: scale bars=100 μm) and % γ-H2AX-positive cells were quantified (*B*: mean ± SD; n=3). ****P<0.0001.

Figure 5.

1035
Fig. 5. ATM-based DNA damage response regulates HSATII RNA expression. (*A*) ATM KD inhibits the induction
of HSATII RNA by HCMV infection or zeocin treatment. ARPE-19 cells were transfected with ATM-, ATR- or
PRKDC-specific siRNAs or NT siRNA as a control. After 48 h, cells were infected with TB40/E-GFP-epi (1
$TCID_{50}$/cell) or zeocin (100 μg/mL). RNA samples were collected 24 h later. RT-qPCR was performed to quantify
HSATII, ATM, ATR and PRKDC transcripts, using GAPDH as an internal control. Data are presented as a fold
change (mean ± SD; n=3-10). (*B*) HSATII RNA is induced less in A-T(-) cells than in A-T(+). A-T(+) and A-T(-)
cells were uninfected, infected with TB40/E-GFP (1 $TCID_{50}$/cell), or treated with zeocin (100 μg/mL) or H_2O as a
solvent control. RNA samples were collected 24 h later. RT-qPCR was performed to quantify HSATII, CDKN1A
and DINO transcripts, using GAPDH as an internal control. Data are presented as a fold change (mean ± SD;
n=3). (*C*) ATM kinase inhibitors do not block the induction of HSATII RNA by zeocin. Left panel: ARPE-19 cells
were treated with various concentrations of Ku-55933 or AZ31. At 24 hpt, cell viability was assessed. Data is
presented as % viable cells, (mean ± SD; n=3). Right panel: ARPE-19 cells were treated with DMSO, Ku-55933
(20 μM) or AZ31 (20 μM) for 2 h before zeocin (100 μg/mL) was added. RNA samples were collected 24 hpt, and
RT-qPCR was performed to quantify HSATII, CDKN1A and DINO transcripts, using GAPDH was used as an
internal control. Data are presented as a fold change (mean ± SD; n=3). (*D*) Ectopic expression of kinase-dead
ATM (ATM*mut*) can enhance the induction of HSATII RNA in A-T(-) cells by zeocin. Left panel: A-T(-) cells were
transfected with GFP-, ATM*wt*- or ATM*mut*-expressing plasmids. After 48 h, cells were exposed to zeocin (100
μg/mL) and protein samples were collected 24 h later. Proteins were assayed by Western blot analysis performed
using antibodies recognizing ATM, GFP, phospho-Chk2 (T68), Chk2 and actin as a loading control. Right panel:
A-T(-) cells were transfected with GFP-, ATM*wt*- or ATM*mut*-expressing plasmids and untransfected cultures
served as controls. After 48 h, cells were treated with zeocin (100 μg/mL) and RNA samples were collected 24 h
later. RT-qPCR was performed to quantify HSATII. GAPDH was used as an internal control. Data are presented
as a fold change (mean ± SD. n=4). *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

Figure 6.

Fig. 6. HSATII RNA expression is induced via a Chk1/2- and p53-independent pathway. (*A*) CHEK1 or CHEK2 KD doesn't block HSATII RNA induction by infection or zeocin. ARPE-19 cells were transfected with CHEK1- or CHEK2-specific siRNA or NT siRNA as a control. After 48 h, cells were infected with TB40/E-GFP-epi (1 $TCID_{50}$/cell) or zeocin (100 μg/mL). RNA samples were collected 24 h later. RT-qPCR was performed to quantify HSATII, CHEK1 and CHEK2 transcripts, using GAPDH as an internal control. Data are presented as a fold change (mean ± SD; n=4). (*B* and *C*) p53 KD does not block HSATII RNA induction following HCMV infection or zeocin treatment. ARPE-19 cells were transfected with TP53 siRNAs or NT siRNA as a control. After 48 h, cells were infected with TB40/E-GFP-epi (1 TCID/cell) or zeocin (100 μg/mL), and RNA samples were collected 24 h (*B*) or 120 h (*C*) later. RT-qPCR was performed using specific primers to HSATII, TP53, CDKN1A or DINO transcripts. GAPDH was used as an internal control. Data are presented as a fold change (mean ± SD; n=3-4). *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$.

Figure 7.

Fig. 7. Breast cancer cells differ in HSATII RNA levels and sensitivity to HCMV infection and zeocin treatment. (*A*) Breast cancer cell lines contain HSATII RNA when grown in adherent (2-dimensional) culture. Cell lines were grown to ~80% confluency and RNA samples were collected. RT-qPCR was performed to quantify HSATII RNA, using GAPDH as an internal control. Data are presented as a fold change (mean ± SD; n=3). (*B*) HCMV infection elevates HSATII RNA levels in breast cancer cell lines. MCF-7, MDA-MB-361, MDA-MB-231, BT-549 and SUM1315MO2 cells were infected with TB40/E-GFP-epi (4 TCID$_{50}$/cell) and RNA samples were collected at 24 hpi. RT-qPCR was performed to quantify HSATII RNA, using GAPDH as an internal control. Data are presented

1154 as a fold change (mean ± SD; n=3-5). (*C*) Zeocin treatment elevates HSATII RNA levels in breast cancer cell
1155 lines. MCF-7, MCF-10A, MDA-MB-361 and SUM1315MO2 cells were treated with zeocin (200 μg/mL). RNA
1156 samples were collected at 24 and 96 hpt and RT-qPCR was performed to quantify HSATII RNA, using GAPDH as
1157 an internal control. Data are presented as a fold change (mean ± SD; n=3). *P<0.05, **P<0.01, ***P<0.001,
1158 ****P<0.0001.

Figure 8.

Fig. 8. DNA damage-induced HSATII RNA enhances motility and proliferation of breast cancer cells. (*A*) Zeocin

treatment increases migration of cells with low basal levels of HSATII RNA. ARPE-1, MCF-7 and SUM315MO2

cells were treated with zeocin (200 μg/mL) for 24 h. Cells were transferred onto transwell inserts, and 24 h later,

migrated cells were washed, fixed and nuclei stained. The graph presents a number of cells (with their indicated

means) that migrated through a transwell per a field of view (FOV) from biological replicates. *n*=11. (*B*) Transwell

assays demonstrated that HSATII KD reduces migration of breast cancer cells. MDA-MB-231, BT-549 and

SUM1315MO2 cells were transfected with NT-LNA or HSATII-LNAs. After 48 hpt, equal number of cells was

transferred onto transwell inserts. 24 h later, migrated cells were washed, fixed and nuclei stained. The graph

presents a number of cells (with their indicated means) migrated through a transwell per FOV from biological

replicates. n=12. (*C*) Wound-healing assays demonstrated that HSATII KD reduces migration of breast cancer

cells. MDA-MB-231, BT-549 and SUM1315MO2 cells were transfected with NT-LNA or HSATII-LNAs. After 48

hpt, wounds were created and their closure was monitored at indicated times. Data from biological replicates are

presented as a percent of remaining wound width mean ± SD. n=20. (*D*) HSATII RNA KD reduces the rate of

breast cancer cell proliferation in 2D culture. MDA-MB-231, BT-549 and SUM1315MO2 cells were transfected

with NT-LNA or HSATII-LNAs. After 48 hpi, equal number of cells were seeded and cell proliferation was

monitored at the indicated times. Cell proliferation is presented as an increase in a measured absorbance (AU –

arbitrary units). n=3. *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$.