**Title:** Neuronal timescales are functionally dynamic and shaped by cortical microarchitecture

**Authors:** Richard Gao[1*], Ruud L. van den Brink[5], Thomas Pfeffer[6], Bradley Voytek[1-4]

**Affiliations:**
[1]Department of Cognitive Science,
[2]Halıcıoğlu Data Science Institute,
[3]Neurosciences Graduate Program,
[4]Kavli Institute for Brain and Mind, University of California, San Diego, La Jolla, CA, USA 92093
[5]Department of Neurophysiology and Pathophysiology, University Medical Center Hamburg-
Eppendorf, Hamburg, Germany
[6]Center for Brain and Cognition, Computational Neuroscience Group, Universitat Pompeu
Fabra, 08018 Barcelona, Spain.

* Correspondence: r.dg.gao@gmail.com

**Abstract**

Complex cognitive functions such as working memory and decision-making require information
maintenance over many timescales, from transient sensory stimuli to long-term contextual cues.
While theoretical accounts predict the emergence of a corresponding hierarchy of neuronal
timescales, direct electrophysiological evidence across the human cortex is lacking. Here, we
infer neuronal timescales from invasive intracranial recordings. Timescales increase along the
principal sensorimotor-to-association axis across the entire human cortex, and scale with single-
unit timescales within macaques. Cortex-wide transcriptomic analysis shows direct alignment
between timescales and expression of excitation- and inhibition-related genes, as well as genes
specific to voltage-gated transmembrane ion transporters. Finally, neuronal timescales are
functionally dynamic: prefrontal cortex timescales expand during working memory maintenance
and predict individual performance, while cortex-wide timescales compress with aging. Thus,
neuronal timescales follow cytoarchitectonic gradients across the human cortex, and are
relevant for cognition in both short- and long-terms, bridging microcircuit physiology with
macroscale dynamics and behavior.

**Introduction**

Human brain regions are broadly specialized for different aspects of behavior and cognition. For example, primary sensory neurons are tightly coupled to changes in the environment, firing rapidly to the onset and removal of a stimulus, and showing characteristically short intrinsic timescales (Ogawa and Komatsu, 2010; Runyan et al., 2017). In contrast, neurons in cortical association (or transmodal) regions, such as the prefrontal cortex (PFC), can sustain their activity for many seconds when a person is engaged in working memory (Zylberberg and Strowbridge, 2017), decision-making (Gold and Shadlen, 2007), and hierarchical reasoning (Sarafyazd and Jazayeri, 2019). This persistent activity in the absence of immediate sensory stimuli reflects longer neuronal timescales, which is thought to result from neural attractor states (Wang, 2002; Wimmer et al., 2014) shaped by NMDA-mediated recurrent excitation and fast feedback inhibition (Wang, 1999, 2008), with contributions from other synaptic and cell-intrinsic properties (Duarte and Morrison, 2019; Gjorgjieva et al., 2016).

Recent studies have shown that variations in many such microarchitectural features follow continuous and coinciding gradients along a sensory-to-association axis across the cortex, including cortical thickness, cell density, and distribution of excitatory and inhibitory neurons (Hilgetag and Goulas, 2020; Huntenburg et al., 2018; Wang, 2020). In particular, grey matter myelination (Glasser and Van Essen, 2011), which indexes anatomical hierarchy, varies with the expression of numerous genes related to microcircuit function, such as NMDA receptor and inhibitory cell-type marker genes (Burt et al., 2018). Functionally, specialization of the human cortex, as well as structural and functional connectivity (Margulies et al., 2016), also follow similar macroscopic gradients. In addition to the broad differentiation between sensory and association cortices, there is evidence for a finer hierarchical organization within the frontal cortex (Sarafyazd and Jazayeri, 2019). For example, the anterior-most parts of the PFC are responsible for long timescale goal-planning behavior (Badre and D'Esposito, 2009; Voytek et al., 2015a), while healthy aging is associated with a shift in these gradients such that older adults become more reliant on higher-level association regions to compensate for altered lower-level cortical functioning (Davis et al., 2008).

Despite convergent observations of continuous cortical gradients in structural features and cognitive specialization, there is no direct evidence for a similar gradient of neuronal timescales across the human cortex. Such a gradient of neuronal dynamics is predicted to be a natural consequence of macroscopic variations in synaptic connectivity and microarchitectural features (Chaudhuri et al., 2015; Duarte et al., 2017; Huntenburg et al., 2018; Wang, 2020), and would be a primary candidate for how functional specialization emerges as a result of hierarchical temporal processing (Kiebel et al., 2008). Single-unit recordings in rodents and non-human primates hint at a hierarchy of timescales that increase, or expand, progressively along a posterior-to-anterior axis (Murray et al., 2014; Runyan et al., 2017; Wasmuht et al., 2018), while intracranial recordings and functional neuroimaging data collected during perceptual and cognitive tasks suggest likewise in humans (Baldassano et al., 2017; Honey et al., 2012; Lerner et al., 2011; Watanabe et al., 2019). However, these data are either sparsely sampled across the cortex or do not measure neuronal activity at the cellular and synaptic level directly, prohibiting the full construction of an electrophysiological timescale gradient across the human

76    cortex. As a result, while whole-cortex data of transcriptomic and anatomical variations exist, we
77    cannot take advantage of them to dissect the contributions of synaptic, cellular, and circuit
78    connectivity in shaping fast neuronal timescales, nor ask whether regional timescales are
79    dynamic and relevant for human cognition (Fig. 1A).
80

**Fig. 1. Schematic of study and timescale inference technique. (A)** in this study, we infer neuronal timescales from intracranial field potential recordings, which reflect integrated synaptic and transmembrane current fluctuations over large neural populations(Buzsáki et al., 2012). Combining multiple open-access datasets (Table S1), we link timescales to known human anatomical hierarchy, dissect its cellular and physiological basis via transcriptomic analysis, and demonstrate its functional modulation during behavior and through aging. **(B)** simulated time series, and their **(C)** autocorrelation functions (ACF), with increasing timescales (decay time constant). **(D)** example ECoG power spectral density (PSD) showing that in frequency domain, timescale is equivalent to the frequency of aperiodic power drop-off ($f_k$, triangle; insets: time series and ACF). **(E)** accurate extraction of timescale parameters from PSDs of simulated time series in **(B)**.

**Results**

**Neuronal timescale can be inferred from frequency domain**
To overcome these limitations, we develop a novel computational method for inferring the timescale of neuronal transmembrane current fluctuations from human intracranial electrocorticography (ECoG) recordings (Fig. 1A, box). Neural time series exhibit variable temporal autocorrelation, or timescales, where future values are partially predictable from past values, and predictability decreases with increasing time lags. To demonstrate the effect of varying autocorrelation, we simulate the aperiodic (non-rhythmic) component of neural field potential recordings by convolving Poisson population spikes with exponentially-decaying synaptic kernels (Fig. 1B). Consistent with previous studies, "neuronal timescale" here is defined as the exponential decay time constant ($\tau$) of the autocorrelation function (ACF) (Murray et al., 2014)—the time it takes for the ACF to decrease by a factor of $e$ (Fig. 1C). Equivalently, we can estimate neuronal timescale from the "characteristic frequency" ($f_k$) of the power spectral density (PSD), especially when the presence of variable 1/f ($\chi$) and oscillatory components can bias timescale inference from the ACF in time-domain (Fig. 1D). In this study, we apply spectral parameterization (Haller et al., 2018) to extract timescales from intracranial recordings, which decomposes neural PSDs into a combination of oscillatory and aperiodic components, where timescale is inferred from the latter. We validate this approach on PSDs computed from simulated neural time series and show that the model-fitted timescales closely match their ground-truth values (Fig. 1E).

**Timescales follow anatomical hierarchy and are 10x faster than spiking timescales**
Applying this technique, we infer a continuous gradient of neuronal timescales across the human cortex and examine its relationship with anatomical hierarchy. We analyze a large dataset of human intracranial (ECoG) recordings of task-free brain activity from 106 epilepsy patients (MNI-iEEG (Frauscher et al., 2018a), see Fig. S1 for electrode coverage), and compare the ECoG-derived timescale gradient to the average T1w/T2w map from the Human Connectome Project, which captures grey matter myelination and indexes the proportion of feedforward vs. feedback connections between cortical regions, defining an anatomical hierarchy (Burt et al., 2018; Glasser and Van Essen, 2011).

Across the human cortex, timescales of fast electrophysiological dynamics (~10-50 ms) predominantly follow a rostrocaudal gradient (Fig. 2A). Consistent with numerous accounts of a principal cortical axis spanning from primary sensory to association regions (Hilgetag and Goulas, 2020; Margulies et al., 2016; Wang, 2020), timescales are shorter in sensorimotor and early visual areas, and longer in association regions, especially posterior parietal, ventral/medial frontal, and medial temporal cortex. Cortical timescales are negatively correlated with T1w/T2w (Fig. 2B, $\rho$ = -0.47, $p$ < 0.001; adjusted for spatial autocorrelation, see Materials and Methods and Fig. S2), such that timescales are shorter in more heavily myelinated (lower-level) cortical regions.

While surface ECoG recordings offer much broader spatial coverage than extracellular single-unit recordings, they are fundamentally different signals: ECoG and field potentials largely reflect

integrated synaptic and other transmembrane currents across many neuronal and glial cells,
rather than putative action potentials from single neurons (Buzsáki et al., 2012) (Fig. 1A, box).
Considering this, we ask whether timescales measured from ECoG are related to single-unit
spiking timescales along the cortical hierarchy. To test this, we extract neuronal timescales from
task-free ECoG recordings in macaques and compare them to a separate dataset of single-unit
spiking timescales (Murray et al., 2014) (Fig. 2C, inset; see Fig. S3 for electrode locations).
Consistent with spiking timescale estimates (Murray et al., 2014; Wasmuht et al., 2018), ECoG
timescales also increase along the macaque cortical hierarchy. While there is a strong
correspondence between spiking and ECoG timescales (Fig. 2C; $\rho = 0.96$, $p < 0.001$)—measured
from independent datasets—across the macaque cortex, ECoG-derived timescales are 10 times
faster than single-unit timescales and are conserved across individual sessions (Fig. 2D). This
suggests that neuronal spiking and transmembrane currents have distinct but related timescales
of fluctuations, and that both are hierarchically organized along the primate cortex.
150

151
**Fig. 2. Timescale increases along the anatomical hierarchy in humans and macaques. (A)**
human cortical timescale gradient (left) falls predominantly along the rostrocaudal axis, similar to
T1w/T2w ratio (right). **(B)** neuronal timescales are negatively correlated with cortical T1w/T2w,
thus increasing along the anatomical hierarchy from sensory to association regions (p-value
adjusted for spatial autocorrelation). **(C)** macaque ECoG timescales track published single-unit
spiking timescales (Murray et al., 2014) in corresponding regions (mean±s.e.m from n=8
sessions); inset: ECoG electrode map of one animal. **(D)** ECoG-derived timescales are
consistently correlated to (left), and an order of magnitude faster than (right), single-unit
timescales across individual sessions. Hollow markers: individual sessions; shapes: animals;
solid circles: grand average from **(C)**.

**Synaptic and ion channel genes shape timescales of neuronal dynamics**

Next, we identify cellular and synaptic mechanisms underlying timescale variations across the human cortex. Theoretical accounts posit that NMDA-mediated recurrent excitation coupled with fast inhibition (Chaudhuri et al., 2015; Wang, 1999, 2008), as well as cell-intrinsic properties (Duarte and Morrison, 2019; Gjorgjieva et al., 2016; Koch et al., 1996), are crucial for shaping neuronal timescales. While *in vitro* and *in vivo* studies in model organisms (van Vugt et al., 2020; Wang et al., 2013) can test these hypotheses at the single-neuron level, causal manipulation and large-scale recording of neuronal networks embedded in the human brain is severely limited. Here, we apply an approach analogous to multimodal single-cell profiling (Bomkamp et al., 2019) and examine the transcriptomic basis of neuronal dynamics at the macroscale.

Leveraging cortex-wide bulk mRNA expression variations (Hawrylycz et al., 2012), we find that the neuronal timescale gradient overlaps with the dominant axis of gene expression across the human cortex ($\rho$ = -0.60, $p$ < 0.001; Fig. 3A and Fig. S4). Consistent with theoretical predictions (Fig. 3B), timescales significantly correlate with the expression of genes encoding for NMDA (GRIN2B) and GABA-A (GABRA3) receptor subunits, voltage-gated sodium (SCN1A) and potassium (KCNA3) ion channel subunits, as well as inhibitory cell-type markers (parvalbumin, PVALB), and genes previously identified to be associated with single-neuron membrane time constants (PRR5) (Bomkamp et al., 2019).

More specifically, *in vitro* electrophysiological studies have shown that, for example, increased expression of receptor subunit 2B extends the NMDA current time course (Flint et al., 1997), while 2A expression shortens it (Monyer et al., 1994). Similarly, the GABA-A receptor time constant lengthens with increasing a3:a1 subunit ratio (Eyre et al., 2012). We show that these relationships are recapitulated at the macroscale, where neuronal timescales positively correlate with GRIN2B and GABRA3 expression, and negatively correlate with GRIN2A and GABRA1. These results demonstrate that timescales of neural dynamics depend on specific receptor subunit combinations with different (de)activation timescales (Duarte et al., 2017; Gjorgjieva et al., 2016), in addition to broad excitation-inhibition interactions (Gao et al., 2017; Wang, 2002, 2020). Notably, almost all genes related to voltage-gated sodium and potassium ion channel alpha-subunits—the main functional subunits—are correlated with timescale, while all inhibitory cell-type markers except parvalbumin have strong positive associations with timescale (Fig. 3C and Fig. S5).

We further test whether single-cell timescale-transcriptomic associations are captured at the macroscale as follows: for a given gene, we can measure how strongly its expression correlates with membrane time constant parameters at the single-cell level using patch-clamp and RNA sequencing (scRNASeq) data (Bomkamp et al., 2019; Tripathy et al., 2017). Analogously, we can measure its macroscopic transcriptomic-timescale correlation using the cortical gradients above. Comparing across these two levels for all previously-identified timescale-related genes (Bomkamp et al., 2019; Tripathy et al., 2017), we find a significant correlation between the

strength of association at the single-cell and macroscale levels (Fig. 3D, horizontal lines; $\rho$ = 0.36 and 0.25, $p$ < 0.001). Furthermore, genes with stronger associations to timescale tend to conserve this relationship across single-cell and macroscale levels (Fig. 3D, separated by macroscale correlation magnitude). Thus, the association between cellular variations in gene expression and cell-intrinsic temporal dynamics is captured at the macroscale, even though scRNAseq and microarray data represent entirely different measurements of gene expression.

While we have shown associations between cortical timescales and genes suspected to influence neuronal dynamics, these data present an opportunity to discover additional novel genes that are functionally related to timescales through a data-driven approach. However, since transcriptomic variation and anatomical hierarchy overlap along a shared macroscopic gradient (Burt et al., 2018; Huntenburg et al., 2018; Margulies et al., 2016), we cannot specify the role certain genes play based on their level of association with timescale alone: gene expression differences across the cortex first result in cell-type and connectivity differences, sculpting the hierarchical organization of cortical anatomy. Consequently, anatomy and cell-intrinsic properties jointly shape neuronal dynamics through connectivity differences (Chaudhuri et al., 2015; Demirtaş et al., 2019) and expression of ion transport proteins with variable activation timescales, respectively. Therefore, we ask whether variation in gene expression still accounts for variation in timescale beyond the principal structural gradient, and if associated genes have known functional roles in biological processes (schematic in Fig. 4E). To do this, we first remove the contribution of anatomical hierarchy by regressing out the T1w/T2w gradient from both timescale and individual gene expression gradients. We then fit partial least squares (PLS) models to simultaneously estimate regression weights for all genes (Whitaker et al., 2016), submitting those with significant associations for gene ontology enrichment analysis (GOEA) (Klopfenstein et al., 2018).

We find that genes highly associated with neuronal timescales are preferentially related to transmembrane ion transporter complexes, as well as GABAergic synapses and chloride channels (Fig. 4F, Table S3 and S4). When restricted to positively-associated genes only (expression increases with timescales), one functional group related to phosphatidate phosphatase activity is uncovered, including the gene PLPPR1, which has been linked to neuronal plasticity(Savaskan et al., 2004). Conversely, genes that are negatively associated with timescale are related to numerous groups involved in the construction and functioning of transmembrane transporters and voltage-gated ion channels, especially potassium and other inorganic cation transporters. The discovery of these gene ontology items suggests that inhibition (Telenczuk et al., 2017)—mediated by GABA and chloride channels—and voltage-gated potassium channels have prominent roles in shaping neuronal timescale dynamics at the macroscale level, beyond what's expected based on the anatomical hierarchy alone.

**Fig. 3. Timescale gradient is linked to expression of genes related to synaptic receptors and transmembrane ion channels across the human cortex. (A)** timescale gradient follows the dominant axis of gene expression variation across the cortex (PC1, arbitrary direction). **(B)** timescale gradient is significantly correlated with expression of genes known to alter synaptic and neuronal membrane time constants, as well as inhibitory cell-type markers, but **(C)** members within a gene family (e.g., NMDA receptor subunits) can be both positively and negatively associated with timescales. **(D)** macroscale timescale-transcriptomic correlation captures association between RNA-sequenced expression of the same genes and single-cell timescale properties fit to patch clamp data (Bomkamp et al., 2019; Tripathy et al., 2017), and the correspondence improves for genes (separated by quintiles) that are more strongly correlated with timescale (horizontal lines: correlation across all genes from (Bomkamp et al., 2019; Tripathy et al., 2017), $\rho$ = 0.36 and 0.25). **(E)** T1w/T2w gradient is regressed out from timescale and gene expression gradients, and a partial least squares (PLS) model is fit to the residual maps. Genes with significant PLS weights are submitted for gene ontology enrichment analysis. **(F)** enriched

genes are primarily linked to transmembrane transporters and GABA-ergic synapses; genes
specifically with strong negative associations further over-represent transmembrane ion
exchange mechanisms, especially voltage-gated potassium and cation transporters. Spatial
correlation p-values in **(A-C)** are adjusted for spatial autocorrelation (see Materials and Methods;
asterisks in **(B,D)** indicate $p$ < 0.05, 0.01, 0.005, and 0.001 respectively; filled circles in **(C,D)**
indicate $p$ < 0.05).

263
264

**Timescales lengthen in working memory and shorten in aging**
Finally, we investigate whether timescales are functionally dynamic and relevant for human
cognition. While previous studies have shown hierarchical segregation of task-relevant
information corresponding to intrinsic timescales of different cortical regions (Baldassano et al.,
2017; Chien and Honey, 2020; Honey et al., 2012; Runyan et al., 2017; Sarafyazd and Jazayeri,
2019; Wasmuht et al., 2018), as well as optimal adaptation of behavioral timescales to match the
environment (Ganupuru et al., 2019; Ossmy et al., 2013), evidence for functionally relevant
changes in regional neuronal timescales is lacking. Here, we examine whether timescales
undergo short- and long-term shifts during working memory maintenance and aging, respectively.

We first analyze human ECoG recordings where participants performed a visuospatial working
memory task that requires a delayed cued response (Fig. 4A) (Johnson et al., 2018a). Neuronal
timescales were extracted for pre-stimulus baseline and memory maintenance delay periods (900
ms, both stimulus-free). Replicating our previous result, we observe that baseline neuronal
timescales follow a hierarchical progression across association regions (Fig. 4B). If neuronal
timescales track the temporal persistence of information in a functional manner, then they should
expand during delay periods. Consistent with our prediction, timescales in all regions are ~20%
longer during delay periods (Fig. 4C; $p$ < 0.005 for all regions). Moreover, timescale changes in
the PFC are significantly correlated with behavior across participants, where longer delay-period
timescales relative to baseline are associated with better working memory performance (Fig. 4D,
$\rho$ = 0.75, $p$ = 0.003). No other spectral features in the recorded brain regions experience
consistent changes from baseline to delay periods while also significantly correlating with
individual performance, including the 1/f-like spectral exponent, narrowband theta (3-8 Hz), and
high-frequency (high gamma; 70-100 Hz) activity power (Fig. S6).

In the long-term, aging is associated with a broad range of functional and structural changes,
such as working memory impairments (Voytek et al., 2015b; Wang et al., 2011), as well as
changes in neuronal dynamics (Voytek and Knight, 2015; Voytek et al., 2015b; Wang et al.,
2011) and cortical structure (Pegasiou et al., 2020; de Villers-Sidani et al., 2010), such as the
loss of slow-deactivating NMDA receptor subunits (Pegasiou et al., 2020). Since neuronal
timescales support working memory maintenance, we specifically predict that timescales
shorten across the lifespan, in agreement with the observed cognitive and structural
deteriorations. To this end, we leverage the wide age range in the MNI-iEEG dataset (13-62
years old) and probe cortical timescales for each participant as a function of age. We observe
that older adults have faster neuronal timescales ($\rho$ = -0.31, $p$ = 0.010; Fig. 4E and Fig. S7; see
Materials and Methods), and that timescales shorten with age in most areas across the cortex ($t$

= -7.04, *p* < 0.001). This timescale compression is especially prominent in sensorimotor,
temporal, and medial frontal regions (Fig. 4F and Fig. S7). These results support our hypothesis
that neuronal timescales, estimated from transmembrane current fluctuations, can rapidly shift
in a functionally relevant manner, as well as slowly—over decades—in healthy aging.

**Fig. 4. Timescales expand during working memory maintenance while tracking performance, and task-free average timescales compress in older adults. (A)** 14 participants with overlapping intracranial coverage performed a visuospatial working memory task, with baseline (pre-stimulus) and delay period data analyzed (PC: parietal, PFC: prefrontal, OFC: orbitofrontal, MTL: medial temporal; n denotes number of subjects). **(B)** baseline timescales follow hierarchical organization within association regions (*: *p* < 0.05; mean±s.e.m. across participants). **(C)** all regions show significant timescale increase during delay period compared to baseline (***: *p* < 0.005, ****: *p* < 0.001, one-sample t-test). **(D)** PFC timescale expansion during delay periods predicts working memory accuracy across participants (mean±s.e.m. across PFC electrodes); inset: correlation between working memory accuracy and timescale change across regions. **(E)** in the MNI-iEEG dataset, participant-average cortical timescales decrease (become faster) with age. **(F)** most cortical parcels show a negative relationship between timescales and age, with the exception of parts of the visual cortex and the temporal poles (one-sample t-test, *t* = -7.04).

## Discussion

Theoretical accounts and converging empirical evidence predict a graded variation of neuronal timescales across the human cortex (Chaudhuri et al., 2015; Huntenburg et al., 2018; Wang, 2020), which reflects functional specialization and implements hierarchical temporal processing crucial for complex cognition (Kiebel et al., 2008). This timescale gradient is thought to emerge

327    as a consequence of cortical variations in cytoarchitecture and microcircuit connectivity, thus
328    linking brain structure to function. In this work, we infer the timescale of non-rhythmic
329    transmembrane current fluctuations from invasive human intracranial recordings and test those
330    predictions explicitly.
331

We find that neuronal timescales vary continuously across the human cortex and coincide with
the anatomical hierarchy, with timescales increasing from primary sensory and motor to
association regions. Timescales inferred from macaque ECoG scale with single-unit spiking
timescales, corroborating the fact that field potential signals mainly reflect fast transmembrane
and synaptic currents (Buzsáki et al., 2012), whose timescales are related to, but distinct from,
single-unit timescales measured in previous studies (Murray et al., 2014; Ogawa and Komatsu,
2010; Wasmuht et al., 2018). Because field potential fluctuations are driven by currents from
both locally generated and distal inputs, our results raise questions on how and when these
timescales interact to shape downstream spiking dynamics.
341

Furthermore, transcriptomic analysis demonstrates the specific roles that transmembrane ion
transporters and synaptic receptors play in establishing the cortical gradient of neuronal
timescales, over and above the degree predicted by the principal structural gradient alone. The
expression of voltage-gated potassium channel, chloride channel, and GABAergic receptor
genes, in particular, are strongly associated with the spatial variation of neuronal timescale.
Remarkably, we find that electrophysiology/transcriptomic relationships discovered at the single-
cell level, through patch-clamp recordings and single-cell RNA sequencing, are recapitulated at
the macroscale between bulk gene expression and timescales inferred from ECoG. Our findings
motivate further studies for investigating the precise roles voltage-gated ion channels and
synaptic inhibition play in shaping functional neuronal timescales through causal manipulations,
complementary to existing lines of research focusing on NMDA activation and recurrent circuit
motifs.
354

Finally, we show that neuronal timescales are not static, but can change both in the short- and
long-term. Transmembrane current timescales across multiple association regions, including
parietal, frontal, and medial temporal cortices, increase during the delay period of a working
memory task, consistent with the emergence of persistent spiking during working memory delay.
Working memory performance across individuals, however, is predicted by the extent of
timescale increase in the PFC only. This further suggests that behavior-relevant neural activity
may be localized despite widespread task-related modulation (Pinto et al., 2019), even at the
level of neuronal membrane fluctuations. In the long-term, we find that neuronal timescale
shortens with age in most cortical regions, linking age-related synaptic, cellular, and connectivity
changes—particularly those that influence neuronal integration timescale—to the compensatory
posterior-to-anterior shift of functional specialization in healthy aging (Davis et al., 2008).
366

Overall, we identify consistent and converging patterns between transcriptomics, anatomy,
dynamics, and function across multiple datasets of different modalities from different individuals
and multiple species. As a result, evidence for these relationships can be supplemented by
more targeted approaches such as imaging of receptor metabolism. Furthermore, the

introduction and validation of a novel method for inferring timescales from macroscale
electrophysiological recordings potentially allows for the non-invasive estimation of neuronal
timescales, using widely accessible tools such as EEG and MEG (Demirtaş et al., 2019). These
results open up many avenues of research for discovering potential relationships between
microscale gene expression and anatomy with the dynamics of neuronal population activity at
the macroscale in humans.
377
**References**

Alexander-Bloch, A.F., Shou, H., Liu, S., Satterthwaite, T.D., Glahn, D.C., Shinohara, R.T.,
Vandekar, S.N., and Raznahan, A. (2018). On testing for spatial correspondence between maps
of human brain structure and function. Neuroimage *178*, 540–551.

Badre, D., and D'Esposito, M. (2009). Is the rostro-caudal axis of the frontal lobe hierarchical?
Nat. Rev. Neurosci. *10*, 659–669.

Baldassano, C., Chen, J., Zadbood, A., Pillow, J.W., Hasson, U., and Norman, K.A. (2017).
Discovering Event Structure in Continuous Narrative Perception and Memory. Neuron *95*, 709–
721.e5.

Bauer, S. (2017). Gene-Category Analysis. In The Gene Ontology Handbook, C. Dessimoz, and
N. Škunca, eds. (New York, NY: Springer New York), pp. 175–188.

Bomkamp, C., Tripathy, S.J., Bengtsson Gonzales, C., Hjerling-Leffler, J., Craig, A.M., and
Pavlidis, P. (2019). Transcriptomic correlates of electrophysiological and morphological diversity
within and across excitatory and inhibitory neuron classes. PLoS Comput. Biol. *15*, e1007113.

Brett, M., Markiewicz, C.J., Hanke, M., Côté, M.-A., Cipollini, B., McCarthy, P., Jarecka, D.,
Cheng, C.P., Halchenko, Y.O., Cottaar, M., et al. (2020). nipy/nibabel: 3.1.0.

Burt, J.B., Demirtaş, M., Eckner, W.J., Navejar, N.M., Ji, J.L., Martin, W.J., Bernacchia, A.,
Anticevic, A., and Murray, J.D. (2018). Hierarchy of transcriptomic specialization across human
cortex captured by structural neuroimaging topography. Nat. Neurosci. *21*, 1251–1259.

Burt, J.B., Helmer, M., Shinn, M., Anticevic, A., and Murray, J.D. (2020). Generative modeling of
brain maps with spatial autocorrelation.

Buzsáki, G., Anastassiou, C.A., and Koch, C. (2012). The origin of extracellular fields and
currents--EEG, ECoG, LFP and spikes. Nat. Rev. Neurosci. *13*, 407–420.

Chaudhuri, R., Knoblauch, K., Gariel, M.-A., Kennedy, H., and Wang, X.-J. (2015). A Large-
Scale Circuit Mechanism for Hierarchical Dynamical Processing in the Primate Cortex. Neuron
*88*, 419–431.

Chien, H.-Y.S., and Honey, C.J. (2020). Constructing and Forgetting Temporal Context in the
Human Cerebral Cortex. Neuron.

Cole, S., Donoghue, T., Gao, R., and Voytek, B. (2019). NeuroDSP: a package for neural digital
signal processing. Journal of Open Source Software *4*, 1272.

Davis, S.W., Dennis, N.A., Daselaar, S.M., Fleck, M.S., and Cabeza, R. (2008). Que PASA?

409 The posterior--anterior shift in aging. Cereb. Cortex *18*, 1201–1209.

410 Demirtaş, M., Burt, J.B., Helmer, M., Ji, J.L., Adkinson, B.D., Glasser, M.F., Van Essen, D.C.,
411 Sotiropoulos, S.N., Anticevic, A., and Murray, J.D. (2019). Hierarchical Heterogeneity across
412 Human Cortex Shapes Large-Scale Neural Dynamics. Neuron *101*, 1181–1194.e13.

413 Duarte, R., and Morrison, A. (2019). Leveraging heterogeneity for neural computation with
414 fading memory in layer 2/3 cortical microcircuits. PLoS Comput. Biol. *15*, e1006781.

415 Duarte, R., Seeholzer, A., Zilles, K., and Morrison, A. (2017). Synaptic patterning and the
416 timescales of cortical dynamics. Curr. Opin. Neurobiol. *43*, 156–165.

417 Eyre, M.D., Renzi, M., Farrant, M., and Nusser, Z. (2012). Setting the time course of inhibitory
418 synaptic currents by mixing multiple GABA(A) receptor α subunit isoforms. J. Neurosci. *32*,
419 5853–5867.

420 Fagerberg, L., Hallström, B.M., Oksvold, P., Kampf, C., Djureinovic, D., Odeberg, J., Habuka,
421 M., Tahmasebpoor, S., Danielsson, A., Edlund, K., et al. (2014). Analysis of the human tissue-
422 specific expression by genome-wide integration of transcriptomics and antibody-based
423 proteomics. Mol. Cell. Proteomics *13*, 397–406.

424 Flint, A.C., Maisch, U.S., Weishaupt, J.H., Kriegstein, A.R., and Monyer, H. (1997). NR2A
425 subunit expression shortens NMDA receptor synaptic currents in developing neocortex. J.
426 Neurosci. *17*, 2469–2476.

427 Frauscher, B., von Ellenrieder, N., Zelmann, R., Doležalová, I., Minotti, L., Olivier, A., Hall, J.,
428 Hoffmann, D., Nguyen, D.K., Kahane, P., et al. (2018a). Atlas of the normal intracranial
429 electroencephalogram: neurophysiological awake activity in different cortical areas. Brain *141*,
430 1130–1144.

431 Frauscher, B., von Ellenrieder, N., Zelmann, R., Rogers, C., Nguyen, D.K., Kahane, P., Dubeau,
432 F., and Gotman, J. (2018b). High-Frequency Oscillations in the Normal Human Brain. Ann.
433 Neurol. *84*, 374–385.

434 Ganupuru, P., Goldring, A.B., Harun, R., and Hanks, T.D. (2019). Flexibility of Timescales of
435 Evidence Evaluation for Decision Making. Curr. Biol. *29*, 2091–2097.e4.

436 Gao, R., Peterson, E.J., and Voytek, B. (2017). Inferring synaptic excitation/inhibition balance
437 from field potentials. Neuroimage *158*, 70–78.

438 Genovese, G., Fromer, M., Stahl, E.A., Ruderfer, D.M., Chambert, K., Landén, M., Moran, J.L.,
439 Purcell, S.M., Sklar, P., Sullivan, P.F., et al. (2016). Increased burden of ultra-rare protein-
440 altering variants among 4,877 individuals with schizophrenia. Nat. Neurosci. *19*, 1433–1441.

441 Gjorgjieva, J., Drion, G., and Marder, E. (2016). Computational implications of biophysical
442 diversity and multiple timescales in neurons and synapses for circuit performance. Curr. Opin.
443 Neurobiol. *37*, 44–52.

444 Glasser, M.F., and Van Essen, D.C. (2011). Mapping human cortical areas in vivo based on
445 myelin content as revealed by T1- and T2-weighted MRI. J. Neurosci. *31*, 11597–11616.

446 Glasser, M.F., Coalson, T.S., Robinson, E.C., Hacker, C.D., Harwell, J., Yacoub, E., Ugurbil, K.,
447 Andersson, J., Beckmann, C.F., Jenkinson, M., et al. (2016). A multi-modal parcellation of

human cerebral cortex. Nature *536*, 171–178.

Gold, J.I., and Shadlen, M.N. (2007). The neural basis of decision making. Annu. Rev. Neurosci.
*30*, 535–574.

Gryglewski, G., Seiger, R., James, G.M., Godbersen, G.M., Komorowski, A., Unterholzner, J.,
Michenthaler, P., Hahn, A., Wadsak, W., Mitterhauser, M., et al. (2018). Spatial analysis and
high resolution mapping of the human whole-brain transcriptome for integrative analysis in
neuroimaging. Neuroimage *176*, 259–267.

Haller, M., Donoghue, T., Peterson, E., Varma, P., Sebastian, P., Gao, R., Noto, T., Knight,
R.T., Shestyuk, A., and Voytek, B. (2018). Parameterizing neural power spectra.

Hawrylycz, M., Miller, J.A., Menon, V., Feng, D., Dolbeare, T., Guillozet-Bongaarts, A.L., Jegga,
A.G., Aronow, B.J., Lee, C.-K., Bernard, A., et al. (2015). Canonical genetic signatures of the
adult human brain. Nat. Neurosci. *18*, 1832–1844.

Hawrylycz, M.J., Lein, E.S., Guillozet-Bongaarts, A.L., Shen, E.H., Ng, L., Miller, J.A., van de
Lagemaat, L.N., Smith, K.A., Ebbert, A., Riley, Z.L., et al. (2012). An anatomically
comprehensive atlas of the adult human brain transcriptome. Nature *489*, 391–399.

Hilgetag, C.C., and Goulas, A. (2020). "Hierarchy" in the organization of brain networks. Philos.
Trans. R. Soc. Lond. B Biol. Sci. *375*, 20190319.

Honey, C.J., Thesen, T., Donner, T.H., Silbert, L.J., Carlson, C.E., Devinsky, O., Doyle, W.K.,
Rubin, N., Heeger, D.J., and Hasson, U. (2012). Slow cortical dynamics and the accumulation of
information over long timescales. Neuron *76*, 423–434.

Huntenburg, J.M., Bazin, P.-L., and Margulies, D.S. (2018). Large-Scale Gradients in Human
Cortical Organization. Trends Cogn. Sci. *22*, 21–31.

Izhikevich, L., Gao, R., Peterson, E., and Voytek, B. (2018). Measuring the average power of
neural oscillations.

Johnson, E. (2018). Intracranial EEG recordings of medial temporal, lateral frontal, and
orbitofrontal regions in 10 human adults performing a visuospatial working memory task
(CRCNS.org).

Johnson, E. (2019). Intracranial EEG recordings of lateral frontal and parietal regions in 7
human adults performing a visuospatial working memory task (CRCNS.org).

Johnson, E.L., Adams, J.N., Solbakk, A.-K., Endestad, T., Larsson, P.G., Ivanovic, J., Meling,
T.R., Lin, J.J., and Knight, R.T. (2018a). Dynamic frontotemporal systems process space and
time in working memory. PLoS Biol. *16*, e2004274.

Johnson, E.L., King-Stephens, D., Weber, P.B., Laxer, K.D., Lin, J.J., and Knight, R.T. (2018b).
Spectral Imprints of Working Memory for Everyday Associations in the Frontoparietal Network.
Front. Syst. Neurosci. *12*, 65.

Kiebel, S.J., Daunizeau, J., and Friston, K.J. (2008). A hierarchy of time-scales and the brain.
PLoS Comput. Biol. *4*, e1000209.

Klopfenstein, D.V., Zhang, L., Pedersen, B.S., Ramírez, F., Warwick Vesztrocy, A., Naldi, A.,

Mungall, C.J., Yunes, J.M., Botvinnik, O., Weigel, M., et al. (2018). GOATOOLS: A Python
library for Gene Ontology analyses. Sci. Rep. *8*, 10872.

Koch, C., Rapp, M., and Segev, I. (1996). A brief history of time (constants). Cereb. Cortex *6*,
93–101.

Lerner, Y., Honey, C.J., Silbert, L.J., and Hasson, U. (2011). Topographic mapping of a
hierarchy of temporal receptive windows using a narrated story. J. Neurosci. *31*, 2906–2915.

Margulies, D.S., Ghosh, S.S., Goulas, A., Falkiewicz, M., Huntenburg, J.M., Langs, G., Bezgin,
G., Eickhoff, S.B., Castellanos, F.X., Petrides, M., et al. (2016). Situating the default-mode
network along a principal gradient of macroscale cortical organization. Proc. Natl. Acad. Sci. U.
S. A. *113*, 12574–12579.

Miller, K.J., Sorensen, L.B., Ojemann, J.G., and den Nijs, M. (2009). Power-law scaling in the
brain surface electric potential. PLoS Comput. Biol. *5*, e1000609.

Monyer, H., Burnashev, N., Laurie, D.J., Sakmann, B., and Seeburg, P.H. (1994).
Developmental and regional expression in the rat brain and functional properties of four NMDA
receptors. Neuron *12*, 529–540.

Mukamel, R., Gelbard, H., Arieli, A., Hasson, U., Fried, I., and Malach, R. (2005). Coupling
between neuronal firing, field potentials, and FMRI in human auditory cortex. Science *309*, 951–
954.

Murray, J.D., Bernacchia, A., Freedman, D.J., Romo, R., Wallis, J.D., Cai, X., Padoa-Schioppa,
C., Pasternak, T., Seo, H., Lee, D., et al. (2014). A hierarchy of intrinsic timescales across
primate cortex. Nat. Neurosci. *17*, 1661–1663.

Nagasaka, Y., Shimoda, K., and Fujii, N. (2011). Multidimensional recording (MDR) and data
sharing: an ecological open research and educational platform for neuroscience. PLoS One *6*,
e22561.

Ogawa, T., and Komatsu, H. (2010). Differential temporal storage capacity in the baseline
activity of neurons in macaque frontal eye field and area V4. J. Neurophysiol. *103*, 2433–2445.

Ossmy, O., Moran, R., Pfeffer, T., Tsetsos, K., Usher, M., and Donner, T.H. (2013). The
timescale of perceptual evidence integration can be adapted to the environment. Curr. Biol. *23*,
981–986.

Pedregosa, F. (2011). Scikit-learn: Machine Learning in Python. J. Mach. Learn. Res. *12*, 2825–
2830.

Pegasiou, C.M., Zolnourian, A., Gomez-Nicola, D., Deinhardt, K., Nicoll, J.A.R., Ahmed, A.I.,
Vajramani, G., Grundy, P., Verhoog, M.B., Mansvelder, H.D., et al. (2020). Age-Dependent
Changes in Synaptic NMDA Receptor Composition in Adult Human Cortical Neurons. Cereb.
Cortex.

Pinto, L., Rajan, K., DePasquale, B., Thiberge, S.Y., Tank, D.W., and Brody, C.D. (2019). Task-
Dependent Changes in the Large-Scale Dynamics and Necessity of Cortical Regions. Neuron
*104*, 810–824.e9.

Podvalny, E., Noy, N., Harel, M., Bickel, S., Chechik, G., Schroeder, C.E., Mehta, A.D.,

525 Tsodyks, M., and Malach, R. (2015). A unifying principle underlying the extracellular field
526 potential spectral responses in the human cortex. J. Neurophysiol. *114*, 505–519.

527 Runyan, C.A., Piasini, E., Panzeri, S., and Harvey, C.D. (2017). Distinct timescales of
528 population coding across cortex. Nature *548*, 92–96.

529 Sarafyazd, M., and Jazayeri, M. (2019). Hierarchical reasoning by neural circuits in the frontal
530 cortex. Science *364*.

531 Savaskan, N.E., Bräuer, A.U., and Nitsch, R. (2004). Molecular cloning and expression
532 regulation of PRG-3, a new member of the plasticity-related gene family. Eur. J. Neurosci. *19*,
533 212–220.

534 Telenczuk, B., Dehghani, N., Le Van Quyen, M., Cash, S.S., Halgren, E., Hatsopoulos, N.G.,
535 and Destexhe, A. (2017). Local field potentials primarily reflect inhibitory neuron activity in
536 human and monkey cortex. Sci. Rep. *7*.

537 Tripathy, S.J., Toker, L., Li, B., Crichlow, C.-L., Tebaykin, D., Mancarci, B.O., and Pavlidis, P.
538 (2017). Transcriptomic correlates of neuron electrophysiological diversity. PLoS Comput. Biol.
539 *13*, e1005814.

540 Vértes, P.E., Rittman, T., Whitaker, K.J., Romero-Garcia, R., Váša, F., Kitzbichler, M.G.,
541 Wagstyl, K., Fonagy, P., Dolan, R.J., Jones, P.B., et al. (2016). Gene transcription profiles
542 associated with inter-modular hubs and connection distance in human functional magnetic
543 resonance imaging networks. Philos. Trans. R. Soc. Lond. B Biol. Sci. *371*.

544 de Villers-Sidani, E., Alzghoul, L., Zhou, X., Simpson, K.L., Lin, R.C.S., and Merzenich, M.M.
545 (2010). Recovery of functional and structural age-related changes in the rat primary auditory
546 cortex with operant training. Proc. Natl. Acad. Sci. U. S. A. *107*, 13900–13905.

547 Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski,
548 E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0: fundamental algorithms for
549 scientific computing in Python. Nat. Methods *17*, 261–272.

550 Vos de Wael, R., Benkarim, O., Paquola, C., Lariviere, S., Royer, J., Tavakol, S., Xu, T., Hong,
551 S.-J., Langs, G., Valk, S., et al. (2020). BrainSpace: a toolbox for the analysis of macroscale
552 gradients in neuroimaging and connectomics datasets. Commun Biol *3*, 103.

553 Voytek, B., and Knight, R.T. (2015). Dynamic network communication as a unifying neural basis
554 for cognition, development, aging, and disease. Biol. Psychiatry *77*, 1089–1097.

555 Voytek, B., Kayser, A.S., Badre, D., Fegen, D., Chang, E.F., Crone, N.E., Parvizi, J., Knight,
556 R.T., and D'Esposito, M. (2015a). Oscillatory dynamics coordinating human frontal networks in
557 support of goal maintenance. Nat. Neurosci. *18*, 1318–1324.

558 Voytek, B., Kramer, M.A., Case, J., Lepage, K.Q., Tempesta, Z.R., Knight, R.T., and Gazzaley,
559 A. (2015b). Age-Related Changes in 1/f Neural Electrophysiological Noise. J. Neurosci. *35*,
560 13257–13265.

561 van Vugt, B., van Kerkoerle, T., Vartak, D., and Roelfsema, P.R. (2020). The Contribution of
562 AMPA and NMDA Receptors to Persistent Firing in the Dorsolateral Prefrontal Cortex in
563 Working Memory. J. Neurosci. *40*, 2458–2470.

Wagner, H.H., and Dray, S. (2015). Generating spatially constrained null models for irregularly
spaced data using M oran spectral randomization methods. Methods Ecol. Evol. *6*, 1169–1178.

Wang, X.J. (1999). Synaptic basis of cortical persistent activity: the importance of NMDA
receptors to working memory. J. Neurosci. *19*, 9587–9603.

Wang, X.-J. (2002). Probabilistic decision making by slow reverberation in cortical circuits.
Neuron *36*, 955–968.

Wang, X.-J. (2008). Decision making in recurrent neuronal circuits. Neuron *60*, 215–234.

Wang, X.-J. (2020). Macroscopic gradients of synaptic excitation and inhibition in the neocortex.
Nat. Rev. Neurosci.

Wang, M., Gamo, N.J., Yang, Y., Jin, L.E., Wang, X.-J., Laubach, M., Mazer, J.A., Lee, D., and
Arnsten, A.F.T. (2011). Neuronal basis of age-related working memory decline. Nature *476*,
210–213.

Wang, M., Yang, Y., Wang, C.-J., Gamo, N.J., Jin, L.E., Mazer, J.A., Morrison, J.H., Wang, X.-
J., and Arnsten, A.F.T. (2013). NMDA receptors subserve persistent neuronal firing during
working memory in dorsolateral prefrontal cortex. Neuron *77*, 736–749.

Wasmuht, D.F., Spaak, E., Buschman, T.J., Miller, E.K., and Stokes, M.G. (2018). Intrinsic
neuronal dynamics predict distinct functional roles during working memory. Nat. Commun. *9*,
3499.

Watanabe, T., Rees, G., and Masuda, N. (2019). Atypical intrinsic neural timescale in autism.
Elife *8*.

Whitaker, K.J., Vértes, P.E., Romero-Garcia, R., Váša, F., Moutoussis, M., Prabhu, G.,
Weiskopf, N., Callaghan, M.F., Wagstyl, K., Rittman, T., et al. (2016). Adolescence is
associated with genomically patterned consolidation of the hubs of the human brain
connectome. Proc. Natl. Acad. Sci. U. S. A. *113*, 9105–9110.

Wimmer, K., Nykamp, D.Q., Constantinidis, C., and Compte, A. (2014). Bump attractor
dynamics in prefrontal cortex explains behavioral precision in spatial working memory. Nat.
Neurosci. *17*, 431–439.

Yanagawa, T., Chao, Z.C., Hasegawa, N., and Fujii, N. (2013). Large-scale information flow in
conscious and unconscious states: an ECoG study in monkeys. PLoS One *8*, e80845.

Zylberberg, J., and Strowbridge, B.W. (2017). Mechanisms of Persistent Activity in Cortical
Circuits: Possible Neural Substrates for Working Memory. Annu. Rev. Neurosci. *40*, 603–627.

595
596

**Acknowledgements**
**Funding:** Natural Sciences and Engineering Research Council of Canada (NSERC PGS-D)
and Katzin Prize (to R.G.). Alexander Von Humboldt Foundation fellowship for post-doctoral
researchers (to R.L.v.d.B). Deutsche Forschungsgemeinschaft (DFG) SFB936/A7/Z3 (to Tobias
H. Donner). Sloan Research Fellowship (FG-2015-66057), the Whitehall Foundation (2017-12-
73), the National Science Foundation under grant BCS-1736028, the NIH National Institute of
General Medical Sciences grant R01GM134363-01, a UC San Diego, Shiley-Marcos
Alzheimer's Disease Research Center (ADRC): Research Training in Alzheimer's Disease
grant, and a Halıcıoğlu Data Science Institute Fellowship (to B.V.).
606
607     **Author contributions:** R.G. and B.V. conception and electrophysiological data analysis. R.G.
608     simulations, anatomical, and gene expression analysis. R.v.d.B and T.P. data preprocessing
609     and conception of anatomical and gene expression analysis. R.G. wrote the manuscript with
610     B.V. All authors edited the manuscript.
611
612     **Competing interests:** the authors declare no competing interests.
613
614     **Data and materials availability:** all data analyzed in this manuscript are from open data
615     sources. All code used for all analyses and plots are publicly available on GitHub at
616     https://github.com/rdgao/field-echos and https://github.com/rudyvdbrink/surface_projection. See
617     Table S1 and S2.

618
619
620
621

## Methods

**Inferring timescale from autocorrelation and power spectral density**

Consistent with previous studies, we define "neuronal timescale" as the exponential decay time constant ($\tau$) of the empirical autocorrelation function (ACF), or lagged correlation (Honey et al., 2012; Murray et al., 2014). $\tau$ can be naively estimated to be the time it takes for the ACF to decrease by a factor of $e$ when there are no additional long-term, scale-free, or oscillatory processes, or by fitting a function of the form $f(t) = e^{-\frac{t}{\tau}}$ and extracting the parameter $\tau$. Equivalently, the power spectral density (PSD) is the Fourier Transform of the ACF via Wiener-Khinchin theorem, and follows a Lorentzian function of the form $L(f) = \frac{A}{k+f^{\chi}}$ for approximately exponential-decay processes, with $\chi = 2$ exactly when the ACF is solely composed of an exponential decay term, though it is often variable and in the range between 2-6 for neural time series (Haller et al., 2018; Miller et al., 2009; Podvalny et al., 2015; Voytek et al., 2015b). Timescale can be computed from the parameter $k$ as $\tau = \frac{1}{2\pi f_k}$, where $f_k \approx k^{1/\chi}$ is approximated to be the frequency at which a bend or knee in the power spectrum occurs and equality holds when $\chi = 2$.

**Computing power spectral density (PSD)**

PSDs are estimated using a modified Welch's method, where short-time windowed Fourier transforms (STFT) are computed from the time series, but the median is taken across time instead of the mean (in conventional Welch's method) to minimize the effect of high-amplitude transients and artifacts (Izhikevich et al., 2018). Custom functions for this can be found in NeuroDSP (Cole et al., 2019), a published and open-source digital signal processing toolbox for neural time series (neurodsp.spectral.compute_spectrum). For simulated data, Neurotycho macaque ECoG, and MNI-iEEG datasets, we use 1-second long Hamming windows with 0.5-s overlap. To estimate single-trial PSDs for the working memory ECoG dataset (Johnson-ECoG (Johnson et al., 2018a, 2018b)), we simply apply Hamming window to 900-ms long epoched time series and compute the squared magnitude of the windowed-Fourier transform.

**Spectral parametrization - Fitting Oscillations and 1/f (FOOOF)**

We apply spectral parameterization (Haller et al., 2018) to extract timescales from PSDs. Briefly, we decompose log-power spectra into a summation of narrowband periodic components—modeled as Gaussians—and an aperiodic component—modeled as a generalized Lorentzian function centered at 0 Hz ($L(f)$ above). For inferring decay timescale, this formalism can be practically advantageous when a strong oscillatory or variable power-law ($\chi$) component is present, as is often the case for neural signals. While oscillatory and power-law components can corrupt naive measurements of $\tau$ as time for the ACF to reach 1/$e$, they can be easily accounted for and ignored in the frequency domain as narrowband peaks and 1/f-exponent fit. We discard the periodic components and infer timescale from the aperiodic component of the PSD. For a complete mathematical description of the model, see (Haller et al., 2018).

**Simulation and validation**

We simulate the aperiodic background component of neural field potential recordings as autocorrelated stochastic processes by convolving Poisson population spikes with exponentially-decaying synaptic kernels with predefined decay time constants (neurodsp.sim.sim_synaptic_current). PSDs of the simulated data are computed and parameterized as described above, and we compare the fitted timescales with their ground-truth values.

**Macaque ECoG and single unit timescales data**

Macaque single-unit timescales are taken directly from values reported in Fig. 1c of (Murray et al., 2014). Whole-brain surface ECoG data (1000Hz sampling rate) is taken from the Neurotycho repository(Nagasaka et al., 2011; Yanagawa et al., 2013), with 8 sessions of 128-channel recordings from two animals (George and Chibi, 4 sessions each). Results reported in Fig. 2 are from ~10 minutes eyes-open resting periods to match the pre-stimulus baseline condition of single-unit experiments. Timescales for individual ECoG channels are extracted and averaged over regions corresponding to single-unit recording areas from(Murray et al., 2014) (Fig. 2C inset and Fig. S3), which are selected visually based on the overlapping cortical map and landmark sulci/gyri. Each region included between 2-4 electrodes (see Fig. S3C for selected ECoG channel indices for each region).

**Statistical analysis for macaque ECoG and spiking timescale**

For each individual recording session, as well as the grand average, Spearman rank correlation was computed between spiking and ECoG timescales. Linear regression models were fit using the python package scipy (Virtanen et al., 2020) (scipy.stats.linregress) and the linear slope was used to compute the scaling coefficient between spiking and ECoG timescales.

**Variations in neuronal timescale, T1/T2 ratio, and mRNA expression across human cortex**

The following sections describe procedures for generating the average cortical gradient maps for neuronal timescale, MR-derived T1w/T2w ratio, and gene expression from the respective raw datasets. All maps were projected onto the 180 left hemisphere parcels of Human Connectome Project's Multimodal Parcellation (Glasser et al., 2016) (HCP-MMP1.0) for comparison, described in the individual sections. All spatial correlations are computed as Spearman rank correlations between maps. Procedure for computing statistical significance while accounting for spatial autocorrelation is described in detail below under the sections **spatial statistics** and **spatial autocorrelation modeling**.

**Neuronal timescale map**

The MNI Open iEEG dataset consists of 1 minute of resting state data across 1772 channels from 106 epilepsy patients (13-62 years old, 58 males and 48 females), recorded using either surface strip/grid or stereoEEG electrodes, and cleaned of visible artifacts (Frauscher et al., 2018a, 2018b). Neuronal timescales were extracted from PSDs of individual channels, and projected from MNI voxel coordinates onto HCP-MMP1.0 surface parcellation as follows:

For each patient, timescale estimated from each electrode was extrapolated to the rest of the
cortex in MNI coordinates using a Gaussian weighting function (confidence mask), $w(r) =$
$e^{-(r^2/\alpha^2)}$, where r is the Euclidean distance between the electrode and a voxel, and α is the
distance scaling constant, chosen here such that a voxel 4mm away has 50% weight (or,
confidence). Timescale at each voxel is computed as a weighted spatial average of timescales
from all electrodes (i) of that patient,

i.e., $\tau_{voxel} = \frac{\sum_i w(r_i) \tau_i}{\sum_i w(r_i)}$.

Similarly, each voxel is assigned a confidence rating that is the maximum of weights over all
electrodes ($w_{voxel}(r_{min})$, of the closest electrode), i.e., a voxel right under an electrode has a
confidence of 1, while a voxel 4mm away from the closest electrode has a confidence of 0.5, etc.
717

Timescales for each HCP-MMP parcel were then computed as the confidence-weighted
arithmetic mean across all voxels that fall within the boundaries of that parcel. HCP-MMP
boundary map is loaded and used for projection using NiBabel (Brett et al., 2020). This results in
a 180 parcels-by-106 patients timescale matrix. A per-parcel confidence matrix of the same
dimensions was computed by taking the maximum confidence over all voxels for each parcel (Fig.
S1A). The average cortical timescale map (gradient) is computed by taking the confidence-
weighted average at each parcel across all participants. Note that this procedure for locally
thresholded and weighted average is different from projection procedures used for the mRNA and
T1w/T2w data due to region-constrained and heterogeneous ECoG electrode sites across
participants. While coverage is sparse and idiosyncratic in individual participants, it does not vary
as a function of age, and when pooling across the entire population, 178 of 180 parcels have at
least one patient with an electrode within 4mm, with the best coverage in later sensorimotor,
temporal, and frontal regions (Fig. S1).
731

**T1w/T2w ratio map**
As a measure of structural cortical hierarchy, we used the ratio between T1- and T2-weighted
structural MRI, referred to as T1w/T2w map in main text, or the myelin map (Burt et al., 2018;
Glasser and Van Essen, 2011). Since there is little variation in the myelin map across individuals,
we used the group average myelin map of the WU-Minn HCP S1200 release (N = 1096, March
1, 2017 release) provided in HCP-MMP1.0 surface space. For correlation with other variables, we
computed the median value per parcel, identical to the procedure for mRNA expression below.
739

**mRNA expression maps**
We used the Allen Human Brain Atlas (AHBA) gene expression dataset (Hawrylycz et al., 2015,
2012) that comprised postmortem samples of 6 donors (1 female, 5 male) that underwent
microarray transcriptional profiling. Spatial maps of mRNA expression were available in
volumetric 2 mm isotropic MNI space, following improved nonlinear registration and whole-brain
prediction using variogram modeling as implemented by (Gryglewski et al., 2018). We used
whole-brain maps available from (Gryglewski et al., 2018) rather than the native sample-wise
values in the AHBA database to prevent bias that could occur due to spatial inhomogeneity of the
sampled locations. In total, 18114 genes were included for analyses that related to the dominant
axis of expression across the genome.

We projected the volumetric mRNA expression data onto the HCP-MMP cortical surface using
the HCP workbench software (v1.3.1 running on Windows OS 10) with the "enclosing" method,
and custom MATLAB code (github.com/rudyvdbrink/surface_projection). The enclosing method
extracts for all vertices on the surface the value from enclosing voxels in the volumetric data.
Alternative projection methods such as trilinear 3D linear interpolation of surrounding voxels, or
ribbon mapping that constructs a polyhedron from each vertex's neighbors on the surface to
compute a weighted mean for the respective vertices, yielded comparable values, but less
complete cortical coverage. Moreover, the enclosing method ensured that no transformation of
the data (non-linear or otherwise) occurred during the projection process and thus the original
values in the volumetric data were preserved.

Next, for each parcel of the left hemisphere in HCP-MMP, we extracted the median vertex-wise
value. We used the median rather than the mean because it reduced the contribution of outliers
in expression values within parcels. Vertices that were not enclosed by voxels that contained data
in volumetric space were not included in the parcel-wise median. This was the case for 539
vertices (1.81% of total vertices). Linear interpolation across empty vertices prior to computing
median parcel-wise values yielded near-identical results ($r$ = 0.95 for reconstructed surfaces).
Lastly, expression values were mean and variance normalized across parcels to facilitate
visualization. Normalization had no effect on spatial correlation between gene expression and
other variables since the spatial distribution of gene expression was left unaltered.

**Spatial statistics**
All correlations between spatial maps (timescale, T1w/T2w, gene principal component, and
individual gene expressions) were computed using Spearman rank correlation. As noted in (Burt
et al., 2018, 2020; Vos de Wael et al., 2020), neural variables vary smoothly and continuously
across the cortical surface, violating the assumption of independent samples. As a result, when
correlating two variables each with non-trivial spatial autocorrelation, the naive p-value is
artificially lowered since it is compared against an inappropriate null hypothesis, i.e., randomly
distributed or shuffled values across space. Instead, a more appropriate null hypothesis
introduces spatial autocorrelation-preserving null maps, which destroys any potential correlation
between two maps while respecting their spatial autocorrelations. For all spatial correlation
analyses, we generated N = 1000 null maps of one variable (timescale map unless otherwise
noted), and the test statistic, Spearman correlation (ρ), is computed against the other variable of
interest to build the null distribution. Two-tailed significance is then computed as the proportion of
the null distribution that is less extreme than the empirical correlation value. All regression lines
were computed by fitting a linear regression to log-timescale and the structural feature maps.

**Spatial autocorrelation modeling**
To generate spatial autocorrelation-preserving null maps, we used Moran's Spectral
Randomization (MSR) (Wagner and Dray, 2015) from the python package BrainSpace (Vos de
Wael et al., 2020). Details of the algorithm can be found in the above references. Briefly, MSR
performs eigendecomposition on a spatial weight matrix of choice, which is taken here to be the

inverse average geodesic distance matrix between all pairs of parcels in HCP-MMP1.0. The
eigenvectors of the weight matrix are then used to generate randomized null feature maps that
preserves the autocorrelation of the empirical map. We used the singleton procedure for null map
generation. All significance values reported (Fig. 2B, Fig. 3A-C) were adjusted using the above
procedure.
799
We also compare two other methods of generating null maps: spatial variogram fitting (Burt et al.,
2020) and spin permutation (Alexander-Bloch et al., 2018). Null maps were generated for
timescale using spatial variogram fitting, while for spin permutation they were generated for
vertex-wise T1w/T2w and gene PC1 maps before parcellation, so as to preserve surface locations
of the parcellation itself. All methods perform similarly, producing comparable spatial
autocorrelation in the null maps, assessed using spatial variogram, as well as null distribution of
spatial correlation coefficients between timescale and T1w/T2w (Fig. S2).
807
**Principal Component Analysis (PCA) of gene expression**
We used scikit-learn (Pedregosa, 2011) PCA (sklearn.decomposition.PCA) to identify the
dominant axes of gene expression variation across the entire AHBA dataset, as well as for brain-
specific genes. PCA was computed on the variance-normalized average gene expression maps,
*X*, an N x P matrix where N = 18114 (or N = 2429 brain-specific) genes, and P = 180 cortical
parcels. Briefly, PCA factorizes *X* such that $X = USV^T$, where *U* and *V* are unitary matrices of
dimensionality N x N and P x P, respectively. *S* is the same dimensionality as *X* and contains non-
negative descending eigenvalues on its main diagonal (Λ). Columns of *V* are defined as the
principal components (PCs), and the dominant axis of gene expression is then defined as the first
column of V, whose proportion of variance explained in the data is the first element of Λ divided
by the sum over Λ. Results for PC1 and PC2-10 are shown in Fig. 3A and Fig. S4, respectively.
819
**Selection of brain-specific genes**
Similar to (Burt et al., 2018; Fagerberg et al., 2014; Genovese et al., 2016), N=2429 brain-specific
genes were selected based on the criteria that expression in brain tissues were 4 times higher
than the median expression across all tissue types, using Supplementary Dataset 1 of (Fagerberg
et al., 2014). PC1 result shown in Fig. 3A is computed from brain-specific genes, though findings
are identical when using all genes ($\rho$ = -0.56 with timescale map, Fig. S4).
826
**Comparison of timescale-transcriptomic association with single-cell timescale genes**
Single-cell timescale genes were selected based on data from Table S3 and Online Table 1 of
(Bomkamp et al., 2019; Tripathy et al., 2017), respectively. Using single-cell RNA sequencing
data and patch-clamp recordings from transgenic mice cortical neurons, these studies identified
genes whose expression significantly correlated with electrophysiological features derived from
generalized linear integrate and fire (GLIF) model fits. We selected genes that were significantly
correlated to membrane time constant (*tau*), input resistance (*Rin* or *ri*), or capacitance (*Cm* or
*cap*) in the referenced data tables, and extracted the level of association between gene
expression and those electrophysiological feature (correlation 'DiscCorr' in (Tripathy et al., 2017)
and linear coefficient "beta_gene" in (Bomkamp et al., 2019)).
837

To compare timescale-gene expression association at the single-cell and macroscale level, we
correlated the single-cell associations extracted above with the spatial correlation coefficient
(macroscale $\rho$) between ECoG timescale and AHBA microarray expression data for those same
genes, restricting to genes with $p < 0.05$ for macroscale correlation (results identical for non-
restrictive gene set). Overall association for all genes, as well as split by quintiles of their absolute
macroscale correlation coefficient, are shown in Fig. 3D. Example "single-cell timescale" genes
shown in Fig. 3B,C are genes showing the highest correlations with those electrophysiology
features reported in Table 2 of (Bomkamp et al., 2019).
846

**T1w/T2w-removed timescale and gene expression residual maps**
To remove anatomical hierarchy as a potential mediating variable in timescale-gene expression
relationships, we linearly regress out the T1w/T2w map from the (log) timescale map and
individual gene expression maps. T1w/T2w was linearly fit to log-timescale, and the error between
T1w/T2w-predicted timescale and empirical timescale was extracted (residual); this identical
procedure was applied to every gene expression map to retrieve the gene residuals. Spatial
autocorrelation-preserving null residual maps were similarly created using MSR.
854

**Partial least squares regression model**
Due to multicollinearity in the high-dimensional gene expression dataset (many more genes than
parcels), we fit a partial least squares model to the timescale map with one output dimension
(sklearn.cross_decomposition.PLSRegression) to estimate regression coefficient for all genes
simultaneously, resulting in N=18114 (or N=2429 brain-specific) PLS weights(Vértes et al., 2016;
Whitaker et al., 2016). To determine significantly associated (or, "enriched") genes, we repeated
the above PLS-fitting procedure 1000 times but replaced the empirical timescale map (or residual
map) with null timescale maps (or residual maps) that preserved its spatial autocorrelation. Genes
whose absolute empirical PLS weight that was greater than 95% of its null weight distribution was
deemed to be enriched, and submitted for gene ontology enrichment analysis.
865

**Gene ontology enrichment analysis (GOEA)**
The Gene Ontology (GO) captures hierarchically structured relationships between GO items
representing aspects of biological processes (BP), cellular components (CC), or molecular
functions (MF). For example, "synaptic signaling", "chemical synaptic transmission", and
"glutamatergic synaptic transmission" are GO items with increasing specificity, with smaller
subsets of genes associated with each function. Each GO item is annotated with a list of genes
that have been linked to that particular process or function. GOEA examines the list of enriched
genes from above to identify GO items that are more associated with those genes than expected
by chance. We used GOATOOLS (Klopfenstein et al., 2018) to perform GOEA programmatically
in python.
876

The list of unranked genes with significant empirical PLS weights was submitted for GOEA as the
"study set", while either the full ABHA list or brain-specific gene list was used as the "reference
set". The output of GOEA is a list of GO terms with annotated genes that are enriched or purified
(i.e., preferentially appearing or missing in the study list, respectively) more often than by chance,
determined by Fisher's exact test.

Enrichment ratio is defined as follows: given a reference set with *N* total genes, and *n* were found
to be significantly associated with timescale (in the study set), for a single GO item with *B* total
genes annotated to it, where b of them overlap with the study set, then $enrichment = \frac{b/n}{B/N}$.
Statistical significance is adjusted for multiple comparisons following Benjamini-Hochberg
procedure (false discovery rate q-value reported in Fig. 3F), and all significant GO items (q < 0.05)
are reported in Fig. 3F, in addition to some example items that did not pass significance threshold.
For a detailed exposition, see (Bauer, 2017). Fig. 3F shows results using brain-specific genes.
The GO items that are significantly associated are similar when using the full gene set, but
typically with larger q-values (Tables S3 and S4) due to a mucher larger set of (non-brain-specific)
genes.

**Working memory ECoG data and analysis**
The CRCNS fcx-2 and fcx-3 datasets include 17 intracranial ECoG recordings in total from
epilepsy patients (10 and 7, respectively) performing the same visuospatial working memory task
(Johnson, 2018, 2019; Johnson et al., 2018a, 2018b). Subject 3 (s3) from fcx-2 was discarded
due to poor data quality upon examination of trial-averaged PSDs (high noise floor near 20 Hz),
while s5 and s7 from fcx-3 correspond to s5 and s8 in fcx-2 and were thus combined. Together,
data from 14 unique participants (22-50 years old, 5 female) were analyzed, with variable and
overlapping coverage in parietal cortex (PC, n=14), prefrontal cortex (PFC, n=13), orbitofrontal
cortex (OFC, n=8), and medial temporal lobe (MTL, n=9). Each channel was annotated as
belonging to one of the above macro regions.

Experimental setup is described in (Johnson, 2018, 2019; Johnson et al., 2018a, 2018b) in detail.
Briefly, following a 1-second pre-trial fixation period (baseline), subjects were instructed to focus
on one of two stimulus contexts ("identity" or "relation" information). Then two shapes were
presented in sequence for 200 ms each. After a 900 or 1150 ms jittered precue delay (delay1),
the test cue appeared for 800 ms, followed by another post-cue delay period of the same length
(delay2). Finally, the response period required participants to perform a 2-alternative forced
choice test based on the test cue, which varied based on trial condition. For our analysis, we
collapsed across the stimulus context conditions and compared neuronal timescales during the
last 900 ms of baseline and delay periods from the epoched data, which were free of visual stimuli,
in order to avoid stimulus-related event-related potential effects. Behavioral accuracy for each
experimental condition was reported for each participant, and we average across both stimulus
context conditions to produce a single working memory accuracy per participant.

Single-trial power spectra were computed for each channel as the squared magnitude of the
Hamming-windowed Fourier Transform. We used 900 ms of data in all 3 periods (pre-trial, delay1,
and delay2). Timescales were estimated by applying spectral parameterization as above, and the
two delay-period estimates were averaged to produce a single delay period value. For
comparison, we computed single-trial theta (3-8 Hz) and high-frequency activity (high gamma
(Mukamel et al., 2005), 70-100 Hz) powers as the mean log-power within those frequency bins,
as well as spectral exponent ($\chi$). Single-trial timescale difference between delay and baseline was
calculated as the difference of the log timescales due to the non-normal distribution of single-trial

timescale estimates. All other neural features were computed by subtracting baseline from the
delay period.

All neural features were then averaged across channels within the same regions, then trials, for
each participant, to produce per-participant region-wise estimates, and finally averaged across all
participants for the regional average in Fig. 4B,C. One-sample two-sided t-tests were used to
determine the statistical significance of timescale change in each region (Fig. 4C), where the null
hypothesis was no change between baseline and delay periods (i.e., delay is 100% of baseline).
Spearman rank correlation was used to determine the relationship between neural activity
(timescale; theta; high-frequency; $\chi$) change and working memory accuracy across participants
(Fig. 4D, Fig. S6).

**Per-subject average cortical timescale across age**
Since electrode coverage in the MNI-iEEG dataset is sparse and non-uniform across participants
(Fig. S1), simply averaging across parcels within individuals to estimate an average cortical
timescale per participant confounds the effect of age with the spatial effect of cortical hierarchy.
Therefore, we instead first normalize each parcel by its max value across all participants before
averaging within participants, excluding those with fewer than 10 valid parcels (71 of 106 subjects
remaining; results hold for a range of threshold values; Fig. S7B). Spearman rank correlation was
used to compute the association between age and average cortical timescale.

**Age-timescale association for individual parcels**
Each cortical parcel had a variable number of participants with valid timescale estimates above
the consistency threshold, so we compute Spearman correlation between age and timescale for
each parcel, but including only those with at least 5 participants (114 of 180 parcels, result holds
for a range of threshold values; Fig. S7C). Spatial effect of age-timescale variation is plotted in
Fig. 4F, where parcels that did not meet the threshold criteria are greyed out. Mean age-timescale
correlation from individual parcels was significantly negative under one-sample t-test.

   **Supplemental Information**

959
**Fig. S1. MNI-iEEG dataset coverage. (A),** per-parcel Gaussian-weighted mask values showing
how close the nearest electrode was to a given HCP-MMP1.0 parcel for each participant.
Brighter means closer, 0.5 corresponds to the nearest electrode being 4 mm away. **(B)**
maximum weight for each parcel across all participants. Most parcels have electrodes very
close by across the entire participant pool. **(C)** the number of HCP-MMP parcels each
participant has above the confidence threshold of 0.5 is uncorrelated with age. **(D)** number of
participants with confidence above threshold at each parcel. Sensorimotor, frontal, and lateral
temporal regions have the highest coverage. **(E)** average age of participants with confidence
above threshold at each parcel. **(F)** age distribution of participants with confidence above
threshold at each parcel. Average age per parcel (red line) is relatively stable while age
distribution varies from parcel to parcel.

**Fig. S2. Comparison of spatial autocorrelation-preserving null map generation methods.**
**(A)** distributions of Spearman correlation values between empirical T1w/T2w map and 1000
spatial-autocorrelation preserving null timescale maps generated using Moran Spectral
Randomization (MSR), spatial variogram fitting (VF), and spin permutation. Red dashed line
denotes correlation between empirical timescale and T1w/T2w maps, p-values indicate two-
tailed significance, i.e., proportion of distribution with values more extreme than empirical
correlation. **(B)** spatial variogram for empirical timescale map (black) and 10 null maps (blue)
generated using MSR and VF. Inset shows distribution of distances between pairs of HCP-MMP
parcels. **(C)** distribution of Spearman correlations between empirical and 1000 null timescale
maps generated using MSR (green) and VF (red), showing similar levels of correlation between
empirical and null maps for both methods.

**C**     indices of ECoG electrodes (1-based indexing)

|  | MT | LIP | LPFC | OFC | ACC | S1 | S2 |
|---|---|---|---|---|---|---|---|
| George | 4, 13, 22 | 10, 11, 20, 21 | 15, 24, 25, 26 | 45, 66 | 52, 57, 58, 59 | 18, 19, 30 | 1, 2, 9, 108 |
| Chibi | 3, 4, 109 | 10, 11 | 15, 25, 25, 26 | 23, 34, 45 | 52, 57, 58, 59 | 9, 19 | 95, 108 |

985    **Fig. S3. Macaque ECoG and single-unit coverage. (A)** locations of 180-electrode ECoG grid

986    from 2 animals in the Neurotycho dataset, colors correspond to locations used for comparison

987    with single-unit timescales. **(B)** single-unit recording locations from Fig. 1a of (Murray et al.,

988    2014). **(C)** electrode indices of the sampled areas from the two animals, corresponding to those

989    colored in **(A).**

991
992 **Fig. S4. Transcriptomic PCA results. (A)** proportion of variance explained by the top 10
993 principal components (PCs) of brain-specific genes (top) and all AHBA genes (bottom). **(B)**
994 absolute Spearman correlation between timescale map and top 10 PCs from brain-specific or
995 full gene dataset. Asterisks indicate resampled significance while accounting for spatial
996 autocorrelation, **** indicate $p < 0.001$.

997
998

999
1000 **Fig. S5. Individual gene correlations from Fig. 3C with gene symbols labeled, and**
1001 **grouped into functional families.**

1002
**Fig. S6. Spectral correlates of working memory performance. (A)** difference between delay
and baseline periods for 1/f-exponent, timescale (same as main Fig. 4C but absolute units on y-
axis, instead of percentage), theta power, and high-frequency power. **(B)** Spearman correlation
between spectral feature difference and working memory accuracy across participants, same
features as in **(A)**. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$ in **(A, B)**. **(C)** scatter plot of other
significantly correlated spectral features from **(B)**.
1009

1010

**Fig. S7. Parameter sensitivity for timescale-aging analysis. (A)** cortex-averaged timescale is
independent of parcel coverage across participants. **(B)** sensitivity analysis for the number of
valid parcels a participant must have in order to be included in analysis for main Fig. 4E (red).
As threshold increases (more stringent), fewer participants satisfy the criteria (right) but
correlation between participant age and timescale remains robust (left). **(C)** sensitivity analysis
for the number of valid participants a parcel must have in order to be included in analysis for
main Fig. 4F. As threshold increases (more stringent), fewer parcels satisfy the criteria (right)
but average correlation across all parcels remains robust (left, error bars denote s.e.m of
distribution as in Fig. 4F).

**Table S1. Summary of open-access datasets used**

| Data | Ref. | Specific Source/ Format Used | Relevant Figures |
|---|---|---|---|
| MNI Open iEEG Atlas | (Frauscher et al., 2018a, 2018b) | | Fig. 2A,B, Fig. 3, Fig. 4 |
| Tw1/T2w map Human Connectome Project | (Glasser and Van Essen, 2011; Glasser et al., 2016) | Release S1200, March 1, 2017 | Fig. 2A,B, Fig. 3D-F |
| Neurotycho macaque ECoG | (Nagasaka et al., 2011; Yanagawa et al., 2013) | Anesthesia datasets, propofol and ketamine (Chibi and Geroge) | Fig. 2C,D |
| Macaque single-unit timescales | (Murray et al., 2014) | Fig. 1 of reference | Fig. 2C,D |
| Whole-cortex interpolated Allen Brain Atlas human gene expression | (Gryglewski et al., 2018; Hawrylycz et al., 2012) | Interpolated maps downloadable from http://www.meduniwien.ac.at/neuroimaging/mRNA.html | Fig. 3 |
| Single-cell timescale-related genes | (Bomkamp et al., 2019; Tripathy et al., 2017) | Table S3 from (Tripathy et al., 2017), Online Table 1 from (Bomkamp et al., 2019) | Fig. 3C,D |
| Human working memory ECoG | (Johnson, 2018, 2019; Johnson et al., 2018a, 2018b) | CRCNS fcx-2 and fcx-3 | Fig. 4A-D |

1021
1022
1023

1024 **Table S2. Reproducing figures from code repository**

| All IPython notebooks: https://github.com/rdgao/field-echos/tree/master/notebooks | |
| --- | --- |
| **Notebook** | **Results** |
| 2a_sim_method_schematic.ipynb | **simulations:** Fig. 1B-E, Fig. S3 |
| 2b_viz_NeuroTycho-SU.ipynb | **macaque timescales:** Fig. 2C,D |
| 3_viz_human_structural.ipynb | **human timescales vs. T1w/T2w and gene expression:** Fig. 2A,B, Fig. 3, Fig. S1, S4, S5; Table. S3 |
| 4b_viz_human_wm.ipynb | **human working memory:** Fig. 4A-D, Fig. S6 |
| 4a_viz_human_aging.ipynb | **human aging:** Fig. 4E,F, Fig. S7 |
| supp_spatialautocorr.ipynb | **spatial autocorrelation-preserving nulls:** Fig. S2 |
| Projection of T1w/T2w and gene expression maps from MNI volumetric coordinates to HCP-MMP1.0 can be found: https://github.com/rudyvdbrink/Surface_projection | |

 **Table S3. Significant items from brain-specific GOEA (Fig. 3F)**

| gene association | ID | e/p | ontology | name | enrichment ratio | p-value (FDR-adjusted) |
|---|---|---|---|---|---|---|
| all | GO:0034702 | e | CC | ion channel complex | 1.959 | 0.008 |
| all | GO:1902495 | e | CC | transmembrane transporter complex | 1.91 | 0.008 |
| all | GO:1990351 | e | CC | transporter complex | 1.91 | 0.008 |
| all | GO:0098982 | e | CC | GABA-ergic synapse | 2.497 | 0.038 |
| all | GO:1902711 | e | CC | GABA-A receptor complex | 4.541 | 0.038 |
| all | GO:0034707 | e | CC | chloride channel complex | 3.385 | 0.038 |
| pos | GO:0008195 | e | MF | phosphatidate phosphatase activity | 13.864 | 0.007 |
| neg | GO:0098660 | e | BP | inorganic ion transmembrane transport | 2.515 | 0.002 |
| neg | GO:0098662 | e | BP | inorganic cation transmembrane transport | 2.529 | 0.007 |
| neg | GO:0098655 | e | BP | cation transmembrane transport | 2.439 | 0.007 |
| neg | GO:0034220 | e | BP | ion transmembrane transport | 2.057 | 0.03 |
| neg | GO:0030001 | e | BP | metal ion transport | 2.239 | 0.036 |
| neg | GO:0071805 | e | BP | potassium ion transmembrane transport | 3.122 | 0.036 |
| neg | GO:0006813 | e | BP | potassium ion transport | 3.081 | 0.037 |
| neg | GO:1902495 | e | CC | transmembrane transporter complex | 2.334 | 0.009 |
| neg | GO:1990351 | e | CC | transporter complex | 2.334 | 0.009 |
| neg | GO:0034702 | e | CC | ion channel complex | 2.36 | 0.009 |
| neg | GO:0098796 | e | CC | membrane protein complex | 2.063 | 0.009 |
| neg | GO:0034703 | e | CC | cation channel complex | 2.379 | 0.03 |
| neg | GO:0005244 | e | MF | voltage-gated ion channel activity | 3.081 | 0.002 |
| neg | GO:0022832 | e | MF | voltage-gated channel activity | 3.081 | 0.002 |
| neg | GO:0046873 | e | MF | metal ion transmembrane transporter activity | 2.453 | 0.002 |
| neg | GO:0022890 | e | MF | inorganic cation transmembrane transporter activity | 2.24 | 0.005 |
| neg | GO:0005216 | e | MF | ion channel activity | 2.289 | 0.006 |
| neg | GO:0008324 | e | MF | cation transmembrane transporter activity | 2.173 | 0.006 |
| neg | GO:0015318 | e | MF | inorganic molecular entity transmembrane transporter activity | 2.04 | 0.006 |
| neg | GO:0015077 | e | MF | monovalent inorganic cation transmembrane transporter activity | 2.535 | 0.006 |
| neg | GO:0015075 | e | MF | ion transmembrane transporter activity | 2.024 | 0.006 |

| neg | GO:0015079 | e | MF | potassium ion transmembrane transporter activity | 3.041 | 0.006 |
|-----|------------|---|----|--------------------------------------------------|-------|-------|
| neg | GO:0005215 | e | MF | transporter activity | 1.883 | 0.006 |
| neg | GO:0022857 | e | MF | transmembrane transporter activity | 1.906 | 0.006 |
| neg | GO:0022836 | e | MF | gated channel activity | 2.301 | 0.006 |
| neg | GO:0015267 | e | MF | channel activity | 2.191 | 0.006 |
| neg | GO:0022803 | e | MF | passive transmembrane transporter activity | 2.191 | 0.006 |
| neg | GO:0005249 | e | MF | voltage-gated potassium channel activity | 3.658 | 0.006 |
| neg | GO:0005261 | e | MF | cation channel activity | 2.353 | 0.009 |
| neg | GO:0005267 | e | MF | potassium channel activity | 3.058 | 0.011 |
| neg | GO:0022843 | e | MF | voltage-gated cation channel activity | 2.744 | 0.022 |

1033    e/p: enriched or purified; BP: biological process; CC: cellular components; MF: molecular
1034    function

1035 **Table S4. Significant items from all-gene GOEA**

| gene association | ID | e/p | ontology | name | enrichment ratio | p-value (FDR-adjusted) |
|---|---|---|---|---|---|---|
| all | GO:0034702 | e | CC | ion channel complex | 1.83 | 0.008 |
| all | GO:1990351 | e | CC | transporter complex | 1.774 | 0.008 |
| all | GO:1902495 | e | CC | transmembrane transporter complex | 1.79 | 0.008 |
| all | GO:0034703 | e | CC | cation channel complex | 1.952 | 0.009 |
| all | GO:0098982 | e | CC | GABA-ergic synapse | 2.468 | 0.048 |
| all | GO:1902711 | e | CC | GABA-A receptor complex | 5.035 | 0.048 |
| pos | GO:0050866 | e | BP | negative regulation of cell activation | 3.596 | 0 |
| pos | GO:0002376 | e | BP | immune system process | 1.629 | 0 |
| pos | GO:0006955 | e | BP | immune response | 1.992 | 0 |
| pos | GO:0002695 | e | BP | negative regulation of leukocyte activation | 3.343 | 0.001 |
| pos | GO:0045087 | e | BP | innate immune response | 2.297 | 0.005 |
| pos | GO:0050865 | e | BP | regulation of cell activation | 2.099 | 0.005 |
| pos | GO:0045321 | e | BP | leukocyte activation | 1.834 | 0.006 |
| pos | GO:0007165 | e | BP | signal transduction | 1.301 | 0.006 |
| pos | GO:0051250 | e | BP | negative regulation of lymphocyte activation | 3.305 | 0.007 |
| pos | GO:0070663 | e | BP | regulation of leukocyte proliferation | 2.82 | 0.007 |
| pos | GO:0002252 | e | BP | immune effector process | 1.778 | 0.009 |
| pos | GO:0050670 | e | BP | regulation of lymphocyte proliferation | 2.823 | 0.009 |
| pos | GO:0032944 | e | BP | regulation of mononuclear cell proliferation | 2.807 | 0.009 |
| pos | GO:0050776 | e | BP | regulation of immune response | 1.787 | 0.011 |
| pos | GO:0002682 | e | BP | regulation of immune system process | 1.571 | 0.015 |
| pos | GO:0046634 | e | BP | regulation of alpha-beta T cell activation | 3.772 | 0.016 |
| pos | GO:0001775 | e | BP | cell activation | 1.709 | 0.016 |
| pos | GO:0032956 | e | BP | regulation of actin cytoskeleton organization | 2.229 | 0.016 |
| pos | GO:0003150 | e | BP | muscular septum morphogenesis | 17.672 | 0.016 |
| pos | GO:0032945 | e | BP | negative regulation of mononuclear cell proliferation | 4.208 | 0.016 |
| pos | GO:0050672 | e | BP | negative regulation of lymphocyte proliferation | 4.208 | 0.016 |
| pos | GO:0006952 | e | BP | defense response | 1.686 | 0.016 |

| pos | GO:0002694 | e | BP | regulation of leukocyte activation | 2.013 | 0.016 |
|-----|------------|---|----|-----------------------------------|-------|-------|
| pos | GO:0002253 | e | BP | activation of immune response | 2.183 | 0.016 |
| pos | GO:0030833 | e | BP | regulation of actin filament polymerization | 2.832 | 0.016 |
| pos | GO:0032970 | e | BP | regulation of actin filament-based process | 2.136 | 0.017 |
| pos | GO:0002684 | e | BP | positive regulation of immune system process | 1.708 | 0.017 |
| pos | GO:0046640 | e | BP | regulation of alpha-beta T cell proliferation | 6.094 | 0.017 |
| pos | GO:0050868 | e | BP | negative regulation of T cell activation | 3.381 | 0.017 |
| pos | GO:0002274 | e | BP | myeloid leukocyte activation | 1.926 | 0.017 |
| pos | GO:0008064 | e | BP | regulation of actin polymerization or depolymerization | 2.697 | 0.017 |
| pos | GO:0030832 | e | BP | regulation of actin filament length | 2.681 | 0.017 |
| pos | GO:0006334 | e | BP | nucleosome assembly | 3.053 | 0.018 |
| pos | GO:0070664 | e | BP | negative regulation of leukocyte proliferation | 3.956 | 0.018 |
| pos | GO:0038096 | e | BP | Fc-gamma receptor signaling pathway involved in phagocytosis | 3.787 | 0.026 |
| pos | GO:0002433 | e | BP | immune response-regulating cell surface receptor signaling pathway involved in phagocytosis | 3.787 | 0.026 |
| pos | GO:0098883 | e | BP | synapse pruning | 10.041 | 0.027 |
| pos | GO:0038094 | e | BP | Fc-gamma receptor signaling pathway | 3.734 | 0.029 |
| pos | GO:0051249 | e | BP | regulation of lymphocyte activation | 2.035 | 0.029 |
| pos | GO:0002431 | e | BP | Fc receptor mediated stimulatory signaling pathway | 3.682 | 0.03 |
| pos | GO:0042116 | e | BP | macrophage activation | 4.734 | 0.03 |
| pos | GO:0110053 | e | BP | regulation of actin filament organization | 2.279 | 0.03 |
| pos | GO:0150064 | e | BP | vertebrate eye-specific patterning | 22.09 | 0.03 |
| pos | GO:0002683 | e | BP | negative regulation of immune system process | 2.008 | 0.03 |
| pos | GO:0051049 | e | BP | regulation of transport | 1.428 | 0.03 |
| pos | GO:0098542 | e | BP | defense response to other organism | 1.811 | 0.033 |
| pos | GO:0150146 | e | BP | cell junction disassembly | 9.204 | 0.033 |
| pos | GO:0016322 | e | BP | neuron remodeling | 9.204 | 0.033 |
| pos | GO:1903038 | e | BP | negative regulation of leukocyte cell-cell adhesion | 3.04 | 0.033 |

| | | | | | | |
|---|---|---|---|---|---|---|
| pos | GO:0007166 | e | BP | cell surface receptor signaling pathway | 1.391 | 0.034 |
| pos | GO:0034728 | e | BP | nucleosome organization | 2.591 | 0.037 |
| pos | GO:0036336 | e | BP | dendritic cell migration | 6.976 | 0.037 |
| pos | GO:0048584 | e | BP | positive regulation of response to stimulus | 1.379 | 0.039 |
| pos | GO:0001774 | e | BP | microglial cell activation | 5.727 | 0.039 |
| pos | GO:0002269 | e | BP | leukocyte activation involved in inflammatory response | 5.727 | 0.039 |
| pos | GO:0050778 | e | BP | positive regulation of immune response | 1.82 | 0.039 |
| pos | GO:2000112 | p | BP | regulation of cellular macromolecule biosynthetic process | 0.718 | 0.017 |
| pos | GO:0051252 | p | BP | regulation of RNA metabolic process | 0.718 | 0.019 |
| pos | GO:0044271 | p | BP | cellular nitrogen compound biosynthetic process | 0.554 | 0.029 |
| pos | GO:0019219 | p | BP | regulation of nucleobase-containing compound metabolic process | 0.737 | 0.029 |
| pos | GO:0090304 | p | BP | nucleic acid metabolic process | 0.654 | 0.037 |
| pos | GO:0032993 | e | CC | protein-DNA complex | 2.829 | 0.016 |
| pos | GO:0000786 | e | CC | nucleosome | 3.488 | 0.016 |
| pos | GO:0005887 | e | CC | integral component of plasma membrane | 1.555 | 0.016 |
| pos | GO:0031226 | e | CC | intrinsic component of plasma membrane | 1.536 | 0.016 |
| pos | GO:0044815 | e | CC | DNA packaging complex | 3.217 | 0.023 |
| pos | GO:0030666 | e | CC | endocytic vesicle membrane | 2.705 | 0.034 |
| pos | GO:0031514 | e | CC | motile cilium | 2.897 | 0.034 |
| pos | GO:0043235 | e | CC | receptor complex | 1.981 | 0.04 |
| pos | GO:0000839 | e | CC | Hrd1p ubiquitin ligase ERAD-L complex | 11.045 | 0.047 |
| pos | GO:0016021 | e | CC | integral component of membrane | 1.232 | 0.047 |
| pos | GO:0005634 | p | CC | nucleus | 0.79 | 0.04 |
| pos | GO:0003676 | p | MF | nucleic acid binding | 0.694 | 0.012 |
| neg | GO:0006813 | e | BP | potassium ion transport | 2.911 | 0.004 |
| neg | GO:0071805 | e | BP | potassium ion transmembrane transport | 2.868 | 0.008 |
| neg | GO:0015079 | e | MF | potassium ion transmembrane transporter activity | 2.888 | 0.001 |
| neg | GO:0015075 | e | MF | ion transmembrane transporter activity | 1.726 | 0.001 |
| neg | GO:0022857 | e | MF | transmembrane transporter activity | 1.649 | 0.001 |

| | | | | | | |
|---|---|---|---|---|---|---|
| neg | GO:0046873 | e | MF | metal ion transmembrane transporter activity | 2.068 | 0.001 |
| neg | GO:0005215 | e | MF | transporter activity | 1.587 | 0.002 |
| neg | GO:0022832 | e | MF | voltage-gated channel activity | 2.468 | 0.006 |
| neg | GO:0005244 | e | MF | voltage-gated ion channel activity | 2.468 | 0.006 |
| neg | GO:0015318 | e | MF | inorganic molecular entity transmembrane transporter activity | 1.662 | 0.008 |
| neg | GO:0001227 | e | MF | DNA-binding transcription repressor activity, RNA polymerase II-specific | 2.227 | 0.011 |
| neg | GO:0001217 | e | MF | DNA-binding transcription repressor activity | 2.218 | 0.011 |
| neg | GO:0022836 | e | MF | gated channel activity | 2.007 | 0.015 |
| neg | GO:0005249 | e | MF | voltage-gated potassium channel activity | 3.126 | 0.015 |
| neg | GO:0015077 | e | MF | monovalent inorganic cation transmembrane transporter activity | 1.916 | 0.022 |
| neg | GO:0005267 | e | MF | potassium channel activity | 2.703 | 0.022 |
| neg | GO:0022890 | e | MF | inorganic cation transmembrane transporter activity | 1.701 | 0.033 |

1036 e/p: enriched or purified; BP: biological process; CC: cellular components; MF: molecular
1037 function