Title:

Changes in arthropod community but not plant quality benefit a specialist herbivore on plant under

reduced water availability

Author

1.   Lin, Po-An*, pbl5066@psu.edu, Department of Entomology, Pennsylvania State University,

State College, PA (**Corresponding author**)

2.   Liu, Chia-Ming*, cobras770920@hotmail.com, Graduate School of Life and Environmental

Sciences, University of Tsukuba, Ibaraki, Japan

3.   Ou, Jia-Ang*, r06b44029@ntu.edu.tw, Institute of Ecology and Evolutionary Biology, National

Taiwan University, Taipei, Taiwan

4.   Sun, Cheng-Han, r06b44010@ntu.edu.tw, Institute of Ecology and Evolutionary Biology,

National Taiwan University, Taipei, Taiwan

5.   Chuang, Wen-Po, wenpo@ntu.edu.tw, Department of Agronomy, National Taiwan University,

Taipei, Taiwan

6.   Ho, Chuan-Kai, ckho@ntu.edu.tw, Institute of Ecology and Evolutionary Biology, National

Taiwan University, Taipei, Taiwan

7.   Kinoshita, Natsuko, kinoshita.natsuko.gf@u.tsukuba.ac.jp, Department of Agro-Bioresources

Science and Technology, Faculty of Life and Environmental Sciences, University of Tsukuba,

Ibaraki, Japan

8.   Felton, Gary W., gwf10@psu.edu, Department of Entomology, Pennsylvania State University,

State College, PA

* these authors contributed equally to the work.

We show that low water availability benefits certain insects through reducing negative species

interactions, and highlight the importance of community factors in plant-insect-environment

interactions.

Declaration of authorship

PAL, KN, CKH, WPC, and GWF conceived the ideas, designed methodology. PAL and JAO

analyzed the data. PAL, CML, and CHS collected the data. PAL, CML, JAO, KN, CKH, WPC, and

GWF wrote the manuscript.

Abstract

Plants grow under reduced water availability can have divergent effects on insect herbivores, in some

instances producing benefits to them. However, the forces mediating these positive impacts remain

mostly unclear. We conducted a manipulative field study using a specialist herbivore *Pieris rapae*,

and its host plant, *Rorippa indica*, in two populations to identify how water availability impacts

overall plant quality and multitrophic interactions. We observed that *R. indica* growing under low

water availability led to higher survival of *P. rapae* larvae. The increase in survival of eggs and larvae

was related to the reduced abundance of other herbivores and natural enemies. Water availability had

differential impacts on members of the herbivore community through changes in plant quality. Low

water availability decreased the quality of *R. indica* to most herbivores as indicated by reduced

abundance in the field and decreased relative growth rate in feeding assays. In contrast, the

performance of *P. rapae* larvae were not affected by differences in sympatric *R. indica* grown under

different water availability. These results indicate that local *P. rapae* possess some physiological

adaptation to overcome fluctuations in host quality. Our findings illustrate that reduced water

availability is beneficial to a specialist herbivore, but detrimental to most other herbivores. Our work

highlights the complex roles of the arthropod communities associated with plants in determining the

impacts of water availability on insect herbivores.

Key words

48 drought, environmental heterogeneity, predation, competition, insect herbivore

Introduction (725)

Water availability is a vital factor influencing individuals, populations, and communities of plants

(Bunker and Carson 2005; Schulze 1986) and ultimately heterotrophic organisms across trophic levels

(Anderegg et al. 2013). Its importance in affecting ecological interactions will likely increase due to

the predicted increase in frequency and severity of drought under current models of climate change (Li

et al. 2009). In addition to large-scale climatic factors, fine-scale topographic heterogeneity also

influences water availability to individual plants, even when the overall climatic conditions are similar

(Murren et al. 2020). This is particularly the case for plants in urban and agricultural environments,

because conditions such as the reduction in tree canopy coverage, increased in light

intensity/temperature, and shallow soil depth all contribute to high evaporation and low water

availability (Dambros et al. 2013; Pickett et al. 2011). Given the many dimensions that influence plant

water access, the variation in plant phenotype due to changes in water availability is likely a key factor

influencing the dynamics and survival of associated insect herbivores.

The influence of water availability on insect herbivores has receive much attention in past literature. A

classic example is the relationship between drought and outbreak of insect herbivores (Mattson and

Haack 1987; White 1974; White 1969). This observation has led to a fruitful area of research and it is

now clear that the impact of water availability on insect herbivores and plants is context-dependent

and varies across different systems (Jamieson et al. 2012; White 2009). However, the main factors that

contribute to this positive effect of low water availability on some insect herbivores remain unclear. A

major limitation in past literature that contribute to this uncertainty was the emphasis on the direct

interaction between plants and insects (English-Loeb et al. 1997). In contrast to a beneficial impact

observed in natural conditions, low water availability generally elicits negative effects on plant quality

(defined as a performance response of insect herbivores, see (Awmack and Leather 2002)) especially

in herbaceous plants (Waring and Cobb 1992). It is likely that factors conttributing to these beneficial

impacts are interactions apart from the direct interaction between the focal herbivore and its host plant.

Low water availability has been documented to reduce the abundance and composition of arthropods

associated with plants (Trotter et al. 2008), and can potentially alter the strength and direction of

multispecies interactions. Changes in arthropod species interactions (e.g., competition, predation)

could affect the fitness of insect herbivores, and have been shown to contribute more to the overall

impacts of environmental changes than the effect of pairwise interactions between plants and insect

herbivores (Ockendon et al. 2014). Although important, few studies investigating the influence of

multitrophic interactions on insect herbivores feeding on plant with different water availability have

been performed (Jamieson et al. 2012).

In this study, we aimed to evaluate how water availability alters the performance of a focal insect

species through direct effects on plant quality and indirect effects mediated by changes in other

arthropods. Our principal hypothesis was that although decreased water availability would reduce the

quality of herbaceous plants to insect herbivores, decline in natural enemies and potential competitors

could be a factor that positively influences insect herbivores. To test our hypothesis, we conducted a

manipulative field study using a wild host, *Rorippa indica* (L.) Hiern, of the specialist herbivore,

cabbage white butterfly (*Pieris rapae*) from two populations in Taiwan and Japan. *R. indica* is a

perennial glabrous herb that is commonly found in highly disturbed areas such as roadsides, gardens,

and field margins throughout the main island of Taiwan (Huang 1996) and also during May to

September in most parts of Japan (Jisaburo□ O□i 1965). *R. indica* individuals with different

phenotypes due to water availability are commonly observed in the field. By manipulating water status

of *R. indica* in the field, in combination with feeding assays in the lab, we aimed to determine (1) the

responses (e.g., preference and performance) of insect herbivores to *R. indica* of different water status,

and (2) the factors (e.g., plant quality and species interactions) that contribute to these responses.

Materials and methods (1423)

Insect and plant

Cabbage white butterflies, *Pieris rapae*, were collected in the experimental field of National Taiwan

University (NTU, Taipei, Taiwan) and University of Tsukuba (UT, Ibaraki, Japan), and kept at 14:10

light:dark cycle, 25-27°C, on cabbage (*Brassica oleracea* var. capitata) and Komatsuna (*B. rapa* var.

Perviridis) respectively. North American *P. rapae* were collected on campus at the Pennsylvania State

University (PSU, University Park, PA, USA) and maintained on kale (*B. oleracea* var. sabellica).

Cabbage looper*, Trichloplusia ni*, corn earworm, *Helicoverpa zea*, and fall armyworm, *Spodoptera*

frugiperda were purchased from Benzon Research (Carlisle, PA, USA) and reared on artificial diet

(Peiffer and Felton 2005). All insects were kept under the same condition described above. Seeds of

Rorippa indica were collected in the experimental fields. Plants were maintained in semi-closed

greenhouse and grown in 3-inch pots (400 mL) with a mixture of peat moss (Kekkila, Finland): perlite

(2-8 mm) (3:1 volume) supplied with fertilizer Osmocote Plus (Everris, Netherland). The water

holding capacity of peat moss was estimated as 195 mL (65% of volume) based on information

provided by Kekkila.

Water treatment and water status quantification

Plants with 7-8 fully expanded leaves were subjected to two levels of water treatments: (1) high water

availability (200 mL day^{-1} pot^{-1}), and (2) low water availability (10 mL day^{-1} pot^{-1}). The water levels

were chosen based on morphological comparisons of plants from a series of water availabilities (10,

30, 50, 100, and 200 mL day^{-1} pot^{-1}) to plants in field growing under moist and dry area. Water was

applied daily at 9 a.m. for 7 days. To quantify the water status of plants (Jones 2007), we measured

shoot biomass, root biomass, shoot water content, and volumetric water content of soil (θ_v). For shoot

biomass and water content, the entire above ground tissue was collected, weighed, dry (70 ℃, 48 h),

and weighed again. For root biomass, below ground tissue was collected after the above ground tissue,

roots were washed and dried (70 ℃, 48 h), and weighed to obtain dry mass. For soil water content,

100 cm^3 of soil sample was collected from each pot avoiding root tissues. Soil was weighed and dried

(70 ℃, 48 h), and weighed again. The water content was calculated using the following formula:

$(\text{mass of water}_{(g)}\ \text{density of soil}_{(g\ cm^{-3})})(\text{mass of soil}_{(g)}\text{density of water}_{(g\ cm^{-3})})^{-1}$. All samples

were collected 3 h after daily water application. For field experiments, water status of *R. indica* was

maintained by the same procedure, accompanied with daily observations at 12, 2, and 4 p.m. to

provide more water if the low water availability treatments exhibited low turgor. When signs of low

turgor were observed, 10 mL of water was added. This procedure prevented most leaves from

irreversible tissue damage. Loss of turgor was not observed for plants growing under high water

availability. The phenotypes were maintained as consistent as possible throughout the field experiment.

(Appendix 1).

Oviposition preference, survival, and relative growth rate of insect herbivores

Individual *R. indica* grown under different water status (in pots) were placed in the experimental field

at 7 a.m., 3 m apart in a square grid (Appendix 2). Ovipositional preferences were determined by

recording the number of eggs of *P. rapae* after the first 72 h. Individual plants were kept in the field

for an additional 7 days before *P. rapae* larvae were collected. Only larvae older than second instar

were collected to ensure that the collected caterpillars came from the recorded eggs. Because *P. rapae*

larvae rarely move away from their host even when the host plant is stressed (based on our rearing

observations) and high levels of egg predation, the survival of *P. rapae* was defined and calculated as:

(caterpillar number) (egg number)$^{-1}$. See "Method: Arthropod community" for additional details on

field experiments.

We analyzed the relative growth rate (RGR) of larvae to determine the quality of *R. indica* to

caterpillars under different water treatments. Larvae of *P. rapae* from Taiwan and Japan were fed with

R. indica from sympatric population. Third instar larvae of *P. rapae* were weighed and placed in 30

c.c. plastic cups with thin layer of 1% agar (5 mL). The larvae were fed with detached mature leaves

of specific water status (either 200 mL day^{-1}, 10 mL day^{-1}, and an additional intermediate water level:

50 mL day^{-1}). Leaves were replaced daily to minimize the potential changes in quality caused by

excision. Larvae were weighed after 3 days. The start and end weights were used to calculate RGR

using the equation: (end weight – start weight) (end weight + start weight)$^{-1}$ days^{-1} (Felton et al. 1989).

We conducted another set of common garden experiment to test the changes in host quality using

insects without previous evolutionary history with *R. indica* from Taiwan and Japan. Plants were

grown in a greenhouse in PSU with 16:8 light cycle at $25°C$. Leaves were collected from two different

water treatments (200 mL day^{-1}, 10 mL day^{-1}) from both populations of *R. indica* and fed to third

instar larvae of caterpillars from North America, including *P. rapae* (specialist), *Trichloplusia ni*

(semi-specialist, cabbage looper), *Helicoverpa zea* (generalist, corn earworm), and *Spodoptera*

frugiperda (generalist, fall armyworm) using method as described above.

Arthropod community

The insect community associated with *R. indica* was determined on the same day by visual

observation. All arthropods on each plant during 7 a.m.-12 p.m. were recorded. All morphospecies

observed in the field were listed in Appendix 3. Field experiments took place in Taiwan during April –

June 2018, and in Japan during July – September 2018. The experiment was replicated three times in

both locations, with one trial in each month, defined as early, middle, and late season. The

experimental fields were located within the campus of NTU (25°00'55.9"N 121°32'25.0"E) and UT

(36°06'51.7"N 140°05'59.7"E), and both were at an elevation of about 900 m^2. The natural community

of plants in the experimental fields was allowed to grow undisturbed. Each trial contained 20-25

individual *R. indica* from each water treatment (total of 40-50 plants). Timing of field experiments

were determined according to governmental weather forecasts in Taiwan and Japan. The experiments

were conducted during a period of 10 days that has the lowest probability of rain in each season. The

incidences of raining were rare, on average 0-1 days per field experiment. In case of rain, plants were

covered with transparent plastic bins and removed after the rain. In addition, all plants were placed on

a round plastic tray to prevent moisture from the soil. To determine parasitism incidence, larvae of *P.*

rapae collected from the field were reared individually using *Brassica oleracea* var. capitata and

Brassica rapa var. Perviridis as food in Taiwan and Japan respectively until pupation. If the caterpillar

died before pupation, the carcass was dissected to determine presence of parasitoids.

Statistical analysis

All statistical analyses were conducted in R version 3.5.3 (R Core Team 2017). Univariate response

variables were modeled with generalized linear models (GLM) with treatment, population, season and

their interaction as fixed effects. We analyzed all count data (egg, caterpillar, herbivore, and natural

enemy) with negative binomial GLM (package: *MASS*) (Venables and Ripley 2013) with biomass of

plants as an offset variable. Survival probabilities of *P. rapae* were modeled with binomial GLM as

the odds ratio between alive (caterpillar count) and dead (egg count - caterpillar count). We excluded

samples with no eggs observed. Dispersion, zero-inflation, residual normality, and homoscedasticity

of all model fits were evaluated with simulation-based residual plots (package: *DHARMa*) (Hartig

2017). Significance of model parameters were tested by Type II rather than Type III Wald's

Chi-squared test (package: *car*) (Fox and Weisberg 2018) to account for unbalanced data (Langsrud

2003).

To examine whether the survival of *P. rapae* could be attributed to abundance of other arthropods, we

pooled samples between treatments and fitted binomial generalized linear mixed-effect models

(GLMM, package: *glmmTMB*) (Brooks et al. 2017) with observation-level random effect to improve

residuals. Abundances of trophic guilds (herbivore and natural enemy, Fig. 4), and the abundances of

individual species (Table 2) were used as fixed effect predictors. Individual GLMM was fitted for

each season in each location. We implemented backwards model selection (likelihood ratio tests) until

all terms were significant (Zuur et al. 2009). Other model diagnostics are performed as described

above. Continuous variables such as water content, biomass, and relative growth rate were analyzed

using ANOVA, followed by Tukey HSD. Diagnostic plots were used to check for model assumptions,

such as equal variance and normality (Zuur et al. 2010). Indicator species analysis was conducted to

ascertain whether specific species were associated with a particular water treatment in a particular

season (packages: *indicspecies* and *vegan*) (Cáceres and Legendre 2009; Oksanen et al. 2010). We

conducted indicator species analysis on presence absence data to partially remove the effects of the

difference in absolute abundance among species as we were more interested in whether each

individual species showed any association towards a particular treatment-season combination.

Permutation test (with 999 permutations) were used to assess the significance of associations between

species and treatment-season combinations. In addition, we applied PCA on log(x+1)-transformed

(improve skew of species abundance distributions) and z-standardized (put equal weights on each

species) species abundances independently for herbivores and natural enemies to help visualize the

changes in species composition between treatments and seasons.

Results (1709)

Water status quantification

Different water treatments led to observable changes in plant phenotype (please see Fig. 1a). Leaf

number and shape were not affected by water availability (Fig. 1b); however, leaf shape differed

between populations (Fig. 1c, $F_{1,29} = 13.03$, $P = 0.001$). Fresh weight of shoot of individuals was

reduced under low water availability (Fig. 1d, $F_{1,29} = 65.29$, $P < 0.001$) and higher in the Taiwanese

population compared to the Japanese population ($F_{1,29} = 22.59$, $P < 0.001$). Dry weight was also

reduced under low water availability (Fig. 1e, $F_{1,29} = 17.05$, $P < 0.001$) and higher in the Taiwanese

population ($F_{1,29} = 8.5$, $P = 0.007$). Water content of shoot was reduced under low water availability

(Fig. 1f, $F_{1,28} = 105.02$, $P < 0.001$), and higher in Taiwanese population ($F_{1,28} = 5.48$, $P = 0.027$).

However, the effect of population was only significantly reduced under low water availability,

indicated by a significant interaction between water and population ($F_{1,28} = 5.48$, $P = 0.027$). The

biomass of root was not affected by water treatments. However, root biomass was higher in Taiwanese

population (Fig. 1g, $F_{1,28}$ = 8.605, P = 0.007). Root: shoot ratio was increased under low water

availability (Fig. 1h, $F_{1,28}$ = 88.17, P < 0.001). Volumetric water content (θ_v) of the soil was 0.53-0.55

for high water availability and 0.09 for low water availability (Fig. 1i, $F_{1,29}$ = 2238.65, P < 0.001).

Oviposition preference, caterpillar number, and survival

Higher quantities of eggs were observed on *R. indica* under high water availability (Fig. 2a, GLM,

water: 200 mL day^{-1}, estimate = 1.04, Z = 4.79, P < 0.001). Egg number on *R. indica* was higher in

Taiwan than in Japan (GLM, region: Taiwan, estimate = 2.2, Z= 10.21, P < 0.001). In addition, egg

number was lower during late season in Taiwan (GLM, season*region: late*Taiwan, estimate = -1.74,

Z= -7.11, P < 0.001).

The number of caterpillars on *R. indica* was overall higher in Taiwan (Fig. 2b, GLM, region: Taiwan,

estimate = 1.25, Z= 3.57, P < 0.001), but lower on *R. indica* under high water availability in Taiwan

(GLM, water*region: 20 mL day^{-1}*Taiwan, estimate = -0.94, Z= -2.66, P = 0.008). The number of

caterpillars were lower during late season (GLM, season: late, estimate = -1.72, Z = -4.19, P < 0.001),

and not affected by water treatment in general.

Survival of *P. rapae* was lower on *R. indica* under high water availability (Fig. 2c, GLM, water: 200

mL day^{-1}, estimate = -2.15, Z = -8.79, P < 0.001), middle, and late seasons (GLM, season: middle,

estimate = -0.63, Z = -2.58, P = 0.01; season: late, estimate = -3.96, Z = -14.95, P < 0.001). However,

the pattern was influenced by several interacting factors. Survival of *P. rapae* was higher on *R. indica*

under high water availability in middle and late season (GLM, water*season: 200 mL day^{-1}*middle,

estimate = 1.57, Z = 10.1, P < 0.001; water*season: 200 mL day^{-1}*late, estimate = 1.31, Z = 6.52, P <

0.001), but lower on *R. indica* under high water availability in Taiwan (GLM, water*region: 200 mL

day^{-1}*late, estimate = -0.53, Z = -2.34, P = 0.019). The survival of *P. rapae* was overall lower in

Taiwan (GLM, location: Taiwan, estimate = -2.74, Z = -10.81, P < 0.001), but significantly higher in

late season (GLM, season*location: late*Taiwan, estimate = 4.2, Z = 17.49, P < 0.001) and lower in

middle season (GLM, season*location: middle*Taiwan, estimate = -0.53, Z = -2.34, P = 0.019).

Relative growth rate of caterpillars

The relative growth rates (RGR) of *P. rapae* larvae were not affected by water status of *R. indica* from

the same geographic origin (Fig. 3a, b). In contrast, populations of *R. indica* from Taiwan and Japan

growing under low water availability led to significant decreases in RGR of *P. rapae* from North

America (Fig. 3c, $F_{1,141}$ = 12.96, P < 0.001). *R. indica* growing under low water availability decreased

RGR of generalist caterpillars. RGR of *H. zea* larvae were lower on *R. indica* growing under low

water availability (Fig. 3d, $F_{1,147} = 25.41$, $P < 0.001$). RGR of *T. ni* larvae were lower on *R. indica*

growing under low water availability (Fig. 3e, $F_{1,128} = 8.76$, $P = 0.004$) and *R. indica* from Taiwan

($F_{1,128} = 8.62$, $P = 0.004$). However, the negative impacts of low water availability were only

significant when *T. ni* larvae fed on *R. indica* from Taiwan ($F_{1,128} = 5.2$, $P = 0.024$). *S. frugiperda*

completely rejected *R. indica* from Taiwan as food source, and we observed a small amount of feeding

on *R. indica* from Japan. The mortality of *S. frugiperda* larvae were significantly lower when *R. indica*

from Taiwan was provided as food (Fig. 3f, GLM, 200 mL day^{-1}: estimate = -1.81, $P < 0.001$).

Abundance and composition of herbivores and natural enemies

Herbivore abundance (Fig. 4a and Fig. 4i) on *R. indica* was higher on *R. indica* growing under high

water availability (GLM, water: 200 mL day^{-1}, estimate = 1.43, $Z = 4.32$, $P < 0.001$), and lower in late

season (GLM, season: late, estimate = -2.61, $Z = -6.54$, $P < 0.001$). The negative association between

herbivore abundance and late season was especially stronger on *R. indica* growing under high water

availability (GLM, water*season: 200 mL day^{-1*}late, estimate = -1.45, $Z = -3.39$, $P < 0.001$), and

became positive during late season in Taiwan (GLM, season*location: late*Taiwan, estimate = 2.69, Z

$= 6.23$, $P < 0.001$).

Natural enemy abundance (Fig. 4e and Fig. 4m) was higher on *R. indica* growing under high water

availability (GLM, water: 200 mL day^{-1}, estimate = 1.98, $Z = 4.21$, $P < 0.001$), but lower during

middle and late season (GLM, season: middle, estimate = -1.54, $Z = -2.28$, $P = 0.022$; season: late,

estimate = -2.04, $Z = -3.26$, $P = 0.001$). Natural enemy abundance was lower in Taiwan compared to

Japan (GLM, region: Taiwan, estimate = -1.84, $Z = -3.24$, $P = 0.001$), but increased drastically in

middle and late season in Taiwan (GLM, season*location: middle*Taiwan, estimate = 2.51, $Z = 5.04$,

$P < 0.001$; season*location: late*Taiwan, estimate = 4.49, $Z = 8.61$, $P < 0.001$).

The composition of invertebrate herbivores and natural enemies associated with *R. indica* changed as

a result of water availability and season. In Taiwan, there was a clear separation of herbivores on *R.*

indica growing under high and low water availability (Fig. 5a). Indicator species analysis revealed that

several specialist herbivores associated significantly with *R. indica* growing under high water

availability (Table 1). In contrast, generalist herbivores had a weaker association with *R. indica*

growing under high water availability. Grasshoppers (Family: Catantopidae, Order: Orthoptera) were

the only generalist that associated significantly with *R. indica* growing under high water availability in

Japan. The only morphospecies associated with *R. indica* growing under low water availability was

spider mites (Tetranychidae) in Taiwan (Table 1). In addition to water treatment, the composition of

herbivores also changed across seasons in Taiwan. Both PCA (Fig. 5b) and indicator species analysis

(Table 1) indicated no strong association between *R. indica* and natural enemies under low water

availability. The association between natural enemies and *R. indica* were stronger in later seasons in

Taiwan under high water availability (Table 1). Many of the natural enemies we observed attacked *P.*

rapae. Based on field observations, several species were observed to cause high mortality of eggs and

larvae of *P. rapae*, such as ants (Formicidae), predatory stink bug (*Eocanthecona concinna*), ladybugs

(Coccinellidae), and spiders (Araneae). One of the surprising finding in Taiwan was a less than 1%

parasitism rate by specialist parasitoid wasp, *Cotesia glomerata*, of *P. rapae* larvae on *R. indica*

compared to *P. rapae* larvae in nearby cabbage fields where parasitism was regularly observed.

In Japan, the separation of herbivore composition between *R. indica* growing under different water

availability was less apparent (Fig. 5c). Based on indicator species analysis, most herbivores were

significantly associated with *R. indica* growing under high water availability (Table 1). Season also

influenced the composition of herbivore on *R. indica*. Only scarlet shieldbug (*E. dominulus*) and

grasshoppers (Catantopidae) sometimes caused noticeable damage, but the level of herbivory in the

field was not comparable to the level of herbivory in Taiwan. For natural enemy compositions, there

were no observable differences between *R. indica* growing under different water availability (Fig. 5d).

According to indicator species analysis some natural enemies were significantly associated with *R.*

indica growing under high water availability in early season (Table 1). However, we did not observe

any predation on *P. rapae* throughout all field experiments.

Survival of *Pieris rapae* and arthropod community

We found significant negative associations between the abundance of herbivores and natural enemies

with the survival of *P. rapae* during several seasons in both locations (Fig. 4). In Taiwan, abundance

of herbivore was negatively associated with the survival of *P. rapae* during early (Fig. 4b, GLMM,

estimate = -1.75, Z = -5.7, $P < 0.001$) and middle season (Fig. 4c, GLMM, estimate = -0.12, Z = -2.41,

$P = 0.016$). Abundance of natural enemy was negatively associated with survival of *P. rapae* during

early (Fig. 4f, GLMM, estimate = -3.64 Z = -10.11, $P < 0.001$) and late season (Fig. 4h, GLMM,

estimate = -0.24, Z = -4.05, $P < 0.001$).

Most of the herbivores observed on *R. indica* showed negative associations with *R. indica* survival

(Table 2). For instance, flea beetles (*P. stiolata*) that heavily attacked well-watered *R. indica* were

negatively associated with survival of *P. rapae* during early season in Taiwan (GLMM, estimate =

-6.50, $P = 0.04$). Two herbivores were found to be positively associated with survival of *P. rapae*,

including scarlet shieldbugs (*E. dominulus*) and Cicadellidae during middle and late season in Taiwan

respectively (GLMM, estimate = 0.99, p = 0.08; GLMM, estimate = 1.31, $P < 0.001$). However, the

scarlet shieldbug (*E. dominulus*) had contrasting associations with the survival of *P. rapae* in different

month (negative association during late season, GLMM, estimate = -0.95, $P < 0.001$), likely due to the

difference in abundance between seasons (Individual plant^{-1}: Middle = 0.136; Late = 0.275). The

analysis of morphospecies in Japan revealed some unexpected associations; for example, Lauxaniidae,

a parasitic mite of aphids, and Syrphidae (during early season, GLMM, estimate = -0.95, $P = 0.021$;

estimate = 0.50, p = 0.001; estimate = -0.31, $P = 0.041$).

Discussion (1258)

As spatial and temporal variation in plant access to water increases in the future (National Research

Council 2011), understanding the impacts of water availability on plants and the associated

heterotrophs are important steps to identify complex effects of water availability on community

interactions. By evaluating interactions across multitrophic levels, we show that plant water

availability differentially impacts specific herbivores and has overall effects on arthropod community

composition. Low water availability reduces overall arthropod abundance associated with the plants,

and this negative impact is linked to decrease in plant quality indicated by reduction in caterpillar

performances. These findings are similar to the common observation that low water availability

affects insect herbivores on herbaceous plant negatively (Waring and Cobb 1992). Notably, we

observed that a specialist herbivore was capable of coping with fluctuations in host plant quality

driven by water availability. The performance of *P. rapae* from Taiwan and Japan revealed a potential

case of local adaptation, where *P. rapae* in Taiwan and Japan performed equally well on sympatric *R.*

indica of different water status. The ability for *P. rapae* to cope with variation in host quality, suggests

the role of host variations in shaping the evolution of physiological capacity of *P. rapae*. Similar local

specialization caused by regional effects and host plants was also documented in other butterfly, such

as *Melitaea cinxia* (Kuussaari et al. 2000).

Even with the physiological capacity to overcome quality variations of plant, *P. rapae* still displayed

preference toward *R. indica* growing under high water availability. As predicted by

preference-performance hypothesis (PPH), oviposition of females should match the performance of

their offspring (Gripenberg et al. 2010; Thompson 1988). Based on the observed decrease in

oviposition on *R. indica* growing under low water availability in both populations, the

preference-performance relationship predicted by PPH was not observed. It has been shown that the

preference of females could be blurred by temporal variation in resource quality (Cronin et al. 2001;

Gripenberg et al. 2007), and thus failure to evolve a proper mechanism to perceive these changes in

host quality for their offspring. Alternatively, this behavior could also be a passive response of female

P. rapae to changes in attractiveness (due to size or chemical cues) of plants under low water

availability. For example, changes in volatile emission patterns are known to be important in

determining the attractiveness of plants to insects (Orre et al. 2010; Pichersky and Gershenzon 2002),

and water availability affects volatile emissions (Copolovici et al. 2014; Scott et al. 2018). It is also

likely that food source stability (Garssen et al. 2014), or preference for food with higher water content

have led to the evolution of preference toward plants under high water availability.

Given the capacity of *P. rapae* to utilize host plants of variable water status, the reduction in other

herbivores and natural enemies on *R. indica* growing under low water availability emerged as an

important factor that increases the survival of *P. rapae*. Although we did not explicitly evaluate the

level of predation and competition, the drastic changes in abundance of herbivores and natural

enemies provides a strong basis in shifting in competition and predation. Several morphospecies that

either consistently observed to prey on *P. rapae* or cause large amounts of damage to *R. indica* in

Taiwan, were linked to observable change in *P. rapae* abundance on *R. indica*. In addition, the overall

low abundance of both herbivores and natural enemies, and species that directly interact with *P. rapae*

in Japan, is a likely reason of the lack of overall association between abundance of arthropods and

survival of *P. rapae* on *R. indica*. This result further supports our conclusion that multitrophic

interaction is an important factor contributing to the benefit of low water availability to *P. rapae*.

A surprising finding that appeared was the positive associations between the survival of *P. rapae* with

two piercing-sucking herbivores in Taiwan. Although this observation contradicted our overall

conclusion that stronger competition reduced the survival of *P. rapae* on *R. indica* growing under high

water availability, it reveals potential divergent defense responses elicited by herbivores in different

feeding guilds. For example, plant defenses induced by piercing-sucking hemipterans are often

salicylic acid-dependent, in contrast to being jasmonic acid-dependent in chewing herbivores. The

antagonistic relationship between these defenses has been frequently shown to benefit chewing

herbivores (De Vos et al. 2005; Kroes et al. 2015; Li et al. 2014; Pieterse et al. 2012; Soler et al. 2012).

We speculate that the positive associations between the survival of *P. rapae* and two piercing-sucking

herbivores were likely mediated by a similar mechanism described above. It has also been shown that

the positive impacts of piercing-sucking herbivores took place only under low density (Kroes et al.

2015)*, which explains the switch in association between survival of *P. rapae* and *E. dominulus* from

positive to negative as the density of *E. dominulus* doubled. Further exploration into mechanisms and

associated tradeoffs in these interactions are needed.

There are some caveats to our study. First, while we generated plants of different water status

morphologically comparable to field individuals, these methods for quantifying water status are

mostly indirect measures of water stress rather than detailed physiological characterizations of plant

stress (Jones 2007). Although they are valuable in detecting plant responses to water limitation, they
are not useful for studies investigating changes in physiological/molecular processes associated with
droughts (Jones 2007). Second, although using detached leaves in caterpillar RGR bioassay has been
shown to reflect chemical changes in leaves (Chung et al. 2013; Wang et al. 2017), we cannot rule out
that detaching leaf itself resulted in alterations in leaf quality. While this method enables us to state
there are changes in the host plants, it may not fully reflect the herbivores performance under natural
feeding conditions. Third, since we only recorded the abundance of herbivore and natural enemies
once in each field experiment, it is possible that arthropod abundance and species richness are
underestimated. Finally, the time period of the field experiment was shorter than the larval stage of *P.*
rapae. As a result, we do not know whether *R. indica* can support the number of larvae observed in
field. However, the likelihood of complete defoliation seems low due to the ability of plants to regrow
and presence of natural enemies under high water availability, and the low quality of plants to other
herbivores under low water availability. The majority of the plants still maintained approximately
50-70% of its biomass after the 10 days field experiment.

The observation that certain herbivores (i.e. *P. rapae*) were unaffected by the reduction in host quality
suggests that addressing focal herbivores without incorporating the natural insect community can alter
interpretations of herbivore success. It can be misleading to focus on few selected insect species and
groups of defense responses because changes in plants under low water availability might have
differential impacts to community members. Some insect herbivores might have better ability to cope
with changes in plant quality than others, while others may be negatively affected by it. Future studies
should aim to include more focal species and aspects of plant-insect interactions in order to capture a
broader scope of how insect and plant communities respond to environmental changes.

In summary, the present study provides insights to the ongoing discussion on how water availability
influences insect herbivores (Jamieson et al. 2012; Mattson and Haack 1987; White 2009). By
studying impact of water availability under the context of arthropod community in addition to the
direct interaction between plants and insect herbivores, we revealed a potential factor in which plants
growing under low water availability can be beneficial to some insects but detrimental to others. The
contradicting impacts of water availability on different arthropod species highlight the complexity and
context dependency of these interactions. The findings also beg the questions of how variation caused
by environmental heterogeneity, including heterogeneity in water availability and other factors, affects
the ecology and evolution of interactions between plants and insect herbivores.

Acknowledgement

We thank the International Agriculture and Development Graduate Program (Pennsylvania State

University) for providing monetary support and feedback on experimental design. We thank Chi-Shun

Chang, Yi-Zhang Wang, Han-Rong Li, Kazumu Kuramitsu, and Kai Han for collecting data. We

would like to express special thanks to Dr. Yooichi Kainoh for arranging space and logistics of

experiment in Japan, Dr. Chin-Wen Tan and Wei-Ting Chen for helpful information on the

experiments in Taiwan, and Dr. Charles Mason and Jagdeep Singh Sidhu for feedbacks on manuscript.

Data availability statement

419   Data in this manuscript will be deposited and made available in the Dryad Digital Repository.

References

421   Anderegg WRL, Kane JM, Anderegg LDL (2013) Consequences of widespread tree mortality triggered by

422   drought and temperature stress. Nat. Clim. Change 3:30-36. doi: 10.1038/nclimate1635

423   Awmack CS, Leather SR (2002) Host plant quality and fecundity in herbivorous insects. Annu. Rev.

424   Entomol. 47:817-844. doi: 10.1146/annurev.ento.47.091201.145300

425   Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, Skaug HJ, Machler M,

426   Bolker BM (2017) glmmTMB balances speed and flexibility among packages for zero-inflated

427   generalized linear mixed modeling. R J. 9:378-400

428   Bunker DE, Carson WP (2005) Drought stress and tropical forest woody seedlings: effect on community

429   structure and composition. J. Ecol. 93:794-806. doi: 10.1111/j.1365-2745.2005.01019.x

430   Cáceres MD, Legendre P (2009) Associations between species and groups of sites: indices and statistical

431   inference. Ecology 90:3566-3574. doi: 10.1890/08-1823.1

432   Chung SH, Rosa C, Scully ED, Peiffer M, Tooker JF, Hoover K, Luthe DS, Felton GW (2013) Herbivore

433   exploits orally secreted bacteria to suppress plant defenses. Proc. Natl. Acad. Sci. U. S. A.

434   110:15728-15733. doi: 10.1073/pnas.1308867110

435   Copolovici L, Kännaste A, Remmel T, Niinemets Ü (2014) Volatile organic compound emissions from

436   *Alnus glutinosa* under interacting drought and herbivory stresses. Environ. Exp. Bot. 100:55-63.

437   doi: 10.1016/j.envexpbot.2013.12.011

438   Cronin JT, Abrahamson WG, Craig TP (2001) Temporal variation in herbivore host-plant preference and

439   performance: constraints on host-plant adaptation. Oikos 93:312-320. doi:

440   10.1034/j.1600-0706.2001.930214.x

441   Dambros CdS, da Silva VNV, Azevedo R, de Morais JW (2013) Road-associated edge effects in Amazonia

442   change termite community composition by modifying environmental conditions. J. Nat. Conserv.

443   21:279-285. doi: https://doi.org/10.1016/j.jnc.2013.02.003

De Vos M, Van Oosten VR, Van Poecke RM, Van Pelt JA, Pozo MJ, Mueller MJ, Buchala AJ, Metraux JP,
Van Loon LC, Dicke M, Pieterse CM (2005) Signal signature and transcriptome changes of
Arabidopsis during pathogen and insect attack. Mol. Plant Microbe Interact. 18:923-937. doi:
10.1094/MPMI-18-0923
English-Loeb G, Stout MJ, Duffey SS (1997) Drought stress in tomatoes: changes in plant chemistry and
potential nonlinear consequences for insect herbivores. Oikos:456-468
Felton GW, Donato K, Del Vecchio RJ, Duffey SS (1989) Activation of plant foliar oxidases by insect
feeding reduces nutritive quality of foliage for noctuid herbivores. J. Chem. Ecol. 15:2667-2694.
doi: 10.1007/BF01014725
Fox J, Weisberg S (2018) An R companion to applied regression. Sage Publications, California
Garssen AG, Verhoeven JTA, Soons MB (2014) Effects of climate-induced increases in summer drought
on riparian plant species: a meta-analysis. Freshwat. Biol. 59:1052-1063. doi: 10.1111/fwb.12328
Gripenberg S, Mayhew PJ, Parnell M, Roslin T (2010) A meta-analysis of preference–performance
relationships in phytophagous insects. Ecol. Lett. 13:383-393. doi:
doi:10.1111/j.1461-0248.2009.01433.x
Gripenberg S, Morriën E, Cudmore A, Salminen J-P, Roslin T (2007) Resource selection by female moths
in a heterogeneous environment: what is a poor girl to do? J. Anim. Ecol. 76:854-865. doi:
10.1111/j.1365-2656.2007.01261.x
Hartig F (2017) DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression models. R
package version 0.1 5:5
Huang TC (1996) Flora of Taiwan. vol 2, 2 edn. Editorial Committee of the Flora of Taiwan, Taipei,
Taiwan, p 766
Jamieson MA, Trowbridge AM, Raffa KF, Lindroth RL (2012) Consequences of climate warming and
altered precipitation patterns for plant-insect and multitrophic interactions. Plant Physiol.
160:1719-1727. doi: 10.1104/pp.112.206524
Jisaburo⬜ O⬜i FGM, Egbert H. Walker (1965) Flora of Japan. Smithsonian Institution
Jones HG (2007) Monitoring plant and soil water status: established and novel methods revisited and their
relevance to studies of drought tolerance. J. Exp. Bot. 58:119-130. doi: 10.1093/jxb/erl118
Kroes A, Van Loon JJA, Dicke M (2015) Density-dependent interference of aphids with caterpillar-induced
defenses in *Arabidopsis*: involvement of phytohormones and ttranscription factors. Plant Cell
Physiol. 56:98-106. doi: 10.1093/pcp/pcu150
Kuussaari M, Singer M, Hanski I (2000) Local specialization and landscape-level influence on host use in
an herbivorous insect. Ecology 81:2177-2187. doi: 10.2307/177106

Langsrud Ø (2003) ANOVA for unbalanced data: use type II instead of type III sums of squares. Stat.

Comput. 13:163-167. doi: 10.1023/A:1023260610025

Li Y, Dicke M, Harvey JA, Gols R (2014) Intra-specific variation in wild *Brassica oleracea* for

aphid-induced plant responses and consequences for caterpillar–parasitoid interactions. Oecologia

174:853-862. doi: 10.1007/s00442-013-2805-6

Li Y, Ye W, Wang M, Yan X (2009) Climate change and drought: a risk assessment of crop-yield impacts.

Clim. Res. 39:31-46. doi: 10.3354/cr00797

Mattson WJ, Haack RA (1987) The role of drought in outbreaks of plant-eating insects. Bioscience

37:110-118. doi: Doi 10.2307/1310365

Murren CJ, Alt CHS, Kohler C, Sancho G (2020) Natural variation on whole‐plant form in the wild is

influenced by multivariate soil nutrient characteristics: natural selection acts on root traits. Am. J.

Bot. doi: 10.1002/ajb2.1420

National Research Council (2011) Advancing the science of climate change. National Academies Press

Ockendon N, Baker DJ, Carr JA, White EC, Almond REA, Amano T, Bertram E, Bradbury RB, Bradley C,

Butchart SHM, Doswald N, Foden W, Gill DJC, Green RE, Sutherland WJ, Tanner EVJ,

Pearce-Higgins JW (2014) Mechanisms underpinning climatic impacts on natural populations:

altered species interactions are more important than direct effects. Global Change Biol.

20:2221-2229. doi: 10.1111/gcb.12559

Oksanen J, Blanchet FG, Kindt R, Legendre P, O'hara R, Simpson GL, Solymos P, Stevens MHH, Wagner

H (2010) Vegan: community ecology package. R package version 1.17-4

Orre GUS, Wratten SD, Jonsson M, Hale RJ (2010) Effects of an herbivore-induced plant volatile on

arthropods from three trophic levels in brassicas. Biol. Control 53:62-67. doi:

10.1016/j.biocontrol.2009.10.010

Peiffer M, Felton GW (2005) The host plant as a factor in the synthesis and secretion of salivary glucose

oxidase in larval *Helicoverpa zea*. Arch. Insect Biochem. Physiol. 58:106-113. doi:

10.1002/arch.20034

Pichersky E, Gershenzon J (2002) The formation and function of plant volatiles: perfumes for pollinator

attraction and defense. Curr. Opin. Plant Biol. 5:237-243. doi: 10.1016/s1369-5266(02)00251-0

Pickett STA, Cadenasso ML, Grove JM, Boone CG, Groffman PM, Irwin E, Kaushal SS, Marshall V,

McGrath BP, Nilon CH, Pouyat RV, Szlavecz K, Troy A, Warren P (2011) Urban ecological

systems: scientific foundations and a decade of progress. J. Environ. Manage. 92:331-362. doi:

https://doi.org/10.1016/j.jenvman.2010.08.022

Pieterse C, Van der Does D, Zamioudis C, Leon-Reyes A, Van Wees S, Schekman R (2012) Hormonal

510   modulation of plant immunity. Annu. Rev. Cell. Dev. Biol. 28:489-521. doi:

511   10.1146/annurev-cellbio-092910-154055

512   R Core Team (2017) R: a language and environment for statistical computing R Foundation for Statistical

513   Computing, Vienna, Austria

514   Schulze E (1986) Whole-plant responses to drought. Funct. Plant Biol. 13:127-141. doi:

515   https://doi.org/10.1071/PP9860127

516   Scott ER, Li X, Kfoury N, Morimoto J, Han W-Y, Ahmed S, Cash SB, Griffin TS, Stepp JR, Robbat A,

517   Orians CM (2018) Interactive effects of drought severity and simulated herbivory on tea (*Camellia*

518   *sinensis*) volatile and non-volatile metabolites. Environ. Exp. Bot. doi:

519   https://doi.org/10.1016/j.envexpbot.2018.10.025

520   Soler R, Badenes-Pérez FR, Broekgaarden C, Zheng S-J, David A, Boland W, Dicke M (2012)

521   Plant-mediated facilitation between a leaf-feeding and a phloem-feeding insect in a brassicaceous

522   plant: from insect performance to gene transcription. Funct. Ecol. 26:156-166. doi:

523   10.1111/j.1365-2435.2011.01902.x

524   Thompson JN (1988) Evolutionary ecology of the relationship between oviposition preference and

525   performance of offspring in phytophagous insects. Entomol. Exp. Appl. 47:3-14. doi:

526   10.1111/j.1570-7458.1988.tb02275.x

527   Trotter R, Cobb N, Whitham T (2008) Arthropod community diversity and trophic structure: a comparison

528   between extremes of plant stress. Ecol. Entomol. 33:1-11. doi: 10.1111/j.1365-2311.2007.00941.x

529   Venables WN, Ripley BD (2013) Modern applied statistics with S-PLUS. Springer Science & Business

530   Media

531   Wang J, Peiffer M, Hoover K, Rosa C, Zeng R, Felton GW (2017) Helicoverpa zea gut-associated bacteria

532   indirectly induce defenses in tomato by triggering a salivary elicitor(s). New Phytol.

533   214:1294-1306. doi: 10.1111/nph.14429

534   Waring GL, Cobb NS (1992) The impact of plant stress on herbivore population dynamics. Insect-plant

535   interactions 4:167-226

536   White TC (1974) A hypothesis to explain outbreaks of looper caterpillars, with special reference to

537   populations of Selidosema suavis in a plantation of Pinus radiata in New Zealand. Oecologia

538   16:279-301. doi: 10.1007/BF00344738

539   White TCR (1969) An index to measure weather-induced stress of trees associated with outbreaks of

540   psyllids in Australia. Ecology 50:905-909. doi: 10.2307/1933707

541   White TCR (2009) Plant vigour versus plant stress: a false dichotomy. Oikos 118:807-808. doi:

542   10.1111/j.1600-0706.2009.17495.x

543   Zuur AF, Ieno EN, Elphick CS (2010) A protocol for data exploration to avoid common statistical

544   problems. Methods Ecol. Evol. 1:3-14. doi: 10.1111/j.2041-210X.2009.00001.x

545   Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM (2009) Mixed effects models and extensions in

546   ecology with R. Springer Science & Business Media

 Figures

 Legends

Fig 1 Quantification of water status on *Rorippa indica*. (**a**) phenotypes after water treatment in a

common garden (2 h after daily water application). (**b**) Box plot of leaf number (N=8, GLM, Poisson).

(**c**) Ratio of length and width of mature leaves (N=8). (**d**) Fresh weight of shoot (N=8). (**e**) Dry weight

of shoot (N=8). (**f**) Shoot water content (N=8). (**g**) Dry mass of root (N=8). (**h**) Root: shoot ratio (g g^{-1},

dry mass) (N=8). (**i**) Soil volumetric water content (θ_v) (N=8). Dots indicate individual observations.

Values indicate mean ± SE. Different letters indicate significant differences between means (ANOVA,

Tukey HSD).

Fig 2 Influence of water status (of *Rorippa indica*), population, and season on *Pieris rapae*. (a)

Ovipositional preferences. (b) Total caterpillars at the end of field experiment, (c) Survival of *P. rapae*

on *R. indica*. Count data were offset by biomass of *R. indica*. Due to higher values of egg, caterpillar,

and herbivore count, data were plotted at the log scale. Transparent points indicate individual

observations; opaque points and error bars indicate mean ± 95% CI.

Fig 3 Relative growth rates of lepidopteran larvae feeding on leaves of *Rorippa indica*. (a) *Pieris*

rapae (Taiwan) and *R. indica* (Taiwan) (N=30). (b) *P. rapae* (Japan) and *R. indica* (Japan) (N=35). (c)

P. rapae (North America) and *R. indica* from both location (Taiwan and Japan) (N=40). (d)

Helicoverpa zea (North America) and *R. indica* from both location (Taiwan and Japan) (N=40). (e)

Trichoplusia ni (North America) and *R. indica* from both location (Taiwan and Japan) (N=40). Values

represent means ± SE. Different letters indicate significant differences between means (ANOVA,

Tukey HSD). (f) *Spodoptera frugiperda* (North America) and *R. indica* from both location (Taiwan

and Japan) (N=40). 1 = dead, 0 = lived. Bar indicated the mean and boxes represent 95% CI (GLM,

binomial).

Fig 4 Abundance of herbivores and natural enemies and its associations with water status of *Rorippa*

indica and survival of *Pieris rapae*. (a) Abundance of herbivore in Taiwan. (b-d) Association between

herbivore abundance and survival of *Pieris rapae* in Taiwan during early, middle, and late season. (e)

Abundance of natural enemy in Taiwan. (f-h) Association between natural enemy abundance and

survival of *Pieris rapae* in Taiwan during early, middle, and late season. (a) Abundance of herbivore

in Japan. (b-d) Association between herbivore abundance and survival of *Pieris rapae* in Japan during

early, middle, and late season. (e) Abundance of natural enemy in Japan. (f-h) Association between

natural enemy abundance and survival of *Pieris rapae* in Japan during early, middle, and late season.

Lines represent model predictions (solid: significant; dashed: non-significant). Grey areas indicate

95% CI. For abundance plots, values indicate estimated mean with 95% CI. All dots indicate

individual observations. Letters indicate significant difference between means (Tukey HSD). * P <

0.05 ** P < 0.01 *** P < 0.001

Fig 5 Herbivore and natural enemy compositions on *Rorippa indica* with different water status across

seasons. (a) Herbivore composition in Taiwan. (b) Natural enemy composition in Taiwan. (c)

Herbivore composition in Japan. (d) Natural enemy composition in Japan. For method please see

Appendix 6. Abbreviation for herbivore: Aph=Aphididae, Api=Apidae, Acr=Acrididae, Aly=Alydidae,

Cat=Catantopidae, Cic=Cicadellidae, Chy=Chrysomelidae, Din=Dinidoridae, Eur=*Eurydema*

dominulus, Gas=Gastropoda, Hel=*Helicoverpa armigera*, Mir=Miridae, Mit=Tetranychidae,

Pen=Pentatomidae, Pha=*Phaedon brassicae*, Phy=*Phyllotreta striolata*, Plu=*Plutella xylostella*,

Pyr=Pyrgomorphidae, Spo=*Spodoptera litura*, Thr=Thripidae. Abbreviation for natural enemy:

Ara=Araneae, Bra=Braconidae, Cer=Ceraphronidae, Chr=Chrysopidae, Coc=Coccinellidae,

Eoc=*Eocanthecona concinna*, Eup=Eupelmidae, Eul=Eulophidae, For=Formicidae, Pte=Pteromalidae,

Syr=Syrphidae, Tac=Tachinidae, Tro=Trombidiidae, Red=Reduviidae.

Fig 6 Graphical summary of relationship between water availability, plant, and insect herbivore.

Female *Peiris rapae* showed ovipositional preferences toward *Rorippa indica* growing under high

water availability. *R. indica* growing under low water availability were associated with lower overall

abundance of arthropods. The changes in species interactions may contribute to the higher survival of

P. rapae larvae on *R. indica* growing under low water availability.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

612   **Fig. 5**

615 **Fig. 6**

618 **Table**

619 **Table 1 Indicator species of water availability in different season**

Species	Treatment	Ind. Val.	P-value
Herbivore			
Japan (early)			
Thripidae	200 mL day^{-1}	0.589	<0.001
Japan (middle)			
Aphididae	200 mL day^{-1}	0.559	<0.001
Eurydema	200 mL day^{-1}	0.551	<0.001
Catantopidae	200 mL day^{-1}	0.596	0.004
Pyrgomorphidae	200 mL day^{-1}	0.363	<0.001
Pentatomidae	200 mL day^{-1}	0.337	0.018
Japan (late)			
Chrysomelidae	200 mL day^{-1}	0.364	0.034
Taiwan (early)			
Phaedon	200 mL day^{-1}	0.675	<0.001
Taiwan (middle)			
Phyllotreta	200 mL day^{-1}	0.555	<0.001
Eurydema	200 mL day^{-1}	0.506	0.025
Catantopidae	200 mL day^{-1}	0.482	0.001
Taiwan (late)			
Plutella	200 mL day^{-1}	0.487	<0.001

Aphididae	200 mL day^{-1}	0.369	<0.001
Tetranychidae	10 mL day^{-1}	0.487	0.028
Natural enemy			
Japan (early)			
Formicidae	200 mL day^{-1}	0.447	<0.001
Syrphidae	200 mL day^{-1}	0.393	0.002
Trombidiidae	200 mL day^{-1}	0.54	<0.001
Taiwan (middle)			
Eupelmidae	200 mL day^{-1}	0.341	0.017
Taiwan (late)			
Formicidae	200 mL day^{-1}	0.43	<0.001
Braconidae	200 mL day^{-1}	0.511	<0.001
Eocanthecona	200 mL day^{-1}	0.555	0.021
Coccinellidae	200 mL day^{-1}	0.675	<0.001
Araneae	200 mL day^{-1}	0.369	0.009
Pteromalidae	200 mL day^{-1}	0.506	0.001
Ceraphronidae	200 mL day^{-1}	0.482	0.04
Others			
Japan (middle)			
Diptera	200 mL day^{-1}	0.693	0.001
Taiwan (middle)			

Lauxaniidae	200 mL day^{-1}	0.437	0.018
Taiwan (late)			
Staphylinidae	200 mL day^{-1}	0.341	0.022

 Table 2 Model summaries of associations between survival of Pieris rapae and morphospecies.

Term	Estimate	SE	Z-value	P-value
Taiwan (Early)				
Intercept	-5.215	1.236	-4.221	<0.001
Phyllotreta	-6.500	3.151	-2.063	0.04
Taiwan (Middle)				
Intercept	-3.544	0.33	-10.729	<0.001
Phyllotreta	-1.049	0.334	-3.141	0.002
Eurydema	0.988	0.374	2.643	0.008
Braconidae	-1.675	0.419	-3.994	<0.001
Tetranychidae	-0.789	0.376	-2.095	0.036
Taiwan (Late)				
Intercept	-1.283	0.171	-7.506	<0.001
Eurydema	-0.953	0.272	-3.503	<0.001
Cicadellidae	1.31	0.29	4.516	<0.001
Eupelmidae	-0.6	0.178	-3.382	0.001
Eocanthecona	-0.441	0.186	-2.371	0.018
Japan (Early)				
Intercept	-0.639	0.194	-3.298	0.001
Lauxaniidae	-0.953	0.412	-2.315	0.021
Trombidiidae	0.499	0.15	3.326	0.001
Japan (Middle)				

Intercept	0.873	0.206	4.238	<0.001
Syrphidae	-0.31	0.151	-2.048	0.041
Japan (Late)				
NA	NA	NA	NA	NA

623