Somatostatin interneurons activated by 5-HT$_{2A}$ receptor suppress slow oscillations in medial entorhinal cortex

Roberto De Filippo [1, 6, *], **Benjamin R. Rost** [2], **Alexander Stumpf** [1], **Claire Cooper**[1], **John J. Tukker** [1,2], **Christoph Harms** [3-5], **Prateep Beed** [1, #], **Dietmar Schmitz** [1, 2, 5, 6, #, *]

[1] Charité–Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin,and Berlin Institute of Health; Neuroscience Research Center, 10117 Berlin, Germany

[2] German Centre for Neurodegenerative Diseases (DZNE), 10117 Berlin, German

[3] Charité–Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health; Department of Experimental Neurology, 10117 Berlin, Germany

[4] Charité–Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health; Center for Stroke Research Berlin, 10117 Berlin, Germany

[5] Charité–Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health; Einstein Center for Neurosciences Berlin, 10117 Berlin, Germany

[6] Charité–Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health; Cluster of Excellence NeuroCure, 10117 Berlin, Germany.

# These authors contributed equally to this work
* Correspondence to: dietmar.schmitz@charite.de and roberto.de-filippo@charite.de

Keywords: serotonin, 5-HT$_{2A}$, somatostatin interneurons, slow oscillations, MDMA, entorhinal cortex.

Abstract

Serotonin (5-HT) is one of the major neuromodulators present in the mammalian brain and has been shown to play a role in multiple physiological processes. The mechanisms by which 5-HT modulates cortical network activity, however, are not yet fully understood. We investigated the effects of 5-HT on slow oscillations (SOs), a synchronized cortical network activity universally present across species. SOs are observed during anesthesia and are considered to be the default cortical activity pattern. We discovered that (±)3,4-methylenedioxymethamphetamine (MDMA) and fenfluramine, two potent 5-HT releasers, inhibit SOs within the entorhinal cortex (EC) in anesthetized mice. Combining opto- and pharmacogenetic manipulations with *in vitro* electrophysiological recordings, we uncovered that somatostatin-expressing (Sst) interneurons activated by the 5-HT$_{2A}$ receptor (5-HT$_{2A}$R)

38 play an important role in the suppression of SOs. Since 5-HT$_{2A}$R signaling is involved in the

39 etiology of different psychiatric disorders and mediates the psychological effects of many

40 psychoactive serotonergic drugs, we propose that the newly discovered link between Sst

41 interneurons and 5-HT will contribute to our understanding of these complex topics.

Introduction

5-HT is one of the most important neuromodulators in the central nervous system. Projections originating from the Raphe nuclei, the brain-stem structure that comprises the majority of 5-HT releasing neurons in the brain, innervate all cortical and sub-cortical areas (Descarries et al., 2010). 5-HT levels in the brain are closely linked to the sleep-wake cycle : the activity of serotonergic raphe neurons is increased during wakefulness, decreased during slow-wave sleep (SWS) and virtually silent during REM sleep (McGinty and Harper, 1976, Oikonomou et al., 2019, Unger et al., 2020). Cortical activity is also influenced by the behavioral state of the animal: SWS is generally associated with "synchronized" patterns of activity, characterized by low-frequency global fluctuations, whereas active wakefulness and REM sleep feature "desynchronized" network activity, in which low-frequency fluctuations are absent. The shifting of cortical networks between different patterns of activity is controlled, at least in part, by neuromodulators (Tukker et al., 2020, Lee and Dan, 2012). For instance, Acetylcholine (ACh) can profoundly alter cortical network activity by inducing desynchronization via activation of Sst interneurons (Chen et al., 2015). However, ACh is not solely responsible for suppressing cortical synchronized activity, as lesions of cholinergic neurons are not sufficient to abolish desynchronization (Kaur et al., 2008). On the other hand, blocking ACh and 5-HT transmission at the same time causes a continuous "synchronized" cortical state, even during active behavior, thus suggesting that 5-HT plays an important role in mediating transitions between different network states (Vanderwolf and Baker, 1986). In agreement with this line of thought, electrical and optogenetic stimulation of the Raphe nuclei reduce low frequency (1-10 Hz) power in the cortex, implying a reduction in neuronal synchronization at these frequencies (Puig et al., 2010, Grandjean et al., 2019). Moreover, optogenetic stimulation of serotonergic neurons is sufficient to awaken mice from SWS (Oikonomou et al., 2019). These data suggest a relationship between 5-HT levels and patterns of cortical activity (Lee and Dan, 2012, Harris and Thiele, 2011). The exact mechanism by which 5-HT modulates network activity in the cortex however, is still not fully understood.

Here, we used electrophysiological techniques together with pharmacology, optogenetics and pharmacogenetics to investigate the effect of 5-HT on SOs, a network oscillation characterized by synchronized transitions (< 1 Hz) between periods of high activity (upstates) and relative quiescence (downstates) (Steriade et al., 1993, Neske, 2015, Isomura et al., 2006). SOs are a global phenomenon observed throughout the cerebral cortex and are considered to be the default emergent activity of cortical networks during SWS and anesthesia (Neske, 2015, Sanchez-Vives et al., 2017, Wolansky et al., 2006). We performed our experiments in

the medial entorhinal cortex (mEC), a region where SOs can be recorded both under anesthesia and in acutely prepared cortical slices (Tahvildari et al., 2012, Beed et al., 2020). Pyramidal neurons located in L3 of mEC provide the excitatory drive underlying each upstate (Namiki et al., 2013, Beed et al., 2020). Similarly to other mammalian cortical areas, mEC comprises different types of inhibitory GABAergic neurons that can be grouped into three main classes according to immunoreactivity: parvalbumin (PV), somatostatin (Sst) and 5-HT$_3$ (Miao et al., 2017, Rudy et al., 2011). Most PV neurons target the soma and the spike initiation zone, have low input resistance and minimal spike frequency adaptation. Sst neurons are divided into two groups, one showing features similar to PV interneurons and a second one (i.e. Martinotti cells) that, in contrast, tend to form synapses onto the dendritic trees of their target cells, have high input resistance and show a considerable adaptation. 5-HT$_3$ neurons are usually located in the superficial layers, have high input resistance and mixed adaptation. While these different classes of interneurons are all depolarized during upstates, PV interneurons receive decidedly the strongest excitation (Tahvildari et al., 2012, Neske et al., 2015a). Recurrent excitation and temporal summation of inputs contribute to the transition from downstate to upstate (Sanchez-Vives et al., 2017, Tukker et al., 2020). This excitation propagates to L5 and differentially entrains L5a and L5b excitatory neurons. L5a neurons do not participate in SOs whereas L5b neurons are steadily synchronized. Both intrinsic and synaptic mechanisms have been implicated in upstate termination (Neske, 2015, Tukker et al., 2020). Activity-dependent K$^+$ channels decrease excitability of neurons over time causing a generalized reduction of facilitation (Neske, 2015, Harris and Thiele, 2011). At the same time, blockage of GABA$_B$ receptors significantly extends upstate duration (Craig et al., 2013) and inhibitory drive has been observed to increase during upstate termination (Lemieux et al., 2015). Sst interneurons, and in particular Martinotti cells, characterized by strongly facilitating synapses (Beierlein et al., 2003, Gibson et al., 1999) have been proposed as an important source of inhibition in the termination of upstates (Melamed et al., 2008, Krishnamurthy et al., 2012).

Our results show that MDMA and fenfluramine (Fen), two substitute amphetamines that induce robust 5-HT release, suppress SOs in anesthetized mice and, concurrently, activate a small group of neurons characterized by an intermediate waveform shape. Using cortical slices *in vitro*, we demonstrate that 5-HT inhibits SOs in the mEC and that Sst interneurons, activated by 5-HT$_{2A}$R, are involved in the modulatory effect of 5-HT. While previous studies have shown that parvalbumin (PV) interneurons are excited by 5-HT$_{2A}$R (Puig et al., 2010,

Athilingam et al., 2017), our results identify cortical Sst interneurons as novel targets of the 5-HT neuromodulatory system via 5-HT$_{2A}$R.

Results

Pharmacological release of 5-HT inhibits SOs in anesthetized mice

We investigated the effect of 5-HT on network activity in anesthetized mice using multisite silicon microelectrodes placed in the mEC L3, a region located in the medial temporal lobe interconnected to a variety of cortical and subcortical areas, including the Raphe nuclei (Figure 1A) (van Strien et al., 2009, Muzerelle et al., 2016). Under urethane anesthesia, EC, like the rest of the cortex, displays SOs (Figure 1B). As expected, we found that upstates were present synchronously in the local field potential (LFP) of all the recording channels (Figure 1–figure supplement 1), and that every upstate coincided with large increases in population spiking activity

Figure 1| MDMA/Fen inhibit SOs in anesthetized mice. (A) Immunocytochemical analysis of an ePet-YFP mouse showing serotonergic fibers in medial entorhinal cortex, horizontal slice (M = medial, L = lateral, P= posterior, A = anterior). Scale bar: 20 μm. (B) LFP (top) and instantaneous population activity (bottom) of a representative *in vivo* recording during SOs (spikes/s units in thousands), cyan rectangles represent detected upstates. (C) 3D visualization of the microelectrode location of the recording shown in E. EC represented in grey. (D) Left: microelectrode tracks (red) of the recording shown in (E). Right: EC position represented in green. (E) Top: Cyan lines represent

(Figure 1B). 5-HT does not cross the blood brain barrier (Hardebo and Owman, 1980),

therefore, to understand the effect of 5-HT on SOs we used MDMA, a potent presynaptic 5-

HT releaser and popular recreational drug (Green et al., 2003) that has shown promising

results in the treatment of post-traumatic stress disorder (PTSD) (Inserra et al., 2021).

Intraperitoneal injections of MDMA (1.25 mg/kg) caused a strong suppression of upstate

incidence (Figure 1E-H), a decrease in power of low frequencies (1-20 Hz) (Figure 1F), and a

reduction of population spiking activity (Figure 1G-I). In addition to 5-HT, MDMA has been

shown to cause the release of other neuromodulators, such as dopamine and noradrenaline

(NE), although to a much lesser extent (Green et al., 2003). To test whether the effect of MDMA

was mediated specifically by 5-HT, we repeated the experiment using Fen (5 mg/kg), a more

selective 5-HT releaser (Rothman and Baumann, 2002, Heifets et al., 2019, Baumann et al.,

2000). Intraperitoneal injection of Fen had a comparably strong suppressive effect on both the

occurrence of upstates and population spiking activity (Figure 1H-I). Given the similarity of the

observed effects, we grouped the results of Fen and MDMA application and found that both

of these drugs significantly decreased LFP power in the delta and theta frequency ranges

(Figure 1–figure supplement 2). Furthermore, the duration and area of upstates were also

significantly reduced (Figure 1–figure supplement 3). These results point to a likely

involvement of 5-HT in modulating ongoing oscillatory activity and suppressing low-frequency

fluctuations.

MDMA and Fen activate a subgroup of cortical neurons

In addition to the LFP signal, we recorded the activity of 355 single units within the EC.

Because of the similar effect of MDMA and Fen on spike rates (Figure 1I), we pooled all units

recorded in both types of experiments. We found that drug injections differentially affected

spiking rates (Figure 2A) of recorded units: while spiking decreased in most units ('non-

activated'), a small group of units ('activated') responded in the opposite fashion (n=31/355,

8,7%).

We then calculated the trough-to-peak (TP) latency of the spike waveforms, which has been

consistently used as a metric to classify units. In accordance with previous studies (Senzai et al., 2019, Roux et al., 2014), we found a clear bimodal distribution of TP latencies distinguishing putative excitatory (Exc) and fast-spiking inhibitory (Int) groups. Analysis of cross-correlograms confirmed the inhibitory nature of a subset of putative FS units (Figure 2– figure supplement 1) (Barthó et al., 2004). The cumulative distribution of TP latencies of 'activated' units was significantly different from both Exc and Int groups (Figure 2B). Specifically, the average TP latency of 'activated' units was situated in between the Int and Exc groups (Figure 2B-D, Figure 2–figure supplement 2), possibly suggesting a non-FS interneuron identity (Trainito et al., 2019, Kvitsiani et al., 2013). Notably the intermediate TP latency of "activated" units is not the result of an equal distribution between high and low values. Units with intermediate TP latency (between 0.4 and 0.6 ms) are unique in showing a 40% to 80% likelihood of being "activated" by drug application (Figure 2B).

Figure 2| Divergent unit responses to MDMA/Fen application. (A) Spike rate of the activated units versus all the other units during MDMA/Fen application (activated: n = 31, Non-activated: n = 324). (B) Top: TP latencies color-coded by group. Middle: cumulative distribution of TP latencies (Kolmogorov-Smirnov test, p Activated vs Int < 10-4, p Activated vs Exc < 10-4). Bottom: bar plot representing probability distribution of TP latencies, on the right y axis dashed line representing the percentage of 'activated' units per TP latency bin. (C) Distribution of units according to trough-to-peak (TP) latencies and repolarization time. Units were classified as putative interneurons (Int, blue) and putative excitatory neurons (Exc,

dark gray) according to a threshold at 0.55 ms; activated units (red) could belong to either
group but were mostly intermediate as shown by the covariance (2 STD) of each group
(Ellipses). Units recorded during control experiments are represented by empty circles. (D)
Waveforms of recorded units (n= 355). Units were divided into "putative excitatory" (black)
and "putative inhibitory" (blue) neurons according to TP latencies. Units activated by either
MDMA or Fen application are represented in red. Inset shows the average waveform for each
group. Scale bars: 0.5 ms.

5-HT suppresses SOs and activates Sst interneurons *via* 5-HT$_{2A}$R *in vitro*

To understand the mechanism underlying the suppression of SOs by MDMA/Fen, we

performed *in vitro* experiments combining electrophysiology and pharmacology. First, we

recorded simultaneously from up to four neurons in the superficial layers of the EC (Figure

3A). Brain slices were perfused with an extracellular solution containing Mg^{2+} and Ca^{2+} in

concentrations similar to physiological conditions. With this method we could reliably detect

SOs reminiscent of the *in vivo* network activity (Tahvildari et al., 2012). Release of 5-HT *in*

vitro, induced by Fen application (200 μM), caused a suppression of SOs similar to what we

observed *in vivo* (Figure 3–figure supplement 1). Likewise, application of low concentrations

of 5-HT (5 μM) resulted in the suppression of SOs (Figure 3B, C). This effect was highly

consistent across different slices and was readily reversible (Figure 3D). Similarly to

spontaneous upstates, electrically evoked upstates (Neske et al., 2015b) were also

suppressed by 5-HT (Figure 3–figure supplement 2). Increasing the stimulation intensity did

not rescue upstate generation, indicating that a lack of excitation alone cannot explain the

suppressive effect of 5-HT on SO. In accordance with these results, we decided to focus

solely on 5-HT, however, we acknowledge that other neurotransmitters might be involved in

the effect of MDMA/Fen we observed *in vivo*.

Suppression of activity can have either an intrinsic or synaptic origin (Turrigiano, 2011). A

substantial subset of EC excitatory neurons is known to express 5-HT$_{1A}$ receptor (5-HT$_{1A}$R)

and hyperpolarize upon 5-HT application via activation of G protein-coupled inwardly

rectifying potassium (GIRK) channels (Schmitz et al., 1998, Chalmers and Watson, 1991). The

suppression of SOs by 5-HT, however, was not influenced by blocking 5-HT$_{1A}$R (WAY 100635,

100 nM) (Figure 3–figure supplement 3A, E). Blocking 5-HT$_{3}$R (tropisetron, 1μM), the receptor

that characterizes one of the three main groups of interneurons, also did not have any impact

on SOs suppression induced by 5-HT. Similarly, application of the 5-HT$_{3}$R agonist m-CPBG

(50 μM) did not have any effect on SOs (Figure 3–figure supplement 4). In contrast, blocking

5-HT$_{2A}$R with the selective antagonist ketanserin (1 μM) (Preller et al., 2018) strongly reduced

the suppression power of 5-HT on SOs from 95 ± 4% to 57 ± 10.1% (Figure 4–figure

supplement 3B, E). The remaining suppression can be possibly explained by the activation of

5-HT$_{1A}$R on excitatory cells, as is reflected by the reduced spiking activity of putative

excitatory cells (Figure 3–figure supplement 5).

Selective activation of 5-HT$_{2A}$R by α-methyl-5-HT (5 µM) mimicked the suppression of SOs

observed after 5-HT wash-in (Figure 3–figure supplement 3C, E). Together, these results point

to the importance of 5-HT$_{2A}$R in the suppression of SOs. 5-HT$_{2A}$R activation is known to cause

an increase in intracellular calcium and consequent depolarization of the resting potential (RP)

(Nichols and Nichols, 2008). Accordingly, after 5-HT application, we found that a small group

of neurons was depolarized (n= 6/48, 12,5%) (Figure 3–figure supplement 6B-C). Using a soft

Figure 3| 5-HT suppresses SOs and activates Sst interneurons. (A) Biocytin staining of
four simultaneously recorded cells shown in (B) WFS1 expression (in red) delimits L2/3
border. (B) Intracellular recordings showing synchronous upstate events in 4 simultaneously
recorded cells before (left) and after (right) 5-HT application. Scale bars: 1: 7.5 mV, 2: 25 mV,
3: 25 mV, 4: 10 mV; 10 s. (C) Upstate raster plot before and after 5-HT application, orange
box represents 5-HT application (n = 17, p < 10^{-4}, Wilcoxon signed rank test). (D)
Representative recording showing the temporary inhibitory effect of 5-HT on SOs in two
simultaneously recorded cells. Scale bars: 5 min, 20 mV. (E) PCA projection plot of all the
cells recorded (n = 48). Cells are color-coded according to group identity: Exc (black), FS
(light blue) or LTS (green). Typical voltage responses to current injection (-150 and +250 pA)
are plotted for each group. Inset shows the average spike waveform for each group. (F)
Representative recording of an excitatory (black) and a low-threshold (green) neuron
simultaneously recorded during 5-HT application. Scale bars: 10 mV, 30 s. (G) Average
change of RP before and after 5-HT application, across excitatory, fast-spiking and low-
threshold neurons (Exc: n = 34, FS: n = 6; LTS: n = 9; p < 10^{-4}, unpaired t test with Holm-Šidák
correction). (H) Biocytin staining of cells recorded in Sst-tdTomato mouse. Biocytin in green,

clustering approach with six electrophysiological parameters (see "Materials and Methods")

we divided the recorded cells into 3 groups: Excitatory (Exc), fast-spiking (FS) and low-

threshold spiking (LTS). Strikingly, the cells excited by 5-HT belonged exclusively to the LTS

group (Figure 3G, Figure 3–figure supplement 6). A substantial proportion of LTS neurons

express Sst (Tremblay et al., 2016, Gibson et al., 1999), therefore we performed targeted

patch-clamp recordings using a mouse line expressing tdTomato specifically in Sst-

expressing interneurons (Figure 3H). A subgroup of Sst interneurons are FS cells (Urban-

Ciecko et al., 2018), only LTS Sst interneurons were considered in the following analysis. Sst

interneurons depolarized upon 5-HT application (n=19, ΔRP: 7.5 ± 1.23 mV) (Figure 3I-L) and

in some cases continued to spike while SOs were suppressed (n= 8/17, 47.05%, mean $_{spiking}$

$_{rate}$ = 3.03 ± 0.39 spikes/s). This effect was blocked by ketanserin (n = 22) (Figure 3L). We

confirmed the presence of 5-HT$_{2A}$R in Sst interneurons using immunohistochemistry in mice

expressing EGFP under the 5-HT$_{2A}$R promoter. In accordance with a previous study we found

the majority of 5-HT$_{2A}$R positive cells in the deep layers of EC, with a peak in L6 (Weber and

Andrade, 2010). We observed that 11.8 ± 2.9 % of the 5-HT$_{2A}$R positive cells in EC colocalized

with Sst (n = 7 mice) (Figure 3–figure supplement 9). These results suggest that Sst

interneurons may provide the synaptic inhibition required for the suppression of SOs.

Sst interneurons mediate the suppression of SOs by 5-HT *in vitro*

To evaluate the contribution of Sst interneurons to the 5-HT-mediated silencing of SOs we

used an opto- and pharmacogenetic approach. First, we transgenically expressed

channelrhodopsin-2 (ChR2) in Sst interneurons (Figure 4A). Light-stimulation of ChR2-

expressing Sst interneurons *in vitro* suppressed SOs consistently (Figure 4D-F). Expectedly,

upstate associated spiking was also diminished (Figure 4C, E, F). At the end of the light

stimulation, spontaneous upstates immediately reoccurred (Figure 4G-H), in line with a critical

role of Sst interneurons in the modulation of SOs (Fanselow et al., 2008, Funk et al., 2017,

Niethard et al., 2018). We acknowledge that activation of PV interneurons can cause a similar

suppression (Zucca et al., 2017). While this experiment establishes the ability of Sst

interneurons to suppress SOs, it does not causally link Sst interneuron activation to the

suppression of SOs induced by 5-HT. Therefore, we generated a transgenic mouse line

carrying a Cre-conditional expression cassette of the pharmacogenetic silencer hM4Di (Figure
4–figure supplement 1) (Armbruster et al., 2007). Homozygous Cre-conditional hM4Di
transgenic mice and Sst-Cre mice were bred to obtain heterozygous Sst-Cre/hM4Di offspring,
which allow specific inhibition of Sst interneuron activity using Clozapine-N-Oxide (CNO).
Following application of 5-HT we observed a strong reduction of upstate incidence, which
was partially restored by subsequent application of CNO (Figure 5 A-B).

Figure 5| **Sst interneurons mediate the effect of 5-HT on SO.** (A) Upstate raster plot during 5-HT and subsequent CNO application. Orange box represents 5-HT, purple boxes represent CNO. Note the appearance of upstates after CNO application. (B) Upstate incidence during 5-HT and 5-HT+CNO application (n=15; p (baseline vs 5-HT) < 10-4, p (baseline vs CNO) = 0.0482, p5-HT vs CNO = 0.0405, Kruskal-Wallis test). Patches represent 95% confidence intervals, lines represent standard deviation.

Activation of 5-HT$_{1A}$R on excitatory cells by 5-HT and the resulting decreased network excitation drive might account for the fact that the incidence of upstates did not completely return to the baseline level upon CNO wash-in. Additionally, in further experiments in which CNO was applied before 5-HT, the significant reduction in upstate incidence typically seen after 5-HT wash-in was not observed (Figure 5–figure supplement 1). CNO did not show any significant effect in both wild-type littermates and hM4Di-PV mice (Figure 5–figure supplement 2), indicating that the results observed are due to its specific effect on the activity of Sst interneurons. In summary, while activation of Sst interneurons either via 5-HT or directly by ChR2 suppresses SOs, the pharmacogenetic inactivation of Sst interneurons weakens the effect of 5-HT on SOs.

Discussion

In this study we show that the substitute amphetamines MDMA and Fen suppress default cortical network oscillations *in vivo* in the mEC. Furthermore, using an opto- and pharmacogenetic approach *in vitro*, we demonstrate that Sst interneurons, activated by 5-HT$_{2A}$R, contribute to this suppression.

Organization of cortical activity is brain state-dependent, ranging continuously from "synchronized" to "desynchronized" states (Harris and Thiele, 2011). SOs are on one end of

this continuum, representing the prototypical synchronized state. Our results, in line with
previous studies (Puig et al., 2010, Grandjean et al., 2019), show that 5-HT can suppress
synchronized cortical activity; in addition, we identify Sst interneurons as contributing to this
suppression.

Sst interneurons have been previously proposed to provide the inhibition necessary for the
termination of upstates, due to their strongly facilitating synapses (Krishnamurthy et al., 2012,
Melamed et al., 2008). Additionally, increased inhibition, as shown both in computational
models and experimental data, can counteract temporal summation of inputs and reduce
correlation due to tracking of shared inputs between inhibitory and excitatory populations
(Sippy and Yuste, 2013, Renart et al., 2010, Stringer et al., 2016). In the case of Sst
interneurons, this would especially impact summation in distal compartments of pyramidal
cells, due to the axonic targeting of Sst cells onto pyramids. Sst interneurons are specifically
known to be sufficient to cause desynchronization in V1 (Chen et al., 2015). While it is well
known that Sst interneurons are potently excited by ACh in various cortical areas (Chen et al.,
2015, Obermayer et al., 2018, Fanselow et al., 2008), including mEC (Desikan et al., 2018),
and NE in frontal cortex (Kawaguchi and Kubota, 1998), our work identifies them as a novel
target of 5-HT regulation via 5-HT$_{2A}$R. Recently it has been proposed that inhibitory
interneurons play a key role in mediating the effect of ACh and NE on cortical state transitions
(Cardin, 2019). Our results add a new level of complexity to this picture.

The excitation of Sst interneurons by 5-HT possibly contributes to the net inhibitory effect of
5-HT release observed in many cortical areas (Grandjean et al., 2019, Seillier et al., 2017,
Azimi et al., 2020), and could explain why the inhibition strength is linearly correlated to 5-
HT$_{2A}$R expression (Grandjean et al., 2019). Giving further support to this idea, Sst interneurons
in the somatosensory cortex show increased cFos levels following 5-HT$_{2A}$R activation (Martin
and Nichols, 2016). Previous studies have reported either 5-HT$_{2A}$R dependent inhibition or 5-
HT$_{2A}$R dependent activation of interneurons in the prefrontal cortex (PFC) (Abi-Saab et al.,
1999, Ashby et al., 1990, Athilingam et al., 2017), piriform cortex (Marek and Aghajanian, 1994,
Sheldon and Aghajanian, 1990), cingulate cortex (Zhou and Hablitz, 1999), cochlear nucleus
(Tang and Trussell, 2017), amygdala (Sengupta et al., 2017), olfactory bulb (Petzold et al.,
2009, Hardy et al., 2005), visual cortex (Michaiel et al., 2019, Azimi et al., 2020) and
hippocampus (Wyskiel and Andrade, 2016). However, none of these studies identified
interneurons using molecular markers, so we do not exclude that different interneuron classes
in other cortical areas might mediate the inhibitory downstream effects of 5-HT$_{2A}$R. For
example, in the PFC a subgroup of PV interneurons has been reported to be activated by this

receptor (Athilingam et al., 2017, Puig et al., 2010). Furthermore, we know that PV neurons can reliably induce up to down state transitions (Zucca et al., 2017). 5-HT modulation is also involved in gain regulation. In the olfactory cortex, 5-HT has a selective subtractive effect on stimulus evoked firing (Lottem et al., 2016), and a recent study has shown in the visual cortex that the reduced gain of evoked responses is dependent on $5\text{-}HT_{2A}R$ activation (Azimi et al., 2020). Intriguingly, Sst interneurons have been shown to regulate subtractive inhibition (Sturgill and Isaacson, 2015, Wilson et al., 2012).

5-HT levels in the brain, similarly to other neuromodulators, are synchronized to the sleep wake cycle, with higher levels present during the waking state (Oikonomou et al., 2019). How does this notion relate to activation of Sst interneurons by $5\text{-}HT_{2A}R$? A limitation of our study is the absence of data in naturalistic conditions, with anesthesia potentially being a confounding factor (Adesnik et al., 2012). We can nonetheless speculate that Sst interneurons should be more active during states with higher 5-HT levels (wake>SWS>REM). A previous study directly measuring the activity of various neuronal classes across different states seems to support this idea. Sst interneurons in the dorsal cortical surface display their highest activity during waking states, lower activity during SWS and lowest activity during REM (Niethard et al., 2016). Following pharmacological release of 5-HT *in* vivo we observed increased spiking activity in a subgroup of neurons whose waveform features are compatible with Sst interneurons. Our *in vitro* data using 5-HT at a concentration commonly used in the field (Gorinski et al., 2019, Wang et al., 2016, Huang et al., 2009) shows that Sst interneurons can be activated by $5\text{-}HT_{2A}R$, however, recordings during natural sleep and wake conditions in combination with optotagging will be needed to demonstrate conclusively that Sst interneurons are activated by 5-HT in physiological conditions *in vivo*. Potential differences between physiological and pharmacological 5-HT release are especially of interest considering the reported efficacy of MDMA in the treatment of PTSD and its recently approved status as breakthrough therapy (Inserra et al., 2021, Mithoefer et al., 2019).

While our data relative to MDMA application supports a role of 5-HT in modulating SOs *in vivo* we cannot exclude that other neuromodulators might play a role. For example, it is known that dopamine can suppress SOs *in* vitro (Mayne et al., 2013). Fen, however, has been reported to selectively increase 5-HT concentration in the brain (Rothman and Baumann, 2002) and has been used previously to specifically disentangle the effect of MDMA on different neuromodulatory systems (Heifets et al., 2019). This, together with our *in vitro* results, suggests that 5-HT is most likely mediating the effect induced by MDMA/Fen.

Besides its involvement in various physiological brain processes, 5-HT is also associated with the etiology of various psychiatric disorders, as are Sst interneurons (Pantazopoulos et al., 2017, Lin and Sibille, 2015). Furthermore, 5-HT is linked to the psychological effect of many psychotropic drugs; specifically, $5\text{-HT}_{2A}\text{R}$ activation has been reported to be essential for the psychological effects induced by various psychedelics (Nichols, 2016), and in the case of MDMA, has been linked to perceptual and emotional alterations (Liechti et al., 2000, Kuypers et al., 2018). Broadband reduction in oscillatory power, triggered by $5\text{-HT}_{2A}\text{R}$, seems to be linked to the subjective effect of serotonergic drugs (Carhart-Harris et al., 2016, Carhart-Harris and Friston, 2019) and has been consistently observed in humans and rodents following administration of MDMA (Frei et al., 2001, Lansbergen et al., 2011) or various other $5\text{-HT}_{2A}\text{R}$ agonists (Kometer et al., 2015, Muthukumaraswamy et al., 2013, Carhart-Harris et al., 2016, Wood et al., 2012). The link between $5\text{-HT}_{2A}\text{R}$ and perception is further supported by the fact that several routinely used antipsychotic drugs are potent $5\text{-HT}_{2A}\text{R}$ antagonists (Marek et al., 2003, Meltzer, 1999). While the most recent attempts to explain the psychological effects of $5\text{-HT}_{2A}\text{R}$ activation focus on the increased spiking of cortical pyramidal neurons in the deep layers (Carhart-Harris and Friston, 2019, Nichols, 2016), our study suggests that Sst interneurons may also play a role. Sst interneurons, in contrast to PV interneurons, form synapses on the dendrites of their target cell (Tremblay et al., 2016). A wealth of evidence suggests that active dendritic processing in cortical pyramidal neurons has a critical influence on sensory perception (Takahashi et al., 2016, Murayama et al., 2009, Smith et al., 2013, Ranganathan et al., 2018) and, in accordance with their unique anatomical properties, Sst interneurons strongly influence dendritic computations and directly modulate perceptual thresholds (Takahashi et al., 2016).

We propose that the novel link between $5\text{-HT}_{2A}\text{R}$ and Sst interneurons might help elucidate the mechanisms underlying a host of psychiatric disorders and contribute to our understanding of how serotonergic drugs exert their psychological effects.

Materials and Methods

All experiments were conducted according to regulations of the Landesamt für Gesundheit und Soziales (Berlin [T 0100/03], Berlin [G0298/18]) and the European legislation (European Directive 2010/63/EU).

Animals

Data for the *in vivo* part of the study was collected from C57Bl/6 mice (aged 6 to 10 weeks). Data for the *in vitro* part was collected from C57Bl/6 (P10-P17), Sst-tdTomato (P10-P30), Sst-Chr2-EYFP (P10-P16), hM4Di-Sst (P10-P15), hM4Di-Sst (+/-) (P10-P15) and hM4Di-PV (P10-P15) mice. Immunostainings to localize 5-HT$_{2A}$R were performed on 5-HT$_{2A}$R-EGFP mice (P20-P90) and immunostainings to localize 5-HT fibers were performed on an ePet-YFP mouse (P35). Sst-Cre mice (RRID:IMSR_JAX:013044) have Cre recombinase targeted to the *Sst* locus; they were obtained from Jackson Laboratory (ME, USA). PV-Cre mice (RRID:IMSR_JAX:008069) have Cre recombinase targeted to the *Pvalb* locus; they were obtained from Charles River Laboratory (MA, USA). tdTomato mice (Ai9, RRID:IMSR_JAX:013044) were obtained from Jackson Laboratory. Chr2-EYFP mice (Ai32, RRID:IMSR_JAX:012569) were obtained from Jackson Laboratory. ePet-Cre mice (RRID:IMSR_JAX:012712) have Cre recombinase targeted to the *Fev* locus; they were obtained from Jackson Laboratory. 5-HT$_{2A}$R-EGFP mice (RRID:MMRRC_010915-UCD) express EGFP reporter protein under the control of the *Htr2a* gene, they were obtained from the Mutant Mouse Resource & Research Centers (MMRRC, CA, USA). Generation of hM4Di mice is described in the paragraph "Generation of Cre-conditional hM4Di mice". The animals were housed in a 12:12 hour light-dark cycle in singularly ventilated cages with *ad libitum* access to food and water. SOs *in vitro* recordings were performed on P12-P16 mice.

Generation of Cre-conditional hM4Di mice

We produced a transgenic mouse line carrying a Cre-conditional hM4Di expression cassette in the Rosa26 locus. The transgene construct was inserted by recombination-mediated cassette exchange (RMCE). RMCE relies on recombination events between attB and attP recognition sites of the RMCE plasmid and genetically modified acceptor embryonic stem (ES) cells, mediated by the integrase of phage phiC31 (Hitz et al., 2007). The RMCE construct is thereby shuttled into the Rosa26 locus of the ES cells, along with a Neomycin resistance cassette (fig. S12A). The acceptor cell line IDG3.2-R26.10-3 (I3) was kindly provided by Ralf

Kühn (GSF National Research Centre for Environment and Health, Institute of Developmental

Genetics, Neuherberg, Germany).

We subcloned a Cre-conditional FLEX (flip-excision) cassette (Schnutgen et al., 2003) into

pRMCE, and inserted a strong CAG promoter (CMV immediate early enhancer/modified

chicken β-actin promoter, from Addgene Plasmid #1378) in front of the FLEX-cassette to

create pRMCE-CAG-Flex. The coding sequence of hM4Di-mKateT was inserted into the FLEX

cassette in reverse orientation to the promoter (fig. S12A). Finally, a rabbit globulin polyA

cassette including stop codons in every reading frame was placed downstream of the FLEX

cassette, in the same direction as hM4Di, in order to prevent unintended transcriptional read-

through from potential endogenous promoters. The construct was completely sequenced

before ES cell electroporation.

Electroporation of the RMCE construct together with a plasmid encoding C31int was

performed by the transgene facility of the 'Research Institute for Experimental Medicine' (FEM,

Charité, Berlin) according to published protocols (Hitz et al., 2009, Hitz et al., 2007).

Recombinant clones were selected by incubation with 140 µg/ml G418 for at least 7 days. To

activate hM4Di expression by recombination of the FLEX switch, selected clones were further

transfected transiently with pCAG-Cre-EGFP using Roti-Fect (Carl Roth, Karlsruhe, Germany).

G418-resistant clones were analyzed by PCR for successful integration and recombination of

the construct (fig. S12B), using the following primers:

GT001　　　　　PGK3'-fw:　　　　　CACGCTTCAAAAGCGCACGTCTG;

GT002　　　　　Neo5'-rev:　　　　　GTTGTGCCCAGTCATAGCCGAATAG;

GT005　　　　　PolyA-fw:　　　　　TTCCTCCTCTCCTGACTACTCC;

GT006　　　　　Rosa3'-rev:　　　　　TAAGCCTGCCCAGAAGACTC;

GT013 hM4Di3'rec-rev: CAGATACTGCGACCTCCCTA

After verification of correct integration and functional FLEX-switch recombination, we

generated chimeras by blastocyst injection of I3 ES cells. Heterozygous offsprings were

mated with a Flpe deleter mouse line in order to remove the neomycin resistance cassette by

Flp-mediated recombination.

Mice homozygous for the Rosa-CAG-FLEX-hM4Di-mKateT allele are viable and fertile and

show now obvious phenotype. Importantly, application of CNO to these mice does not induce

any behavioral effects. Homozygous Cre-conditional hM4Di transgenic mice and Sst-Cre

mice (Taniguchi et al., 2011) were maintained on a C57Bl/6 genetic background and were

bred to obtain heterozygous Sst-Cre / hM4Di offsprings.

Drugs

Urethane (U2500, Merck), Fenfluramine ((+)-Fenfluramine hydrochloride, F112, Merck), 5-HT
(Serotonin creatinine sulfate monohydrate, H7752, Merck), m-CPBG (1-(3-
Chlorophenyl)biguanide hydrochloride, C144, Merck), tropisetron (Tropisetron
hydrochloride,Y0000616, Sigma), WAY-100635 (W108, Merck), α-Methylserotonin (α-
Methylserotonin maleate salt, M110, Merck) MDMA ((±)3,4-
methylenedioxymethamphetamine, 64057-70-1, Merck), CNO (Clozapine N-oxide
dihydrochloride, 6329, Tocris) were dissolved in water for *in vitro* application and in 0.9%
normal saline for *in vivo* application. Ketanserin (Ketanserin (+)-tartrate salt, S006, Merck) was
dissolved in Dimethyl sulfoxide (DMSO).

Surgery and *in vivo* recording

Before recordings mice were briefly anaesthetized with isofluorane (2%) and then injected
intraperitoneally with urethane (1.2 g/kg, Sigma Aldrich, Munich, Germany). The level of
anesthesia was maintained so that hindlimb pinching produced no reflex movement and
supplemental doses of urethane (0.2 g/kg) were delivered as needed. Upon cessation of
reflexes the animals were mounted on a stereotaxic frame (Kopf Instruments, Tujunga,
California), and body temperature was maintained at 38°C. The scalp was removed, and the
skull was cleaned with saline solution. A craniotomy was performed at +3 mm ML, -3 mm AP,
+3.25 mm DV to reach mEC.
Extracellular recordings from EC (Figure 1-2) were performed using a Cambridge Neurotech
(Cambridge, United Kingdom) silicon probe (64-channels (n = 15) or 32-channels (n = 3). The
recording electrode was painted with the fluorescent dye DiI (Thermo Fisher Scientific,
Schwerte, Germany) and then slowly lowered into the craniotomy using micromanipulators
(Luigs&Neumann, Ratingen, Germany) at a 25° angle AP (toward the posterior side of the
brain). The exposed brain was kept moist using saline solution. A ground wire connected to
the amplifier was placed in the saline solution covering the skull to eliminate noise. Brain
signals were recorded using a RHD2000 data acquisition system (Intan Technologies, Los
Angeles, California) and sampled at 20kHz. Recording quality was inspected on-line using the
open-source RHD2000 Interface Software. A supplementary dose of urethane (0.2 g/kg) was
injected right before the start of the recording to standardize anesthesia level across different
experiments and avoid the arise of theta activity in the first 30 minutes of recording.
Recordings began after a 10-minute waiting period in which clear upstates could consistently
be seen at a regular frequency.

In vivo analysis

We selected the channel to use for upstate detection based on the standard deviation (STD) of the trace during baseline (first 5 minutes of recording): the channel with the highest STD was selected, as larger voltage deflection increases detection algorithm accuracy. Given the highly synchronous nature of SOs (Supplementary Fig. 3) the spatial location of the channel selected was not considered. Upstates were detected comparing threshold crossing points in two signals: the delta-band filtered signal (0.5-4 Hz) and the population spike activity. Candidate upstates were identified in the delta-band filtered signal using two dynamic thresholds 'a' and 'b':

$$a = m + \frac{\sigma}{1.5}$$

$$b = m + \frac{\sigma}{0.8}$$

Where σ is the standard deviation of the signal during the first five minutes of recording (baseline) and *m* is the centered moving median calculated using 60 s windows (Matlab function *movmedian*). The median was used instead of the mean to account for non-stationaries in the data. A candidate upstate was identified at first using the threshold crossings of the signal compared to 'a': candidates shorter than 200 ms were deleted and multiple candidates occurring within a window of 300 ms were joined together. Subsequently the threshold 'b' was used to separate upstates occurring in close proximity: if the signal within one candidate crossed the threshold 'b' in more than one point then the candidate upstate was split in two at the midpoint between the two threshold crossings. Candidate upstates were finally confirmed if the population spike activity (calculated in 100 ms windows) within the candidate crossed a threshold of 1 σ (calculated during the baseline).

Units detection and classification

Spike detection was performed offline using the template-based algorithm Kilosort2 (https://github.com/MouseLand/Kilosort2), with the following parameters:

- ops.fshigh = 300
- ops.fsslow = 8000
- ops.minfr_goodchannels = 0
- ops.Th = [8 4]
- ops.lam = 10

- ops.AUCsplit = 0.9
- ops.minFR = 1/1000
- ops.momentum = [20 400]
- ops.sigmaMask = 30
- ops.ThPre = 8
- ops.spkTh = -6
- ops.nfilt_factor = 8
- ops.loc_range = [3 1]
- ops.criterionNoiseChannel = 0.2
- ops.whiteningrange = 32
- ops.ntbuff = 64

Spike sorting in recordings performed with Neuropixel probes was perfomed using ecephys_spike_sorting (https://github.com/AllenInstitute/ecephys_spike_sorting). Manual curation of the results was performed using Phy (https://github.com/cortex-lab/phy). Each Isolated unit satisfied the following two criteria: Refractory period (2 ms) violations < 5%, fraction of spikes below detection threshold (as estimated by a gaussian fit to the distribution of the spike amplitudes) < 15%. Units with negative maximal waveform amplitude were further classified as putative excitatory if the latency (TP latency) was > 0.55 ms or putative inhibitory when TP latency < 0.55 ms. The value 0.55 was chosen in accordance with previous studies (Senzai et al., 2019, Antoine et al., 2019). In pharmacological classification, units were classified as 'activated' if their firing rate in the 25 minutes following drug injection was 2 σ (standard deviation) above the baseline rate for at least 5 minutes. Remaining units were pulled together in the category 'non-activated'.

Cross-correlogram analysis

Cross-correlogram based connectivity analysis was performed for every unit to identify inhibitory connections. Units with a spiking rate smaller than 0.3 spikes/s were discarded from the analysis. We used total spiking probability edges (TPSE) algorithm (https://github.com/biomemsLAB/TSPE) (De Blasi et al., 2019) to identify in a computationally efficient manner putative inhibitory connections between units and all clusters recorded. The parameters used were:

•   d = 0,

•   neg_wins = [2, 3, 4, 5, 6, 7, 8],

•   co_wins = 0,

•   pos_wins = [2, 3, 4, 5, 6],

•   FLAG_NORM = 1.

The connectivity vectors of each unit resulting from TSPE were sorted by inhibition

strength. The top 20 connections were further analyzed using custom Matlab

(RRID:SCR_001622) code. A connection was classified as inhibitory if the cross correlogram

values (x) were smaller than the mean of x by more than one standard deviation ($x <$ mean(x)

– std(x)) in at least 4 consecutive bins (bin size = 1 ms) in a window 4 to 9 ms after the center

of the cross-correlogram.

Slice preparation

We prepared acute near horizontal slices (~15° off the horizontal plane) of the medial

entorhinal cortex (mEC) from C57Bl/6 mice. Animals were decapitated following isoflurane

anesthesia. The brains were quickly removed and placed in ice-cold (~4° C) ACSF (pH 7.4)

containing (in mM) 85 NaCl, 25 NaHCO3, 75 Sucrose, 10 Glucose, 2.5 KCl, 1.25 NaH2PO4,

3.5MgSO4, 0.5 CaCl2, and aerated with 95% O2, 5% CO2. Tissue blocks containing the brain

region of interest were mounted on a vibratome (Leica VT 1200, Leica Microsystems), cut at

400 μm thickness, and incubated at 35 °C for 30 min. The slices were then transferred to

ACSF containing (in mM) 85 NaCl, 25 NaHCO3, 75 Sucrose, 10 Glucose, 2.5 KCl, 1.25

NaH2PO4, 3.5 MgSO4, 0.5 CaCl2. The slices were stored at room temperature in a

submerged chamber for 1-5 hr before being transferred to the recording chamber.

***In vitro* recording**

In order to perform whole-cell recordings slices were transferred to a submersion style

recording chamber located on the stage of an upright, fixed-stage microscope (BX51WI,

Olympus) equipped with a water immersion objective (×60, Olympus) and a near-infrared

charge-coupled device (CCD) camera. The slices were perfused with ACSF (~35 °C bubbled

with 95 % O2-5 % CO2) at 3-5 ml/ min to maintain neuronal health throughout the slice. The

ACSF had the same composition as the incubation solution except for the concentrations of

calcium and magnesium, which were reduced to 1.2 and 1.0 mM, respectively. Recording

electrodes with impedance of 3-5 MΩ were pulled from borosilicate glass capillaries (Harvard

Apparatus, Kent, UK; 1.5 mm OD) using a micropipette electrode puller (DMZ Universal Puller).

The intracellular solution contained the following (in mM): 135 K-gluconate, 6 KCl, 2 MgCl2,

0.2 EGTA, 5 Na2- phosphocreatine, 2 Na2-ATP, 0.5 Na2-GTP, 10 HEPES buffer, and 0.2%

biocytin. The pH was adjusted to 7.2 with KOH. Recordings were performed using Multiclamp

700A/B amplifiers (Molecular Devices, San Jose, California). The seal resistance was >1 GΩ.

Capacitance compensation was maximal and bridge balance adjusted. Access resistance

was constantly monitored. Signals were filtered at 6 kHz, sampled at 20 kHz, and digitized

using the Digidata 1550 and pClamp 10 (Molecular Devices, San Jose, California). Activation

light was delivered by a 460 nm laser (DPSS lasers, Santa Clara, California) using a 460–480

nm bandpass excitation filter. Stimulation consisted of 500 ms pulses at 1 Hz.

Stimulation experiments were performed using a bipolar micro-electrode (glass pipette filled

with ACSF solution, wrapped by a fine grounding wire) connected to an isolated voltage

stimulator (ISO-Flex, A.M.P.I., Israel). A 4x objective (Olympus) was used to visually guide the

stimulating electrode into the mEC. Stimulation power was adjusted to achieve consistent

upstate generation during baseline (> 95%). Each stimulus had a duration of 50 μs and the

inter-stimulus interval was 8-10 seconds.

In vitro analysis

In vitro upstates were detected in Matlab using an algorithm similar to the one described in

the *in vivo analysis* method section. We used a coincident detection in two signals. In

multicellular recordings we used the membrane potential of 2 cells, in single cell recordings

we used membrane potential and the envelope of the gamma filtered trace (50-250 Hz), as

upstates are characterized by an increase in gamma activity (Neske, 2015).

The baseline condition was defined as the last 120 s before drug application, while the post-

drug application condition was defined as the 120 s of recording after drug application (Total

recording duration: 600 s).

Excitatory (Exc), fast-spiking (FS) and low-threshold spiking (LTS) neurons were classified

using Gaussian mixture models (GMM) with a soft clustering approach in Matlab. Input

resistance (R_{in}), Δafter-hyperpolarization (ΔAHP), sag, rheobase, spike width and resting

potential (RP) were extracted from each neuron and used in the classification. The first two

components of the principal component analysis (PCA) were used to fit the data to a Gaussian

mixture model distribution. Initial values were set according to the k-means algorithm with

centroid positioned at x and y position: 5, 0; -15, -15; -15, 10. This centroid were placed
according to the loadings of the PCA to identify 3 clusters with the following main features:

• Cluster 1 (putative Exc): high spike width, low AHP, low rheobase.

• Cluster 2 (putative FS): low spike width, low SAG, high rheobase, low R_{in}.

• Cluster 3 (putative LTS): low spike width, high SAG, high AHP, high R_{in}.

Covariance matrices were diagonal and not shared. Neurons with a posterior probability of
belonging to any of the three clusters of < 90% were discarded from further analysis (1/49).
While the majority of Sst-interneurons display LTS features, a minority (~10%) belong to the
FS group (Urban-Ciecko et al., 2015). To distinguish FS and LTS interneurons in the Sst-Td
Tomato mice, we employed the GMM with posterior probability threshold of 90%. Only LTS
Sst neurons were considered for further analysis.

Histological analysis

For the *post-mortem* electrode track reconstructions of the *in vivo* recordings, mice were not
perfused; rather, brains were extracted from the skull, post-fixed in 4% PFA overnight at 4°C
and afterwards cut with a vibratome (Leica Microsystems, Wetzlar Germany) in 100 µM thick
sequential sagittal slices. Images were taken using a 1.25x objective and stitched together
using the microscope software (BX61, Olympus). Afterwards we used AllenCCF code
(https://github.com/cortex-lab/allenCCF) to identify electrode shank location (Shamash et al.,
2018).
For the anatomical reconstructions of recorded cells *in vitro,* brain slices were fixed with 4%
paraformaldehyde in 0.1 M phosphate buffer (PB) for at least 24 hours at 4°C. After being
washed three times in 0.1 M PBS, slices were then incubated in PBS containing 1% Triton X-
100 and 5% normal goat serum for 4 hr at room temperature (RT). To visualize biocytin-filled
cells we used Streptavidin Alexa 488 conjugate (1:500, Invitrogen Corporation, Carlsbad, CA,
RRID:AB_2315383). WFS1 (1:1000, Rabbit, Proteintech, IL, USA, RRID:AB_2880717) was
used in a subset of experiments to visualize the L2/L3 border, and Sst (1:1000, Rat, Bachem,
Switzerland, RRID:AB_2890072) was used in the 5-HT$_{2A}$R localization analysis. Slices were
incubated with primary antibodies for 48 hours at RT. After rinsing two times in PBS, sections
were incubated in the PBS solution containing 0.5% Triton X-100, Alexa fluor 488, Alexa fluor
555 and Alexa fluor 647 (Invitrogen Corporation, Carlsbad, CA) according to the number of

antibodies used. Slices were mounted in Fluoroshield (Sigma-Aldrich) under coverslips 2-3 hr after incubation with the secondary antibodies and stored at 4 °C.

Labeled cells were visualized using 20x or 40x objectives on a confocal microscope system (SP8, Leica, RRID:SCR_018169). For the 5-HT$_{2A}$R localization analysis, images of the whole EC were acquired and stitched together using the auto stitching method, without smoothing. Z stacks were acquired every 30 µM. The image stacks obtained were registered and combined in Fiji (RRID:SCR_002285) to form a montage of the sections. Cell counting was executed using the Fiji multi-point tool. X-Y-Z coordinates of each 5-HT$_{2A}$R-EGFP positive cell were exported to Matlab and subsequently, using custom written code in Matlab, we semi-automatically inspected each cell for colocalization between EGFP(5-HT$_{2A}$R) and Sst.

Statistical Analysis

All datasets were tested to determine normality of the distribution either using D'Agostino-Pearson omnibus normality test or Shapiro-Wilk normality test. Student's t-test and one-way ANOVA were used for testing mean differences in normally distributed data. Wilcoxon matched-pairs signed rank test and Kruskal-Wallis were used for non-normally distributed datasets. Dunn-Sidak multiple comparison test was used to compare datasets with 3 or more groups. Kolmogorov-Smirnov test was used to compare cumulative distributions. Statistical analysis was performed using Prism (6.01, RRID:SCR_002798) and Matlab. All data are expressed as mean ± SEM. Asterisks in figures represent p-values smaller than 0.05, unless stated otherwise in the legend.

Supplementary Figures

 Figure 1–figure supplement 1. *In vivo* upstate spatial features.

 (A) Microelectrode implant location and microelectrode features. 64 channels ($n_{animals}$= 15,

 n_{shanks} = 4) and 32 channels ($n_{animals}$ = 3, n_{shanks} = 2) microelectrodes were used in this study,

 analysis shown in this figure excludes data recorded with 32 channels probe due to the

 different spatial configuration of the channels. **(B)** Average upstate voltage deflection for each

 channel of the microelectrode. For experiments with drug application (either MDMA or Fen),

 only baseline upstates were taken in account. **(C)** Left: Average upstate voltage deflection

 grouped by shank. Right: Average upstate voltage deflection grouped by depths (right). Insets

 show the normalized correlation between averages in the two different groups.

Figure 1–figure supplement 2. LFP power analysis for saline and MDMA/Fen injection.

(A) Average normalized LFP power during baseline and saline injection. **(B)** Average LFP power during baseline and MDMA/Fen injection. **(C)** Box plots of normalized LFP power per frequency band: delta (Control: 0.98 ± 0.01, MDMA/Fen: 0.82 ± 0.04, $p = 0.019$, Mann-Whitney test), theta (Control: 0.90 ± 0.01 µV, MDMA/Fen: 0.76 ± 0.04, $p = 0.046$, Mann-Whitney test), alpha (Control: 0.77 ± 0.01, MDMA/ Fen: 0.45 ± 0.03, $p > 0.05$, Mann-Whitney test), beta(Control: 0.49 ± 0.02, MDMA/ Fen: 0.30 ± 0.03, $p > 0.05$, Mann-Whitney test), and gamma (Control: 0.28 ± 0.02, MDMA/ Fen: 0.25 ± 0.02, $p > 0.5$, Mann-Whitney test).

Figure 1–figure supplement 3. *In vivo* **upstate metrics for saline and MDMA/Fen injection.**

(A) Average upstate voltage deflection for control and MDMA/Fen, patches represent 1 standard deviation. Scale bars: 0.5 seconds, 200 μV. **(B)** Violin plots of duration (control: 0.98 ± 0.02 s, MDMA/ Fen: 0.69 ± 0.04 s, p < 0.001, Mann-Whitney test). **(C)** Violin plots of peak 1 amplitude (control: 434.3 ± 13.09 μV, MDMA/Fen: 248.1 ± 10.49 μV, p < 0.001, Mann-Whitney test). **(D)** Violin plots of peak 2 amplitude (control: 232.4 ± 3.53 μV, MDMA/ Fen: 187.3 ± 7.40 μV, p < 0.001, Mann-Whitney test). **(E)** Violin plots of area (control: 185.6 ± 4.14 μV•s, MDMA/ Fen: 187.3 ± 7.40 μV, p < 0.001, Mann-Whitney test). **(F)** Violin plots of inter-event intervals (control: 5.55 ± 0.26 s, MDMA/ Fen: 12.02 ± 1.75 2, p < 0.001, Mann-Whitney test).

 Figure 2–figure supplement 1. Cross-correlogram (CCG) based connectivity analysis.
 (A) Units are plotted according to TP latencies and repolarization time and color-coded
 according to the number of inhibitory connections detected. Units displaying a TP latency <
 0.55 ms were classified as putative inhibitory interneurons ('Putative int'), the remaining units
 were classified as putative excitatory neurons ('Putative exc'). Inhibitory connections were
 detected using Total Spiking Probability Edges (TSPE) (See Supplementary materials).
 Putative interneurons had a 40.0 % chance of displaying at least one inhibitory connection in
 the CCGs with an average number of 3.38 ± 0.68 inhibitory connections while putative
 excitatory cells had a 0.33 % chance of displaying inhibitory connections. Units with a spiking

rate lower than 0.3 spikes/s were discarded from the analysis. Units from control experiments
are included in the analysis. **(B)** Connectivity scheme of one putative inhibitory unit (source
unit, black circle) displaying 10 inhibitory connections with surrounding clusters. Grey rhombi
represent recording channels on the probe with the tip of the shanks pointing north. Each
colored circle represents the location of an inhibited unit. Waveforms of the inhibited units are
plotted nearby the location using the same color. Inset shows the location of the source unit
on the probe. **(C)** CCGs of the connections displayed in (B) using the same color code. Solid
lines represent mean, dashed lines represent 1 standard deviation.

B

Gof metrics	Excitatory	Inhibitory	Activated
sse	0.6764	0.3036	0.5860
rsquare	0.9340	0.9669	0.9353
dfe	97	97	97
adjrsquare	0.9320	0.9659	0.9333

Figure 2–figure supplement 2. TP latency density distributions.
(A) Dashed lines represent kernel density estimations of probability density functions of the
TP latencies of putative inhibitory (blue), putative excitatory (black) and 'activated' units. Solid
lines represent gaussian fitting curves for each group. Peak inhibitory gaussian: 0.31 ms, peak
excitatory gaussian: 0.81 ms and peak 'activated' gaussian: 0.56 ms. **(B)** Goodness of fit
metrics for each gaussian fit. Sse = Sum of squares due to error, rsquare = R-squared
(coefficient of determination), dfe = Degrees of freedom in the error, adjrsquare = Degree-of-
freedom adjusted coefficient of determination.

Figure 3–figure supplement 1. Effect of Fen on SOs *in vitro*.

(A) Top: upstate raster plot during Fen application. Bottom: Histogram of upstate incidence

before and after Fen application (n= 6, baseline: 1.13 ± 0.17 upstates/10s, Fen: 0.04 ± 0.03

upstates/10s). **(B)** Top: upstate raster plot during ketanserin + Fen application. Bottom:

Histogram of upstate incidence before and after ketanserin + Fen application (n= 5, baseline:

1.18 ± 0.07 upstates/10s, Fen: 1.03 ± 0.17 upstates/10s). **(C)** Upstate metrics during baseline

and ketanserin + Fen. Left: violin plots of inter-event interval (baseline = 9.46 ± 0.28 s,

ketanserin + Fen = 9.63 ± 0.74 s). Middle: violin plots of inter-event upstate duration (baseline

= 2.13 ± 0.08 s, ketanserin + Fen = 2.21 ± 0.05 s). Right: violin plots of upstates area (baseline

= 45.55 ± 2.45 mV•s, ketanserin + Fen = 39.45 ± 2.92 mV•s).

Figure 3–figure supplement 2. 5-HT suppresses evoked upstates *in vitro*.
(A) Experimental protocol: recording and stimulation electrode were placed in mEC layer 3,
stimulation electrode was located towards the lateral side of the slice. **(B)** Effect of electrical
stimulation before (black) and after 5-HT application (orange). 5-HT consistently suppressed
spiking. Increasing the stimulation power up to 10x (n = 40/80 in 4 slices) had no rescue effect.
Top: voltage responses to electrical stimulation of a representative neuron. Middle: summary
spike raster plot before and after 5-HT application. Bottom: spike rate line histogram **(C)**
Scatter plot showing area (top, n = 8 neurons, mean $_{control}$ = 16.22 ± 0.80, , mean$_{5-HT}$ = 1.24 ±
0.97, p < 10^{-4}, Wilcoxon matched-pairs signed rank test) and duration (bottom, n = 8 neurons,
mean $_{control}$ = 1.92 ± 0.07, , mean$_{5-HT}$ = 0.24 ± 0.01, , p < 10^{-4}, Wilcoxon matched-pairs signed
rank test) of evoked upstates before (black) and after 5-HT application (orange). **(D)**
Representative voltage responses to 1 second 4 Hz stimulation following wash-in (left) and
wash-out (right) of 5-HT. 5-HT prevents spiking from input summation.

Figure 3–figure supplement 3. 5-HT$_{2A}$Rs are involved in 5-HT mediated SOs suppression *in vitro*.

(A) Top: upstate raster plot during application of WAY 100635 (5-HT$_{1A}$ antagonist) + 5-HT. Bottom: change in RP in putative excitatory cells after application of WAY 100635 (5-HT$_{1A}$ antagonist) + 5-HT (n = 25 cells). **(B)** Top: upstate raster plot during application of ketanserin (5-HT$_{2A}$ antagonist) + 5-HT. Bottom: change in RP in putative excitatory cells after application of ketanserin (5-HT$_{2A}$ antagonist) + 5-HT (n = 21 cells). **(C)** Top: upstate raster plot during application of α-methyl-5-HT (5-HT$_2$ agonist). Bottom: change in RP in putative excitatory cells after application of α-methyl-5-HT (5-HT$_2$ agonist) (n = 11 cells). **(D)** Dot plot showing change in RP for each pharmacological condition (5-HT: -4.52 ± 0.64 mV, WAY + 5-HT: -2.09 ± 0.47 mV, ketanserin + 5-HT: -3.68 ± 0.60 mV and α-methyl-5-HT: -1.67 ± 1.13 mV; p = 0.0329,

766   Kruskal-Wallis with Dunn's multiple comparisons test). **(E)** Dot plot showing percentage
767   reduction of upstate incidence for each pharmacological condition (5-HT: 95 ± 4 %, WAY +
768   5-HT: 100 ± 0 %, ketanserin + 5-HT: 57 ± 10.1 % and α-methyl-5-HT: 100 ± 0 %; $p < 10^{-4}$,
769   Kruskal-Wallis with Dunn's multiple comparisons test).

Figure 3–figure supplement 4. 5-HT$_3$ receptor is not involved in 5-HT mediated SOs suppression.

(A) Top: upstate raster plot during application of m-CPBG (5-HT3 agonist). Bottom: histogram of upstate incidence before and after m-CPBG application (n= 6, baseline: 1.04 ± 0.19 upstates/10s, m-CPBG: 1.11 ± 0.21 upstates/10s). Patches represent 95% confidence interval, lines represent standard deviation. (B) Top: upstate raster plot during application of tropisetron (5-HT3 antagonist) + 5-HT. Bottom: histogram of upstate incidence before and after tropisetron + 5-HT application (n= 6, baseline: 1.23 ± 0.18 upstates/10s, tropisetron + 5-HT: 0.01 ± 0.01 upstates/10s). patches represent 95% confidence interval, lines represent standard deviation.

Figure 3–figure supplement 5. *In vitro* **upstates metrics during baseline and ketanserin + 5-HT application.**

(A) Box plot showing spiking rate before and after ketanserin + 5-HT (baseline: 0.38 ± 0.14 spikes/s, ketanserin + 5-HT: 0.15 ± 0.06 spikes/s, p= 0.011, Mann-Whitney test). patches represent 95% confidence interval, lines represent standard deviation. **(B)** Violin plots of upstate inter-event interval (baseline = 10.92 ± 0.34 s, ketanserin+5-HT = 15.28 ± 1.34 s, p = 0.071, Mann-Whitney test). **(C)** Violin plots of upstate duration (baseline = 3.11 ± 0.16 s, ketanserin +5-HT = 1.91 ± 0.15 s, p < 0.001, Mann-Whitney test). **(D)** Violin plots of upstate area (baseline = 31.55 ± 1.13 mV•s, ketanserin+5-HT = 28.16 ± 1.91 mV•s, p = 0.0825, Mann-Whitney test).

Figure 3–figure supplement 6. LTS neurons are depolarized by 5-HT.
(A) PCA projection plot of all the cells recorded. Cells are color-coded according to group
identity (posterior probability > 0.9): excitatory (black), fast-spiking (blue), low-threshold
spiking (green). Red circles represent PCA loadings. **(B)** PCA projection plot color-coded
according to ΔRP after 5-HT application. Inset shows a recording from one LTS neuron during
5-HT application. Scale bars: 20 mV, 25 s. Dotted line showing -70 mV. **(C)** PCA projection
plot color-coded according to Δspikes/s after 5-HT application. **(D)** Posterior probability of
being classified as Exc, FS or LTS.

A

B

	LTS	FS	Exc
Rin	277.97 ± 57.52	69.38 ± 13.20	135.77 ± 18.47
ΔAHP	10.68 ± 2.38	3.78 ± 4.46	-0.42 ± 1.00
SAG	16.70 ± 5.03	1.17 ± 1.16	6.84 ± 2.40
Rheobase	61.11 ± 14.40	266.67 ± 60.23	76.47 ± 22.37
Spike width	0.71 ± 0.11	0.52 ± 0.09	1.43 ± 0.17
RP	-65.03 ± 3.87	-69.13 ± 8.36	-64.80 ± 2.08

Figure 3–figure supplement 7. Excitatory, fast-spiking and LTS cells have unique sets of
electrophysiological features.
(A) Box plot showing the values of input resistance (R_{in}), delta after-hyperpolarization (ΔAHP),
SAG, RP, rheobase and spike width of excitatory (Exc, black), fast-spiking (FS, blue) and low-
threshold spiking (LTS, green) cells ($n_{EXC} = 33$, $n_{FS} = 6$, $n_{LTS} = 9$; asterisk means $p < 0.05$, double

asterisk means p < 0.01). patches represent 95% confidence interval, lines represent standard
deviation. **(B)** Table showing values plotted in (A).

Figure 3–figure supplement 8. Classification of cells recorded in Sst-tdTomato mice.
(A) PCA projection plot of cells recorded in Sst-tdTomato mice. Cells are color-coded
according to group identity (posterior probability > 0.9): fast-spiking (blue) or low-threshold
spiking (green). Red circles represent PCA loadings. **(B)** Posterior probability of being
classified as FS. **(C)** Posterior probability of being classified as LTS.

 Figure 3–figure supplement 9. Spatial localization of 5-HT$_{2A}$R positive cells in EC.

(A) 3D visualization of EC (purple). **(B)** 3D localization of all the 5-HT$_{2A}$R positive cells detected
in EC using same perspective as (A). **(C** and **D)** Co-localization of 5-HT$_{2A}$R and Sst, arrows
point to colocalized cells (scale bar: 100 μm, n $_{animals}$= 7, total number of 5-HT$_{2A}$R positive cells
= 3570, average number of 5-HT$_{2A}$R positive cells per animal = 510 ± 80.32). **(E)** Spatial
distribution of 5-HT$_{2A}$R positive cells and colocalized cells along the 3 dimensions depicted in
(A) (Z dimension centered on the midline).

(**A**) The coding sequence of hM4Di-mKate flanked by two opposing *loxP* and *lox2272* sites

was placed in reverse orientation to the CAG- promoter in the pRMCE. In the acceptor ES

cells the ROSA26 allele harbours a PGK promoter driving the hygromycin selection marker,

flanked by two attP sites. RMCE by C31int replaces the hygromycin resistance by the

neomycin resistance of the donor vector. Location of primer binding sites in the Rosa26-

hM4Di locus is indicated by green triangles. (**B**) Identification of successful genomic

integration events and Cre-mediated inversion of the hM4Di coding sequence by PCRs. PCR

1 and PCR 2 test for correct integration of the 5' and 3' end of the construct into the ROSA26

locus. The lower band in PCR 1 results from the Neomycin resistance cassette of the feeder

cells in the ES cell culture. PCR 3 tests for successful recombination of the FLEX site by Cre.

A successful Cre-mediated recombination of the FLEX cassette was observed for clone 1,

resulting in an 826 bp product in PCR 3. C: control cells (not electroporated), H: H_2O input.

Figure 5–figure supplement 1. CNO application prevents 5-HT mediated upstates
suppression.
(A) Upstate raster plot during CNO and subsequent 5_HT application. Orange box represents
5-HT, purple boxes represent CNO. CNO significantly prevents 5-HT-induced suppression of
upstates. **(B)** Upstate incidence during 5-HT, CNO and 5-HT+CNO application (n=5, baseline:
1.13 ± 0.13 upstates/10s, CNO: 1.13 ± 0.10 upstates/10s, 5-HT: 0.93 ± 0.24 upstates/10s).
Patches represent 95% confidence intervals, lines represent standard deviation.

References

ABI-SAAB, W. M., BUBSER, M., ROTH, R. H. & DEUTCH, A. Y. 1999. 5-HT2 receptor regulation of extracellular GABA levels in the prefrontal cortex. *Neuropsychopharmacology,* 20**,** 92-6.

ADESNIK, H., BRUNS, W., TANIGUCHI, H., HUANG, Z. J. & SCANZIANI, M. 2012. A neural circuit for spatial summation in visual cortex. *Nature,* 490**,** 226-231.

ANTOINE, M. W., LANGBERG, T., SCHNEPEL, P. & FELDMAN, D. E. 2019. Increased Excitation-Inhibition Ratio Stabilizes Synapse and Circuit Excitability in Four Autism Mouse Models. *Neuron,* 101**,** 648-661.e4.

ARMBRUSTER, B. N., LI, X., PAUSCH, M. H., HERLITZE, S. & ROTH, B. L. 2007. Evolving the lock to fit the key to create a family of G protein-coupled receptors potently activated by an inert ligand. *Proc Natl Acad Sci U S A,* 104**,** 5163-8.

ASHBY, C. R., JR., JIANG, L. H., KASSER, R. J. & WANG, R. Y. 1990. Electrophysiological characterization of 5-hydroxytryptamine2 receptors in the rat medial prefrontal cortex. *J Pharmacol Exp Ther,* 252**,** 171-8.

ATHILINGAM, J. C., BEN-SHALOM, R., KEESHEN, C. M., SOHAL, V. S. & BENDER, K. J. 2017. Serotonin enhances excitability and gamma frequency temporal integration in mouse prefrontal fast-spiking interneurons. *eLife,* 6**,** e31991.

AZIMI, Z., BARZAN, R., SPOIDA, K., SURDIN, T., WOLLENWEBER, P., MARK, M. D., HERLITZE, S. & JANCKE, D. 2020. Separable gain control of ongoing and evoked activity in the visual cortex by serotonergic input. *eLife,* 9**,** e53552.

BARTHÓ, P., HIRASE, H., MONCONDUIT, L., ZUGARO, M., HARRIS, K. D. & BUZSÁKI, G. 2004. Characterization of neocortical principal cells and interneurons by network interactions and extracellular features. *J Neurophysiol,* 92**,** 600-8.

BAUMANN, M. H., AYESTAS, M. A., DERSCH, C. M., BROCKINGTON, A., RICE, K. C. & ROTHMAN, R. B. 2000. Effects of phentermine and fenfluramine on extracellular dopamine and serotonin in rat nucleus accumbens: therapeutic implications. *Synapse,* 36**,** 102-13.

BEED, P., DE FILIPPO, R., HOLMAN, C., JOHENNING, F. W., LEIBOLD, C., CAPUTI, A., MONYER, H. & SCHMITZ, D. 2020. Layer 3 Pyramidal Cells in the Medial Entorhinal Cortex Orchestrate Up-Down States and Entrain the Deep Layers Differentially. *Cell Reports,* 33**,** 108470.

BEIERLEIN, M., GIBSON, J. R. & CONNORS, B. W. 2003. Two dynamically distinct inhibitory networks in layer 4 of the neocortex. *J Neurophysiol,* 90**,** 2987-3000.

CARDIN, J. A. 2019. Functional flexibility in cortical circuits. *Curr Opin Neurobiol,* 58**,** 175-180.

CARHART-HARRIS, R. L. & FRISTON, K. J. 2019. REBUS and the Anarchic Brain: Toward a Unified Model of the Brain Action of Psychedelics. *Pharmacol Rev,* 71**,** 316-344.

CARHART-HARRIS, R. L., MUTHUKUMARASWAMY, S., ROSEMAN, L., KAELEN, M.,
DROOG, W., MURPHY, K., TAGLIAZUCCHI, E., SCHENBERG, E. E., NEST, T.,
ORBAN, C., LEECH, R., WILLIAMS, L. T., WILLIAMS, T. M., BOLSTRIDGE, M., SESSA,
B., MCGONIGLE, J., SERENO, M. I., NICHOLS, D., HELLYER, P. J., HOBDEN, P.,
EVANS, J., SINGH, K. D., WISE, R. G., CURRAN, H. V., FEILDING, A. & NUTT, D. J.
2016. Neural correlates of the LSD experience revealed by multimodal neuroimaging.
Proc Natl Acad Sci U S A, 113**,** 4853-8.

CARHART-HARRIS, R. L. & NUTT, D. J. 2017. Serotonin and brain function: a tale of two
receptors. *J Psychopharmacol***,** 269881117725915.

CHALMERS, D. T. & WATSON, S. J. 1991. Comparative anatomical distribution of 5-HT1A
receptor mRNA and 5-HT1A binding in rat brain--a combined in situ hybridisation/in
vitro receptor autoradiographic study. *Brain Res,* 561**,** 51-60.

CHEN, N., SUGIHARA, H. & SUR, M. 2015. An acetylcholine-activated microcircuit drives
temporal dynamics of cortical activity. *Nat Neurosci,* 18**,** 892-902.

CRAIG, M. T., MAYNE, E. W., BETTLER, B., PAULSEN, O. & MCBAIN, C. J. 2013. Distinct
roles of GABAB1a- and GABAB1b-containing GABAB receptors in spontaneous and
evoked termination of persistent cortical activity. *J Physiol,* 591**,** 835-43.

DE BLASI, S., CIBA, M., BAHMER, A. & THIELEMANN, C. 2019. Total spiking probability
edges: A cross-correlation based method for effective connectivity estimation of
cortical spiking neurons. *J Neurosci Methods,* 312**,** 169-181.

DESCARRIES, L., RIAD, M. & PARENT, M. 2010. CHAPTER 1.4 - Ultrastructure of the
Serotonin Innervation in the Mammalian Central Nervous System. *In:* MÜLLER, C. P. &
JACOBS, B. L. (eds.) *Handbook of Behavioral Neuroscience.* Elsevier.

DESIKAN, S., KOSER, D. E., NEITZ, A. & MONYER, H. 2018. Target selectivity of septal
cholinergic neurons in the medial and lateral entorhinal cortex. *Proceedings of the
National Academy of Sciences,* 115**,** E2644-E2652.

FANSELOW, E. E., RICHARDSON, K. A. & CONNORS, B. W. 2008. Selective, state-
dependent activation of somatostatin-expressing inhibitory interneurons in mouse
neocortex. *J Neurophysiol,* 100**,** 2640-52.

FREI, E., GAMMA, A., PASCUAL-MARQUI, R., LEHMANN, D., HELL, D. & VOLLENWEIDER,
F. X. 2001. Localization of MDMA-induced brain activity in healthy volunteers using low
resolution brain electromagnetic tomography (LORETA). *Hum Brain Mapp,* 14**,** 152-65.

FUNK, C. M., PEELMAN, K., BELLESI, M., MARSHALL, W., CIRELLI, C. & TONONI, G. 2017.
Role of Somatostatin-Positive Cortical Interneurons in the Generation of Sleep Slow
Waves. *J Neurosci,* 37**,** 9132-9148.

GIBSON, J. R., BEIERLEIN, M. & CONNORS, B. W. 1999. Two networks of electrically coupled
inhibitory neurons in neocortex. *Nature,* 402**,** 75-9.

GOŁEMBIOWSKA, K., JURCZAK, A., KAMIŃSKA, K., NOWORYTA-SOKOŁOWSKA, K. &
GÓRSKA, A. 2016. Effect of Some Psychoactive Drugs Used as 'Legal Highs' on Brain
Neurotransmitters. *Neurotox Res,* 29**,** 394-407.

GORINSKI, N., BIJATA, M., PRASAD, S., WIRTH, A., ABDEL GALIL, D., ZEUG, A., BAZOVKINA, D., KONDAUROVA, E., KULIKOVA, E., ILCHIBAEVA, T., ZAREBA-KOZIOL, M., PAPALEO, F., SCHEGGIA, D., KOCHLAMAZASHVILI, G., DITYATEV, A., SMYTH, I., KRZYSTYNIAK, A., WLODARCZYK, J., RICHTER, D. W., STREKALOVA, T., SIGRIST, S., BANG, C., HOBUS, L., FIEDLER, J., THUM, T., NAUMENKO, V. S., PANDEY, G. & PONIMASKIN, E. 2019. Attenuated palmitoylation of serotonin receptor 5-HT1A affects receptor function and contributes to depression-like behaviors. *Nature Communications,* 10**,** 3924.

GRANDJEAN, J., CORCOBA, A., KAHN, M. C., UPTON, A. L., DENERIS, E. S., SEIFRITZ, E., HELMCHEN, F., MANN, E. O., RUDIN, M. & SAAB, B. J. 2019. A brain-wide functional map of the serotonergic responses to acute stress and fluoxetine. *Nat Commun,* 10**,** 350.

GREEN, A. R., MECHAN, A. O., ELLIOTT, J. M., O'SHEA, E. & COLADO, M. I. 2003. The Pharmacology and Clinical Pharmacology of 3,4-Methylenedioxymethamphetamine (MDMA, "Ecstasy"). *Pharmacological Reviews,* 55**,** 463-508.

HARDEBO, J. E. & OWMAN, C. 1980. Barrier mechanisms for neurotransmitter monoamines and their precursors at the blood-brain interface. *Ann Neurol,* 8**,** 1-31.

HARDY, A., PALOUZIER-PAULIGNAN, B., DUCHAMP, A., ROYET, J. P. & DUCHAMP-VIRET, P. 2005. 5-Hydroxytryptamine action in the rat olfactory bulb: in vitro electrophysiological patch-clamp recordings of juxtaglomerular and mitral cells. *Neuroscience,* 131**,** 717-31.

HARRIS, K. D. & THIELE, A. 2011. Cortical state and attention. *Nat Rev Neurosci,* 12**,** 509-23.

HEIFETS, B. D., SALGADO, J. S., TAYLOR, M. D., HOERBELT, P., CARDOZO PINTO, D. F., STEINBERG, E. E., WALSH, J. J., SZE, J. Y. & MALENKA, R. C. 2019. Distinct neural mechanisms for the prosocial and rewarding properties of MDMA. *Sci Transl Med,* 11.

HITZ, C., STEUBER-BUCHBERGER, P., DELIC, S., WURST, W. & KUHN, R. 2009. Generation of shRNA transgenic mice. *Methods Mol Biol,* 530**,** 101-29.

HITZ, C., WURST, W. & KUHN, R. 2007. Conditional brain-specific knockdown of MAPK using Cre/loxP regulated RNA interference. *Nucleic Acids Res,* 35**,** e90.

HUANG, C.-C., LIANG, Y.-C., LEE, C.-C., WU, M.-Y. & HSU, K.-S. 2009. Repeated Cocaine Administration Decreases 5-HT2A Receptor-Mediated Serotonergic Enhancement of Synaptic Activity in Rat Medial Prefrontal Cortex. *Neuropsychopharmacology,* 34**,** 1979-1992.

INSERRA, A., DE GREGORIO, D. & GOBBI, G. 2021. Psychedelics in Psychiatry: Neuroplastic, Immunomodulatory, and Neurotransmitter Mechanisms. *Pharmacol Rev,* 73**,** 202-277.

ISOMURA, Y., SIROTA, A., OZEN, S., MONTGOMERY, S., MIZUSEKI, K., HENZE, D. A. & BUZSAKI, G. 2006. Integration and segregation of activity in entorhinal-hippocampal subregions by neocortical slow oscillations. *Neuron,* 52**,** 871-82.

JUN, J. J., STEINMETZ, N. A., SIEGLE, J. H., DENMAN, D. J., BAUZA, M., BARBARITS, B., LEE, A. K., ANASTASSIOU, C. A., ANDREI, A., AYDIN, Ç., BARBIC, M., BLANCHE, T. J., BONIN, V., COUTO, J., DUTTA, B., GRATIY, S. L., GUTNISKY, D. A., HÄUSSER,

M., KARSH, B., LEDOCHOWITSCH, P., LOPEZ, C. M., MITELUT, C., MUSA, S., OKUN,
M., PACHITARIU, M., PUTZEYS, J., RICH, P. D., ROSSANT, C., SUN, W.-L.,
SVOBODA, K., CARANDINI, M., HARRIS, K. D., KOCH, C., O'KEEFE, J. & HARRIS, T.
D. 2017. Fully integrated silicon probes for high-density recording of neural activity.
Nature, 551**,** 232-236.

KAUR, S., JUNEK, A., BLACK, M. A. & SEMBA, K. 2008. Effects of ibotenate and 192IgG-
saporin lesions of the nucleus basalis magnocellularis/substantia innominata on
spontaneous sleep and wake states and on recovery sleep after sleep deprivation in
rats. *J Neurosci,* 28**,** 491-504.

KAWAGUCHI, Y. & KUBOTA, Y. 1998. Neurochemical features and synaptic connections of
large physiologically-identified GABAergic cells in the rat frontal cortex. *Neuroscience,*
85**,** 677-701.

KOMETER, M., POKORNY, T., SEIFRITZ, E. & VOLLEINWEIDER, F. X. 2015. Psilocybin-
induced spiritual experiences and insightfulness are associated with synchronization
of neuronal oscillations. *Psychopharmacology (Berl),* 232**,** 3663-76.

KRISHNAMURTHY, P., SILBERBERG, G. & LANSNER, A. 2012. A Cortical Attractor Network
with Martinotti Cells Driven by Facilitating Synapses. *PLOS ONE,* 7**,** e30752.

KUYPERS, K. P. C., DE LA TORRE, R., FARRE, M., PIZARRO, N., XICOTA, L. & RAMAEKERS,
J. G. 2018. MDMA-induced indifference to negative sounds is mediated by the 5-HT2A
receptor. *Psychopharmacology (Berl),* 235**,** 481-490.

KVITSIANI, D., RANADE, S., HANGYA, B., TANIGUCHI, H., HUANG, J. Z. & KEPECS, A. 2013.
Distinct behavioural and network correlates of two interneuron types in prefrontal
cortex. *Nature,* 498**,** 363-6.

LANSBERGEN, M. M., DUMONT, G. J. H., VAN GERVEN, J. M. A., BUITELAAR, J. K. &
VERKES, R.-J. 2011. Acute effects of MDMA (3,4-methylenedioxymethamphetamine)
on EEG oscillations: alone and in combination with ethanol or THC (delta-9-
tetrahydrocannabinol). *Psychopharmacology,* 213**,** 745-756.

LEE, S. H. & DAN, Y. 2012. Neuromodulation of brain states. *Neuron,* 76**,** 209-22.

LEMIEUX, M., CHAUVETTE, S. & TIMOFEEV, I. 2015. Neocortical inhibitory activities and
long-range afferents contribute to the synchronous onset of silent states of the
neocortical slow oscillation. *J Neurophysiol,* 113**,** 768-79.

LIECHTI, M. E., SAUR, M. R., GAMMA, A., HELL, D. & VOLLENWEIDER, F. X. 2000.
Psychological and Physiological Effects of MDMA ("Ecstasy") after Pretreatment with
the 5-HT2 Antagonist Ketanserin in Healthy Humans. *Neuropsychopharmacology,* 23**,**
396-404.

LIN, L. C. & SIBILLE, E. 2015. Somatostatin, neuronal vulnerability and behavioral
emotionality. *Mol Psychiatry,* 20**,** 377-87.

LOTTEM, E., LÖRINCZ, M. L. & MAINEN, Z. F. 2016. Optogenetic Activation of Dorsal Raphe
Serotonin Neurons Rapidly Inhibits Spontaneous But Not Odor-Evoked Activity in
Olfactory Cortex. *J Neurosci,* 36**,** 7-18.

MAREK, G. J. & AGHAJANIAN, G. K. 1994. Excitation of interneurons in piriform cortex by 5-hydroxytryptamine: blockade by MDL 100,907, a highly selective 5-HT2A receptor antagonist. *Eur J Pharmacol,* 259**,** 137-41.

MAREK, G. J., CARPENTER, L. L., MCDOUGLE, C. J. & PRICE, L. H. 2003. Synergistic action of 5-HT2A antagonists and selective serotonin reuptake inhibitors in neuropsychiatric disorders. *Neuropsychopharmacology,* 28**,** 402-12.

MARTIN, D. A. & NICHOLS, C. D. 2016. Psychedelics Recruit Multiple Cellular Types and Produce Complex Transcriptional Responses Within the Brain. *EBioMedicine,* 11**,** 262-277.

MAYNE, E. W., CRAIG, M. T., MCBAIN, C. J. & PAULSEN, O. 2013. Dopamine suppresses persistent network activity via D(1) -like dopamine receptors in rat medial entorhinal cortex. *Eur J Neurosci,* 37**,** 1242-7.

MCGINTY, D. J. & HARPER, R. M. 1976. Dorsal raphe neurons: depression of firing during sleep in cats. *Brain Research,* 101**,** 569-575.

MELAMED, O., BARAK, O., SILBERBERG, G., MARKRAM, H. & TSODYKS, M. 2008. Slow oscillations in neural networks with facilitating synapses. *J Comput Neurosci,* 25**,** 308-16.

MELTZER, H. Y. 1999. The role of serotonin in antipsychotic drug action. *Neuropsychopharmacology,* 21**,** 106s-115s.

MIAO, C., CAO, Q., MOSER, M. B. & MOSER, E. I. 2017. Parvalbumin and Somatostatin Interneurons Control Different Space-Coding Networks in the Medial Entorhinal Cortex. *Cell,* 171**,** 507-521.e17.

MICHAIEL, A. M., PARKER, P. R. L. & NIELL, C. M. 2019. A Hallucinogenic Serotonin-2A Receptor Agonist Reduces Visual Response Gain and Alters Temporal Dynamics in Mouse V1. *Cell Rep,* 26**,** 3475-3483.e4.

MITHOEFER, M. C., FEDUCCIA, A. A., JEROME, L., MITHOEFER, A., WAGNER, M., WALSH, Z., HAMILTON, S., YAZAR-KLOSINSKI, B., EMERSON, A. & DOBLIN, R. 2019. MDMA-assisted psychotherapy for treatment of PTSD: study design and rationale for phase 3 trials based on pooled analysis of six phase 2 randomized controlled trials. *Psychopharmacology (Berl),* 236**,** 2735-2745.

MURAYAMA, M., PÉREZ-GARCI, E., NEVIAN, T., BOCK, T., SENN, W. & LARKUM, M. E. 2009. Dendritic encoding of sensory stimuli controlled by deep cortical interneurons. *Nature,* 457**,** 1137-1141.

MUTHUKUMARASWAMY, S. D., CARHART-HARRIS, R. L., MORAN, R. J., BROOKES, M. J., WILLIAMS, T. M., ERRTIZOE, D., SESSA, B., PAPADOPOULOS, A., BOLSTRIDGE, M., SINGH, K. D., FEILDING, A., FRISTON, K. J. & NUTT, D. J. 2013. Broadband cortical desynchronization underlies the human psychedelic state. *J Neurosci,* 33**,** 15171-83.

MUZERELLE, A., SCOTTO-LOMASSESE, S., BERNARD, J. F., SOIZA-REILLY, M. & GASPAR, P. 2016. Conditional anterograde tracing reveals distinct targeting of individual serotonin cell groups (B5–B9) to the forebrain and brainstem. *Brain Structure and Function,* 221**,** 535-561.

NAMIKI, S., NORIMOTO, H., KOBAYASHI, C., NAKATANI, K., MATSUKI, N. & IKEGAYA, Y. 2013. Layer III neurons control synchronized waves in the immature cerebral cortex. *J Neurosci,* 33**,** 987-1001.

NESKE, G., PATRICK, S. & CONNORS, B. 2015a. Contributions of Diverse Excitatory and Inhibitory Neurons to Recurrent Network Activity in Cerebral Cortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience,* 35**,** 1089-105.

NESKE, G. T. 2015. The Slow Oscillation in Cortical and Thalamic Networks: Mechanisms and Functions. *Front Neural Circuits,* 9**,** 88.

NESKE, G. T., PATRICK, S. L. & CONNORS, B. W. 2015b. Contributions of diverse excitatory and inhibitory neurons to recurrent network activity in cerebral cortex. *J Neurosci,* 35**,** 1089-105.

NICHOLS, D. E. 2016. Psychedelics. *Pharmacol Rev,* 68**,** 264-355.

NICHOLS, D. E. & NICHOLS, C. D. 2008. Serotonin Receptors. *Chemical Reviews,* 108**,** 1614-1641.

NIETHARD, N., HASEGAWA, M., ITOKAZU, T., OYANEDEL, C. N., BORN, J. & SATO, T. R. 2016. Sleep-Stage-Specific Regulation of Cortical Excitation and Inhibition. *Curr Biol,* 26**,** 2739-2749.

NIETHARD, N., NGO, H. V., EHRLICH, I. & BORN, J. 2018. Cortical circuit activity underlying sleep slow oscillations and spindles. *Proc Natl Acad Sci U S A,* 115**,** E9220-e9229.

OBERMAYER, J., HEISTEK, T. S., KERKHOFS, A., GORIOUNOVA, N. A., KROON, T., BAAYEN, J. C., IDEMA, S., TESTA-SILVA, G., COUEY, J. J. & MANSVELDER, H. D. 2018. Lateral inhibition by Martinotti interneurons is facilitated by cholinergic inputs in human and mouse neocortex. *Nat Commun,* 9**,** 4101.

OHMURA, Y., IZUMI, T., YAMAGUCHI, T., TSUTSUI-KIMURA, I., YOSHIDA, T. & YOSHIOKA, M. 2010. The serotonergic projection from the median raphe nucleus to the ventral hippocampus is involved in the retrieval of fear memory through the corticotropin-releasing factor type 2 receptor. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology,* 35**,** 1271-1278.

OIKONOMOU, G., ALTERMATT, M., ZHANG, R. W., COUGHLIN, G. M., MONTZ, C., GRADINARU, V. & PROBER, D. A. 2019. The Serotonergic Raphe Promote Sleep in Zebrafish and Mice. *Neuron,* 103**,** 686-701.e8.

PANTAZOPOULOS, H., WISEMAN, J. T., MARKOTA, M., EHRENFELD, L. & BERRETTA, S. 2017. Decreased Numbers of Somatostatin-Expressing Neurons in the Amygdala of Subjects With Bipolar Disorder or Schizophrenia: Relationship to Circadian Rhythms. *Biol Psychiatry,* 81**,** 536-547.

PEROUTKA, S. J. & SNYDER, S. H. 1979. Multiple serotonin receptors: differential binding of [3H]5-hydroxytryptamine, [3H]lysergic acid diethylamide and [3H]spiroperidol. *Molecular pharmacology,* 16**,** 687-699.

PETZOLD, G. C., HAGIWARA, A. & MURTHY, V. N. 2009. Serotonergic modulation of odor input to the mammalian olfactory bulb. *Nature Neuroscience,* 12**,** 784-791.

PRELLER, K. H., BURT, J. B., JI, J. L., SCHLEIFER, C. H., ADKINSON, B. D., STAMPFLI, P., SEIFRITZ, E., REPOVS, G., KRYSTAL, J. H., MURRAY, J. D., VOLLENWEIDER, F. X. & ANTICEVIC, A. 2018. Changes in global and thalamic brain connectivity in LSD-induced altered states of consciousness are attributable to the 5-HT2A receptor. *Elife,* 7.

PUIG, M. V., WATAKABE, A., USHIMARU, M., YAMAMORI, T. & KAWAGUCHI, Y. 2010. Serotonin modulates fast-spiking interneuron and synchronous activity in the rat prefrontal cortex through 5-HT1A and 5-HT2A receptors. *J Neurosci,* 30, 2211-22.

RANADE, S. P. & MAINEN, Z. F. 2009. Transient Firing of Dorsal Raphe Neurons Encodes Diverse and Specific Sensory, Motor, and Reward Events. *Journal of Neurophysiology,* 102, 3026-3037.

RANGANATHAN, G. N., APOSTOLIDES, P. F., HARNETT, M. T., XU, N. L., DRUCKMANN, S. & MAGEE, J. C. 2018. Active dendritic integration and mixed neocortical network representations during an adaptive sensing behavior. *Nat Neurosci,* 21, 1583-1590.

RENART, A., DE LA ROCHA, J., BARTHO, P., HOLLENDER, L., PARGA, N., REYES, A. & HARRIS, K. D. 2010. The asynchronous state in cortical circuits. *Science,* 327, 587-90.

ROTHMAN, R. B. & BAUMANN, M. H. 2002. Therapeutic and adverse actions of serotonin transporter substrates. *Pharmacol Ther,* 95, 73-88.

ROUX, L., STARK, E., SJULSON, L. & BUZSAKI, G. 2014. In vivo optogenetic identification and manipulation of GABAergic interneuron subtypes. *Curr Opin Neurobiol,* 26, 88-95.

RUDY, B., FISHELL, G., LEE, S. & HJERLING-LEFFLER, J. 2011. Three groups of interneurons account for nearly 100% of neocortical GABAergic neurons. *Dev Neurobiol,* 71, 45-61.

SAKAI, K. & CROCHET, S. 2001. Differentiation of presumed serotonergic dorsal raphe neurons in relation to behavior and wake–sleep states. *Neuroscience,* 104, 1141-1155.

SANCHEZ-VIVES, M. V., MASSIMINI, M. & MATTIA, M. 2017. Shaping the Default Activity Pattern of the Cortical Network. *Neuron,* 94, 993-1001.

SCHMITZ, D., GLOVELI, T., EMPSON, R. M., DRAGUHN, A. & HEINEMANN, U. 1998. Serotonin reduces synaptic excitation in the superficial medial entorhinal cortex of the rat via a presynaptic mechanism. *J Physiol,* 508 (Pt 1), 119-29.

SCHNUTGEN, F., DOERFLINGER, N., CALLEJA, C., WENDLING, O., CHAMBON, P. & GHYSELINCK, N. B. 2003. A directional strategy for monitoring Cre-mediated recombination at the cellular level in the mouse. *Nat Biotechnol,* 21, 562-5.

SCOTT, M. M., WYLIE, C. J., LERCH, J. K., MURPHY, R., LOBUR, K., HERLITZE, S., JIANG, W., CONLON, R. A., STROWBRIDGE, B. W. & DENERIS, E. S. 2005. A genetic approach to access serotonin neurons for in vivo and in vitro studies. *Proceedings of the National Academy of Sciences of the United States of America,* 102, 16472-16477.

SEILLIER, L., LORENZ, C., KAWAGUCHI, K., OTT, T., NIEDER, A., POURRIAHI, P. & NIENBORG, H. 2017. Serotonin Decreases the Gain of Visual Responses in Awake Macaque V1. *J Neurosci,* 37, 11390-11405.

SENGUPTA, A., BOCCHIO, M., BANNERMAN, D. M., SHARP, T. & CAPOGNA, M. 2017. Control of Amygdala Circuits by 5-HT Neurons via 5-HT and Glutamate Cotransmission. *The Journal of neuroscience : the official journal of the Society for Neuroscience,* 37**,** 1785-1796.

SENZAI, Y., FERNANDEZ-RUIZ, A. & BUZSAKI, G. 2019. Layer-Specific Physiological Features and Interlaminar Interactions in the Primary Visual Cortex of the Mouse. *Neuron,* 101**,** 500-513 e5.

SHAMASH, P., CARANDINI, M., HARRIS, K. & STEINMETZ, N. 2018. A tool for analyzing electrode tracks from slice histology. *bioRxiv***,** 447995.

SHELDON, P. W. & AGHAJANIAN, G. K. 1990. Serotonin (5-HT) induces IPSPs in pyramidal layer cells of rat piriform cortex: evidence for the involvement of a 5-HT2-activated interneuron. *Brain Res,* 506**,** 62-9.

SIPPY, T. & YUSTE, R. 2013. Decorrelating action of inhibition in neocortical networks. *J Neurosci,* 33**,** 9813-30.

SMITH, S. L., SMITH, I. T., BRANCO, T. & HÄUSSER, M. 2013. Dendritic spikes enhance stimulus selectivity in cortical neurons in vivo. *Nature,* 503**,** 115-120.

STERIADE, M., NUNEZ, A. & AMZICA, F. 1993. A novel slow (< 1 Hz) oscillation of neocortical neurons in vivo: depolarizing and hyperpolarizing components. *J Neurosci,* 13**,** 3252-65.

STRINGER, C., PACHITARIU, M., STEINMETZ, N. A., OKUN, M., BARTHO, P., HARRIS, K. D., SAHANI, M. & LESICA, N. A. 2016. Inhibitory control of correlated intrinsic variability in cortical networks. *Elife,* 5.

STURGILL, J. F. & ISAACSON, J. S. 2015. Somatostatin cells regulate sensory response fidelity via subtractive inhibition in olfactory cortex. *Nat Neurosci,* 18**,** 531-5.

TAHVILDARI, B., WOLFEL, M., DUQUE, A. & MCCORMICK, D. A. 2012. Selective functional interactions between excitatory and inhibitory cortical neurons and differential contribution to persistent activity of the slow oscillation. *J Neurosci,* 32**,** 12165-79.

TAKAHASHI, N., OERTNER, T. G., HEGEMANN, P. & LARKUM, M. E. 2016. Active cortical dendrites modulate perception. *Science,* 354**,** 1587-1590.

TANG, Z.-Q. & TRUSSELL, L. O. 2017. Serotonergic Modulation of Sensory Representation in a Central Multisensory Circuit Is Pathway Specific. *Cell reports,* 20**,** 1844-1854.

TANIGUCHI, H., HE, M., WU, P., KIM, S., PAIK, R., SUGINO, K., KVITSIANI, D., FU, Y., LU, J., LIN, Y., MIYOSHI, G., SHIMA, Y., FISHELL, G., NELSON, S. B. & HUANG, Z. J. 2011. A resource of Cre driver lines for genetic targeting of GABAergic neurons in cerebral cortex. *Neuron,* 71**,** 995-1013.

TRAINITO, C., VON NICOLAI, C., MILLER, E. K. & SIEGEL, M. 2019. Extracellular Spike Waveform Dissociates Four Functionally Distinct Cell Classes in Primate Cortex. *Curr Biol*.

TREMBLAY, R., LEE, S. & RUDY, B. 2016. GABAergic Interneurons in the Neocortex: From Cellular Properties to Circuits. *Neuron,* 91**,** 260-92.

TUKKER, J. J., BEED, P., SCHMITZ, D., LARKUM, M. E. & SACHDEV, R. N. S. 2020. Up and Down States and Memory Consolidation Across Somatosensory, Entorhinal, and Hippocampal Cortices. *Frontiers in Systems Neuroscience,* 14.

TURRIGIANO, G. 2011. Too many cooks? Intrinsic and synaptic homeostatic mechanisms in cortical circuit refinement. *Annu Rev Neurosci,* 34**,** 89-103.

UDO DE HAES, J. I., HARADA, N., ELSINGA, P. H., MAGUIRE, R. P. & TSUKADA, H. 2006. Effect of fenfluramine-induced increases in serotonin release on [18F]MPPF binding: a continuous infusion PET study in conscious monkeys. *Synapse,* 59**,** 18-26.

UNGER, E. K., KELLER, J. P., ALTERMATT, M., LIANG, R., MATSUI, A., DONG, C., HON, O. J., YAO, Z., SUN, J., BANALA, S., FLANIGAN, M. E., JAFFE, D. A., HARTANTO, S., CARLEN, J., MIZUNO, G. O., BORDEN, P. M., SHIVANGE, A. V., CAMERON, L. P., SINNING, S., UNDERHILL, S. M., OLSON, D. E., AMARA, S. G., TEMPLE LANG, D., RUDNICK, G., MARVIN, J. S., LAVIS, L. D., LESTER, H. A., ALVAREZ, V. A., FISHER, A. J., PRESCHER, J. A., KASH, T. L., YAROV-YAROVOY, V., GRADINARU, V., LOOGER, L. L. & TIAN, L. 2020. Directed Evolution of a Selective and Sensitive Serotonin Sensor via Machine Learning. *Cell,* 183**,** 1986-2002.e26.

URBAN-CIECKO, J., FANSELOW, E. E. & BARTH, A. L. 2015. Neocortical somatostatin neurons reversibly silence excitatory transmission via GABAb receptors. *Curr Biol,* 25**,** 722-31.

URBAN-CIECKO, J., JOUHANNEAU, J. S., MYAL, S. E., POULET, J. F. A. & BARTH, A. L. 2018. Precisely Timed Nicotinic Activation Drives SST Inhibition in Neocortical Circuits. *Neuron,* 97**,** 611-625.e5.

VAN STRIEN, N. M., CAPPAERT, N. L. & WITTER, M. P. 2009. The anatomy of memory: an interactive overview of the parahippocampal-hippocampal network. *Nat Rev Neurosci,* 10**,** 272-82.

VANDERWOLF, C. H. & BAKER, G. B. 1986. Evidence that serotonin mediates non-cholinergic neocortical low voltage fast activity, non-cholinergic hippocampal rhythmical slow activity and contributes to intelligent behavior. *Brain Res,* 374**,** 342-56.

VERTES, R. P. & LINLEY, S. B. 2007. Comparison of projections of the dorsal and median raphe nuclei, with some functional considerations. *International Congress Series,* 1304**,** 98-120.

WANG, H., HU, L., LIU, C., SU, Z., WANG, L., PAN, G., GUO, Y. & HE, J. 2016. 5-HT2 receptors mediate functional modulation of GABAa receptors and inhibitory synaptic transmissions in human iPS-derived neurons. *Scientific Reports,* 6**,** 20033.

WEBER, E. T. & ANDRADE, R. 2010. Htr2a Gene and 5-HT(2A) Receptor Expression in the Cerebral Cortex Studied Using Genetically Modified Mice. *Front Neurosci,* 4.

WILSON, N. R., RUNYAN, C. A., WANG, F. L. & SUR, M. 2012. Division and subtraction by distinct cortical inhibitory networks in vivo. *Nature,* 488**,** 343-348.

1207 WOLANSKY, T., CLEMENT, E. A., PETERS, S. R., PALCZAK, M. A. & DICKSON, C. T. 2006. Hippocampal Slow Oscillation: A Novel EEG State and Its Coordination with Ongoing Neocortical Activity. *The Journal of Neuroscience,* 26**,** 6213.

1210 WOOD, J., KIM, Y. & MOGHADDAM, B. 2012. Disruption of prefrontal cortex large scale neuronal activity by different classes of psychotomimetic drugs. *J Neurosci,* 32**,** 3022-31.

1213 WYSKIEL, D. R. & ANDRADE, R. 2016. Serotonin excites hippocampal CA1 GABAergic interneurons at the stratum radiatum-stratum lacunosum moleculare border. *Hippocampus,* 26**,** 1107-14.

1216 ZHOU, F.-M. & HABLITZ, J. J. 1999. Activation of Serotonin Receptors Modulates Synaptic Transmission in Rat Cerebral Cortex. *Journal of Neurophysiology,* 82**,** 2989-2999.

1218 ZUCCA, S., D'URSO, G., PASQUALE, V., VECCHIA, D., PICA, G., BOVETTI, S., MORETTI, C., VARANI, S., MOLANO-MAZÓN, M., CHIAPPALONE, M., PANZERI, S. & FELLIN, T. 2017. An inhibitory gate for state transition in cortex. *eLife,* 6**,** e26177.