Multi-contrast Anatomical Subcortical Structures Parcellation

Pierre-Louis Bazin[1,2], Anneke Alkemade[1], Martijn J. Mulder[1,3], Amanda G. Henry[4], Birte U. Forstmann[1]

[1] Integrative Model-based Cognitive Neuroscience research unit, University of Amsterdam, Amsterdam, The Netherlands; [2] Max-Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany; [3] Psychology Department, Utrecht University, Utrecht, The Netherlands; [4] Faculty of Archaeology, Leiden University, Leiden, The Netherlands.

Abstract

The human subcortex is comprised of more than 450 individual nuclei which lie deep in the brain. Due to their small size and close proximity, up until now only 7% have been depicted in standard MRI atlases. Thus, the human subcortex can largely be considered as terra incognita. Here we present a new open source parcellation algorithm to map terra incognita automatically. The new algorithm has been tested on 17 prominent subcortical structures based on a large quantitative MRI dataset at 7 Tesla. The algorithm is carefully validated against expert human raters, available in Open Source, and can easily be extended to other subcortical structures and applied to any quantitative MRI dataset. In sum, we hope this novel parcellation algorithm will facilitate functional and structural neuroimaging research into small subcortical nuclei and help to chart terra incognita.

Introduction

Subcortical brain structures are often neglected in neuroimaging studies due to their small size, limited inter-regional contrast, and weak signal-to-noise ratio in functional imaging (Forstmann et al., 2017a; Johansen-Berg, 2013). Yet, these small and diverse structures are prominent nodes in functional networks (Marquand et al., 2017; Ji et al., 2019), and they undergo pathological alterations already at early stages of neurodegenerative diseases (Andersen et al., 2014; Koshiyama et al., 2018). Deep brain stimulation surgery, originally performed to reduce motor symptoms in essential tremors, is now a promising therapeutic option in later stages of Parkinson's disease and movement disorders, as well as refractory psychiatric illnesses in obsessive-compulsive disorder, anorexia, or depression (Forstmann et al., 2017b, Mosley et al., 2018). Evolutionary genetics even uncovered that in modern humans, Neanderthal-inherited alleles were preferentially down-regulated in subcortical and cerebellar regions compared to other brain regions (McCoy et al., 2017), suggesting these structures to be essential in making us specifically human.

Despite their importance, these areas are particularly difficult to image. Furthermore, the size, shape and location of these brain regions changes with development and aging (Fjell et al., 2013; Keuken et al. 2013; Yeatman et al., 2014; Herting et al., 2018). Experience-based plasticity continuously remodels myelin (Tardif et al., 2016; Hill et al., 2018; Turner, 2019), iron and other magnetic substances accumulate with age or pathology (Andersen et al., 2014; Zhang et al., 2018), both bringing changes in the magnetic resonance imaging (MRI) appearance of subcortical regions with diverse tissue characteristics (Draganski et al., 2011; Keuken et al., 2017).

Thus, mapping the structure and function of the subcortex is a major endeavor as well as a major challenge for human neuroscience. Extensive work available from animal brain models unfortunately does not translate in a straightforward way to human subcortical anatomy nor does it shed much light on its involvement in human cognition (Steiner and Tseng, 2017). Besides serious difficulties in obtaining adequate measures of subcortical neural activity in functional MRI (de Hollander et al., 2017; Miletic et al., 2020), atlases and techniques for labeling accurately and

reliably individual subcortical structures have also been scarce (Frazier et al., 2005; Chakravarty et al., 2006; Ahsan et al., 2007; Yelnik et al., 2007; Qiu et al., 2010; Patenaude et al., 2011), typically labelling the thalamus, striatum (or its subdivision into caudate and putamen), and globus pallidus (internal and external segments combined), sometimes the amygdala. However, recent advances in anatomical MRI, combining multiple contrasts and/or quantitative MRI mapping and utilizing the higher resolution achievable with 7 Tesla (7T) and above have started to reduce the gap, each mapping a few additional structures or sub-structures, primarily the iron-rich substantia nigra, red nucleus and sub-thalamic nucleus (Keuken et al., 2014; Xiao et al., 2015; Visser et al., 2016a; Visser et al., 2016b; Wang et al., 2016; Makowski et al., 2018; Ewert et al., 2018; Iglesias et al., 2018; Pauli et al., 2018; Sitek et al., 2019). While these efforts generated valuable atlases, they do not yet enable to identify many subcortical structures in individual subjects. Manual delineation, on the other hand, requires extensive manual labor from highly trained experts which cannot be easily applied to large cohorts or clinical settings.

Here, we propose a new automated parcellation technique to identify and label 17 individual subcortical structures of varying size and composition in individual subjects, based on a large quantitative 7T MRI database (Alkemade et al., under review). The algorithm follows a Bayesian multi-object approach similar in essence to previous efforts (Fischl et al., 2002; Iglesias et al., 2013; Visser et al., 2016a; Garzon et al., 2018), combining shape priors, intensity distribution models, spatial relationships and global constraints. The main innovation of our approach is to explicitly estimate interfaces between subcortical structures based on a joint model derived from signed distance functions. Modeling interfaces in addition to the structure itself provides a rich basis to encode relationships and anatomical knowledge in shape and intensity priors. A voxel-wise Markovian diffusion regularizes the combined priors for each defined interface, lowering the imaging noise. Finally, the voxel-wise posteriors for the different structures and interfaces are further combined into global anatomical parcels by topology correction and region growing taking into account volumetric priors, which regularizes parcellation results further in smaller nuclei with low or heterogeneous contrast. To validate the results from this new method, in a thorough comparison with expert manual labeling, we show that the proposed method provides results very close from manual raters in many structures and exhibit reasonable bias across the adult lifespan. The method can easily be extended to new structures, can be applied to any quantitative MRI dataset and is available in Open Source as part of Nighres (Huntenburg et al., 2018), a neuroimage analysis package aimed at high resolution neuroimaging.

Results

The parcellation method presented here, Multi-contrast Anatomical Subcortical Structure Parcellation (MASSP), has been trained to parcellate the following 17 structures: striatum (Str), thalamus (Tha), lateral, 3rd and 4th ventricles (LV, 3V, 4V), amygdala (Amg), globus pallidus internal segment (GPi) and external segment (GPe), SN, STN, red nucleus (RN), ventral tegmental area (VTA), fornix (fx), internal capsule (ic), periaqueductal gray (PAG), pedunculopontine nucleus (PPN), and claustrum (Cl), see Fig. 1. These structures include the most commonly defined subcortical regions (Str, Tha, Amg, LV), the main iron-rich nuclei (GPi, GPe, RN, SN, STN), as well as smaller, less studied areas (VTA, PAG, PPN, Cl), white matter structures (ic, fx) and the central ventricles (3V, 4V).

Figure 1: The 17 subcortical structures currently included in the parcellation algorithm in axial (A), sagittal (B) and coronal (C) views.

MASSP uses a data set of ten expert delineations as a basis for its modeling. In a leave-one-out validation study comparing performance with the manual raters, MASSP performed above 95% of the level of quality of the raters for Str, Tha, 4V, GPe, SN, RN, VTA, ic in terms of Dice overlap, the most stringent of the quality measures (see Fig. 2 and 3 and Table 1). Several of the smaller structures have lower overlap ratios likely due to their smaller size (GPi, STN, PAG, PPN). Structures with an elongated shape (fx, Cl) remain challenging, due to the fact that small differences in location can substantially reduce overlap (Bazin et al., 2016). Despite these challenges, when comparing the dilated Dice scores, all structures were above 75% of overlap, with most reaching over 90% of the manual raters ability. Note that the Dice coefficient is very sensitive to size, as smaller structures will have lower overlap ratios for the same number of misclassified voxels. The dilated Dice coefficient is more representative of the variability regardless of size, as the smaller structures can reach high levels of overlap, both in manual and automated parcellations (see Table 1). The average surface distance confirms these results, showing values generally between one and two voxels of distance at a resolution of 0.7 mm, except in the cases of Amg, LV, fx, PPN, and Cl. These structures are generally more variable (LV), elongated (fx, Cl), or have a particularly low contrast with neighboring regions (Amg, PPN).

Figure 2: Leave-one-out validation of the structures parcellated by MASSP, compared to the most expert of the two human raters. Scores for the left and right side are computed separately then combined into box-and-whisker plots.

Figure 3: Inter-rater variability for the human expert raters. Scores for the left and right side are computed separately then combined into box-and-whisker plots.

Table 1: Mean overlap and distance measures for the leave-one-out validation

	Str	STN	SN	RN	GPi	GPe	Tha	LV	3V	4V	Amg	ic	VTA	fx	PAG	PPN	Cl
Dice overlap																	
MASSP	0.893	0.648	0.805	0.870	0.702	0.800	0.867	0.849	0.741	0.869	0.723	0.745	0.570	0.527	0.641	0.496	0.485
manual	0.897	0.800	0.841	0.875	0.762	0.813	0.877	0.907	0.797	0.882	0.866	0.732	0.574	0.823	0.791	0.665	0.727
ratio	0.995	0.811	0.957	0.996	0.925	0.987	0.989	0.936	0.936	0.988	0.836	1.020	0.994	0.641	0.814	0.754	0.664
Dilated overlap																	
MASSP	0.982	0.919	0.977	0.991	0.909	0.956	0.970	0.929	0.890	0.951	0.891	0.915	0.863	0.756	0.897	0.795	0.789
manual	0.987	0.988	0.985	0.995	0.953	0.972	0.970	0.967	0.944	0.961	0.978	0.924	0.818	0.957	0.960	0.910	0.914
ratio	0.995	0.930	0.992	0.995	0.955	0.984	1.000	0.961	0.946	0.991	0.911	0.992	1.059	0.790	0.935	0.879	0.863
Boundary distance																	
MASSP	0.750	0.911	0.676	0.491	1.140	0.863	1.058	2.690	0.994	0.817	1.476	1.275	1.074	2.950	0.955	1.484	1.685
manual	0.723	0.508	0.571	0.482	0.902	0.804	0.971	0.615	0.637	0.671	0.779	1.045	1.204	0.703	0.555	0.801	0.670
ratio	0.971	0.590	0.852	0.996	0.861	0.943	0.916	0.277	0.662	1.020	0.553	0.834	1.161	0.287	0.610	0.619	0.465

A common concern of brain parcellation methods is the risk of biases, as they are typically built from a small number of manual delineations. Our data set is part of a large scale study of the subcortex, for which we obtained manual delineations of the STN, SN, RN, GPe, and GPi on 105 subjects over the adult lifespan (18-80 year old, see Alkemade et al., under review for details). We

used the priors from our 10 subjects to delineate all structures, and performed the same analysis as above on the entire cohort, separated into age groups (18-40, 41-60, 61-80). Results are given in Table 2 and Fig. 4.

Figure 4: MASSP parcellation scores over the adult lifespan. Circles show individual data points, linear regression is indicated by a straight line, and 95% confidence interval is given as the shaded area. Regression coefficients are indicated when significant (p-value<0.01).

Table 2: Mean overlap and distance measures for each age group

	STN			SN			RN			GPi			GPe		
Dice overlap	18-40	41-60	61-80	18-40	41-60	61-80	18-40	41-60	61-80	18-40	41-60	61-80	18-40	41-60	61-80
MASSP	0.647	0.572	0.516	0.803	0.794	0.735	0.873	0.858	0.829	0.678	0.622	0.606	0.788	0.736	0.711
manual	0.772	0.704	0.714	0.849	0.846	0.803	0.910	0.900	0.902	0.782	0.785	0.767	0.817	0.795	0.800
ratio	0.842	0.793	0.725	0.947	0.939	0.915	0.961	0.964	0.919	0.886	0.806	0.809	0.966	0.941	0.892
Dilated overlap	18-40	41-60	61-80	18-40	41-60	61-80	18-40	41-60	61-80	18-40	41-60	61-80	18-40	41-60	61-80
MASSP	0.922	0.863	0.806	0.971	0.964	0.928	0.992	0.984	0.970	0.870	0.824	0.816	0.944	0.911	0.898
manual	0.968	0.908	0.946	0.985	0.986	0.963	0.997	0.986	0.998	0.947	0.950	0.941	0.974	0.962	0.962
ratio	0.953	0.918	0.853	0.986	0.977	0.963	0.995	1.004	0.973	0.924	0.872	0.871	0.970	0.948	0.935
Boundary distance	18-40	41-60	61-80	18-40	41-60	61-80	18-40	41-60	61-80	18-40	41-60	61-80	18-40	41-60	61-80
MASSP	0.918	1.287	2.136	0.712	0.742	0.942	0.484	0.558	0.646	1.936	1.651	1.693	0.940	1.135	1.214
manual	0.608	1.153	0.704	0.567	0.584	0.706	0.350	0.528	0.377	0.866	0.902	0.875	0.777	0.824	0.824
ratio	0.675	0.719	0.574	0.837	0.839	0.811	0.737	0.866	0.627	0.746	0.602	0.554	0.857	0.760	0.721

Examining this age bias, we can see a decrease in performance in both the automated and manual delineations of STN, SN, and GPe, which can be explained by the changes in MRI contrast associated with changing myelination, iron deposition, and brain atrophy (Draganski et al., 2011; Daugherty et al. 2013; Fjell et al., 2013; Keuken et al., 2017). For STN and GPi, the automated method showed an additional decrease in performance which indicate that the algorithm priors are more representative of the younger subjects. Variability also increased with age, likely reflecting the increasing heterogeneity of the older compared to younger subjects. Such age biases are structure specific, and not solely related to contrast changes: very little bias was observed for SN and RN, while their contrast changes substantially due to iron deposition (see Fig. 6 below). This bias does affect estimated structure volumes, but measured quantities inside each structure such as quantitative MRI metrics remain largely unaffected, even when age biases are present for the parcellations (see Fig. 5).

Figure 5: Regression of volume (log scale), R1, R2*, and QSM MRI parameters estimated using manual delineations versus MASSP automated parcellations.Circles show individual data points,

linear regression is indicated by a straight line, and 95% confidence interval is given as the shaded area. Regression coefficients are indicated when significant (p-value<0.01).

Finally, we used the delineations on the entire cohort to compute structure volumes, R1, R2* and QSM values for different age groups, extending our previous work based on manual delineations on a different data set (Keuken et al., 2017; Forstmann et al., 2014). Results are given in Table 3, describing average volumes and quantitative MRI parameters for young, middle-aged, and older subjects for the 17 subcortical structures.

Table 3: Mean volume and quantitative MRI values for each age group

	Str	STN	SN	RN	GPi	GPe	Tha	LV	3V	4V	Amg	ic	VTA	fx	PAG	PPN	Cl
Volume (mm³)																	
18-40	10656	118	566	253	567	1366	7112	7524	1895	1391	1315	4335	254	1632	250	193	843
41-60	10572	124	583	256	586	1403	7492	8850	2024	1408	1363	4495	264	1808	255	195	830
61-80	10734	130	584	260	586	1397	7463	9142	2023	1407	1321	4407	272	1910	259	192	829
qR1 (Hz)																	
18-40	0.647	0.949	0.857	0.928	0.868	0.850	0.761	0.332	0.346	0.274	0.546	0.906	0.819	0.714	0.654	0.779	0.650
41-60	0.662	0.968	0.893	0.939	0.879	0.856	0.758	0.278	0.315	0.269	0.559	0.904	0.833	0.671	0.653	0.771	0.664
61-80	0.648	0.952	0.882	0.903	0.860	0.830	0.743	0.273	0.300	0.270	0.552	0.884	0.814	0.638	0.647	0.764	0.669
qR2*(Hz)																	
18-40	43.8	67.1	67.8	63.2	75.2	79.6	38.1	14.7	18.9	9.0	25.5	36.8	39.2	37.4	25.9	32.7	32.6
41-60	50.4	74.1	74.1	77.1	80.2	87.9	40.3	8.4	12.4	11.7	28.1	38.7	42.8	37.4	28.0	33.4	36.9
61-80	51.8	77.0	72.5	73.8	77.8	87.0	40.1	8.5	10.2	12.0	30.1	39.6	52.6	35.7	28.4	34.2	35.4
QSM (ppm)																	
18-40	0.0329	0.0609	0.0738	0.0717	0.1015	0.1150	0.0138	0.0130	0.0100	0.0279	0.0036	-0.0234	0.0241	0.0079	0.0119	0.0135	-0.0122
41-60	0.0400	0.0647	0.0713	0.0829	0.0984	0.1241	0.0134	0.0115	0.0025	0.0234	0.0085	-0.0226	0.0201	0.0079	0.0089	0.0099	-0.0110
61-80	0.0411	0.0705	0.0610	0.0738	0.0925	0.1249	0.0064	0.0089	-0.0034	0.0236	0.0061	-0.0243	0.0177	0.0100	0.0039	0.0096	-0.0091

Discussion

Our goal with the MASSP algorithm was to provide a fully automated method to delineate as many subcortical structures as possible on high resolution structural MRI now available on 7T scanners. We modeled 17 distinct structures, taking into account location, shape, volume, and quantitative MRI contrasts to provide individual subject parcellations. Based on our results, we can be confident that the automated parcellation technique performs comparably to human experts, providing delineations within one or two voxels of the structure boundaries (dilated Dice overlap over 75% for all structures, including in aging groups). Results were nearly indistinguishable from expert delineations for eight major structures (Str, Tha, 4V, GPe, SN, RN, VTA, ic), and smaller structures retain high levels of overlap, comparable to trained human raters. This parcellation includes the most commonly defined structures (Str, Tha, SN, RN, STN) with overlap scores comparable to those previously reported (Garzon et al., 2018; Visser et al., 2016a; Visser et al., 2016b; Iglesias et al., 2013; Chakravarty et al., 2013; Patenaude et al., 2011). More importantly, it also includes structures seldom or never before considered in MRI atlases and parcellation methods, such as GPe, GPi, VTA, 3V, 4V, ic, fx, PAG, PPN, Cl. The technique handles structures of varying sizes well, as indicated by dilated overlap and boundary distance. Additional structures can be added, if they can be reliably delineated by expert raters on single-subject MRI at achievable resolutions. Some enhancement techniques such as building a multi-subject template (Pauli et al., 2018) or adding a denoising step (Bazin et al., 2019) may be beneficial.

Age biases are present both in expert manual delineations and automated parcellation techniques. Age trajectories in volume and quantitative MR parameters indicate systematic shifts in contrast intensities and an increasing variability with age, thus indicating that age-matched priors or an explicit modeling of age effects may be beneficial to further improve parcellation quality. These results also point to exercising caution when applying automated parcellation methods to study morphometry in diseased populations, where measured differences may include biases. They also

point out that while volume is indeed affected by such biases, MRI-based measures are much more robust. Interestingly, biases also exist in expert delineations: when the size or shape of a structure is refined in neuroanatomical studies, experts may become more or less conservative in their delineations. Automated methods provide a more objective measure in such case, as the source of their bias is explicitly encoded in the atlas prior delineations and computational model.

Another interesting aspect of our validation study is the observation that dilated overlap, i.e., the overlap of structures up to one voxel, provided a measure of accuracy largely independent of size, for automated or manual delineations. Imprecision in the range of one voxel in the boundary is to be expected due to partial voluming which impacts Dice overlap. The dilated overlap measure is a better representative of performance, and indicates that conservative or inclusive versions of the subcortical regions can be obtained by eroding or dilating the estimated boundary by a single voxel. Such masks may be useful when separating functional MRI signals between neighboring nuclei or when locating smaller features inside a structure. Additionally, the Bayesian estimation framework provides voxel-wise probability values, which can also be used to further weight the contribution of each voxel within a region in subsequent analyses.

In summary, our method provides fast and accurate parcellation for subcortical structures of varying size, taking advantage of the high resolution offered by 7T and the specificity of quantitative MRI. The algorithm is based on an explicit model of structures given in a Bayesian framework, and is free of tuning parameters. Given a different set of regions of interest or different populations, new priors can be automatically generated and used as the basis for the algorithm. If more MRI contrasts are available, the method can also be augmented to take them into account. The main requirement for the technique is a set of manual delineations of all the structures of interest in a small group of representative subjects. Performance may further improve with the number of included structures, as the number of distinct interfaces increases, refining in particular the intensity priors. In future works, we plan to include more structures or sub-structures and model the effects of age on the priors. We hope that the method, available in open source, will help neuroscience researchers to include more subcortical regions in their structural and functional imaging studies.

Materials and Methods

Data acquisition
Our parcellation method has been developed for the MP2RAGEME sequence (Caan et al., 2018). Briefly, the MP2RAGEME consists of two interleaved MPRAGEs with different inversions and four echoes in the second inversion. Based on these images, one can estimate quantitative MR parameters of R1, R2* and QSM. In this work, we used the following sequence parameters: inversion times $TI_{1,2}$ = 670 ms, 3675.4 ms; echo times TE1= 3 ms, $TE_{2,1-4}$ = 3, 11.5, 19, 28.5 ms; flip angles $FA_{1,2}$ = 4°, 4°; $TRGRE_{1,2}$ = 6.2 ms, 31 ms; bandwidth = 404.9 MHz; TRMP2RAGE = 6778 ms; SENSE acceleration factor = 2; FOV = 205 x 205 x 164mm; acquired voxel size = 0.70 x 0.7 x 0.7 mm; acquisition matrix was 292 x 290; reconstructed voxel size = 0.64 x 0.64 x 0.7 mm; turbo factor (TFE) = 150 resulting in 176 shots; total acquisition time = 19.53 min.

T1-maps were computed using a look-up table (Marques et al., 2010). T2*-maps were computed by least-squares fitting of the exponential signal decay over the multi-echo images of the second inversion. R1 and R2* maps were obtained as the inverse of T1 and T2*. For QSM, phase maps were pre-processed using iHARPERELLA (integrated phase unwrapping and background phase removal using the Laplacian) of which the QSM images were computed using LSQR (Li et al., 2014). Skull information was removed through creation of a binary mask using FSL's brain extraction tool on the reconstructed uniform T1-weighted image and then applied to the quantitative contrasts (Smith et al., 2002). As all images were acquired as part of a single sequence, no co-registration of the quantitative maps was required (see Fig.6).

Figure 6: MP2RAGEME maps and delineations: quantitative R1 (left), quantitative R2* (middle), QSM (right). Manual delineations for the 17 structures of interest are overlaid on all images.

Anatomical structure delineations

Manual delineations of subcortical structures were performed by two raters trained by an expert anatomist, according to protocols optimized to use the better contrast or combination of contrasts for each structure and to ensure a consistent approach across raters. The following 17 structures were defined on a group of ten subjects (average age 24.4, 8 female): striatum (Str), thalamus (Tha), lateral, 3rd and 4th ventricles (LV, 3V, 4V), amygdala (Amg), globus pallidus internal segment (GPi) and external segment (GPe), SN, STN, red nucleus (RN), ventral tegmental area (VTA), fornix (fx), internal capsule (ic), periaqueductal gray (PAG), pedunculopontine nucleus (PPN), and claustrum (Cl). Separate masks for left and right hemisphere were delineated except for 3V, 4V, and fx. In the following the algorithm treats each side separately, resulting in a total of 31 distinct structures (see Fig. 1).

Anatomical interface priors

In order to inform the algorithm, we built a series of priors derived from the manual delineations. Each subject was first co-registered to a MP2RAGEME anatomical template built from 105 subjects co-aligned with the MNI2009b atlas (Fonov et al., 2011) with the SyN algorithm of ANTs (Avants et al., 2008) using successively rigid, affine, and non-linear transformations, high levels of regularisation as recommended for the subcortex (Ewert et al., 2019) and mutual information as cost function.

The first computed prior is a prior of anatomical interfaces, recording the most likely location of boundaries between the different structures, defined as follows. Given two delineated

structures i, j, let φ_i, φ_j be the signed distance functions to their respective boundary, i.e. $\varphi_i(x)$ is the distance to the boundary of i at x, with a negative sign inside the structure. Then we define the interface $B_{i|j}$ with the distance function $d_{i|j}$:

$$d_{i|j}(x) = \min(\varphi_i(x), \varphi_j(x) - \delta, 0),$$

where δ is a scale parameter for the thickness of the interface. These interfaces functions are not symmetrical, as the intensity inside i next to j is generally different from the intensity inside j next to i. Based on this definition, the prior for a given interface based on N manual delineations is given by:

$$P(x \in B_{i|j}) \sim \frac{1}{\sqrt{2\pi\sigma_{i|j}^2(x)}} \exp -\frac{1}{2}\frac{\mu_{i|j}^2(x)}{\sigma_{i|j}^2(x)},$$

$$\mu_{i|j}(x) = \frac{1}{N}\sum_{n \in N} d_{i|j,n}(x), \ \sigma_{i|j}(x) = \sqrt{\frac{1}{N}\sum_{n \in N}\left(d_{i|j,n}(x) - \mu_{i|j}(x)\right)^2} + \delta.$$

These probability functions are calculated for all possible configurations including $i|i$, which represent the inside of each structure. We thus have a total of N^2 functions, but only a few are non-zero at a given voxel x, and we may keep only the 16 largest values to account for any number of interfaces in 3D (Bazin et al., 2007a). Finally, we need to scale the prior to be globally consistent with the priors below by assuming that the 95[th] percentile of the highest kept $P(x \in B_{i|j})$ values have a probability of 0.95. The resulting interface prior is shown in Fig. 7A.

axial view   coronal view   sagittal view

Figure 7: Anatomical interface (A) and skeleton (B) priors derived from the 10 manually delineated subjects.

Anatomical skeleton priors

Next, we defined priors for the skeleton of each structure, representing their essential shape regardless of exact boundaries (Blum, 1973). As we are mostly interested in the most likely components of the skeleton or medial axis S_i, we follow a simple method to estimate its location:

$$S_i = \left\{ x, |\nabla \varphi_i(x)| < \frac{1}{2} \right\}$$

We define as $s_i(x)$ the signed distance function of this discrete skeleton, and define prior probabilities as above:

$$P(x \in S_i) \sim \frac{1}{\sqrt{2\pi\sigma_i^2(x)}} \exp -\frac{1}{2}\frac{\mu_i^2(x)}{\sigma_i^2(x)},$$

$$\mu_i(x) = \frac{1}{N} \sum_{n \in N} s_{i,n}(x), \sigma_{i|j}(x) = \sqrt{\frac{1}{N} \sum_{n \in N} (s_{i,n}(x) - \mu_i(x))^2} + \delta.$$

The skeletons are defined inside each structure, which implies $P(x \in S_i) \leq P(x \in B_{i|i})$. To respect this relationship, we scale $P(x \in S_i)$ with the same factor as $P(x \in B_{i|i})$ but use $\sqrt{P(x \in S_i)}$ when combining probabilities during the estimation stage. The obtained anatomical skeleton priors are given on Fig. 7B.

Interface intensity priors

While anatomical priors already provide rich information, they are largely independent of the underlying MRI. From the co-aligned quantitative MRI maps and manual delineations, we defined intensity priors for every interface $i|j$, in the form of intensity histograms to ensure a flexible representation of intensity distributions. Given a quantitative contrast $R_n(x)$, we built a histogram $H_{i|j,n}$ for each subject n and interface $i|j$. Histograms have 200 bins covering the entire intensity range within a radius of 10 mm from any of the delineated structures. To obtain an average histogram, we combine each histogram with a weighting function $w_n(x)$ giving the likelihood of the subject's intensity measurement compared to the group:

$$w_{i|j,n}(x) = P(x \in B_{i|j})\frac{1}{\sqrt{2\pi\sigma_R^2}} \exp -\frac{1}{2}\frac{(R_n(x) - \mu_R(x))^2}{\sigma_R(x)^2},$$

where $\mu_R(x)$ is the median of the $R_n(x)$ values at x, and $\sigma_R(x)$ is 1.349 times the inter-quartile range of $R_n(x)$. These are robust estimators of the mean and standard deviation, used here to avoid biases by intensity outliers. To further combine the R1, R2*, and QSM contrasts we take the geometric mean of the histogram probabilities: $H_{i|j}(x) = \prod_R H_{i|j}(R(x))^{1/3}$.

Global volume priors

The last type of priors extracted from manual delineations are volume priors for each of the structure. Here we assume a log-normal distribution for the volumes V_i and simply estimate the mean $\mu_{V,i}$ and standard deviation $\sigma_{V,i}$ of $\log V_{i,n}$ over the subjects.

Voxel-wise posterior probabilities

When parcellating a new subject, we first co-register its R1, R2*, and QSM maps jointly to the template and use the inverse transformation to deform the anatomical priors into subject space. Then we derive voxel-wise posteriors as follows:

$$P(x \in B_{i|j}|R(x)) \sim P(x \in B_{i|j})H_{i|j}(x) \text{ if } i \neq j,$$
$$\text{and}$$
$$P(x \in B_{i|i}|S_i(x), R(x)) \sim \max(P(x \in B_{i|j}), P(x \in S_j)^{1/2})H_{i|j}(x)$$

Once again we should compute all possible combinations, but due to the multiplication of the priors we can restrict ourselves to the 16 highest probabilities previously estimated. To balance the contribution of the anatomical priors and the intensity histograms, we also need to normalize the intensity priors sampled on the subject's intensities. We use the same approach, namely assuming that the 95[th] percentile of the highest kept $H_{i|j}(x)$ values have a probability of 0.95, separately for each contrast. The voxel-wise parcellation and posteriors obtained are shown in Fig. 8A.

axial view   coronal view   sagittal view

Figure 8: Successive parcellation results: A) voxel-wise posteriors and parcellation, B) diffused posteriors and parcellation, C) topology-corrected posteriors and final region-growing parcellation.

Markovian diffusion

The voxel-wise posteriors are independent from each other, and do not reflect the continuous nature of the structures. The next step is to combine information from neighboring voxels. We define a sparse Markov Random Field model for the posteriors:

$$P(x \in B_{i|j}|R,S,C) = \sum_{y \in C(x)} P(y \sim x|R)P(y \in B_{i|j}|R,S,C),$$

with $P(y \sim x|R) = \prod_R \exp -(R(y) - R(y))^2/2\sigma_R^2$, where σ_R is the median of the standard deviations $\sigma_{i|j,R}$ of the contrast histograms $H_{i|j}(R(x))$. The neighborhood $C(x)$ is defined as x itself and the four 26-connected neighboring voxels with highest probability $P(y \sim x|R)$, thus representing the neighbors most likely to be connected to x. The model is similar to a diffusion process and can be estimated with an iterated conditional modes (ICM) approach, updating sequentially the probabilities (Li, 2009):

$$P(x \in B_{i|j}|R,S,C) \leftarrow \sum_{y \in C(x)} P(y \sim x|R)P(y \in B_{i|j}|R,S,C)$$

from the initial voxel-wise posteriors until the ratio of changed parcellation labels decreases below 0.1, typically within 50-80 iterations. The diffused probabilities and parcellation are shown in Fig. 8B.

Topology correction

The final step of the parcellation algorithm takes a global view of the individual structures, growing from the highest posterior values inside toward the boundaries. This region growing approach makes the implicit assumption that posterior maps should be monotonically decreasing from inside to outside, which is not necessarily the case. Therefore, we perform first a topology correction step on the individual structure posteriors $P(x \in i|R,B,S,C) = \max_{i|j} P(x \in B_{i|j}|R,S,C)$ with a fast marching algorithm (Bazin et al., 2007b). While the corrected posterior is very similar to the original one (see Fig. 8C), it ensures that all regions obtained by growing to a threshold have spherical object topology.

Anatomical region growing

Last, we turn the posteriors into optimized parcellations, by growing them concurrently (to avoid overlaps) until the target volume for each structure is reached. Given the volume $V_i(R,B,S,C)$ of the parcellation of the diffused and topology-corrected posteriors, we define the following target volume:

$$\hat{V}_i = P(V_i|\mu_{V,i},\sigma_{V,i})V_i + (1 - P(V_i|\mu_{V,i},\sigma_{V,i}))\exp \mu_{V,i},$$

taking a weighted average of the volume estimated from the data and the prior volume. This approach ensures that even in extreme cases where some structures have low posteriors, they are still able to grow to a plausible size. The region growing algorithm is driven from the most likely voxels, defined as $P(x \in i|R,B,S,C) - \max_{j \neq i} P(x \in j|R,B,S,C)$, and further modulated to follow isocontours of the skeleton prior:

$$P(x \leftarrow y) \sim P(y \in i|R,B,S,C) - max_{j \neq i}P(y \in j|R,B,S,C) - |P(y \in S_i) - P(x \in S_i)|.$$

Directionality of internal structures is a useful tool for understanding mechanical function in bones (Maquer et al., 2015). Here, we adapt this concept by using the skeleton isocontours as a

representation of internal directionality, maintaining the intrinsic shape of structures. Thus, voxels with highest probability compared to the other structures and with similar distance to the internal skeleton are preferentially selected. The final parcellation is given in Fig. 8C.

Validation metrics

To validate the method, we compared the MASSP results and the expert delineations with the following three measures:

1. the Dice overlap coefficient (Dice, 1945) $D(A, B) = \frac{2A \cap B}{A+B}$, which measures the strict overlap between voxels in both delineation;
2. the dilated Dice overlap coefficient $dD(A, B) = \frac{A \cup d(B) + B \cup d(A)}{A+B}$, where $d(.)$ is a dilation of the delineation by one voxel, which measures the overlap between delineations allowing for one voxel of uncertainty;
3. the average surface distance $asd(A, B)$, measuring the average distance between voxels on the surface boundary of the first delineation to the other one and reciprocally, which measures the distance between both delineations.

We computed all three measures for the manual delineations from the two independent raters, as well as the ratio of overlaps (automated over manual) and distances (manual over automated) to compare both performances, as detailed in the Results section.

Software implementation

The proposed method, Multi-contrast Anatomical Subcortical Structure Parcellation (MASSP), has been implemented as part of the Nighres toolbox (Huntenburg et al., 2018), using python and java for optimized processing. The software is available in open source from https://github.com/nighres/nighres (release-1.3.0) and https://github.com/IMCN-UvA/imcn-imaging. A complete parcellation pipeline is included with the Nighres examples. Computations take under 30 minutes per subject on a modern workstation.

Acknowledgments

We would like to thank Josephine Groot, Nikita Berendonk, Nicky Lute for their help collecting the AHEAD database, and Wietske van der Zwaag and Matthan Caan for their help in setting up the MP2RAGEME sequence. We also thank Steven Miletić and Dagmar Timmann for stimulating discussions around this topic, and all undergraduate students who contributed to the manual delineations. This work was supported by a NWO Vici grant (BF) and a NWO STW grant (AA, BF).

Competing Interests

We have no competing interest to declare regarding this work.

References

Ahsan, R.L., Allom, R., Gousias, I.S., Habib, H., Turkheimer, F.E., Free, S., Lemieux, L., Myers, R., Duncan, J.S., Brooks, D.J., 2007. Volumes, spatial extents and a probabilistic atlas of the human basal ganglia and thalamus. NeuroImage 38, 261–270. https://doi.org/10.1016/j.neuroimage.2007.06.004

Alkemade, A., Mulder, M.J., Groot, J.M., Isaacs, B.R., van Berendonk, N., Lute, N., Isherwood, S.J.S., Bazin, P.-L., Forstmann, B.U., 2020. The Amsterdam Ultra-high field adult lifespan database (AHEAD): A freely available multimodal 7 Tesla submillimeter magnetic resonance imaging database. Under review.

Andersen, H.H., Johnsen, K.B., Moos, T., 2014. Iron deposits in the chronically inflamed central nervous system and contributes to neurodegeneration. Cellular and Molecular Life Sciences 71, 1607–1622. https://doi.org/10.1007/s00018-013-1509-8

Avants, B., Epstein, C., Grossman, M., Gee, J., 2008. Symmetric diffeomorphic image registration with cross-correlation: Evaluating automated labeling of elderly and neurodegenerative brain. Medical Image Analysis 12, 26–41. https://doi.org/10.1016/j.media.2007.06.004

Bazin, P.-L., Ellingsen, L.M., Pham, D.L., 2007a. Digital Homeomorphisms in Deformable Registration, in: Karssemeijer, N., Lelieveldt, B. (Eds.), Information Processing in Medical Imaging. Lecture Notes in Computer Science, vol 4584, pp. 211–222. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-540-73273-0_18

Bazin, P.-L., Pham, D.L., 2007b. Topology correction of segmented medical images using a fast marching algorithm. Computer Methods and Programs in Biomedicine 88, 182–190. https://doi.org/10.1016/j.cmpb.2007.08.006

Bazin, P.-L., Plessis, V., Fan, A.P., Villringer, A., Gauthier, C.J., 2016. Vessel segmentation from quantitative susceptibility maps for local oxygenation venography, in: 2016 IEEE 13th International Symposium on Biomedical Imaging (ISBI), pp. 1135–1138. https://doi.org/10.1109/ISBI.2016.7493466

Bazin, P.-L., Alkemade, A., van der Zwaag, W., Caan, M., Mulder, M., Forstmann, B.U., 2019. Denoising High-Field Multi-Dimensional MRI With Local Complex PCA. Frontiers in Neuroscience 13. https://doi.org/10.3389/fnins.2019.01066

Blum, H., 1973. Biological shape and visual science (part I). Journal of Theoretical Biology 38, 205–287. https://doi.org/10.1016/0022-5193(73)90175-6

Caan, M., Bazin, P.-L., Marques, J.P., de Hollander, G., Dumoulin, S.O., Zwaag, W., 2019. MP2RAGEME: T_1, T_2^*, and QSM mapping in one sequence at 7 tesla. Human Brain Mapping 40, 1786–1798. https://doi.org/10.1002/hbm.24490

Chakravarty, M.M., Bertrand, G., Hodge, C.P., Sadikot, A.F., Collins, D.L., 2006. The creation of a brain atlas for image guided neurosurgery using serial histological data. NeuroImage 30, 359–376. https://doi.org/10.1016/j.neuroimage.2005.09.041

Chakravarty, M.M., Steadman, P., van Eede, M.C., Calcott, R.D., Gu, V., Shaw, P., Raznahan, A., Collins, D.L., Lerch, J.P., 2013. Performing label-fusion-based segmentation using multiple

automatically generated templates: MAGeT Brain: Label Fusion Segmentation Using Automatically Generated Templates. Human Brain Mapping 34, 2635–2654. https://doi.org/10.1002/hbm.22092

Daugherty, A., Raz, N., 2013. Age-related differences in iron content of subcortical nuclei observed in vivo: A meta-analysis. NeuroImage 70, 113–121. https://doi.org/10.1016/j.neuroimage.2012.12.040

de Hollander, G., Keuken, M.C., van der Zwaag, W., Forstmann, B.U., Trampel, R., 2017. Comparing functional MRI protocols for small, iron-rich basal ganglia nuclei such as the subthalamic nucleus at 7 T and 3 T: Optimizing fMRI for Subcortical Nuclei. Human Brain Mapping 38, 3226–3248. https://doi.org/10.1002/hbm.23586

Dice, L.R., 1945. Measures of the Amount of Ecologic Association Between Species. Ecology 26, 297–302. https://doi.org/10.2307/1932409

Draganski, B., Ashburner, J., Hutton, C., Kherif, F., Frackowiak, R.S.J., Helms, G., Weiskopf, N., 2011. Regional specificity of MRI contrast parameter changes in normal ageing revealed by voxel-based quantification (VBQ). NeuroImage 55, 1423–1434. https://doi.org/10.1016/j.neuroimage.2011.01.052

Ewert, S., Plettig, P., Li, N., Chakravarty, M.M., Collins, D.L., Herrington, T.M., Kühn, A.A., Horn, A., 2018. Toward defining deep brain stimulation targets in MNI space: A subcortical atlas based on multimodal MRI, histology and structural connectivity. NeuroImage 170, 271–282. https://doi.org/10.1016/j.neuroimage.2017.05.015

Ewert, S., Horn, A., Finkel, F., Li, N., Kühn, A.A., Herrington, T.M., 2019. Optimization and comparative evaluation of nonlinear deformation algorithms for atlas-based segmentation of DBS target nuclei. NeuroImage 184, 586–598. https://doi.org/10.1016/j.neuroimage.2018.09.061

Fischl, B., Salat, D.H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., van der Kouwe, A., Killiany, R., Kennedy, D., Klaveness, S., Montillo, A., Makris, N., Rosen, B., Dale, A.M., 2002. Whole Brain Segmentation. Neuron 33, 341–355. https://doi.org/10.1016/S0896-6273(02)00569-X

Fjell, A.M., Westlye, L.T., Grydeland, H., Amlien, I., Espeseth, T., Reinvang, I., Raz, N., Holland, D., Dale, A.M., Walhovd, K.B., 2013. Critical ages in the life course of the adult brain: nonlinear subcortical aging. Neurobiology of Aging 34, 2239–2247. https://doi.org/10.1016/j.neurobiolaging.2013.04.006

Fonov, V., Evans, A.C., Botteron, K., Almli, C.R., McKinstry, R.C., Collins, D.L., 2011. Unbiased average age-appropriate atlases for pediatric studies. NeuroImage 54, 313–327. https://doi.org/10.1016/j.neuroimage.2010.07.033

Forstmann, B.U., Keuken, M.C., Schafer, A., Bazin, P.-L., Alkemade, A., Turner, R., 2014. Multi-modal ultra-high resolution structural 7-Tesla MRI data repository. Scientific Data 1. https://doi.org/10.1038/sdata.2014.50

Forstmann, B.U., de Hollander, G., van Maanen, L., Alkemade, A., Keuken, M.C., 2017a. Towards a mechanistic understanding of the human subcortex. Nature Reviews Neuroscience 18, 57–65. https://doi.org/10.1038/nrn.2016.163

Forstmann, B.U., Isaacs, B.R., Temel, Y., 2017b. Ultra High Field MRI-Guided Deep Brain Stimulation. Trends in Biotechnology 35, 904–907. https://doi.org/10.1016/j.tibtech.2017.06.010

Frazier, J.A., Chiu, S., Breeze, J.L., Makris, N., Lange, N., Kennedy, D.N., Herbert, M.R., Bent, E.K., Koneru, V.K., Dieterich, M.E., Hodge, S.M., Rauch, S.L., Grant, P.E., Cohen, B.M., Seidman, L.J., Caviness, V.S., Biederman, J., 2005. Structural Brain Magnetic Resonance Imaging of Limbic and Thalamic Volumes in Pediatric Bipolar Disorder. American Journal of Psychiatry 162, 1256–1265. https://doi.org/10.1176/appi.ajp.162.7.1256

Garzón, B., Sitnikov, R., Bäckman, L., Kalpouzos, G., 2018. Automated segmentation of midbrain structures with high iron content. NeuroImage 170, 199–209. https://doi.org/10.1016/j.neuroimage.2017.06.016

Herting, M.M., Johnson, C., Mills, K.L., Vijayakumar, N., Dennison, M., Liu, C., Goddings, A.-L., Dahl, R.E., Sowell, E.R., Whittle, S., Allen, N.B., Tamnes, C.K., 2018. Development of subcortical volumes across adolescence in males and females: A multisample study of longitudinal changes. NeuroImage 172, 194–205. https://doi.org/10.1016/j.neuroimage.2018.01.020

Hill, R.A., Li, A.M., Grutzendler, J., 2018. Lifelong cortical myelin plasticity and age-related degeneration in the live mammalian brain. Nature Neuroscience 21, 683–695. https://doi.org/10.1038/s41593-018-0120-6

Huntenburg, J.M., Steele, C.J., Bazin, P.-L., 2018. Nighres: processing tools for high-resolution neuroimaging. GigaScience 7. https://doi.org/10.1093/gigascience/giy082

Iglesias, J.E., Sabuncu, M.R., Van Leemput, K., 2013. A unified framework for cross-modality multi-atlas segmentation of brain MRI. Medical Image Analysis 17, 1181–1191. https://doi.org/10.1016/j.media.2013.08.001

Iglesias, J.E., Insausti, R., Lerma-Usabiaga, G., Bocchetta, M., Van Leemput, K., Greve, D.N., van der Kouwe, A., Fischl, B., Caballero-Gaudes, C., Paz-Alonso, P.M., 2018. A probabilistic atlas of the human thalamic nuclei combining ex vivo MRI and histology. NeuroImage 183, 314–326. https://doi.org/10.1016/j.neuroimage.2018.08.012

Ji, J.L., Spronk, M., Kulkarni, K., Repovš, G., Anticevic, A., Cole, M.W., 2019. Mapping the human brain's cortical-subcortical functional network organization. NeuroImage 185, 35–57. https://doi.org/10.1016/j.neuroimage.2018.10.006

Johansen-Berg, H., 2013. Human connectomics — What will the future demand? NeuroImage 80, 541–544. https://doi.org/10.1016/j.neuroimage.2013.05.082

Keuken, M.C., Bazin, P.-L., Schafer, A., Neumann, J., Turner, R., Forstmann, B.U., 2013. Ultra-High 7T MRI of Structural Age-Related Changes of the Subthalamic Nucleus. Journal of Neuroscience 33, 4896–4900. https://doi.org/10.1523/JNEUROSCI.3241-12.2013

Keuken, M.C., Bazin, P.-L., Backhouse, K., Beekhuizen, S., Himmer, L., Kandola, A., Lafeber, J.J., Prochazkova, L., Trutti, A., Schäfer, A., Turner, R., Forstmann, B.U., 2017. Effects of aging on T1, T2*, and QSM MRI values in the subcortex. Brain Structure and Function 222, 2487–2505. https://doi.org/10.1007/s00429-016-1352-4

Koshiyama, D., Fukunaga, M., Okada, N., Yamashita, F., Yamamori, H., Yasuda, Y., Fujimoto, M., Ohi, K., Fujino, H., Watanabe, Y., Kasai, K., Hashimoto, R., 2018. Role of subcortical structures on cognitive and social function in schizophrenia. Scientific Reports 8. https://doi.org/10.1038/s41598-017-18950-2

Li, Stan Z., 2009. Markov Random Field Modeling in Image Analysis, Advances in Pattern Recognition. Springer London, London. https://doi.org/10.1007/978-1-84800-279-1

Makowski, C., Béland, S., Kostopoulos, P., Bhagwat, N., Devenyi, G.A., Malla, A.K., Joober, R., Lepage, M., Chakravarty, M.M., 2018. Evaluating accuracy of striatal, pallidal, and thalamic segmentation methods: Comparing automated approaches to manual delineation. NeuroImage 170, 182–198. https://doi.org/10.1016/j.neuroimage.2017.02.069

Marquand, A.F., Haak, K.V., Beckmann, C.F., 2017. Functional corticostriatal connection topographies predict goal-directed behaviour in humans. Nature Human Behaviour 1. https://doi.org/10.1038/s41562-017-0146

Marques, J.P., Kober, T., Krueger, G., van der Zwaag, W., Van de Moortele, P.-F., Gruetter, R., 2010. MP2RAGE, a self bias-field corrected sequence for improved segmentation and T1-mapping at high field. NeuroImage 49, 1271–1281. https://doi.org/10.1016/j.neuroimage.2009.10.002

McCoy, R.C., Wakefield, J., Akey, J.M., 2017. Impacts of Neanderthal-Introgressed Sequences on the Landscape of Human Gene Expression. Cell 168, 916-927.e12. https://doi.org/10.1016/j.cell.2017.01.038

Miletić, S., Bazin, P.-L., Weiskopf, N., van der Zwaag, W., Forstmann, B.U., Trampel, R., 2020. fMRI protocol optimization for simultaneously studying small subcortical and cortical areas at 7T. NeuroImage, under review.

Mosley, P.E., Smith, D., Coyne, T., Silburn, P., Breakspear, M., Perry, A., 2018. The site of stimulation moderates neuropsychiatric symptoms after subthalamic deep brain stimulation for Parkinson's disease. NeuroImage: Clinical 18, 996–1006. https://doi.org/10.1016/j.nicl.2018.03.009

Patenaude, B., Smith, S.M., Kennedy, D.N., Jenkinson, M., 2011. A Bayesian model of shape and appearance for subcortical brain segmentation. NeuroImage 56, 907–922. https://doi.org/10.1016/j.neuroimage.2011.02.046

Pauli, W.M., Nili, A.N., Tyszka, J.M., 2018. A high-resolution probabilistic in vivo atlas of human subcortical brain nuclei. Scientific Data 5. https://doi.org/10.1038/sdata.2018.63

Qiu, A., Brown, T., Fischl, B., Ma, J., Miller, M.I., 2010. Atlas Generation for Subcortical and Ventricular Structures With Its Applications in Shape Analysis. IEEE Transactions on Image Processing 19, 1539–1547. https://doi.org/10.1109/TIP.2010.2042099

Sitek, K.R., Gulban, O.F., Calabrese, E., Johnson, G.A., Lage-Castellanos, A., Moerel, M., Ghosh, S.S., De Martino, F., 2019. Mapping the human subcortical auditory system using histology, postmortem MRI and in vivo MRI at 7T. eLife 8. https://doi.org/10.7554/eLife.48932

Smith, S.M., 2002. Fast robust automated brain extraction. Human Brain Mapping 17, 143–155. https://doi.org/10.1002/hbm.10062

Steiner, Heinz, Tseng, Kuei Y. (Eds.), 2017. Handbook of Basal Ganglia Structure and Function, Second Edition, Handbook of Behavioral Neuroscience. Elsevier. https://doi.org/10.1016/C2014-0-02359-3

Tardif, C.L., Gauthier, C.J., Steele, C.J., Bazin, P.-L., Schäfer, A., Schaefer, A., Turner, R., Villringer, A., 2016. Advanced MRI techniques to improve our understanding of experience-induced neuroplasticity. NeuroImage 131, 55–72. https://doi.org/10.1016/j.neuroimage.2015.08.047

Turner, R., 2019. Myelin and Modeling: Bootstrapping Cortical Microcircuits. Frontiers in Neural Circuits 13. https://doi.org/10.3389/fncir.2019.00034

Visser, E., Keuken, M.C., Douaud, G., Gaura, V., Bachoud-Levi, A.-C., Remy, P., Forstmann, B.U., Jenkinson, M., 2016a. Automatic segmentation of the striatum and globus pallidus using MIST: Multimodal Image Segmentation Tool. NeuroImage 125, 479–497. https://doi.org/10.1016/j.neuroimage.2015.10.013

Visser, E., Keuken, M.C., Forstmann, B.U., Jenkinson, M., 2016b. Automated segmentation of the substantia nigra, subthalamic nucleus and red nucleus in 7 T data at young and old age. NeuroImage 139, 324–336. https://doi.org/10.1016/j.neuroimage.2016.06.039

Wang, B.T., Poirier, S., Guo, T., Parrent, A.G., Peters, T.M., Khan, A.R., 2016. Generation and evaluation of an ultra-high-field atlas with applications in DBS planning, in: Styner, M.A., Angelini, E.D. (Eds.), . Presented at the SPIE Medical Imaging, San Diego, California, United States, p. 97840H. https://doi.org/10.1117/12.2217126

Xiao, Y., Fonov, V., Bériault, S., Subaie, F.A., Chakravarty, M.M., Sadikot, A.F., Pike, G.B., Collins, D.L., 2015. Multi-contrast unbiased MRI atlas of a Parkinson's disease population. International Journal of Computer Assisted Radiology and Surgery 10, 329–341. https://doi.org/10.1007/s11548-014-1068-y

Yeatman, J.D., Wandell, B.A., Mezer, A.A., 2014. Lifespan maturation and degeneration of human brain white matter. Nature Communications 5. https://doi.org/10.1038/ncomms5932

Yelnik, J., Bardinet, E., Dormont, D., Malandain, G., Ourselin, S., Tandé, D., Karachi, C., Ayache, N., Cornu, P., Agid, Y., 2007. A three-dimensional, histological and deformable atlas of the human basal ganglia. I. Atlas construction based on immunohistochemical and MRI data. NeuroImage 34, 618–638. https://doi.org/10.1016/j.neuroimage.2006.09.026

Zhang, Y., Wei, H., Cronin, M.J., He, N., Yan, F., Liu, C., 2018. Longitudinal atlas for normative human brain development and aging over the lifespan using quantitative susceptibility mapping. NeuroImage 171, 176–189. https://doi.org/10.1016/j.neuroimage.2018.01.008