Azole use in agriculture reduces *Aspergillus fumigatus* abundance but does not alter its population structure

Amelia E. Barber[1,2] #, Jennifer Riedel[3] #, Tongta Sae-Ong[1], Kang Kang[1], Werner Brabetz[4], Gianni Panagiotou[1,5], Holger B. Deising[3], Oliver Kurzai[1,2] *

# Denotes equal contribution

* Address correspondence to: okurzai@hygiene.uni-wuerzburg.de

Running title: The impact of azole use in agriculture on *Aspergillus fumigatus*

Keywords: *Aspergillus fumigatus, Aspergillus*, antifungal resistance, azole, fungicide, population genomics

Author affiliations:

[1] Leibniz Institute of Natural Product Research and Infection Biology – Hans Knöll Institute. Jena, Germany.

[2] Institute for Hygiene and Microbiology. University of Würzburg. Würzburg, Germany

[3] Institute for Agriculture and Nutritional Sciences. Martin Luther University Halle-Wittenberg. Halle (Saale), Germany.

[4] Biotype GmbH. Dresden, Germany.

[5] Systems Biology and Bioinformatics Group. School of Biological Sciences. Faculty of Sciences. The University of Hong Kong. Hong Kong, China.

ABSTRACT

Antibiotic resistance is an increasing threat to human health. In the case of *Aspergillus fumigatus*, which is both an environmental saprobe and an opportunistic human fungal pathogen, resistance is suggested to arise from fungicide use in agriculture, as the azoles used for plant protection are almost identical to the frontline antifungals used clinically. However, limiting azole fungicide use on crop fields to preserve their activity for clinical use could threaten the global food supply via a reduction in yield. In this study we clarify the link between fungicide use on crop fields and resistance in a prototypical human pathogen through systematic soil sampling on farms in Germany and surveying fields before and after azole application. We observe a reduction in the abundance of *A. fumigatus* on fields following azole treatment—a finding that was not observed on an organic control field not applying azoles. However, the overall resistance frequency among agricultural isolates is low, with only 1-3% of isolates from 2016-2018 displaying resistance to medical azoles. Isolates collected after the growing season and azole exposure show a subtle, but consistent decrease in susceptibility to medical and agricultural azoles. Whole genome sequencing indicates that, despite the decrease in fungal abundance, azole application does not significantly affect the population structure and genetic diversity of *A. fumigatus* in fields. Given the low observed resistance rate among agricultural isolates, as well the lack of genomic impact following azole application, we do not find evidence that azole use on crops is significantly driving resistance in *A. fumigatus*.

IMPORTANCE

Antibiotic resistance is an increasing threat to human health. In the case of

Aspergillus fumigatus, which is an environmental fungus that also causes life-

threatening infections in humans, antimicrobial resistance is suggested to arise from

fungicide use in agriculture, as the chemicals used for plant protection are almost

identical to the antifungals used clinically. However, removing azole fungicides from crop

fields threatens the global food supply via a reduction in yield. In this study, we survey

crop fields before and after fungicide application, and find that while azole fungicide

application reduces the abundance of *A. fumigatus* in the soil, we do not find evidence

that fungicide use is significantly driving resistance in this organism and context.

INTRODUCTION

Aspergillus fumigatus is a globally-distributed fungus responsible for an estimated

300,000 case of invasive disease and more than 10 million cases of chronic and allergic

disease globally each year (1). Humans inhale the infectious particles, or spores, on a

daily basis, but are actually an accidental host for the fungus whose primary niche is soil

and decaying vegetation. Management and prophylaxis against aspergillosis relies

largely on the azole class of antifungals with voriconazole and isavuconazole

recommended as first line therapy (2). Unfortunately, clinical resistance to the azoles in

A. fumigatus is an increasing problem, with some medical centers reporting rates as

high as 30% in specific patient populations, and similarly high rates for environmentally-

isolated *A. fumigatus* (3, 4). Regrettably, the mortality for resistant infections is upwards

of 90% in some patient populations (5-7). While resistance can evolve during patient

therapy (8, 9), the emergence of resistance in *A. fumigatus* has mainly been linked to

the use of azoles in agriculture, as structurally similar and mechanistically

indistinguishable compounds are heavily used for plant protection (10). This resistance

has been described as collateral damage as *A. fumigatus* is not a plant pathogen that is

being directly targeted by fungicide treatments (11). The triazoles were first released for

agriculture in 1973, well before they were first introduced to human medicine in the early

1990s, and are currently the most widely-utilized antifungal compound group in

agriculture due to their systemic distribution in treated plants, high efficiency and broad

spectrum of target pathogens (12, 13). Crops, particularly cereals and fruits, are sprayed

multiple times each growing season at a recommended dose of 100 g/hectare to control

powdery mildew, rust, septoria leaf blotch, and other phytopathogenic fungi (14).

Currently, there are 32 azoles commercially available for plant protection (15), but only

five in regular use in human medicine (16).

The most common azole resistance mechanism in *A. fumigatus* occurs via

mutations in the target protein of azole fungicides, sterol 14α-demethylase (CYP51A,

syn ERG11), a key enzyme of the ergosterol biosynthesis pathway. In *A. fumigatus*, the

dominant resistant mechanism among both environmental and clinical isolates is a 34 bp

tandem repeat (TR_{34}) in the *cyp51a* promoter coupled with a leucine to histidine

substitution (L98H) in the amino acid coding sequence—the net effect of which is an

increase in gene expression as well as an alteration in both the stability of the target

enzyme and the interaction between the protein heme cofactor of *cyp51a* and the azole

ligand (17-19). Additional mutations that have been identified to confer azole resistance

in *A. fumigatus* include other variations of the tandem repeat, such as

TR_{46}/Y121F/T289A and TR_{53}, as well as other point mutations in the *cyp51a* coding

sequence (20, 21).

Disease-causing fungi are responsible for roughly 20% of crop yield loss, with a

further 10% loss postharvest (16), so the use of azoles is critical for securing the food

supply. However, this must be balanced against the need to preserve the activity of the

azoles for clinical use and, as such, there is an urgent need to identify the contributions

of azole fungicide use on food crops to the development of resistance in *A. fumigatus*.

We address this through systematic soil sampling conducted on 10 agricultural sites in

Germany over a three-year period—including conventionally-managed fields applying

azoles fungicides, as well as those practicing organic agriculture that do not use these

compounds. In the largest published *A. fumigatus* sequencing effort to date, and the first

to focus on the fungus in its natural niche, we also use whole genome sequencings

(WGS) to examine the impact of azole fungicides on the population genetics of 64

agriculturally-isolated *A. fumigatus* isolates.

RESULTS

Variable abundance of *Aspergillus fumigatus* on agricultural sites in Germany

To examine the depth distribution of *A. fumigatus* in agricultural soils, we

collected soil samples from a test field at five cm intervals down to a depth of 30 cm

below the surface. *A. fumigatus* was most abundant in the top five cm of soil and was

not significantly observed below a depth of 15 cm (Fig 1A). As a result, only the top 5 cm

layer of soil was collected in subsequent soil samples.

To identify whether our sampling areas are representative of azole and fungicide

usage for Germany as a whole, we calculated the fraction of each administrative district

that is theoretically treated with azoles in the context of agriculture using publicly

available data. Overall, 51% of Germany is designated as agricultural land (see Material

& Methods for data sources). However, not all this land is sprayed with azoles. For example, meadow or pasture land is rarely treated with fungicides. Additionally, approximately 7% of agriculture in Germany utilizes organic farming and does not apply azoles, with a range of 3-15% between the different federal states. Using land use data on arable farm land and permanent crop areas for each district in Germany, we calculated that the mean fraction of potentially azole-treated area in Germany is 32% (for the districts where data is available), with a range of 1-68% (Fig 1B). The districts where we performed our soil sampling were almost all above the average for Germany, with a mean of 52% potentially azole-treated hectares and a range of 30-65%.

To examine the inter- and intra-field variability in the density of *A. fumigatus* on agricultural fields, soil samples were taken from nine conventional and eight organic fields before the growing season in 2017. Conventional fields were also sampled again after the vegetative period and application of azoles. In total, 2875 soil samples were taken between 2016 and 2018 (Supplemental Table 1). The predominant crops being grown were cereals such as wheat and barley, but several apple orchards were also sampled. Of the fields sampled during this period in 2017, 67% of the soil samples taken were positive for *A. fumigatus* with a large range between fields (28-100%) (Fig 1C). We also observed a large degree of variation in the mean CFU/g of soil between different fields, with some fields having 30x higher *A. fumigatus* density over others (0.7 – 18.8 CFU/g).

To examine the stability of *A. fumigatus* population sizes in agricultural soil, we investigated the same farms a year later in the spring of 2018 and repeated the soil sampling on eight conventional and seven organic fields. Due to crop rotation, it was not possible to sample the identical fields as the previous year, except for the apple

orchards on Farms H and L. During the 2018 sampling period, an overall lower proportion of samples were positive for *A. fumigatus* (51% with a range of 10-96% between different fields) (Fig 1D), but the mean CFU/g for the 1000 samples was similar to the previous year (5.18 CFU/g soil for 2017 compared to 5.74 CFU/g for 2018). As in the previous year, there was a large variability in the mean CFU of *A. fumigatus* present between fields. Interestingly, the farms with the highest mean *A. fumigatus* CFU/g soil in 2017 (*e.g.* Farms B and D) were also the highest in 2018, and the farm with the lowest amount of *A. fumigatus* in 2017 (Farm K) was also the lowest in 2018, suggesting that the amount of *A. fumigatus* is consistent year-to-year (Fig 1E). To investigate potential factors that might support a higher abundance of *A. fumigatus*, we compared the total organic carbon (TOC) content of a random selection of samples with the CFU/g *A. fumigatus,* but did not detect a clear relationship between the two (Supplemental Fig 1). Altogether, we observed a non-uniform distribution of *A. fumigatus* in soil samples taken from the same field and a large degree of heterogeneity between fields, but the relative amounts of *A. fumigatus* remained largely consistent year-to-year.

Azole application reduces *A. fumigatus* abundance

To examine the impact of azole fungicides on *A. fumigatus* in agricultural soil, we performed additional soil sampling on the conventional fields surveyed in the spring at the end of the vegetative period and after several months of fungicidal crop protection. A schematic illustration of the soil sampling and fungicide application time lines for 2017 and 2018 can be found in Supplemental Figures 2A and B. Unfortunately, the fungicide history for Farms H and L was not available to us. Fields were typically treated with fungicides twice during the growing period and every application recorded contained at

least one azole [Note: fungicides are often applied as commercially-available cocktails of different chemicals].

When comparing the amount of *A. fumigatus* on fields before azole exposure to after azole exposure, we detected a significant reduction in the CFU/g soil on the majority of fields in 2017 (Fig 2A), even though it is not being directly targeted as a plant pathogen. A similar trend was observed in 2018 comparing *A. fumigatus* density before and after azole treatment, but the results were less dramatic (Fig 2B), possibly due to the extreme heat and drought experienced in Germany during the growing season.

To confirm that the reduction in agricultural *A. fumigatus* populations was the result of azole application, and not a seasonal effect from comparing April to July, in 2017 we took monthly soil samples from a conventional field and an organic field not treated with azoles as a control. Samples were taken beginning in April, before azoles were applied to the conventional field, through the harvest period in July, and then additional samples were taken in October and November to allow for a period without azole fungicide application. From April to July, the conventional field was sprayed with azoles every 3-5 weeks. When comparing the abundance of *A. fumigatus* on the organic field, we did not observe any significant differences between the abundance recorded monthly between April and July (Fig 2C). However, the conventional field showed a significant reduction in abundance between April and May, corresponding to the beginning of the azole application period, and this reduction was maintained through the rest of the azole application period (Fig 2D). In summary, we observed a reduction in *A. fumigatus* abundance on conventional fields following the application of azole fungicides, suggesting that this selection pressure might be imposing a population bottleneck in this niche.

Reduced susceptibility to agricultural azoles in populations isolated after the growing season and azole exposure

To assess the susceptibility of *A. fumigatus* to commonly-applied agricultural

azoles, approximately 20 isolates per field and time period from 2017 and 2018 were

screened for their ability to grow at a set concentration of difenoconazole and

tebuconazole (n = 435 for 2017 and 342 for 2018). To limit potentially clonal isolates

from skewing the results, a maximum of two isolates per soil sample were included for

testing. As there are no established breakpoints for defining resistance to these

compounds in *A. fumigatus*, we selected concentrations that mimicked MIC90 values for

these azoles (1 mg/L for difenoconazole and 2 mg/L for tebuconazole). When examining

conventional and organic farms in the spring, we observed a wide range in the fraction

of isolates per field that were able to grow when challenged with agricultural azoles. For

difenoconazole, this ranged from 10-65% per field in 2017 (n = 17 fields, 320 isolates in

total) and 0-50% in 2018 (n = 15 fields, 261 isolates in total) (Fig 3A and Supplemental

Table 2). For tebuconazole, the rates ranged from 0-25% in 2017 and 0-20% in 2018

(Fig 3B and Supplemental Tables 2). We did not detect any significant differences in the

rates between conventional and organic fields in the spring or between fields growing

different crops (cereals or apples).

Given that we observed a reduction in the *A. fumigatus* population size on

conventional fields following azole application, we wanted to examine the effect of azole

fungicides on the local susceptibility following the vegetative period and several months

of azole application. We observed an increase in the proportion of isolates that were

able to grow at the test concentrations of difenoconazole (1 mg/L) and tebuconazole

(2mg/L) for fields sampled after azole exposure compared to the same field in the spring prior to azole application (Fig 3C, D, and Supplemental Table 3). We detected a 1.67-fold increase in the proportion of isolates that were resistant to our test concentration of difenoconazole in 2017 with a range from a -1.4 to 4.5-fold for individual fields (Fig 3C). In 2018, this increased to a 2.85-fold with a range of 1-4.5-fold change increase for individual fields. We also saw a similar increase in the fraction of isolates with reduced susceptibility to our test concentration of tebuconazole after azole exposure compared to the before. In 2017, we observed a 4.9-fold in number of isolates that were able to grow at the test concentration after azole exposure vs. before exposure with a range of 0-25.0% for individual fields (Fig 3D). In 2018, we detected a more modest 1.8-fold with range of -5 to 18.2% for individual fields.

For the fields growing cereals, we were not able to sample the same fields over subsequent years due to crop rotation and the fields not being in use the following year. However, we were able to compare the same apple fields in both 2017 and 2018. We tracked the local susceptibility to agricultural azoles in these fields over two consecutive years at two time points: in the spring prior to azole application and just prior to harvest after ≈ 3 months of azole application. The fraction of isolates that were able to grow in the presence of 1 mg/L difenoconazole or 2 mg/L tebuconazole was, in general, low in the spring, and increased after azole application (Fig 3E, Supplemental Table 3). Interestingly, the spring of 2018, the proportion had returned to a level comparable to what we observed in the spring of 2017—indicating that the reduced susceptibility is transient and recedes when the selective pressure of the azoles is removed. In summary, we found a wide range of susceptibilities to agricultural azoles between different fields, but a consistent decrease in susceptibility following the growing season

and azole exposure. However, this change is seemingly transient, and the *A. fumigatus*

populations from azole-treated fields typically returned to what they were prior to azole

application by the following spring.

<u>Resistance to medical azoles is agricultural *A. fumigatus* isolates</u>

We next examined our isolate collection for resistance to medical azoles and

determined the proportion that would be considered clinically resistant. Using the

VIPcheck™ agar-based screening method followed by broth microdilution for isolates

showing growth on agar-containing wells (22, 23), we determined that only a very small

fraction of *A. fumigatus* isolated from agriculture showed resistance to itraconazole,

voriconazole, or posaconazole in 2016-2018 (Figure 3F). The overall resistance rate to

itraconazole among all isolates collected was highest, with 3.0% (11/333) of isolates

being resistant in 2016, 0.7% in 2017 (4/460), and 0.6% (2/322) in 2018. We observed

lower resistance rates for posaconazole and voriconazole, with only 2.1% (7/333) being

resistant in 2016, 0.7% (4/460) in 2017, and 0.0% (0/322) in 2018.

To quantify what mutations were responsible for azole resistance in the

population analyzed, we genotyped the *cyp51a* locus of all resistant isolates using

Sanger sequencing (n = 18). This gene is the target of the azoles and the major source

of resistance among both environmental and clinical *A. fumigatus*. We found that the

most dominant DNA alteration observed was the well-characterized TR$_{34}$/L98H mutation

(Fig 3G). This *cyp51a* genotype accounted for 6/12 (50%) of resistant isolates in 2016

and 3/4 (75%) in 2017. In 2018, we identified only two resistant isolates among the 322

screened and both had wild type *cyp51a* loci. However, both of these isolates were only

weakly resistant to itraconazole, but not other azoles, with itraconazole MIC values right

at the breakpoint of 2-4 mg/L. For comparison, resistant isolates from other years had MIC values of >8 mg/L. In total, the genetic cause of resistance remained unknown for 8 isolates from 2016-18.

Since the majority of fields had no resistant isolates, it was not possible to effectively compare resistance rates among agricultural *A. fumigatus* for medical azoles before and after azole treatment. In lieu of this, we examined the MIC distribution for isolates collected before and after the growing season and azole application. Examining 40 isolates from the before period and 40 isolates from after azole exposure, we observed a shift in the MIC distribution towards higher MICs for all azoles examined, both medical and agricultural (Fig 3H). However, the median MICs remained unchanged, indicating that the majority of the population following azole exposure does not exhibit a change in MIC. Taken together, these results indicate that the rate of resistance to clinical azoles among environmental *A. fumigatus* isolated from agricultural environments is overall low and that exposure to agricultural azoles only causes a minor increase in MIC values for some of the population, with the majority of isolates left unchanged.

No population structure observed for *A. fumigatus* in agricultural fields

To better understand the population structure of *A. fumigatus* in the environment and how it is impacted by the azoles, we performed whole genome sequencing on isolates from four conventional farms collected before and after azole exposure. 64 isolates were sequenced by Illumina paired-end sequencing, representing eight isolates per farm and time point. Isolates from each field and time point were randomly selected, with a maximum of two per soil sample to avoid sequencing of clonal isolates. Raw

reads were checked for quality and then aligned to the Af293 reference genome, resulting in a median depth of coverage after mapping of 31.5X (range 10-90X) and a median genome coverage median of 94% (range 92.8-96.7%). Consistent with what has been observed among sequenced clinical strains (24), isolates differed from the Af293 reference by a median of 84,690 single nucleotide variants (SNVs) or 2.88 SNVs/kb, with a range of 65,854-146,055 SNVs. The analysis of copy-number variations (CNVs) identified 8,277 unique CNVs in total, with a median of 3,115 CNVs per isolate (range 1,666-4,532 CNVs). These CNVs were further delimited into a median of 2,589 deletions (range 1,247-4,405) and 5,687 insertions (range 3,872-7,030).

A maximum-likelihood phylogeny based on SNVs indicated no population stratification among isolates from different farms and regions of Germany (Fig 4). To more directly assess the association between genetic and geographic distance, we performed a Mantel test correlating a geographic distance matrix with the F_{ST} genetic distance matrix, and observed no significant association between the two. When considering the azole-resistance status of the isolates, the two itraconazole-resistant TR$_{34}$/L98H isolates sequenced clustered next to each other, despite originating from separate farms, while the third itraconazole-resistant isolate with an undefined resistance mechanism was on a distinct branch. Despite being the nearest sequenced neighbors, the two TR$_{34}$/L98H isolates were genetically distinct, each possessing 19,439 and 60,841 unique SNVs not shared by the other isolate, along with 67,196 common SNVs relative to Af293.

Comparative analysis of molecular variance (AMOVA) indicated that the majority of the variation seen among the 64 isolates came from the population as a whole (within sample) (94.8%) and between samples (5.0%) (Table 1). There was no significant

molecular variance between farms (0.2%), with the exception of modest variation

between Farm B and Farm C (1.2% of variation observed). Weighted Weir and

Cockerham's fixation indexes (F_{ST}) for each farm were essentially zero, indicating an

interbreeding, panmimetic population with no separation between farms (Supplemental

Fig 3A). V_{ST} values estimating population differentiation based on copy number variation

also indicated no subdivision among the farms (Supplemental Fig 3B). To examine the

genetic diversity within farms, nucleotide diversity (the average number of nucleotide

differences per site for all possible pairs in the population or π), and the number of

polymorphic sites (Watterson estimator or θ) were calculated along 5 kb windows with a

500 bp step size for each farm population. Farm C showed the greatest intra-farm

diversity, while Farm E showed the smallest (Supplemental Fig 3C-D). Taken together,

there was no population differentiation between *A. fumigatus* isolates from different

farms in Germany, a finding in line with the fungus' capacity for aerosol dispersal.

Changes in population genetics following azole exposure vary by field

Given the observed reduction in overall *A. fumigatus* abundance after azole

treatment, we examined the populations for changes in genetic diversity and evidence of

selective sweeps in *A. fumigatus* field populations. Using 5 kb sliding windows with a 500

bp step size, nucleotide diversity (π) was calculated for the individual farms before and

after azole treatment. No clear trend was seen regarding changes in nucleotide diversity

between the time points. The isolates from Farms B and E showed increased diversity

following azole application, while nucleotide diversity decreased on Farms C and D (Fig

5A). We also calculated the number of segregating sites (θ) for the same 5 kb windows

and observed the same lack of consensus. Farms B and C showed similar values of θ

before as compared to after azole application, while Farm D showed a dramatic decrease in θ following azole application (Fig 5B). AMOVA analysis on isolates from before and after the vegetative period and azole exposure did not indicate any significant molecular variation between the time periods, with the majority of the variation being between and within samples (Table 2). Finally, we measured Tajima's D to test neutrality along 5 kb windows, where negative values indicate less variation than expected and are indicative of a selective sweep. Positive values denote a population that is more heterogenous than would be expected and suggest either a sudden population contraction or balancing selection. Overall, the bulk of the Tamija's D values were close to neutral and there was no clear trend between farms, indicating that there was no genomic signature of a population bottleneck or selective sweep following azole exposure (Fig 5C). The median Tajima's D was roughly zero for Farm B and increased slightly to 0.37 following azole exposure prior to azole application, and the same direction shift was seen for Farm D, but the starting Tajima's D was negative at the time point before azole exposure (-0.53 to 0.53) (Fig 5C). Conversely, the median Tajima's D for farms C and E shifted from positive to negative after the growing season and azole exposure (0.75 to -0.57 for field C and 0.61 to -0.10 for Field E). Taken together, these results indicate that despite the reduction in abundance of *A. fumigatus* on agricultural fields following azole application, we were unable to detect any marked changes at the population level.

DISCUSSION

The use of azole fungicides for plant protection has been previously suggested as a driver of clinical resistance in the environmental saprobe and human pathogen

A. fumigatus, as well as the emergence of new fungal pathogens such as *Candida auris*

(11, 25, 26).   However, direct evidence linking azole use in agriculture and clinical

resistance is missing. Additionally, delineation of specific roles for the use of azoles in

crops vs. ornamental plants such as flower bulbs has not been defined and is important

to make, as limiting azole fungicide use would significantly impact disease control and

yield in many crops. In this study, we provide the first systematic investigation of the

impact of azole fungicides on the ecology and resistance status of a human pathogen,

A. fumigatus. Through analysis of 2,875 soil samples over a three year period in central

Germany, we found an overall low incidence of *A. fumigatus* isolates that would be

considered clinically resistant (1-3%). However, we observed a reduced abundance of

the fungus following the growing season and azole application, which coincided with a

modest, but consistent decrease in azole susceptibility. Interestingly, this susceptibility

was transient and reset by the following spring. We also assessed the influence of azole

application on *A. fumigatus* population dynamics by WGS and were unable to find a

clear impact on the population structure or genetic diversity.

Despite sampling on fields that were actively treated with azole fungicides, and in

regions of Germany with above average fungicide exposure, only 1-3% of *A. fumigatus*

isolates collected were resistant to medical azoles. This incidence is in agreement with

clinically-reported frequencies in Germany and other European countries, where the rate

of triazole resistance ranged from 0.6% to 4.2% (3.2% in Germany) (27, 28). The

environmental resistance rates reported for Europe, however, have been much broader.

Some studies have found environmental resistance frequencies approaching 20%, while

others have reported virtually no incidence of resistant environmental isolates (29-34).

Part of these differences could be attributed to differences in methodology, such as the

usage of azole selection during the isolation procedure or inclusion of potentially clonal

isolates from a given sample to skew the data. In the current study, we avoided azole

selection during the isolation procedure, as well as only allowed two isolates per soil

sample, to avoid potentially clonal isolates that influence the results. Another potentially

contributing factor could be that while all technically "environmental" isolates, rural or

agricultural settings could be a completely different niche than an urban flower garden—

a supposition supported by a recent study where rural areas in the UK had much lower

resistance rates (1.1%) than urban locations (13.8%) (33).

Another important and novel finding from this study is that isolates from after the

vegetative period and azole exposure consistently showed decreased azole

susceptibilities to difenoconazole and tebuconazole, as well as the subtle MIC shift to

both agricultural and medical azoles. However, one shortcoming to our study is that we

did not collect isolates from organic fields at a matching time point for susceptibility

testing, so we cannot exclude that the changes observed in azole susceptibility are not

also influenced by seasonal changes. The observation that changes in susceptibility are

transient and reset in the period between the end of the growth period and the following

spring on the two field is intriguing and worthy of further study. This transformation could

either be the result of a naïve population coming in via aerosol dispersal, or a

consequence of isolates acquiring only transient resistance, a phenomena previously

observed in other plant pathogenic fungi, as well as the human pathogen *Mucor*

circinelloides (35-37).

We observed a wide range in the CFU/g soil within and between farms during

annual spring sampling, but the mean level for each farm was stable between years for

reasons that remain undiscovered. Our mean *A. fumigatus* CFU/g soil was in line with

what was reported recently for abundance in wheat grain, maize silage, and fruit waste

(38). However, this abundance is several magnitudes lower than that reported for *A.*

fumigatus in flower bulb waste and green material waste in this same study, where it

was not uncommon to isolate 10_4 CFU/g. Importantly, we demonstrated a reduction in

the *A. fumigatus* population size on fields in the context of azole application for plant

protection. Despite this apparent population contraction, there were no substantial or

consistent changes in the population genomics of *A. fumigatus* on the fields we

sampled. One possible confounding factor in comparing time points before and after

azole exposure could be the significant aerial dispersion of the fungus. Once the

effective fungicide concentration decreased sufficiently, but before we collected samples

after azole application, new isolates could be introduced to the field and add their

genetic diversity.

Our study also provides the first WGS-based study focused on *A. fumigatus* in its

natural niche, the environment. Previous studies have primarily concentrated on clinical

isolates, with particular priority given to resistant strains (24, 39). Even while sampling

within the same field, we found a large degree of genetic diversity, where the majority of

diversity came from within samples. We also did not observe a defined population

structure or separation between farms or regions. The degree to which this

environmental diversity is recapitulated in clinical isolates, and whether there are

enrichments for particular subgroups in the transition from environment to clinic is an

interesting question for further study.

Given our low observed resistance rate among agricultural isolates, as well the lack of genomic impact of azole application on the fields surveyed, our study does not find evidence that azole fungicide use in crop agriculture significantly contributes to resistance in *A. fumigatus,* nor should be necessarily removed from use in this context due to their crucial role in global food production. Instead, we propose a refined model, where azole resistance in *A. fumigatus* is being driven by the use of these compounds in the cultivation of flowers and other ornamentals. It is common practice to dip flower bulbs directly in fungicide stocks prior to storage and shipment, and their use in this setting should be more carefully considered and managed (40, 41). This revised model is supported by the recent identification of flower bulb and green waste as environmental "hotspots" for resistance selection, which possess both high overall fungal colony counts as well as increased incidence of resistant *A. fumigatus* isolates compared to soil (38). Unfortunately, due to massive aerial dispersal of *A. fumigatus* conidia, the use of azoles in any hotspot can lead to the worldwide distribution of resistant strains.

MATERIALS AND METHODS

Site selection and soil sampling

During 2016 to 2018, soil sampling was conducted on agricultural sites in the federal states of Thuringia, Saxony-Anhalt, and Saxony with the approval of the land owner and/or relevant ministries. The majority of fields sampled were growing cereals such as wheat or barley, but some apple orchards were sampled as well (see Supplemental Table 1 for full details). Farms were arbitrarily assigned an alphabetic identifier (A-L) and specific fields a numeric identifier as described in the supplementary Materials &

Methods. Due to crop rotation, the same field could not be surveyed over subsequent

years, with the exception of the apple orchards. Soil samples were collected at

beginning of the vegetation period on the conventional and organic cultivated sites and

after azole application an additional sampling was carried out on the conventional sites.

In general, 50 soil samples per site and type of farming (conventional or organic) were

collected, with a total of approximately 1000 soil samples per year. Soil samples were

selected to best cover the field with a minimum distance of 1 m between samples. For

each sample, the top layer of soil was collected by a metal spatula into a sterile sample

cup and refrigerated until processing.

<u>Soil processing and isolation of *A. fumigatus*</u>

3 g of soil from each sample cup was weighed out and resuspended in 8 mL 0.2M NaCl

containing 1% Tween20. Samples were vortexed vigorously and then left to settle until a

phase separation became apparent. 2 mL of the upper phase was transferred to a new

tube for plating onto Sabouraud Glucose Agar (SGA) containing 50 µg/mL

Chloramphenicol (Sigma-Aldrich, Taufkirchen, Germany). Of this 2 mL, 150 µL was

plated onto one plate and the remaining volume was plated onto a second plate in order

to adjust for variable fungal concentrations in samples. Plates were then incubated at

50°C for 5 days to select for *A. fumigatus,* which is unique among *Aspergillus* spp in its

ability to grow at this temperature. On day 5, the incubator temperature was reduced to

42°C to allow for sporulation and plates grown for another 2 days. The number of *A.*

fumigatus colonies was counted and up to two colonies per soil sample transferred to

new plates for isolation.

 Antifungal susceptibility testing

Quick screening of susceptibility to itraconazole, voriconazole, and posaconazole were

assessed using the agar-based VIPcheck™ Assay (Mediaproducts BV, Groningen,

Netherlands) following the manufacturer's directions. For testing, isolates were grown on

SGA for 2-4 days at 37°C. Plates were then swabbed with a damp, sterile cotton swab

to prepare a conidial suspension of 0.5-2 McFarland. 25 µL of this suspension was then

plated onto the 4 wells of the VIPcheck plate containing 4 mg/L itraconazole, 2 mg/L

voriconazole, 0.5 mg/L posaconazole, or a control well containing no drug. To assess

susceptibility to agricultural azoles during isolates collected during 2017 and 2018,

RPMI+2% glucose agar plates containing either 1 mg/L difenoconazole or 2 mg/L

tebuconazole were prepared as in (42). Resistance was defined as significant inhibition

of germination and hyphal growth compared to the no drug control. Isolates that showed

resistance to any of the medical azoles in the VIPcheck assay were subject to broth

microdilution following EUCAST methodology.

DNA extraction

For WGS and PCR-based amplification, isolates were grown shaking in SG broth at

37°C and genomic DNA isolated using the Quick-DNA Fungal/Bacterial miniprep kit

(Zymo Research, Ivrine, CA) according to the manufacturer's suggested protocol.

cyp51a genotyping

The *cyp51a* coding sequence and upstream region containing the tandem repeat was

amplified using the primers described in Supplemental Table 4. Cleaned up PCR

products were sequenced and the *cyp51a* genotype determined using FunResDB (https://elbe.hki-jena.de/FunResDb/index.php).

Genome sequencing, quality assessment, and alignment

Library preparation and 2x150bp paired-end sequencing was performed on a NextSeq 500 v2 by LGC Genomics (Berlin, Germany) following the manufacturer's recommended protocols. Sequence data quality control and filtering were performed using an in-house script and FastQC (v0.11.5). Quality reads were mapped to the *A. fumigatus* Af293 reference genome (version 2015-09-27; retrieved from FungiDB (43)) using BWA-MEM (version 0.7.8-r779-dirty) (44). PCR duplicates were marked using MarkDuplicate from Picard version 2.18.25 embeded in the Genome Analysis Toolkit (GATK; version 4.1.0.0). All WGS samples included for analysis possessed greater than 10 fold genome coverage after mapping and more than 90% of reads mapped to the reference genome.

Variant identification and SNV-based phylogeny

Short variants, including single nucleotide variants (SNVs) and short insertions and deletions (InDels) were detected using GATK Haplotype Caller following their recommended best practices for single calling (45). Copy number variants (CNVs) were identified using Control-FREEC (46). For the phylogenetic analysis, nucleotide consensus sequences were extracted from vcf files using VCFtools (47) and an in-house script used to translate nucleotide to protein-coding sequences. Multiple sequence alignment was performed using MUSCLE v3.8.31 (48) using 7,771 conserved core genes and an approximately-maximum-likelihood phylogeny constructed using

FastTree2 (version 2.1.10) (49). Interactive Tree of Life (iTOL) v4 was used for

visualization (50).

<u>Genetic diversity analyses</u>

Analysis of molecular variance (AMOVA) was determined using the R package ade4

(nrep = 999). Nucleotide diversity (π) was calculated by VCFtools (version 0.1.6) using

5kbp windows with a step size of 500bp. Nucleotide polymorphism (θ) and Tajima's D

were calculated using ANGSD (51) using a window size of 5 kbp and a step size of 500

bp. Weighted Weir and Cockerham's F_{ST} values were calculated using VCFtools, while

V_{ST} values were calculated as in (52). A Mantel test correlating geographic distance

matrices with pairwise FST matrices was performed using the R package ade4 (nrep =

9999).

<u>Estimation of fungicide treatment areas and rates in Germany</u>

The fraction of each district theoretically treated with fungicides was calculated using

publicly reported data from 2016 available from the German Statistical Offices

(statistikportal.de) using the following equation:

535
$$\frac{(Ha_{Ackerland} + Ha_{Dauerkulturen}) \times Fraction_{Conventional}}{Ha_{Total}}$$

The total area per district was obtained from Table 33111-01-02-4: Ground area by

actual use. The sum of arable farmland and permeant crop areas was calculated for

each administrative district using Table 41141-01-01-4: Farms and their agricultural

used area by crop type. To accommodate that some percentage of this area is cultivated

under organic agriculture methods, the hectares of crop land was then multiplied by the

fraction of non-organic agriculture for the federal state in which the district is located to

estimate the number of hectares potentially treated with fungicides (as calculated using

data on "Farms, agricultural areas, and workers"; accessed on November 1, 2019 on

https://www.statistikportal.de/node/254). Unfortunately, no data was available on the

breakdown of agricultural methods at the district level to allow for more exact estimation.

Finally, this value of estimated treated area per district was divided by the total area of

the district for visualization as a choropleth map.

<u>Box and whisker plots</u>

Box and whisker plots presented in this paper are in the style of Tukey, where the bold

line indicates the 50th percentile, and the hinges represent the 25th and 75th percentiles.

The lower whisker extends from the lower hinge to the lowest datum within 1.5

interquartile range (IQR), while the upper whisker represents the highest datum still

within 1.5 IQR. Outliers are marked with points.

<u>Data and isolate availability</u>

Isolates generated within this study were submitted to and are publicly available in the

Jena Microbial Resource Collection. Raw FASTQ files were uploaded to the NCBI

Sequence Read Archive and are publicly available under BioProject XXXXXXX.

ACKNOWLEDGEMENTS

The authors thank Ivo Schliebner for helpful scientific discussions and his input

establishing the network of farms sampled in this study, Anja Kunert for scientific advice,

and Matthew Blango for thoughtful comments and discussions regarding this

manuscript. We also acknowledge Antonia Lange for her experimental support in the isolation of environmental strains from 2016. G.P. would like to thank the Deutsche Forschungsgemeinschaft (DFG) CRC/Transregio 124 "Pathogenic fungi and their human host: Networks of interaction", subproject INF & B5.

AUTHOR CONTRIBUTIONS

O.K. and H.B.D. conceived the study. A.E.B. and J.B. collected samples and performed experiments. W.B. provided reagents and experimental advice. A.E.B., J.B., T.S.E., K.K, G.P., H.B.D., and O.K. analyzed the data and interpreted the results. A.E.B. wrote the manuscript with input from all authors.

FUNDING

This project was supported by The Federal Ministry for Education and Science (Bundesministerium für Bildung und Forschung) within the framework of InfectControl 2020 (Projects FINAR and FINAR 2.0, grants 03ZZ0809 and 03ZZ0834). The work of the German National Reference Center for Invasive Fungal Infections is supported by the Robert Koch Institute from funds provided by the German Ministry of Health (grant 1369-240).

REFERENCES

1. Leading International Fungal Education (LIFE): Invasive Aspergillosis. http://www.life-worldwide.org/fungal-diseases/invasive-aspergillosis.

2. Patterson TF, Iii RT, Denning DW, Fishman JA, Hadley S, Herbrecht R, Kontoyiannis DP, Marr KA, Morrison VA, Nguyen MH, Segal BH, Steinbach WJ, Stevens DA, Walsh TJ, Wingard JR. 2016. Practice Guidelines for the Diagnosis and Management of Aspergillosis : 2016 Update by the Infectious Diseases Society of America. Clinical Infectious Diseases doi:10.1093/cid/ciw326:1-60.

3. Buil JB, Snelders E, Denardi LB, Melchers WJG, Verweij PE. 2019. Trends in Azole Resistance in Aspergillus fumigatus, the Netherlands, 1994-2016. Emerg Infect Dis 25:176-178.

4. Steinmann J, Hamprecht A, Vehreschild MJ, Cornely OA, Buchheidt D, Spiess B, Koldehoff M, Buer J, Meis JF, Rath PM. 2015. Emergence of azole-resistant invasive aspergillosis in HSCT recipients in Germany. J Antimicrob Chemother 70:1522-6.

5. Chowdhary A, Kathuria S, Xu J, Meis JF. 2013. Emergence of Azole-Resistant Aspergillus fumigatus Strains due to Agricultural Azole Use Creates an Increasing Threat to Human Health. PLoS Pathogens 9:3-7.

6. Chowdhary A, Kathuria S, Xu J, Sharma C, Sundar G, Singh PK, Gaur SN, Hagen F, Klaassen CH, Meis JF. 2012. Clonal expansion and emergence of environmental multiple-triazole-resistant Aspergillus fumigatus strains carrying the TR(3)(4)/L98H mutations in the cyp51A gene in India. PLoS One 7:e52871.

7. Meis JF, Chowdhary A, Rhodes JL, Fisher MC, Verweij PE. 2016. Clinical implications of globally emerging azole resistance in Aspergillus fumigatus. Philos Trans R Soc Lond B Biol Sci 371.

8. Ballard E, Melchers WJG, Zoll J, Brown AJP, Verweij PE, Warris A. 2018. In-host microevolution of Aspergillus fumigatus: A phenotypic and genotypic analysis. Fungal Genet Biol 113:1-13.

9. Hare RK, Gertsen JB, Astvad KMT, Degn KB, Lokke A, Stegger M, Andersen PS, Kristensen L, Arendrup MC. 2019. In Vivo Selection of a Unique Tandem Repeat Mediated Azole Resistance Mechanism (TR120) in Aspergillus fumigatus cyp51A, Denmark. Emerg Infect Dis 25:577-580.

10. Berger S, El Chazli Y, Babu AF, Coste AT. 2017. Azole Resistance in Aspergillus fumigatus: A Consequence of Antifungal Use in Agriculture? Front Microbiol 8:1024.

11. Verweij PE, Snelders E, Kema GHJ, Mellado E, Melchers WJG. 2009. Azole resistance in Aspergillus fumigatus: a side-effect of environmental fungicide use? The Lancet Infectious Diseases 9:789-795.

12. Morton V, Staub T. 2008. A Short History of Fungicides. American Phytopathological Society Net Online Feature doi:American Phytopathological Society.

13. Maertens JA. 2004. History of the development of azole derivatives. Clin Microbiol Infect 10 Suppl 1:1-10.

14. Price CL, Parker JE, Warrilow AG, Kelly DE, Kelly SL. 2015. Azole fungicides - understanding resistance mechanisms in agricultural fungal pathogens. Pest Manag Sci 71:1054-8.

15. Anonymous. 2019. FRAC Code List 2019: Fungal control agents sorted by cross resistance pattern and mode of action. Fungicide Resistance Action Committee.

16. Fisher MC, Hawkins NJ, Sanglard D, Gurr SJ. 2018. Worldwide emergence of resistance to antifungal drugs challenges human health and food security. Science 360:739-742.

17. Mellado E, Garcia-Effron G, Alcazar-Fuoli L, Melchers WJ, Verweij PE, Cuenca-Estrella M, Rodriguez-Tudela JL. 2007. A new Aspergillus fumigatus resistance mechanism conferring in vitro cross-resistance to azole antifungals involves a combination of cyp51A alterations. Antimicrob Agents Chemother 51:1897-904.

18.   Liu M, Zheng N, Li D, Zheng H, Zhang L, Ge H, Liu W. 2016. cyp51A-based mechanism
of azole resistance in Aspergillus fumigatus: Illustration by a new 3D Structural Model of
Aspergillus fumigatus CYP51A protein. Med Mycol 54:400-8.
19.   Nash A, Rhodes J. 2018. Simulations of CYP51A from Aspergillus fumigatus in a model
bilayer provide insights into triazole drug resistance. Med Mycol 56:361-373.
20.   Zhang J, Snelders E, Zwaan BJ, Schoustra SE, Meis JF, van Dijk K, Hagen F, van der
Beek MT, Kampinga GA, Zoll J, Melchers WJG, Verweij PE, Debets AJM. 2017. A Novel
Environmental Azole Resistance Mutation in Aspergillus fumigatus and a Possible Role
of Sexual Reproduction in Its Emergence. MBio 8.
21.   Vermeulen E, Maertens J, Schoemans H, Lagrou K. 2012. Azole-resistant Aspergillus
fumigatus due to TR46/Y121F/T289A mutation emerging in Belgium, July 2012. Euro
Surveill 17.
22.   Arendrup MC, Verweij PE, Mouton JW, Lagrou K, Meletiadis J. 2017. Multicentre
validation of 4-well azole agar plates as a screening method for detection of clinically
relevant azole-resistant Aspergillus fumigatus. J Antimicrob Chemother 72:3325-3333.
23.   Buil JB, van der Lee HAL, Rijs A, Zoll J, Hovestadt J, Melchers WJG, Verweij PE. 2017.
Single-Center Evaluation of an Agar-Based Screening for Azole Resistance in
Aspergillus fumigatus by Using VIPcheck. Antimicrob Agents Chemother 61.
24.   Garcia-Rubio R, Monzon S, Alcazar-Fuoli L, Cuesta I, Mellado E. 2018. Genome-Wide
Comparative Analysis of Aspergillus fumigatus Strains: The Reference Genome as a
Matter of Concern. Genes (Basel) 9.
25.   Bowyer P, Denning DW. 2014. Environmental fungicides and triazole resistance in
Aspergillus. Pest Management Science 70:173-178.
26.   Rhodes J. 2019. Rapid Worldwide Emergence of Pathogenic Fungi. Cell Host Microbe
26:12-14.
27.   Resendiz Sharpe A, Lagrou K, Meis JF, Chowdhary A, Lockhart SR, Verweij PE, group
IEARSw. 2018. Triazole resistance surveillance in Aspergillus fumigatus. Med Mycol
56:83-92.
28.   Bader O, Weig M, Reichard U, Lugert R, Kuhns M, Christner M, Held J, Peter S,
Schumacher U, Buchheidt D, Tintelnot K, Gross U, MykoLabNet DP. 2013. cyp51A-
Based mechanisms of Aspergillus fumigatus azole drug resistance present in clinical
samples from Germany. Antimicrob Agents Chemother 57:3513-7.
29.   Bader O, Tünnermann J, Dudakova A, Tangwattanachuleeporn M, Weig M, Groß U,
Hoberg N, Geibel S, Vogel E, Büntzel J, Springer J, Lehning LY, Schädel C, Antweiler E,
Metzger L, Zautner A, Buchheidt D, Spiess B, Hamprecht A, Steinmann J, Rößler S,
Wiegmann S, Klingebiel S, Loock AC, Hegewald J, Hassenpflug M, Aurin A, Szymczak J,
Diffloth N, Kuhns M. 2015. Environmental isolates of azole-resistant Aspergillus
fumigatus in Germany. Antimicrobial Agents and Chemotherapy 59:4356-4359.
30.   Alvarez-Moreno C, Lavergne RA, Hagen F, Morio F, Meis JF, Le Pape P. 2019.
Fungicide-driven alterations in azole-resistant Aspergillus fumigatus are related to
vegetable crops in Colombia, South America. Mycologia 111:217-224.
31.   Prigitano A, Venier V, Cogliati M, De Lorenzis G, Esposto MC, Tortorano AM. 2014.
Azole-resistant aspergillus fumigatus in the environment of Northern Italy, May 2011 to
June 2012. Eurosurveillance 19:1-7.
32.   Mortensen KL, Mellado E, Lass-Fl??rl C, Rodriguez-Tudela JL, Johansen HK, Arendrup
MC. 2010. Environmental study of azole-resistant Aspergillus fumigatus and other
aspergilli in Austria, Denmark, and Spain. Antimicrobial Agents and Chemotherapy
54:4545-4549.
33.   Sewell TR, Zhang Y, Brackin AP, Shelton JMG, Rhodes J, Fisher MC. 2019. Elevated
prevalence of azole resistant Aspergillus fumigatus in urban versus rural environments in
the United Kingdom. Antimicrob Agents Chemother doi:10.1128/AAC.00548-19.

34. Jeanvoine A, Rocchi S, Reboux G, Crini N, Crini G, Millon L. 2017. Azole-resistant
Aspergillus fumigatus in sawmills of Eastern France. J Appl Microbiol
doi:10.1111/jam.13488.
35. Karaoglanidis GS, Thanassoulopoulos CC. 2002. Phenotypic Instability of Cercospora
beticola Sacc. Strains Expressing Resistance to the Sterol Demethylation-Inhibiting
(DMI) Fungicide Flutriafol after Cold Exposure. Journal of Phytopathology 150:692-696.
36. Cox KD, Bryson PK, Schnabel G. 2007. Instability of Propiconazole Resistance and
Fitness in Monilinia fructicola. Phytopathology 97:448-53.
37. Calo S, Shertz-Wall C, Lee SC, Bastidas RJ, Nicolas FE, Granek JA, Mieczkowski P,
Torres-Martinez S, Ruiz-Vazquez RM, Cardenas ME, Heitman J. 2014. Antifungal drug
resistance evoked via RNAi-dependent epimutations. Nature 513:555-8.
38. Schoustra SE, Debets AJM, Rijs A, Zhang J, Snelders E, Leendertse PC, Melchers
WJG, Rietveld AG, Zwaan BJ, Verweij PE. 2019. Environmental Hotspots for Azole
Resistance Selection of Aspergillus fumigatus, the Netherlands. Emerg Infect Dis
25:1347-1353.
39. Abdolrasouli A, Rhodes J, Beale MA, Hagen F, Rogers TR, Chowdhary A, Meis JF,
Armstrong-James D, Fisher MC. 2015. Genomic Context of Azole Resistance Mutations
in Aspergillus fumigatus Determined Using Whole-Genome Sequencing. MBio 6:e00536.
40. Dunne K, Hagen F, Pomeroy N, Meis JF, Rogers TR. 2017. Intercountry Transfer of
Triazole-Resistant Aspergillus fumigatus on Plant Bulbs. Clin Infect Dis 65:147-149.
41. Alvarez-Moreno C, Lavergne RA, Hagen F, Morio F, Meis JF, Le Pape P. 2017. Azole-
resistant Aspergillus fumigatus harboring TR34/L98H, TR46/Y121F/T289A and TR53
mutations related to flower fields in Colombia. Sci Rep 7:45631.
42. Guinea J, Verweij PE, Meletiadis J, Mouton JW, Barchiesi F, Arendrup MC,
Subcommittee on Antifungal Susceptibility Testing of the EECfAST. 2018. How to:
EUCAST recommendations on the screening procedure E.Def 10.1 for the detection of
azole resistance in Aspergillus fumigatus isolates using four-well azole-containing agar
plates. Clin Microbiol Infect doi:10.1016/j.cmi.2018.09.008.
43. Basenko EY, Pulman JA, Shanmugasundram A, Harb OS, Crouch K, Starns D,
Warrenfeltz S, Aurrecoechea C, Stoeckert CJ, Jr., Kissinger JC, Roos DS, Hertz-Fowler
C. 2018. FungiDB: An Integrated Bioinformatic Resource for Fungi and Oomycetes. J
Fungi (Basel) 4.
44. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics 25:1754-60.
45. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K,
Altshuler D, Gabriel S, Daly M, DePristo MA. 2010. The Genome Analysis Toolkit: a
MapReduce framework for analyzing next-generation DNA sequencing data. Genome
Res 20:1297-303.
46. Boeva V, Popova T, Bleakley K, Chiche P, Cappo J, Schleiermacher G, Janoueix-
Lerosey I, Delattre O, Barillot E. 2012. Control-FREEC: a tool for assessing copy number
and allelic content using next-generation sequencing data. Bioinformatics 28:423-5.
47. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE,
Lunter G, Marth GT, Sherry ST, McVean G, Durbin R, Genomes Project Analysis G.
2011. The variant call format and VCFtools. Bioinformatics 27:2156-8.
48. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res 32:1792-7.
49. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood
trees for large alignments. PLoS One 5:e9490.
50. Letunic I, Bork P. 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. Nucleic Acids Res 47:W256-W259.

51. Korneliussen TS, Moltke I, Albrechtsen A, Nielsen R. 2013. Calculation of Tajima's D and
other neutrality test statistics from low depth next-generation sequencing data. BMC
Bioinformatics 14:289.
52. Redon R, Ishikawa S, Fitch KR, Feuk L, Perry GH, Andrews TD, Fiegler H, Shapero MH,
Carson AR, Chen W, Cho EK, Dallaire S, Freeman JL, Gonzalez JR, Gratacos M, Huang
J, Kalaitzopoulos D, Komura D, MacDonald JR, Marshall CR, Mei R, Montgomery L,
Nishimura K, Okamura K, Shen F, Somerville MJ, Tchinda J, Valsesia A, Woodwark C,
Yang F, Zhang J, Zerjal T, Zhang J, Armengol L, Conrad DF, Estivill X, Tyler-Smith C,
Carter NP, Aburatani H, Lee C, Jones KW, Scherer SW, Hurles ME. 2006. Global
variation in copy number in the human genome. Nature 444:444-54.

Table 1. Analysis of molecular variance (AMOVA) between farms. Calculated using
isolates from before and after azole application. n = 16 isolates per farm.

	Total variation	Variation within samples (%)	p-value	Variation between samples (%)	p-value	Variation between farms (%)	p-value
All farms	289735.7	274622.7 (94.8)	<0.001	14629 (5)	<0.001	484 (0.2)	0.303
B vs. C	147358.0	139257.7 (94.5)	<0.001	6386.2 (4.3)	<0.001	1714.1 (1.2)	0.028
B vs. D	190425.2	181470 (95.3)	<0.001	9002.4 (4.7)	<0.001	-47.2 (0.0)	0.429
B vs. E	162874.7	155745.5 (95.6)	<0.001	6879.3 (4.2)	<0.001	249.9 (0.2)	0.211
C vs. D	130625.7	122692.8 (93.9)	<0.001	7932 (6.1)	<0.001	1 (0.0)	0.355
C vs. E	105529.1	99521.4 (94.3)	<0.001	6098.3 (5.8)	<0.001	-90.6 (-0.1)	0.414
D vs. E	143684.1	135847.0 (94.5)	<0.001	8382.7 (5.8)	<0.001	-545.6 (0.4)	0.890

752 **Table 2.** Analysis of molecular variance between isolates collected before and after
753 azole exposure. n = 16 isolates per farm field (eight pre-azole and eight post-azole).

	Total variation	Variation within samples (%)	p-value	Variation between samples (%)	p-value	Variation between time periods (%)	p-value
Farm B	106712.9	103295.3 (96.8)	<0.001	3729.5 (3.5)	<0.001	-311.9 (-0.3)	0.617
Farm C	48438.0	45393.1 (93.7)	<0.001	3341.7 (6.9)	<0.001	-296.8 (-0.6)	0.593
Farm D	85724.6	80522.3 (93.9)	<0.001	5592 (6.5)	<0.001	-389.7 (-0.5)	0.564
Farm E	59366.5	55941.4 (94.2)	<0.001	3001.2 (5.1)	<0.001	423.9 (0.7)	0.149

Figure 1. Abundance of *A. fumigatus* in the soil of conventional and organic farms.
(A) CFU/g *A. fumigatus* at varying soil depths. n = 10 samples per depth. (B) Estimated fungicide treatment rates and areas in Germany. The fraction of each district that is theoretically treated with fungicides was calculated using land use and organic agriculture share data reported by the Statistical Office of Germany in December 2016. Districts where no data on land use was available are shaded grey. (C-D) Abundance of A. fumigatus in the spring as measured by CFU/g soil in the spring of 2017 (C) and 2018 (D). For 2017, boxplots represent n = 50 soil samples per field for farms A, B, C, E, F, H, K and n = 25 for farms D, G, L. For 2018, n = 50 soil samples per field for farms A, B, C, D, E, G, H, K and n = 25 for farms F and L. (E) Comparison of the mean CFU/g soil on farms between 2017 and 2018.

Figure 2. Abundance of *A. fumigatus* in the soil of conventional farms before and after the vegetative period and azole application.

(A, B) CFU/g *A. fumigatus* on conventional fields sampled in April, prior to the vegetative period and azole application (orange), and in July, after the vegetative period and three months of azole application (gray) in 2017 (A) and 2018 (B). *, $P \leq 0.05$; ** $P \leq 0.01$; *** $P \leq 0.001$ as determined by Mann-Whitney U test. (A) Boxplots represent n = 50 samples per field and time point for farms A, B, C, E, H and n = 25 samples for farms D and L. (B) n = 50 samples per field and time point for farms A, B, C, D, E, H and n = 25 samples for farm L. (C-D) CFU/g A. fumigatus during the months of April, May, June, July, October and November of a conventional field applying azoles from May-July (D) and an organic field not applying azole fungicides (C). Bars represent mean ± SEM of 50 soil samples per month. No significant difference was found in abundance between the months of April, May, June, and July for the organic field using a Kruskal-Wallis test. In contrast, we found a significant difference (P = 0.004) for the abundances in this time period on the conventional field. Pairwise Wilcoxon signed-rank test confirmed a significant difference between April and all subsequent months until the cessation of azoles.

Figure 3. Azole resistance among agricultural *A. fumigatus*.
(A-B) Fraction of the ≈ 20 isolates tested per field that grow at 1 mg/L difenoconazole (A) and 2 mg/L tebuconazole (B). For 2017, n = 340 isolates from 9 conventional and 8 organic fields. For 2018, n = 213 isolates from 8 conventional and 7 organic fields (full summary in Supplemental Tables 2 & 3). (C-D) Comparison of the proportion of isolates that grow at the 1 mg/L difenoconazole (C) and 2 mg/L tebuconazole (D) before and after the vegetative period and azole application. For 2017, n = 275 isolates from 7 fields and for 2018, n = 261 isolates from 8 fields; n ≈ 20 isolates per field tested (full summary in Supplemental Tables 4 & 5). Positive values indicate an increased proportion for after compared to before; negative values indicate a decreased proportion for after compared to before. P values calculated by Wilcoxon signed-rank test between before and after values for each field. (E) Local resistance/tolerance in agricultural *A. fumigatus* populations shifts in response to azole application. Fraction of isolates that can grow at 1 mg/L difenoconazole (top) and 2 mg/L tebuconazole (bottom). n = 12-20 isolates/field and time point. (F) Medical azole resistance rates among agriculturally-isolated *A. fumigatus* isolates. Isolates were screened for potential azole resistance using the VIPcheck agar-based screening. Resistance was confirmed and MICs determined via EUCAST broth microdilution testing. (G) cyp51a genotypes of isolates resistant to one or more medical azole. (H) MIC of agricultural *A. fumigatus* isolated before and after azole exposure from 2 fields (n = 20 before and n = 20 after from each field, n = 79 in total).

Figure 4. Phylogeny of agricultural _A. fumigatus_ isolates from before and after the vegetative period and azole application.

From top to bottom, the colored bars indicate: 1) the farm where the isolate was collected, 2) the collection period, 3) voriconazole resistance (susceptible, intermediate, or resistant, according to EUCAST definitions), 4) posaconazole resistance, 5) itraconazole resistance, and 6) the presence of the cyp51a TR34/L98H allele. _A. fischerei_ is indicated as an out group and the two _A. fumigatus_ reference strains, Af293 and A1163 (CEA10), are also marked. Branches with support values less than 0.9 are marked in red.

Nucleotide Diversity

A

Nucleotide Polymorphism

B

Tajima's D

C

Figure 5. Genetic diversity among isolates from before and after the vegetative period and azole exposure from four farms.

(A-C) Nucleotide diversity (π) (A), nucleotide polymorphism (Watterson estimator or θ) (B), and Tajima's D (C) along 5 kb windows before and after the vegetative period and azole exposure. n = 8 isolates per farm and time point, 64 in total.

Supplemental Figure 1. *A. fumigatus* **abundance does not correlate with total organic carbon content of soil.** n = 170 soil samples (10 per field from 17 fields on 10 different farms).

A
2017

B
2018

Supplemental Figure 2. Schematic representation of the sampling time points and fungicide applications.

(A, B) Schematic representation of the sampling time points and fungicide applications for the fields sampled in 2017 (A) and 2018 (B). Soil sampling time points are marked in dark grey, while fungicide applications are noted in a combination of red and light grey indicating the relative proportion of azoles within the fungicide cocktail applied. Fungicide history was not available for fields on Farms H and L.

A

B

C

D

Supplemental Figure 3. Population genetic of farms using isolates from before and after azole exposure.

n = 16 isolates per farm. (A-B) Heat map representing weighted Weir and Cockerham F_{ST} values (A) and V_{ST} values (B) between farms. Blue colors represent more similarity between populations, while red represents greater population differentiation. (C-D) Comparison of nucleotide diversity (π) (C) and nucleotide polymorphism (Watterson estimator or θ) (D) between farms. Values calculated from 5 kb sliding windows.

SUPPLEMENTAL MATERIALS

SUPPLEMENTAL MATERIALS & METHODS

Additional information on field samling and the naming of fields

Six of the sampled farms practiced biological (organic) measures of land management without azole application, while seven practiced integrated pest management, including application of azole fungicides. To differentiate the year the sampling occurred, a corresponding suffix was added to the Field ID (*e.g.* D-2-17 identifies the second field sampled on farm D in the year 2017). Due to crop rotation, the same field could not be surveyed over subsequent years, nor was it possible to always sample the same number of fields from each site each year. To avoid confusion, when we returned to farms the following year, fields were assigned new numbers starting with where the labeling system left off the previous year. (*e.g.* on farm D, two field were sampled in 2017, so in 2018 the labeling system started with D-3). Fields that were sampled over subsequent years, which was only possible for apple orchards, maintained the same numbering system (*e.g.* L-4-17 and L-4-18 are the same fields). In the case where two fields of the same agricultural type on the same farm were sampled, the number of soil samples was divided between the two fields and 25 samples per field were collected.

Assessment of TOC content

Soil samples were air dried and sieved through a 2 mm stainless steel mesh to remove larger organic particles and stones. Samples were then manually ground to a fine powder and divided into two parts. The first fraction (approximately 1000 mg) was used to measure total carbon (TC) content by total combustion with oxygen at 1000°C using an elemental macro-analyzer (Elementar Vario Max Cube, Langenselbold, Germany). The second fraction (appr. 200 mg) was used to determine the total inorganic fraction (TIC) present by volumetrically

adding hydrochloric acid and recording the volume of CO_2 produced (also using an Elementar Vario Max Cube, Langenselbold, Germany). The final total organic carbon (TOC) content in the soil was calculated by subtracting the total carbon (TC) from the total inorganic carbon (TIC).

Supplemental Table 1. Summary of the agricultural sites surveyed 2016- 2018.

Farm	Type of Agriculture	Crop	Soil samples (n)			*A. fumigatus* isolates (n)		
			2016	2017	2018	2016	2017	2018
A	Conventional and organic	Cereal	150	150	150	232	94	81
B	Conventional	Cereal	100	100	100	114	266	335
C	Conventional	Cereal	100	100	100	29	123	37
D	Conventional	Cereal	100	100	100	65	228	287
E	Conventional	Cereal	100	100	100	61	131	165
F	Organic	Cereal	50	50	25	73	87	15
G	Organic	Cereal	50	50	50	29	87	13
H	Conventional and organic	Apple	100	150	150	158	301	421
K	Organic	Cereal	50	50	50	45	26	6
L	Conventional and organic	Apple	100	150	150	91	184	162
		Total	900	1000	975	897	1527	1522

Supplemental Table 2. Resistance summary of the fields sampled during the spring of 2017 (A) and 2018 (B).

(A)

Type of agriculture	Field	Crop	n tested	DIF (1 mg/L)	TEB (2 mg/L)	ITR (4 mg/L)	VOR (2 mg/L)	POS (0.5 mg/L)
				Proportion of isolates that grow at the concentration indicated				
Conventional	A-1-17	Cereal	20	0.15	0.00	0.00	0.00	0.00
	B-1-17	Cereal	20	0.25	0.05	0.00	0.00	0.00
	C-1-17	Cereal	20	0.10	0.00	0.00	0.00	0.00
	D-1-17	Cereal	20	0.25	0.00	0.00	0.00	0.00
	D-2-17	Cereal	20	0.40	0.05	0.00	0.00	0.00
	E-1-17	Cereal	20	0.15	0.05	0.00	0.00	0.00
	H-1-17	Apple	20	0.15	0.05	0.00	0.00	0.00
	L-1-17	Apple	20	0.15	0.00	0.00	0.00	0.00
	L-4-17	Apple	20	0.65	0.00	0.00	0.00	0.00
Organic	A-2-17	Cereal	20	0.35	0.15	0.05	0.00	0.00
	F-1-17	Cereal	20	0.25	0.00	0.00	0.00	0.00
	G-1-17	Cereal	20	0.25	0.10	0.05	0.00	0.00
	G-2-17	Cereal	20	0.15	0.00	0.00	0.00	0.00
	K-1-17	Cereal	20	0.25	0.00	0.00	0.00	0.00
	H-2-17	Apple	20	0.25	0.00	0.00	0.00	0.00
	L-2-17	Apple	20	0.10	0.05	0.00	0.00	0.00
	L-3-17	Apple	20	0.35	0.25	0.00	0.00	0.00

(B)

Type of agriculture	Field	Crop	n tested	DIF (1 mg/L)	TEB (2 mg/L)	ITR (4 mg/L)	VOR (2 mg/L)	POS (0.5 mg/L)
				Proportion of isolates that grow at the concentration indicated				
Conventional	A-3-18	Cereal	11	0.27	0.00	0.00	0.00	0.00
	B-2-18	Cereal	20	0.10	0.05	0.00	0.00	0.00
	C-2-18	Cereal	11	0.09	0.00	0.00	0.00	0.00
	D-3-18	Cereal	20	0.10	0.05	0.00	0.00	0.00
	E-2-18	Cereal	20	0.10	0.10	0.00	0.00	0.00
	H-1-18	Apple	20	0.20	0.10	0.00	0.00	0.00
	L-1-18	Apple	18	0.11	0.00	0.00	0.00	0.00
	L-4-18	Apple	12	0.17	0.00	0.00	0.00	0.00
Organic	A-4-18	Cereal	20	0.40	0.20	0.00	0.00	0.00
	F-2-18	Cereal	10	0.40	0.00	0.00	0.00	0.00
	G-3-18	Cereal	12	0.50	0.00	0.00	0.00	0.00
	K-2-18	Cereal	5	0.00	0.00	0.00	0.00	0.00
	H-2-18	Apple	20	0.20	0.05	0.00	0.00	0.00
	L-2-18	Apple	6	0.17	0.00	0.00	0.00	0.00
	L-3-18	Apple	8	0.38	0.00	0.00	0.00	0.00

Supplemental Table 3. Resistance summary of the fields sampled before and after the vegetative period and azole application in 2017 (A) and 2018 (B).

(A)

Field	Crop	Time Period	n tested	DIF (1 mg/L)	TEB (2 mg/L)	ITR (4 mg/L)	VOR (2 mg/L)	POS (0.5 mg/L)
				Proportion of isolates that grow at the concentration indicated				
B-1-17	Cereal	Before	20	0.25	0.05	0.00	0.00	0.00
		After	20	0.50	0.15	0.00	0.00	0.00
C-1-17	Cereal	Before	20	0.10	0.00	0.00	0.00	0.00
		After	20	0.45	0.15	0.10	0.00	0.00
D-1-17	Cereal	Before	20	0.25	0.00	0.00	0.00	0.00
		After	20	0.50	0.25	0.00	0.00	0.00
D-2-17	Cereal	Before	20	0.40	0.05	0.00	0.00	0.00
		After	20	0.35	0.05	0.00	0.00	0.00
E-1-17	Cereal	Before	20	0.15	0.05	0.00	0.00	0.00
		After	15	0.47	0.07	0.00	0.00	0.00
H-1-17	Apple	Before	20	0.15	0.05	0.00	0.00	0.00
		After	20	0.55	0.20	0.00	0.00	0.00
L-4-17	Apple	Before	20	0.65	0.00	0.00	0.00	0.00
		After	20	0.50	0.10	0.00	0.00	0.00

(B)

Field	Crop	Time Period	n tested	DIF (1 mg/L)	TEB (2 mg/L)	ITR (4 mg/L)	VOR (2 mg/L)	POS (0.5 mg/L)
				Proportion of isolates that grow at the concentration indicated				
A-3-18	Cereal	Before	11	0.27	0.00	0.00	0.00	0.00
		After	11	0.27	0.18	0.00	0.00	0.00
B-2-18	Cereal	Before	20	0.10	0.05	0.00	0.00	0.00
		After	20	0.25	0.10	0.05	0.00	0.00
C-2-18	Cereal	Before	11	0.09	0.00	0.00	0.00	0.00
		After	16	0.25	0.06	0.00	0.00	0.00
D-3-18	Cereal	Before	20	0.10	0.05	0.00	0.00	0.00
		After	20	0.35	0.10	0.05	0.00	0.00
H-1-18	Apple	Before	20	0.20	0.10	0.00	0.00	0.00
		After	20	0.50	0.10	0.00	0.00	0.00
L-1-18	Apple	Before	18	0.11	0.00	0.00	0.00	0.00
		After	10	0.50	0.00	0.00	0.00	0.00
L-4-18	Apple	Before	12	0.17	0.00	0.00	0.00	0.00
		After	12	0.58	0.08	0.00	0.00	0.00

Supplemental Table 4. Primers used for amplifying and sequencing *cyp51a*.

Primer	Sequence	Use
cyp51a_F	5'-CGTAGCAAGGGAGAAGGAAA	Amplification and sequencing
cyp51a_R	5'-CACCTATTCCGATCACACCA	Amplification and sequencing
cyp51a_internal	5'-ATGTCAATGCGGAAGAGGTC	Sequencing