Metformin increases tauroursodeoxycholic acid levels to improve insulin

resistance in diet-induced obese mice

Ya Zhang [a,b,1], Yang Cheng [a,1], Jian Liu [a,1], Dan He [a], Jihui Zuo [a], Liping Yan [a],

Ronald W. Thring [b], Mingjiang Wu [a], Yitian Gao [a,*], Haibin Tong [a,*]

[a] College of Life and Environmental Science, Wenzhou University, Wenzhou 325035,

China;

[b] Environmental Science and Environmental Engineering, University of Northern

British Columbia, Prince George, BC, Canada;

*Correspondence: tonghaibin@gmail.com (H.T.); gyt@wzu.edu.cn (Y.G.); Tel.: +

86-577-86689078

[1] These authors contributed equally to this study.

Abstract

Metformin is widely used to surmount insulin resistance (IR) and type 2 diabetes. Evidence indicates that metformin improves insulin resistance associated with gut microbiota, but the underlying mechanism remains unclear. In the present study, metformin effectively improved insulin sensitivity and alleviated liver inflammation and oxidative stress in high-fat diet (HFD)-fed mice. Metabolomics analysis showed that metformin increased tauroursodeoxycholic acid (TUDCA) levels both in intestinal content and liver by reducing the production and activity of bile salt hydrolase (BSH). We further found that TUDCA was able to antagonize with KEAP1 to prevent its binding to Nrf2 and activate Nrf2/ARE pathway, thereby reducing intracellular ROS and improving insulin signaling. Moreover, metformin increased the proportion of *Akkermanisia muciniphlia* in the HFD-fed mice, while *in vitro* growth curve test confirmed that it's TUDCA, not metformin, promoted the proliferation of *A. muciniphlia*. Subsequently, TUDCA administration could effectively ameliorate insulin resistance, activate hepatic Nrf2/ARE pathways, and increase the abundance of intestinal *A. muciniphlia* in *ob/ob* mice. These findings reveal that metformin remodels the gut microbiota, reduces oxidative stress and enhances insulin sensitivity partly due to increasing the production of TUDCA. This provides a novel mechanism by which metformin alleviates diet-induced insulin resistance and improves metabolism.

Keywords: Metformin; Gut microbiota; Insulin resistance; TUDCA; Nrf2

Abbreviations

T2DM, type 2 diabetes mellitus; AMPK, AMP-activated protein kinase; IR, insulin

resistance; BA, bile acids; ER, endoplasmic reticulum; BSH, bile salt hydrolase;

NASH, nonalcoholic steatohepatitis; ROS, reactive oxygen species; Nrf2, nuclear

factor-erythroid-2-related factor 2; KEAP1, Kelch-like ECH-associated protein 1;

TUDCA, tauroursodeoxycholic acid; IRS-1, insulin receptor substrate 1; Akt, protein

kinase B; 4-HNE, 4-hydroxynonenal; IPGTT, intraperitoneal glucose tolerance test;

IPITT, intraperitoneal insulin tolerance test; FFA, free fatty acids; FBG, fasting blood

glucose; FINS, fasting serum insulin; HOMA-IR, homeostasis model

assessment-estimated insulin resistance; 2-NBDG,

2-deoxy-2-[(7-nitro-2,1,3-benzoxadiazol-4-yl) amino]-D-glucose; NQO1,

NAD(P)H:quinone oxidoreductase-1; HO-1, heme oxygenase 1; IL-1β, interleukin-1

β; Il-6, interleukin-6; TNF-α, tumor necrosis factor-α; CYP7a1, cholesterol

7α-hydroxylase; BAAT, bile acid-CoA:amino acid N-acyltransferase; 7α/β-HSDH,

7α/β-hydroxysteroid dehydrogenase.

Introduction

The prevalence of obesity and overweight have been on the rise worldwide[1]. Current predictions indicate that more than 1 billion people will be obese by 2030[2]. Visceral obesity precedes the development of metabolic complications and chronic diseases such as type 2 diabetes mellitus (T2DM)[3], hepatic steatosis[4] and cardiovascular diseases[5]. Metformin has been the first-line anti-diabetic drug for more than 60 years because of its excellent hypoglycemic effect and safety profile[1]. Although several studies have explored the mechanisms by which metformin maintains glucose homeostasis, findings from such studies are inconsistent. The prevailing view is that metformin activates AMP-activated protein kinase (AMPK) in the liver, similar to the principle of caloric restriction that relieves symptoms of diabetes[6]. However, metformin is an orally administered drug that reaches high concentrations in the intestine with much lower serum concentrations[7], thus, the possibility that its metabolic therapeutic mechanism might be due in part to actions in the intestine cannot be ignored. Accumulated evidence shows that a high-fat diet (HFD) reduces the abundance of *Lactobacilli* in the small intestine. However, this can be counteracted by metformin, which increases the proportion of probiotics, thus, relieving the symptoms of HFD-induced diabetes[8]. Although metformin may play a hypoglycemic effect by improving the gut microbiota, oral administration of antibiotics to diabetic mice impairs the hypoglycemic ability of metformin[9]. In addition, while metformin can increase the abundance of *Escherichia coli*, it can as

well reduce the abundance of *Intestinibacter*[10]. This can further influence diverse

biological processes such as the synthesis of branched-chain amino acid and the

secretion of GLP-1, thereby improving glucose tolerance and alleviating insulin

resistance (IR)[10,11]. These findings suggest that modulation of gut microbiota is

involved in the hypoglycemic mechanism of metformin.

Bile acids (BAs), which are produced in the liver and serve as important

ingredients of the digestive fluid in the intestine, enter the small intestine through the

biliary system and participate in fat digestion and absorption[12]. Gathered evidence

demonstrates that BAs could improve the intestinal homeostasis by modulating the

gut microbiota, thus relieving metabolic syndrome[13]. Recent studies also suggest that

BAs can participate in the liver-gut axis circulation and the improvement of

HFD-initiated metabolic disorders by inhibiting the proliferation of pathogenic

bacteria[14]. By modulating the gut microbiota, BAs have also been reported to induce

multiple beneficial effects on the intestinal lumen and intestinal walls such as the

suppression of intestinal inflammatory responses[15], the resolution of endoplasmic

reticulum (ER) stress in the intestinal epithelial cells underlying the pathology of

inflammatory bowel disease[16] and the improvement of gut barrier dysfunction[17-19].

These findings suggest that there is a complex relationship among BAs, gut

microbiota and metabolic syndrome.

Previous studies have reported the association of BAs with oxidative stress,

which has been shown to treat metabolic dysfunction by acting as an endogenous

chemical chaperone to protect cells against ER stress[20]. Intestinal micro-environment

(including inflammation status, the function of the epithelial tight junction and gut

microbiota) plays an essential role in the progression of obese-induced IR[21]. The

presence of gut microbiota and BAs in the intestine may be closely associated with

intestinal metabolic state[22,23]. Analyses from metagenomic sequencing and

metabolomics show that metformin treatment increases the levels of the conjugated

bile acid in the gut by decreasing the abundance and bile salt hydrolase (BSH) activity

of *Bifidobacterium* species in the intestines of individuals with T2DM[24,25]. BAs are

signal molecules that control the dynamic balance between energy metabolism and

liver protection. It has been reported that BAs change with obesity, IR[26] and

nonalcoholic steatohepatitis (NASH)[27]. However, the mechanism by which BAs

alleviate HFD-induced IR and the underlying role played by the gut microbiota during

this process are not clear.

Accumulating evidence has clearly indicated that oxidative stress plays a major

role in the pathological process of IR[28]. What's more, mitochondrial dysfunction,

reactive oxygen species (ROS) over production, and lipid peroxidation have been

found in the liver of Zucker rats with T2DM[29]. Increased oxidative stress seems to be

a deleterious factor leading to IR and impaired glucose tolerance in T2DM[30]. Nuclear

factor-erythroid-2-related factor 2 (Nrf2), tightly interacts with Kelch-like

ECH-associated protein 1 (KEAP1), an important transcription factor responsible for

inducing phase II detoxifying and antioxidant enzymes, is a key player in the

antioxidant response and glucose metabolism[31,32]. Recent study has confirmed

KEAP1-Nrf2 system as a critical target for preventing the onset of IR[33], and bile acids

have a great influence on the Nrf2/ARE antioxidant signal pathway[34]. This study

investigated how metformin prevents the development of IR in HFD-induced obese

mice by modulating the dysbiosis of gut microbiota and alteration metabolites.

Exploring the mechanism behind this improves our knowledge of gut microbiota and

metabolic interactions underlying the anti-insulin resistance effects of metformin.

Results

1. Metformin alleviates HFD-induced insulin resistance and oxidative stress

The administration of metformin significantly alleviated diet-induced increase of

body weight (**Fig. 1A**), without affecting food intake (**Fig. S1A**). The reduced liver

index and subcutaneous fat index (**Fig. S1B)** as well as H&E staining of subcutaneous

adipose tissue (**Fig. S1C**) and number of adipocytes (**Fig. S1D)** suggested that

metformin prevented fat deposition in HFD-fed mice. The diet-induced dyslipidemia

was effectively prevented by metformin as indicated by the lower levels of T-CHO,

TG and FFA in both the liver and the serum (**Fig. S1E-G**). Further, metformin

significantly improved HFD-feeding impaired glucose tolerance and insulin resistance

determined by IPGTT (**Fig. 1B**) and IPITT (**Fig. 1C**), respectively. Marked elevation

in fasting serum insulin (FINS) was observed in the HFD group. Notably, metformin

treatment tended to reverse the increase in FINS (**Fig. 1D**) as well as reduce

HOMA-IR (**Fig. 1E**), an indicator for evaluating the level of insulin resistance. In

addition, the level of phosphorylated Akt at Ser473 was significantly reduced in the

liver of the HFD-induced obese mice, but the level was restored in the HFD+Met

group (**Fig. 1F**). Contrary to the phosphorylated Akt, the level of phosphorylated

IRS-1 at Ser307 was significantly elevated in the HFD group, whereas this level was

reduced in the HFD+Met group (**Fig. 1F**). These results indicate that the preventive

effect of metformin on HFD-induced obesity resulted in improved glucose

homeostasis and insulin sensitivity.

Accumulated evidence indicated that oxidative stress is involved in the

development of HFD- induced IR, thus we further measured the oxidative stress

levels after metformin treatment. The results showed that the activity of CAT in the

serum and the liver was markedly lower in the HFD group compared with the NCD

group (**Fig. 2A**), the ratio of GSH/GSSG was also reduced (**Fig. 2B**). After metformin

intervention, the activity of CAT and the ratio of GSH/GSSG were significantly

increased. The accumulation of lipid peroxide MDA in liver, caused by HFD, was

significantly reduced in the metformin group (**Fig. 2C**). Nrf2/ARE is a classic

antioxidant-signaling pathway and KEAP1 is a key protein that mediates the

degradation of Nrf2. HFD feeding significantly inhibited Nrf2/ARE signaling

whereas metformin significantly increased the expression of Nrf2 as well as

phosphorylated Nrf2 (**Fig. 2D**). Nrf2/ARE signaling regulates the expression of a

variety of detoxification and antioxidant enzymes. As shown in **Fig. 2E**, Metformin

significantly up-regulated the mRNA levels of *Nqo1* and *Ho1*, which are essential

genes for balancing oxidative stress. Also, metformin prevented metabolic

inflammation in the liver and ileum, leading to a reduced gene expression of *Il-1β*,

Il-6 and *Tnf-α* in the ileum (**Fig. S2A,B**). Metformin also prevented the release of

endotoxin LPS in serum (**Fig. S2C**). Meanwhile, the phosphorylated levels of p65 and

IκB were decreased in the livers of metformin-treated mice compared with the

HFD-fed mice (**Fig. S2D**), suggesting that the HFD-induced activation of NF-κB

signaling was suppressed by metformin. These results indicate that metformin can

improve HFD-induced insulin resistance, oxidative stress and inflammation.

2. Metformin prevents obesity-driven dysbiosis of gut microbiota and reshapes

intestinal metabolites

Faecal samples were collected for gut microbiota analysis by 16S rRNA

amplicon sequencing. Following all quality trimming and checking, three dataset with

highly qualified reads were collected for subsequent analysis. Our data showed that

the numbers of unique OTUs in the NCD, HFD, and HFD+Met groups were 372, 350

and 374, respectively. The coverage of sequencing results was nearly complete for all

sequences in the three groups, demonstrating sufficient sequencing depth for

follow-up analyses. The metformin-treated group significantly increased the Ace and

Chao1 indices compared with the HFD group (**Table 1**). Metformin-treated mice,

however, clustered partially apart from HFD-fed mice samples, suggesting important

changes in gut microbial profile (**Fig. 3A, Fig. S3A**). Metformin treatment prevented

HFD-induced decrease in Bacteroidetes and the increase in the

Firmicutes/Bacteroidetes ratio (**Fig. S3B,C**), two hallmarks of obesity-driven

dysbiosis. Based on FunGuild and FAPROTAX analysis of flora function, the HFD

group showed a decrease in the proportion of the chemoheterotrophy and the

fermentation bacteria when compared with the NCD group (**Fig. S3D**). However,

metformin supplementation reshaped the structure of gut microbiota,

chemoheterotrophy and fermentation improved in the HFD+Met group. Heatmap

presents the clustering of bacterial communities with their relative abundances at the

genus level (**Fig. 3B**). Compared with the NCD group, unique high-abundance

bacteria in the HFD group are mainly concentrated in *Bifidobacterium, Roseburia,*

Lachnoclostridum, Alloprevotella and *Mucispirilum*. However, metformin decreased

the abundance of these bacteria, making the metformin-treated group comparable to

the NCD group. In addition, metformin specifically increased the abundance of

Butyricimonas, Parasulterella, Parabacteroides and *Akkermansia*. Consistent with

the decrease in Firmicutes/Bacteroidetes ratio, the difference between the HFD group

and the HFD+Met group is mainly reflected in the decrease in the abundance of

Firmicutes and the increase in the abundance of Bacteroidetes (**Fig. S3E**). Compared

with the NCD and HFD+Met groups, *c_Erysipelotrichia, f_proteobacteria,*

g_Turicibacter, o_Bifidobacteriales and *s_Clostridium* in the HFD group were

significantly up-regulated, whereas f_*Muribaculaceae, g_parabacteroides* and

f_tannerellaceae were significantly up-regulated in the HFD+Met group. Therefore,

metformin can increase the abundance of Bacteriodia at the phylum level and

Akkermansia at the genus level, and modulate HFD-induced dysbiosis of gut

microbiota.

The intestinal metabolite profiles were further measured using lipid

chromatography and mass spectrometry (LC-MS) to reveal the influences of

metformin on the intestinal metabolites. Metformin caused differential changes of

intestinal metabolites in HFD-fed mice evidenced by the scatter plots of OPLS-DA

(**Fig. 3C**). Compared to HFD group, metformin up-regulated 182 of intestinal

metabolites and down-regulated 118 metabolites in the HFD+Met group (**Fig. S4A**).

A total of 85 significantly differential metabolites between HFD and HFD+Met

groups were shown in **Fig. 3D**. The HFD+Met group effectively increases the content

of taurine, choline and TUDCA in the intestine compared with the HFD group (**Fig.**

S4B). To further determine the source of TUDCA, the TUDCA content both in serum

and liver was detected. We found that TUDCA in the serum and liver of the

HFD+Met group was significantly increased compared with that in the HFD group

(**Fig. S4C**). In the HFD+Met group, the gene expression of *Cyp7a1* and *Baat*, the

important enzymes involved in the biosynthesis of TUDCA, was significantly

up-regulated compared with that in the HFD group (**Fig. S4D**). In addition, BSH

secreted by *Bifidobacterium* and *Lactobacillus* could degrade the conjugated bile

acids, such as TUDCA. The relative abundance of these two bacteria was

significantly reduced in the HFD+SFF group compared to that in the HFD group, with

a concomitant reduction both in content (**Fig. 3E**) and activity (**Fig. 3F**) of BSH. It is

noteworthy that the reduced BSH activity was not caused by metformin (**Fig. 3G**).

Fig. 4A,**B** showed the correlation between intestinal metabolism and the gut

microbiome at the phylum and species level, respectively, after the intervention of

metformin in HFD-fed mice. TUDCA was highly associated with *A. muciniphila*

species belonging to the *Verrucomicrobia* phylum. Interestingly, through the analysis

of growth curves, it was not metformin but TUDCA significantly accelerated the

proliferation of *A. muciniphila in vitro* (**Fig. 4**C). These data suggest that metformin

may indirectly improve insulin resistance caused by a high-fat diet through

modulation of the intestinal abundance of *A. muciniphila* and content of TUDCA.

3. TUDCA improves palmitic acid (PA)-induced insulin resistance and lipid

peroxidation

Further study was carried out to determine whether TUDCA could alleviate

hepatic insulin resistance using a PA-treated HepG2 cell model. PA exposure (200

µM) resulted in a decrease in glucose uptake of HepG2 cells, demonstrated by the

reduction of fluorescence density in the cytoplasm (**Fig. 5A**) and the high

concentration of glucose in the medium (**Fig. 5B**). In addition, PA decreased the level

of phosphorylated Akt at Ser473 and increased phosphorylated IRS-1 at Ser307 (**Fig.**

5C), thus, revealing a typical symptoms of insulin resistance. As shown in **Fig. 5A,B**,

the impaired glucose uptake caused by PA was significantly alleviated by TUDCA,

excluding choline and taurine (**Fig. S5A-C**). Moreover, the suppressed

phosphorylation level of Akt in PA-treated HepG2 cells was elevated by TUDCA,

while the phosphorylated IRS-1 at Ser307 was also significantly reduced (**Fig. 5C**).

Consistent with the above results, in primary mouse hepatocytes, the PA-imparied

glucose uptake was also effectively alleviated and the phosphorylated Akt was

increased by TUDCA (**Fig. 5D-F**). These findings confirmed that TUDCA could

improve hepatic glucose intolerance and enhance insulin sensitivity of PA-treated

hepatocyte.

Homeostasis imbalances on glucose metabolism often cause inflammation and

oxidative stress damage in hepatic cells. Thus, we further investigated whether

TUDCA could alleviate these pathological processes. As shown in **Fig. 6A,** PA could

increase ROS generation in primary mouse hepatocytes and HepG2 cells, while

TUDCA significantly decreased PA-induced ROS production. On the other hand, the

ratio of GSH/GSSG was significantly increased by TUDCA (**Fig. 6B**). In addition,

pro-inflammatory cytokines such as *Tnf-α*, *Il-1β* and *Il-6* were also significantly

reduced by TUDCA treatment at mRNA level (**Fig. S6**).

We further investigated whether TUDCA alleviates PA-induced insulin

resistance through the Nrf2/ARE signaling pathway. The transcriptional activity of

Nrf2 was determined by measuring the mRNA levels of its representative downstream

target genes. TUDCA significantly up-regulated the expression of Nrf2 as well as its

phosphorylation level, suppressed by PA (**Fig. 6C, Fig. S7**). It also up-regulated the

expression of antioxidant genes, *Ho-1*, *Nqo1*, and *Gpx4*, downstream of the Nrf2/ARE

signaling pathway (**Fig. 6D**). We found that TUDCA was unable to alleviate

PA-impaired glucose uptake in HepG2 cells that interfered with Nrf2 expression, as

well as the primary hepatocytes from Nrf2 knockout mice (**Fig. 6E,F, Fig. S8A,B**).

The physical interaction between Nrf2 and KEAP1 was reduced by TUDCA (**Fig.**

6G), suggesting that TUDCA potentially antagonizes with KEAP1 to keep Nrf2 from

protein degradation by ubiquitination-proteasome. Binding to the Kelch domain of

KEAP1 to achieve the inhibition of KEAP1-Nrf2 interaction is a well-documented

mechanism for Nrf2 activation, thus the molecular docking analysis was performed to

explore the interaction between TUDCA and the Kelch domain of KEAP1. The

theoretical three-dimensional binding mode of the complex with the lowest docking

energy is illustrated in **Fig. 6H**. The simulation results revealed that TUDCA was able

to interact with the primary amino acid residues on the active site of KEAP1. TUDCA

was positioned at the hydrophobic pocket, surrounded by the residues Val-418,

Leu-557, Val-514, Val-561, and Thr-560, forming a stable hydrophobic binding. All

those interactions helped TUDCA to anchor in the binding site of KEAP1, and the

estimated binding energy of KEAP1-TUDCA complex was found to be -10.21

kcal/mol. The docking simulation provided supportive evidence for TUDCA induced

Nrf2 activation by allowing us to predict the binding site in Kelch domain.

Further, *ob/ob* mice were employed to validate the effect of TUDCA in

alleviating insulin resistance and lipid accumulation *in vivo*. Compared with *ob/ob*

mice without treatment, the *ob/ob* mice administrated with TUDCA showed

remarkable decrease in body weight, FBG, FINS, as well as HOMA-IR (**Fig. 7A-E**).

TUDCA treatment substantially restored glucose intolerance (**Fig. 7F-I**) and insulin

resistance (**Fig. 7J**) in *ob/ob* mice. Moreover, similar to WT mice, *ob/ob* mice

received TUDCA showed significant amelioration of total cholesterol and triglyceride

both in serum and liver (**Fig. S9A,B**). TUDCA significantly improved Nrf2

translocation into the hepatic nucleus (**Fig. 7J**), and up-regulated its downstream

antioxidant genes *Nqo1* and *Ho-1* (**Fig. 7K**), suggesting that the Nrf2/ARE signaling

pathway was activated by TUDCA. In addition, the abundance of *A. muciniphila* in

ob/ob+TUDCA group was significantly increased compared to *ob/ob* group (**Fig. 7L**).

The gene experssion of pro-inflammatory factors (*Il-1β, Il-6,* and *Tnf-α*) in ileum were

also decresed in *ob/ob*+TUDCA group (**Fig. S9C**). Together, these results suggest

that TUDCA may significantly activate the Nrf2/ARE signaling, thereby enhancing

antioxidant and anti-inflammatory capacity as well as alleviating insulin resistance *in*

vitro and *in vivo*.

Discussion

Metabolic syndromes such as obesity, NAFLD and T2DM are increasingly

prevalent globally, especially in recent decades[2]. Metformin, the most widely used

anti-diabetic drug worldwide, has been reported to favorably influence metabolic and

cellular processes, alleviate insulin resistance[3,4]; however, the mechanism is still not

clear. Recent studies have revealed that the development of metabolic disease

strongly correlates to gut microbiota and metabolites. Multiple studies have

investigated the impact of metformin on gut microbiota, metabolites and related host

targets which regulate glucose metabolism[35]. We explored the potential mechanism

involving the gut microbiota and intestinal metabolites to reveal how metformin

improves diet-induced obesity and insulin resistance. We established that the daily

administration of metformin strongly ameliorated the obesity-related interruption of

gut microbiota. The bile acid TUDCA was maintained at a high level in the intestine,

which in turn promoted the proliferation of *A. muciniphila*, improved the integrity of

the intestinal wall and reduced intestinal inflammation. Furthermore, TUDCA was

identified as an activator of Nrf2/ARE signaling in hepatocyte that decreased

oxidative stress imbalance, while improving hepatic inflammation and insulin

resistance.

There is a variety of microbiota in the human intestine, which is involved in

body immunity, infection and nutrient metabolism, and they are closely associated

with the occurrence of diabetes. The gut microbiota imbalance in composition and

diversity can result in the release of bacterial endotoxin. This could lead to chronic

inflammation and destruction of the normal physiological functions of tissues and

organs. Several studies have confirmed that chronic inflammation can cause insulin

resistance and subsequently diabetes. The decrease in Firmicutes/Bacteroidetes ratio

is considered to be an indicator of improved metabolism by metformin therapy[35,36].

LDA analysis showed that the decreased abundance of Firmicutes by metformin could

be due to the decrease in *Erysipelotrichia* species, which belongs to the Firmicutes

phylum. The role of *Erysipelotrichia* species in human disease may be well

established in studies investigating metabolic disorders[37]. For instance, a previous

study showed a bloom of species belonging to the *Erysipelotrichaceae* family

(initially classified as *Mollicutes*) in diet-induced obese animals[38]. Within the host,

Erysipelotrichia species appear to be highly immunogenic[39], however, in surplus, they

induce intestinal inflammation. Studies have shown that species in

Erysipelotrichaceae are positively correlated with TNFα and IL-6 in the intestine and

the liver[40].

In contrast, most of the bacteria up-regulated by metformin were probiotics. For

example, species in *Muribacaceae* are involved in the L-citrulline nitrogen metabolic

cycle. They scavenge free radicals, improve immunity and maintain blood glucose.

Metformin also up-regulates the abundance of *Parabacteroides*, including *P.*

distasonis and *P. goldsteinii*. Liu et al.[41] found that *P. distasonis* could transform bile

acids, activate intestinal gluconeogenesis through succinic acid and improve the host

metabolic disorders. In another study, Wu et al.[42] recorded that *P. goldsteinii* reduces

systemic inflammation and increases insulin sensitivity by maintaining intestinal

integrity. *Akkermansia muciniphila* is known to effectively alleviate intestinal

inflammation caused by obesity[43]. Patients with intestinal obesity have been shown to

have a significantly reduced abundance of *A. muciniphila* compared with healthy

individuals. Numerous studies have demonstrated that metformin could up-regulate

the abundance of *A. muciniphila* in HFD-fed mice intestine, maintain the integrity of

the intestinal wall as well as effectively improve intestinal inflammation[44]. The

abundance of *A. muciniphila* is also significantly up-regulated in the HFD + Met

group. As a mucin-degrading bacterium, the use of *A. muciniphila* has been shown to

reduce inflammation, maintain the intestinal barrier and increase levels of

endocannabinoids secreted by intestinal peptides[45]. We speculate that the

transformation of gut microbiota by metformin is the defining factor in improving

chronic inflammation in the intestine and even the whole body. Metformin can

effectively inhibit the number of pathogenic bacteria and promote the proliferation of

probiotics. A suppression of pathogenic bacteria coupled with a proliferation of

probiotics may synergistically reduce intestinal inflammation and maintain the

integrity and permeability of the intestinal wall. Furthermore, this may lead to a

decrease in serum LPS and improvement of systemic inflammation caused by

HFD-induced obesity.

The "dialogue" between the gut microbiota and the organism mainly depends on

microbial metabolites. By producing various enzymes for biochemical metabolic

pathways of the intestinal microbes, gut microbiota can perform diverse metabolic

activities such as the metabolism of amino acids, carbohydrates and bile acids, as well

as the formation of a co-metabolic relationship with the host. Bile acids produced by

the gut microbiota are essential in the regulation of glycolipids and energy. Studies

have shown that in human primary hepatocytes, bile acids promote the β-oxidation of

fatty acids by activating FXR[46] and up-regulate the expression of apolipoprotein CII

gene to reduce the triglyceride content in the liver[47].

In our present study, metabolomics data showed that TUDCA in the intestine of

HFD+Met group increased significantly. Previous studies revealed that metformin

treatment increased the levels of conjugated bile acid in the gut by decreasing the

abundance of *Bifidobacterium* and its bile salt hydrolase (BSH) activity in the

intestines of individuals with T2DM[14,25]. BSH, secreted by intestinal bacteria,

decomposes TUDCA in the intestine and metabolizes TUDCA to taurine and UDCA.

The data from gut microbiota showed that metformin reduced the intestinal bacterial

abundance of *Bifidobacterium* and *Lactobacillus* in the mice fed a high-fat diet, while

Bifidobacterium was the main BSH-producing bacterium. In addition, the content and

activity of BSH in the mouse intestine of HFD+Met group were significantly reduced,

which was not related to metformin treatment. We hypothesize that high levels of

TUDCA are caused by a reduction in BSH-producing bacteria due to the remodeling

of the gut microbiota by metformin. *In vitro*, TCDCA can be transformed into

TUDCA by 7α/β-hydroxysteroid dehydrogenase (7α/β-HSDH)[12]. Data from the gut

microbiota also revealed that metformin did not affect the abundance of *Eubacterium*

and *Clostridium*, the major 7α/β-HSDH-producing bacteria. Therefore, the increase of

TUDCA in the intestine is purportedly not due to the inhibitory effect of metformin

on BSH-producing bacteria. According to the biosynthesis of bile acids in liver, the

conjugated bile acid TUDCA can be generated in the liver through a series of

catalytic reactions and enzymes such as CYP7A1 and BAAT[12,18]. CYP7A1 is a

rate-limiting enzyme that catalyzes the breakdown of cholesterol into bile acids in the

liver[12], whereas BAAT is a vital enzyme that catalyzes the amidation reaction of

taurine and UDCA[48]. Interestingly, we found that the gene expressions of *Cyp7a1* and

Baat were significantly enhanced by metformin, which might further promote the

synthesis of TUDCA in the liver. Thus, we suppose that the increase of TUDCA in

the intestine might due to the hepatic biosynthesis by metformin treatment.

TUDCA has been shown to treat NAFLD by acting as an endogenous chemical

chaperone; to protect cells against ER stress and to reduce liver steatosis[20]. Moreover,

TUDCA can alleviate dextran sulfate sodium-induced colitis in mice[49]. *In vitro*, we

found that TUDCA antagonizes with KEAP1 to keep Nrf2 from protein degradation

to activate Nrf2/ARE signaling pathway. This is reminiscent of the expression pattern

of Nrf2, a crucial stress regulator to reduce ROS, which is a key factor in sensitizing

insulin signaling[30,50]. It has also been confirmed that up-regulation of the Nrf2

expression both by genetic manipulation and pharmacological interference, can

significantly decrease the content of ROS and ameliorate IR and T2DM[31-34]. In the

mice supplemented with metformin, the significant up-regulation in the expression of

Nrf2 and its downstream target genes (*Gclc*, *Cat* and *Ho-1*) occurred, which further

alleviated IR and over-oxidation[51,52]. Conversely, the stress-resistant effect of

TUDCA was dramatically repressed by the inhibition of the Nrf2/ARE signaling. We

also found that TUDCA can effectively reduce the inflammation and oxidative stress

as well as decreasing the accumulation of lipid peroxide *in vitro* and *in vivo*. This

might be a novel mechanism for metformin in the treatment of insulin resistance and

diabetes.

Overall, in our present study, metformin was found to increase TUDCA by

reducing the production of BSH, which antagonized the interaction between KEAP1

and Nrf2, prevented Nrf2 degradation and promoted its nuclear translocation.

TUDCA activated Nrf2/ARE pathway and increased the gene transcription of

antioxidant enzyme, further reduced oxidative stress and sensitized insulin signaling.

The increase of TUDCA in the intestine corresponded to the increase in the

abundance of *A. muciniphlia*. Metformin remodeled the gut microbiota by promoting

probiotics such as *Muribacaceae,* and *Parabacteroides*, and inhibiting pathogenic

bacteria such as *Erysipelotrichia* and *Helicobacter*, which maintained intestinal

integrity and reduced inflammation. However, for a better understanding of the

crucial mechanisms involved in the long-term treatment with metformin in humans,

the role of the intestine-liver axis should be explored. Since TUDCA is a novel

effector molecule of metformin in the gut, plays an important role in alleviating

metabolic syndrome.

Materials and methods

Chemicals and diet

Metformin (purity $\geq$ 98%), choline (purity $\geq$ 98%), taurine (purity $\geq$ 99%) and

palmitic acid (PA, purity ≥ 99%) were obtained from Sigma-Aldrich (St. Louis, MO,

USA). Tauroursodeoxycholic acid (TUDCA, purity ≥ 99%) was purchased from

Selleck (Houston, Texas, USA). Both the normal chow diet (NCD, containing 10%

fat by energy) and high-fat diet (HFD, containing 60% fat by energy) were purchased

from Beijing HFK Bio-Technology Co., Ltd. (Beijing, China). Antibodies against Akt,

phospho-Akt (Ser473), IRS-1, phospho-IRS-1 (Ser307), KEAP1, GAPDH, lamin B1

and β-tubulin were obtained from Cell Signaling Technology (Beverly, MA, USA).

Nrf2, phospho-Nrf2 (Ser40), and 4-HNE were purchased from Abcam (Cambridge,

UK). Horseradish peroxidase (HRP)–conjugated secondary antibody was obtained

from Santa Cruz Biotechnology (Santa Cruz, USA).

Animal experiments

All male mice (ICR mice, 7 weeks old, body weight 25 ± 2 g; C57BL/6J mice, 7

weeks old, body weight 20 ± 2 g; *ob/ob* mice, 7 weeks old, body weight 40 ± 2 g;

$Nrf2^{-/-}$ mice, 7 weeks old, body weight 20 ± 2 g) were housed in ventilated cages

(three animals per cage) at the SPF facility of Wenzhou University under controlled

environmental conditions (temperature 22 ± 2°C; relative humidity 60–70%) with free

access to standard laboratory chow and tap water. The mice were maintained on a

regular 12/12 h light/dark cycle.

Mice were acclimatized to their environment for 1 week before the experiments.

Mice were randomly allocated into three groups (n = 9 for each group). A co-worker

blinded to the experimental protocol randomized animals into these groups. As for the

pharmacological activity of metformin, one group of ICR mice was fed with NCD

and the other two groups were fed with HFD. After 12 weeks of HFD feeding, one

group of HFD-fed mice was administered metformin (200 mg/kg body weight), once

daily (HFD+Met group) for 6 weeks, whereas the NCD-fed mice (NCD group) and

other group of HFD-fed mice (HFD group) were treated with an equal volume of

saline for 6 weeks. The whole study lasted 18 weeks, during which the body weight,

water consumption and food intake were measured every week. At week 18,

intraperitoneal glucose tolerance test (IPGTT) and intraperitoneal insulin tolerance

test (IPITT) were performed as previously described[1]. As for the pharmacological

activity of TUDCA, one group of *ob/ob* mice was administered TUDCA (300 mg/kg

body weight), whereas the wild-type C57BL/6J mice and *ob/ob* mice in the other two

groups were treated with an equal volume of saline for 8 weeks as control.

Fresh feces were collected and stored immediately at 80°C for subsequent

analysis. At the end of the trial, after overnight fasting for 12 h, blood samples were

collected, and serum was isolated by centrifugation at $1000 \times g$ for 15 min at 4°C, and

stored at −80°C for further assay. Tissues, including the adipose tissue, liver and

ileum, were weighed; one portion of the tissues was fixed with 10% formaldehyde for

histological analysis, and the other portion was immediately frozen in liquid nitrogen

for further analysis.

Organ index were calculated by the following formula:

$\text{BMI} = \text{body weight (kg)}/\text{body length (m)}^2$

Liver index = wet weight of liver/bw* 100%

Kidney index = wet weight of kidney/bw* 100%

Pancreas index = wet weight of pancreas/bw* 100%

Subcutaneous fat index = wet weight of adipose tissue/bw * 100%

Biochemical analysis

Total cholesterol (T-CHO), triacylglycerol (TG), free fatty acids (FFA), fasting

blood glucose (FBG), fasting serum insulin (FINS), acid phosphatase (ACP), alkaline

phosphatase (AKP), alanine aminotransferase (ALT), aspartate aminotransferase

(AST), catalase (CAT), malondialdehyde (MDA), lipopolysaccharides (LPS) were

determined by biochemical kits purchased from Jiancheng Bioengineering Institute

(Nanjing, China). The bile salt hydrolase (BSH) content was determined by

biochemical kit purchased from Jianglia Bioengineering Institute (Shanghai, China).

The ratio of reduced glutathione (GSH) and oxidized glutathione (GSSG) were

determined by a GSH:GSSG kit (Jiancheng Bioengineering Institute, Nanjing, China).

Homeostasis model assessment-estimated insulin resistance (HOMA-IR) was

calculated using the following formula:

HOMA-IR = FBG (mmol/L) × FINS (mU)/22.5

Hematoxylin and eosin staining

Adipose tissue were fixed in 10% formaldehyde overnight, paraffin-embedded,

sectioned (4-μm thickness, 3–5 sections/specimen) and stained with hematoxylin and

eosin (H&E) for histological analysis. Digital images of H&E stained sections were

acquired with a Nikon Eclipse Ti microscope at ×400 magnifications (Ti-E/U/S,

Japan). Image J software (National Institutes of Health, USA) was used to count

adipocytes.

Cell culture

498   Human liver hepatocellular carcinoma (HepG2) cells were cultured in MEM

499   (Sigma, St. Louis, MO, USA) supplemented with 10% FBS, penicillin and

500   streptomycin. Cells were maintained in 5% CO_2 at 37°C.

Isolation and culture of primary hepatocyte

Male C57/B6J and $Nrf2^{-/-}$ mice (8 weeks old) were used for isolating primary

hepatocyte. Hepatocytes were isolated by a two-step collagenase perfusion technique.

Briefly, the inferior vena cava was cannulated with angiocatheter and the portal vein

was cut. The liver was perfused via the inferior vena cava with 100 mL of PBS at

37 ℃, followed by perfusion with 100 mL of collagenase type IV (Wellington) in

HBSS containing Ca^{2+} and Mg^{2+} (GIBCO). After the liver was digested, it was

dissected out and cut into small pieces and passed through a 100 μm strainer (Falcon).

Hepatocytes were separated from non-parenchymal cells by low-speed centrifugation,

and further purified by Percoll gradient centrifugation (50% v/v, Sigma). Cells were

plated at a density of 0.3×10^6 on 6-well collagen-coated plate. Hepatocytes were

allowed to recover overnight and experiments were started 24 h post isolation.

siRNA transfection

Nrf2 was knocked-down by RNA interference (RNAi) using the following 19-bp

(including a 2-deoxynucleotide overhang) siRNAs (Origene, Beijing, China): *Nrf2*,

SR321100A-AUUGAUGUUUCUGAUCUAUCACUTT;

SR321100B-GUCAGUAUGUUGAAUCAGUAGUUTC;

SR321100C-CCAGUCUUCAUUGCUACUAAUCAGG. Stealth RNAi (Origene,

Beijing, China) was used as a negative control (siCont). For transfection, cells were

seeded on a six-well plate, grown to ~80% confluence and transfected with siRNA

duplexes using Lipofectamine 3000 (Invitrogen, Camarillo, CA, USA) according to

the manufacturer's recommendations. After incubation for 48 h, the expression level

of Nrf2 protein was detected by western blotting.

Glucose uptake assay

Glucose uptake assay was carried out according to the previous study with some

modifications[53]. Cells were plated at 1×10^4/well in 96-well plates and used at

subconfluence after 24 h of preincubation. For experiments, all culture medium was

removed from each well and replaced with 100 μM fluorescent

2-Deoxy-2-[(7-nitro-2,1,3-benzoxadiazol-4-yl) amino]-D-glucose (2-NBDG) in

serum-free medium and incubated for 30 min. Subsequently, cells were washed thrice

with PBS and then the fluorescence intensity was determined with a fluorescence

microplate reader (Ex/Em, 488/520 nm). The glucose concentration in the medium

supernatant was determined by a glucose detection kit (Jiancheng Bioengineering

Institute, Nanjing, China).

Determination of ROS levels

536   Intercellular ROS levels were measured using a ROS Assay Kit (Beyotime,

537 Shanghai, China) according to the manufacturer's protocol. In brief, the

538 dichlorodihydrofluorescein diacetate (DCFH-DA; 10 mM) supplied in the kit was

539 diluted to 10 μM in serum-free medium. Subsequently, cells were pretreated with

540 TUDCA (20 μM) for 2 hour at a density of 3×10^3/96-well. Next, 200 μM PA

541 solution was added and co-incubated for 24 hours. The supernatant was then replaced

542 with serum-free medium. The DCFH-DA solution (10 μM) was added to the cells for

543 30 minutes at 37°C, and they were then resuspended with PBS (0.1 mM) after

544 washing twice with the indicated inducers. Fluorescence intensity was detected using

545 a hybrid/multi-modereaders (BioTek instruments Inc., Vermont, USA) to determine

546 ROS levels.

547 **Quantitative RT-PCR**

548   Total mRNA was isolated from tissue samples using TRIzol reagent (TAKARA,

549 Tokyo, Japan) and was reverse-transcribed into cDNA using a high-capacity cDNA

550 reverse transcription kit (TAKARA, Tokyo, Japan) according to the manufacturer's

551 protocol. The mRNA levels were quantified with quantitative PCR (qPCR) using

552 SYBR Green (Qiagen, Hilden, Germany). Amplification was performed on a

553 LightCycler480 *q*RT-PCR system (Roche, Mannheim, Germany) under the following

554 reactions: 95°C for 15 min, followed by 40 cycles at 95°C for 10 s, 60°C for 20 s, and

555 72°C for 20 s. The relative mRNA levels of target genes were normalized to the

556 expression of β-actin calculated using $2^{-\Delta\Delta ct}$ method. The primer pairs used in this

study are listed in **Supplementary table**.

Western blotting

Samples were homogenized with ice-cold RIPA lysis buffer containing protease

and phosphatase inhibitors (Beyotime, Shanghai, China). The homogenates were

centrifuged at 10 000 × g for 20 min at 4°C to remove the insoluble tissue debris. The

protein concentration in the supernatant was determined using a BCA protein assay

kit (Beyotime, Shanghai, China). Equal amount of protein for each group were

denatured in boiling water for 5 min. Aliquots (40 μg) of protein samples were

subjected to 10% SDS-PAGE and transferred to PVDF membranes (Millipore,

Bedford, MA, USA). After blocking with 5% nonfat milk (dissolved in TBST) for 1 h,

the membranes were incubated with the indicated antibodies at 4°C overnight,

followed by incubation with the appropriate HRP-conjugated second antibodies for 1

h at room temperature. Chemiluminescent detection was performed using the ECL

Plus Western blotting reagent (TransStart, Beijing, China). Semi-quantitative analysis

for densitometry of each band was performed using ImageJ software.

Co-immunoprecipitation

For co-immunoprecipitation (Co-IP), the whole protein lysates prepared from

HepG2 cells were extracted in a RIPA lysis buffer. Briefly, cell lysates were

incubated with Nrf2 or KEAP1 antibodies for 2 h at room temperature, followed by

binding of antigen antibody complexes to protein A/G magnetic beads (Thermo

scientific, Rockford, USA) for 1 h. Then, the antigen/antibody immunocomplex was

stripped by boiling with 5× loading sample buffer for western blotting incubated with

KEAP1, Nrf2 and GAPDH antibodies.

Methodologies of gut microbiota by 16S rRNA amplicon sequencing

Sample preparation: Fæcal samples were freshly collected at week 16 and

immediately stored at -80°C. Total fecal DNA was extracted using CTAB/SDS

method. The V4 region of the 16S rRNA was amplified using the universal primers

515F and 806R. All PCR reactions were carried out with Phusion® High-Fidelity PCR

Master Mix (New England Biolabs), and the mixture PCR products were purified

with GeneJET Gel Extraction Kit (Thermo Scientific). Sequencing libraries were

generated using Ion Plus Fragment Library Kit (Thermo Scientific) following

manufacturer's recommendations. The library quality was assessed on the Qubit 2.0

Fluorometer (Thermo Scientific). At last, the library was sequenced on an Ion S5™

XL platform (Thermo Scientific) and 400 bp/600 bp single end reads were generated.

Data acquisition: Paired-end reads were merged using FLASH (V1.2.7), and the

Raw fastq files were processed by QIIME (version 1.7.0). Sequence analysis was

performed by Uparse software (v7.0.1001), and Sequences with $\geqslant$ 97% identity

were assigned to the same Operational Taxonomic Units (OTUs).

Data analysis: To compare the compositional OTUs of the gut microbiota in each

group, a Venn diagram was constructed using R packages (version 3.1.0) as

previously documented. Mothur software packages (version V.1.30.1) were used to

calculate the value of Chao1, Ace, Simpson index and Shannon index for evaluation

of the community richness and community diversity. Pair-group method with

arithmetic means (UPGMA) clustering was generated using the average linkage and

conducted by QIIME software (V1.7.0). Linear discriminant analysis (LDA) was

carried out to determine the highly dimensional gut microbes and characteristics

associated with NCD mice, HFD mice and HFD+Met mice.

Methodologies of non-targeted metabolome analysis

Metabolites extraction from colon contents: Colon contents were added ddH2O

(4°C) and mixed. 100 mg of sample was extracted with 1000 μL of pre-cooled

methanol (-20°C). After centrifugation, the supernatant was evaporated and finally

dissolved in 400 μL methanol aqueous solution (1:1, 4°C). For the quality control

(QC) samples, 20 μL of extract was taken from each sample and mixed. These QC

samples were used to monitor deviations of the analytical results from these pool

mixtures and compare them to the errors caused by the analytical instrument itself.

And the rest of the samples were used for LC-MS detection.

UPLC Conditions: Chromatographic separation was accomplished in an Acquity

UPLC system equipped with an ACQUITY UPLC® HSS T3 (150 × 2.1 mm, 1.8 μm,

Waters) column maintained at 4°C. The temperature of the auto sampler was 4°C.

Gradient elution of analytes was carried out with 0.1% formic acid in water (A) and

0.1% formic acid in acetonitrile (B) at a flow rate of 0.25 mL/min. Injection of 5 μL

of each sample was done after equilibration. An increasing linear gradient of solvent

B (v/v) was used as follows: 0~1 min, 2% B; 1~9.5 min, 2%~50% B; 9.5~14 min,

50%~98% B; 14~15 min, 98% B; 15~15.5 min, 98%~2% B; 15.5~17 min, 2%.

Mass spectrometry conditions: The ESI-MS^n experiments were executed on the

Thermo LTQ Orbitrap XL mass spectrometer with the spray voltage of 4.8 kV and

-4.5 kV in positive and negative modes, respectively. Sheath gas and auxiliary gas

were set at 45 and 15 arbitrary units, respectively. The capillary temperature was

325°C. The voltages of capillary and tube were 35 V and 50 V, -15 V and -50 V in

positive and negative modes, respectively. The Orbitrap analyzer scanned over a mass

range of m/z 89-1000 for full scan at a mass resolution of 60000. Data dependent

acquisition (DDA) MS/MS experiments were performed with CID scan. The

normalized collision energy was 30 eV. Dynamic exclusion was implemented with a

repeat count of 2, and exclusion duration of 15 s.

Data processing: UPLC-QTOF-MS raw data were analyzed with MarkerLynx

Application Manager 4.1 (Waters Corp.). The matrix from UPLC-QTOF-MS was

introduced into SIMCA-P 11.0 software (Umetrics) and standardized to a mean of 0

and variance of 1, according to the formula [X - mean(X)]/ std (X), for multivariate

statistical analysis. The t test with false discovery rate correction was used to measure

the significance of each metabolite. Partial least-squared discriminant analysis

(PLS-DA) and orthogonal partial least-squared discriminant analysis (OPLS-DA)

were conducted to identify the metabolite discrimination between the two group

samples. Differential metabolites were defined with variable importance in the

projection (VIP) > 1.0 obtained from OPLS-DA and P values less than 0.05 obtained

from t test. Differential metabolites were tentatively identified by database matching,

i.e., Human Metabolome Database (HMDB) (http://www.hmdb.ca), Metlin

(http://metlin.scripps.edu), massbank (http://www.massbank.jp/), LipidMaps

(http://www.lipidmaps.org), mzclound (https://www.mzcloud.org). Heatmaps of

differential metabolites among all groups were obtained based on spearman

correlation and cluster analyses.

Growth curve of *Akkermansia muciniphila*

Akkermansia muciniphila strain ATCC BAA-835 was purchased from the Beina

Biologicals, Inc. (Beijing, China). Bacteria were cultured in BHI medium in tubes at

37°C in an anaerobic chamber (Whitley A35 Workstation 2.5, Don Whitley Scientific,

UK). To acquire the growth curve of *A. muciniphila*, the medium in test tubes

supplemented with or without 10 mM metformin (or 20 μM TUDCA) was placed in

an anaerobic bag to eliminate oxygen for 24 h. The test tubes were then inoculated

with bacteria and incubated at 37°C in an anaerobic bag. The OD_{600} of the cultures

was measured every 3 hours.

Determination of hepatic/serum TUDCA

The contents of TUDCA in tissues were determined by high-performance liquid

chromatography (HPLC). Briefly, 200 mg of frozen tissue was homogenized with a

Qiagen TissueLyserII (Germantown, MD, USA) in 1 mL of PBS to prepare the tissue

homogenates. The impurities were removed through a 0.22 μm filter, then the filtrate

was precipitated by methanol and 10 μL of supernatant was analyzed using an Agilent

1290 HPLC system (Santa Clara, CA) equipped with a Hypersil ODS-2 column (5 µm,

4.6 × 250 mm; Waters, USA). The supernatant was detected at the wavelength of 210

nm, eluted with the mobile phase of 0.03 M phosphate buffer solution (pH 4.4) and

methanol (32:68, v/v) at a flow rate of 1.0 mL/min.

Determination of bile salt hydrolase activity *in vitro*

The enzyme activity of BSH was determined using an amino acid chromogenic

method. Briefly, mouse faeces were collected and the faecal suspension was prepared.

10 µL of sample (containing 500 µg of protein) was incubated with 180 µL of sodium

phosphate buffer and 10 µL of TDCA (final concentration 10 mM) at 37°C for 30 min.

50 µL of the mixture was mixed with 50 µL of 15% TCA and centrifuged at 10,000g

for 10 min. Mix 20 µL of supernatant with 80 µL of water, add 1.9 mL of ninhydrin

(0.5 mL 1% ninhydrin, 1.2 mL glycerol, 0.2 mL 0.5 M citrate buffer, pH 5.5) and

boil for 15 min. Cool the reaction tube, measure the absorbance at 570 nm and

calculate the BSH enzyme activity by substituting the formula for the taurine standard

curve.

Molecular docking analysis

To investigate the probable binding of TUDCA to KEAP1 as the potential

inhibitor, the automated docking studies were carried out using AutoDock vina 1.1.2

package. The X-ray crystal structure of the human KEAP1 (PDB ID: 6LRZ) was

downloaded from RCSB Protein Data Bank. The 3D structure of TUDCA (ZINC ID:

3914813) was downloaded from ZINC. The AutoDockTools 1.5.6 package was

employed to generate the docking input files. The molecular docking simulation

protein of Keap1 was prepared by removing water molecules and bound ligands. The

binding site of the KEAP1 was identified as centre_x: -9.944, centre_y: -38.993, and

centre_z: 3.869 with dimensions size_x: 85, size_y: 85, and size_z: 85. The

best-scoring (i.e.,with the lowest docking energy) pose as judged by the Vina docking

score was chosen and further analyzed using PyMoL 2.4.0 software.

Statistical analysis

Statistical comparisons among experimental groups were analyzed by one-way

ANOVA and Duncan's multiple-comparison test using the SPSS software (Version

21.0. Armonk, NY: IBM Corp.). *P* values less than 0.05 were considered statistically

significant.

Acknowledgements

We gratefully thank Qiaojuan Li and Lingfeng Hou (Wenzhou University) for

their assistance in animal experiments. We gratefully thank Alan K Chang (Wenzhou

University) for helpful discussion and for revising the language of the manuscript.

Financial support

This work was financially supported by the National Natural Science Foundation

of China (81872952 and 41876197), the National Key Research and Development

Project (2018YFD0901503), the Science and Technology Program of Wenzhou

(ZY2019013).

Conflict of interest

The authors declare no conflict of interest.

Authors' contributions

Y.Z., Y.C., J.L, D.H., J.Z. and L.Y performed the research. H.T., Y.Z. and Y.G.

designed this study. R.T. and M.W. provided technical assistance. H.T. and Y.Z.

analyzed the data and wrote the manuscript. R.T. and Y.G. revised the manuscript.

H.T. and Y.G. had primary responsibility for final content. All authors have read and

approved the final manuscript.

References

1. Muoio DM, Newgard CB. Mechanisms of disease: Molecular and metabolic mechanisms of insulin resistance and beta-cell failure in type 2 diabetes. *Nat. Rev. Mol. Cell Bio.* 2008; 9: 193-205.

2. Estes C, Razavi H, Loomba R, Younossi Z, Sanyal AJ. Modeling the epidemic of nonalcoholic fatty liver disease demonstrates an exponential increase in burden of disease. Hepatology. 2018; 67(1): 123-133.

3. Shen J, Obin MS, Zhao L. The gut microbiota, obesity and insulin resistance. *Mol. Aspec. Medic.* 2013; 34(1): 39-58.

4. Bäckhed F, Manchester JK, Semenkovich CF, Gordon JI. Mechanisms underlying the resistance to diet-induced obesity in germ-free mice. *P. Natl. Acda. Sci. USA.* 2007;104(3): 979-984.

5. Cho NH, Shaw JE, Karuranga S, Huang Y, Rocha Fernandes JD, Ohlrogge AW, Malanda B. IDF diabetes atlas: global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Res. Clin. Pract.* 2018; 138: 271-281.

6. Zhou G, Myers R, Li Y, Chen Y, Shen X, Fenyk-Melody J, Wu M, Ventre J, Doebber T, Fujii N, Musi N, Hirshman MF, Goodyear LJ, Moller DE. Role of AMP-activated protein kinase in mechanism of metformin action. *J. Clin. Invest.* 2001; 108: 1167-1174.

7. Lee H, Ko G. Effect of metformin on metabolic improvement and gut microbiota. *Appl. Enviro. Micorb.* 2014; 80: 5935-5943.

8.   Zhang X, Zhao Y, Xu J, Xue ZS, Zhang MH, Pang XY, Zhang XJ, Zhao LP.

Modulation of gut microbiota by berberine and metformin during the treatment of

high-fat diet-induced obesity in rats. *Sci. Rep.* 2015; 5: 14405.

9.   Zhou ZY, Ren LW, Zhan P, Yang HY, Chai DD, Yu ZW. Metformin exerts

glucose-lowering action in high-fat fed mice via attenuating endotoxemia and

enhancing insulin signaling. *Acta Pharmacol. Sin.* 2016; 37: 1063-1075.

10. Kristoffer F, Falk H, Trine N, Gwen F, Emmanuelle LC, Shinichi S, Edi P,

Sara VS, Valborg G, Helle KP, Manimozhiyan A, Karsten K, Anita YV, Henrik

V, Rajna H, Paul IC, Jens RK, Junhua L, Torben J, Florence L, Joël D, Søren B,

Jeroen R, Torben H, Jun W, Dusko ES, Peer B, and Oluf P. Disentangling the

effects of type 2 diabetes and metformin on the human gut microbiota. *Nature.*

2015; 528(7581): 262–266.

11. Pedersen HK, Gudmundsdottir V, Nielsen HB, Hyotylainen T, Nielsen T,

Jensen BA, Forslund K, Hildebrand F, Prifti E, Falony G, Le Chatelier E,

Levenez F, Doré J, Mattila I, Plichta DR, Pöhö P, Hellgren LI, Arumugam M,

Sunagawa S, Vieira-Silva S, Jørgensen T, Holm JB, Trošt K; MetaHIT

Consortium, Kristiansen K, Brix S, Raes J, Wang J, Hansen T, Bork P, Brunak S,

Oresic M, Ehrlich SD, Pedersen O. Human gut microbes impact host serum

metabolome and insulin sensitivity. *Nature.* 2016; 535(7612): 376-381.

12. Jia W, Xie GX, Jia WP. Bile acid-microbiota crosstalk in gastrointestinal

inflammation and carcinogenesis. *Nat. Rev. Gastroenterol Hepatol.* 2018; 15(2):

111-128.

13. Yao L, Seaton SC, Ndousse-Fetter S, Adhikari AA, DiBenedetto N, Mina AI,

Banks AS, Bry L, Devlin AS. A selective gut bacterial bile salt hydrolase alters

host metabolism. *eLife*. 2018; 17(7): e37182.

14. Sun LL, Xie C, Wang G, Wu Y, Wu Q, Wang XM, Liu J, Deng YY, Xia JL,

Bo C, Zhang SY, Chuyu Y, Lian G, Zhang XJ, Zhang H, Bisson WH, Shi JM,

Gao XX, Ge PP, Liu CH, Krausz WK, Nicholis RG, Cai JW, Rimal BP, Patterson

A, Wang X, Gonzalez FJ, Jiang CT. Gut microbiota and intestinal FXR mediate

the clinical benefits of metformin. *Nat. Med.* 2018; 24(12): 1919-1929.

15. Martínez-Moya P, Romero-Calvo I, Requena P, Hernández-Chirlaque C,

Aranda CJ, González R, Zarzuelo A, Suárez MD, Martínez-Augustin O, Marín JJ,

de Medina FS. Dose-dependent antiinflammatory effect of ursodeoxycholic acid

in experimental colitis. *Int. Immunopharmacol.* 2013; 15(2): 372–380.

16. Berger E, Haller D. Structure-function analysis of the tertiary bile acid

TUDCA for the resolution of endoplasmic reticulum stress in intestinal epithelial

cells. *Biochem. Biophys. Res. Commun.* 2011; 409: 610–615.

17. Stenman LK, Holma R, Gylling H, Korpela R. Genetically obese mice do not

show increased gut permeability or faecal bile acid hydrophobicity. *Br. J. Nutr.*

2013; 110: 1157–1164.

18. Arab JP, Karpen SJ, Dawson PA, Arrese M, Trauner M. Bile acids and

nonalcoholic fatty liver disease: Molecular insights and therapeutic perspectives.

Hepatology. 2017; 65: 350–362.

19. Wang WJ, Zhao JF, Gui WF, Sun D, Dai HJ, Xiao L, Chu HK, Fan D, Zhu QJ, Bernd S, Huang K, Yang L, and Hou XH. Tauroursodeoxycholic acid inhibits intestinal inflammation and barrier disruption in mice with non-alcoholic fatty liver disease. *Brit. J. Pharmacol.* 2018; 175: 469–484.

20. Xie Q, Khaoustov VI, Chung CC, Sohn J, Krishnan B, Lewis DE, Yoffe B. Effect of tauroursodeoxycholic acid on endoplasmic reticulum stress-induced caspase-12 activation. *Hepatology.* 2002; 36(3): 592–601.

21. Fondevila C, Shen XD, Tsuchiyashi S, Yamashita K, Csizmadia E, Lassman C, Busuttil RW, Kupiec-Weglinski JM, Bach FH. Biliverdin therapy protects rat livers from ischemia and reperfusion injury. *Hepatology.* 2004; 40: 1333–1341.

22. Nakao A, Otterbein LE, Overhaus M, Sarady JK, Tsung A, Kimizuka K, Nalesnik MA, Kaizu T, Uchiyama T, Liu F, Murase N, Bauer AJ, Bach FH, Biliverdin protects the functional integrity of a transplanted syngeneic small bowel. *Gastroenterology.* 2004; 127: 595–606.

23. Ester GS, Maria JP, Nikolaj SN, Oscar B, Maria JM, Elisa L, Maria AS, Jose JGM. Protective role of biliverdin against bile acid-induced oxidative stress in liver cells. *Free Radical Bio. Med.* 2016; 97: 466–477.

24. Zhang M, Feng R, Yang M, Qian C, Wang Z, Liu W, Ma J. Effects of metformin, acarbose, and sitagliptin monotherapy on gut microbiota in Zucker diabetic fatty rats. *BMJ Open Diab. Res. Ca.* 2019; 7(1): e000717.

25. Nova E, Pérez de Heredia F, Gómez-Martínez S, Marcos A. The Role of

Probiotics on the Microbiota: Effect on Obesity. *Nutr. Clin. Pract.* 2016; 31(3):

387-400.

26. Zhang WP, Chen LH, Feng H, Wang W, Cai Y, Qi F, Tao XF, Liu J, Shen

YJ, Ren XF, Chen X, Xu JM, Shen YX. Rifampicin-induced injury in HepG2

cells is alleviated by TUDCA via increasing bile acid transporters expression and

enhancing the Nrf2- mediated adaptive response. *Free Radical Bio. Med.* 2017;

112: 24–35.

27. Tangvarasittichai S. Oxidative stress, insulin resistance, dyslipidemia and

type 2 diabetes mellitus. *World J. Diabetes.* 2015; 6(3): 456–480.

28. Maiese K. New insights for oxidative stress and diabetes mellitus. *Oxid. Med.*

Cell Longev. 2015; 2015: e875961.

29. Raza H, John A, Howarth FC. Increased oxidative stress and mitochondrial

dysfunction in zucker diabetic rat liver and brain. *Cell Physiol. Biochem.* 2015;

35(3): 1241–1251.

30. Wan HWN, Kwak MK, Makpol S, Wan NWS, Mohd Yusof YA. Piper betle

induces phase I & II genes through Nrf2/ARE signaling pathway in mouse

embryonic fibroblasts derived from wild type and Nrf2 knockout cells, BMC

Complement. *Altern. Med*. 2014; 14: 72.

31. Uruno A, Yagishita Y, Yamamoto M. The Keap1-Nrf2 system and diabetes

mellitus. *Arch. Biochem. Biophys*. 2015; 566: 76–84.

32. Uruno A, Furusawa Y, Yagishita Y, Fukutomi T, Muramatsu H, Negishi T,

Sugawara A, Kensler TW, Yamamoto M. The Keap1-Nrf2 system prevents onset

of diabetes mellitus. *Mol. Cell Biol.* 2013, 33(15): 2996–3010.

33. Peng DF, Lu H, Zhu SM, Zhou ZJ, Hu TL, Chen Z, Zaika A, El-Rifai W.

NRF2 antioxidant response protects against acidic bile salts- induced oxidative

stress and DNA damage in esophageal cells. *Cancer Lett.* 2019; 458: 46–55.

34. Kamisako T, Tanaka Y, Kishino Y, Ikeda T, Yamamoto K, Masuda S,

Ogawa H. Role of Nrf2 in the alteration of cholesterol and bile acid

metabolismrelated gene expression by dietary cholesterol in high fat fed mice. *J.*

Clin. Biochem. Nutr. 2014; 54(2): 90–94.

35. Canfora EE, Meex R, Venema K, and Blaak EE. Gut microbial metabolites

in obesity, NAFLD and T2DM. *Nat. Rev. Endocrinol.* 2019; 15: 261–273.

36. Kaakoush NO. Insights into the Role of erysipelotrichaceae in the human

host. *Front. Cell. Infect. Microbiol.* 2015; 20: 84.

37. Turnbaugh PJ, Backhed F, Fulton L, and Gordon JI. Marked alterations in

the distal gut microbiome linked to diet-induced obesity. *Cell Host Microbe.*

2008; 3(4): 213–223.

38. Craven M, Egan CE, Dowd SE, McDonough SP, Dogan B, Denkers EY.

Inflammation drives dysbiosis and bacterial invasion in murine models of ileal

Crohn's disease. *PLoS ONE.* 2012; 7: e41594.

39. Ff

842 40.

843 41. Wang K, Liao MF, Zhou N, Bao L, Ma K, Zheng ZY, Wang YJ, Liu C,

844 Wang WZ, Wang J, Liu SJ, Liu HW. *Parabacteroides distasonis* alleviates

845 obesity and metabolic dysfunctions via production of succinate and secondary

846 bile acids. *Cell Rep.* 2019; 26(1): 222-235.

847 42. Wu TR, Lin CS, Chang CJ, Lin TL, Martel J, Ko YF, Ojcius MD, Lu CC,

848 Young DJ, Lai HC. Gut commensal *Parabacteroides goldsteinii* plays a

849 predominant role in the anti-obesity effects of polysaccharides isolated from

850 *Hirsutella sinensis*. *Gut.* 2019; 68(2): 248–262.

851 43. Anhê FF, Roy D, Pilon G, Dudonne S, Matamoros S, Varin T, Marette A. A

852 polyphenol-rich cranberry extract protects from diet-induced obesity, insulin

853 resistance and intestinal inflammation in association with increased *akkermansia*

854 *spp*. population in the gut microbiota of mice. *Gut.* 2015; 64(6): 872–883.

855 44. Chelakkot C, Choi Y, Kim DK, Park HT, Ghim J, Kwon Y, Jeon J, Kim MS,

856 Jee YK, Gho YS, Park HS, Kim YK, Ryu SH. *Akkermansia muciniphila*-derived

857 extracellular vesicles influence gut permeability through the regulation of tight

858 junctions. *Exp. Mol. Med*. 2018; 50(2): e450.

859 45. Everard A, Belzer C, Geurts L, Ouwerkerk JP, Druart C, Bindels LB, Guiot

860 Y, Derrien M, Muccioli GG, Delzenne NM, de Vos WM, Cani PD. Cross-talk

861 between Akkermansia muciniphila and intestinal epithelium controls diet-induced

862 obesity. *Proc. Natl. Acad. Sci. U S A*. 2013; 110(22):9066-9071.

46. Manley S, Ding WX. Role of farnesoid X receptor and bile acids in alcoholic

liver disease. *Acta Pharm. Sin. B.* 2015; 5(2): 158–167.

47. Kast HR, Nguyen CM, Sinal CJ, Jones SA, Laffitte BA, Reue K, Gonzalez

FJ, Willson TM, Edwards PA. Farnesoid X-activated receptor induces

apolipoprotein C-II transcription: a molecular mechanism linking plasma

triglyceride levels to bile acids. *Mol. Endocrinol.* 2001; 15(10):1720-1728.

48. Chiang YLJ. Bile acids: regulation of synthesis. *J. Lipid Res.* 2009; 50(10):

1955-1966.

49. Rodriguez J, Hiel S, Delzenne NM. Metformin: old friend, new ways of

action-implication of the gut microbiome? *Curr. Opin. Clin. Nutr. Metab. Care.*

2018; 21(4): 294-301.

50. Wible RS, Ramanathan C, Sutter CH, Olesen KM, Kensler TW, Liu AC,

Sutter TR. NRF2 regulates core and stabilizing circadian clock loops, coupling

redox and timekeeping in Mus musculus. Elife. 2018 Feb 26;7:e31656.

51. Sun CC, Lai YN, Wang WH, Xu XM, Li XQ, Wang H, Zheng JY, Zheng JQ.

Metformin Ameliorates Gestational Diabetes Mellitus-Induced Endothelial

Dysfunction via Downregulation of p65 and Upregulation of Nrf2. *Front*

Pharmacol. 2020; 11: e575390.

52. Abdelsamia EM, Khaleel SA, Balah A, Abdel Baky NA. Curcumin augments

the cardioprotective effect of metformin in an experimental model of type I

diabetes mellitus; Impact of Nrf2/HO-1 and JAK/STAT pathways. Biomed

Pharmacother. 2019; 109: 2136-2144.

53. Zou CH, Wang YJ, Zhen ZF. 2-NBDG as a fluorescent indicator for direct

glucose uptake measurement. *J. Biochem. Biophys. Methods.* 2005; 64: 207–215.

Figure captions

Figure 1. Metformin prevents obesity and insulin resistance in HFD-fed mice. (A)

Body weight was shown throughout 6 weeks. (B,C) Mice, fasted overnight and fasted

for 6 hours, were injected with glucose (2 g/kg in saline) or insulin (0.75 IU/kg in

saline) for IPGTT and IPITT, respectively. (B) IPGTT and (B') AUC of IPGTT; (C)

IPITT and (C') slope of IPITT. (D) FINS and (E) HOMA-IR. (F) Protein expression

of Akt, phosphorylated Akt at Ser473, IRS-1, phosphorylated IRS-1 at Ser307 and

quantitative analysis for the densitometry. Data are expressed as mean ± SD. *$P <$

0.05, **$P < 0.01$ and ***$P < 0.001$ for HFD vs HFD+Met. AUC, area under the curve;

FINS, fasting serum insulin; HOMA-IR, homeostatic model assessment for insulin

resistance; IPGTT, intraperitoneal glucose tolerance test; IPITT, intraperitoneal

insulin tolerance test; Met, metformin.

Figure 2. Metformin ameliorates HFD-induced oxidative stress. (A-C) Serum or

liver homogenate were used to quantify the representative of oxidative stress: (A)

CAT, (B) GSH:GSSG ratio, and (C) MDA. (D) Protein expression of total Nrf2,

phosphorylated Nrf2 and (D', D'') quantitative analysis for the densitometry. (E)

Quantitative RT-PCR analysis of *Nrf2*, *Keap1*, *Nqo1* and *Ho-1* are shown. Data are

expressed as mean ± SD. *$P < 0.05$, **$P < 0.01$ and ***$P < 0.001$ for HFD vs

HFD+Met. CAT, catalase; GSH, glutathione; GSSG, oxidized glutathione; MDA,

malondialdehyde.

Figure 3. Genus-based comparison of gut microbiota and non-targeted

metabolome analysis. (A) Principal coordinates analysis. (B) Heatmap showing

clustering of bacterial communities with their relative abundances at the genus level,

the bacterial communities or OTUs were clustered phylogenetically by

neighbor-joining method and the libraries were clustered by profile pattern using

Unifrac analysis. Relative abundances (log values) of microbial genus are displayed

in a red-to-blue color code (high to low abundance). (C) OPLS-DA score plot of

non-targeted metabolome. (D) Heatmap of 85 significantly differential metabolites. (E)

BSH content and (F) BSH activity in faeces of the three groups of mice. (G) Fecal

suspensions were prepared by collecting fresh faeces, adding metformin to final

concentrations of 5, 10 and 20 mg/mL, incubating for 30 min and then measuring

BSH activity. Data are expressed as mean ± SD. *$P < 0.05$, **$P < 0.01$ and ***$P <$

0.001 for HFD *vs* HFD+Met. PCoA, principal coordinates analysis; OPLS-DA,

orthogonal partial least-squared discriminant analysis; Met, metformin; BSH, bile salt

hydrolase.

Figure 4. Correlation heatmap for gut microorganisms and metabolites. (A-B)

The correlation coefficients of the differential metabolites and gut microbiota were

calculated by Pearson's algorithm using R language, and heat maps were created

based on the correlation coefficients. Matrix heatmap showing the correlation

between gut microbiota and metabolome at (A) Phylum and (B) Species levels,

respectively. (C) Growth curve of *A. muciniphila* in the presence of metformin or

TUDCA. Met, metformin; TUDCA, tauroursodeoxycholic acid.

Figure 5. TUDCA alleviates insulin resistance in PA-treated hepatocytes. (A-C)

HepG2 cells were pre-treated with 200 μM PA for 24 h and then incubated with

TUDCA (10 or 20 μM) for 12 h. (A) Glucose uptake and (B) glucose in the medium

supernatant. (C) Protein levels of p-Akt, Akt, p-IRS-1, IRS-1 and (C', C'')

quantitative analysis for the densitometry. (D-F) Primary mouse hepatocytes were

pre-treated with 200 μM PA for 24 h and then incubated with TUDCA (10 or 20 μM)

for 4 h. (D) Glucose uptake and (E) glucose in the medium supernatant. (F) Protein

levels of p-Akt, Akt, p-IRS-1, IRS-1 and (F', F'') quantitative analysis for the

densitometry. Data are expressed as the mean ± SD. *$P < 0.05$, **$P < 0.01$ and ***P

< 0.001. PA, palmitic acid; TUDCA, tauroursodeoxycholic acid.

Figure 6. TUDCA relieves PA-induced insulin resistance via activating

Nrf2/ARE signaling pathway. (A) Intracellular ROS content; (B) ratio of

GSH/GSSG; (C) protein levels of p-Nrf2 and Nrf2 were determined by western

blotting in primary mouse hepatocytes and HepG2 cells. (D) Quantitative RT-PCR

analysis of *Nrf2*, *Ho-1*, *Nqo1* and *Gpx4* in primary mouse hepatocytes and HepG2

cells. Glucose uptake by cells and glucose content of the medium supernatant

measured in Nrf2 knock-down HepG2 cells (E) and primary hepatocytes from Nrf2$^{-/-}$

mice (F). (G) Immunoprecipitation. HepG2 cells were treated with or without

TUDCA, and then lysates were immunoprecipitated with the indicated antibodies,

followed by western blotting. (H) Docking of TUDCA with Kelch domain of KEAP1.

The residues of KEAP1 were represented using sticks or surface structures; TUDCA

was shown in pink; the dashed lines (blue) represent hydrogen-bonding interactions.

Data are expressed as mean ± SD. *$P < 0.05$, **$P < 0.01$ and ***$P < 0.001$. ROS,

reactive oxygen species; GSH, glutathione; GSSG, oxidized glutathione; PA, palmitic

acid; TUDCA, tauroursodeoxycholic acid.

Figure 7. TUDCA improves obesity and insulin resistance in *ob/ob* mice. (A-E)

Wild-type C57BL/6J mice and *ob/ob* mice was administered with or without TUDCA

for 8 weeks. (A) Body size, (B) body weight, serum levels of (C) FBG and (D) FINS,

(E) HOMA-IR. (F-I) Mice, fasted overnight and fasted for 6 hours, were injected with

glucose (2 g/kg in saline) or insulin (0.75 IU/kg in saline) for IPGTT and IPITT,

respectively. (F) IPGTT, (G) AUC of IPGTT, (H) IPITT and (I) slope of IPITT are

shown. (G) Protein expression of p-Akt, Akt, Nrf2 in nuclear and cytoplasm. (K)

Quantitative RT-PCR analysis of *Nqo1* and *Ho-1*. (L) Abundance of *A. muciniphila*.

Data are expressed as mean ± SD. *$P < 0.05$, **$P < 0.01$ and ***$P < 0.001$ for *ob/ob*

vs *ob/ob*+TUDCA. AUC, area under the curve; FBG, fasting blood glucose; FINS,

fasting serum insulin; HOMA-IR, homeostatic model assessment for insulin

resistance; IPGTT, intraperitoneal glucose tolerance test; IPITT, intraperitoneal

insulin tolerance test; TUDCA, tauroursodeoxycholic acid.

Table 1. Diversity of gut microbiota in HFD and HFD+Met treated mice

Groups	Reads	OTUs	Observed species	Ace	Chao1	Shannon	Simpson
NCD	78316±4945	372±35	330±21	345±17	350±21	5.10±0.67	0.92±0.04
HFD	76154±7745	350±40	312±20	325±24	326±24	5.38±0.33	0.95±0.02
HFD+Met	80004±421	374±21	335±20*	342±22*	348±20*	5.24±0.27	0.95±0.02

970 Data are expressed as mean ± SD. *P < 0.05 compared with HFD.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7