Metformin alleviates insulin resistance in obese-induced mice via remodeling gut

microbiota and intestinal metabolites

Ya Zhang [a,b,1], Yang Cheng [a,1], Jihui Zuo [a,1], Liping Yan [a], Qiaojuan Li [a], Ronald W.

Thring [b], Yue Yang [a], Yitian Gao [a,*], Mingjiang Wu [a,*], Haibin Tong [a,*]

[a] College of Life and Environmental Science, Wenzhou University, Wenzhou 325035,

China;

[b] Environmental Science and Environmental Engineering, University of Northern

British Columbia, Prince George, BC, Canada;

*Correspondence: tonghaibin@gmail.com (H.T.); wmj@wzu.edu.cn (M.W.);

gyt@wzu.edu.cn (Y.G.); Tel.: + 86-577-86689078

[1] These authors contributed equally to this study.

Abstract

Metformin is widely used to surmount insulin resistance (IR) and diabetes.

Evidence indicates that metformin remodels gut microbiota but the underlying

mechanism remain unclear. Present results showed that metformin effectively

improved insulin sensitivity and alleviated liver inflammation and oxidative stress in

high fat diet (HFD)-induced mice. Metabolomics analysis showed that metformin

increased tauroursodeoxycholic acid (TUDCA) production by increasing the

expression of bile acid synthase *Cyp7a1* and *Baat*. In the palmitic acid (PA)-induced

cell, TUDCA activated Nrf2/ARE pathway, thereby reducing intracellular ROS and

improving insulin signaling. Further gut microbiota analysis showed that metformin

increased the proportion of *Akkermanisia muciniphlia* in the HFD-fed mice, while

TUDCA promoted the proliferation of *A. muciniphlia* but metformin did not. These

findings reveal that metformin remodels the gut microbiota, reduces oxidative stress

and enhances insulin sensitivity by increasing the production of TUDCA. This

provides a novel mechanism by which metformin alleviates diet-induced IR and

improves metabolism.

Keywords: Metformin; Gut microbiota; Insulin resistance; Metabolomics; TUDCA;

Nrf2

Introduction

The prevalence of obesity and overweight have been on the rise worldwide[1,2]. Current predictions indicate that more than 1 billion people will be obese by 2030[3]. Visceral obesity precedes the development of metabolic complications and chronic diseases such as type 2 diabetes (T2D)[4], hepatic steatosis[5] and cardiovascular diseases[6]. Metformin has been the first-line anti-diabetic drug for more than 60 years because of its excellent hypoglycemic effect and safety profile[1,7]. Although several studies have explored the mechanisms by which metformin maintains glucose homeostasis, findings from such studies are inconsistent. The prevailing view is that metformin activates AMP-activated protein kinase (AMPK) in the liver, similar to the principle of caloric restriction that relieves symptoms of diabetes[8,9]. However, metformin is an orally administered drug that reaches high concentrations in the intestine with much lower serum concentrations[10], thus, the possibility that its metabolic therapeutic mechanism might be due in part to actions in the intestine cannot be ignored. Accumulated evidence shows that a high-fat diet (HFD) reduces the abundance of *Lactobacilli* in the small intestine. However, this can be counteracted by metformin, which increases the proportion of probiotics, thus, relieving the symptoms of HFD-induced diabetes[11]. Although metformin may play a hypoglycemic effect by improving the gut microbiota, oral administration of antibiotics to diabetic mice impairs the hypoglycemic ability of metformin[12]. In addition, while metformin can increase the abundance of *Escherichia coli*, it can as well reduce the abundance of *Intestinibacter*. This can further influence diverse biological processes such as the synthesis of branched-chain amino acid and the secretion of GLP-1, thereby improving glucose tolerance and alleviating insulin resistance (IR)[13,14]. These findings suggest that modulation of gut microbiota is involved in the hypoglycemic mechanism of metformin.

Bile acids (BAs), which are produced in the liver and serve as important ingredients of the digestive fluid in the intestine, enter the small intestine through the

biliary system and participate in fat digestion and absorption[15,16]. Gathered evidence

demonstrates that BAs could improve the intestinal homeostasis by modulating the

gut microbiota, thus relieving metabolic syndrome[17]. Recent studies also suggest that

BAs can participate in the liver-gut axis circulation and the improvement of

HFD-initiated metabolic disorders by inhibiting the proliferation of pathogenic

bacteria[18,19]. By modulating the gut microbiota, BAs have also been reported to

induce multiple beneficial effects on the intestinal lumen and intestinal walls such as

the suppression of intestinal inflammatory responses[20,21], the resolution of

endoplasmic reticulum (ER) stress in the intestinal epithelial cells underlying the

pathology of inflammatory bowel disease[22] and the improvement of gut barrier

dysfunction[23-25]. These findings suggest that there is a complex relationship between

BAs, gut microbiota and metabolic syndrome.

Previous studies have reported the association of BAs with oxidative stress,

which has been shown to treat metabolic dysfunction by acting as an endogenous

chemical chaperone to protect cells against ER stress[26]. Intestinal microenvironment

(including inflammation status, the function of the epithelial tight junction and gut

microbiota) plays an essential role in the progression of obese-induced IR. The

presence of gut microbiota and BAs in the intestine may be closely associated with

intestinal metabolic state. Analyses from metagenomic sequencing and metabolomics

show that metformin treatment increases the levels of the conjugated bile acid in the

gut by decreasing the abundance and bile salt hydrolase (BSH) activity of

Bifidobacterium species in the intestines of individuals with T2D[27-31]. BAs are signal

molecules that control the dynamic balance between energy metabolism and liver

protection. It has been reported that BAs change with obesity, IR[32] and nonalcoholic

steatohepatitis (NASH)[25]. However, the mechanism by which BAs alleviate

HFD-induced IR and the underlying role played by the gut microbiota during this

process are not clear.

This study investigated how metformin prevents the development of IR in

HFD-induced obese mice by modulating the dysbiosis of gut microbiota and

alteration metabolites. Exploring the mechanism behind this improves our knowledge

of gut microbiota and metabolic interactions underlying the anti-insulin resistance

effects of metformin.

Methods

Chemicals and diet

Metformin (purity $\geq$ 98%), choline (purity $\geq$ 98%) and taurine (purity $\geq$ 99%)

were obtained from Sigma-Aldrich (St. Louis, MO, USA). Tauroursodeoxycholic acid

(TUDCA, purity $\geq$ 99%) was purchased from Selleck (Houston, Texas, USA). Both

the normal chow diet (NCD, containing 10% fat by energy) and high-fat diet (HFD,

containing 60% fat by energy) were purchased from Beijing HFK Bio-Technology

Co., Ltd. (Beijing, China). Antibodies against Akt, phospho-Akt (Ser473), IRS-1,

phospho-IRS-1 (Ser307), AMPK, phospho-AMPK (Thr172), Keap1, IκB,

phospho-IκB (Ser32), p65, phospho-p65 (Ser536), GAPDH and β-tubulin were

obtained from Cell Signaling Technology (Beverly, MA, USA). Nrf2, phospho-Nrf2

(Ser40), and 4-HNE were purchased from Abcam (Cambridge, UK). Horseradish

peroxidase (HRP)–conjugated secondary antibody was obtained from Santa Cruz

Biotechnology (Santa Cruz, USA).

Animal experiments

Male ICR mice (7 weeks old, body weight 20 ± 1 g) were housed in ventilated

cages (four animals per cage) at the SPF facility of Wenzhou University under

controlled environmental conditions (temperature 22 ± 2°C; relative humidity

60–70%) with free access to standard laboratory chow and tap water. The mice were

maintained on a regular 12/12 h light/dark cycle. All animal care and experimental

procedures were approved by the Animal Care Ministry of Health and were

performed in accordance with the guide for the Care and Use of Experimental

Animals of Wenzhou University.

Mice were acclimatized to their environment for 1 week before the experiments.

Mice were randomly allocated into three groups (n = 9 for each group). A co-worker

blinded to the experimental protocol randomized animals into these groups. One

group of animals was fed with NCD and the other two groups were fed with HFD.

After 12 weeks of HFD feeding, one group of HFD-fed mice was administered

metformin (200 mg/kg body weight), once daily (HFD+Met group) for 6 weeks,

whereas the NCD-fed mice (NCD group) and other group of HFD-fed mice (HFD

group) were treated with an equal volume of saline for 6 weeks. The whole study

lasted 18 weeks, during which the body weight, water consumption and food intake

were measured every week. At week 18, intraperitoneal glucose tolerance test (IPGTT)

and intraperitoneal insulin tolerance test (IPITT) were performed as previously

described[33].

Fresh feces were collected and stored immediately at 80°C for subsequent

analysis. At the end of the trial, after overnight fasting for 12 h, blood samples were

collected, and serum was isolated by centrifugation at $1000 \times g$ for 15 min at 4°C, and

stored at −80°C for further assay. Tissues, including the adipose tissue, liver and

ileum, were weighed; one portion of the tissues was fixed with 10% formaldehyde for

histological analysis, and the other portion was immediately frozen in liquid nitrogen

for further analysis.

Total cholesterol (T-CHO), triacylglycerol (TG), free fatty acids (FFA),

high-density lipoprotein (HDL-c), low-density lipoprotein (LDL-c), fasting blood

glucose (FBG), fasting serum insulin (FINS), catalase (CAT), malondialdehyde

(MDA), aspartate aminotransferase (AST), alanine aminotransferase (ALT), acid

phosphatase (ACP) and alkaline phosphatase (AKP), mice-lipopolysaccharides (LPS)

were determined by biochemical kits purchased from Jiancheng Bioengineering

Institute (Nanjing, China). The ratio of reduced glutathione (GSH) and oxidized

glutathione (GSSG) were determined by a GSH:GSSG kit (Jiancheng Bioengineering Institute, Nanjing, China). Homeostasis model assessment-estimated insulin resistance (HOMA-IR) was calculated using the following formula:

HOMA-IR = FBG (mmol/L) × FINS (mU)/22.5

Hematoxylin and eosin staining

Samples were fixed in 10% formaldehyde overnight, paraffin-embedded, sectioned (4-μm thickness, 3–5 sections/specimen) and stained with hematoxylin and eosin (H & E) for histological analysis. Digital images of H & E stained sections were acquired with a Nikon Eclipse Ti microscope at ×400 magnifications (Ti-E/U/S, Japan). Image J software (National Institutes of Health, USA) was used to count adipocytes.

Cell culture

Human liver hepatocellular carcinoma (HepG2) cells were cultured in MEM (Sigma, St. Louis, MO, USA) supplemented with 10% FBS (Gibco, South American), penicillin and streptomycin. Cells were maintained in 5% CO_2 at 37°C.

siRNA Transfection

Nrf2 was knocked-down by RNA interference (RNAi) using the following 19-bp (including a 2-deoxynucleotide overhang) siRNAs (Origene, Beijing, China): *Nrf2*, SR321100A-AUUGAUGUUUCUGAUCUAUCACUTT; SR321100B-GUCAGUAUGUUGAAUCAGUAGUUTC; SR321100C-CCAGUCUUCAUUGCUACUAAUCAGG. Stealth RNAi (Origene, Beijing, China) was used as a negative control (siCont). For transfection, cells were seeded on a six-well plate, grown to ~80% confluence and transfected with siRNA duplexes using Lipofectamine 3000 (Invitrogen, Camarillo, CA, USA) according to the manufacturer's recommendations. After incubation for 48 h, the expression level of Nrf2 protein was detected by western blot.

Glucose uptake assay

Cells were plated at 1×10^4/well in 96-well plates and used at subconfluence after 24 h of preincubation. For experiments, all culture medium was removed from each well and replaced with 100 µM fluorescent 2-Deoxy-2-[(7-nitro-2,1,3-benzoxadiazol-4-yl) amino]-D-glucose (2-NBDG)[34] in serum-free medium and incubated for 30 min. Subsequently, cells were washed thrice with PBS and then the fluorescence intensity was determined with a fluorescence microplate reader (Ex/Em, 488/520 nm). The glucose concentration in the medium supernatant was determined by a glucose detection kit (Jiancheng Bioengineering Institute, Nanjing, China) following the manufacturer's instructions.

Flow cytometry assays

The reactive oxygen species (ROS) level was evaluated by flow cytometry (FACSCanto II, BD, USA) using the DCFDA cellular reactive oxygen species detection assay kit (Byotime, Shanghai, China) according to the manufacturer's protocol. Cells were briefly stained with 2′,7′-dichlorofluorescein diacetate (DCFA) at 37°C for 30 min, washed with 1× PBS buffer, and the signal was read at an excitation of 485 nm and an emission of 535 nm.

Quantitative RT-PCR

Total mRNA was isolated from tissue samples using TRIzol reagent (TAKARA, Tokyo, Japan) and was reverse-transcribed into cDNA using a high-capacity cDNA reverse transcription kit (TAKARA, Tokyo, Japan) according to the manufacturer's protocol. The mRNA levels were quantified with quantitative PCR (qPCR) using SYBR Green (Qiagen, Hilden, Germany). Amplification was performed on a LightCycler480 qRT-PCR system (Roche, Mannheim, Germany) under the following reactions: 95°C for 15 min, followed by 40 cycles at 95°C for 10 s, 60°C for 20 s, and 72°C for 20 s. The relative mRNA levels of target genes were normalized to the expression of GAPDH calculated using $2^{-\Delta\Delta ct}$ method. The primer pairs used in this

study are listed in Supporting table.

Western blot analysis

Samples were homogenized with ice-cold RIPA lysis buffer containing protease

and phosphatase inhibitors (Beyotime, Shanghai, China). The homogenates were

centrifuged at 10 000 × g for 20 min at 4°C to remove the insoluble tissue debris. The

protein concentration in the supernatant was determined using a BCA protein assay

kit (Beyotime, Shanghai, China) according to the manufacturer's protocol. Equal

amount of protein for each group were denatured in boiling water for 5 min. Aliquots

(40 μg) of protein samples were subjected to 10% SDS-PAGE and transferred to

PVDF membranes (Millipore, Bedford, MA, USA). After blocking with 5% nonfat

milk (dissolved in TBST) for 1 h, the membranes were incubated with the indicated

antibodies at 4°C overnight, followed by incubation with the appropriate

HRP-conjugated second antibodies for 1 h at room temperature. Chemiluminescent

detection was performed using the ECL Plus Western blotting reagent (TransStart,

Beijing, China). Semi-quantitative analysis for densitometry of each band was

performed using ImageJ software.

Analysis of gut microbiota by 16S rRNA amplicon sequencing

As described previously[11], total bacterial DNA was extracted from the collected

feces using CTAB/SDS method. Illumina MiSeq (Novogene Bioinformatics

Technology Co., Ltd.) was used to analyze the 16S rRNA genes in the DNA sample

to determine the composition and diversity of bacteria. Sample processing, data

acquisition and analysis were described in full detail in Supplementary

methodologies.

Non-targeted metabolome analysis

A non-targeted metabolome analysis of the colon contents was performed with a

liquid chromatography-tandem mass spectrometry (LC-MS) method (Waters

ACQUITY UPLC (Waters, America) coupled to a Thermo LTQ Orbitrap XL system

(Thermo, America)). Sample processing, data acquisition, UPLC conditions and data

analysis were described in full detail in the Supplementary methodologies.

Growth curve of *Akkermansia muciniphila*

Akkermansia muciniphila strain ATCC BAA-835 was purchased from the

Beinan (Beijing, China). Bacteria were cultured in BHI medium in tubes at 37°C in an

anaerobic chamber (Whitley A35 Workstation 2.5, Don Whitley Scientific, UK). To

acquire the growth curve of *A. muciniphila*, the medium in test tubes supplemented

with or without 10 mM metformin (or 200 μM TUDCA) was placed in an anaerobic

bag to eliminate oxygen for 24 h. The test tubes were then inoculated with bacteria

and incubated at 37°C in an anaerobic bag. The OD_{600} of the cultures was measured

every 3 hours.

Determination of hepatic TUDCA

The contents of TUDCA in hepatic tissues were determined by

high-performance liquid chromatography (HPLC). Briefly, 200 mg of frozen tissue

was homogenized with a TissueLyserII (Qiagen, Germantown, MD, USA) in 1 mL of

PBS to prepare the tissue homogenates. The impurities were removed through a 0.22

μm filter, then the filtrate was precipitated by methanol and 10 μL of supernatant was

analyzed using an Agilent 1290 HPLC system (Santa Clara, CA) equipped with a

Hypersil ODS-2 column (5 μm, 4.6×250 mm; Waters, USA). The supernatant was

detected at the wavelength of 210 nm, eluted with the mobile phase of 0.03 M

phosphate buffer solution (pH 4.4) and methanol (32:68, v/v) at a flow rate of 1.0

mL/min.

Statistical analysis

Statistical comparisons among experimental groups were analyzed by one-way

ANOVA and Duncan's multiple-comparison test using the SPSS software (Version

21.0. Armonk, NY: IBM Corp.). *P* values less than 0.05 were considered statistically

significant.

Results

1. Metformin prevents diet-induced obesity and increases insulin sensitivity

To assess the efficacy of metformin on the treatment of obese mice with IR,

animals were divided into the NCD, HFD and HFD + Met groups. Daily oral

administration of metformin significantly alleviated diet-induced body weight

increase from day 14 onwards (**Figure 1A**). At the end of metformin treatment, the

weight gain and BMI were also significantly decreased (**Figure 1B, Supplementary**

Figure 1a), without affecting food intake (**Figure 1C**). Metformin prevented fat

deposition in the subcutaneous adipose, the adipose index (**Supplementary Figure**

1b) and H & E staining of adipose tissue (**Figure 1D).** Adiposity numbers showed

that metformin effectively ameliorated adipocyte hypertrophy (**Figure 1E**). The

indexes of the pancreas and kidney were not affected by metformin administration.

The liver index was significantly reduced in metformin-treated mice (**Supplementary**

Figure 1b). The diet-induced hepatic steatosis and dyslipidemia were effectively

prevented by metformin treatment as indicated by the lower concentration of T-CHO,

TG and FFA in both the liver and the serum of the HFD + Met group compared with

the HFD group (**Figure 1 F-H**). We also found that metformin could effectively

prevent hepatocyte damage. This was evidenced by the lower levels of AST, ALT,

ACP and AKP in the serum (**Supplementary Figure 2**).

Glycemic homeostasis was determined by IPGTT and IPITT. As shown in

Figure 2A, the oral administration of glucose increased the blood glucose (BG) level

within 30 min in the HFD-fed mice, which remained high for the subsequent 90 min.

This indicated that HFD feeding markedly impaired glucose tolerance. In addition, the

BG of the NCD group peaked within 30 min but dropped to the normal level after 120

min. Metformin significantly inhibited the rise of BG level and enhanced glucose

clearance in the HFD-fed mice after oral administration of glucose. The results of

IPGTT were expressed as the area under the curve (AUC) of BG over 120 min. The

AUC of the HFD + Met group was significantly decreased compared with that of the

HFD group (**Figure 2B**, $P < 0.05$). The improvement of insulin resistance (IR) was

also supported by the results of IPITT. After intraperitoneally injected insulin at 0.75

U/kg, the BG levels at 30, 60, and 120 min revealed that the mice in the HFD + Met

group took more advantage of insulin compared with the mice in the HFD group

(**Figure 2C**). The significant difference ($P < 0.05$) of the slope between the HFD

group and the HFD + Met group suggested that metformin increased the utilization of

insulin and improved insulin sensitivity in the HFD-fed mice (**Figure 2D**). Moreover,

the FBG level in the HFD group was significantly higher compared to that in the

NCD group. Compared with the HFD group, the mice treated with metformin had a

lower FBG level (**Figures 2E**). Marked elevation in fasting serum lisulin (FINS) was

observed in the HFD group compared with the NCD group. Notably, metformin

treatment tended to reverse the increase in FINS (**Figure 2F**). HOMA-IR is an

indicator for evaluating the level of insulin resistance in an individual. Based on the

HOMA-IR analysis, we found that the HOMA-IR of the HFD + Met group was

comparable to that of the NCD group but significantly lower compared with that of

the HFD group (**Figure 2G**).

PI3K/Akt and insulin receptor substrate-1 (IRS-1) are the main signaling

molecules in the signaling pathway of the insulin receptor. The level of

phosphorylated Akt at Ser473 was significantly reduced in the liver of the

HFD-induced obese mice, but the level was restored in the HFD + Met group (**Figure**

2H). Contrary to the phosphorylated Akt, the level of phosphorylated IRS-1 at Ser307

was significantly elevated in the HFD group, whereas in the HFD + Met group, the

level was reduced. These results indicate that the preventive effect of metformin on

HFD-induced obesity resulted in improved glucose homeostasis and insulin

sensitivity.

2. Metformin alleviates HFD-induced inflammation and oxidative stress

As shown in **Figure 3A** and **3B**, metformin prevented metabolic inflammation in the liver and ileum, leading to an increased tendency towards reduced gene expression of interleukin-1β (*Il-1β*), interleukin-6 (*Il-6*) and tumor necrosis factor-α (*Tnf-α*) in the ileum. However, there   was no effect on interleukin-10 (*Il-10*). Metformin also prevented the release of endotoxin, which was indicated by lower levels of LPS in serum (**Figure 3C**). Meanwhile, the phosphorylated levels of p65 and IκB were decreased in the livers of metformin-treated mice compared with the HFD-fed mice (**Figure 3D**). This indicated that the HFD-induced activation of NF-κB signaling was suppressed by metformin.

To investigate the involvement of oxidative stress in the development of HFD-induced IR, we first measured the activity of the antioxidant enzyme in both the plasma and the liver. The results showed that the activities of CAT in the serum and the liver were markedly lower in the HFD group compared with the NCD group (**Figure 4A**). In addition, the ratio of GSH/GSSG was reduced (**Figure 4C**). After metformin intervention, the activities of CAT and the ratio of GSH/GSSG were significantly increased. The accumulation of lipid peroxide, caused by HFD, was improved in the metformin group (**Figure 4B**). This was exhibited by the significantly reduced MDA level in the liver. Nrf2/ARE is a classic antioxidant-signaling pathway and KEAP1 is a key protein that mediates the degradation of Nrf2. HFD feeding significantly inhibited Nrf2/ARE signaling (**Figure 4D**) whereas metformin significantly increased the expression of Nrf2. Nrf2/ARE signaling regulates the expression of a variety of detoxification and antioxidant enzymes. As shown in **Figure 4E**, Metformin significantly up-regulated the mRNA levels of *Nqo1* and *Ho1*, which are essential genes for balancing oxidative stress. These results indicate that metformin can improve intestinal and hepatic inflammation, oxidative stress and lipid peroxide accumulation in HFD-fed mice.

3. Metformin prevents obesity-driven dysbiosis of gut microbiota and reshapes intestinal metabolites

Faecal samples were collected for gut microbiota analysis by 16S rRNA amplicon sequencing. Following all quality trimming and checking, a dataset consisting of 80004, highly qualified reads, was collected for subsequent analysis. Our data showed that the numbers of unique OTUs in the NCD, HFD, and HFD + Met groups were 372, 350 and 374, respectively. The coverage of sequencing results was nearly complete for all sequences in the three groups, demonstrating sufficient sequencing depth for follow-up analyses. The metformin-treated group significantly increased the Ace and Chao1 indices compared with the HFD group (**Table 1**). Metformin-treated mice, however, clustered partially apart from HFD-fed mice samples, suggesting important changes in metformin gut microbial profile (**Figure 5A**). Metformin treatment prevented HFD-induced decrease in Bacteroidetes and the increase in the Firmicutes/Bacteroidetes ratio (**Figure 5B** and **C**), two hallmarks of obesity-driven dysbiosis. Based on FunGuild and FAPROTAX analysis of flora function, the HFD group showed a decrease in the proportion of the chemoheterotrophy and the fermentation bacteria when compared with the NCD group (**Figure 5D**). However, after metformin supplementation reshaped the structure of gut microbiota, chemoheterotrophy and fermentation improved in the HFD + Met group. Heatmap presents the clustering of bacterial communities with their relative abundances at the genus level (**Figure 6A)**. Compared with the NCD group, unique high-abundance bacteria in the HFD group are mainly concentrated in *Bifidobacterium, Roseburia, Lachnoclostridum, Alloprevotella* and *Mucispirilum*. However, metformin decreased the abundance of these bacteria, making the metformin-treated group comparable to the NCD group. In addition, metformin specifically increased the abundance of *Butyricimonas, Parasulterella, Parabacteroides* and *Akkermansia*. Consistent with the decrease in F/B ratio as shown in **Figure 5C**, the difference between the HFD group and the HFD + Met group is

mainly reflected in the decrease in the abundance of Firmicutes and the increase in the abundance of Bacteroidetes (**Figure 6B**). Compared with the NCD and HFD + Met groups, *c_Erysipelotrichia, f_proteobacteria, g_Turicibacter, o_Bifidobacteriales* and *s_Clostridium* in the HFD group were significantly up-regulated, whereas f_*Muribaculaceae, g_parabacteroides* and *f_tannerellaceae* were significantly up-regulated in the HFD + Met group. Therefore, metformin can increase the abundance of Bacteriodia at the phylum level and *Parabacteroides* at the genus level, and modulate HFD-induced dysbiosis of gut microbiota.

The intestinal metabolite profiles were further measured using lipid chromatography and mass spectrometry (LC-MS) to reveal the influences of metformin on the intestinal metabolites. When comparing the HFD group and HFD + Met group, metabolites were dictated by metformin. For instance, in the HFD group, 118 up-regulated metabolites were identified, whereas 182 down-regulated metabolites were established in the HFD + Met group (**Figure 7A-C**). The HFD + Met group effectively increases the content of taurine, choline and TUDCA in the intestine compared with the HFD group (**Figure 7D**). This is consistent with previous reports[18]. Collectively, these results showed that metformin administration altered the intestinal metabolites and the gut microbiota. **Figures 8A** and **8B** showed the correlation between intestinal metabolism and the gut microbiome at the phylum and species level after the intervention of metformin in HFD-fed mice. TUDCA was highly associated with *A. muciniphila* species belonging to the *Verrucomicrobia* phylum. Interestingly, through the analysis of growth curves, it was not metformin but TUDCA significantly accelerated the proliferation of *A. muciniphila in vitro* (**Figure 8**C). These data suggest that metformin may indirectly improve insulin resistance caused by a high-fat diet through modulation of the structure of gut microbiota and metabolites.

4. TUDCA improves PA-induced insulin resistance and lipid peroxidation

Further study was carried out to determine whether TUDCA could alleviate

hepatic insulin resistance using a PA-treated HepG2 cell model. PA exposure (200

μM) resulted in a decrease in glucose uptake in HepG2 cells. This was demonstrated

by the reduction of fluorescence density in the cytoplasm (**Figure 9A**) and the high

concentration of glucose in the medium (**Figure 9B**). In addition, PA decreased the

level of phosphorylated Akt at Ser473 and increased phosphorylated IRS-1 level at

Ser307, thus, revealing a typical symptoms of insulin resistance (**Supplementary**

Figure 3). As shown in **Figure 9A** and **9B**, the impaired glucose uptake caused by PA

was significantly alleviated by TUDCA, excluding choline and taurine

(**Supplementary Figure 4**). Moreover, the suppressed phosphorylation level of Akt

in PA-treated HepG2 cells was elevated by TUDCA, while the phosphorylated IRS-1

level at Ser307 was also significantly reduced (**Figure 9C**). Note that, TUDCA

treatment at high doses may produce toxic side effects and fail to exert insulin

sensitization. These findings confirmed that TUDCA could improve hepatic glucose

intolerance and enhance insulin sensitivity of PA-treated hepatocyte in a

metformin-independent manner.

Homeostasis imbalances on glucose metabolism often cause inflammation and

oxidative stress damage in hepatic cells. Thus, we further investigated whether

TUDCA could alleviate these pathological processes. The intracellular ROS content

was visualized by using the DCF-DA dye and flow cytometric platform. **Figure 10A**

showed that PA could increase ROS generation in the HepG2 cells. Interestingly,

TUDCA significantly decreased PA-induced ROS generation. On the other hand, the

ratio of GSH/GSSG was significantly increased by TUDCA (**Figure 10B**). In addition,

pro-inflammatory cytokines such as *Tnf-α*, *Il-1β* and *Il-6* were also significantly

reduced by TUDCA treatment at mRNA level (**Supplementary Figure 5**).

We further investigated whether TUDCA alleviates PA-induced insulin

resistance through the Nrf2/ARE signaling pathway. The transcriptional activity of

Nrf2 was determined by measuring the mRNA levels of its representative downstream

target genes. TUDCA significantly up-regulated the expression of Nrf2, suppressed by PA (**Figure 10D**). It also up-regulated the expression of antioxidant genes, *Ho-1*, *Nqo1*, *Cat* and *Gpx4*, downstream of the Nrf2/ARE signaling pathway (**Figure 10D**). We found that TUDCA was unable to alleviate PA-induced insulin resistance in HepG2 cells that interfered with Nrf2 expression (**Supplementary Figure 6**, **10E & F**). Together, the results suggest that TUDCA may significantly activate the Nrf2/ARE signaling, thereby enhancing antioxidant capacity, and alleviating insulin resistance in the HepG2 cells caused by PA.

To further determine the source of TUDCA, we detected the TUDCA content in the liver of the HFD and the HFD + Met groups. As shown in **Figure 11A**, TUDCA in the liver of the HFD + Met group (5.281 ± 0.745 nmol/g) was significantly increased compared with that in the HFD group (2.992 ± 0.691 nmol/g). In the HFD + Met group, the gene expression of *Cyp7a1* and *Baat*, the important enzymes involved in the biosynthesis of TUDCA, was significantly up-regulated compared with that in the HFD group (**Figure 11B**). This suggests that metformin can promote the production of TUDCA in the liver by up-regulating the enzymes that limit the rate of TUDCA biosynthesis, and *Cyp7a1* and *Baat* expression.

Discussion

Metabolic syndromes such as obesity, nonalcoholic fatty liver disease and T2D are increasingly prevalent globally, especially in recent decades[2]. Metformin, the most widely used anti-diabetic drug worldwide, has been reported to favorably influence metabolic and cellular processes. Metformin is also shown to be closely associated with the development of insulin resistance[35,36]; however, the mechanism is still not clear. Recent studies have revealed that the development of metabolic disease strongly correlates to the accumulation of gut microbiota and metabolites. Multiple studies have investigated the impact of metformin on the gut microbiota, metabolism and related host targets on the regulation of glucose metabolism[37,38]. We explored the

potential mechanism involving the gut microbiota and intestinal metabolites to reveal

how metformin improves diet-induced obesity and insulin resistance. We established

that the daily administration of metformin strongly ameliorated the obesity-related

interruption of gut microbiota. The bile acid TUDCA was maintained at a high level

in the intestine, which in turn promoted the proliferation of *A. muciniphila*, improved

the integrity of the intestinal wall and reduced intestinal inflammation. Furthermore,

TUDCA was identified as an activator of Nrf2/ARE signaling in hepatocyte that

decreased oxidative stress imbalance, while improving hepatic inflammation and

insulin resistance in HFD-induced obese mice.

T2D is both an inflammatory and a natural immune disease. There is a variety of

microbiota in the human intestine, which is involved in body immunity, infection and

nutrient metabolism, and they are closely associated with the occurrence of diabetes.

The gut microbiota imbalance in composition and diversity can result in the release of

bacterial endotoxin. This could lead to chronic inflammation and destruction of the

normal physiological functions of tissues and organs. Several studies have confirmed

that chronic inflammation can cause insulin resistance and subsequently diabetes. The

decrease in Firmicutes/Bacteroidetes ratio is considered to be an indicator of

improved metabolism by metformin therapy[39,40]. LDA analysis showed that the

decreased abundance of Firmicutes by metformin could be due to the decrease in

Erysipelotrichia species, which belongs to the Firmicutes phylum. The role of

Erysipelotrichia species in human disease may be well established in studies

investigating metabolic disorders[41]. For instance, a previous study showed a bloom of

species belonging to the *Erysipelotrichaceae* family (initially classified as *Mollicutes*)

in diet-induced obese animals[42]. Within the host, *Erysipelotrichia* species appear to be

highly immunogenic[43], however, in surplus, they induce intestinal inflammation.

Studies have shown that species in *Erysipelotrichaceae* are positively correlated with

TNFα and IL-6 in the intestine and the liver[44].

In contrast, most of the bacteria up-regulated by metformin were probiotics. For

example, species in *Muribacaceae* are involved in the L-citrulline nitrogen metabolic

cycle. They scavenge free radicals, improve immunity and maintain blood glucose.

Metformin also up-regulates the abundance of *Parabacteroides*, including *P.*

distasonis and *P. goldsteinii*. Liu et al.[45] found that *P. distasonis* could transform bile

acids, activate intestinal gluconeogenesis through succinic acid and improve the host

metabolic disorders. In another study, Wu et al.[46] recorded that *P. goldsteinii* reduces

systemic inflammation and increases insulin sensitivity by maintaining intestinal

integrity. *Akkermansia muciniphila* is known to effectively alleviate intestinal

inflammation caused by obesity. Patients with intestinal obesity have been shown to

have a significantly reduced abundance of *A. muciniphila* compared with healthy

individuals. Numerous studies have demonstrated that metformin could up-regulate

the abundance of *A. muciniphila* in HFD-fed mice intestine, maintain the integrity of

the intestinal wall as well as effectively improve intestinal inflammation[47]. The

abundance of *A. muciniphila* is also significantly upregulated in the HFD + Met group.

As a mucin-degrading bacterium, the use of *A. muciniphila* has been shown to reduce

inflammation, maintain the intestinal barrier and increase levels of endocannabinoids

secreted by intestinal peptides[48]. We speculate that the transformation of gut

microbiota by metformin is the defining factor in improving chronic inflammation in

the intestine and even the whole body. Metformin can effectively inhibit the number

of pathogenic bacteria and promote the proliferation of probiotics. A suppression of

pathogenic bacteria coupled with a proliferation of probiotics may synergistically

reduce intestinal inflammation and maintain the integrity and permeability of the

intestinal wall. Furthermore, this may lead to a decrease in serum LPS and

improvement of systemic inflammation caused by HFD-induced obesity.

The "dialogue" between the gut microbiota and the organism mainly depends on

microbial metabolites. By producing various enzymes for biochemical metabolic

pathways of the intestinal microbes, gut microbiota can perform diverse metabolic

activities such as the metabolism of amino acids, carbohydrates and bile acids, as well

as the formation of a co-metabolic relationship with the host. Bile acids produced by

the gut microbiota are essential in the regulation of glycolipids and energy. Studies

have shown that in human primary hepatocytes, bile acids promote the β-oxidation of

fatty acids by activating FXR[18,49] and up-regulate the expression of apolipoprotein CII

gene to reduce the triglyceride content in the liver[21].

As a conjugated bile acid derivative, TUDCA has been shown to treat

nonalcoholic fatty liver disease (NAFLD) by acting as an endogenous chemical

chaperone; to protect cells against ER stress[26,50,51] and to reduce liver steatosis[52].

Moreover, TUDCA can alleviate dextran sulfate sodium-induced colitis in mice[53,54].

In vitro, we found that TUDCA could act as an agonist of the Nrf2/ARE signaling

pathway. This is reminiscent of the expression pattern of Nrf2, a crucial stress

regulator to reduce the ROS, which is a key factor in sensitizing insulin signaling[55-57].

It has also been confirmed that up-regulation of the Nrf2 expression both by genetic

manipulation and pharmacological interference, can significantly decrease the content

of ROS and ameliorate IR and T2D[58,59]. In the mice supplement with metformin, the

significant up-regulation in the expression of *Nrf2* and its downstream target genes

(*Gclc*, *Cat* and *Ho-1*) occurred, which further alleviated IR and over-oxidation.

Conversely, the stress-resistant effect of TUDCA was dramatically repressed by the

inhibition of the Nrf2/ARE signaling. *In vitro* experiments have found that TUDCA

have similar effects, which can effectively reduce the inflammation and oxidative

stress caused by PA as well as decreasing the accumulation of lipid peroxide. This

might be a novel mechanism for metformin in the treatment of insulin resistance and

diabetes.

In our present study, metabolomics data showed that TUDCA in the intestinal

contents of the HFD + Met group increased significantly. Previous studies revealed

that metformin treatment increased the levels of the conjugated bile acid in the gut by

decreasing the abundance of species of *Bifidobacterium* and its bile salt hydrolase

(BSH) activity in the intestines of individuals with T2D[17,30,31]. Bile salt hydrolase,

secreted by intestinal bacteria, decomposes TUDCA in the intestine and metabolizes

TUDCA to taurine and UDCA. However, data from the gut microbiota showed that

metformin did not affect the abundance of *Bifidobacterium* and *Lactobacillus*, the

major BSH-producing bacteria. In addition, we found that the content of taurine in the

intestine increased significantly. *In vitro*, TCDCA can be transformed into TUDCA

by 7α/β-hydroxysteroid dehydrogenase (7α/β-HSDH)[15,35,60]. Data from the gut

microbiota also revealed that metformin did not affect the abundance of *Eubacterium*

and *Clostridium*, the major 7α/β-HSDH-producing bacteria. Therefore, the increase in

TUDCA in the intestine is purportedly not due to the inhibitory effect of metformin

on BSH-producing bacteria. Further analysis of the biosynthetic site of bile acids and

the liver confirmed that the conjugated bile acid TUDCA was generated in the liver

through a series of catalytic reactions and enzymes such as CYP7A1 and BAAT.

CYP7A1 is a rate-limiting enzyme that catalyzes the breakdown of cholesterol into

bile acids in the liver[15,61] whereas BAAT is a vital enzyme that catalyzes the

amidation reaction of taurine and UDCA[46,62]. Interestingly, we found that the gene

expressions of *Cyp7a1* and *Baat* were significantly enhanced in the HFD + Met group,

which might further promote the synthesis of TUDCA in the liver. HPLC analysis

also showed an increase in the content of TUDCA in the livers of the HFD + Met

group. We suppose that the increase in TUDCA biosynthesis may be the primary

cause of the increase in TUDCA in the intestine.

Overall, in our present study, metformin was found to increase the bile acid

TUDCA production by increasing bile acid synthase *Cyp7a1* and *Baat.* As an

activator of the Nrf2/ARE signaling pathway, TUDCA promotes Nrf2

phosphorylation at Ser40. It also increases the gene transcription of its downstream

antioxidant enzyme, reduces oxidative stress and sensitizes insulin signaling; thereby

alleviating insulin resistance caused by HFD. The increase of TUDCA levels in the

intestine corresponded to the increase in the abundance of *A. muciniphlia.* Metformin

remodels the gut microbiota by promoting probiotics such as *Muribacaceae,* and *Parabacteroides*, and inhibiting pathogenic bacteria such as *Erysipelotrichia* and *Helicobacter*. This maintains intestinal integrity and reduces inflammation induced by HFD. However, for a better understanding of the crucial mechanisms involved in the long-term treatment with metformin in humans, the role of the intestine-liver axis should be explored. Since *A. muciniphlia* and/or TUDCA are the novel target of metformin in the gut, they are potential targets for the treatment of metabolic conditions in humans.

Conflict of Interest

The authors declare no conflict of interest.

Author Contributions

Y.Z., Y.C., L.Y and J.Z. performed the research. H.T., Y.G. and M.W. designed research. Y.Z., Q.L. and S.W. provided technical assistance. H.T. and Y.Z. analyzed the data. Y.Z. and H.T. wrote the paper. H.T., Y.G. and M.W. had primary responsibility for final content. All authors have read and approved the final manuscript.

Acknowledgements

We thank Chujun Lin, Yufeng Wu, Lingfeng Hou, Huijing Jiang (Wenzhou University) for their assistance in animal experiments. We thank Alan K Chang (Wenzhou University) for helpful discussion and for revising the language of the manuscript.

Funding

This work was financially supported by the National Natural Science Foundation of China (41876197 and 81872952), the National Key Research and Development Project (2018YFD0901503), the Natural Science Foundation of Zhejiang Province (LY18C020006), the Science and Technology Program of Wenzhou (ZY2019013 and Y20180210).

References

1. Muoio DM, Newgard CB. Mechanisms of disease:Molecular and metabolic mechanisms of insulin resistance and beta-cell failure in type 2 diabetes. *Nat. Rev.Mol. Cell Bio.* 2008; 9, 193-205. doi: 10.1038/nrm2327

2. Igel LI, Saunders KH, Fins JJ. Why weight? An analytic review of obesity management, diabetes prevention, and cardiovascular risk reduction. *Cur. Athero. Repor.* 2018; 20: 39. doi: 10.1007/s11883-018-0740-z

3. Abarca-Gómez L, Abdeen ZA, Hamid ZA, Dika Z, Ivkovic V, Jelakovic A, Jelakovic B, Kos J, Jureša V, Majer M, et al. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million children, adolescents, and adults. *Lancet.* 2017; 390: 2627-2632. doi: 10.1016/s0140-6736(17)32129-3

4. Shen J, Obin MS, Zhao L. The gut microbiota, obesity and insulin resistance. *Mol. Aspec. of Medic.* 2013; 34(1):39-58. doi: 10.1016/j.mam.2012.11.001

5. Bäckhed F, Manchester JK, Semenkovich CF, Gordon JI. Mechanisms underlying the resistance to diet-induced obesity in germ-free mice. *P Natl. Acda. Sci. USA.* 2007;104(3):979-984. doi: 10.1073/pnas.0605374104

6. Cho NH, Shaw JE, Karuranga S, Huang Y, Rocha Fernandes JD, Ohlrogge AW, Malanda B. IDF diabetes atlas: global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes. Res. Clin. Pract.* 2018; 138: 271-281. doi: 10.1016/j.diabres.2018.02.023

7. Graham R. Hardie DG. Pearson RE. The mechanisms of action of metformin. *Diabetologia.* 2017; 60:1577-1585. doi: 10.1007/s00125-017-4342-z

8. Zhou G, Myers R, Li Y, Chen Y, Shen X, Fenyk-Melody J, Wu M, Ventre J, Doebber T, Fujii N, Musi N, Hirshman MF, Goodyear LJ, Moller DE. Role of AMP-activated protein kinase in mechanism of metformin action. *J Clin Invest.* 2001; 108:1167-74. doi: 10.1172/JCI13505

9. Hawley SA, Ross FA, Chevtzoff C, Green KA, Evans A, Fogarty S,
Towler MC, Brown LJ, Ogunbayo OA, Evans AM, Hardie DG. Use of cells ex-
pressing gamma subunit variants to identify diverse mechanisms of AMPK
activation. *Cell Metab.* 2010; 11:554-565. doi: 10.1016/j.cmet.2010.04.001

10. Lee H, Ko G. Effect of metformin on metabolic improvement and gut
microbiota. *Appl. Enviro. Micorb.* 2014; 80:5935-5943. doi:
10.1128/AEM.01357-14

11. Zhang X, Zhao Y, Xu J, Xue ZS, Zhang MH, Pang XY, Zhang XJ, Zhao LP.
Modulation of gut microbiota by berberine and metformin during the treatment of
high-fat diet-induced obesity in rats. *Sci Rep.* 2015; 5:14405. doi:
10.1038/srep14405

12. Zhou ZY, Ren LW, Zhan P, Yang HY, Chai DD, Yu ZW. Metformin exerts
glucose-lowering action in high-fat fed mice via attenuating endotoxemia and
enhancing insulin signaling. *Acta Pharmacologica Sinica.* 2016; 37: 1063-75. doi:
10.1038/aps.2016.21

13. Kristoffer F, Falk H, Trine N, Gwen F, Emmanuelle LC, Shinichi S, Edi P,
Sara VS, Valborg G, Helle KP, Manimozhiyan A, Karsten K, Anita YV, Henrik
V, Rajna H, Paul IC, Jens RK, Junhua L, Torben J, Florence L, Joël D, Søren B,
Jeroen R, Torben H, Jun W, Dusko ES, Peer B, and Oluf P. Disentangling the
effects of type 2 diabetes and metformin on the human gut microbiota. *Nature.*
2015; 528(7581): 262–266. doi: 10.1038/nature15766

14. Pedersen HK, Gudmundsdottir V, Nielsen HB, Hyotylainen T, Nielsen T,
Jensen BA, Forslund K, Hildebrand F, Prifti E, Falony G, Le Chatelier E,
Levenez F, Doré J, Mattila I, Plichta DR, Pöhö P, Hellgren LI, Arumugam M,
Sunagawa S, Vieira-Silva S, Jørgensen T, Holm JB, Trošt K; MetaHIT
Consortium, Kristiansen K, Brix S, Raes J, Wang J, Hansen T, Bork P, Brunak S,
Oresic M, Ehrlich SD, Pedersen O. Human gut microbes impact host serum
metabolome and insulin sensitivity. *Nature.* 2016; 535(7612):376-381. doi:

10.1038/nature18646

15. Jia W, Xie G, Jia W. Bile acid-microbiota crosstalk in gastrointestinal

inflammation and carcinogenesis. *Nat. Rev. Gastroenterol Hepatol.*

2018;15(2):111-128. doi: 10.1038/nrgastro.2017.119

16. Schroeder BO and Backhed F. Signals from the gut microbiota to distant

organs in physiology and disease. *Nat. Med.* 2016; 22: 1079-1089. doi:

10.1038/nm.4185

17. Guo JL, Han X, Tan HY, Huang WD, You YL, Zhan JC. Blueberry extract

improves obesity through regulation of the gut microbiota and bile acids via

pathways involving FXR and TGR5. iScience. 2019; 19:676-690. doi: 10.1016 /

j.isci.2019.08.020

18. Sun LL , Xie C, Wang G, Wu Y, Wu Q, Wang XM, Liu J, Deng YY, Xia JL,

Bo C, Zhang SY, Chuyu Y, Lian G, Zhang XJ, Zhang H, Bisson WH, Shi JM,

Gao XX, Ge PP, Liu CH, Krausz WK, Nicholis RG, Cai JW, Rimal BP, Patterson

A, Wang X, Gonzalez FJ, Jiang CT. Gut microbiota and intestinal FXR mediate

the clinical benefits of metformin. *Nat Med.* 2018; 24(12): 1919-1929. doi:

10.1038/s41591-018-0222-4

19. Lee H, Lee Y, Kim J, An J, Lee S, Kong H, Song Y, Lee CK, Kim K.

Modulation of the gut microbiota by metformin improves metabolic profiles in

aged obese mice. *Gut Microbes.* 2018; 9(2): 155-165. doi:

10.1080/19490976.2017.1405209

20. Martínez-Moya P, Romero-Calvo I, Requena P, Hernández-Chirlaque C,

Aranda CJ, González R, Zarzuelo A, Suárez MD, Martínez-Augustin O, Marín JJ,

de Medina FS. Dose-dependent antiinflammatory effect of ursodeoxycholic acid

in experimental colitis. *Int Immunopharmacol.* 2013; 15(2): 372–380. doi:

10.1016/j.intimp.2012.11.017.

21. Renga B, Mencarelli A, Cipriani S, D'Amore C, Carino A, Bruno A,

Francisci D, Zampella A, Distrutti E, Fiorucci S. The bile acid sensor FXR is

required for immune-regulatory activities of TLR-9 in intestinal inflammation. *PLoS One.* 2013; 8(1): e54472. doi: 10.1371/journal.pone.0054472

22. Berger E, Haller D. Structure-function analysis of the tertiary bile acid TUDCA for the resolution of endoplasmic reticulum stress in intestinal epithelial cells. *Biochem Biophys Res Commun.* 2011; 409: 610–615. doi: 10.1016/j.bbrc.2011.05.043

23. Stenman LK, Holma R, Gylling H, Korpela R. Genetically obese mice do not show increased gut permeability or faecal bile acid hydrophobicity. *Br J Nutr.* 2013; 110: 1157–64. doi: 10.1017/S000711451300024X

24. Arab JP, Karpen SJ, Dawson PA, Arrese M, Trauner M. Bile acids and nonalcoholic fatty liver disease: Molecular insights and therapeutic perspectives. *Hepatology.* 2017; 65: 350–362. doi: 10.1002/hep.28709

25. Wang WJ , Zhao JF, Gui WF, Sun D, Dai HJ, Xiao L, Chu HK, Fan D, Zhu QJ, Bernd S, Huang K, Yang L, and Hou XH. Tauroursodeoxycholic acid inhibits intestinal inflammation and barrier disruption in mice with non-alcoholic fatty liver disease. *British J of Pharmaco.* 2018; 175: 469–484. doi: 10.1111/bph.14095

26. Xie Q, Khaoustov VI, Chung CC, Sohn J, Krishnan B, Lewis DE, Yoffe B. Effect of tauroursodeoxycholic acid on endoplasmic reticulum stress-induced caspase-12 activation. *Hepatology.* 2002; 36(3): 592–601. doi: 10.1053/jhep.2002.35441

27. Fondevila C, Shen XD, Tsuchiyashi S, Yamashita K, Csizmadia E, Lassman C, Busuttil RW, Kupiec-Weglinski JM, Bach FH. Biliverdin therapy protects rat livers from ischemia and reperfusion injury. *Hepatology.* 2004; 40: 1333–1341. doi: 10.1002/hep.20480

28. Nakao A, Otterbein LE, Overhaus M, Sarady JK, Tsung A, Kimizuka K, Nalesnik MA, Kaizu T, Uchiyama T, Liu F, Murase N, Bauer AJ, Bach FH, Biliverdin protects the functional integrity of a transplanted syngeneic small

bowel. *Gastroenterology.* 2004; 127: 595–606. doi: 10.1053/j.gastro.2004.05.059

29. Ester GS, Maria JP, Nikolaj SN, Oscar B, Maria JM, Elisa L, Maria AS, Jose

JGM. Protective role of biliverdin against bile acid-induced oxidative stress in

liver cells. *Free Radical Biology and Medicine.* 2016; 97: 466–477. doi:

10.1016/j.freeradbiomed.2016.06.016

30. Zhang M, Feng R, Yang M, Qian C, Wang Z, Liu W, Ma J. Effects of

metformin, acarbose, and sitagliptin monotherapy on gut microbiota in Zucker

diabetic fatty rats. *BMJ Open Diabetes Res Care.* 2019; 7(1):e000717. doi:

10.1136/bmjdrc-2019-000717

31. Nova E, Pérez de Heredia F, Gómez-Martínez S, Marcos A. The Role of

Probiotics on the Microbiota: Effect on Obesity. *Nutr Clin Pract.* 2016;

31(3):387-400. doi: 10.1177/0884533615620350

32. Zhang WP , Chen LH , Feng H, Wang W, Cai Y, Qi F, Tao XF, Liu J, Shen

YJ, Ren XF, Chen X, Xu JM, Shen YX. Rifampicin-induced injury in HepG2

cells is alleviated by TUDCA via increasing bile acid transporters expression and

enhancing the Nrf2- mediated adaptive response. *Free Radical Biology and*

Medicine. 2017; 112: 24–35. doi: 10.1016/j.freeradbiomed.2017.07.003

33. Navarrete J, Vásquez B, Del SM. Morphoquantitative analysis of the Ileum

of C57BL/6 mice (Mus musculus) fed with a high-fat diet. *Int J Clin Exp Pathol.*

2015; 8(11):14649-14657.

34. Zou CH, Wang YJ, Zhen ZF. 2-NBDG as a fluorescent indicator for direct

glucose uptake measurement. *J. Biochem. Biophys. Methods.* 2005; 64: 207–215.

doi: 10.1016/j.jbbm.2005.08.001

35. Lien F, Berthier A, Bouchaert E, Gheeraert C, Alexandre J, Porez G, Prawitt

J, Dehondt H, Ploton M, Colin S, Lucas A, Patrice A, Pattou F, Diemer H, Van

Dorsselaer A, Rachez C, Kamilic J, Groen AK, Staels B, Lefebvre P. Metformin

interferes with bile acid homeostasis through AMPK-FXR crosstalk. *J. Clin.*

Invest. 2014; 124(3): 1037–1051. doi: 10.1172/JCI68815.

36. Bailey CJ, Wilcock C and Scarpello JH. Metformin and the intestine. *Diabetologia.* 2008; 51: 1552–1553. doi: 10.1007/s00125-008-1053-5

37. Canfora EE, Meex R, Venema K, and Blaak EE. Gut microbial metabolites in obesity, NAFLD and T2DM. *Nat. Rev. Endocrinol.* 2019; 15: 261–273. doi: 10.1038/s41574-019-0156-z

38. Wu H, Esteve E, Tremaroli V, Khan MT, Caesar R, Holm LM, Sthlman M, Olsson L, Serino M, Planas-Fèlix M, Xifra G, Mercader JM, Torrents D, Burcelin R, Ricart W, Perkins R, Fernàndez-Real JM, Bäckhed F. Metformin alters the gut microbiome of individuals with treatment-naive type 2 diabetes, contributing to the therapeutic effects of the drug. *Nat Med.* 2017; 23 (7) :850-868. doi: 10.1038/nm.4345

39. Nova E, Pérez de Heredia F, Gómez-Martínez S, Marcos A. The Role of Probiotics on the Microbiota: Effect on Obesity. *Nutr Clin Pract.* 2016; 31(3): 387-400. doi: 10.1177/0884533615620350

40. Sircana A, Framarin L, Leone N, Berrutti M, Castellino F, Parente R, Michieli FD, Paschetta E, Musso G. Altered gut microbiota in type 2 diabetes: just a coincidence? *Curr. Diab. Rep.* 2018; 18(10): 98. doi: 10.1007/s11892-018-1057-6

41. Kaakoush NO. Insights into the Role of Erysipelotrichaceae in the Human Host. *Front Cell Infect Microbiol.* 2015; 20; 5: 84. doi: 10.3389/fcimb.2015.00084

42. Turnbaugh PJ, Backhed F, Fulton L, and Gordon JI. Marked alterations in the distal gut microbiome linked to diet-induced obesity. *Cell Host Microbe.* 2008; 3(4): 213–223. doi: 10.1016/j.chom.2008.02.015

43. Craven M, Egan CE, Dowd SE, McDonough SP, Dogan B, Denkers EY. Inflammation drives dysbiosis and bacterial invasion in murine models of ileal Crohn's disease. *PLoS ONE.* 2012; 7:e41594. doi: 10.1371/journal.pone.0041594

44. Nguyen TL, Vieira-Silva S, Liston A, and Raes J. How informative is the

mouse for human gut microbiota research? *Dis. Model Mech.* 2015; 8: 1–16. doi: 10.1242/dmm.017400.

45. Tsung-Ru Wu, Chuan-Sheng Lin, Chih-Jung Chang Tzu-Lung Lin, Jan Martel, Yun-Fei Ko, David M Ojcius, Chia-Chen Lu, John D Young,1 Hsin-Chih Lai. Gut commensal *Parabacteroides goldsteinii* plays a predominant role in the anti-obesity effects of polysaccharides isolated from *Hirsutella sinensis*. *Gut* 2018;0:1–14. doi: 10.1136/gutjnl-2017-315458

46. Manley S, Ding WX. Role of farnesoid X receptor and bile acids in alcoholic liver disease. *Acta Pharmaceutica Sinica B.* 2015;5(2):158–167. doi: 10.1016/j.apsb.2014.12.011

47. Anhê FF, Roy D, Pilon G, Dudonne S, Matamoros S, Varin T, Marette A. A polyphenol-rich cranberry extract protects from diet-induced obesity, insulin resistance and intestinal inflammation in association with increased *akkermansia spp*. population in the gut microbiota of mice. *Gut.* 2015; 64(6): 872–883. doi:10.1136/gutjnl-2014-307142

48. Rodriguez J, Hiel S, Delzenne NM. Metformin: old friend, new ways of action-implication of the gut microbiome? *Curr Opin Clin Nutr Metab Care.* 2018; 21(4): 294-301. doi: 10.1097/MCO.0000000000000468

49. Everard A, Belzer C, Geurts L, Ouwerkerk JP, Druart C, Bindels LB, Guiot Y, Derrien M, Muccioli GG, Delzenne NM, de Vos WM, Cani PD. Cross-talk between *Akkermansia muciniphila* and intestinal epithelium controls diet-induced obesity. *Proc Natl Acad Sci USA.* 2013; 110(22):9066-9071. doi: 10.1073/pnas.1219451110

50. Dempsey JL, Wang D, Siginir G, Fei Q, Raftery D, Gu H, Yue Cui J. Pharmacological Activation of PXR and CAR Downregulates Distinct Bile Acid-Metabolizing Intestinal Bacteria and Alters Bile Acid Homeostasis. *Toxicol Sci.* 2019;168(1):40-60. doi: 10.1093/toxsci/kfy271

51. Wahlstrom A, Sayin SI, Marschall HU, Backhed F. Intestinal crosstalk

between bile acids and microbiota and its impact on host metabolism. *Cell Metab.* 2016; 24, 41–50. doi: 10.1016/j.cmet.2016.05.005

52. Li F, Jiang CT, Krausz KW, Li YF, Albert I, Hao HP, Fabre KM, Mitchell JB, Patterson A, Gonzalez FJ. Microbiome remodelling leads to inhibition of intestinal farnesoid X receptor signalling and decreased obesity. *Nat. Commun.* 2013; 4: 2384. doi: 10.1038/ncomms3384

53. Choi YJ, Shin HS, Choi HS, Park JW, Jo I, Oh ES, Lee KY, Lee BH, Johnson R, Kang DH. Uric acid induces fat accumulation via generation of endoplasmic reticulum stress and SREBP-1c activation in hepatocytes. *Lab Invest.* 2014; 94(10): 1114–1125. doi: 10.1038/labinvest.2014.98

54. Legry V, Van Rooyen DM, Lambert B, Sempoux C, Poekes L, Español-Suñer R, Molendi-Coste O, Horsmans Y, Farrell GC, Leclercq IA. Endoplasmic reticulum stress does not contribute to steatohepatitis in obese and insulin-resistant high-fat-diet-fed foz/foz mice. *Clin Sci (Lond).* 2014; 127(7):507-518. doi: 10.1042/CS20140026

55. Cao YN, Xue Y, Xue L, Jiang XL, Wang XL, Zhang ZJ, Yang J, Lu JL, Zhang CX, Wang WQ, Ning G. Hepatic menin recruits SIRT1 to control liver steatosis through histone deacetylation. *J Hepatol.* 2013; 59(6): 1299-1306. doi: 10.1016/j.jhep.2013.07.011

56. Fazakerley DJ, Chaudhuri R, Yang PY, Maghzal GJ, Thomas KC, Krycer JR, Humphrey SJ, Parker BL, Fisher-Wellman KH, Meoli CC, Hoffman NJ, Diskin C, Burchfield JG, Cowley MJ, Kaplan W, Modrusan Z, Kolumam G, Yang KYL, Chen DL, Samocha-Bonet D, Greenfield JR, Hoehn JL, Stocker R, James DE. Mitochondrial CoQ deficiency is a common driver of mitochondrial oxidants and insulin resistance. *eLife* . 2018; 7:e32111. doi: 10.7554/eLife.32111

57. Hancock M, Hafstad AD, Nabeebaccus AA, Catibog N, Logan A, Smyrnias L, Hansen SS, Lanner J, Schröder K, Murphy MP, Shah MA, Zhang M. Myocardial NADPH oxidase-4 regulates the physiological response to acute

808 exercise. *eLife*. 2018;7:e41044. doi: 10.7554/eLife.41044

809 58. Hu H, Fan X, Guo Q, Wei X, Yang D, Zhang B, Liu J, Wu Q, Oh Y, Feng Y,

810 Chen K, Hou L, Gu N. Silicon dioxide nanoparticles induce insulin resistance

811 through endoplasmic reticulum stress and generation of reactive oxygen species.

812 *Part Fibre Toxicol.* 2019;16(1):41. doi: 10.1186/s12989-019-0327-z

813 59. You L, Yang C, Du Y, Liu Y, Chen G, Sai N, Dong X, Yin X, Ni J. Matrine

814 Exerts Hepatotoxic Effects via the ROS-Dependent Mitochondrial Apoptosis

815 Pathway and Inhibition of Nrf2-Mediated Antioxidant Response. *Oxid Med Cell*

816 *Longev.* 2019; 2019:1045345. doi: 10.1155/2019/1045345

817 60. Takafumi S, Shiori S, Keiichiro H, Eriko N, Eri HK, Ayaka Y, Liam B,

818 Nobuyuki T, Hiroshi S, Masayuki Y. Hyperactivation of Nrf2 in early tubular

819 development induces nephrogenic diabetes insipidus. *Nat Commun.* 2017;

820 8:14577. doi: 10.1038/ncomms14577

821 61. Manuel M, Katrin F, and Elke HH. Activation of Nrf2 signaling by natural

822 products-can it alleviate diabetes? *Biotechnol Adv.* 2018; 36(6): 1738–1767. doi:

823 10.1016/j.biotechadv.2017.12.015

824 62. Annika SA, Emily T, Brig M, Shaji C, Hannah AN, Yunzhao T, Jed WF,

825 Jonathan MJD, Claes BW, Nils W, Morey WH, Stephen HF, Hindrik M, Anders

826 HR. Sulforaphane reduces hepatic glucose production and improves glucose

827 control in patients with type 2 diabetes. *Sci Transl Med.* 2017; 9(394). pii:

828 eaah4477. doi: 10.1126/scitranslmed.aah4477

 Figure captions

Figure 1. Metformin prevents obesity and systemic metabolic disorders in obese

mice induced by HFD. Boy weight (A), weight gain (B), food intake (C), H&E

staining of epididymal adipose tissue (D, 400×), number of adipocytes (E), levels of

total cholesterol and triacylglycerol in liver (F) and serum (G), and serum FFA (H)

are shown. Data are expressed as the mean ± SD. $*P < 0.05$, $**P < 0.01$ and $***P <$

0.001.

Figure 2. Metformin improves glucose homeostasis and insulin sensitivity in

diet-induced obese mice. IPGTT (A); AUC of IPGTT (B); IPITT (C); slope of IPITT

(D) measured at the end of the experimental period. Serum level of FBG (E), FINS

(F), and HOMA-IR (G). Protein expression of Akt, phosphorylated Akt, IRS-1,

phosphorylated IRS-1 Ser307 and quantitative analysis for the densitometry of

pIRS-1 and p-Akt protein level(H). Data are expressed as mean ± SD. $*P < 0.05$, $**P$

< 0.01 and $***P < 0.001$.

Figure 3. Metformin blunts intestinal and liver inflammation. Determination of

inflammatory factors in liver (A) and ileum (B) by qPCR and serum LPS (C) by kit.

Expression levels of IκB, phosphorylated IκB, P65, and phosphorylated P65 as

determined by western blot and quantitative analysis for the densitometry of p-IκB

and p-P65 protein level (D). Data are expressed as mean ± SD. $*P < 0.05$, $**P < 0.01$

and $***P < 0.001$.

Figure 4. Metformin ameliorates HFD-induced oxidative stress. Representative

markers of oxidative stress level: CAT (A), MDA (B), and GSH:GSSG ratio in the

liver (C). Protein expression of total Nrf2, phosphorylated Nrf2, total AMPK, and

phosphorylated AMPK; quantitative analysis for the densitometry of the p-Nrf2 and

p-AMPK protein (D). Quantitative RT-PCR analysis of *Nrf2, Keap1* and

representative downstream genes (*Nqo1* and *Ho-1*) (E). Data are expressed as mean ±

SD. $*P < 0.05$, $**P < 0.01$ and $***P < 0.001$.

Figure 5. Metformin remodels fecal microbial structure and communities.

Microbial composition with cluster at the phylum level (A); relative abundance of gut

microbiota at the phylum level (B); ratio of Bacteroidetes to Firmicutes (C); function

analysis of gut microbiota (D). Data are represented as mean ± SD. *$P < 0.05$ and **P

< 0.01.

Figure 6. Genus-based comparison of microbial similarities and differences.

Heatmap showing clustering of bacterial communities with their relative abundances

at the genus level (A). The bacterial communities or OTUs were clustered

phylogenetically by neighbor-joining method and the libraries were clustered by

profile pattern using Unifrac analysis. Relative abundances (log values) of microbial

genus are displayed in a red-to-blue color code (high to low abundance). Linear

discriminant analysis (LDA) score for taxa differing between treatment groups (B).

LDA scores with threshold > 3 indicated a higher relative abundance in the

corresponding group than in the other two groups.

Figure 7. Differential metabolites determined by non-targeted metabolomics.

Metabolite clustering heat map (A); OPLS-DA score plot (B), in the OPLS-DA score

plot, each data point represents one sample, and the distance between points in the

plot indicates the similarity between samples; number of different metabolites

between each group (C), the major upregulated differential metabolites (D).

Figure 8. Correlation heatmap for gut microorganisms and metabolites. Matrix

heatmap showing the correlation between gut microbiota and metabolome at Phylum

(A) and Species (B) levels. Growth curve of *A. muciniphila* in the presence of

metformin or TUDCA (C).

Figure 9. TUDCA alleviates insulin resistance caused by PA. Glucose uptake (A)

and the blood glucose remaining in supernatant (B). Protein levels of p-Akt, Akt,

pIRS-1 Ser307, IRS-1 were determined by western blot (C). Data are expressed as the

mean ± SD. *$P < 0.05$, **$P < 0.01$ and ***$P < 0.001$.

Figure 10. TUDCA relieves PA-induced insulin resistance via activating
Nrf2/ARE signaling pathway. Intracellular ROS content (A); the ratio of
GSH/GSSG (B); Protein levels of p-Nrf2 and Nrf2 were determined by western blot
(C); Quantitative RT-PCR analysis of *Nrf2*, representative Nrf2 downstream genes
Ho-1 and *Nqo1* (D); Under Nrf2 knock-down, glucose uptake (E) and blood glucose
in the medium supernatant (F) are shown. Data are expressed as mean ± SD. *$P <$
0.05, **$P < 0.01$ and ***$P < 0.001$.

Figure 11. Metformin increases hepatic TUDCA. The content of TUDCA in the
liver was determined using HPLC (A), and the relative mRNA levels of bile acid
synthetase (*Cyp7a1*, *Cyp27a1*, *Baat*) determined by qPCR (B).

Table 1. Diversity of gut microbiota in HFD and HFD+Met treated mice

Groups	Reads	OTUs	Observed species	Ace	Chao1	Shannon	Simpson
NCD	78316±4945	372±35	330±21	345±17	350±21	5.10±0.67	0.92±0.04
HFD	76154±7745	350±40	312±20	325±24	326±24	5.38±0.33	0.95±0.02
HFD+Met	80004±421	374±21	335±20*	342±22*	348±20*	5.24±0.27	0.95±0.02

896　　Data are expressed as mean ± SD.

897　　∗ Indicates significantly different from the HFD group at the *P* < 0.05 level.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Figure 11