Local chromatin context dictates the genetic determinants of the heterochromatin spreading reaction.

R.A. Greenstein[1,2,†], Henry Ng[1,2,†], Ramon R. Barrales[3,#], Catherine Tan[5,6], Sigurd Braun[3,4], and Bassem Al-Sady[1,*]

[1]Department of Microbiology &Immunology, George Williams Hooper Foundation, University of California San Francisco, San Francisco, CA 94143, USA, [2]TETRAD graduate program, University of California San Francisco, San Francisco, CA 94143, USA, [3], Biomedical Center, Department of Physiological Chemistry, Ludwig-Maximilians-University of Munich, Planegg-Martinsried, Germany, [4]International Max Planck Research School for Molecular and Cellular Life Sciences, Martinsried, Germany, [#]current address: Centro Andaluz de Biología del Desarrollo, Universidad Pablo de Olavide de Sevilla-Consejo Superior de Investigaciones Científicas-Junta de Andalucía, Sevilla, Spain. [5]Department of Cell and Tissue Biology, University of California San Francisco, San Francisco, CA 94143, USA, [6]Biomedical Sciences graduate program, University of California San Francisco, San Francisco, CA 94143, USA

† these authors contributed equally to this work.

[*]correspondence: bassem.al-sady@ucsf.edu

Abstract

Heterochromatin spreading, the expansion of gene-silencing structures from DNA-encoded nucleation sites, occurs in distinct chromatin contexts. Spreading re-establishes gene-poor constitutive heterochromatin every cell cycle, but also invades gene-rich euchromatin *de novo* to steer fate decisions. Unlike heterochromatin nucleation and assembly, the determinants of the spreading process remain poorly understood. Our heterochromatin spreading sensor separately records nucleation site-proximal, and distal, heterochromatin gene silencing. By screening a nuclear function gene deletion library in fission yeast, we identified regulators that alter the propensity, both positively and negatively, of a nucleation site to spread heterochromatin. Critically, the involvement of many regulators is conditioned by the chromatin context within which spreading occurs. We find spreading, but not nucleation, within constitutive heterochromatin, requires distinct Clr6 histone deacetylase complexes. However, spreading is universally antagonized by a suite of chromatin remodelers. Our results disentangle the machineries that control lateral heterochromatin spreading from those that instruct DNA-directed assembly.

Introduction

Cellular specification requires the genome to be partitioned into regions of activity and inactivity such that only genes appropriate for a given cellular state are available for expression. This requires the formation and propagation, in time and space, of gene-repressive heterochromatin structures. Heterochromatin is most commonly seeded by DNA-directed nucleation (Hall et al. 2002; Reyes-Turcu et al. 2011), and then propagates across the chromosome by a DNA-sequence indifferent process termed spreading, to repress genes expression in the underlying regions. Silencing is instructed by assembly factors, such as HP1, that recognize heterochromatic chemical modifications (Jacobs et al. 2001; Lachner et al. 2001). The spreading of silencing structures occurs in very different chromatin contexts, which may intrinsically promote or antagonize this process: (1) Constitutive heterochromatin, which is generally gene-poor and as such depleted of activities associated with active genes known to antagonize heterochromatin (Scott et al. 2006; Greenstein et al. 2019). Its maintenance through replication is aided by the inheritance of nucleosomes bearing heterochromatic marks (Alabert et al. 2015). This inheritance promotes modification of nearby nucleosomes due to the "read-write" positive feedback intrinsic to heterochromatin histone modifiers (Zhang et al. 2008; Al-Sady et al. 2013; Ragunathan et al. 2015). (2) Conversely, during differentiation, heterochromatin is either seeded at new nucleation sites or invades gene-rich euchromatin *de-novo* from existing nucleation sites (Wen et al. 2009; Zhu et al. 2013; Zylicz et al. 2015; Zylicz et al. 2018; Nicetto and Zaret 2019). In either case, it encounters active chromatin that lacks repressive marks and can specifically antagonize heterochromatin (Greenstein et al. 2019). Thus, during this initial invasion process, heterochromatin spreading cannot benefit from the inheritance through replication of pre-existing marked nucleosomes. Beyond the differences between active and inactive chromatin, it remains unclear whether distinct nucleation elements require different regulators

to enact efficient spreading outward from those sites. Nonetheless, we recently described that distinct

nucleation elements trigger divergent types of spreading behavior (Greenstein et al. 2018).

Over the past four decades, forward and reverse genetic screens in fission yeast have established

an exhaustive list of factors required for heterochromatin silencing and heterochromatin nucleation.

Some of these nucleation mechanisms include repeat sequences that instruct specialized RNAi-

machinery to process noncoding (nc) RNAs involved in targeting the histone methyl transferase Clr4

(Moazed 2009); signals within nascent transcripts that trigger heterochromatin island formation (Zofall

et al. 2012); and shelterin-dependent pathways (Wang et al. 2016; Zofall et al. 2016). However, those

genetic screens have neither directly addressed factors that specifically regulate the spreading process,

nor whether such factors act in general or context-specific settings. With our previously established

fluorescent reporter-based heterochromatin spreading sensor (HSS) we can segregate the central output

of heterochromatin (gene silencing) from the spatial control of the reaction (spreading) (Greenstein et al.

2018; Greenstein et al. 2019). This allows us to address the following questions: (1) Are there known or

novel regulators of heterochromatin that primarily regulate spreading, versus nucleation? (2) Does

spreading over chromatin with distinct characteristics, such as gene density or nucleosome arrangement,

require different sets of regulators? (3) Does the set of regulators required for efficient spreading depend

on the type of nucleator that seeds it - for example nucleators using transcription and ncRNA pathways

versus direct tethering of heterochromatic factors?

To address these questions, we conducted a series of reverse genetic screens in fission yeast,

using a custom nuclear function gene deletion library in four heterochromatin contexts. These include

derivations of the fission yeast mating type (MAT) locus, a gene-poor constitutive heterochromatin

region contained by *IR-L* and *IR-R* boundaries and nucleated by at least two elements: (1) *cenH,*

homologous to pericentromeric *dh* and *dg* elements, which rely on ncRNA pathways, including RNAi,

and (2) *REIII*, a sequence element that directly recruits heterochromatin factors via the stress-response transcription factors Atf1 and Pcr1 (Jia et al. 2004; Kim et al. 2004). We also queried an ectopic heterochromatin domain that is embedded in gene rich euchromatin. This domain is nucleated by an ectopically inserted *dh* element fragment proximal to the *ura4* locus (Canzio et al. 2013; Marina et al. 2013; Greenstein et al. 2018).

We find that the genetic requirements for promotion and containment of heterochromatin spreading diverge significantly between different chromatin contexts, and to some degree also between different types of nucleators. However, despite these context-dependent differences for spreading, common themes also emerge from this work: (1) Sub-complexes of the Clr6 histone deacetylase (HDAC) complex, in particular the Fkh2-associated Clr6 complex I″, appear to promote spreading, but not nucleation, at multiple heterochromatin loci. (2) At both euchromatic and heterochromatic loci, spreading is antagonized by a diverse set of nucleosome remodelers, in particular Ino80 and Swr1C. (3) Unexpectedly, members of an AP3 adaptor complex, normally involved in membrane traffic, are moderate positive regulators of spreading in hetero- and euchromatin. Together, these results specifically pinpoint the machineries required to regulate the spatial expansion of heterochromatin domains independent of the initial seeding by DNA-encoded nucleation sites.

Results

Our previously developed HSS relies on three transcriptionally-encoded fluorescent protein-coding genes that collectively report single cell measurements of heterochromatin formation via flow cytometry, while normalizing for transcriptional and translational noise (Al-Sady et al. 2016; Greenstein et al. 2018). It provides separate, quantitative recordings of nucleation-proximal ("green") and distal ("orange") events at a heterochromatin site over large populations of isogenic cells (typically N >20,000) (**Figure 1A**). In contrast to the singular readout employed by auxotrophy-dependent reporter

gene silencing assays, the HSS assay provides a "multidimensional snapshot" that distinguishes changes

in heterochromatin nucleation and spreading, and additionally permits tracking of emerging multimodal

cell populations and unique population distributions. This is an important conceptual advance compared

to traditional methods.

With this ability to observe the entire population distribution pattern from any isogenic

background, we sought first to investigate the requirements for heterochromatin activity for both

nucleation and distal spreading, within either constitutive heterochromatin or gene-rich euchromatic

regions. Further, we queried whether different types of heterochromatic nucleation sequences utilize

similar or different sets of regulators to nucleate and/or spread the resulting heterochromatin structures

outwards. To address the latter, we explored three different chromatin contexts, derived from the

constitutive heterochromatic mating type (MAT) locus, each containing an embedded HSS (**Figure 1A**)

(Greenstein et al. 2018): wild type, with the *cenH* and *REIII* nucleating DNA elements uncompromised,

and two *MAT* variants that contained mutations in either the *cenH* or *REIII* elements (**Figure 1A**).

Mutations in these DNA elements limit initiation of heterochromatin spreading to one nucleator

(Greenstein 2018). To address differences in chromatin context in addition to variants of the constitutive

MAT locus, we examined heterochromatin formation at the euchromatic *ura4* locus*,* where

heterochromatin spreading is ectopically driven by the upstream insertion of a pericentromeric *dh* DNA

element (Marina et al. 2013; Greenstein et al. 2018). We refer to this chromatin context as *ECT*

(ect̲opic). When analyzed by flow cytometry, *WT MAT* and *MAT ΔcenH* populations appear largely

fully nucleated with near-complete local spreading, as evidenced by population density in the bottom

left in the 2D density histogram (**Figure 1C**,**D** (Greenstein et al. 2018)). *MAT ΔREIII* and *ECT* cell

populations, while mostly nucleated, display a stochastic distribution of spreading states, evidenced by a

vertical distribution on the left of the 2D density histogram (**Figure 1E,F** (Greenstein et al. 2018)). The

distribution of cells in a strong loss of silencing mutant, *Δclr3,* is shown for all chromatin contexts next

to the wild-type parents (**Figure 1C-F**). To identify potential regulators of the spreading reaction, we

conducted a genetic screen by crossing a deletion library of ~400 nuclear function genes (**Table 1**,

Figure 1 Supplement 1) to reporter strains with the HSS integrated in the four chromatin contexts

described above. We quantified the fluorescence signal from the three reporters in each mutant by flow

cytometry, with approximately 20k-30k cells per mutant, depending on growth conditions.

To visualize the pattern of heterochromatin domain activity across the isogenic mutant

populations, we first employed t-distributed stochastic neighbor embedding (t-SNE). For all the mutants

in each chromatin context, we converted the normalized 2D distribution of cells into a numerical vector

representing cell density in 25 isometric plot regions as different parameters (**Figure 1B**). The t-SNE

model was built by calculating the similarity between different cell distributions and comparing these 25

parameters from all mutants and wild type controls in the four chromatin contexts (**Figure 1B**).

We plotted the t-SNE model with all the four chromatin contexts and respective mutants,

coloring each mutant by its parental chromatin context (**Figure 1G**). We also colored the mutants by the

median nucleation ("green", **Figure 1H**) or spreading ("orange", **Figure 1I**) reporter expression values.

As expected, the majority of mutants did not strongly deviate from their parent and broadly clustered

together into a "neighborhood" by chromatin context. This is evident for *ECT*, for which the parent

strain has "green" and "orange" in a less repressed state, particularly compared to *WT MAT* and *MAT*

ΔcenH. Within each neighborhood, the distribution of "orange" expression, especially for *MAT ΔREIII*

and *ECT*, is graded from above to below the expression level of the parent(s), revealing a continuum of

mutants with enhanced or abrogated spreading. We could not find mutants that display more repression

than the parental strains of *MAT ΔcenH,* which are highly repressed in the OFF state, as previously

described (Grewal and Klar 1996; Greenstein et al. 2018). However, we did observe mutants located

out of the area of their chromatin context-driven "neighborhood". First, the major known

heterochromatin mutants, *Δclr4, Δswi6, Δclr3* among others, from each chromatin context formed a

cluster with high expression of "green" and "orange" (**Figure 1I** enlarged region, exemplified by *Δclr3*),

segregating from the rest of the population. Second, we observed mutants, such as *Δcdt2, Δepe1* and

Δchp1, that segregate out of neighborhood only for selected chromatin contexts, indicating specificity

(highlighted in **Figure 1I**). The t-SNE analysis visualized the relationship of all four chromatin contexts,

and mutants therein, with respect to their nucleation and spreading behavior, directly revealing the

graded nature of mutant phenotypes. This is particularly the case with respect to spreading in *ECT* and

MAT ΔREIII neighborhoods (**Figure 1I**)

However, in the t-SNE analysis the phenotype patterns are weighted by the intrinsic behavior of

each parent's chromatin context. To quantify how much each mutation impacted the heterochromatin

state in each chromatin context, we performed Earth Mover's Distance analysis (EMD, **Figure 1J** see

also materials and methods and (Orlova et al. 2016)). We express the contribution of each mutant

relative to the parental isolates by a quotient of their respective EMDs to *Δclr4*, which is completely

deficient in heterochromatin assembly and serves as a fixed reference point for all chromatin contexts.

We represent a subset of the mutants' EMD values in a heatmap (**Figure 1K**). Some mutants contributed

similarly in different chromatin contexts, such as *Δclr3* and *Δrik1,* which display strong de-repression,

and *Δair1,* which displays an intermediate de-repression (Thon et al. 1999; Keller et al. 2010). However,

many mutants had differential contributions in each chromatin context. As an example, *Δpob3,* a

mutation in the FACT complex (Lejeune et al. 2007), had weaker phenotypes where nucleation is solely

ncRNA-driven (*MAT ΔREIII* and *ECT*) and stronger phenotypes when *REIII,* an ncRNA-independent

element, is intact (*WT MAT* and *MAT ΔcenH*) (**Figure 1K**). Interestingly, the TRAMP subunit mutant

Δcid14, had more pronounced roles in *MAT* contexts over *ECT* (**Figure 1K**). Beyond these more subtle

differences, we noticed a large divergence between *ECT* and other contexts. This is evident from very

low pairwise correlations of $EMD^{mut/par}$ values between all the mutants for *ECT* against the *MAT*

contexts (**Figure 1, Supplement 2 C,E,F**). Conversely, we observe much higher correlations between

all pairwise combinations of the three *MAT* chromatin contexts (**Figure 1, Supplement 2 A,B,D**). The

divergence between *MAT* and *ECT* contexts is especially true for the RNAi pathway. *Δchp1* and *Δtas3*

(members of RITS complex), *Δdsh1* (RNAi factor), *Δcid12* (member of RDRC complex), and *Δsaf1*

(RNAi associated splicing factor (Bayne et al. 2014)) all have very strong effects only in *ECT.*

The above results are consistent with previous reports of *WT MAT* and ectopic reporters (Hall et

al. 2002), which are respectively independent and dependent on RNAi for heterochromatin maintenance.

Surprisingly, *MAT ΔREIII,* which similar to *ECT* solely relies on a *dh*-homologous *cenH* element for

H3K9me nucleation, also behaved independent of RITS for maintenance (**Figure 1K**). *MAT* has another

H3K9me-independent silencing element, *REII* (Hansen et al. 2011), which is located upstream of

"green". However, this element appears to act quite locally (Ayoub et al. 2000; Hansen et al. 2011).

Moreover, we showed removal of *REII* had no effect on expression of HSS reporters in *MAT ΔcenH*

(Greenstein et al. 2018). Thus, we do not believe *REII* accounts for the difference observed between

MAT ΔREIII and *ECT*.

Similar to RNAi factors, the heterochromatin-antagonizing *epe1* pathway had strikingly different

impacts on the *MAT* chromatin contexts compared to *ECT* (**Figure 1G**). *Δepe1* results in destabilized

heterochromatin at *ECT,* but the deletion has little effect at *MAT* chromatin contexts. Heterochromatin in

these contexts instead is sensitive to mutations in the Epe1-turnover pathway, exemplified by *cdt2*, the

specificity factor for the Epe1-targeting E3 ubiquitin ligase (Braun et al. 2011). In contrast to loss of

silencing mutants, we also identified mutations that induce hyper-repressed states ($EMD^{mut/par} > 1$) in

multiple chromatin contexts with some differences in contribution to hyper-repression. These included

Δyap9, Δsnf22, Δswc5 and *Δiec3* (see below).

Overall, the EMD analysis enabled us to systematically and quantitatively discern how each

mutant contributes to the combined nucleation and spreading heterochromatin state within each

chromatin context. The results suggest that the genetic circuitry for heterochromatin maintenance differs

significantly between naïve sites in euchromatin and constitutive loci. This is consistent with the tightly

repressed *MAT ΔcenH* or the redundantly nucleated *WT MAT*. But the fact that *MAT ΔREIII* also

diverged strongly from *ECT* in its dependence on genes for heterochromatin nucleation and spreading

was more surprising, because both contexts are driven exclusively by a similar ncRNA nucleators, *cenH*

and *dh* (Hansen et al. 2006), and display stochastic spreading over similar ranges (**Figure 1C, F**).

Further we previously showed that they both exhibit dynamically unstable heterochromatin over time

(Greenstein et al. 2018). The finding that even heterochromatin domains with very similar nucleation

and spreading dynamics rely on different regulators for their maintenance, suggests that such context-

specific requirements for heterochromatin assembly are likely common across the genome.

The above analyses addressed the genetic requirements for heterochromatin silencing, including

both nucleation and distal spreading, in different chromatin contexts. Next, we aimed to identify

regulators that are specific to heterochromatin spreading but function independent of nucleation. To do

so, we employed the capability of the HSS to segregate those activities. We isolated cell populations that

reside within a "green"-off gate which represents cells with heterochromatin fully assembled at the

nucleation site (see methods, and (Greenstein et al. 2018)). To quantify increased or decreased spreading

in a given mutant, we calculated a $\text{Grid}_n^{\text{mut/par}}$ metric (described in Methods), which tracks the changes

of cell distributions in "orange" expression within the "green"-off gate (**Figure 2A**). To isolate gain or

loss of spreading mutants (hits) for further analysis, we only considered the top 15% of mutants in which

Grid$_n^{mut/par}$ values were also above 2 standard deviations from the mean of the replicate parent isolates.

(**Figure 2 Supplement 1**). With these gene hits isolated, we proceeded to analyze their relationships

within and across chromatin contexts.

We first examined the degree to which spreading modulators are shared between chromatin

contexts via upset plots (**Figure 2B&C**). Conceptually similar to a Venn diagram, this analysis allows

rapid visualization of the degree of overlap between sets, with the number of shared genes plotted as a

bar graph and the sets each bar represents annotated below the plot. The upset plot for loss of spreading

phenotypes (*i.e.* genes that promote spreading, **Figure 2B**) showed that exceedingly few genes, three out

of 164 unique genes found as hits, are shared across all chromatin contexts (*csn1, rrp17, apm3*). This

result emphasizes the specific impact that each chromatin context has on heterochromatin spreading.

Seven genes were shared across all the MAT locus chromatin contexts. In contrast, 111 genes

contributed only to one chromatin context. The degree to which genes contributed positively towards

spreading, and the degree of overlap across chromatin contexts is shown in bar-graphs of Grid$_n^{mut/par}$ in

Figure 2D-G. We additionally show the 2D density histogram of the screen mutants for the top two loss

of spreading hits for each chromatin context in **Figure 2H-K**. The top two hits for *MAT ΔcenH* were

gad8 and *cdt2* (**Figure 2E&I**) The significant contribution of *cdt2* suggests that Epe1specifically limits

spreading in *MAT ΔcenH* compared to other chromatin contexts. Spreading in *WT MAT* and *MATΔREIII*

was most dependent on *fkh2,* while the second strongest hits were *rrp6* and *prw1,* respectively (**Figure**

2D,F,H,J and see below). Prw1 and Fkh2 form parts of Clr6 complexes (see below), while Rrp6 is a

central part of the exosome, required for gene silencing in multiple ncRNA-dependent pathways (Buhler

et al. 2007). The top hit in *ECT* was the Csn1 COP9 signalosome subunit, which has been implicated in

neddylation of cullin-based E3 ligases and may operate in a similar pathway as Cdt2 (Bayne et al. 2014).

ECT was also highly sensitive to *Δpht1* (**Figure 2G**) which codes for H2A.Z, normally associated with

antagonizing spreading (Meneghini et al. 2003).

We also observed a significant number of genes that showed gain of spreading in *WT MAT, MAT*

ΔREIII and *ECT* (**Figure 2C**). We could not examine *MAT ΔcenH* for this phenotype because this

chromatin context is highly repressed in the OFF state as reported previously (Grewal and Klar 1996;

Greenstein et al. 2018). Six out of 98 genes found as hits are shared across these three chromatin

contexts (**Figure 2C**, **Figure 2 Supplement 2**). The 2D density histograms of the screen mutants are

shown for top two hits per chromatin context (**Figure 2 Supplement 2D-F**). Validating the approach,

we found *leo1,* a gene previously impacted in spreading control across boundaries (Verrier et al. 2015),

as a moderate gain of spreading hit. *ECT* displayed the largest fraction, 43 out of 58 of spreading-

antagonizing genes that are unique to one context. In contrast, we found 11 out of 43 genes unique to

MAT ΔREIII, which displays a very similar spatio-temporal spreading behavior to *ECT.* This likely

reflects that even though heterochromatin can assemble in a euchromatin context, spreading is under

multiple layers of constraint in this setting.

To understand which nuclear pathways, as opposed to individual genes, direct or antagonize

heterochromatin spreading across backgrounds, we performed a protein complex-level analysis. Using

the Gene Ontology (GO) protein complex annotations from Pombase (Lock et al. 2018), we annotated

the GO complex membership of each screen hit and tabulated the frequency ("counts") of each GO

complex per background in both loss of spreading (loss) and gain of spreading (gain) categories. Using

these GO complex counts, we generated a heatmap for all the protein complexes represented by hits in

our screen (**Figure 3A**). In order to assess the similarities between chromatin contexts (*WT MAT, MAT*

ΔcenH, MAT ΔREIII, ECT) and categories (loss or gain of spreading) we performed unsupervised

clustering on the heatmap columns. Broadly, the hit categories (loss or gain of spreading) clustered

together, (**Figure 3A**, bottom), revealing relative similarity at the GO complex level. The exception

however was the *ECT* "loss" set of hits. Consistent with our gene-level analysis (**Figure 2B-G, Figure 2**

Supplement 2), complexes that contribute most strongly to "loss" and "gain" categories are different in

ECT compared to the *MAT* variants.

Overall, we identified three common trends within and between categories: (1) a role for

antagonizing spreading by chromatin remodelers, (2) a role of histone deacetylase complexes (HDACs),

in particular Clr6 sub-complexes, in promoting spreading, and (3) a role for a small number of additional

known and novel spreading regulators, including the AP-3 adaptor and COP9 signalosome complex,

which promoted spreading across all backgrounds.

As evidenced by the heatmap (**Figure 3A, top right**), chromatin remodeling complexes are

strongly represented in the gain of spreading hit category, including the Swr1C, Ino80, SWI/SNF, and

RSC-type complexes. To explore this further, we assessed which protein components were contributing

to these GO complex counts. For all the genes annotated to a given chromatin remodeling complex and

present in our screen, we displayed whether they were identified as a hit (blue) or not (grey) (**Figure**

3B). Indeed, we found that the large majority of the gene hits annotated fall within the "gain" category

across backgrounds. The manner in which the cell distributions change in these screen hits is evident

from the 2D density histograms (**Figure 3 Supplement 1**). While complex-specific heterochromatin

antagonizing activities have previously been ascribed to chromatin remodelers, such as Swr1C

(Meneghini et al. 2003) and Ino80 (Xue et al. 2015), we observe a broad involvement across remodelers.

The specific contributions of these remodeling complexes to destabilizing heterochromatin spreading are

interesting candidates for future studies.

Most of the GO complexes we found to be required for spreading have been implicated in

chromatin regulation or silencing pathways specifically (**Figure 3A**). One notable exception was two

predicted subunits of the AP-3 adaptor complex, Apm3 and Apl5, with roles either in all chromatin

contexts (Apm3) or only in *MAT* contexts (Apl5). We were intrigued by the discovery of an AP-3

adaptor protein complex normally associated with vesicular traffic. While the spreading phenotype is

moderate, *apm3* is one of only three genes that display a loss of spreading phenotype across all contexts,

along with *csn1* and *rrp17* encoding a key member of the COP9 signalosome and a putative rRNA

exonuclease, respectively. Thus, we sought to examine whether this phenotype correlates with a

H3K9me2 spreading defect and assessed H3K9me2 levels at the constitutive and facultative

heterochromatin loci by chromatin immunoprecipitation followed by qPCR (ChIP-qPCR). Consistent

with its moderate phenotype, *Δapm3* had mildly reduced H3K9me2 within the MAT locus, but not at

pericentromeres and subtelomeres (**Figure 3, Supplement 2A-C**). Importantly we find a role for Apm3

in H3K9me2 accumulation at heterochromatin islands, including *mei4, ssn4* and *mcp7* (**Figure 3**

Supplement 2D), indicating a role in facultative heterochromatin. Since AP-3 adaptor proteins are

cytoplasmic, we tested whether a fraction of Apm3 and Apl5 is also targeted to the nucleus. We

expressed Apm3 and Apl5 as fusions with SF-GFP and visualized them together with Swi6:E2C, which

marks heterochromatin foci in the nucleus (**Figure 3 Supplement 2E**). Notably, we found that

Apm3:SF-GFP is distributed broadly in both the cytosol and nucleus, whereas the related Apl5 protein

appeared to be excluded from the nucleus (**Figure 3 Supplement 2F**). Together, these data indicate that

Apm3 may represent a novel regulator of heterochromatin spreading, which is also supported by the

finding that it physically interacts with the heterochromatin regulators Fft3 (Lee et al. 2017) and Epe1

(Wang et al. 2013).

The relationship between histone deacetylation and gene silencing is well described. Three classes of HDACs exist, which have partially redundant and non-overlapping functions in the formation of heterochromatin domains and gene silencing. Sir2 belongs to the class III HDAC of the sirtuin family (Shankaranarayana et al. 2003). Clr3 belongs to class II and is a member of the SHREC complex (Sugiyama et al. 2007). Clr6 belongs to class I and is part of several sub-complexes, contributing to both heterochromatic and euchromatic gene regulation (Grewal et al. 1998; Nicolas et al. 2007). We find here that unlike class III and II HDACs, sub-complexes of the Clr6 family including the Rpd3S, Rpd3L-Expanded, and Clr6 I″ contribute exclusively to spreading and not nucleation (**Figure 1, 2** and **Figure 3A**, top of heatmap). In particular, we noticed that the forkhead transcription factor Fkh2, was identified as a common spreading regulator in all *MAT* HSS strains. Despite not being formally annotated to the Clr6 I″ complex by GO terms, Fkh2 was previously described as a member of this sub-complex (Zilio et al. 2014). We included Fkh2 as a member of Clr6 I″ in further analysis for this reason. Analogous to the analysis for nucleosome remodeling complexes, we represented the HDAC components that were identified as hits (**Figure 3C**). The population distributions of HSS reporter fluorescence in these Clr6 complex mutants are shown in **Figure 3 Supplement 2**. We find an interesting bifurcation in the contribution of Clr6 sub-complexes towards "loss" or "gain" categories. Clr6 I″ and Clr6S (Complex II) positively contributed to spreading ("loss"), while several members of the Rpd3L-Expanded complex antagonized spreading and were found as hits in the "gain" category. This includes a subset belonging to the Set3 Complex (Set3, Hif2, Hos2, Snt1). Overall, this suggests that Clr6 I″ and Clr6S HDAC complexes specifically promote heterochromatin spreading in addition to their described roles in transcriptional gene silencing.

Given the strong initial phenotype of Crl6 I″ subunits in spreading (**Figure 2**), we further explored their role here. Amongst the known members of Clr6 sub-complexes (**Figure 4A**), we chose

three subunits to validate our results: two Clr6 I″-specific members, Fkh2 and Png3, and the shared Clr6

core subunit Prw1. We validated the phenotypes by *de novo* single and double deletions for these genes

in the three *MAT* HSS reporter backgrounds (**Figure 4B-C, Figure 4 Supplement 1 A,B**) and

performed three-color flow cytometry to record fluorescence.

We first sought to validate the phenotype of these deletions in the *WT MAT* and *MAT ΔREIII*

HSS, which were most prominent in the screen. We found a similar phenotype for the single *Δprw1* and

the *Δfkh2 Δprw1* double mutant at *MAT ΔREIII,* corroborating the notion that Fkh2 is a Clr6 component

and acts in the same pathway as Prw1 (**Figure 4C**, bottom panels). However, we noted that the

phenotype of *Δprw1* was weaker than *Δfkh2* in the *WT MAT* background; nonetheless, also here the

double mutant showed a non-additive phenotype (**Figure 4C**, top panels). We additionally aimed to

validate these three deletions in the *MAT ΔcenH* HSS reporter background (**Figure 4 Supplement 1C**).

While this strain remains strongly repressed in the majority of the population, we detected a noticeable

increase in reporter signal in *Δfkh2* and *Δprw1*.

The above results, evidencing an impact on silencing at the spreading ("orange"), but not

nucleation ("green") reporter, predicted that the chromatin state may also be affected primarily at

nucleation-distal sites in Clr6 I″ complex mutants. We therefore next examined the chromatin state by

ChIP-qPCR analysis of the H3K9me2 mark at *MATΔREIII* and other heterochromatin loci. First, we

examined the MAT locus. Here we observed strong reductions of H3K9me2 signal at the "orange"

spreading reporter and also more nucleation-distal targets, in *Δfkh2, Δprw1*, and the *Δfkh2Δprw1* double

mutant (**Figure 4D**). There is no increase in severity of the spreading defect in the *Δfkh2Δprw1* double

mutant relative to the two single mutations, although further decreases in H3K9me2 should have been

detectable (see euchromatic target *mtd1,* **Figure 4D**). This further supports that Fkh2 and Prw1 act

together to promote spreading. However, consistent with the flow data, we did not observe strong

H3K9me2 reductions at the nucleation "green" reporter (**Figure 4D**), which is embedded in the *cenH*

nucleation element, except for a mild reduction in *Δfkh2*. We obtained a similar result at the related

pericentromeric *dh* and *dg* nucleation elements (**Figure 4 Supplement 1D**). We also find defects in

H3K9me2 spreading in Clr6 1″ subunit mutants at loci in addition to *MAT*. In particular, we observed

reduced H3K9me2 accumulation at subtelomeric targets in chromosomes I and II in *Δprw1* and *fkh2*,

with a stronger effect in *Δprw1* (**Figure 1E**). At the facultative heterochromatin islands *mei4, ssm4,* and

mcp7, *Δfkh2, Δprw1*, and *Δfkh2Δprw1* have similar effects as at subtelomeres (**Figure 4F**).

The *Δpng3* has a moderate to mild phenotype in *WT-MAT* and *MAT ΔREIII*, respectively (**Figure**

4 Supplement 1A,B). Consistent with this phenotype, we can observe a small change in H3K9me2

accumulation only distal to the "orange" reporter (**Figure 4D**) and a moderate effect at all

heterochromatin islands tested (**Figure 4F**), suggesting that Png3 plays a less prominent role than Fkh2

within Clr6 I″. These results evidence a defect of heterochromatin assembly in Clr6 I″ subunit mutants

primarily at distal, but not nucleation sites, such as *cenH* or *REIII*, indicating a surprising

heterochromatin-spreading specific role of Clr6 I″.

Discussion

The formation of a heterochromatin domain requires three interconnected steps, nucleation,

assembly of silencing structures, and the lateral spreading from DNA-sequence driven nucleation sites.

While the nucleation reaction has been well described, it has not been fully resolved which parts of the

genetic circuitry discovered to date are required for spreading. Prior studies on spreading have focused

primarily on factors that restrain heterochromatin formation across boundaries, which include Leo1,

Mst2, Epe1, Bdf2*,* among others (Ayoub et al. 2003; Wang et al. 2013; Verrier et al. 2015; Wang et al.

2015). Here, our ability to separate requirements for nucleation and distal spreading within

heterochromatin domains allowed us to pinpoint which factors are necessary to enable the spreading

reaction and factors that constrain it genome-wide. A key finding from this work is the requirement of

variants of the Clr6 HDAC complex specifically in the spreading reaction, in addition to the antagonism

by a broad class of chromatin remodelers, including Ino80, Swr1C, SWI/SNF and RSC (**Figure 3**).

HDACs have long been implicated in heterochromatin function generally, and here we were able

to distinguish which major HDACs regulate heterochromatin broadly, versus spreading specifically. The

Clr3 HDAC and associated SHREC complex is required for silencing, likely via its ability to repress

nucleosome turnover (Aygun et al. 2013), maintain nucleosome occupancy (Sugiyama et al. 2007;

Garcia et al. 2010), and remove H3K14 acetylation known to antagonize heterochromatin assembly

(Wirén et al. 2005). We find that SHREC mutants completely lose heterochromatic silencing (**Figure

1C-F, K**). Similarly, the Sir2 HDAC is broadly implicated in heterochromatin nucleation and assembly

(Shankaranarayana et al. 2003; Alper et al. 2013), and we also observe near-complete silencing loss in

Δsir2 (**Figure 1 Supplement 2**). Clr6 complexes have been implicated in suppression of antisense

transcription globally, particularly complex II/Rpd3S (Nicolas et al. 2007; Yamane et al. 2011) and the

maintenance of heterochromatic gene silencing at major constitutive sites (Grewal et al. 1998), via the

recruitment by HP1/Swi6 (Fischer et al. 2009). Remarkably, we find here that particular Clr6

subcomplexes are specifically required for distal spreading within constitutive heterochromatin and

heterochromatin islands (**Figure 3,4**). Clr6 sub-complex activity is only marginally required for

maintaining repression at nucleation-proximal sites. This is consistent with the finding that the *clr6-1*

allele has only small impacts on transcription of the *cenH* nucleator-encoded ncRNAs (Yamane et al.

2011). We find the recently characterized Fkh2-associated complex I″ (Zilio et al. 2014) to be a central

player in promoting distal spread (**Figure 3,4**). This complex contains the core of complex I, Nts1,

Png3, Mug165 and associates with Fkh2. Not all these subunits contributed equally to regulating

spreading in all chromatin contexts, with Fkh2 having the strongest effect (**Figure 3, 4, Figure 3**

Supplement 3 and **Figure 4 Supplement 1**). The double mutant analysis indicates that Fkh2 and Prw1

act together in promoting spreading (**Figure 4C**), which can be inferred from non-additive phenotype in

MAT ΔREIII. The mutants are also non-additive with respect to silencing in *WT MAT*, however *Δprw1*

and *Δfkh2Δprw1* phenotypes are weaker than *Δfkh2.* This type of behavior, i.e. the partial suppression of

phenotype by additional deletion of complex components, has been observed for silencing defects of

protein complexes (Barrales et al. 2016). In addition to Clr6 I″, the Alp13 and Cph1 subunits, which are

assigned to the separate complex II/Rpd3S (Nicolas et al. 2007) associated with global deacetylation,

also had positive impacts on spreading. This may indicate that (1) the composition of different Clr6

subcomplexes *in vivo* is either dynamic, or (2) a version of Rpd3S, jointly with a complex typified by

Fkh2, cooperate in promoting distal spreading of heterochromatin. Interestingly, the Set3-submodule

that typifies the Rpd3L-Expanded complex (Shevchenko et al. 2008), has a distinct spreading-

antagonizing behavior (**Figure 3C**). This contrasts with a mild positive role of the Set3 complex at

pericentromeres, which was proposed to be mediated indirectly, via transcriptional regulation of ClrC

H3K9 methylase complex genes (Yu et al. 2016).

We do not believe that the spreading-specific role of Clr6 complexes at the MAT locus is

mediated primarily by Asf/HIRA (Yamane et al. 2011), since Asf/HIRA subunits Hip1, Hip3 and Slm9

have mild to no phenotype for spreading in *MAT* contexts. Asf/HIRA mutant phenotypes were more

pronounced in *ECT*, which implies less reliance on Clr6 for spreading in *ECT* (**Figure 2**)*. The

spreading-specific role of specific Clr6 complexes may be encoded by their recruitment mechanisms.

HP1/Swi6, which recruits Clr6 (Fischer et al. 2009), is a known spreading regulator (Hall et al. 2002;

Canzio et al. 2011). Further, the Fkh2 transcription factor, which plays roles in origin coordination and

clustering in budding yeast (Knott et al. 2012), may play a key role in either directing Clr6 to distal sites

or ensuring continued association through the spreading process, which would explain its dominant roles

versus Png3 and other Clr6 I″ subcomplex-specific subunits (**Figure 4**, **Figure 4 Supplement 1**). The

precise role of Fkh2 will be the subject of further study.

Chromatin remodelers across several classes antagonize spreading. While Ino80 and Swr1C

have been linked to heterochromatin containment by specific mechanisms in budding yeast either by

H2A.Z exchange or preventing the invasion of euchromatin (Meneghini et al. 2003; Xue et al. 2015),

our results appear to indicate a more widespread effect of remodeling activities on spreading. This is

because in addition to Ino80 and Swr1C, the major remodeling complexes RSC and SWI/SNF also

contributed to spreading antagonism. A more general way remodelers have been implicated in

heterochromatin function is creating nucleosome free regions (NFRs, (Lorch and Kornberg 2017)) that

antagonize heterochromatin. Since NFRs may be roadblocks to spreading (Garcia et al. 2010;

Lantermann et al. 2010), it is possible that remodelers employ this mechanism to restrain

heterochromatin spreading globally. In addition, remodelers such as SWI/SNF and RSC destabilize

nucleosomes generally (Narlikar et al. 2001; Rowe and Narlikar 2010), leading to increased turnover

(Rawal et al. 2018), which would antagonize heterochromatin formation. This increased turnover may

be tolerated at ncRNA nucleation sites, where it is at near euchromatic levels (Greenstein et al. 2018),

likely due to ncRNA transcription (Volpe et al. 2002; Noma et al. 2004). This would indicate that

regulation of nucleosome stability has a particular significance at distal, but not nucleation sites.

Further, we found that related ncRNA nucleators in different chromatin environments require

distinct factors for spreading. Similarly, spreading from qualitatively different nucleators within the

same environment, namely *REIII* and *cenH,* also differ in their sensitivity to different mutants. The

significant overlap in factors between *WT MAT* and *MAT ΔREIII* indicates that heterochromatin

formation at MAT is dominated by the ncRNA nucleator *cenH,* in agreement with our previous findings

(Greenstein et al. 2018). The *REIII* element, which nucleates heterochromatin independent of ncRNA

(Jia et al. 2004), had different requirements. ncRNA-independent spreading is strongly antagonized by

the Epe1 pathway, and uniquely promoted by the MTOR pathway Gad8 kinase, partially consistent with

a previous report implicating Gad8 for MAT silencing (Cohen et al. 2018). The biggest difference was

between the MAT contexts and *ECT*. Some of the factors unique to *ECT* have been implicated in

inhibiting spreading across boundaries, such as Leo1 (Verrier et al. 2015), or heterochromatin stability

and spreading generally, such as HIRA (Yamane et al. 2011). However, these factors do not

significantly contribute to spreading within constitutive heterochromatin. The significant vulnerability of

ECT, compared to the similarly behaving *MAT ΔREIII* context, could be accounted by the following

possibilities: (1) The loss of Epe1 may impair RNAi specifically at *ECT,* nucleated by a pericentromeric

dh nucleator*,* but not *MAT ΔREIII,* which relies on the *dh*-homologous *cenH* nucleator (Trewick et al.

2007; Braun et al. 2011). (2) Alternatively, this Epe1-dependence may indicate that a naïve euchromatic

context is less able to compete for heterochromatin factors. *Δepe1* is known to induce heterochromatin

domain expansion at constitutive heterochromatin sites (Ayoub et al. 2003; Trewick et al. 2007; Braun

et al. 2011; Wang et al. 2015) and also heterochromatin islands (Zofall et al. 2012; Wang et al. 2015;

Greenstein et al. 2019), likely depleting factors available for *ECT.* This alternative would hold important

implications for heterochromatin formation in euchromatic contexts, which occurs throughout

differentiation in animal and plant systems (Ringrose and Paro 2007; Schmitz and Amasino 2007; Zylicz

et al. 2018). Of note, only the *ECT* context appeared to strongly require Hip1, and moderately Slm9, for

efficient spreading, which code for a key subunits of the HIRA H3/H4 chaperone. HIRA has been

implicated in stabilizing heterochromatic nucleosomes (Yamane et al. 2011) and it is possible this

requirement additionally highlights the challenge faced by heterochromatic domains expanding within

gene-rich chromatin, known to destabilize nucleosomes via transcription-associated processes.

In this work, we defined how regulation of heterochromatin silencing and nucleation differ in fundamental ways from distal spreading. While similar nucleation elements likely rely on a common set of machinery, the success of heterochromatin spreading appears much more sensitive to the chromatin context, particularly if euchromatic regions are targeted for *de novo* silencing. This finding has important implications for directing gene silencing of new loci appropriately, as cells change states in differentiation. In these situations, regions that are previously in a transcriptionally active state are invaded by heterochromatin and will have to compete for factors in a dosage limited system (Eissenberg et al. 1992; Nakayama et al. 2000).

Acknowledgements

We thank Sy E Redding, Douglas Myers-Turnbull and Kamir Hiam for helpful discussions on data

acquisition, analysis, and interpretation, and Arthur Molines for help with microscopy experiments. This

work was supported by grants from the National Institutes of Health (DP2GM123484) and the UCSF

Program for Breakthrough Biomedical Research (partially funded by the Sandler Foundation) to BA-S,

and the ARCS Foundation Scholarship and Hooper Graduate Fellowship to RAG. HN and CT were

supported by National Science Foundation Graduate Research Fellowships (grant number 1650113).

This work was supported by grants awarded to SB from the European Union Network of Excellence

EpiGeneSys (HEALTH-2010-257082). SB is Member of the Collaborative Research Center 1064

(Project-ID 213249687) funded by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation) and acknowledges infrastructural support. Flow cytometry and FACS data were generated

in the UCSF Parnassus Flow Cytometry Core which is supported by the Diabetes Research Center

(DRC) grants NIH P30 DK063720.

Materials & Methods

Strains and strain construction

Mutant Generation for Genetic Screen

For the ectopic locus HSS reporter strain, the screen was performed essentially as described (Greenstein et al. 2019). Briefly, the parent HSS reporter strain was crossed to a ~400 gene deletion mini-library (see table 1) primarily consisting of subset of the Bioneer haploid deletion and several independently validated mutants. Crosses were performed as described (Verrier et al. 2015; Barrales et al. 2016; Greenstein et al. 2019) using a RoToR HDA colony pinning robot (Singer) for ECT while for the MAT HSS reporter strains, crosses were generated using a 96 well manual pinner. In addition, for the MAT HSS strains three Δclr4 mutant isolates and six individual parent isolates from each genomic context were included as controls. Crosses for the MAT HSS strains were performed on ME media for 3d at 27°C, while for the ectopic HSS strain crosses were performed using SPAS media for 4d at room temperature. For all strains, crosses were incubated for 4-5d at 42°C to retain spores, while removing unmated haploid and diploid cells. For MAT HSS strains, spores were germinated on YES medium supplemented with G418 and hygromycin B. The ectopic locus HSS spores were germinated on YES medium supplemented with G418, hygromycin B, and nourseothricin. The resulting colonies were pinned into YES liquid medium for overnight growth and then prepared for flow cytometry as described below.

Validation Strain and Plasmid Construction

Plasmid constructs for gene knockout validation were generated by *in vivo* recombination as described (Greenstein et al. 2018; Greenstein et al. 2019). *S. pombe* transformants were selected as described (Greenstein et al. 2018). For microscopy, *hygMX* super-folder GFP (SFGFP) constructs for C-terminal tagging that we described previously (Al-Sady 2016) were amplified with 175bp ultramer primers with homology to *apm3* or *apl5* and transformed into a Swi6:E2C *kanMX* strain. Apm3:SFGFP;Swi6:E2C and Apl5:SFGFP;Swi6:E2C strains were selected on hygromycin B and G418. Integrations and gene knockout were confirmed by PCR.

All strains used for this study beyond the individual deletion library mutants are listed in Table 2.

Flow Cytometry

Flow cytometry data collection and normalization for genetic screen

In preparation for flow cytometry, overnight cultures were diluted to OD = 0.1 (approximately a 1:40 dilution) in rich media (YES) and incubated at 32°C with shaking of rpm for 4–6 hours. For the ectopic locus HSS strains, flow cytometry was performed essentially as described (Greenstein et al. 2018; Greenstein et al. 2019). For the MAT locus HSS strains, flow cytometry was performed using a Fortessa X20 Dual instrument (Becton Dickinson) attached with high throughput sampler (HTS) module. With a threshold of 30,000 events, samples sizes ranged from ~1000 to 30,000 cells depending on strain growth. Fluorescence detection and compensation, and data analysis were as described (Al-Sady et al. 2016; Greenstein et al. 2018; Greenstein et al. 2019). The R scripts for the screen analysis is included as a text file.

Flow cytometry data collection and normalization for validation

For validation flow cytometry experiments, cells were grown as described (Greenstein et al. 2018;
Greenstein et al. 2019) with the exception that cells were diluted into YES medium and grown 5-8 hours
before measurement. Flow cytometry was performed as above. Depending on strain growth and the
volume collected per experiment, fluorescence values were measured for ~20,000-100,000 cells per
replicate. Fluorescence detection, compensation, and data analysis were as described (Al-Sady et al.
2016; Greenstein et al. 2018; Greenstein et al. 2019)

<u>2D-density histogram plots with ON and OFF boundary guidelines.</u>
2D-density histogram plots (Al-Sady et al. 2016; Greenstein et al. 2018; Greenstein et al. 2019) were
generated as described previously, (see (Greenstein et al. 2018) methods for Figure 4) with the following
exceptions: For MAT locus strains, the guide-lines for boundary values of "off" and "on" states were
determined using median of a Red-Only control plus 3 times the median absolute deviation (MAD) and
median of Δ*clr4* minus 2 times the MAD value respectively. For the ECT strain, the "on" boundary
guideline was calculated by median of Δ*clr4* minus one MAD value. The boundary guideline for the
ECT "off" state was determined by adjusting the raw red-channel values for cells from of a no-color
control strain analyzed on the same flow cytometry run to simulate a Red-Only control strain by adding
the median value of the respective Δ*clr4* strain to the red value of each cell. The resulting adjusted data
was used to calculate the "off" cutoff by median +3MAD as above. Validation flow cytometry plots
were generated using the ggplot2 R package (Wickham 2016).

t-distributed stochastic neighbor embedding (t-SNE) analysis

Since our data varied in sample size, we transformed normalized fluorescent data into 5x5 density grids.
The percentage of population residing in each of the 25 grids is represented by one variable. Increasing
the number of grids to 8x8 or 10x10 did not significantly alter results. The following input settings were
used before t-SNE reduction: concatenated distributions, 1000 iterations, 60 perplexity, 0 Theta. t-SNE
reduction were conducted using the Rtsne package (Krijthe 2015) and the generated t-SNE model was
plotted using the R packages ggplot2 (Wickham 2016).

Earth Mover's Distance (EMD) Analysis

EMD is a distance measure between two multi-dimensional distributions to evaluate the dissimilarity.
EMD calculates the minimal amount of work to match the two equisized distribution (Rubner et al.
1998) The normalized fluorescent data are transformed into a 15x15 density grid as above. To calculate
the fraction of change between mutant and parent population, we calculated a metric of $EMD^{mut/par}$ by
taking the log2 transformation of the quotient of the EMD between mutant to Δ*clr4* distributions
(EMD^{mut}) by the EMD between parent to Δ*clr4* distributions (EMD^{par}). For MAT HSS strains, the three
Δ*clr4* and six parent control isolates were respectively combined and transformed into one reference
population matrix. Two-dimensional EMD between any pair of distributions in Euclidean distance are
calculated using functions from the R package emdist (Urbanek and Rubner 2012).

We validated the 1000 cell cutoff for the EMD analysis by iteratively down-sampling parent strains,
calculating the EMD value to the Δ*clr4* distribution, and comparing the resulting values across
iterations. For each HSS parent context, all wells containing parent isolates (N=6 for MAT locus
backgrounds, N=1 for ECT) were combined into one pool from which a random sample was drawn 100
times for each N = 1000, and 5000-30,000 in 5000 cell increments. For each iteration of each N number
of cells, the EMD value of the down-sampled population was calculated relative to the respective Δ*clr4*

population. We calculated the mean and SD for the 100 iterations of for each N and determined that even at an N of 1000 cells in the most broadly distributed background (ECT) the SD of EMD values was <2% of the EMD value of the parent calculated from its entire distribution. Given this analysis, we conclude that while the EMD values for wells with lower cell counts will likely have higher error than those with more cells, this error is not likely to have major effects on the calculated metric.

$EMD^{mut/par}$ data were plotted in a heatmap using functions from the R package superheat (Barter and Yu 2018). Any values < -2 were converted to -2 and interpreted as such for the heatmap.

Correlation Analysis

The correlation between the $EMD^{mut/par}$ data of different pairs of genomic contexts were calculated using a linear regression model. Sir2, ClrC and SHREC are essential heterochromatin assembly factor, and their mutants have very low $EMD^{mut/par}$. These mutants were disregarded from the model fitting because they would drive the correlation. Pearson correlation coefficients and 95% confidence interval were calculated using ggscatter functions from R package ggplot2 (Wickham 2016) and ggpubr (Kassambara 2020).

Spreading Analysis

Nucleated cells were extracted using a "green"-off gate, using median of a "red"-Only control plus 2 times the SD. Enrichment of cell populations in particular "orange" fluorescence ranges ($Grid_n$) are calculated as $Grid_n^{mut/par}$: fraction of mutant population is divided by the fraction of parent population in one grid. The intervals of "orange" fluorescence used in grids are determined by: median+2SD of "orange"-OFF cells, median-1SD of "orange"-ON cells and the median of the two. To evaluate gain of spreading phenotype, enrichment in Grid 1 in *WT MAT, MAT ΔREIII and ECT* were calculated. To evaluate loss of spreading phenotype, enrichment in Grid 3 and 4 in *WT MAT* and *MAT ΔcenH,* as well as Grid 4 in *MAT ΔREIII and ECT* were calculated. The distribution of the $Grid_n^{mut/par}$ were plotted as histogram with annotation of 85th percentile and median+2SD from parent isolates. Gene hit lists comprised mutations above median and 2SD within the 85th percentile. Upset plots were generated using the R package UpSetR (Conway et al. 2017). Barplots were plotted using the R packages ggplot2 (Wickham 2016).

GO Complex and Sub-Complex analysis

generating the heatmap count data
GO Complexes – Based on the GO Complex annotations [link] (2019) retrieved from pombase (Lock et al. 2018), GO complex membership was determined for genes identified as hits for each strain background and hit category (gain/loss). Using functions from the R package dplyr (Wickham et al. 2020), gene names were converted to systematic ID numbers and these systematic IDs were queried against the GO complex annotation table. The number of times a GO complex appeared per background and hit category was tabulated. Genes can be associated with any number of GO complexes depending on their annotations. However, any particular gene was only counted once per GO complex despite potentially being annotated to that GO complex by more than one evidence code. The unique list of GO complexes for all hits was determined and a matrix was computed representing the number of times each

GO complex (row) was identified per strain/hit category (column). This counts matrix was used to

generate the GO complex heatmap in Figure 3A, described below.

Sub-complex analysis – Genes annotated to the seven complexes in Figure 3B,C were obtained from

pombase (Lock et al. 2018). *fkh2* was added to the Clr6 I″ complex given the protein contacts described

previously (Zilio et al. 2014). For the unique set of genes per panel it was determined if each gene was

identified as a hit in each strain background/hit category combination. The data was summarized in a

counts matrix where rows represent the unique list of genes per panel and columns represent the strain

background / hit category. The counts matrix for each set of genes was used to generate the heatmaps in

Figure 3B, C as described below.

generating the heatmap clustering

Using the R package ComplexHeatmap (Gu et al. 2016), both row and column dendrogram and

clustering were generated using hierarchical clustering. Based on an optimal Silhouette score, the strain

background / hit category (columns) were clustered into 3 (Figure 3A). The dendrogram representing

complexes (Figure 3A) in rows were not separated because validations of the clustering by connectivity,

Dunn index or Silhouette score were inconclusive. Clustering validations were conducted using the R

package clValid (Brock et al. 2008).

Chromatin Immunoprecipitation and Quantification

Chromatin Immunoprecipitation (ChIP) was performed essentially as described (Greenstein et al. 2018;

Greenstein et al. 2019) Bulk populations of cells for were grown overnight to saturation in YES

medium. The following morning, cultures were diluted to OD 0.1 in 25mL YES and grown for 8h at

32°C and 225rpm. Based on OD measurements, 60×10^6 cells were fixed and processed for ChIP as

previously described (Greenstein et al. 2018) without the addition of W303 carrier. Cleared chromatin

for each ChIP sample was incubated with 1μL of anti-H3K9me2 antibody (Abcam, ab1220) overnight

after a small fraction was retained as Input/WCE. DNAs were quantified by RT-qPCR and percent

immunoprecipitation (%IP, ChIP DNA/Input DNA*100) was calculated as described (Greenstein et al.

2018). Data for %IP was plotted using the R packages ggplot2 (Wickham 2016) and dplyr (Wickham et

al. 2020).

Microscopy

Swi6:E2C; Apl5:SFGFP and Swi6:E2C; Apm3:SFGFP cells were grown is YS media as described.

Slides (ibidi, Cat. No. 80606) were pre-coated with 100 mg/mL lectin (Sigma-Aldrich, Cat. No. L1395)

diluted in water by adding lectin solution to slide for 1 min. and removing supernatant. Log-phase

growing cells were applied to the slide and excess cells were rinsed off with YS. Cells were immediately

imaged with a 60x objective (CFI Plan Apochromat VC 60XC WI) on a Nikon TI-E equipped with a

spinning-disk confocal head (CSU10, Yokogawa) and an EM-CCD camera (Hammamatsu). Cells were

imaged in brightfield and additionally excited with 488nm (SFGFP) and 561nm (E2C) lasers. Emission

was collected using a 510/50 band-pass filter for GFP emission and a 600/50 band-pass filter for E2C

emission. For the SFGFP and E2C channels, z-stacks were obtained at 0.3μm/slice for 11 slices total.

An overlay of the maximum z-projections for SFGFP and E2C channels are shown separately from the

brightfield images. Brightness and contrast were adjusted in ImageJ to clearly show both Swi6 and

Apl5/Apm5 signals in the overlay. At least 2 isolates were imaged to confirm localization patterns.

Data Availability
642
643 Screen flow cytometry data and analysis scripts will be made available upon publication of the
644 manuscript.
645

References

646 **References**

647 Al-Sady B, Greenstein RA, El-Samad HJ, Braun S, Madhani HD. 2016. Sensitive and Quantitative Three-Color
648  Protein Imaging in Fission Yeast Using Spectrally Diverse, Recoded Fluorescent Proteins with
649  Experimentally-Characterized In Vivo Maturation Kinetics. *PloS one* **11**: e0159292.
650 Al-Sady B, Madhani HD, Narlikar GJ. 2013. Division of labor between the chromodomains of HP1 and Suv39
651  methylase enables coordination of heterochromatin spread. *Mol Cell* **51**: 80-91.
652 Alabert C, Barth TK, Reveron-Gomez N, Sidoli S, Schmidt A, Jensen ON, Imhof A, Groth A. 2015. Two distinct
653  modes for propagation of histone PTMs across the cell cycle. *Genes & development* **29**: 585-590.
654 Alper BJ, Job G, Yadav RK, Shanker S, Lowe BR, Partridge JF. 2013. Sir2 is required for Clr4 to initiate
655  centromeric heterochromatin assembly in fission yeast. *The EMBO journal* **32**: 2321-2335.
656 Aygun O, Mehta S, Grewal SI. 2013. HDAC-mediated suppression of histone turnover promotes epigenetic
657  stability of heterochromatin. *Nature structural & molecular biology* **20**: 547-554.
658 Ayoub N, Goldshmidt I, Lyakhovetsky R, Cohen A. 2000. A fission yeast repression element cooperates with
659  centromere-like sequences and defines a mat silent domain boundary. *Genetics* **156**: 983-994.
660 Ayoub N, Noma K-i, Isaac S, Kahan T, Grewal SIS, Cohen A. 2003. A novel jmjC domain protein modulates
661  heterochromatization in fission yeast. *Molecular and cellular biology* **23**: 4356-4370.
662 Barrales RR, Forn M, Georgescu PR, Sarkadi Z, Braun S. 2016. Control of heterochromatin localization and
663  silencing by the nuclear membrane protein Lem2. *Genes & development* **30**: 133-148.
664 Barter RL, Yu B. 2018. Superheat: An R package for creating beautiful and extendable heatmaps for visualizing
665  complex data. *Journal of computational and graphical statistics : a joint publication of American
666  Statistical Association, Institute of Mathematical Statistics, Interface Foundation of North America* **27**:
667  910-922.
668 Bayne EH, Bijos DA, White SA, de Lima Alves F, Rappsilber J, Allshire RC. 2014. A systematic genetic screen
669  identifies new factors influencing centromeric heterochromatin integrity in fission yeast. *Genome biology*
670  **15**: 481.
671 Braun S, Garcia JF, Rowley M, Rougemaille M, Shankar S, Madhani HD. 2011. The Cul4-Ddb1(Cdt)(2)
672  ubiquitin ligase inhibits invasion of a boundary-associated antisilencing factor into heterochromatin. *Cell*
673  **144**: 41-54.
674 Brock G, Pihur V, Datta S, Datta S. 2008. clValid: An R Package for Cluster Validation *Journal of Statiscal
675  Software* **25**: http://www.jstatsoft.org/v25/i04.
676 Buhler M, Haas W, Gygi SP, Moazed D. 2007. RNAi-dependent and -independent RNA turnover mechanisms
677  contribute to heterochromatic gene silencing. *Cell* **129**: 707-721.
678 Canzio D, Chang EY, Shankar S, Kuchenbecker KM, Simon MD, Madhani HD, Narlikar GJ, Al-Sady B. 2011.
679  Chromodomain-mediated oligomerization of HP1 suggests a nucleosome-bridging mechanism for
680  heterochromatin assembly. *Mol Cell* **41**: 67-81.
681 Canzio D, Liao M, Naber N, Pate E, Larson A, Wu S, Marina DB, Garcia JF, Madhani HD, Cooke R et al. 2013.
682  A conformational switch in HP1 releases auto-inhibition to drive heterochromatin assembly. *Nature* **496**:
683  377-381.
684 Cohen A, Habib A, Laor D, Yadav S, Kupiec M, Weisman R. 2018. TOR complex 2 in fission yeast is required
685  for chromatin-mediated gene silencing and assembly of heterochromatic domains at subtelomeres. *The
686  Journal of biological chemistry* **293**: 8138-8150.
687 Consortium TGO. 2019. The Gene Ontology Resource: 20 years and still GOing strong. *Nucleic acids research*
688  **47**: D330-d338.
689 Conway JR, Lex A, Gehlenborg N. 2017. UpSetR: an R package for the visualization of intersecting sets and their
690  properties. *Bioinformatics (Oxford, England)* **33**: 2938-2940.
691 Eissenberg JC, Morris GD, Reuter G, Hartnett T. 1992. The heterochromatin-associated protein HP-1 is an
692  essential protein in Drosophila with dosage-dependent effects on position-effect variegation. *Genetics*
693  **131**: 345-352.

Fischer T, Cui B, Dhakshnamoorthy J, Zhou M, Rubin C, Zofall M, Veenstra TD, Grewal SIS. 2009. Diverse
roles of HP1 proteins in heterochromatin assembly and functions in fission yeast. *Proc Natl Acad Sci USA*
106: 8998-9003.
Garcia JF, Dumesic PA, Hartley PD, El-Samad H, Madhani HD. 2010. Combinatorial, site-specific requirement
for heterochromatic silencing factors in the elimination of nucleosome-free regions. *Genes &*
development **24**: 1758-1771.
Greenstein RA, Barrales RR, Sanchez NA, Bisanz JE, Braun S, Al-Sady B. 2019. Set1/COMPASS repels
heterochromatin invasion at euchromatic sites by disrupting Suv39/Clr4 activity and nucleosome stability.
Genes & development **34**: 99-117.
Greenstein RA, Jones SK, Spivey EC, Rybarski JR, Finkelstein IJ, Al-Sady B. 2018. Noncoding RNA-nucleated
heterochromatin spreading is intrinsically labile and requires accessory elements for epigenetic stability.
eLife **7**.
Grewal SIS, Bonaduce MJ, Klar AJS. 1998. Histone deacetylase homologs regulate epigenetic inheritance of
transcriptional silencing and chromosome segregation in fission yeast. *Genetics* **150**: 563-576.
Grewal SIS, Klar AJS. 1996. Chromosomal inheritance of epigenetic states in fission yeast during mitosis and
meiosis. *Cell* **86**: 95-101.
Gu Z, Eils R, Schlesner M. 2016. HilbertCurve: an R/Bioconductor package for high-resolution visualization of
genomic data. *Bioinformatics (Oxford, England)* **32**: 2372-2374.
Hall IM, Shankaranarayana GD, Noma KI, Ayoub N, Cohen A, Grewal SIS. 2002. Establishment and
maintenance of a heterochromatin domain. *Science* **297**: 2232-2237.
Hansen KR, Hazan I, Shanker S, Watt S, Verhein-Hansen J, Bahler J, Martienssen RA, Partridge JF, Cohen A,
Thon G. 2011. H3K9me-independent gene silencing in fission yeast heterochromatin by Clr5 and histone
deacetylases. *PLoS genetics* **7**: e1001268.
Hansen KR, Ibarra PT, Thon G. 2006. Evolutionary-conserved telomere-linked helicase genes of fission yeast are
repressed by silencing factors, RNAi components and the telomere-binding protein Taz1. *Nucleic acids*
research **34**: 78-88.
Jacobs SA, Taverna SD, Zhang YN, Briggs SD, Li JM, Eissenberg JC, Allis CD, Khorasanizadeh S. 2001.
Specificity of the HP1 chromo domain for the methylated N-terminus of histone H3. *Embo Journal* **20**:
5232-5241.
Jia S, Noma K, Grewal SI. 2004. RNAi-independent heterochromatin nucleation by the stress-activated
ATF/CREB family proteins. *Science* **304**: 1971-1976.
Kassambara A. 2020. ggpubr: 'ggplot2' Based Publication Ready Plots. https://CRAN.R-
project.org/package=ggpubr.
Keller C, Woolcock K, Hess D, Bühler M. 2010. Proteomic and functional analysis of the noncanonical poly(A)
polymerase Cid14. *RNA (New York, NY)* **16**: 1124-1129.
Kim HS, Choi ES, Shin JA, Jang YK, Park SD. 2004. Regulation of Swi6/HP1-dependent heterochromatin
assembly by cooperation of components of the mitogen-activated protein kinase pathway and a histone
deacetylase Clr6. *The Journal of biological chemistry* **279**: 42850-42859.
Knott SR, Peace JM, Ostrow AZ, Gan Y, Rex AE, Viggiani CJ, Tavaré S, Aparicio OM. 2012. Forkhead
transcription factors establish origin timing and long-range clustering in S. cerevisiae. *Cell* **148**: 99-111.
Krijthe JH. 2015. Rtsne: T-Distributed Stochastic Neighbor Embedding using a Barnes-Hut Implementation.
https://github.com/jkrijthe/Rtsne.
Lachner M, O'Carroll N, Rea S, Mechtler K, Jenuwein T. 2001. Methylation of histone H3 lysine 9 creates a
binding site for HP1 proteins. *Nature* **410**: 116-120.
Lantermann AB, Straub T, Stralfors A, Yuan GC, Ekwall K, Korber P. 2010. Schizosaccharomyces pombe
genome-wide nucleosome mapping reveals positioning mechanisms distinct from those of Saccharomyces
cerevisiae. *Nature structural & molecular biology* **17**: 251-U215.
Lee J, Choi ES, Seo HD, Kang K, Gilmore JM, Florens L, Washburn MP, Choe J, Workman JL, Lee D. 2017.
Chromatin remodeller Fun30(Fft3) induces nucleosome disassembly to facilitate RNA polymerase II
elongation. *Nature communications* **8**: 14527.

Lejeune E, Bortfeld M, White SA, Pidoux AL, Ekwall K, Allshire RC, Ladurner AG. 2007. The chromatin-
remodeling factor FACT contributes to centromeric heterochromatin independently of RNAi. *Current*
biology : CB **17**: 1219-1224.
Lock A, Rutherford K, Harris MA, Hayles J, Oliver SG, Bähler J, Wood V. 2018. PomBase 2018: user-driven
reimplementation of the fission yeast database provides rapid and intuitive access to diverse,
interconnected information. *Nucleic acids research* **47**: D821-D827.
Lorch Y, Kornberg RD. 2017. Chromatin-remodeling for transcription. *Quarterly reviews of biophysics* **50**: e5.
Marina DB, Shankar S, Natarajan P, Finn KJ, Madhani HD. 2013. A conserved ncRNA-binding protein recruits
silencing factors to heterochromatin through an RNAi-independent mechanism. *Genes & development* **27**:
1851-1856.
Meneghini MD, Wu M, Madhani HD. 2003. Conserved histone variant H2A.Z protects euchromatin from the
ectopic spread of silent heterochromatin. *Cell* **112**: 725-736.
Moazed D. 2009. Small RNAs in transcriptional gene silencing and genome defence. *Nature* **457**: 413-420.
Nakayama J, Klar AJ, Grewal SI. 2000. A chromodomain protein, Swi6, performs imprinting functions in fission
yeast during mitosis and meiosis. *Cell* **101**: 307-317.
Narlikar GJ, Phelan ML, Kingston RE. 2001. Generation and Interconversion of Multiple Distinct Nucleosomal
States as a Mechanism for Catalyzing Chromatin Fluidity. *Mol Cell* **8**: 1219-1230.
Nicetto D, Zaret KS. 2019. Role of H3K9me3 heterochromatin in cell identity establishment and maintenance.
Current opinion in genetics & development **55**: 1-10.
Nicolas E, Yamada T, Cam HP, FitzGerald PC, Kobayashi R, Grewal SIS. 2007. Distinct roles of HDAC
complexes in promoter silencing, antisense suppression and DNA damage protection. *Nature structural &*
molecular biology **14**: 372-380.
Noma K, Sugiyama T, Cam H, Verdel A, Zofall M, Jia S, Moazed D, Grewal SI. 2004. RITS acts in cis to
promote RNA interference-mediated transcriptional and post-transcriptional silencing. *Nature genetics*
36: 1174-1180.
Orlova DY, Zimmerman N, Meehan S, Meehan C, Waters J, Ghosn EE, Filatenkov A, Kolyagin GA, Gernez Y,
Tsuda S et al. 2016. Earth Mover's Distance (EMD): A True Metric for Comparing Biomarker Expression
Levels in Cell Populations. *PloS one* **11**: e0151859.
Ragunathan K, Jih G, Moazed D. 2015. Epigenetics. Epigenetic inheritance uncoupled from sequence-specific
recruitment. *Science* **348**: 1258699.
Rawal Y, Chereji RV, Qiu H, Ananthakrishnan S, Govind CK, Clark DJ, Hinnebusch AG. 2018. SWI/SNF and
RSC cooperate to reposition and evict promoter nucleosomes at highly expressed genes in yeast. *Genes &*
development **32**: 695-710.
Reyes-Turcu FE, Zhang K, Zofall M, Chen E, Grewal SI. 2011. Defects in RNA quality control factors reveal
RNAi-independent nucleation of heterochromatin. *Nat Struct Mol Biol* **18**: 1132-1138.
Ringrose L, Paro R. 2007. Polycomb/Trithorax response elements and epigenetic memory of cell identity.
Development **134**: 223-232.
Rowe CE, Narlikar GJ. 2010. The ATP-dependent remodeler RSC transfers histone dimers and octamers through
the rapid formation of an unstable encounter intermediate. *Biochemistry* **49**: 9882-9890.
Rubner Y, Tomasi C, Guibas LJ. 1998. A metric for distributions with applications to image databases. . in *IEEE*
International Conference on Computer Vision, pp. 59-66.
Schmitz RJ, Amasino RM. 2007. Vernalization: a model for investigating epigenetics and eukaryotic gene
regulation in plants. *Biochimica et biophysica acta* **1769**: 269-275.
Scott KC, Merrett SL, Willard HF. 2006. A heterochromatin barrier partitions the fission yeast centromere into
discrete chromatin domains. *Current biology : CB* **16**: 119-129.
Shankaranarayana GD, Motamedi MR, Moazed D, Grewal SIS. 2003. Sir2 regulates histone H3 lysine 9
methylation and heterochromatin assembly in fission yeast. *Current Biology* **13**: 1240-1246.
Shevchenko A, Roguev A, Schaft D, Buchanan L, Habermann B, Sakalar C, Thomas H, Krogan NJ, Shevchenko
A, Stewart AF. 2008. Chromatin Central: towards the comparative proteome by accurate mapping of the
yeast proteomic environment. *Genome biology* **9**: R167.

Sugiyama T, Cam HP, Sugiyama R, Noma K-i, Zofall M, Kobayashi R, Grewal SIS. 2007. SHREC, an effector
complex for heterochromatic transcriptional silencing. *Cell* **128**: 491-504.
Thon G, Bjerling KP, Nielsen IS. 1999. Localization and properties of a silencing element near the mat3-M
mating-type cassette of Schizosaccharomyces pombe. *Genetics* **151**: 945-963.
Trewick SC, Minc E, Antonelli R, Urano T, Allshire RC. 2007. The JmjC domain protein Epe1 prevents
unregulated assembly and disassembly of heterochromatin. *The EMBO journal* **26**: 4670-4682.
Urbanek S, Rubner Y. 2012. emdist: earth mover's distance. https://CRAN.R-project.org/package=emdist.
Verrier L, Taglini F, Barrales RR, Webb S, Urano T, Braun S, Bayne EH. 2015. Global regulation of
heterochromatin spreading by Leo1. *Open biology* **5**.
Volpe TA, Kidner C, Hall IM, Teng G, Grewal SIS, Martienssen RA. 2002. Regulation of heterochromatic
silencing and histone H3 lysine-9 methylation by RNAi. *Science* **297**: 1833-1837.
Wang J, Cohen AL, Letian A, Tadeo X, Moresco JJ, Liu J, Yates JR, 3rd, Qiao F, Jia S. 2016. The proper
connection between shelterin components is required for telomeric heterochromatin assembly. *Genes &*
development **30**: 827-839.
Wang J, Reddy BD, Jia S. 2015. Rapid epigenetic adaptation to uncontrolled heterochromatin spreading. *eLife* **4**.
Wang J, Tadeo X, Hou H, Tu PG, Thompson J, Yates JR, 3rd, Jia S. 2013. Epe1 recruits BET family
bromodomain protein Bdf2 to establish heterochromatin boundaries. *Genes & development* **27**: 1886-
1902.
Wen B, Wu H, Shinkai Y, Irizarry RA, Feinberg AP. 2009. Large histone H3 lysine 9 dimethylated chromatin
blocks distinguish differentiated from embryonic stem cells. *Nature genetics* **41**: 246-250.
Wickham H. 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer
Wickham H, François R, Henry L, Müller K. 2020. dplyr: A Grammar of Data Manipulation. in *R package*
version 0 8 4 https://CRAN.R-project.org/package=dplyr.
Wirén M, Silverstein RA, Sinha I, Walfridsson J, Lee H-M, Laurenson P, Pillus L, Robyr D, Grunstein M, Ekwall
K. 2005. Genomewide analysis of nucleosome density histone acetylation and HDAC function in fission
yeast. *The EMBO journal* **24**: 2906-2918.
Xue Y, Van C, Pradhan SK, Su T, Gehrke J, Kuryan BG, Kitada T, Vashisht A, Tran N, Wohlschlegel J et al.
2015. The Ino80 complex prevents invasion of euchromatin into silent chromatin. *Genes & development*
29: 350-355.
Yamane K, Mizuguchi T, Cui B, Zofall M, Noma K, Grewal SI. 2011. Asf1/HIRA facilitate global histone
deacetylation and associate with HP1 to promote nucleosome occupancy at heterochromatic loci. *Mol*
Cell **41**: 56-66.
Yu Y, Zhou H, Deng X, Wang W, Lu H. 2016. Set3 contributes to heterochromatin integrity by promoting
transcription of subunits of Clr4-Rik1-Cul4 histone methyltransferase complex in fission yeast. *Sci Rep* **6**:
31752.
Zhang K, Mosch K, Fischle W, Grewal SIS. 2008. Roles of the Clr4 methyltransferase complex in nucleation,
spreading and maintenance of heterochromatin. *Nature structural & molecular biology* **15**: 381-388.
Zhu J, Adli M, Zou JY, Verstappen G, Coyne M, Zhang X, Durham T, Miri M, Deshpande V, De Jager PL et al.
2013. Genome-wide chromatin state transitions associated with developmental and environmental cues.
Cell **152**: 642-654.
Zilio N, Codlin S, Vashisht AA, Bitton DA, Head SR, Wohlschlegel JA, Bähler J, Boddy MN. 2014. A novel
histone deacetylase complex in the control of transcription and genome stability. *Molecular and cellular*
biology **34**: 3500-3514.
Zofall M, Smith DR, Mizuguchi T, Dhakshnamoorthy J, Grewal SIS. 2016. Taz1-Shelterin Promotes Facultative
Heterochromatin Assembly at Chromosome-Internal Sites Containing Late Replication Origins. *Mol Cell*
62: 862-874.
Zofall M, Yamanaka S, Reyes-Turcu FE, Zhang K, Rubin C, Grewal SI. 2012. RNA elimination machinery
targeting meiotic mRNAs promotes facultative heterochromatin formation. *Science* **335**: 96-100.
Zylicz JJ, Borensztein M, Wong FC, Huang Y, Lee C, Dietmann S, Surani MA. 2018. G9a regulates temporal
preimplantation developmental program and lineage segregation in blastocyst. *eLife* **7**.

844 Zylicz JJ, Dietmann S, Gunesdogan U, Hackett JA, Cougot D, Lee C, Surani MA. 2015. Chromatin dynamics and
845   the role of G9a in gene regulation and enhancer silencing during early mouse development. *eLife* **4**.

      FIGURE LEGENDS

Figure 1: A genetic screen based on a suite of fluorescent reporters identifies context-
dependent positive and negative regulators of heterochromatin function.

A. TOP: Overview of heterochromatin spreading sensor (HSS, Greenstein 2018). Three
transcriptionally encoded fluorescent protein genes are integrated in the genome. *SFGFP* ("green")
proximal to the nucleation site allows identification of heterochromatin nucleation; *mKO2* ("orange")
distal to the nucleation site allows identification of heterochromatin spreading. *3xE2C* ("red") in a
euchromatin region normalizes cell-to-cell noise. BOTTOM: The endogenous mating type locus (*MAT*)
and ectopically heterochromatic *ura4* locus (Greenstein 2018) were examined with the HSS in the
screen. *Bona fide* mutations of the nucleators, *cenH* and *REIII*, in *MAT* were made to limit nucleation to
occur from one site.

B. Overview of data processing for t-SNE. The multidimensional fluorescence data is linearized before
subjection to t-SNE. Two-dimensional "orange"/"red" *v.* "green"/"red" density plots are broken down into
25 grids in an unbiased manner, and normalized cell counts of each grid are used as parameters for
each mutant in the t-SNE processing. The 25 parameters of all mutants from all chromatin contexts
were embedded into a model two-dimensional t-SNE space.

C.-F. 2D-density hexbin plots of the wild-type parent or *Δclr3* mutant in the (**C**) *WT MAT,* (**D**) *MAT*
ΔcenH, (**E**)*MAT ΔREIII,* and (**F**) *ECT* background. Hexagonal bins are colored from light grey to black
indicating low to high density of cells per bin. Blue lines indicate boundary guidelines for the fully
repressed state and red lines indicate boundary guidelines values for fully expressed state (see
methods for treatment of MAT and ECT strains)

G-I. t-SNE visualization of all mutants across all four chromatin contexts. Each data point represents a
mutant, the fill color represents the (**G**) chromatin context of the mutant, or median (**H**) "green" or (**I**)
"orange" fluorescence of the entire mutant population. In (**H**) the parent isolates of each background are
depicted in individual colors indicated in the key. In (**I**), selected mutants are shown, with the chromatin
contexts highlighted with the same colors as (C-F). An enlarged region to highlight the *Δclr3* mutants is
shown to the right.

J. To linearize the multidimensional fluorescence data, the earth mover's distance (EMD) between each
mutant and a *Δclr4* mutant is calculated. EMD$^{mut/par}$ is computed by dividing each mutant EMD (EMD^{mut})
by the respective parent EMD (EMD^{par}) and transforming the quotient by log2.

K. Heatmaps depicting EMD$^{mut/par}$ of indicated mutants in each chromatin context. Any values <-2 were
converted to -2 and interpreted as such. Crossed-out boxes indicate mutants excluded from the
analysis due to growth defect or low sample size.

Figure 2: Identification of heterochromatin spreading regulators in different chromatin contexts.

A. Overview of the spreading-specific analysis with mock distributions of cells and grids indicated. To
segregate spreading from nucleation or silencing phenotypes, "green"-off populations (successful
nucleation events) are isolated. Within these populations, enrichment of cell populations in particular
"orange" fluorescence ranges (Grid$_n$) are calculated as Grid$_n^{mut/par}$. *E.g.* to identify loss of spreading
mutants in *WT MAT*, Grid$_{3+4}^{mut/par}$ is calculated as percentage of mutant population divided by
percentage of parent population in Grid$_{3+4}$ (green, blue). The Grids used for analysis of gain and loss of

spreading in the four chromatin contexts are indicated. Grid$_{3+4}$ was used for *WT MAT* and *MAT ΔcenH,*
given the highly repressed nature of this chromatin context.

B.-C. Upset plots indicating the frequency of (**B**) Loss of Spreading, or (**C**) Gain of Spreading gene hits
appearing in one or multiple chromatin contexts. For each bar, the chromatin context(s) with shared
phenotypes for the underlying gene hits is indicated below the plot. The inset indicates the total number
gene hits in each chromatin context of the same phenotype.

D.-G. Bar graphs representing the Grid$_n^{mut/par}$ and number of chromatin context(s) of gene hits with Loss
of Spreading phenotype from (**D**) *WT MAT*, (**E**) *MAT ΔcenH,* (**F**) *MAT ΔREIII* and (**G**) *ECT* respectively.
The genes are ranked in descending order of Grid$_n^{mut/par}$ and the color of each bar represents the
number of backgrounds the mutant show the same phenotype.

H.-I. 2D-density hexbin plots of the top two loss of spreading gene hits for all 4 chromatin contexts, (**H**)
WT MAT, (**I**) *MAT ΔcenH,* (**J**) *MAT ΔREIII,* and (**K**) *ECT*. Plotting as in Figure 1, C-F. For wild-type
comparison, see Figure 1 C-F.

Figure 3: Heterochromatin spreading is regulated by sets of unique and common protein
complexes across different chromatin contexts.

A. Heatmap of GO complex annotations for hits in each category and strain. Rows, representing GO
complexes annotated to genes within the screen that were identified as hits, are arranged via
hierarchical clustering. Columns are defined by the hit category (loss of spreading – white; gain of
spreading – black), and each screen chromatin context is indicated at the top. The columns were
clustered by hierarchical clustering and the tree was cut to define 3 clusters.

B. Hit table of complex members for Swr1C, Ino80, SWI/SNF, and RSC-type complexes. Components
identified as a hit in either the "gain" or "loss" category for each background are marked blue. Columns
are defined and ordered as in (**A**). The proteins present in each complex are annotated at the right with
the presence of color indicating membership of that protein in the complex.

C. Hit table of complex members for Rpd3L-Expanded, Rpd3S, and Clr6 complex I" as in (**B**). Columns
are defined and ordered as in (**A**) Proteins present in each complex are annotated as in (**B**).

Figure 4: Clr6 Complex I" regulates heterochromatin spreading at constitutive and facultative
heterochromatin loci.

A. Cartoon of Clr6 complexes (Clr6L, the Clr6LExpanded modules, Clr6 subcomplex 1", and Clr6S).
Subunits not in the screen are shown in grey.

B. Scheme for generation of deletion mutants. Double mutant of fkh2 and prw1 was generated by both
cross and *de novo* deletion of *prw1* in *Δfkh2*.

C. TOP PANELS: 2D-density hexbin plots for WT and Clr6 I" mutants in the *WT MAT* strain
background. Hexagonal bins are colored from light grey to black indicating low to high density of cells
per bin. A rug plot is included on the X and Y axes indicating the 1D density for each color. Rug lines
are colored with partial transparency to assist with visualization of density changes. Blue lines indicate
boundary guidelines for the fully repressed state and red lines indicate boundary guidelines values for
fully expressed state as in Figure 1.

BOTTOM PANELS: 2D-density hexbin plots with rug as above for WT and Clr6 I" mutants in the *MAT*
ΔREIII strain background.

D.-F. ChIP-qPCR quantification of H3K9me2 signal in the *MAT ΔREIII* strain at constitutive and
facultative heterochromatin regions. Error bars represent 1SD from three biological replicate isolates.
Individual values are plotted for each isolate. The WT data is additionally replicated in Figure 3
Supplement 2 as these experiments were performed together. SD; standard deviation. Telomere 1L
primer distances are from the end of the assembled genomic sequence.

Figure 1 Supplement 1: Screen overview.

A. Schematic of the screen to identify genes that contribute to heterochromatin nucleation and
spreading. A custom nuclear function deletion library (Table 1) was mated with four different reporter
strains (*WT MAT, MAT ΔcenH, MAT ΔREIII and ECT*). The fluorescences of "green", "orange" and
"red" for each mutant cell within each background are recorded by flow cytometry.

Figure 1 Supplement 2: Pairwise plots depicting the comparisons of mutant EMD$^{mut/par}$.

Pairwise plots depicting the EMD$^{mut/par}$ comparisons of (**A**) *MAT ΔcenH* with *WT MAT,* (**B**) *MAT ΔREIII*
with *WT MAT,* (**C**) *ECT* with *WT MAT,* (**D**) *MAT ΔREIII* with *MAT ΔcenH,* (**E**) *ECT* with *MAT ΔcenH* and
(**F**) *ECT* with *MAT ΔREIII*. Pearson correlation coefficients (ρ), determined without SHREC, ClrC
mutants and *Δsir2*, were calculated and correlations plotted for A, B and D.

Figure 2 Supplement 1: The distribution of Grid$_n$$^{mut/par}$ of all mutants in the loss or gain of
spreading hit categories.

A.-D. Histograms representing the distribution of Grid$_n$$^{mut/par}$ of (**A**) *WT MAT*, (**B**) *MAT ΔcenH,* (**C**) *MAT*
ΔREIII and (**D**) *ECT* in the loss of spreading hit category. The red line indicates two standard deviation
above median of parent isolates. The dashed blue line indicates the 85th percentile.

E.-G. Histograms representing the distribution of Grid$_n$$^{mut/par}$ of (**E**) *WT MAT,* (**F**) *MAT ΔREIII* and (**G**)
ECT in the gain of spreading hit category. The red line indicates two standard deviation above median
of parent isolates. The dashed blue line indicates the 85th percentile.

Figure 2 Supplement 2: Grid$_n$$^{mut/par}$ and overlapped chromatin contexts of identified gain of
spreading mutants

A.-C. Bar graphs representing the Grid$_n$$^{mut/par}$ and number of overlapped chromatin context(s) of gene
hits with Gain of Spreading phenotype in (**A**) *WT MAT*, (**B**) *MAT ΔREIII* and (**C**) *ECT* chromatin
contexts respectively. The genes are ranked in descending order of Grid$_n$$^{mut/par}$ and the color of each bar
represents the number of backgrounds the mutant show the same phenotype.

D.-F. 2D-density hexbin plots of the top two gain of spreading gene hits for all 3 chromatin contexts, (**D**)
WT MAT, (**E**) *MAT ΔREIII,* and (**F**) *ECT*. Plotting as in Figure 1, C-F. For wild-type comparison, see
Figure 1 C-F.

Figure 3 Supplement 1: Original 2D density histograms for chromatin remodeler gain of
spreading hits.

A. Chromatin remodeler gain of spreading gene hits for *MAT ΔREIII.* Original 2D density histograms for
all the gain of spreading gene hits from Figure 3B are shown. **B.** Chromatin remodeler gain of
spreading gene hits for *ECT.* Original 2D density histograms for all the gain of spreading gene hits from
Figure 3B are shown. GO complex annotations are indicated next to each mutant. 2D density
histograms as in Figure 1 C-F. Original *MAT ΔREIII* and *ECT* parents shown in Figure 1 are reproduced
here again (with transparency) for comparison. Mutants within each chromatin next are shown in
descending order of their $Grid_n^{mut/par}$ values.

1000
1001   **Figure 3 Supplement 2: *Δapm3* has a mild heterochromatin spreading phenotype and the**
1002   **protein is distributed in the cytosol and nucleus.**

1003

A.-D. ChIP-qPCR quantification of H3K9me2 signal in the *MAT ΔREIII* strain at constitutive and
facultative heterochromatin regions in wild-type and an independently generated *Δapm3* alleles. Error
bars represent 1SD from three biological replicate isolates. Individual values are plotted for each
isolate. The WT data is additionally replicated in Figure 4 as these experiments were performed
together.

E. Apm3:SFGFP is distributed in the cytosol and nucleus. Apm3:SFGFP was expressed from its native
locus and co-expressed with Swi6:E2C. Swi6:E2C labels nuclear heterochromatin. Z-projection
overlays of the Apm3:SFGFP and Swi6:E2C on top, and a brightfield image on the bottom.

F. Apl5:SFGFP is largely nuclear excluded. Apl5:SFGFP was expressed from its native locus and co-
expressed with Swi6:E2C. Swi6:E2C labels nuclear heterochromatin. Z-projection overlays of the
Apl5:SFGFP and Swi6:E2C on top, and a brightfield image on the bottom.

1015
1016
1017   **Figure 3 Supplement 3: Original 2D density histogram for all Clr6 complex subunit mutants in**
1018   ***MAT ΔREIII.***
1019
1020   Original 2D density histograms of all Clr6 complexes gene mutants corresponding to Figure 3C in *MAT*
1021   *ΔREIII* context. Original *MAT ΔREIII* wild type parent and *Δfkh2* and *Δprw1* mutants shown in Figures 1
1022   and 4 are reproduced here again (with transparency) for comparison.

1023

1024   **Figure 4 Supplement 1: The effect of Clr6 I" in the *MAT ΔcenH* background.**
1025
1026   **A.** 2D-density hexbin plots with rug as in Figure 4C for *Δpng3* and WT control run on the same day in
1027   the *WT MAT* strain background.
1028
1029   **B.** 2D-density hexbin plots with rug as in Figure 4C for *Δpng3* and WT control run on the same day in
1030   the *MATΔREIII* strain background.
1031
1032   **C.** Scatter plot with rug for WT and Clr6 I" mutants in the *MAT ΔcenH* strain background. Cell are
1033   plotted as individual points versus summarized in 2D density hexbin plots for increased resolution.
1034   Points and rug lines are colored with partial transparency to assist with visualization of density changes.
1035
1036   **D.** ChIP-qPCR quantification of H3K9me2 signal in the *MAT ΔREIII* strain at the pericentromeric
1037   heterochromatin region. Error bars represent 1SD from three biological replicate isolates. Individual
1038   values are plotted for each isolate. SD; standard deviation.
1039

Figure 1

Figure 2

Figure 3

Figure 4

Figure 1 - Supplement 1

A

Figure 1 - Supplement 2

Figure 2 - Supplement 1

Figure 2 - Supplement 2

A **Gain of Spreading**

Figure 3 - Supplement 1

A *MAT ΔREIII* - Chromatin Remodelers

B *ECT* - Chromatin Remodelers

Figure 3 - Supplement 2

A pericentromere

%IP H3K9me2

dh *dg*

B *MAT* locus

"green" "orange" *MAT distal* *mtd1p*

C subtelomere

tel1L 13kb tel1L 30kb

D heterochromatin islands

mei4 *ssm4* *mcp7*

genotype
WT
apm3

E Apm3:SFGFP Swi6:E2C

brightfield

8 μm

F Apl5:SFGFP Swi6:E2C

brightfield

8 μm

Figure 3 - Supplement 3

MAT △REIII - Clr6 Subcomplex Components

Figure 4 Supplement 1

A wild-type parent | Δpng3

B wild-type parent | Δpng3

C wild-type parent | Δfkh2 | Δprw1 | Δpng3

D pericentromere

Table 1: Nuclear function deletion library

Table 1: Nuclear function gene deletion library

	Systematic ID	Symbol	Description
1	SPAC1002.05c	jmj2	histone demethylase Jmj2
2	SPAC1006.03c	red1	RNA elimination defective protein Red1
3	SPAC1039.05c	klf1	transcription factor, zf-fungal binuclear cluster type Klf1
4	SPAC1071.02	mms19	CIA machinery protein Mms19
5	SPAC1071.06	arp9	SWI/SNF and RSC complex subunit Arp9
6	SPAC10F6.08c	nht1	Ino80 complex HMG box subunit Nht1
7	SPAC10F6.11c	atg17	autophagy associated protein kinase activator Atg17
8	SPAC1142.03c	swi2	Swi5 complex subunit Swi2
9	SPAC1142.08	fhl1	forkhead transcription factor Fhl1
10	SPAC11D3.07c	toe4	transcription factor, zf-fungal binuclear cluster type(predicted)
11	SPAC11D3.16c		Schizosaccharomyces specific protein
12	SPAC11E3.01c	swr1	SNF2 family ATP-dependent DNA helicase Swr1
13	SPAC11H11.01	sst6	ESCRT I complex subunit Vps23
14	SPAC11H11.05c	fta6	Mis6-Sim4 complex Fta6
15	SPAC12B10.10	nod1	medial cortical node Gef2-related protein protein Nod1
16	SPAC12G12.13c	cid14	TRAMP complex poly(A) polymerase subunit Cid14
17	SPAC139.03	toe2	transcription factor, zf-fungal binuclear cluster type (predicted)
18	SPAC139.06	hat1	histone acetyltransferase Hat1
19	SPAC1399.05c	toe1	transcription factor, zf-fungal binuclear cluster type
20	SPAC13A11.04c	ubp8	SAGA complex ubiquitin C-terminal hydrolase Ubp8
21	SPAC13D6.02c	byr3	translational activator, zf-CCHC type zinc finger protein (predicted)
22	SPAC13G6.01c	rad8	ubiquitin-protein ligase E3/ ATP-dependent DNA helicase Rad8
23	SPAC144.02	iec1	Ino80 complex subunit Iec1
24	SPAC144.05		DNA-dependent ATPase/ ubiquitin-protein ligase E3 (predicted)
25	SPAC144.06	apl5	AP-3 adaptor complex subunit Apl5 (predicted)
26	SPAC144.14	klp8	kinesin-like protein Klp8
27	SPAC14C4.06c	nab2	poly(A) binding protein Nab2 (predicted)
28	SPAC14C4.12c	laf1	Clr6 L associated factor 1 Laf1
29	SPAC14C4.13	rad17	RFC related checkpoint protein Rad17
30	SPAC1556.01c	rad50	DNA repair protein Rad50
31	SPAC15A10.11	ubr11	UBR ubiquitin-protein ligase E3 Ubr11
32	SPAC15A10.15	sgo2	inner centromere protein, shugoshin Sgo2
33	SPAC1610.01	saf5	splicing factor Saf5
34	SPAC1610.02c	mrpl1	mitochondrial ribosomal protein subunit L1 (predicted)
35	SPAC1687.05	pli1	SUMO E3 ligase Pli1
36	SPAC1687.09	irs4	autophagy/CVT pathway ENTH/VHS domain protein Irs4 (predicted)
37	SPAC16A10.03c		ubiquitin-protein ligase E3 involved in vesicle docking Pep5/Vps11-like (predicted)
38	SPAC16A10.07c	taz1	shelterin complex subunit Taz1
39	SPAC16C9.04c	mot2	CCR4-Not complex ubiquitin-protein ligase E3 subunit Mot2
40	SPAC16C9.05	cph1	Clr6 histone deacetylase associated PHD protein-1 Cph1
41	SPAC16E8.12c	png3	ING family homolog Png3 (predicted)
42	SPAC1751.01c	gti1	gluconate transmembrane transporter inducer Gti1
43	SPAC1782.05	ypa2	protein phosphatase type 2A regulator, PTPA family Ypa2
44	SPAC1782.08c	rex3	exonuclease Rex3 (predicted)
45	SPAC1782.09c	clp1	Cdc14-related protein phosphatase Clp1/Flp1
46	SPAC1783.05	hrp1	ATP-dependent DNA helicase Hrp1
47	SPAC17A2.12	rrp1	ATP-dependent DNA helicase/ ubiquitin-protein ligase E3 (predicted)
48	SPAC17G8.05	med20	mediator complex subunit Med20
49	SPAC17G8.07	yaf9	YEATS family histone acetyltransferase subunit Yaf9
50	SPAC17G8.09	shg1	Set1C complex subunit Shg1
51	SPAC17G8.10c	dma1	mitotic spindle checkpoint ubiquitin ligase Dma1
52	SPAC17G8.13c	mst2	histone acetyltransferase Mst2
53	SPAC17H9.10c	ddb1	Cul4-RING E3 adaptor Ddb1
54	SPAC17H9.19c	cdt2	WD repeat protein Cdt2
55	SPAC1805.14		Schizosaccharomyces specific protein
56	SPAC1805.15c	pub2	HECT-type ubiquitin-protein ligase E3 Pub2
57	SPAC1851.03	ckb1	CK2 family regulatory subunit Ckb1
58	SPAC18G6.02c	chp1	heterochromatin (HP1) family chromodomain protein Chp1
59	SPAC18G6.10	lem2	LEM domain nuclear inner membrane protein Heh1/Lem2
60	SPAC18G6.13		Schizosaccharomyces specific protein
61	SPAC1952.05	gcn5	SAGA complex histone acetyltransferase catalytic subunit Gcn5
62	SPAC19A8.10	rfp1	SUMO-targeted ubiquitin-protein ligase subunit Rfp1
63	SPAC19D5.06c	din1	RNA pyrophosphohydrolase Din1
64	SPAC19D5.11c	ctf8	Ctf18 RFC-like complex subunit Ctf8
65	SPAC19E9.02	fin1	serine/threonine protein kinase, NIMA related Fin1
66	SPAC19G12.13c	poz1	shelterin complex subunit Poz1
67	SPAC19G12.17	erh1	enhancer of rudimentary homolog Erh1
68	SPAC1B3.17	clr2	chromatin silencing protein Clr2
69	SPAC1D4.09c	rtf2	replication termination factor Rtf2
70	SPAC1D4.11c	lkh1	dual specificity protein kinase Lkh1
71	SPAC1F3.01	rrp6	exosome 3'-5' exoribonuclease subunit Rrp6
72	SPAC1F3.06c	spo15	mitotic and mieotic spindle pole body protein Spo15

73	SPAC1F7.01c	spt6	nucleosome remodeling protein Spt6
74	SPAC20G4.04c	hus1	checkpoint clamp complex protein Hus1
75	SPAC20G8.08c	fft1	SMARCAD1 family ATP-dependent DNA helicase Fft1 (predicted)
76	SPAC20H4.03c	tfs1	general transcription elongation factor TFIIS
77	SPAC20H4.10	ufd2	ubiquitin-protein ligase E4 Ufd2 (predicted)
78	SPAC21E11.03c	pcr1	transcription factor Pcr1
79	SPAC21E11.05c	cyp8	cyclophilin family peptidyl-prolyl cis-trans isomerase Cyp8
80	SPAC222.04c	ies6	Ino80 complex subunit Ies6
81	SPAC222.15	meu13	Tat binding protein 1(TBP-1)-interacting protein (TBPIP) homolog (predicted)
82	SPAC22A12.01c	pso2	DNA 5' exonuclease (predicted)
83	SPAC22E12.11c	set3	histone lysine methyltransferase Set3
84	SPAC22E12.19	snt1	Set3 complex subunit Snt1
85	SPAC22F3.02	atf31	transcription factor Atf31
86	SPAC22F3.09c	res2	MBF transcription factor complex subunit Res2
87	SPAC22F8.12c	shf1	small histone ubiquitination factor Shf1
88	SPAC22H12.02	tfg3	TFIID, TFIIF, Ino80, SWI/SNF, and NuA3 complex subunit Tfg3
89	SPAC23A1.07		ubiquitin-protein ligase E3 (predicted)
90	SPAC23C11.08	php3	CCAAT-binding factor complex subunit Php3
91	SPAC23C11.15	pst2	Clr6 histone deacetylase complex subunit Pst2
92	SPAC23C4.03	hrk1	haspin related kinase Hrk1
93	SPAC23D3.01	pdp3	PWWP domain protein, involved in chromatin remodeling (predicted)
94	SPAC23D3.09	arp42	SWI/SNF and RSC complex subunit Arp42
95	SPAC23E2.01	fep1	iron-sensing transcription factor Fep1
96	SPAC23E2.03c	ste7	arrestin family meiotic suppressor protein Ste7
97	SPAC23G3.04	ies4	Ino80 complex subunit Ies4
98	SPAC23G3.07c	snf30	SWI/SNF complex subunit Snf30
99	SPAC23G3.08c	ubp7	ubiquitin C-terminal hydrolase Ubp7
100	SPAC23G3.10c	ssr3	SWI/SNF and RSC complex subunit Ssr3
101	SPAC23H3.05c	swd1	Set1C complex subunit Swd1
102	SPAC23H4.12	alp13	MRG family Clr6 histone deacetylase complex subunit Alp13
103	SPAC24B11.10c	cfh1	SEL1/TPR repeat protein Cfh1 (predicted)
104	SPAC25A8.01c	fft3	SMARCAD1 family ATP-dependent DNA helicase Fft3
105	SPAC25H1.02	jmj1	histone demethylase Jmj1 (predicted)
106	SPAC26H5.03	pcf2	CAF assembly factor (CAF-1) complex subunit B, Pcf2
107	SPAC29A4.09		rRNA exonuclease Rrp17 (predicted)
108	SPAC29A4.18	prw1	Clr6 histone deacetylase complex subunit Prw1
109	SPAC29B12.02c	set2	histone lysine H3-K36 methyltransferase Set2
110	SPAC29B12.03	spd1	ribonucleotide reductase (RNR) inhibitor
111	SPAC29B12.06c	rcd1	CCR4-Not complex RNA-binding protein subunit Rcd1
112	SPAC29B12.08	clr5	Clr5 protein
113	SPAC2C4.07c	dis32	3'-5'-exoribonuclease activity Dis3L2
114	SPAC2F3.15	lsk1	P-TEFb-associated cyclin-dependent protein kinase Lsk1
115	SPAC2F3.16		ubiquitin-protein ligase E3, implicated in DNA repair (predicted)
116	SPAC2F7.07c	cph2	Clr6 histone deacetylase associated PHD protein Cph2
117	SPAC2F7.08c	snf5	SWI/SNF complex subunit Snf5
118	SPAC2G11.05c	rim20	BRO1 domain protein Rim20
119	SPAC2G11.10c	uba42	thiosulfate sulfurtransferase, URM1 activating enzyme E1-type Uba42 (predicted)
120	SPAC30D11.07	nth1	DNA endonuclease III
121	SPAC31A2.09c	apm4	AP-2 adaptor complex mu subunit Apm4 (predicted)
122	SPAC31A2.16	gef2	RhoGEF Gef2
123	SPAC31G5.09c	spk1	MAP kinase Spk1
124	SPAC31G5.19	abo1	ATPase with bromodomain protein
125	SPAC323.03c		Schizosaccharomyces specific protein
126	SPAC328.05	hrb1	RNA-binding protein involved in export of mRNAs Hrb1 (predicted)
127	SPAC32A11.03c	phx1	stationary phase-specific homeobox transcription factor Phx1
128	SPAC343.04c	gid7	GID complex subunit Gid7 (predicted)
129	SPAC343.11c	msc1	Swr1 complex subunit Msc1
130	SPAC343.18	rfp2	SUMO-targeted ubiquitin-protein ligase subunit Rfp2
131	SPAC3A11.05c	kms1	meiotic spindle pole body KASH domain protein Kms1
132	SPAC3C7.08c	elf1	AAA family ATPase Elf1
133	SPAC3F10.10c	map3	pheromone M-factor receptor Map3
134	SPAC3F10.12c		transcription factor (predicted)
135	SPAC3G6.01	hrp3	ATP-dependent DNA helicase Hrp3
136	SPAC3G6.06c	rad2	FEN-1 endonuclease Rad2
137	SPAC3G6.11	chl1	ATP-dependent DNA helicase Chl1 (predicted)
138	SPAC3G9.07c	hos2	histone deacetylase (class I) Hos2
139	SPAC3H1.11	hsr1	transcription factor Hsr1
140	SPAC3H1.12c	snt2	Lid2 complex PHD finger subunit Snt2
141	SPAC3H8.08c		transcription factor (predicted)
142	SPAC4A8.09c	cwf21	complexed with Cdc5 protein Cwf21
143	SPAC4F8.11	sea2	SEA complex WD repeat subunit Sea2 (predicted)
144	SPAC4G9.06c	chz1	histone H2A-H2B dimer chaperone Chz1 (predicted)
145	SPAC4H3.02c	swc3	Swr1 complex subunit Swc3
146	SPAC4H3.05	srs2	ATP-dependent DNA helicase, UvrD subfamily

147	SPAC56F8.16	esc1	transcription factor Esc1 (predicted)
148	SPAC57A10.09c	nhp6	High-mobility group non-histone chromatin protein (predicted)
149	SPAC5D6.02c	mug165	Clr6 histone deacetylase complex subunit Mug165
150	SPAC5D6.08c	mes1	meiotic APC inhibitor Mes1
151	SPAC630.14c	tup12	transcriptional corepressor Tup12
152	SPAC631.02	bdf2	BET family double bromodomain protein Bdf2
153	SPAC637.09	rex1	3'-5'- exoribonuclease Rex1 (predicted)
154	SPAC644.14c	rad51	RecA family recombinase Rad51/Rhp51
155	SPAC664.01c	swi6	heterochromatin (HP1) family chromodomain protein Swi6
156	SPAC664.02c	arp8	Ino80 complex actin-like protein Arp8
157	SPAC664.07c	rad9	checkpoint clamp complex protein Rad9
158	SPAC664.15	caf4	CCR4-Not complex subunit Caf4/Mdv1 (predicted)
159	SPAC688.06c	slx4	structure-specific endonuclease subunit Slx4
160	SPAC694.06c	mrc1	claspin, Mrc1
161	SPAC6B12.05c	ies2	Ino80 complex subunit Ies2
162	SPAC6B12.07c		ubiquitin-protein ligase E3 with SPX domain, human LORNRF1 ortholog (predicted)
163	SPAC6B12.14c		conserved fungal protein
164	SPAC6B12.16	meu26	DUF4451 family conserved fungal protein
165	SPAC6F12.09	rdp1	RNA-directed RNA polymerase Rdp1
166	SPAC6F6.09	eaf6	Mst2/NuA4 histone acetyltransferase complex subunit Eaf6
167	SPAC6G9.03c	mug183	histone H3.3 H4 heterotetramer chaperone Rtt106-like (predicted)
168	SPAC6G9.10c	sen1	ATP-dependent 5' to 3' DNA/RNA helicase Sen1
169	SPAC6G9.16c	xrc4	XRCC4 nonhomologous end joining factor Xrc4
170	SPAC7D4.04	atg11	autophagy associated protein Atg11
171	SPAC7D4.14c	iss10	NURS complex subunit Iss10
172	SPAC821.07c	moc3	transcription factor Moc3
173	SPAC823.03	ppk15	serine/threonine protein kinase Ppk15 (predicted)
174	SPAC824.04	swd22	mRNA cleavage and polyadenylation specificity factor complex, WD repeat protein Swd22
175	SPAC890.07c	rmt1	type I protein arginine N-methyltransferase Rmt1
176	SPAC8C9.14	prr1	transcription factor Prr1
177	SPAC8C9.17c	spc34	DASH complex subunit Spc34
178	SPAC8F11.03	msh3	MutS protein homolog 3
179	SPAC9E9.08	rad26	ATRIP, ATR checkpoint kinase regulatory subunit Rad26
180	SPAC9E9.10c	cbh1	CENP-B homolog Cbh1
181	SPAP14E8.02	tos4	chromatin binding FHA domain protein Tos4 (predicted)
182	SPAP27G11.15	slx1	structure-specific endonuclease catalytic subunit Slx1
183	SPAP32A8.03c	bop1	ubiquitin-protein ligase E3, human RNF126 ortholog (predicted)
184	SPAP8A3.02c	ofd2	histone H2A dioxygenase Ofd2
185	SPAPB1E7.02c	mcl1	DNA polymerase alpha accessory factor Mcl1
186	SPAPB24D3.01	toe3	transcription factor (predicted)
187	SPAPB2B4.03	cig2	G1/S-specific B-type cyclin Cig2
188	SPBC1105.04c	cbp1	CENP-B homolog
189	SPBC119.08	pmk1	MAP kinase Pmk1
190	SPBC119.14	rti1	Rad22 homolog Rti1
191	SPBC1198.11c	reb1	RNA polymerase I transcription termination factor/ RNA polymerase II transcription factor Reb1
192	SPBC11B10.05c	rsp1	random septum position protein, DNAJ domain protein Rsp1
193	SPBC11B10.08		WW domain containing conserved fungal protein
194	SPBC11B10.10c	pht1	histone H2A variant H2A.Z, Pht1
195	SPBC1347.07	rex2	RNA exonuclease (predicted)
196	SPBC13E7.08c	leo1	RNA polymerase II associated Paf1 complex subunit Leo1
197	SPBC13G1.08c	ash2	Ash2-trithorax family protein
198	SPBC146.06c	fan1	DNA repair protein Fan1
199	SPBC14C8.17c	spt8	SAGA complex subunit Spt8
200	SPBC14F5.07	doa10	ER ubiquitin-protein ligase E3 Doa10 (predicted)
201	SPBC15C4.01c	oca3	TPR repeat protein Oca3/ ER membrane protein complex Ecm2 (predicted)
202	SPBC15C4.06c		ubiquitin-protein ligase E3 Meu34, human RNF13 family homolog, unknown biological role (predicted)
203	SPBC15D4.03	slm9	histone H3.3 H4 chaperone, hira family Slm9
204	SPBC1604.09c	rex4	exoribonuclease Rex4 (predicted)
205	SPBC1604.16c		RNA-binding protein, G-patch type, human GPANK1 ortholog
206	SPBC1685.08	cti6	histone deacetylase complex ubiquitin-like protein ligase subunit Cti6
207	SPBC16A3.07c	nrm1	MBF complex corepressor Nrm1
208	SPBC16A3.19	eaf7	histone acetyltransferase complex subunit Eaf7
209	SPBC16D10.07c	sir2	Sirtuin family histone deacetylase Sir2
210	SPBC16E9.11c	pub3	HECT-type ubiquitin-protein ligase E3 Pub3 (predicted)
211	SPBC16E9.12c	pab2	poly(A) binding protein Pab2
212	SPBC16G5.03	mrz1	ubiquitin-protein ligase E3/SUMO transferase, Topors, possibly associated with DNA damage (predicted)
213	SPBC16G5.15c	fkh2	forkhead transcription factor Fkh2
214	SPBC16G5.17		transcription factor, zf-fungal binuclear cluster type (predicted)
215	SPBC1703.04	mlh1	MutL family protein Mlh1 (predicted)
216	SPBC1703.14c	top1	DNA topoisomerase I
217	SPBC1709.11c	png2	ING family histone acetyltransferase complex PHD-type zinc finger subunit Png2
218	SPBC1711.14	rec15	meiotic recombination protein Rec15
219	SPBC1718.02	hop1	linear element associated protein Hop1
220	SPBC1734.06	rhp18	Rad18 homolog ubiquitin protein ligase E3, Rhp18

221	SPBC1734.15	rsc4	RSC complex subunit Rsc4
222	SPBC1773.16c		transcription factor, zf-fungal binuclear cluster type(predicted)
223	SPBC1778.10c	ppk21	serine/threonine protein kinase Ppk21 (predicted)
224	SPBC17D11.04c	nto1	histone acetyltransferase complex PHD finger subunit Nto1 (predicted)
225	SPBC17G9.05	rct1	cyclophilin family peptidyl-prolyl cis-trans isomerase, RRM-containing Rct1
226	SPBC18H10.06c	swd2	Set1C complex subunit Swd2.1
227	SPBC18H10.15	cdk11	serine/threonine protein kinase Cdk11
228	SPBC19C7.02	ubr1	N-end-recognizing protein, UBR ubiquitin-protein ligase E3 Ubr1
229	SPBC1A4.03c	top2	DNA topoisomerase II
230	SPBC1D7.03	clg1	cyclin-like protein involved in autophagy Clg1 (predicted)
231	SPBC1D7.04	mlo3	RNA binding protein Mlo3
232	SPBC20F10.05	nrl1	RNAi-mediated silencing protein, human NRDE2 ortholog Nrl1
233	SPBC20F10.10	psl1	cyclin pho85 family Psl1 (predicted)
234	SPBC215.03c	csn1	COP9/signalosome complex subunit Csn1
235	SPBC215.06c		nucleolar RNA-binding protein, human LYAR homolog, implicated in transcriptional regulation
236	SPBC215.07c	pdp2	PWWP domain protein Pdp2 (predicted)
237	SPBC216.05	rad3	ATR checkpoint kinase Rad3
238	SPBC216.06c	swi1	replication fork protection complex subunit Swi1
239	SPBC21B10.13c	yox1	MBF complex corepressor Yox1
240	SPBC21C3.02c	dep1	Sds3-like family protein Dep1
241	SPBC21C3.20c	git1	C2 domain protein Git1
242	SPBC21D10.09c	rkr1	RQC complex ubiquitin-protein ligase E3 Rkr1 (predicted)
243	SPBC21D10.10	bdc1	bromodomain protein Bdc1
244	SPBC23E6.02	rrp2	ATP-dependent DNA helicase/ ubiquitin-protein ligase E3 (predicted)
245	SPBC23E6.09	ssn6	transcriptional corepressor Ssn6
246	SPBC23G7.13c		plasma membrane urea transmembrane transporter (predicted)
247	SPBC24C6.05	sec28	coatomer epsilon subunit (predicted)
248	SPBC25B2.08		Schizosaccharomyces pombe specific protein
249	SPBC26H8.09c	snf59	SWI/SNF complex subunit Snf59
250	SPBC28E12.02		RNA-binding protein
251	SPBC28F2.07	sfr1	Swi five-dependent recombination mediator Sfr1
252	SPBC28F2.10c	ngg1	SAGA complex subunit Ngg1/Ada3
253	SPBC29A10.03c	pcf1	CAF assembly factor (CAF-1) complex large subunit Pcf1
254	SPBC29A10.05	exo1	exonuclease I Exo1
255	SPBC29A10.14	rec8	meiotic cohesin complex subunit Rec8
256	SPBC29A3.03c	gid2	GID complex ubiquitin-protein ligase E3 subunit Gid2/Rmd5 (predicted)
257	SPBC29A3.05	vps71	Swr1 complex subunit Vps71
258	SPBC29A3.13	pdp1	PWWP domain protein Pdp1
259	SPBC29B5.01	atf1	transcription factor, Atf-CREB family Atf1
260	SPBC2A9.04c	san1	sir antagonist, ubiquitin-protein ligase E3
261	SPBC2D10.11c	nap2	histone H2A-H2B chaperone Nap2
262	SPBC2D10.17	clr1	SHREC complex intermodule linker subunit Clr1
263	SPBC2F12.09c	atf21	transcription factor, Atf-CREB family Atf21
264	SPBC2F12.12c	cay1	cactin, spliceosome complex subunit
265	SPBC2G2.06c	apl1	AP-2 adaptor complex beta subunit Apl1 (predicted)
266	SPBC2G2.14	csi1	mitotic centromere-SPB clustering protein Csi1
267	SPBC2G5.02c	ckb2	CK2 family regulatory subunit Ckb2 (predicted)
268	SPBC30B4.04c	sol1	SWI/SNF complex subunit Sol1
269	SPBC30D10.10c	tor1	serine/threonine protein kinase Tor1
270	SPBC31F10.07	lsb5	actin cortical patch component Lsb5 (predicted)
271	SPBC31F10.10c	mub1	Armadillo-type fold protein, zf-MYND type zinc finger protein, Mub1-Rad6-Ubr2 ubiquitin ligase complex Mub1 (predicted)
272	SPBC31F10.13c	hip1	histone H3.3 H4 chaperone, hira family Hip1
273	SPBC31F10.14c	hip3	HIRA interacting protein Hip3
274	SPBC32F12.07c		membrane associated ubiquitin-protein ligase E3, MARCH family (predicted)
275	SPBC32H8.06	mug93	TPR repeat protein, meiotically spliced
276	SPBC337.03	rhn1	RNA polymerase II transcription termination factor homolog
277	SPBC342.05	crb2	DNA repair protein Rad9 homolog Crb2
278	SPBC342.06c	rtt109	RTT109 family histone lysine acetyltransferase
279	SPBC354.03	swd3	WD repeat protein Swd3
280	SPBC354.05c	sre2	membrane-tethered transcription factor Sre2
281	SPBC365.10	arp5	Ino80 complex actin-like protein Arp5
282	SPBC36B7.05c	pib1	endosomal and vacuolar ubiquitin-protein ligase E3/phosphatidylinositol(3)-phosphate binding protein Pib1
283	SPBC36B7.08c	ccp1	histone chaperone, CENP-A nucleosome disassembly Ccp1
284	SPBC3B8.02	php5	CCAAT-binding factor complex subunit Php5
285	SPBC3D6.04c	mad1	mitotic spindle checkpoint protein Mad1
286	SPBC3D6.09	dpb4	DNA polymerase epsilon subunit Dpb4
287	SPBC3H7.13	far10	SIP/FAR complex FHA domain subunit Far10/Csc1
288	SPBC4.05	mlo2	ubiquitin protein ligase E3 component human N-recognin 7 homolog Mlo2
289	SPBC428.06c	rxt2	histone deacetylase complex subunit Rxt2
290	SPBC428.08c	clr4	histone lysine H3 methyltransferase Clr4
291	SPBC4B4.03	rsc1	RSC complex subunit Rsc1
292	SPBC4C3.12	1-Sep	forkhead transcription factor Sep1
293	SPBC530.08		membrane-tethered transcription factor (predicted)
294	SPBC530.14c	dsk1	SR protein-specific kinase Dsk1

295	SPBC543.07	pek1	MAP kinase kinase Pek1
296	SPBC56F2.03	arp10	dynactin complex actin-like protein Arp10 (predicted)
297	SPBC56F2.05c		transcription factor (predicted)
298	SPBC582.04c	dsh1	RNAi protein, Dsh1
299	SPBC582.06c	mcp6	horsetail movement protein Hrs1/Mcp6
300	SPBC609.05	pob3	histone H2A-H2B chaperone, FACT complex subunit Pob3
301	SPBC651.11c	apm3	AP-3 adaptor complex subunit Apm3 (predicted)
302	SPBC660.06		WW domain containing conserved fungal protein
303	SPBC660.14	mik1	mitotic inhibitor kinase Mik1
304	SPBC6B1.04	mde4	microtubule-site clamp monopolin complex subunit Mde4
305	SPBC6B1.06c	ubp14	Lys48-specific deubiquitinase Ubp14
306	SPBC725.02	mpr1	histidine-containing response regulator phosphotransferase Mpr1
307	SPBC725.11c	php2	CCAAT-binding factor complex subunit Php2
308	SPBC776.02c	dis2	serine/threonine protein phosphatase PP1, Dis2
309	SPBC776.16	mis20	centromere protein Mis20/Eic2
310	SPBC800.03	clr3	histone deacetylase (class II) Clr3
311	SPBC83.03c	tas3	RITS complex subunit 3
312	SPBC902.02c	ctf18	Ctf18 RFC-like complex subunit Ctf18
313	SPBC902.04	rmn1	RNA-binding protein
314	SPBC902.06	mto2	gamma tubulin complex linker Mto2
315	SPBP16F5.03c	tra1	SAGA complex phosphatidylinositol pseudokinase Tra1
316	SPBP22H7.05c	abo2	ATPase with bromodomain protein (predicted)
317	SPBP23A10.05	ssr4	SWI/SNF and RSC complex subunit Ssr4
318	SPBP35G2.08c	air1	TRAMP complex zinc knuckle subunit Air1
319	SPBP35G2.10	mit1	SHREC complex ATP-dependent DNA helicase subunit Mit1
320	SPBP35G2.13c	swc2	Swr1 complex subunit Swc2
321	SPBP8B7.07c	set6	histone lysine methyltransferase Set6 (predicted)
322	SPBP8B7.23	rnf10	ubiquitin-protein ligase E3 (predicted)
323	SPBP8B7.28c	stc1	CLRC ubiquitin ligase complex linker protein, LIM-like Stc1
324	SPCC1020.12c	xap5	xap-5-like protein
325	SPCC11E10.08	rik1	CLRC ubiquitin ligase complex WD repeat protein Rik1
326	SPCC1223.13	cbf12	CBF1/Su(H)/LAG-1 family transcription factor Cbf12
327	SPCC1235.05c	fft2	SMARCAD1 family ATP-dependent DNA helicase Fft2 (predicted)
328	SPCC1235.09	hif2	Set3 complex subunit Hif2
329	SPCC1235.12c	mug146	Schizosaccharomyces specific protein Mug46
330	SPCC1259.04	iec3	Ino80 complex subunit Iec3
331	SPCC1259.07	rxt3	transcriptional regulatory protein Rxt3
332	SPCC126.02c	pku70	Ku domain protein Pku70
333	SPCC126.04c	sgf73	SAGA complex deubiquitinating submodule subunit Sgf73
334	SPCC126.07c	asr1	ubiquitin-protein ligase E3 Asr1 (predicted)
335	SPCC126.11c		RNA-binding protein, rrm type
336	SPCC126.13c	sap18	splicing factor Sap18 (predicted)
337	SPCC132.02	hst2	Sirtuin family histone deacetylase Hst2
338	SPCC1393.02c	spt2	non-specific DNA binding protein Spt2 (predicted)
339	SPCC1393.05	ers1	RNA-silencing factor Ers1
340	SPCC1442.13c	sqs2	R3H and G-patch domain protein Sqs2
341	SPCC1450.02	bdf1	Swr1 complex bromodomain subunit Bdf1
342	SPCC1450.03	utp502	ribonucleoprotein (RNP) complex Utp502 (predicted)
343	SPCC1494.03	arz1	human RAP1 GTPase-GDP dissociation stimulator ortholog, Zfs1 target number 1
344	SPCC162.11c	urk1	uridine kinase/uracil phosphoribosyltransferase (predicted)
345	SPCC1620.14c	snf22	ATP-dependent DNA helicase Snf22
346	SPCC1682.13	laf2	Clr6 associated factor 2, Laf2
347	SPCC16C4.11	pef1	Pho85/PhoA-like cyclin-dependent kinase Pef1
348	SPCC1739.03	hrr1	Helicase Required for RNAi-mediated heterochromatin assembly Hrr1
349	SPCC1739.05	set5	histone lysine methyltransferase Set5 (predicted)
350	SPCC1739.07	cti1	exosome C1D family subunit Cti1
351	SPCC1739.12	ppe1	serine/threonine protein phosphatase Ppe1
352	SPCC1753.03c	rec7	meiotic recombination protein Rec7
353	SPCC1840.04	pca1	metacaspase Pca1
354	SPCC188.07	ccq1	shelterin complex HEAT repeat subunit Ccq1
355	SPCC188.13c	dcr1	dicer
356	SPCC18B5.03	wee1	M phase inhibitor protein kinase Wee1
357	SPCC18B5.07c	nup61	nucleoporin Nup61
358	SPCC24B10.07	gad8	AGC family protein kinase Gad8
359	SPCC24B10.08c	ada2	SAGA complex subunit Ada2
360	SPCC24B10.14c	xlf1	XRCC4-like nonhomologous end joining factor, Cernunnon Xlf1/Nej1
361	SPCC24B10.19c	nts1	Clr6 histone deacetylase complex subunit Nts1
362	SPCC297.03	ssp1	Ca2+/calmodulin-dependent (CaMMK)-like protein kinase Ssp1
363	SPCC297.04c	set7	histone lysine H3-K37 methyltransferase Set7
364	SPCC2H8.05c	dbl1	double strand break localizing Dbl1
365	SPCC306.04c	set1	histone lysine H3-K4 methyltransferase Set1
366	SPCC31H12.08c	ccr4	CCR4-Not complex 3'-5'-exoribonuclease subunit Ccr4
367	SPCC330.01c	rhp16	Rad16 homolog ATP-dependent DNA helicase/ ubiquitin protein ligase E3 Rhp16
368	SPCC330.02	rhp7	Rad7 homolog Rhp7

369	SPCC338.16	pof3	F-box protein Pof3
370	SPCC364.02c	bis1	splicing factor Bis1
371	SPCC364.06	nap1	histone H2A-H2B chaperone Nap1
372	SPCC417.07c	mto1	gamma tubulin complex linker Mto1
373	SPCC417.09c		transcription factor (predicted)
374	SPCC4B3.12	set9	histone lysine H4-K20 methyltransferase Set9
375	SPCC4G3.15c	not2	CCR4-Not complex NOT box subunit Not2
376	SPCC4G3.19	alp16	gamma tubulin complex subunit Alp16
377	SPCC548.05c	dbl5	ubiquitin-protein ligase E3 Dbl5
378	SPCC550.12	arp6	actin-like protein Arp6
379	SPCC550.15c	rei1	ribosome biogenesis protein Rei1 (predicted)
380	SPCC553.04	cyp9	WD repeat containing cyclophilin family peptidyl-prolyl cis-trans isomerase Cyp9 (predicted)
381	SPCC576.13	swc5	Swr1 complex subunit Swc5
382	SPCC594.05c	spf1	Set1C ubiquitin-protein ligase E3 subunit Spf1
383	SPCC61.02	spt3	SAGA complex subunit Spt3
384	SPCC613.12c	raf1	CLRC ubiquitin ligase complex WD repeat subunit Raf1/Dos1
385	SPCC622.15c		Schizosaccharomyces specific protein
386	SPCC622.16c	epe1	Jmjc domain chromatin associated protein Epe1
387	SPCC622.19	jmj4	peptidyl-lysine 3-dioxygenase activity jmj4 (predicted)
388	SPCC645.13	bye1	transcription elongation regulator Bye1 (predicted)
389	SPCC663.11	saf1	splicing associated factor Saf1
390	SPCC663.12	cid12	poly(A) polymerase Cid12
391	SPCC736.08	cbf11	CBF1/Su(H)/LAG-1 family transcription factor Cbf11
392	SPCC736.11	ago1	argonaute
393	SPCC757.09c	rnc1	KH domain RNA-binding protein Rnc1
394	SPCC895.06	elp2	elongator complex WD repeat protein Elp2 (predicted)
395	SPCC895.07	alp14	TOG/XMAP215 microtubule plus end tracking polymerase Alp14
396	SPCC970.07c	raf2	CLRC ubiquitin ligase complex subunit Raf2

Table 2: Strain table

Strain	Genotype
PAS075	Locus2::*ade6p::3xE2C:hygMX* at Locus2 (between SPBC1711.11 and SPBC1711.12)
PM003	Wild-type strain: h(+); ura4-D18; leu1-32; ade6-M216; his7-366
PM006	972 h- wild-type
PAS193	ΔK::*ade6p:*mKO2; *ade6p:*SF-GFP between *REIII* and mat3M; *ade6p:*3xE2C: *hygMX* at Locus2; *clr4::kanMX,* h(-)
PAS216	*cenH*::*ade6p:*SF-GFP(Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C: *hygMX* at Locus2; *clr4::kanMX,* h90
PAS217	*cenH*: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C: *hygMX* at Locus2, h90
PAS231	*ura4::natMX:dh:ade6p:*SF-GFP, *ade6p:*mKO2 3 kb, *leu1::ade6p:*3xE2C: *hygMX*
PAS331	*cenH*:: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C:*hygMX* at Locus2; Δ*REIII*::REIII(Δs1, Δs2) in *clr4::kanMX,* h90
PAS332	*cenH*:: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C:*hygMX* at Locus2; Δ*REIII*::REIII(Δs1, Δs2), h90
PAS482	ΔK::*ade6p:*mKO2; *ade6p:* SF-GFP between *REIII* and mat3M; *ade6p:*3xE2C: *hygMX* at Locus2, h(-); 'OFF' allele
PAS795	*cenH*: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C: *hygMX* at Locus2; *fkh2::natMX*
PAS796	*cenH*: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C: *hygMX* at Locus2; *prw1::kanMX*
PAS797	*cenH*: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C: *hygMX* at Locus2; *png3::kanMX*
PAS798	*cenH*:: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C:*hygMX* at Locus2; Δ*REIII*::REIII(Δs1, Δs2); *fkh2::natMX*
PAS799	*cenH*:: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C:*hygMX* at Locus2; Δ*REIII*::REIII(Δs1, Δs2); *prw1::kanMX*
PAS800	*cenH*:: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C:*hygMX* at Locus2; Δ*REIII*::REIII(Δs1, Δs2); *png3::kanMX*
PAS803	ΔK::*ade6p:*mKO2; *ade6p:* SF-GFP between *REIII* and mat3M; *ade6p:*3xE2C: *hygMX* at Locus2; 'OFF' allele; *fkh2::natMX*
PAS804	ΔK::*ade6p:*mKO2; *ade6p:* SF-GFP between *REIII* and mat3M; *ade6p:*3xE2C: *hygMX* at Locus2; 'OFF' allele; *prw1::kanMX*
PAS805	ΔK::*ade6p:*mKO2; *ade6p:* SF-GFP between *REIII* and mat3M; *ade6p:*3xE2C: *hygMX* at Locus2; 'OFF' allele; *png3::kanMX*
PAS808	*cenH*: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C: *hygMX* at Locus2; *fkh2::natMX; prw1::kanMX* by cross
PAS809	*cenH*: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C: *hygMX* at Locus2; *fkh2::natMX; prw1::kanMX* by sequential knockout
PAS810	*cenH*:: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C:*hygMX* at Locus2; Δ*REIII*::REIII(Δs1, Δs2); *fkh2::natMX; prw1::kanMX* by cross
PAS811	*cenH*:: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C:*hygMX* at Locus2; Δ*REIII*::REIII(Δs1, Δs2); *fkh2::natMX; prw1::kanMX* by sequential knockout
PAS813	*cenH*:: *ade6p:*SF-GFP (Kint2); *mat3m(Eco*RV):: *ade6p:*mKO2; *ade6p:*3xE2C:*hygMX* at Locus2; Δ*REIII*::REIII(Δs1, Δs2); *apm3::natMX*
PAS816	apl5:SF-GFP:*hygMX*; Swi6:E2C:*kanMX*
PAS817	apm3:SF-GFP:*hygMX;* Swi6:E2C:*kanMX*