An adolescent sensitive period of heightened frontal top-down projection activity promotes local circuit integration and attentional behavior

Elisa M Nabel [1,2,3,4,5,6], Yury Garkun [1,2,3,4,5,6], Hiroyuki Koike [1,2,3,4,5], Masato Sadahiro [1,2,3,4,5], Ana Liang[1,2,3,4,5], Kevin J Norman[1,2,3,4,5], Giulia Taccheri[1,2,3,4,5], Michael Demars [1,2,3,4,5], Susanna Im[1,2,3,4,5], Keaven Caro[1,2,3,4,5], Sarah Lopez[1,2,3,4,5], Julia Bateh[1,2,3,4,5], Patrick R. Hof [2,3,5], Roger L. Clem [1,2,5], Hirofumi Morishita [*,1,2,3,4,5]

[1]Department of Psychiatry, [2]Nash Family Department of Neuroscience, [3]Department of Ophthalmology, [4]Mindich Child Health and Development Institute, [5]Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, New York, NY 10029 USA

[6]These authors contributed equally.

*Corresponding Author and Lead Contact: Hirofumi Morishita, M.D., Ph.D.

One Gustave L. Levy Place Box 1230, New York, NY 10029 USA,

Tel.: +1 212 824 8975

Fax: +1 646 537 9584

E-mail address: hirofumi.morishita@mssm.edu

Abstract

Frontal top-down cortical neurons projecting to sensory cortical regions are well-positioned to integrate long-range inputs with local circuitry in frontal cortex to implement top-down attentional control of sensory regions. How adolescence contributes to the maturation of top-down neurons and associated local/long-range input balance, and the establishment of attentional control is poorly understood. Here we combine projection-specific electrophysiological and rabies-mediated input mapping in mice to uncover adolescence as a developmental stage when frontal top-down neurons projecting from the anterior cingulate to visual cortex are highly functionally integrated into local excitatory circuitry and have heightened activity compared to adulthood. Chemogenetic suppression of top-down neuron activity selectively during adolescence, but not later periods, produces long-lasting visual attentional behavior deficits, and results in excessive loss of local excitatory inputs in adulthood. Our study reveals an adolescent sensitive period when top-down neurons integrate local circuits with long-range connectivity to produce attentional behavior.

Main Text

Introduction

Cognitive function relies on cohesive activity of large-scale networks of functionally interconnected yet structurally separate cortical regions. Corticocortical projections facilitate such network dynamics by integrating information from various inputs, not only from local cortical microcircuits but also long-range cortical and subcortical connections, and then transmitting this information to other cortical regions [1]. Developing a balance of local and long-range inputs in a cortical region is necessary for neuronal processing underlying mature cognition. Altricial species, including both humans and mice, are born with relatively immature brains that follow a postnatal developmental trajectory before adult cognitive capacities are established. During development, initial axonal and dendritic branching is followed by a stage of synaptic overproduction and reduction to an adult plateau level [2]. Particularly in the frontal cortex, this postnatal maturational process is integral for corticocortical projection neurons to integrate local inputs with long-range distal inputs carrying sensory, motor, and cognitive information and consequently to execute essential complex behaviors. Notably, deficits in frontal cortex long-range connectivity are pervasively reported in neurodevelopmental disorders and psychiatric disorders with developmental origins [3-8], and often emerge following childhood and adolescence [9-12]. Yet limited knowledge of normal processes governing corticocortical circuit development hinders further pathophysiological insight into these conditions. Clinical efforts to diagnose, treat, and monitor such diseases will remain limited to symptomatic approaches until a basic physiological understanding of the maturational processes and their vulnerabilities are better established [13].

Of particular significance among corticocortical projections is the top-down feedback projection from frontal cortex to sensory visual cortex [14,15]. Non-human primate studies demonstrate that a frontal projection to visual cortical areas executes attention controlled downstream visual processing [14]. In mice, the anterior cingulate cortex area (ACA) and adjacent secondary motor cortex area (MOs) of the frontal cortex has been shown to project to visual cortical areas including primary visual cortex (VIS) [16-18]. Optogenetic terminal stimulation of ACA neurons projecting to VIS (ACA_{VIS}) in VIS can enhance the signal to noise ratio of neural response to preferred visual orientation stimuli and therefore increase the visual cortex ability to detect visual features—a hallmark feature of visual attention [16]. Of note, disruption of functional connectivity between frontal

cortex and visual cortex during visual attention tasks is implicated in several neurodevelopmental disorders, including schizophrenia, autism spectrum disorder, and attention deficit hyperactivity disorder [3,19-21]. Here we aimed to identify the circuit basis of adolescent maturation of mouse top-down ACA_{VIS} projection neurons and interrogate the contribution of this developmental window to adult circuitry and cognitive behavior.

Among various cognitive behaviors, we focused on visual attentional behavior by utilizing the five-choice serial reaction time task (5CSRTT), which is known to require ACA activity [22], to interrogate the functionality of ACA_{VIS} in the context of visual attention. Back-translated from the human continuous performance attention task in humans, the 5CSRTT has been extensively used to examine visual attention in rodents [23-27]. The task requires the subject to touch one of five screens when it flashes in order to receive a reward. Because the stimulus is presented pseudorandomly across these five visual areas, the subjects must divide their attention across the areas. The number of times the subject touches the screen that flashed, in relationship to missed and incorrect touches, reflects a level of attentional processing. Further, taxing of attentional capacity is demonstrated if performance worsens with shorter flash duration. Alongside these primary measures of attentional performance, additional 5CSRTT basic task parameters provide relatively independent indices of cognitive control[28]. The 5CSRTT is currently being applied in drug discovery efforts, particularly toward identifying novel drug treatments for conditions associated with attention deficits. Here, we integrated electrophysiology, input mapping, chemogenetics, and a translational visual attentional behavioral task in circuit-specific manner in mice, and revealed an adolescent sensitive period when top-down cortical circuits refine their local excitatory inputs in an activity-dependent manner to establish adult visual attention behavior.

Results

Greater adolescent excitatory drive onto frontal top-down ACA$_{VIS}$ projection neurons diminishes by adulthood

We first characterized the localization of top-down projection neurons projecting from frontal cortex to VIS in the adult mice. Retrograde viral labeling of these projection neurons to VIS revealed that a majority of labeled medial frontal neurons arise from dorsal ACA (ACAd) (65%), followed by adjacent secondary motor area (MOs) (27%) **(Figure S1 A-C)**. Quantification of cortical layer distribution of these neurons showed approximately 81% arising from layers V/VI and 19% arising from layers II/III in adulthood **(Figure S1 D)**. In the following we thus focused on characterizing top-down projection neurons from ACA to VIS (ACA$_{VIS}$ projection neurons).

We next assessed changes in synaptic drive onto top-down ACA$_{VIS}$ projection neurons during postnatal development across adolescent age. Adolescence is characterized by various behavioral and physiological changes that differs substantially from both infancy and adulthood, and broadly correspond to a period after the completion of weaning (p22-25) before the beginning of adulthood (p55-65)[29,30]. Of note, adolescent period around p32 is characterized by an increase in social play and affiliative behaviors [31]. We performed whole-cell patch clamp recordings of miniature excitatory (mEPSC) and inhibitory (mIPSC) postsynaptic currents from ACA$_{VIS}$ projection neurons in juvenile (~p20: p18-p22), adolescent (~p32: p29-p37), and adult (~p78: p65-p100) mice. Fluorescent retrobeads were injected into the VIS 4-6 days prior to slice experiments to visualize ACA$_{VIS}$ projection neurons **(Figure 1A).** While there was no significant change in mEPSC frequency between the juvenile to adolescent period, we observed a significant decrease in mEPSC frequency from adolescence to adulthood **(Figure 1B and D)**, suggesting adolescence as a period of heightened excitatory drive. The cumulative distribution of mEPSC amplitudes shifted toward higher values between juvenile and adolescent groups with no changes in distribution between adolescents and adults, and mean amplitude of mEPSC also did not differ between the adolescent period and adulthood **(Figure 1B and E)**. mIPSC frequency showed no statistically significant changes across ages **(Figure 1C and F)**. Similar to mEPSC, the cumulative distribution of mIPSC amplitudes shifted toward higher values between juvenile and adolescent stages with no changes in distribution between adolescents and adults, and mean amplitude of mIPSC did not differ among three groups **(Figure 1E and G)**. Overall, our data show a selective shift in the frequency of excitatory postsynaptic currents between the

adolescent and adult groups with no changes in amplitude after adolescence. In contrast, mIPSC frequency and

amplitude showed no significant changes across age groups. Collectively, our study highlights adolescence as

a period of greater adolescent excitatory drive onto frontal top-down ACA_{VIS} projection neurons.

Local presynaptic inputs onto top-down ACA_{VIS} projection neurons are more extensive during

adolescence

To characterize this shift in excitatory drive between adolescence and adulthood further, we next sought

to identify the input source onto frontal top-down ACA_{VIS} neurons present during adolescence that is absent by

adulthood. We employed rabies mediated monosynaptic input mapping that would allow us to detect brain

regions providing inputs to frontal ACA_{VIS} projection neurons and to quantify if neural input from these brain

regions changed across development. We used a pseudotyped and genetically modified rabies virus

(RbVdG(EnvA)-eGFP) that allows visualization of "starter" cells by requiring the cognate receptor, avian tumor

virus receptor A (TVA), to be expressed for cell entry and rabies glycoprotein (RbG) for cell exit [32]. Specifically,

this system restricts expression of GFP only in "starter" ACA_{VIS} projection neurons expressing TVA and their

presynaptic inputs. ACA_{VIS} circuit specific TVA and RbG expression was conferred by an intersectional viral

approach that introduced Cre-dependent TVA and RbG vectors into the ACA and retrograde canine adenovirus

type 2 (CAV-2) Cre in VIS. This approach targeted ACA_{VIS} neurons mainly in deep layers consistent with the

characteristic of CAV-2 transduction pattern[33], but also some in superficial layers comparable to the labeling

pattern of retrobeads or retro AAV2 **(Figure 2A)**. After allowing 2.5 weeks for gene expression, monosynaptic

input labeling was accomplished by injection of RbVdG(EnvA)-eGFP. We examined expression 7 days later and

estimate that monosynaptic input labeling at adolescent and adult time points represent connectivity around ages

p34 and p80 respectively **(Figure 2B)**. Because the p12 and p60 brain differed in size when stereotactic surgery

was performed, we first characterized and compared the distribution of starter cells initiating the retrograde

transfer of rabies virus, identified as double-labelled mCherry+ eGFP+ **(Figure 2C)** between adolescent and

adult groups and found no difference in their anterior-posterior distribution **(Figure 2D)**, or in the absolute number

of starter cells per animal across the two age groups **(Figure 2E)**. An additional experiment verified that the

distribution of ACA_{VIS} neurons across the cortical layers does not change across the age groups **(Figure S1D)**.

Consequently, we were able to directly compare the number of monosynaptic input neurons from each

anatomical location between adolescent and adult groups by normalizing to the number of starter cells for each

animal.

In each animal, all monosynaptic input neurons for evenly sampled coronal slices were counted and

assigned to the corresponding brain regions of the Allen Brain Atlas hierarchical neuroanatomic ontology

(http://mouse.brain-map.org/static/atlas) **(Table1)**. To allow a direct comparison of input number and distribution

across animals and developmental groups, for each animal, inputs were normalized by dividing by the number

of starter cells. At the highest hierarchical level of organization, we observed more inputs in the cortical plate

(CTXpl) in the adolescent group with no differences observed in the brainstem, where all inputs were limited to

the diencephalon (BS/IB), the cerebral nuclei (CNU), and the cortical subplate (CTXsp). No inputs from the

midbrain (MB) or hindbrain (HB) were detected **(Figure 2F)**. At the next nested level of brain regions, we found

that changes in the CTXpl were contained entirely within the isocortex (ISO) **(Figure 2G)**. No developmental

differences within the BS/IB and CNU were found **(Figure 2H and 2I)**. Within the ISO, only two brain regions,

the ACA and anatomically adjacent secondary motor cortex (MOs) showed a significant difference between the

adolescent and adult groups **(Figure 2J)**. Specifically, the ACA and MOs provided more inputs at adolescence,

suggesting that these local inputs are reduced by adulthood **(Figure 2K)**. Although this system is noted to have

local, non-RbG-dependent labeling [34], we found that the amount of labeling in the presence of RbG **(Figure S2A)**

was much greater than an independent experiment in which RbG was excluded, and that there was no difference

in leak between adolescent and adult groups **(Figure S2B).** This was additionally confirmed by an independent

experiment in which RVdG(EnvA)-eGFP was injected in VIS for uptake by TVA expressed at axon terminals to

avoid local leak **(Figure S2C)**.

As these data revealed a reduction of input source following adolescence, we next set out to determine

the impact of this structural switch on top-down ACA_{VIS} projection integration into its neural networks. We

calculated the relative percentage of total inputs coming from "local" ACA and MOs areas compared to all other

"long-range" distal brain regions for each animal. Our analysis revealed that the relative ratio of local to long-

range distal inputs is significantly higher in adolescents (5.21:1) compared to adults (1.74:1) **(Figure 2L)**.

Together these results support a model that top-down ACA_{VIS} neurons receive greater local inputs during

adolescence than adulthood, which in turn increase a relative contribution of long-range distal connectivity into

adulthood.

Synaptic strength of local excitatory inputs onto top-down ACAvis neurons is greater in adolescence.

Our patch-clamp recordings (**Figure 1**) and rabies mediated mapping of monosynaptic input onto ACA$_{VIS}$

projection neurons (**Figure 2**) collectively point to the model that ACA$_{VIS}$ neurons are more extensively connected

with local frontal excitatory networks during adolescence than adulthood. As rabies mapping cannot fully rule out

non-specific labeling within the local injection area [34] as well as an activity-dependent labelling [35], we next set

out to perform non-rabies based approach to directly examine local functional connectivity. To this end, we

combined optogenetics and slice patch-clamp recording by applying Cre- and Flippase (Flp)- dependent

intersectional viral strategy called INTRSECT [36] to allow optogenetic interrogation of local inputs selectively

without direct activation of ACA$_{VIS}$ projection neurons (**Figure 3A**). This approach can overcome the challenge

of assessing the very low probability of connectivity between pairs of ACA pyramidal neurons (2% in recent study

[37]) using conventional paired patch clamp recordings. We infused retrograde Cre encoding virus in VIS to

retrogradely label the soma of ACA$_{VIS}$ projection neurons in ACA, and also introduced codon-optimized Flp (Flpo)

and FlpON/CreOff-ChR2 vectors [36] into the ACA to express ChR2 selectively in Flpo-expressing ACA neurons

yet actively avoid labeling ACA$_{VIS}$ projection neurons expressing Cre. AAV infusion was performed around p12

or p43 for adolescence and adult recordings respectively. After waiting ~18 days for ChR2 accumulation, we

performed whole-cell patch-clamp recordings. The spread and intensity of YFP signal of analyzed slices show

comparable levels of ChR2 expression between groups (**Figure 3B, C**; see details in methods section). Patch-

clamp recordings were obtained from visually identified mCherry+ACA$_{VIS}$ neurons in acute ACA slices obtained

from adolescent and adult mice. Brief pulses of blue light (0.5 ms) successfully activated local inputs onto ACA$_{VIS}$

neurons and elicited reliable EPSCs in every ACA$_{VIS}$ neuron recorded. The latency from light onset to the onset

of EPSCs in ACA$_{VIS}$ neurons was 5.2 ms on average due to synaptic delay, suggesting monosynaptic

connectivity. In our experiments, we found no cells with the latency below 2 ms which are considered ChR2-

expressing.

First, to assess evoked excitatory monosynaptic transmission from local inputs, we isolated monosynaptic

connections with tetrodotoxin (TTX, 1μM), 4-AP (100 μM) and picrotoxin (100 μM). Light intensity was adjusted

to elicit stable response that didn't change with the additional increase of the stimulus intensity. We found that

the amplitude of evoked EPSC was higher in adolescent animals in comparison to the adults (**Figure 3D**).

Differences in the amplitude of evoked responses could be due to changes in presynaptic release probability,

changes in synaptic strength or synapse number. To investigate more thoroughly, we next used paired

stimulation to evoke EPSCs and quantify possible changes in short-term dynamics. Experiments were performed

in the absence of TTX, 4-AP and picrotoxin to preserve presynaptic action potentials. In response to paired

stimulation, we observed short-term depression of local inputs to ACA$_{VIS}$ neurons (**Figure 3E**). In agreement with

the published data [38], magnitude of short-term depression depended on the interstimulus interval with the higher

depression at shorter intervals. Paired-pulse analysis did not show significant interaction between age and inter-

pulse interval (**Figure 3E**), suggesting that the change in EPSCs could be due to changes in synaptic strength

or synapse number. To further differentiate between these possibilities, we recorded EPSCs in ACSF where

extracellular calcium (Ca^{2+}) was replaced with strontium (Sr^{2+}, 3 mM) in the presence of TTX, 4-AP and picrotoxin

(**Figure 3F**). Sr^{2+} desynchronizes glutamate release and allows the resolution of quantal synaptic events. The

resultant local input-specific qEPSC provides a measure of unitary synaptic strength (amplitude) and a number

of connections (frequency) [39] [40] Optogenetic stimulation of local ACA excitatory axons revealed higher amount

of quantal synaptic events in adolescent animals in comparison to adult (**Figure 3F**). In contrast, we found no

difference in the amplitude of light-evoked quantal events (**Figure 3F**). Collectively, these results suggest that,

during adolescence, the local excitatory drive onto ACA$_{VIS}$ neurons is higher in adolescence than in adulthood

likely due to the increased synaptic number from local excitatory inputs.

Heightened *in vivo* activity of top-down ACA$_{VIS}$ neurons during adolescence

As ACA$_{VIS}$ neurons are more extensively connected with local frontal excitatory networks during

adolescence than adulthood, we next investigated if ACA$_{VIS}$ projection neurons themselves have heightened

activity during adolescence compared to adulthood **(Figure 4A)**. To evaluate ACA$_{VIS}$ neuron activity *in vivo*, we

expressed channelrhodopsin-2 (ChR2) as an optogenetic tag to allow efficient identification of individual ACA$_{VIS}$

neurons amongst a population of neurons by using a blue light (473 nm) as a search stimulus during *in vivo*

electrophysiological recordings **(Figure 4C)**. We targeted expression of ChR2-GFP in ACA$_{VIS}$ projection neurons

by injecting retrograde AAV2-ChR2-GFP into the VIS **(Figure 4AB)**. After >2 weeks of viral incorporation, the

mice were subjected to *In vivo* electrophysiological recordings at adolescent (~p36: p33-39) and adulthood

(~p79: p76-82). Using this technique, detected units were confirmed as ACA$_{VIS}$ neurons if they also responded

to optogenetic stimulation within 3 ms upon blue light stimulation **(Figure 4C-E)**. We found that there was a

significantly greater spontaneous firing rate of ACA$_{VIS}$ neurons *in vivo* during adolescence compared to adulthood

(Figure 4FG). Taken together, our findings suggest that ACA$_{VIS}$ projection neurons are not only highly functionally

integrated into local excitatory circuitry within ACA **(Figure 1-3)**, but also ACA$_{VIS}$ neurons themselves have

heightened activity during adolescence.

Suppression of top-down ACA$_{VIS}$ neuron activity during adolescence produces an adult visual attention

deficit

We next examined whether this transient developmental stage of heightened ACA$_{VIS}$ neuron activity had

significance for adult function. Because ACA$_{VIS}$ projection neurons can enhance the ability to detect visual

features in VIS—a hallmark feature of visual attention [16], we aimed to determine if the activity of top-down ACA$_{VIS}$

projection neurons during this adolescent period is required to establish adult visual attentional behavior. To this

end, we perturbed ACA$_{VIS}$ neuron activity during adolescence (p25-34) or late-adolescent (p45-54) periods by

chemogenetically suppressing ACA$_{VIS}$ neuron activity during these developmental windows and examined its

long-lasting consequence on adult visual attentional behavior. Visual attention capacity was tested by the

5CSRTT[28]. A use of this unimodal visual attention task is particularly relevant for our study which focuses on the

maturation of visual top-down ACA$_{VIS}$ projection neurons, given that a previous chemogenetic silencing study

demonstrated that 5CSRTT requires proper activity of ACA neurons in mice [22]. In contrast, a cross-modal

attention task, in which animals select between conflicting visual and auditory stimuli, does not require ACA or

VIS activity [41]. 5CSRTT assays sustain visual attention by requiring mice to maintain divided attention across 5

areas for the presentation of a brief light stimulus that must be nose-poked to release a reward [42] **(Figure S4)**.

Prior studies have shown that attention deficits can be measured quantitatively as the combined error rate of an

increase in omitted responses to light stimulus and a decrease in accurate responses [25]. Our study was

conducted in automated, standardized Bussey-Saksida touch-screen operant chambers that are highly analogous to Cambridge Neuropsychological Test Automated Battery for humans and non-human primates [43-46], which enables systematic cognitive assessment with easily standardized outputs that can facilitate data dissemination[47].

We introduced inhibitory designer receptor exclusively activated by a designer drug (iDREADD)[48] into visual top-down ACA_{VIS} projection neurons **(Figure 5A)** using an intersectional Cre-dependent viral approach that introduced Cre-dependent DREADD vector into frontal cortex and retrograde Cre encoding virus in VIS (**Figure S3**), and suppressed top-down neuron activity through Clozapine-N-oxide (CNO) delivered intraperitoneally twice daily during the adolescent period of heightened local connectivity from p25 through p34 **(Figure 5C)**. While our intersectional approach does not limit the expression of DREADD to frontal projection to primary sensory cortices due to collaterals to association sensory cortices [49], and therefore cannot rule out the contribution of these collaterals to behavior effects, a previous study demonstrated that stimulation of ACA_{VIS} projection terminals within primary sensory cortex (VIS) is sufficient to impact visual gain [16]. To test the temporal specificity of this developmental perturbation, we also delivered CNO during the late-adolescent period (p45-p54) in a separate group of mice with top-down visual projection neuron iDREADD expression **(Figure 5C)**. As another control, we also examined the effect of suppressing an activity of an adolescent frontal population projecting to primary auditory cortex (AUD), which are located at secondary and primary motor cortical area (MOs/p) adjacent to ACAd [49,50] (MOs_{AUD} projection neurons) **(Figure 5BC)**. To confirm that neurons virally transfected at p12 were responsive to CNO during the adolescent period, we performed slice patch clamp recordings. Whole-cell current clamp recordings were obtained from mCherry expressing ACA_{VIS} neurons in ACA slices. A steady depolarizing current of 1.5-times rheobase (the minimal current to evoke action potential) was applied to initiate firing of ACA_{VIS} neurons following by application of CNO **(Figure S5A)**. CNO application (10 µM) decreased the membrane potential **(Figure S5B)** and inhibited spontaneous firing of projection neurons **(Figure S5C)**.

During 5CSRTT testing, we observed a selective increase in error rate in mice with top-down ACA_{VIS} iDREADD that received CNO during adolescence compared to the group that received CNO during the late-adolescent period, or the MOs_{AUD} iDREADD group that received CNO during the same adolescent period **(Figure**

5D). No differences in correct response latency **(Figure 5E)**, trial number **(Figure 5F)**, premature responses

(Figure 5G), or perseverative responses **(Figure 5H)** were observed, suggesting a selective visual attention

deficit. An increase error rate did not reflect a decrease in motivation because the CNO group as they displayed

no differences in latency to collect the reward during the task **(Figure 5I).** Additionally, in an independent

progressive ratio assay of motivation, no differences in break point to attempt release of the liquid reward **(Figure**

5J). Further, no differences in acquisition of the 5CSRTT or baseline training performance prior to testing were

observed **(Figure S6)**. While it is challenging to truly dissociate deficits in attention and arousal, which are

intertwined concepts, if overall arousal was compromising 5CSRTT error rate, we would additionally expect to

see a decrease in the trial number attempted. Additionally, overall decreased arousal would lead to decreased

breakpoint in our separate testing of motivation with progressive ratio. As arousal state should affect multiple

measures of cognitive performance, and we observe deficits specific to error rate, this rather supports an

attentional deficit over general arousal deficit. Together, these data point to adolescence as an important

sensitive window for top-down ACA$_{VIS}$ projection neurons to drive activity-dependent maturation of visual

attentional behavior in adulthood.

Suppression of top-down ACA$_{VIS}$ projection neuron activity during adolescence leads to excessive loss

of local excitatory drive onto ACA$_{VIS}$ neurons in adulthood

We next set out to gain mechanistic insights into the later adult visual attentional deficits caused by

transient adolescent suppression of top-down ACA$_{VIS}$ neuron activity. To this end, we performed optogenetic

interrogation of local inputs onto ACA$_{VIS}$ neuron by patch-clamp recordings to examine the impact of adolescent

chemogenetic suppression of ACA$_{VIS}$ neuron activity to local excitatory connectivity which showed greater level

during adolescence compared to adulthood in contrast to long-range excitatory inputs which did not differ

between adolescence and adulthood. To virally express iDREADD-mCherry or mCherry in maturing ACA$_{VIS}$

projection neurons, we infused retrograde Cre encoding virus in VIS and Cre-dependent iDREADD-mCherry or

mCherry encoding virus in ACA around p12. We then virally introduced Flpo and FlpON/CreOff-ChR2 vectors

into the ACA to express ChR2 selectively in Flpo-expressing ACA neurons yet actively avoid labeling ACA$_{VIS}$

neurons expressing Cre at p52-p80. 'Adolescence iDREADD' and 'adolescence mCherry' groups experienced

CNO injections during adolescence (p25-34) where 'adolescence mCherry' group was used as a control both for

the effect of CNO and surgery/injections. 'Late-adolescent iDREADD' group experienced CNO from p45 to p54

to test the temporal specificity of ACA suppression (**Figure 6A**). In all experimental groups, patch-clamp

recordings were performed from ACA$_{VIS}$ neurons in acute ACA slices obtained from adult mice (p85.7+/-2.3 in

average (p72-107)). The spread and intensity of YFP signal of analyzed slices show comparable levels of ChR2

expression between groups (**Figure 6B, C:** see details in methods section).

To assess consequence of developmental suppression of ACA$_{VIS}$ neuron activity on local excitatory drive

onto ACA$_{VIS}$ neurons in adulthood, we recorded quantal EPSCs in the presence of strontium (Sr^{2+}, 3 mM), TTX,

4-AP and picrotoxin in a bath (**Figure 6D**). Optogenetic stimulation of local inputs onto ACA$_{VIS}$ projection neurons

revealed decreased frequency of quantal synaptic events in adolescent iDREADD animals in comparison to

adult iDREADD or adolescent mCherry groups following the light stimulation (**Figure 6D**). No differences were

found in the amplitude of quantal events (**Figure 6D**). We also did not observe difference in paired-pulse ratio

(**Figure S7**). These results strongly suggest that adult visual attention deficit caused by adolescent suppression

of top-down ACA$_{VIS}$ neuron activity parallels the excessive reduction of local excitatory inputs onto ACA$_{VIS}$

neurons.

Next, to examine a structural correlate of our findings, we virally introduced a novel bright monomeric

fluorescent protein (mHyperGFP1) that allows visualization of dendritic spines without immunohistochemical

enhancement (Campbell et al unpublished) together with iDREADD selectively in ACA$_{VIS}$ neurons (**Figure S8A**)

and suppressed activity in iDREADD expressing neurons by CNO administration during adolescence from p25

to p34 (**Figure S8B**).   Injections volumes for DREADD and mHyperGFP1 were optimized so that some

mHyperGF1P labelled neurons were co-infected with inhibitory DREADD (mCherry+), and thus allowed for within

animal comparison with DREADD negative (mHyperGFP+ mCherry-) ACA$_{VIS}$ neurons (**Figure S8C and S8D**)

while also ruling out any potential DREADD independent CNO effect [51]. Adolescent chemogenetic suppression

produced an excessive decrease in adult total spine density mCherry+ suppressed ACA$_{VIS}$ neurons compared

to mCherry-control neurons selectively in proximal dendrites (**Figure S8E**). Of note, a previous study

demonstrated that proximal dendrites of layer V/VI neocortical pyramidal neurons preferentially receive local

inputs from surrounding layer II/III neurons [52], while different long-range inputs target select dendritic locations

across dendrites [52,53]. Observed loss of spines in proximal dendrites is consistent with the optogenetics-assisted functional data showing an excessive loss of local excitatory drive onto ACA$_{VIS}$ projection neurons **(Figure 6)**. While future investigations are warranted to examine whether the impact of adolescent suppression of ACA$_{VIS}$ neuron activity is limited to local excitatory inputs or also extends to long-range inputs, our study collectively suggests that adolescent top-down ACA$_{VIS}$ projection neuron activity is required to maintain their local excitatory inputs and support adult visual attentional behavior.

Discussion

In this study, we examined the maturation of frontal top-down corticocortical projection neurons, and demonstrated an adolescence as a time of elevated local excitatory input drive and heightened activity of top-down projection neurons prior to a transition into an adult state with increased long-range contribution through local circuit pruning. Transient perturbation of top-down projection neuron activity at this adolescent window produced long-lasting attentional behavior deficits in adulthood and is accompanied by excessive loss of local inputs. Collectively, our data suggests that heightened adolescent activity is essential for a frontal top-down projection to maintain local inputs and establish attention in adulthood **(Figure S9)**.

Adolescence as a window of heightened frontal top-down neuron activity and associated local inputs

Our circuit-specific patch recording, rabies input mapping, and optogenetic interrogations collectively identified an adolescent period of extensive local excitatory drive onto ACA$_{VIS}$ projection neurons. Circuit-specific patch recordings revealed greater excitatory but not inhibitory drive during adolescence **(Figure 1)**. Rabies-mediated circuit mapping further revealed the input source to be limited to local brain regions **(Figure 2).** As rabies mapping cannot fully rule out non-specific labeling [34] as well as an activity-dependent labelling [35] in the immediate injection area, we further performed non-rabies-based optogenetic interrogation which directly and independently confirmed a heightened local excitatory drive on ACA$_{VIS}$ projection neurons during adolescence **(Figure 3)**. Importantly*, in vivo* electrophysiological recordings revealed that the activity of ACA$_{VIS}$ neurons themselves are also heightened during adolescence neurons in conjunction with the highlighted input drive **(Figure 4)**. A postnatal synaptic pruning is stereotypical across several cortical regions [54,55], but how this developmental event impacts input-output relationship was poorly characterized. This study relates such observations to circuit rewiring for a corticocortical projection and provides new knowledge that this process is driven by local events. These findings also provide a novel conceptual basis for considering similar phenomenon observed in human frontal cortex, in which synaptic density heightens during childhood [56,57].

While the developmental changes revealed by this study were restricted to local excitatory input, we found a relative impact of local input pruning on long-range circuitry. Our rabies input mapping identified that approximately one in every three inputs onto top-down projections arise from a distal brain region in adults, a

significantly higher inputs ratio than the one in every six inputs received during adolescence. Notably, the

absolute number of these long-range connections did not change **(Figure 3E-I)**. Instead, the post-adolescent

loss of local excitatory input tipped the scale toward greater relative long-range, rather than local, engagement

(Figure 3K). Forming long-range synapses is more challenging and energetically intensive than establishing

local connections in which processes can more easily extend and retract [58-60]. The identified reduction in local

drive may serve as a more efficient and effective manner of increasing the contribution of long-range connectivity.

Further, as local inputs tend to preferentially target proximal dendrite [52], removal of irrelevant local circuits may

aid somatic transmission from some of long-range inputs targeting distal dendrites. We propose a model that

this post-adolescent shift in balance between local and long-range inputs is a key developmental milestone for

top-down projection to effectively mediate attentional behavior. Future studies are warranted to examine to what

extent these findings in ACA_{VIS} projection neurons are applicable to other prefrontal neurons located in different

layers and/or projecting to different brain regions.

Linking adolescent top-down neuron activity to adult attentional circuity and behavior

What is the purpose of heightening top-down neuron activity and associated greater local inputs if they are

eventually diminished? At the circuit level, top-down neurons that experienced adolescent chemogenetic

suppression showed over-reduction of local excitatory inputs **(Figure 6)** and total spine density on proximal

dendrites in adulthood **(Figure S8)**. We therefore speculate that adolescent heightened activity of top-down

projection is required to allow robust activity-dependent post-adolescent synaptic refinement from a pool of

enough number of local inputs to maintain adequate local connectivity into adulthood. Based on previous reports

examining the role of activity in circuit formation, we speculate that highly-active local connectivity is stabilized,

while insufficiently active contacts are eliminated or fail to be maintained [61 62 63]. Our findings also suggest that

physiological developmental reduction local connectivity onto ACA_{VIS} neurons following adolescence into

adulthood is likely mediated by distinct mechanisms and requires future investigations. While the current study

focuses on input integration to ACA_{VIS} projection neurons in frontal cortex, it is also important to examine in future

studies if adolescent ACA_{VIS} projection neuron activity can additionally impact downstream ACA_{VIS} projection

outputs onto VIS neurons.

At the behavioral level, our study demonstrates that disruption of frontal top-down projection neuron activity

during adolescence affects adult attentional behavior **(Figure 5)**. We speculate that an excessive local pruning,

as produced by adolescent chemogenetic suppression may fail to effectively drive adult ACA_{VIS} neurons when

attentional adjustment is necessary. Heightened activity of ACA_{VIS} neurons in conjunction with an elevated

excitatory local input drive at the adolescent period may be therefore essential for the maturation of top-down

neurons and association connections within the frontal cortex to play a role in attentional processing and sensory

cue-guided actions [25,64,65] as well as to effectively control distal sensory cortical neurons[14,15]. It should be noted

that our study did not directly manipulate adolescent local excitatory inputs due to the lack of currently available

techniques to selectively chronically manipulate selective input neurons. Development of new techniques will

allow in future studies to directly examine the contribution of heightened local excitatory inputs onto ACA_{VIS} circuit

maturation and associated attentional behavior. Future studies using additional cognitive tasks is also necessary

to examine the generalizability and selectivity of adolescent ACA_{VIS} activity manipulations to cognitive behaviors.

Implications to neurodevelopmental and psychiatric disorders

Our study provides an experimental support to a hypothesized framework that disrupted local vs long-range

balance in frontal cortex may underlie cognitive deficits in psychiatric disorders [66-68], and stresses a need for

future studies to examine the balance between local and long-range connectivity across development in the

context of psychiatric disorders. Our transient adolescent manipulation produced deficits later in life, also points

to a possible developmental mechanism for deficits in corticocortical functional connectivity, and attentional

function. Disruption of circuit activity during the essential adolescent period could decrease frontal dendritic spine

density **(Figure S8)**, as is observed in post-mortem studies of schizophrenia and other neuropsychiatric

disorders [69,70]. Our study also proposes that failure to achieve this developmental milestone of activity-dependent

local input refinement may produce attention deficits. Heterogeneous high-risk gene variations or environmental

insults at this critical developmental stage could converge on failures of proper circuit rewiring and thus produce

common functional deficits—as is observed across attention deficits in autism, schizophrenia, and ADHD [3,19-21].

Clinical approaches that protect and elicit sufficient excitatory drive and top-down projection activity during this

adolescent sensitive period could ameliorate visual attention deficits in neurodevelopmental disorders. Further,

interventions promoting activity-dependent spine formation may have therapeutic value for adult deficits.

Interrogating these possibilities in mouse models carrying these circuit deficits would provide valuable preclinical

insight.

Author Contributions

E.N., Y.G., H.K., and H.M. designed research; E.N., Y.G., H.K., M.S., A.L., K.N., G.T., S.I., K.C., S.L, M.P.D, J.B. performed experiments; E.N., Y.G., H.K., A.L., G.T., S.I., K.C., and H.M. analyzed data; P.R.H., R.C., H.M. supervised the research, E.N., Y.G. and H.M. wrote the manuscript with contributions from all other co-authors. The authors declare no conflicts of interest.

Acknowledgements

This work was supported by National Institute of Health F30 MH111143 to E.N., R21 MH106919, R21 NS105119, R01 MH119523 to H.M., T32 DA007135 to M.P.D., Seaver Foundation to E.N. We thank Milo Smith for assisting statistical analysis, Annalee Tacuri for assisting histological analysis, Dr. Mark Baxter for his expertise on touch-screen behavior system, Dr. Yasmin Hurd and Dr. Michael Michaelides for their expertise on DREADD. The authors declare no competing financial interests or conflict of interest.

References

Harris, A. Z. & Gordon, J. A. Long-range neural synchrony in behavior. *Annual review of neuroscience* **38**, 171-194, doi:10.1146/annurev-neuro-071714-034111 (2015).

Purves, D. & Lichtman, J. W. *Principles of neural development.* (Sinauer Associates, 1985).

Roiser, J. P. *et al.* Dysconnectivity in the frontoparietal attention network in schizophrenia. *Frontiers in psychiatry* **4**, 176, doi:10.3389/fpsyt.2013.00176 (2013).

Kikuchi, M. *et al.* Reduced long-range functional connectivity in young children with autism spectrum disorder. *Soc Cogn Affect Neurosci* **10**, 248-254, doi:10.1093/scan/nsu049 (2015).

Tomasi, D. & Volkow, N. D. Abnormal functional connectivity in children with attention-deficit/hyperactivity disorder. *Biological psychiatry* **71**, 443-450, doi:10.1016/j.biopsych.2011.11.003 (2012).

Zalesky, A. *et al.* Delayed Development of Brain Connectivity in Adolescents With Schizophrenia and Their Unaffected Siblings. *JAMA psychiatry* **72**, 900-908, doi:10.1001/jamapsychiatry.2015.0226 (2015).

Barttfeld, P. *et al.* A big-world network in ASD: dynamical connectivity analysis reflects a deficit in long-range connections and an excess of short-range connections. *Neuropsychologia* **49**, 254-263, doi:10.1016/j.neuropsychologia.2010.11.024 (2011).

Liston, C., Malter Cohen, M., Teslovich, T., Levenson, D. & Casey, B. J. Atypical prefrontal connectivity in attention-deficit/hyperactivity disorder: pathway to disease or pathological end point? *Biological psychiatry* **69**, 1168-1177, doi:10.1016/j.biopsych.2011.03.022 (2011).

Paus, T., Keshavan, M. & Giedd, J. N. Why do many psychiatric disorders emerge during adolescence? *Nature reviews. Neuroscience* **9**, 947-957, doi:10.1038/nrn2513 (2008).

Lee, F. S. *et al.* Mental health. Adolescent mental health--opportunity and obligation. *Science* **346**, 547-549, doi:10.1126/science.1260497 (2014).

Casey, B. J., Glatt, C. E. & Lee, F. S. Treating the Developing versus Developed Brain: Translating Preclinical Mouse and Human Studies. *Neuron* **86**, 1358-1368, doi:10.1016/j.neuron.2015.05.020 (2015).

Uhlhaas, P. J. & Singer, W. The development of neural synchrony and large-scale cortical networks during adolescence: relevance for the pathophysiology of schizophrenia and neurodevelopmental hypothesis. *Schizophrenia bulletin* **37**, 514-523, doi:10.1093/schbul/sbr034 (2011).

Sahin, M. & Sur, M. Genes, circuits, and precision therapies for autism and related neurodevelopmental disorders. *Science* **350**, doi:10.1126/science.aab3897 (2015).

Moore, T. & Zirnsak, M. Neural Mechanisms of Selective Visual Attention. *Annu Rev Psychol* **68**, 47-72, doi:10.1146/annurev-psych-122414-033400 (2017).

Knudsen, E. I. Fundamental components of attention. *Annual review of neuroscience* **30**, 57-78, doi:10.1146/annurev.neuro.30.051606.094256 (2007).

Zhang, S. *et al.* Long-range and local circuits for top-down modulation of visual cortex processing. *Science* **345**, 660-665, doi:10.1126/science.1254126 (2014).

17   Leinweber, M., Ward, D. R., Sobczak, J. M., Attinger, A. & Keller, G. B. A Sensorimotor Circuit in Mouse
Cortex for Visual Flow Predictions. *Neuron* **95**, 1420-1432 e1425, doi:10.1016/j.neuron.2017.08.036
(2017).

18   Itokazu, T. *et al.* Streamlined sensory motor communication through cortical reciprocal connectivity in a
visually guided eye movement task. *Nature communications* **9**, 338, doi:10.1038/s41467-017-02501-4
(2018).

19   Mazaheri, A. *et al.* Functional disconnection of frontal cortex and visual cortex in attention-
deficit/hyperactivity disorder. *Biological psychiatry* **67**, 617-623, doi:10.1016/j.biopsych.2009.11.022
(2010).

20   Keehn B, B. L., Palmer E, Lincoln AJ, Müller RA. Functional brain organization for visual search in ASD.
Austism Research **5**, 314-330, doi:10.1002/aur.1245 (2008).

21   Minshew, N. J. & Keller, T. A. The nature of brain dysfunction in autism: functional brain imaging studies.
Current opinion in neurology **23**, 124-130, doi:10.1097/WCO.0b013e32833782d4 (2010).

22   Koike, H. *et al.* Chemogenetic Inactivation of Dorsal Anterior Cingulate Cortex Neurons Disrupts
Attentional Behavior in Mouse. *Neuropsychopharmacology : official publication of the American College*
of Neuropsychopharmacology **41**, 1014-1023, doi:10.1038/npp.2015.229 (2016).

23   Tan, Z. *et al.* Dynamic ErbB4 Activity in Hippocampal-Prefrontal Synchrony and Top-Down Attention in
Rodents. *Neuron* **98**, 380-393 e384, doi:10.1016/j.neuron.2018.03.018 (2018).

24   White, M. G. *et al.* Anterior Cingulate Cortex Input to the Claustrum Is Required for Top-Down Action
Control. *Cell reports* **22**, 84-95, doi:10.1016/j.celrep.2017.12.023 (2018).

25   Kim, H., Ahrlund-Richter, S., Wang, X., Deisseroth, K. & Carlen, M. Prefrontal Parvalbumin Neurons in
Control of Attention. *Cell* **164**, 208-218, doi:10.1016/j.cell.2015.11.038 (2016).

26   Counotte, D. S. *et al.* Lasting synaptic changes underlie attention deficits caused by nicotine exposure
during adolescence. *Nature neuroscience* **14**, 417-419, doi:10.1038/nn.2770 (2011).

27   Guillem, K. *et al.* Nicotinic acetylcholine receptor beta2 subunits in the medial prefrontal cortex control
attention. *Science* **333**, 888-891, doi:10.1126/science.1207079 (2011).

28   Robbins, T. W. The 5-choice serial reaction time task: behavioural pharmacology and functional
neurochemistry. *Psychopharmacology* **163**, 362-380, doi:10.1007/s00213-002-1154-7 (2002).

29   Brust, V., Schindler, P. M. & Lewejohann, L. Lifetime development of behavioural phenotype in the house
mouse (Mus musculus). *Front Zool* **12 Suppl 1**, S17, doi:10.1186/1742-9994-12-S1-S17 (2015).

30   Spear, L. P. The adolescent brain and age-related behavioral manifestations. *Neurosci Biobehav Rev*
24, 417-463, doi:10.1016/s0149-7634(00)00014-2 (2000).

31   Laviola, G., Macri, S., Morley-Fletcher, S. & Adriani, W. Risk-taking behavior in adolescent mice:
psychobiological determinants and early epigenetic influence. *Neurosci Biobehav Rev* **27**, 19-31,
doi:10.1016/s0149-7634(03)00006-x (2003).

32   Wickersham, I. R. *et al.* Monosynaptic restriction of transsynaptic tracing from single, genetically targeted
neurons. *Neuron* **53**, 639-647, doi:10.1016/j.neuron.2007.01.033 (2007).

33 Huh, C. Y. L., Peach, J. P., Bennett, C., Vega, R. M. & Hestrin, S. Feature-Specific Organization of
Feedback Pathways in Mouse Visual Cortex. *Current biology : CB* **28**, 114-120 e115,
doi:10.1016/j.cub.2017.11.056 (2018).

34 Miyamichi, K. *et al.* Dissecting local circuits: parvalbumin interneurons underlie broad feedback control
of olfactory bulb output. *Neuron* **80**, 1232-1245, doi:10.1016/j.neuron.2013.08.027 (2013).

35 Beier, K. T. *et al.* Rabies screen reveals GPe control of cocaine-triggered plasticity. *Nature* **549**, 345-350,
doi:10.1038/nature23888 (2017).

36 Fenno, L. E. *et al.* Targeting cells with single vectors using multiple-feature Boolean logic. *Nat Methods*
11, 763-772, doi:10.1038/nmeth.2996 (2014).

37 Blom, S. M., Pfister, J. P., Santello, M., Senn, W. & Nevian, T. Nerve injury-induced neuropathic pain
causes disinhibition of the anterior cingulate cortex. *The Journal of neuroscience : the official journal of*
the Society for Neuroscience **34**, 5754-5764, doi:10.1523/JNEUROSCI.3667-13.2014 (2014).

38 Regehr, W. G. Short-term presynaptic plasticity. *Cold Spring Harb Perspect Biol* **4**, a005702,
doi:10.1101/cshperspect.a005702 (2012).

39 Rajkovich, K. E. *et al.* Experience-Dependent and Differential Regulation of Local and Long-Range
Excitatory Neocortical Circuits by Postsynaptic Mef2c. *Neuron* **93**, 48-56,
doi:10.1016/j.neuron.2016.11.022 (2017).

40 McGarry, L. M. & Carter, A. G. Prefrontal Cortex Drives Distinct Projection Neurons in the Basolateral
Amygdala. *Cell reports* **21**, 1426-1433, doi:10.1016/j.celrep.2017.10.046 (2017).

41 Wimmer, R. D. *et al.* Thalamic control of sensory selection in divided attention. *Nature* **526**, 705-709,
doi:10.1038/nature15398 (2015).

42 Bari, A., Dalley, J. W. & Robbins, T. W. The application of the 5-choice serial reaction time task for the
assessment of visual attentional processes and impulse control in rats. *Nature protocols* **3**, 759-767,
doi:10.1038/nprot.2008.41 (2008).

43 Romberg, C., Mattson, M. P., Mughal, M. R., Bussey, T. J. & Saksida, L. M. Impaired attention in the
3xTgAD mouse model of Alzheimer's disease: rescue by donepezil (Aricept). *The Journal of*
neuroscience : the official journal of the Society for Neuroscience **31**, 3500-3507,
doi:10.1523/JNEUROSCI.5242-10.2011 (2011).

44 Mar, A. C. *et al.* The touchscreen operant platform for assessing executive function in rats and mice.
Nature protocols **8**, 1985-2005, doi:10.1038/nprot.2013.123 (2013).

45 Bussey, T. J. *et al.* New translational assays for preclinical modelling of cognition in schizophrenia: the
touchscreen testing method for mice and rats. *Neuropharmacology* **62**, 1191-1203,
doi:10.1016/j.neuropharm.2011.04.011 (2012).

46 Silverman, J. L., Gastrell, P. T., Karras, M. N., Solomon, M. & Crawley, J. N. Cognitive Abilities on
Transitive Inference Using a Novel Touchscreen Technology for Mice. *Cerebral cortex*,
doi:10.1093/cercor/bht293 (2013).

47 Beraldo, F. H. *et al.* MouseBytes, an open-access high-throughput pipeline and database for rodent
touchscreen-based cognitive assessment. *Elife* **8**, doi:10.7554/eLife.49630 (2019).

48 Zhu, H. & Roth, B. L. DREADD: A Chemogenetic GPCR Signaling Platform. *International Journal of*
Neuropsychopharmacology **18**, pyu007-pyu007, doi:10.1093/ijnp/pyu007 (2015).

49   Zhang, S. *et al.* Organization of long-range inputs and outputs of frontal cortex for top-down control.
Nature neuroscience **19**, 1733-1742, doi:10.1038/nn.4417 (2016).

50   Nelson, A. *et al.* A circuit for motor cortical modulation of auditory cortical activity. *The Journal of*
neuroscience : the official journal of the Society for Neuroscience **33**, 14342-14353,
doi:10.1523/JNEUROSCI.2275-13.2013 (2013).

51   Gomez, J. L. *et al.* Chemogenetics revealed: DREADD occupancy and activation via converted
clozapine. *Science* **357**, 503-507, doi:10.1126/science.aan2475 (2017).

52   Petreanu, L., Mao, T., Sternson, S. M. & Svoboda, K. The subcellular organization of neocortical
excitatory connections. *Nature* **457**, 1142-1145, doi:10.1038/nature07709 (2009).

53   Little, J. P. & Carter, A. G. Subcellular synaptic connectivity of layer 2 pyramidal neurons in the medial
prefrontal cortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **32**,
12808-12819, doi:10.1523/JNEUROSCI.1616-12.2012 (2012).

54   Markham, J. A., Mullins, S. E. & Koenig, J. I. Periadolescent maturation of the prefrontal cortex is sex-
specific and is disrupted by prenatal stress. *The Journal of comparative neurology* **521**, 1828-1843,
doi:10.1002/cne.23262 (2013).

55   Koss, W. A., Belden, C. E., Hristov, A. D. & Juraska, J. M. Dendritic remodeling in the adolescent medial
prefrontal cortex and the basolateral amygdala of male and female rats. *Synapse* **68**, 61-72,
doi:10.1002/syn.21716 (2014).

56   Petanjek, Z. *et al.* Extraordinary neoteny of synaptic spines in the human prefrontal cortex. *Proceedings*
of the National Academy of Sciences of the United States of America **108**, 13281-13286,
doi:10.1073/pnas.1105108108 (2011).

57   Huttenlocher, P. R. Synaptic density in human frontal cortex - developmental changes and effects of
aging. *Brain Res* **163**, 195-205 (1979).

58   Latora, V. & Marchiori, M. Efficient behavior of small-world networks. *Phys Rev Lett* **87**, 198701,
doi:10.1103/PhysRevLett.87.198701 (2001).

59   Kaiser, M. & Hilgetag, C. C. Nonoptimal component placement, but short processing paths, due to long-
distance projections in neural systems. *PLoS computational biology* **2**, e95,
doi:10.1371/journal.pcbi.0020095 (2006).

60   Nicosia, V., Vertes, P. E., Schafer, W. R., Latora, V. & Bullmore, E. T. Phase transition in the economically
modeled growth of a cellular nervous system. *Proceedings of the National Academy of Sciences of the*
United States of America **110**, 7880-7885, doi:10.1073/pnas.1300753110 (2013).

61   Katz, L. C. & Shatz, C. J. Synaptic activity and the construction of cortical circuits. *Science* **274**, 1133-
1138 (1996).

62   Hua, J. Y. & Smith, S. J. Neural activity and the dynamics of central nervous system development. *Nature*
neuroscience **7**, 327-332, doi:10.1038/nn1218 (2004).

63   Selemon, L. D. A role for synaptic plasticity in the adolescent development of executive function.
Translational psychiatry **3**, e238, doi:10.1038/tp.2013.7 (2013).

64   Barthas, F. & Kwan, A. C. Secondary Motor Cortex: Where 'Sensory' Meets 'Motor' in the Rodent Frontal
Cortex. *Trends in neurosciences* **40**, 181-193, doi:10.1016/j.tins.2016.11.006 (2017).

Nakajima, M., Schmitt, L. I. & Halassa, M. M. Prefrontal Cortex Regulates Sensory Filtering through a
Basal Ganglia-to-Thalamus Pathway. *Neuron* **103**, 445-458 e410, doi:10.1016/j.neuron.2019.05.026
(2019).

66   Stephan, K. E., Friston, K. J. & Frith, C. D. Dysconnection in schizophrenia: from abnormal synaptic
plasticity to failures of self-monitoring. *Schizophrenia bulletin* **35**, 509-527, doi:10.1093/schbul/sbn176
(2009).

67   Northoff, G. & Sibille, E. Why are cortical GABA neurons relevant to internal focus in depression? A cross-
level model linking cellular, biochemical and neural network findings. *Molecular psychiatry* **19**, 966-977,
doi:10.1038/mp.2014.68 (2014).

68   Khan, S. *et al.* Local and long-range functional connectivity is reduced in concert in autism spectrum
disorders. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 3107-
3112, doi:10.1073/pnas.1214533110 (2013).

69   Roberts, R. C., Barksdale, K. A., Roche, J. K. & Lahti, A. C. Decreased synaptic and mitochondrial density
in the postmortem anterior cingulate cortex in schizophrenia. *Schizophrenia research* **168**, 543-553,
doi:10.1016/j.schres.2015.07.016 (2015).

70   Osimo, E. F., Beck, K., Reis Marques, T. & Howes, O. D. Synaptic loss in schizophrenia: a meta-analysis
and systematic review of synaptic protein and mRNA measures. *Molecular psychiatry* **24**, 549-561,
doi:10.1038/s41380-018-0041-5 (2019).

71   Kuhlman, S. J. *et al.* A disinhibitory microcircuit initiates critical-period plasticity in the visual cortex.
Nature **501**, 543-546, doi:10.1038/nature12485 (2013).

72   Bukhari, N. *et al.* Unmasking Proteolytic Activity for Adult Visual Cortex Plasticity by the Removal of
Lynx1. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **35**, 12693-12702,
doi:10.1523/JNEUROSCI.4315-14.2015 (2015).

73   Heath, C. J., Bussey, T. J. & Saksida, L. M. Motivational assessment of mice using the touchscreen
operant testing system: effects of dopaminergic drugs. *Psychopharmacology* **232**, 4043-4057,
doi:10.1007/s00213-015-4009-8 (2015).

74   Dumitriu, D., Rodriguez, A. & Morrison, J. H. High-throughput, detailed, cell-specific neuroanatomy of
dendritic spines using microinjection and confocal microscopy. *Nature protocols* **6**, 1391-1411,
doi:10.1038/nprot.2011.389 (2011).

75   Christoffel, D. J. *et al.* IkappaB kinase regulates social defeat stress-induced synaptic and behavioral
plasticity. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **31**, 314-321,
doi:10.1523/JNEUROSCI.4763-10.2011 (2011).

76   Rodriguez, A., Ehlenberger, D. B., Dickstein, D. L., Hof, P. R. & Wearne, S. L. Automated three-
dimensional detection and shape classification of dendritic spines from fluorescence microscopy images.
PloS one **3**, e1997, doi:10.1371/journal.pone.0001997 (2008).

77   Rodriguez, A., Ehlenberger, D. B., Hof, P. R. & Wearne, S. L. Rayburst sampling, an algorithm for
automated three-dimensional shape analysis from laser scanning microscopy images. *Nature protocols*
1, 2152-2161, doi:10.1038/nprot.2006.313 (2006).

Figures and Figure Legends

Figure 1: Frontal top-down ACA$_{VIS}$ projection neurons receive greater excitatory drive during adolescence. A. Developmental changes in excitatory and inhibitory synaptic drive onto top-down ACA$_{VIS}$ projection neurons were assessed cross-sectionally. Slice patch-clamp recordings were performed from ACA$_{VIS}$ projection neurons, fluorescently labeled by retrobeads injected in the VIS, on ACA/MOs brain slices at p20.3±-0.3 in average (p18-22) for Juvenile group, at p32.7±0.4 in average (p29-37) for Adolescent group, and at p78.1±4.3 in average (p65-110) for Adult group. Miniature excitatory postsynaptic currents (mEPSC) and

miniature inhibitory postsynaptic currents (mIPSC) were recorded in the presence of TTX at −70 mV and 0 mV

respectively. **B-C.** Representative traces of mEPSC (**B**) and mIPSC (**C**) measured from ACA$_{VIS}$ neurons in three

developmental age groups (for mEPSC: Juvenile: n=23 cells from 5 mice, Adolescent: n=27 cells from 11 mice,

Adult: n=15 cells from 6 mice; for mIPSC: Juvenile: n=25 cells from 5 mice, Adolescent: n=26 cells from 11 mice,

Adult: n=17 cells from 6 mice). Cumulative distribution (top) and averaged data (bottom) of mEPSC frequency

(**D**) and mEPSC amplitude (**E**) in three age groups. Average mEPSC frequency were significantly higher in

adolescent mice compared to adult (one-way ANOVA p=0.021, post hoc Tukey test $p_{Juv-Adoles}$=0.43, *$p_{Adoles-}$

$_{Adult}$=0.016). While average mEPSC amplitude shows overall age effect, there was no significant difference

between adolescence and adulthood (one-way ANOVA p=0.014; post hoc Tukey test $p_{Juv-Adoles}$=0.24, $p_{Adoles-}$

$_{Adult}$=0.23). Cumulative distribution (top) and averaged data (bottom) of mIPSC frequency (**F**) and mEPSC

amplitude (**G**) in three age groups show trending yet no significant difference in average mIPSC frequency (one-

way ANOVA p=0.054), and no difference in average mIPSC amplitude (p=0.18 one-way ANOVA).

See related Figure S1.

Figure 2: Frontal top-down ACA_VIS projection neurons receive greater local presynaptic input during adolescence compared to adulthood. A. Schematic overview of intersectional viral injection approach

enabling labeling of presynaptic input neurons specifically onto top-down ACA$_{VIS}$ projection neurons. Cre-dependent AAVs expressing TVA-mcherry and Rb-G were injected into the ACA and retrograde CAV-Cre into VIS. After 2.5 weeks, pseudotyped rabies virus was injected into the ACA to allow for uptake of rabies virus in top-down neurons only to express GFP over one week. **B.** Injection timeline for two age groups. AAV injections were initiated at p12 to capture inputs during adolescence (p38) and at p60 to capture inputs during adulthood (p84). **C.** Top-down starter cells and local inputs were visualized in ACA. Left: Magenta "helper" TVA-mcherry infected cells permit rabies entry. White "starter" AAV TVA-mcherry + Rbs-GFP co-infected cells allow rabies to retrograde back one synapse onto green monosynaptic input cells. Right: High magnification of boxed area. Scale Bars =100 μm. **D.** The anterior posterior distribution of starter cells is consistent across adolescent (n=7 mice) and adult groups (n=6 mice). (Linear mixed model with age and animal as independent effects; p=0.98). **E.** The absolute number of starter cells per animal are comparable across the two age groups (*t*-test, p=0.6521, n=7 [adolescent], 6 mice [adult]). **F-J.** Absolute total number of input neurons in adolescence and adulthood normalized to starter cell count per animal. See **Table 1** for abbreviations. **F.** Compared to adult mice (n=6 mice), adolescent mice (n=7 mice) show a larger number of input cells at the highest hierarchical level of neuroanatomical organization selectively in the cortical plate (CTXpl) (two-way ANOVA post hoc analysis, ***p<0.0001). No inputs were detected from the midbrain or hindbrain for either group. **G.** Within the CTXpl, adolescent mice showed a larger number of input cells selectivity in the isocortex (ISO) (two-way ANOVA post hoc analysis, *** p=0.0036), and not in the olfactory area (p>0.99) or hippocampus (p>0.99). **H.** Adolescent and adult mice show the same number of inputs coming from sub-regions of the diencephalon (two-way ANOVA post hoc analysis, p=0.81, p>0.99, p>0.99, p>0.99, p>0.99) and **I.** cerebral nuclei (two-way ANOVA post hoc analysis, p=0.92, 0.62). **J.** In the isocortex, more input neurons are found selectively in the secondary motor area (MOs) (two-way ANOVA post hoc analysis, ***p<0.0001) and anterior cingulate cortex (ACA) (two-way ANOVA post hoc analysis, ***p<0.0001). **K.** Representative image of ACA and MOs in adolescent vs adult group (Scale bar 500μm). **L.** Local:Long-Range input ratio is higher in adolescence compared to adults (*t*-test, *p=0.0253, n=7,6 mice). See related Figure S2.

Figure 3: Greater adolescent local excitatory synaptic inputs onto frontal ACA_{VIS} projection neurons. **A.**
Schematic of intersectional viral approach to test local excitatory synaptic input specifically onto ACA_{VIS}
projection neurons in adolescence and adulthood. (*left*) Injections of retrograde AAV2-mCherry-Cre into the VIS
labeled ACA_{VIS} projection neurons in ACA with red fluorescence, while injections of AAV-Flpo and AAV-
CreOff/FlpOn-ChR2-YFP into ACA enable ChR2-YFP expression in all but ACA_{VIS} projection neurons. (*middle*)
AAV injections were done at p12.8±0.5 in average (p11-15) for Adolescent group and at p43.0±1.9 in average
(p38-58) for Adult group. Slice patch clamp recordings were performed at p33.5±0.7 in average (p31-37) for
Adolescent group and at p71.8±4.2 in average (p64-99) for Adult group. (*right*) Local inputs were stimulated *in
situ* by activating ChR2 using wide-field LED. **B.** (*left*) Labeling of local inputs (green) and ACA_{VIS} projection
neurons (red) in ACA. (*right*) High magnification of boxed area showing mCherry+ ACA_{VIS} neuron does not
express ChR2-YFP and is surrounded by ChR2+YFP terminals. Scale bars: 200 µm on the left and 20 µm on

the right. **C.** Quantification of the spread and intensity of the YFP signal shows no significant differences between Adolescent and Adult groups (area: p=0.75, mean intensity: p=0.07, *t* test, n=11 slices from 10 mice for adolescent and n=12 slices from 7 mice for adult groups). **D.** Light-evoked monosynaptic EPSC traces (*top*) and mean amplitude (*bottom*) for two developmental age groups (*p=0.047 *t* test, n=10 cells from 5 mice for adolescent and n=11 cells from 6 mice for adult groups). Blue dashes indicate the time of light stimulation. **E.** Short-term dynamics of local synaptic inputs onto ACA$_{VIS}$ neurons. Representative EPSC evoked by pair of light stimuli (*top*) and average paired pulse ratio (*bottom*) (EPSC2/EPSC1 Adolescent vs. EPSC2/EPSC1 Adult, n=12 cells from 4 mice for adolescent and n=11 cells from 4 mice for adult, two-way ANOVA, time x age: p=0.596). **F.** Examples of light-evoked quantal EPSC (qEPSC) in ACA$_{VIS}$ neurons in the presence of strontium (*top*) to produce delayed asynchronous quantal release of neurotransmitters. Arrows indicate detected qEPSCs. (*bottom*) Averaged qEPSC amplitude and frequency for Adolescent and Adult groups (frequency: *p= 0.045, amplitude: p=0.12, *t* test, n=13 cells from 4 mice for adolescent and n=12 cells from 5 mice for adult).

Figure 4: Greater firing activity of frontal top-down ACAvis projection neurons *in vivo* during adolescence. A. Schematic of viral approach to express Channelrhodopsin-2 (ChR2) in frontal top-down ACAvis projection neurons for *in vivo* electrophysiological recordings of optically tagged ACAvis neurons in adolescent and adult mice. (*left*) Injections of retrograde AAV2-ChR2-GFP into the VIS enable ChR2-GFP expression in cells projecting to VIS including ACAvis projection neurons. (*right*) AAV injections were done at p20 for Adolescent group and at p63.3±0.7 in average (p62-64) for Adult group. Recordings were performed at p35.8±1.2 in average (p33-39) for Adolescent group and at p79.0±1.7 in average (p76-82) for Adult group. **B.** Representative images of viral GFP+ACAvis neurons in ACA following injection of retroAAV2-ChR2-GFP into the VIS. Scale: 200 μm. **C.** Schematic of *in vivo* electrophysiology setup with optical tagging. ChR2 has been

used as an optogenetic tag to allow identification of ACAvis projection neurons (green) *in vivo* based on whether they can be activated by optical stimulation. **D.** Representative overlapping spike waveforms of optically-tagged ACAvis projection neuron recorded with (blue trace) and without (gray trace) optic stimulation. **E.** Raster plot and peristimulus time histogram showing time locked optogenetic activation of a representative ACAvis neuron within 2 ms. Laser (wavelength 473 nm, fiber diameter 105 μm) was delivered using an optic fiber coupled to the multichannel extracellular recording electrode. The fiber tip was positioned immediately above the ACA recording site. ACAvis neurons were identified with a light pulse (1 ms, 5Hz) stimulus. **F.** Cumulative distribution of spontaneous firing frequency of ACAvis projection neurons in adolescent and adult animals (blue line, n= 231 cells from 5 adolescent mice, black line, n= 78 cells from 3 adult mice, ****P<0.0001, Kolmogorov-Smirnov test). **G.** Averaged firing activity of ACAvis projection neurons in adolescent and adult animals (blue, n=231 cells from 5 adolescent mice, black line, n= 78 cells from 3 adult mice, ****P<0.0001).

Figure 5: Suppression of frontal top-down ACA_VIS projection neuron activity during adolescence selectively produces a visual attention deficit in adulthood. A-B. *(left)* Schematic of intersectional viral approach to introduce hM4D(Gi)-inhibitory DREADD (iDREADD) selectively into **(A)** frontal visual top-down ACA_VIS projection neurons, or (**B**) frontal auditory top-down MOs_AUD projection neurons by introducing Cre-dependent iDREADD vectors into the frontal cortex and retrograde Cre encoding virus in VIS. Viruses were

injected at p12 to allow for expression of iDREADD by p25. (*right*) Representative images of frontal cortex showing labeling of (**A**) mCherry+ACA$_{VIS}$ projection neurons and their terminals in layer1 of VIS or (**B**) mCherry+MOs$_{AUD}$ projection neurons in frontal cortex. **C.** Overview of experimental timeline. Mice were injected with 10 mg/kg CNO in sterile saline twice a day from p25 to p34 or from p45 to p54 and started training on the five-choice serial reaction time task (5CSRTT) at adulthood (p60). See **Figure S2** for a detailed overview of the 5CSRTT. Mice were subsequently tested for motivation with the progressive ratio task. **D.** CNO-mediated suppression of frontal top-down projection neuron increases adult attentional error rate selectively in adolescent visual top-down suppression condition (two-way RM ANOVA, *p= 0.0429, Newman-Keuls multiple comparisons test: *p< 0.05: adolescent vs late-adolescent visual top-down suppression (n=10, 8 mice), *p< 0.05: adolescent visual top-down suppression vs adolescent auditory top-down suppression (n=10,8 mice)). **E-I.** Adolescent CNO administration does not alter other measures on the 5CSRTT in adult mice (n=10,8,8 mice): **E.** Correct response latency (one-way ANOVA, p= 0.1331), **F.** Trial number (one-way ANOVA, p= 0.2820) **G.** Premature responses (one-way ANOVA, p= 0.2495) **H.** Perseverative responses (one-way ANOVA, p=0.7321) **I.** Reward collection latency (one-way ANOVA, p=0.5171) **J.** No differences in breakpoint were observed in the progressive ratio task (one-way ANOVA, p= 0.8454, n=7,8,8 mice). See related Figures S3-S6.

Figure 6: Chemogenetic suppression of frontal top-down ACA_VIS projection neuron activity during adolescence results in their excessive loss of local excitatory inputs. A. Experimental approach to test the impact of ACA_VIS neuron activity during adolescent or late-adolescent period on maintaining local excitatory input drive onto ACA_VIS neurons into adulthood through optogenetic interrogation of local excitatory input onto ACA_VIS

neurons. (*left*) RetroAAV2-cre was injected into the VIS, while AAV-DIO- hM4D(Gi)-(iDREADD)-mCherry or AAV-DIO-mCherry was injected into ACA at p12 to selectively express iDREADD-mCherry or mCherry in ACA$_{VIS}$ neurons by p25. (*middle*) Mice were injected with 10 mg/kg CNO in normal saline twice a day from p25 to p34 in adolescent iDREADD and adolescent mCherry groups, or from p45 to p54 in late-adolescence iDREADD group. Additional injections of AAV-Flpo and AAV-CreOff/FlpOn-ChR2-YFP in ACA at p60.4±2.2 in average (p52-p80) enable ChR2-GFP expression in all but ACA$_{VIS}$ neurons in ACA. (*right*) Slice patch clamp recordings were performed from ACA$_{VIS}$ neuron at p85.7±2.3 in average (p72-107) while optogenetically activating ChR2-expressing local excitatory inputs. **B.** (*left*) Representative image showing ChR2-YFP expressing local inputs (green) and mCherry expressing ACA$_{VIS}$ neurons (red) in ACA. (*right*) High magnification of boxed area showing mCherry+ACA$_{VIS}$ neuron does not express ChR2-YFP and is surrounded by ChR2+ terminals. Scale bars: 200 μm on the left and 20 μm on the right. **C.** The quantification of the spread and intensity of the YFP signal showed no significant differences among groups (area: p=0.88, one-way ANOVA; mean intensity: p=0.16, one-way ANOVA, n=13 slices from 9 mice for adolescent iDREADD. n=15 slices from 10 mice for adolescent mCherry, and n=7 slices from 5 mice for late-adolescent iDREADD). **D.** Examples of light-evoked EPSC in ACA$_{VIS}$ neurons in the presence of strontium to produced delayed asynchronous release of neurotransmitters for each groups. Arrows indicate detected qEPSCs. **E-F.** Averaged qEPSC **(E)** amplitude and **(F)** frequency for adolescent iDREADD, adolescent mCherry and late-adolescent iDREADD groups (frequency:**p<0.01 one-way ANOVA, followed by post hoc Tukey test **p$_{Adol.iDREADD-Adol.mCherry}$=0.009, **p$_{Adol.iDREADD-Lateadol.iDREADD}$=0.002 ; mean amplitude: p=0.27 one-way ANOVA, n=15 cells from 4 mice for adolescent iDREADD. n=15 cells from 4 mice for adolescent mCherry, and n=16 cells from 4 mice for late-adolescent iDREADD). See related Figure S7.

Taxonomic Table of Abbreviations

CTXpl Cortical Plate
          ISO   Isocortex
                       FRP   Frontal Pole
                       AI   Insular Cortex
                       ORB   Orbitofrontal Cortex
                       ILA   Infralimbic Cortex
                       PL   Prelimbic Cortex
          Local MOs   Secondary Motor Cortex
          Local ACA   Anterior Cingulate Cortex
                       MOp   Primary Motor Cortex
                       SS   Somatosensory Cortex
                       GU   Gustatory Area
                       RSP   Retrosplenal Cortex
                       VISC   Visceral Area
                       AUD   Auditory Cortex
                       TEa   Temporal Cortex
                       ECT   Entorhinnal Cortex
                       PERI   Perirhinal Cortex
                       PTLp   Posterior Parietal Cortex
                       VIS   Visual Cortex
          OLF   Olfactory Areas
          HIP   Hippocampal Formation
BS/IB Brain Stem
          Interbrain (diencephalon)

                       HYP   Hypothalamus
                       TH   Thalamus
                       DOR   Dorsal Thalamus
                       EPI   Epithalamus
                       RT   Thalamic Reticular
                                   Nucleus
                       GENv   Ventral Thalamus
          MB   Midbrain
          HB   Hindbrain
CNU   Cerebral Nuclei
          STR   Striatum
          PAL   Pallidum
CTXsp Cortical Subplate

Table 1. Hierarchical table of abbreviations for brain regions. Indented regions are subregions of the

outdented region above.

Methods

Animals: Male C57Bl/6 mice (Charles River) were group-housed under a standard 12 hr light/dark cycle in a temperature and humidity-controlled vivarium and were provided with food and water *ad libitum*. Adolescent mice were shipped at 10 days after birth (P10) and weaned at P21. Adult mice were shipped at P56. During behavior testing, mice were allowed access to water for 2 hr each day and maintained approximately 85-90% of respective baseline free feeding weight. All animal protocols were approved by IACUC at the Icahn School of Medicine at Mount Sinai.

Drugs: Clozipine-N-Oxide dihydrochloride (CAS2250025-93-3, Tocris Biosciences) was injected intraperitoneally (i.p.) two times a day at 10mg/kg in sterile saline for 10 days from p25 to p34, or from p45 to p54 following the previously validated protocol in mouse cortex[22,71]

Stereotactic Surgery: Mice were anesthetized with 2% isoflurane and head-fixed on in a mouse stereotactic apparatus (Narishige International USA Inc.) equipped with a heating pad. Mice were isolated post-surgery until fully-awake and immediately returned to their home cage. VIS injection sites relative to lambda are: AP +0.0 mm, ML +3.0 mm, DV -0.4 mm; AP +0.1 mm, ML +2.85 mm, DV -0.4 mm; AP +0.1 mm, ML +3.15 mm, DV -0.4 mm. ACA injections sites relative to bregma are: AP +1.7 mm, ML +0.2 mm, DV -0.7 mm; AP +1.1 mm, ML +0.2 mm, DV -0.7 mm; AP +0.4 mm, ML +0.2 mm, DV -0.7 mm. MOs injection sites relative to lambda are: AP +0.5 mm, ML +0.4 mm, DV -1.0 mm; AP +0.0 mm, ML +0.4 mm, DV -1.0 mm; AP -0.5 mm, ML +0.4 mm, DV -1.0 mm. For patch-clamp recordings, 500 nl of fluorescent retrobeads (Lumafluor, Inc) were injected into the VIS 4 to 6 days prior to patch recordings to allow visualization of top-down projection neurons. 350 nl of AAV5-hSyn-eGFP-WPRE (University of Pennsylvania Viral Core) was infused into VIS for the basic characterization of ACA$_{VIS}$ projection neurons. For rabies-mediated monosynaptic input mapping, 500 nl of a 1:1 volume mixture of AAV8-CA-FLEX-RG and AAV8-EF1a-FLEX-TVAmCherry (University of North Carolina Viral Core) were injected unilaterally in the ACA and 500 nl of CAV2-Cre (Institut de Genetique Moleculaire de Montpellier) were injected in VIS. For somatic introduction of rabies, 2.5 weeks later, 500 nl of pseudotyped EnvA RVdG-GFP (Salk Institute Viral Core) was injected into the ACA for local axonal uptake of rabies or VIS for distal axonal uptake of rabies.

Mice were euthanized 7 days later. For behavioral studies, 500 nL of AAV8-DIO-hM4D(Gi)-mCherry and 500 nl CAV2-Cre were injected in the ACA/MOs and VIS, respectively. For the dendritic spine study, a total of 500 nl of AAV8-DIO-hM4D(Gi)-mCherry (University of North Carolina Viral Core) and AAV1-EF1a-DIO-hyperGFP2-WPRE (Custom Prep, Boston Children's Hospital) at a ratio of 2.5:1 was injected in the ACA and 500 nl CAV2-Cre (Institut de Génétique Moleculaire de Montpellier) in VIS sites at p12. For optogenetics-based patch clamp recordings, rAAV2-Ef1a-mCherry-IRES-Cre (Addgene) was injected into VIS while AAVDJD/hSyn-CreOff/FlpOn-ChR2-YFP and AAV5/EF1a-FLPO-WPRE (both from University of North Carolina Viral Core) were injected in ACA at p11-15 for adolescent group and at p38-58 for late-adolescent group. For the developmental experiments with iDREADD, rAAV2-CAG-Cre-WPRE (Boston Children's Hospital) was injected into VIS while AAV8-hSyn-DIO-hM4D(Gi)(iDREADD)-mCherry or AAV8-hSyn-DIO-mCherry (Addgene) was injected into ACA at p12-13. Additional injections of AAV5/EF1a-Flpo and AAVDJD-hSyn-CreOff/FlpOn-ChR2- YFP in ACA were at p52-p80. For the *in vivo* electrophysiology experiment where ACAvis neurons were optogenetically tagged and recorded, a virus expressing channelrhodopsin-2 (rAAV2-Syn-ChR2(H134R)-GFP: Addgene) was injected into the VIS.

Tissue preparation and Histology: Animals underwent transcardial perfusion with phosphate-buffered saline (PBS) followed by 4% paraformaldehyde (PFA) in PBS. Brains from rabies input mapping and DREADD validation were dissected, postfixed in 4% PFA for 3 h, and cryoprotected in 30% sucrose in PBS for 24-48 h before embedding in Optimum Cutting Temperature (OCT, Tissue Tek) and sectioning into 35 μm-thick coronal slices using a cryostat (CM3050, Leica). For rabies input mapping, free-floating sections were washed in PBS before undergoing immunohistochemistry by 1hour incubation in blocking solution (1% bovine serum albumin in 0.1% Triton X-100 in PBS) followed by overnight incubation with rabbit anti-eGFP antibody (1:1000 in blocking solution; Life Technologies) at room temperature. Sections were washed 3 times with blocking solution followed by incubation with Alexa 488-conjugated goat anti-rabbit antibody (Life Technologies) for 2.5 h, washed 3 times in blocking solution, and mounted onto slides with DAPI Fluoromount-G (Southern Biotech). For spine analysis, brains were post-fixed in 4% PFA overnight and coronally sectioned at 75 μm on a Vibratome (Leica VT1000 S). All slices were then mounted on Superfrost Plus slides using Fluoromount-G mounting medium (Southern

Biotech) and coverslipped. DREADD injections were validated using a LSM780 confocal microscope (Zeiss) in

reference to the Paxinos and Franklin mouse atlas. For the viral spread validation of mice used for behavioral

experiments, mice completed behavior testing underwent transcardial perfusion with PBS followed by 4% PFA

in PBS, with overnight postfixation. Brains were coronally sectioned on a Leica CM3050 S cryostat (Leica

Microsystems, Buffalo Grove, IL) at 60 μm for immunohistochemistry. Slices were collected at nine specific

Bregma areas (2.10, 1.70, 1.18, 0.74, 0.14, -0.34, -0.82, -1.34, -1.82) to analyze the anterior-posterior spread of

virally infected cells. Images were acquired using the EVOS FL Imaging System (Thermo Fisher Scientific).

Slices were imaged using a 4x lens for mCherry and to ensure consistency across imaging sessions, power was

set at 100%. Using Excel, all groups were assigned five levels representing increasing levels of signal intensity.

The first level (L1) represented the minimum number of mice showing signal in a given area, namely n=1. The

second level (L2) represented the 25th percentile of mice with overlapping signal in a given area, the third level

(L3) represented the 50th percentile, and the fourth level (L4) represented the 75th percentile. The fifth level (L5)

represented the maximum number of mice showing overlapping expression in a given area, namely the total

number of mice in the group. Using the GNU Image Manipulation Program (GIMP), areas with mCherry signal

were delineated on templates taken from the Paxinos and Franklin mouse atlas. Signal intensity was determined

by the quartile and the number of mice with overlapping viral expression in a given area.

Slice Electrophysiology: Animals were decapitated under isofluorane anesthesia. Brains were quickly

removed and transferred into ice-cold artificial cerebrospinal fluid (ACSF) of the following composition (in mM):

210.3 sucrose, 11 glucose, 2.5 KCl, 1 NaH_2PO_4, 26.2 $NaHCO_3$, 0.5 $CaCl_2$, and 4 $MgCl_2$. Acute coronal slices of

ACA (300 μm) contained both hemispheres. Slices were allowed to recover for 40 min at room temperature in

the same solution, but with reduced sucrose (105.2 mM) and addition of NaCl (59.5 mM). Following recovery,

slices were maintained at room temperature in standard ACSF composed of the following (in mM): 119 NaCl,

2.5 KCl, 1 NaH_2PO_4, 26.2 $NaHCO_3$, 11 glucose, 2 $CaCl_2$, and 2 $MgCl_2$.

Slices were visualized under upright differential interference contrast microscope equipped with high power

LED-coupled light source used for the identification of fluorescently-labeled cells and optogenetic stimulation

(Prizmatix). Patch clamp recordings were made from ACA_{VIS} projection neurons fluorescently labeled with

retrobeads or mCherry. For voltage clamp recordings, borosilicate glass electrodes (5–7 MΩ) were filled with the

internal solution containing (in mM): 120 Cs-methanesulfonate, 10 HEPES, 0.5 EGTA, 8 NaCl, 4 Mg-ATP, 1 QX-

314, 10 Na-phosphocreatine, and 0.4 Na-GTP (pH 7.25, 295 mOsm). Current clamp recordings were obtained

with the internal solution containing (in mM): 127.5 K-methanesulfonate, 10 HEPES, 5 KCl, 5 Na-

phosphocreatine, 2 MgCl$_2$, 2 Mg-ATP, 0.6 EGTA, and 0.3 Na-GTP (pH 7.25, 295 mOsm). Signals were low-pass

filtered at 3 kHz and sampled at 20 kHz. Neurons were included in the analysis if input resistance, series

resistance and membrane potential did not change more than 10% during the course of recordings. Miniature

inhibitory and excitatory postsynaptic currents were isolated by holding the neuron at the reversal potential for

excitatory or inhibitory currents. mIPSC and mEPSC were recorded in the presence of TTX (1 μM; Abcam) in

the bath solution to block action potentials. After acquisition of a stable baseline (10 min), miniature postsynaptic

events were recorded 3-5 min for each potential.

For iDREADD validation spontaneous firing rate was measured in a current clamp mode. Recordings were

performed in normal ACSF with a depolarizing current of 1.6±0.04 times rheobase injected to the cell. In order

to establish steady-state firing activity of recorded neurons, we slowly increased the depolarizing current over

the course of 2-3 minutes to elicit low level of sustained spontaneous firing. The recordings were initiated after

the adjustment of the depolarizing current. Firing rate was quantified as the average instantaneous firing

frequency measured over 3 min before and following CNO (10 μM) perfusion.

To selectively activate local inputs to frontal ACA$_{VIS}$ projection neurons, we used slices being introduced

with ChR2-GFP construct in ACA using the intersectional viral approach [36]. ChR2 was activated with a 0.25-0.5

ms light pulse delivered through the objective and synchronized with electrophysiological recordings. At the

beginning of each experiment an input/output curve was established for each neuron to determine the intensity

of the stimulation that evoked EPSC reliably at the top of the curve to ensure activation of all ChR2-expressing

terminals within the field of view. The power of light stimuli was quantified using an optical power meter

(Thorlabs). The range of intensities at the tissue level was 0 to 3.9 mW/mm^2. For all optogenetic experiments we

set the intensity of the stimulus at the plateau level where additional increase in the light intensity didn't affect

the amplitude of the light-evoked responses. Short-term dynamics were tested with twin pulses separated by

100, 250 and 500 ms. Stimuli were given every 20 sec and four responses were averaged at each interpulse

interval. Paired-pulse recordings were conducted in standard ACSF. To limit the impact of disynaptic excitation,

paired pulse ratio was determined as a ratio of the slope of $EPSC_2$ to the slope of $EPSC_1$. To isolate

monosynaptic connections, eEPSCs were recorded in the presence of TTX (1 µM), 4-AP (100 µM) and picrotoxin

(100 µM). Quantal EPSC were measures in ACSF where $CaCl_2$ was replaced with 3 mM $SrCl_2$ in addition to

TTX, 4-AP and picrotoxin. Quantal EPSCs were detected and analyzed using MiniAnalysis (Synaptosoft Inc.).

The detection threshold for events was set at 2x root mean square noise measured before stimulation and

quantified by Noise Analysis tool built in MiniAnalysis. Evoked quantal EPSCs were detected in 30-400 msec

from the stimulus onset, acquired for not less than 3 sweeps and averaged.

To assess consistency of ChR2-GFP expression across preparations, the spread and intensity of YFP

expression was quantified for slices used in patch-clamp experiments. Following the recording, slices were fixed

in phosphate buffer saline containing 4% PFA and stored at 4ºC. Slices were subsequently washed three times

for 5 min each in PBS and mounted on microscope slides (Brain Research Laboratories) using Fluoromount-G

mounting medium (Southern Biotech). Images were obtained with an EVO FL Imaging System equipped with 4x

objective to visualize entire ACA with the adjacent cortical areas simultaneously. Images were analyzed using

ImageJ (NIH) software. Images were thresholded at mean intensity plus two standard deviation of the

background fluorescent signal. The background fluorescent signal was calculated as an intensity of fluorescence

in motor/somatosensory areas lacking YFP expression. The spread and intensity of YFP expression were

quantified as the area and mean intensity of YFP signal respectively. Recordings were excluded from the

analysis if obtained from slices in which the YFP signal deviated more than two standard deviation from the

average intensity or area. This exclusion criteria led to exclude 1 slice from adolescent iDREADD group. This

allowed for comparisons of recordings obtained from slices with similar levels of ChR2 expression in ACA.

Monosynaptic Input Mapping: Samples were imaged using a LSM780 confocal microscope (Zeiss). All slices

around the ACA injection regions were stained for starter cell analysis. One out of every 8 serial sections was

stained for input analysis. ImageJ (National Institute of Health) were used to process and analyze all images.

Starter cells were defined as both mCherry+ (>50% of soma pixels above 2 SD the mean image mCherry

fluorescence intensity) and eGFP+ (>50% of soma pixels above 3SD the mean image GFP fluorescence

intensity). Input cells were defined as eGFP+ (>50% of soma pixels above 3 SD the mean image eGFP

fluorescence intensity). To determine the brain region location for all input and starter cells, all brain slices were

registered to corresponding Allen Brain Institute coronal maps using anatomical landmarks visualized by DAPI

counterstaining and tissue autofluorescence. In a small minority of cases, assignment of input neurons to specific

brain nuclei may be approximate if eGFP+ cell bodies were located on borders between regions, or when

anatomical markers were lacking between directly adjacent regions. However, quantitative analyses of input

tracing results were performed on anatomical classifications that were at least one hierarchical level broader (as

specified by the Allen Brain Atlas) than the discrete brain regions input cells were assigned to. Notably in many

of these cases, directly adjacent nuclei fall together into the same hierarchical group. Both eGFP+ input neurons

and eGFP+ mCherry+ starter cells were manually counted using the "Cell Counter" plug-in in ImageJ. For the

input analysis, we normalized the total number of input neurons in each brain to the total number of starter cells

because the efficiency of rabies uptake by helper cells differed slightly across animals. The investigator

performing these analyses remained agnostic of the group identities of animals through the duration of this

analysis.

***In vivo* Extracellular Electrophysiology**: Preparatory surgery leading to, and recording itself was performed

initially under Nembutal/chlorprothixene anesthesia and then maintained with isoflurane [72]. Optogenetically

tagged ACAvis neurons were identified using a laser (wavelength 473 nm, 1ms duration, 5 Hz) search stimulus

emitted and delivered through the optic fiber (diameter 105 µm) coupled with a linear sixteen-channel silicon

probes with 177 µm^2 recording sites (NeuroNexus Technologies, Ann Arbor, MI) spaced 50 µm apart and

oriented immediately above the ACA cortical surface. The power at the fiber-optic tip was approximately 10 mW.

The signal detected from the probe was amplified and thresholded, and unit-sorted using an OmniPlex Neural

Recording Data Acquisition System (Plexon). A template online sorting method was used to capture spikes as

units. A PCA cluster-based spike shape template-based sorting method was used. Visually identifiable subsets

of waveforms form clusters on 2D PCA space. Spike shape templates were selected by manually drawing

selection contours around PCA clusters. The selected spikes were then automatically averaged and used to

create templates, which were then used to sort incoming baseline spikes of similar shape and placement in PCA

space into units. After sorting for optogenetically responsive units (ACAvis neurons expressing ChR2), the

optogenetic stimulus was switched off. Then baseline spike activity was recorded from the sorted units for 3 mins

followed by 1min of recordings upon turning on a laser for optogenetic tagging. To ensure single-unit isolation,

the waveforms of recorded units were further examined using Offline Sorter (Plexon). To analyze the

electrophysiology data, firing rate was computed by custom made MATLAB (MathWorks) script during 3 min

laser off session mentioned above. Only sorted units which showed time-locked spiking response within 2 ms

upon laser pulse stimulation as well as whose spike width time (trough-to-peak time) is more than 400 μs were

included in the analysis. For each animal, single units were recorded in each of the AP -0.3 mm, 0.0 mm, +0.3

mm, +0.6 mm, ML ±0.25 mm, DV -1.0 mm vertical penetrations across ACA.

Behavior: Five-Choice Serial Reaction Time Task (5CSRTT): Behavior was conducted in black plastic

trapezoid Bussey–Saksida touch-screen chambers (walls 20-cm high × 18-cm wide (at screen-magazine) × 24-

cm wide (at screen) × 6-cm wide (at magazine) (Lafayette Instrument). Stimuli were displayed on a touch-

sensitive screen (12.1-inch, screen resolution 600 × 800) divided into five response windows by a black plastic

mask (4.0 × 4.0-cm, positioned centrally with windows spaced 1.0 cm apart, 1.5 cm above the floor) fitted in front

of the touchscreen. Schedules were designed and data were collected and analyzed using ABET II Touch

software (Lafayette Instrument). The inputs and outputs of the multiple chambers were controlled by

WhiskerServer software (Lafayette Instrument). Before training on 5CSRTT, mice were initially trained to touch

the screen. Mice were acclimated to the chambers for 3 days in 30-min sessions. During this habituation phase,

the food magazine was illuminated and diluted sweetened condensed milk (Eagle Brand) was dispensed every

40 s. Mice were then trained to touch the response windows. If the mouse touched the stimulus, the milk reward

was delivered in conjunction with a tone and magazine light. Touches to non-stimuli had no consequence. After

reaching criterion on this phase (20 touches in 30 min for 2 consecutive days), mice moved onto 5CSRTT training

phase. Mice were tested 5 days a week, 100 trials a day (or up to 30 min). Each trial began with the illumination

of the magazine light. When the mouse made a nose poke in the food magazine, the stimulus was delivered

after an intertrial interval (ITI) period of 5 s. If a mouse touched the screen during this ITI period, the response

was recorded as premature and the mouse was punished with a 5-s time-out (house light on). After the time-out

period, the magazine light illumination and house light switch off signaled onset of the next trial. After the ITI

period, a stimulus appeared randomly in one of the five response windows for a set stimulus duration (this varied

from 32 to 2 s). A limited-hold period followed by the stimulus duration was 5 s, during which the stimulus was

absent but the mouse was still able to respond to the location. Responses during stimulus presence and limited

holding period could be recorded either as correct (touching the stimulus window) or incorrect (touching any

other windows). A correct response was rewarded with a tone, and milk delivery, indicated by the illumination of

the magazine light. A failure to respond to any window over the stimulus and limited-hold period was counted as

an omission. Incorrect responses and omissions were punished with a 5- time-out. In addition, animals could

make perseverative responses that are screen touches after a correct or incorrect response. Animals started at

stimulus duration of 32 seconds. With a goal to baseline mice at a stimulus duration of 2 s, the stimulus duration

was sequentially reduced from 32, 16, 8, 4, to 2 s. Animals had to reach a criterion (>50 trials, >80% accuracy,

<20% omissions) over 2 consecutive days to pass from one stage to the next. After reaching baseline criterion

with the 2 s stimulus duration (5 consecutive days), mice were challenged with an increased attentional demand

by reducing the stimulus duration to 2, 1.5, 1, and 0.8 s (reduced stimulus test). They then underwent 4 days of

testing. Attention and response control were assessed by measuring the following performance: percentage

error (100 x (incorrect responses + omissions))/(correct responses + incorrect responses + omissions),

percentage premature response (100 × premature responses/(omissions + correct responses + incorrect

responses), percentage perseverative response (100 × perseverative responses /(correct responses + incorrect

responses), latency to correct response, and latency to reward collection after correct choices.

    Progressive Ratio Task: Behavior was conducted following completion of the 5CSRTT in touch-screen

chambers based on a previous report [73] with minor modifications. In initial operant training, mice were trained to

nose-touch the visual stimulus (white square, 100% luminance) at the center screen one time in each trial to

receive a diluted sweetened condensed milk (Eagle Brand) reward as dispensed in the 5CSRTT. Touches to

non-stimuli had no consequence. After reaching criterion of at least 30 completed trials in 30 mins, mice moved

onto fixed ratio training. In fixed ratio training, mice were trained to nose-touch the visual stimulus two, three,

then five times to get a reward. Mice were required to reach criterion for one day for fixed ratio of 2:1 and 3:1,

and for three days for fixed ratio 5:1 before proceeding to progressive ratio training and testing. Task parameters

were identical for progressive ratio training, except that completion of each trial, the number of touches required

to receive a reward was incremented on a linear +4 basis (i.e. 1, 5, 9, etc.). If no stimulus response or magazine

entry in the presence of a delivered reward was detected for 5 minutes, the session ended and the animal was

removed from the chamber. Otherwise the session ended after 30 min. Mice had to achieve a stabile "breakpoint"

defined as the number of target location responses in the last successfully completed trial for 2 days before

proceeding to progressive ratio testing. In progressive ratio testing, mice performed the progressive ratio task

using the same parameters as progressive ratio training, but were allowed 60 min to complete the session. Mice

were baselined on Monday with progressive ratio training to ensure consistent performance and tested for four

consecutive days. Breakpoint across the four days was averaged to give each mouse on breakpoint score.

Dendritic Spine Imaging and Characterization: Images were acquired using an Upright LSM 780 Confocal

microscope (Carl Zeiss). A non-biased, whole neuron approach was taken to characterize dendritic spines.

Specifically, two neurons per animal were imaged on both the apical and basal dendrites. To qualify for spine

analysis, whole neurons were selected that met the following requirements: (1) the neuron had to be filled and

display spines at least up to 200 μm away from the soma on both apical and basal dendrites and (2) display no

overlapping of dendritic processes with other neuron processes. For the chemogenetic study, mCherry+ and

mCherry- neurons were defined as having >50% of soma pixels above 3 SD and below 2 SD the mean image

mCherry fluorescence intensity, respectively. Images were taken of sections from two basal dendrites and two

apical dendrites from each neuron and binned into 4 distance groups from the soma: 0-50 μm, 50-100 μm, 100-

150 μm, and 150-200 μm. Dendritic segments were imaged using a 100X lens (numerical aperture 1.4; Carl

Zeiss) and a zoom of 3.0. Images were taken with a resolution of 1024 x 300, pixel dwell time was 1.58 μm/s,

and the line average was set to 8. Pixel size was 0.03um in the x–y plane and 0.01 μm in the z plane. To assure

consistency of imaging across difference confocal sessions, power was consistently set to 3.0%, and gain was

adjusted within the range of 600-800 units to achieve consistent light intensity values within the same set of

neurons.

Images were deconvolved using a 3D resolution enhancement with AutoDeblur software (Media Cybernetics)

and then run through the dynamic range filter in Neuron-Studio [74-77]. To detect dendritic spines from background

noise accurately, puncta were only counted as spines if they moved into and out of the z-plane with the dendritic

branch. Spine density was calculated as the total spine count/dendritic length. All imaging and analysis were all

performed by an author who remained agnostic to the group conditions until the analysis was completed.

Statistical Methods: Linear mixed modeling approaches using the packages LmerTest (v. 2.0.32), lme4 (v.

1.1.12), and lsmeans (v. 2.25) in the R programing language (v. 3.2.2) were user for analysis of starter cell

distribution. Groups were modeled as fixed effects and animals were modeled as random effects (this factor

considered independent and randomly sampled from their respective populations) to determine the impact of

group on location of starter cell.

All other statistical analyses were performed using Prism 6.0 (Graphpad). In electrophysiological recordings

differences in amplitude and frequency of postsynaptic currents as well as the spread and intensity of YFP

expression were tested by a one-way repeated measures analysis of variance (ANOVA) followed by post hoc t

test with Tukey's multiple comparisons test, while the effect of CNO on membrane potential was determined with

two-tailed paired T-tests. Paired pulse ratio analysis was tested by two-way repeated measures ANOVA. For

input mapping and ratio comparisons, input numbers were first normalized to the number of starter cells within

each animal before analyses. Input mapping analyses were completed by a two-way repeated ANOVA with brain

regions as within-subjects factors, followed by post hoc Bonferroni's multiple comparisons test. Input ratios and

locally restricted rabies experiment were both analyzed using an unpaired T-test. 5CSRTT data (error) were

analyzed using a two-way repeated measures analysis of variance (ANOVA) with and stimulus duration (2, 1.5,

1, 0.8 second) as within-subjects factors, followed by Newman-Keuls multiple comparisons test. In the analysis

of other 5CSRTT data and progressive ratio testing, an unpaired t-test was used. Dendritic spine data were

analyzed with ANOVA tests with two-way repeated measures with distance from soma (0-50 μm, 50-100 μm,

100-150 μm, 150-200 μm) as within-subjects factors, followed by post hoc Sidak's multiple comparisons test. All

data are expressed as means ± SEM.