Morphological Cell Profiling of SARS-CoV-2 Infection Identifies Drug Repurposing Candidates for COVID-19

Carmen Mirabelli[1,*], Jesse W. Wotring[2,*], Charles J. Zhang[2,†], Sean M. McCarty[2,†], Reid Fursmidt[3,4,†], Tristan Frum[5,] Namrata S. Kadambi[3], Anya T. Amin[3], Teresa R. O'Meara[1], Carla D. Pretto[1], Jason R. Spence[3,5], Jessie Huang[6,7], Konstantinos D. Alysandratos[6,7], Darrell N. Kotton[6,7], Samuel K. Handelman[3,4], Christiane E. Wobus[1], Kevin J. Weatherwax[4,8,9], George A. Mashour[4,8,10], Matthew J. O'Meara[11‡], Jonathan Z. Sexton[2,3,4,8‡]

[1]Department of Microbiology and Immunology, University of Michigan Medical School, Ann Arbor, MI, 48109, USA
[2]Department of Medicinal Chemistry, College of Pharmacy, University of Michigan, Ann Arbor, MI, 48109, USA
[3]Department of Internal Medicine, Gastroenterology, Michigan Medicine at the University of Michigan, Ann Arbor, MI, 48109, USA
[4]U-M Center for Drug Repurposing, University of Michigan, Ann Arbor, MI, 48109, USA
[5]Department of Cell and Developmental Biology, University of Michigan, Ann Arbor, MI, 48109, USA
[6]Center for Regenerative Medicine of Boston University and Boston Medical Center, Boston, MA, 02118, USA
[7]The Pulmonary Center and Department of Medicine, Boston University School of Medicine, Boston, MA, 02118, USA
[8]Michigan Institute for Clinical and Health Research (MICHR), University of Michigan, Ann Arbor, MI, 48109, USA
[9]College of Pharmacy, University of Michigan, Ann Arbor, MI 48109, USA
[10]Department of Anesthesiology, Michigan Medicine at the University of Michigan, Ann Arbor, MI, 48109, USA
[11]Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI, 48109, USA

[*]These authors contributed equally to this work.
[†]These authors contributed equally to this work.
[‡]Co-Senior Authors

Abbreviations:

MOI: multiplicity of infection
UMAP: uniform manifold approximation and projection
COVID-19: Coronavirus Disease-2019
MOA: mechanism of action
ROI: region of interest
iAEC2: induced pluripotent stem cell (iPSC)-derived alveolar epithelial type 2 cells
HCQ: hydroxychloroquine

Conflicts of interest

The authors declare no conflicts of interest.

ABSTRACT

The global spread of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), and

the associated disease COVID-19, requires therapeutic interventions that can be rapidly identified

and translated to clinical care. Unfortunately, traditional drug discovery methods have a >90%

failure rate and can take 10-15 years from target identification to clinical use. In contrast, drug

repurposing can significantly accelerate translation. We developed a quantitative high-throughput

screen to identify efficacious single agents and combination therapies against SARS-CoV-2.

Quantitative high-content morphological profiling was coupled with an AI-based machine learning

strategy to classify features of cells for infection and stress. From a library of 1,425 FDA-approved

compounds and clinical candidates, we identified 17 dose-responsive compounds with antiviral

efficacy. In particular, we discovered that lactoferrin is an effective inhibitor of SARS-CoV-2

infection with an IC_{50} of 308 nM and that it potentiates the efficacy of both remdesivir and

hydroxychloroquine. Lactoferrin also stimulates an antiviral host cell response and retains

inhibitory activity in iPSC-derived alveolar epithelial cells, a model for the primary site of infection.

Given its safety profile in humans, these data suggest that lactoferrin is a readily translatable

therapeutic option for COVID-19. Additionally, several commonly prescribed drugs were found to

exacerbate viral infection and warrant follow up studies. We conclude that morphological profiling

for drug repurposing is an effective strategy for the selection and optimization of drugs and drug

combinations as viable therapeutic options for COVID-19 pandemic and other emerging infectious

diseases.

MAIN

SARS-CoV-2 is an enveloped, positive-sense, single-stranded RNA betacoronavirus that emerged in Wuhan, China in November 2019 and rapidly developed into a global pandemic. The associated disease, COVID-19, has an array of symptoms, ranging from flu-like illness and gastrointestinal distress[1,2] to acute respiratory distress syndrome, heart arrhythmias, strokes, and death[3,4]. Drug repurposing has played an important role in the search for COVID-19 therapies. Recently, the FDA issued emergency approval of remdesivir, a nucleoside inhibitor prodrug developed for Ebola virus treatment[5], and hydroxychloroquine, an aminoquinoline derivative first developed in the 1940s for the treatment of malaria, for patients with severe COVID-19. However, there are no established prophylactic strategies or direct antiviral treatments available to limit SARS-CoV-2 infections and to prevent/cure the associated disease COVID-19.

Repurposing of FDA-approved drugs is a promising strategy for identifying rapidly deployable treatments for COVID-19. Benefits of repurposing include known safety profiles, robust supply chains, and a short time-frame necessary for development[6]. Additionally, approved drugs serve as chemical probes to understand the biology of viral infection and can help make new associations between COVID-19 and molecular targets/pathways that influence pathogenesis of the disease. A complementary approach to standard *in vitro* antiviral assays is high-content imaging-based morphological cell profiling. Using morphological cell profiling, it is possible to identify pathways and molecular targets underlying infection, thus allowing for targeted screening around a biological process or targeting of host processes that limit viral infection.

Here, we developed a pipeline for quantitative high-throughput image-based screening of SARS-CoV-2 infection. We leveraged machine learning approaches to create an assay metric that accurately and robustly identifies features that predict antiviral efficacy. From this, we identified

several FDA-approved drugs and clinical candidates with unique antiviral activity. We further

demonstrated that one of our most promising hits, lactoferrin, inhibits viral entry and replication,

enhances antiviral host cell response, and potentiates the effects of remdesivir and

hydroxychloroquine. Furthermore, we identified currently prescribed drugs that exacerbate viral

infectivity. As a confirmatory step, efficacy of lead drugs was validated in a highly physiologically

relevant organotypic and biomimetic human model system for bronchial epithelium. Collectively,

we present evidence that morphological profiling can be used to characterize the viral life cycle in

vitro and robustly identify new potential therapeutics against SARS-CoV-2 infection.

Morphological profiling reveals unique features associated with SARS-CoV-2 infection

To determine the optimal cell line and appropriate endpoint for antiviral drug screening, we

assessed SARS-CoV-2 infectivity in previously reported permissive cell lines: Vero E6, Caco-2,

and Huh7[7]. Viral growth kinetics at a multiplicity of infection (MOI) of 0.2 revealed that Vero E6,

Caco-2, and Huh7 cells supported viral infection, with peak viral titers at 48 hours post infection

(hrs p.i.) (Supplementary Figure 1a/b). Although the viral load was higher in Vero E6 cells, Huh7

were selected for our morphological drug screen as a human cell line that expresses both ACE2

and TMPRSS2, which are the primary entry factors for SARS-CoV-2[8]. Infection was detectable in

Huh7 cells at an MOI as low as 0.004 at 48 hrs p.i. (Supplementary Figure 1c), which highlights

the high sensitivity of image-based screening. To identify compounds that inhibit or exacerbate

infection, we selected an MOI of 0.2, leading to a baseline infectivity rate of 20%.

Morphological cell profiling was enabled through multiplexed staining and automated high-content

fluorescence microscopy. Our multiplexed dye set included markers for SARS-CoV-2

nucleocapsid protein (NP), nuclei (Hoechst 33342), neutral lipids (HCS LipidTox Green), and cell

boundaries (HCS CellMask Orange). These fluorescent probes were chosen to capture a wide

variety of cellular features relevant to viral infectivity, including nuclear morphology, nuclear

texture, cytoplasmic and cytoskeletal features, and indicators of cell health.

We observed several prominent features associated with SARS-CoV-2 infection in Huh7 cells

including: the formation of syncytia, small/round cells with a prominent negative viral staining in

the nucleus, increased nucleoli count (Supplementary Figure 1d), and cytoplasmic protrusions

(Figure 1a). These protrusions are consistent with what has been observed in other cell lines

infected with SARS-CoV-2, including Vero-E6 and Caco-2.[9]

In order to quantify the unique morphological features associated with viral infection in Huh7 cells,

we used the image segmentation and analysis software CellProfiler to measure 660 cellular

features on a per-cell basis. Features included measurements of intensity, radial distribution, and

texture for each fluorescent channel.[10] Infected cells were initially distinguished from uninfected

cells by the presence of NP signal, however through morphological profiling we determined that

there were several different manifestations of the infection within the cell population.

The different stages of infection are described by their divergence of intensity and morphological

features in any of the fluorescent channels. To efficiently display the diversity of natural

morphologies of infected cells, cellular features were dimensionally reduced via the non-linear

uniform manifold approximation and projection (UMAP) to embed 2.47 million infected cells into

2-dimensions. Clusters of cells were identified based on isolated clusters or regional high-density

peaks[11] (Figure 1b, Supplementary Figure 2). In the UMAP embedding, we identified 12 regions

of interest (ROI) with high cell density. Inspection of cell images within ROIs revealed distinct

morphologic characteristics that include punctate NP signal (ROI 11), individually infected cells

with high NP intensity at the periphery (ROI 9), cells in multinucleated syncytia (ROI 8) and

infected cells with protrusions wrapping around uninfected cells (ROI 6). Inhibition of viral

progression is the aim for antiviral therapeutics, therefore compounds that impede any of these

phenotypic stages can potentially be useful for treatment of SARS-CoV-2 infection. To leverage

the spatial resolution of imaging-based screening, we used a multivariate scoring system that

incorporated both intensity and morphometric measurements to determine antiviral efficacy rather

than single measurements based on viral staining intensity or rescuing a cytopathic effect.

Phenotypic features describing the different stages of infection in Huh7 cells were used to

generate a machine learning pipeline for antiviral drug screening

Machine learning identifies FDA-approved molecules with antiviral activity against SARS-

CoV-2

To identify compounds with antiviral activity against SARS-CoV-2, we screened a library of 1,425

FDA-approved compounds and rationally included clinical candidates (Supplementary File 1) in

quantitative high-throughput screening (qHTS) at five concentrations (50 nM, 250 nM, 500 nM,

1000 nM and 2000 nM) in Huh7 cells. Compounds were assessed for their antiviral activity using

a CellProfiler-based image analysis pipeline and a random forest classification algorithm to

identify infected cells and quantify their morphological characteristics (Figure 2a). The random

forest classifier leveraged 660 unique cellular features including measurements of intensity,

texture and radial distribution for each fluorescent channel (nuclei, cytoplasm, lipid, virus). From

the qHTS, 132 drugs were selected as active with consistent decreases in viral infectivity in at

least three of the tested concentrations as well as minimal cytotoxicity.

In confirmatory screening, 10-point, two-fold dilution dose-response experiments were performed

in triplicate on the 132 qHTS hits, with validation of dose-responsive efficacy for 17 compounds

below 1 μM potency (Supplementary Table 1 and Figure 2b). These hits include eleven that are

novel *in vitro* observations (bosutinib, domperidone, entecavir, fedratinib, ipratropium bromide,

lacoferrin, lomitapide, metoclopramide, S1RA, thioguanine, and Z-FA-FMK), and six that have

been previously identified to have antiviral activity (amiodarone, verapamil, gilteritinib,

clofazimine[12,13], niclosamide[14], and remdesivir). Amiodarone, gilterinib, lomitapide, thioguanidine

and Z-FA-FMK retained activity in a traditional CPE-based antiviral assay in Vero E6

(Supplementary Table 1). In addition to antiviral drug hits, we also identified several compounds

that appear to exacerbate SARS-CoV-2 infection, including trametinib, binimetinib and

cobimetinib -potent MEK inhibitors used to treat metastatic melanoma- and the Parkinson's

disease drugs carbidopa, methyldopa and levodopa (Supplementary Figure 3).

Lactoferrin blocks SARS-CoV-2 replication at different stages of the viral cycle

One of the most efficacious hits identified from our screen was lactoferrin, a protein found in milk

and other secretory fluids[15]. We determined that lactoferrin has dose-dependent antiviral activity

through a range of MOIs (Figure 3a and b). Previous work on lactoferrin in the context of infection

with SARS-CoV-1 suggests that it blocks viral entry by binding heparan sulfate proteoglycans that

are important for early viral attachment[16]. Lactoferrin blocks SARS-CoV-2 infection via entry

inhibition (Supplementary Figure 4) and also has significant antiviral activity when added 1 or 24

hrs p.i. (Figure 3b), suggesting multimodal efficacy. Lactoferrin has been proposed to enhance

innate interferon responses to limit viral replication within host cells[17]. Upon treatment, we

observed a dose-dependent reduction of viral replication (Figure 3c), which was consistent with

elevated mRNA levels of IFNβ and interferon-stimulated genes (ISG15, MX1, Viperin and IFITM3)

in lactoferrin-treated Huh7 cells (Figure 3d). Interestingly, we detected a robust antiviral effect by

both holo and apolactoferrin (human and bovine), the latter being the component of widely

available dietary supplements. To rule out a mode of action that involved a general iron depletion

mechanism, we tested the protein transferrin and found that it was devoid of any anti-SARS-CoV-

2 activity at the highest concentration of 2.3 μM (Figure 3e).

A clinically effective strategy for antiviral therapies uses a combinatorial (or "drug cocktail")

approach, where compounds with varying mechanisms of action are concomitantly used to target

different stages in the viral life cycle and to minimize the risk of acquired drug resistance from

single-agent selective pressure. This is especially true for RNA viruses, which are highly variable

and can develop drug-resistance[18]. Given the pronounced single-agent efficacy of lactoferrin, we

tested whether combinations with remdesivir or hydroxycholoroquine could improve the overall

antiviral activity. We found that lactoferrin potentiates the efficacy of both remdesivir (Figure 3f

and Supplementary Figure 5a) and hydroxychloroquine (Figure 3f and Supplementary Figure 5b),

which are currently explored treatments for SARS-CoV-2 infection. Therefore, combination

therapy with lactoferrin could be beneficial in the management of the COVID-19 pandemic by

reducing toxicity (e.g., hydroxycholorquine) or consumption (e.g., remdesivir).

Lead compounds demonstrate efficacy in iPSC-derived model of bronchial epithelium

To evaluate the translatability of our identified lead compounds, we used a biomimetic model of

bronchial epithelium, iPSC-derived alveolar epithelial type 2 cells (iAEC2s)[19]. Surfactant protein

C positive (SFTPC+) epithelial cells were previously used to model other lung diseases in place

of primary AEC2s[20]. The advantage of using iPSC-derived AEC2s is in the development of

physiologically relevant heterogeneous airway cell populations that includes alveolar type II cells,

that are involved in COVID-19 pathogenesis[21]. We demonstrated that iAECs are permissive to

infection with an MOI of 10, resulting in 50-60% infected cells. Acetylated tubulin staining revealed

variable cytoskeleton structures, reminiscent of different cell types, and interesting protrusions

that co-stained with viral NP-protein. The morphology of infected cells has key differences

compared to the other cell types used in our study; particularly, the proportion of individually

infected cells are greater than viral syncytia (Figure 4b). Remarkably, even at a high MOI of 10,

dose-responsive antiviral activity was observed with bovine lactoferrin (IC_{50} = 45 nM), human

lactoferrin (IC_{50} = 466 nM), S1RA (IC_{50} = 1 μM), and remdesivir (IC_{50} = 18 nM) (Figure 4a). At lower

MOIs, the infection rate in iAEC2s was more variable. However, with a lower MOI of 0.2 the dose

dependent antiviral efficacy was observed for amiodarone, lomitapide, ipratropium bromide,

clofazimine, gilteritinib and fedratinib (Supplementary Figure 6). This physiologically relevant

model is a proxy of human lung tissue and serves as an intermediate model to further validate

clinical potential of our identified lead compounds prior to *in vivo* studies.

DISCUSSION

In this study, we developed an experimental workflow based on high-content imaging and

morphological profiling that allows for rapid screening of FDA-approved compounds, leveraging

machine learning to score compounds based on many different cellular features relevant to

infection. Our morphological profiling analysis and corresponding UMAP embedding

demonstrated that SARS-CoV-2 infection in Huh7 cells is variable and highlights the necessity of

multivariate scoring systems to assess efficacy. From our SARS-CoV-2 screening, we identified

17 FDA-approved compounds that limit infection *in vitro*. Of these, six were previously reported

and serve as a benchmark validation of our endpoints and experimental approach, and eleven

were hitherto unknown. We demonstrate that this approach is versatile (i.e., it can be applied to

both transformed and more physiologically-relevant non-transformed cell lines) and can identify

the emergent properties of the infection as well as novel phenotypes that can be perturbed

through chemical inhibition.

Importantly, our study identified drugs that implicate new molecular targets/pathways in the

pathogenesis of SARS-CoV-2 and produce clinically testable and readily translatable hypotheses.

As an example, we observed dose-dependent antiviral activities of metoclopramide and

domperidone, two potent dopamine receptor D2 antagonists used to treat gastroesophageal

reflux disease and prevent other gastrointestinal symptoms, including nausea and vomiting[22].

Gastrointestinal symptoms have been increasingly reported in more than half of the patients

infected by SARS-CoV-2[2]. Notably, investigational drugs like hydroxychloroquine, lopinavir-

ritonavir, tocilizumab and others can be associated with gastrointestinal and hepatic adverse

events and hence are not ideal for patients already experiencing severe GI symptoms[23].

Metoclopramide and domperidone therefore represent a dual-target therapeutic option for

COVID-19 patients. In contrast, the pro-dopaminergic drugs carbidopa, levodopa, and

methyldopa promote infection, suggesting that the dopamine pathway may contribute to adverse

infection outcomes. Additionally, all FDA-approved MEK inhibitors exacerbate viral infection 3-

fold indicating a putative role of MEK in SARS-CoV-2 pathogenesis. These *in vitro* observations

should be validated through clinical research that examines whether concomitant presence of

drug and SARS-CoV-2 infection worsen COVID-19 symptoms.

As most FDA-approved drugs are optimized against human molecular targets, our screen helped

identify crucial host factors involved in SARS-CoV-2 infection. Z-FA-FMK, an irreversible inhibitor

of cysteine proteases, including cathepsins B, L, and S[24], exhibited potent antiviral activity. A

recent report using a pseudovirus indicated cathepsin L is an entry factor of SARS-CoV-2[25]. The

antiviral effect of Z-FA-FMK suggests that cathepsin L is a requirement also in the context of

SARS-CoV-2 infection and suggests that this molecule could be a useful investigational tool to

study virus entry. Similarly, fedratinib, approved by the FDA in 2019 for myeloproliferative

neoplasms[26], is an orally bioavailable semi-selective JAK2 inhibitor. JAK-inhibitors have been

proposed for COVID-19 to specifically inhibit TH17-mediated inflammatory responses. JAK-

inhibitors have been proposed for COVID-19 treatment to specifically inhibit TH17-mediated

inflammatory response[27,28] and to block numb-associated kinase responsible for clathrin-

mediated viral endocytosis[29]. Several JAK-inhibitors are currently evaluated in clinical trials for

COVID-19 management, including with baricitinib[30], jakotinib (ChiCTR2000030170), and

ruxolitinib (ChiCTR2000029580). For their inhibitory effect on innate immune response at the

cellular level, JAK-inhibitors could serve as useful tools in the future to elucidate the involvement

of the innate immune response in SARS-CoV-2 infection.

The sigma receptors (SigmaR1/R2) are permissive chaperones that mediate endoplasmic

reticulum stress response and lipid homeostasis[31], processes that have been implicated in early

stages of hepatitis C viral infection in Huh7 cells[32] and coronavirus pathogenesis[33]. We identified

two sigma receptor modulators amiodarone[34], and S1RA[35] with potent antiviral activity,

demonstrating IC_{50} values of 52 nM and 222 nM, respectively, with limited cell toxicity. Amiodarone

is approved for treatment of arrhythmias but, like hydroxychloroquine, has potent cardiotoxic side

effects through inhibition of the hERG ion channel[36] that limit therapeutic potential. S1RA has

completed phase II clinical trials for the treatment of neuropathic pain[37,38]. Although Gordon *et al.*

identified several other sigmaR1/R2 modulators that inhibited SARS-CoV-2 infection in Vero-E6

cells, antiviral activity for S1RA was not observed[9]. This suggests that the activity of S1RA is

dependent on host cell factors specific to each cell line and, promisingly, that human cells may

be more responsive to this compound, as observed in iAEC2s (Figure 5a).

Most noteworthy, our screen demonstrates lactoferrin as a SARS-CoV-2 inhibitor *in vitro* with

multimodal efficacy. We showed that lactoferrin strongly inhibits cellular binding of SARS-CoV-2

and is therefore a promising therapy for pre- and post-exposure prophylaxis. We also showed

dose-dependent efficacy in multiple cell types, including a non-transformed and clinically relevant

iPSC-derived model of alveolar epithelium. Lactoferrin gene expression has been shown

previously to be highly upregulated in response to SARS-CoV-1 infection[39] and, in addition to

enhancing natural killer cell and neutrophil activity, lactoferrin blocks viral entry through binding

to heparan sulfate proteoglycans. Lactoferrin retains anti-SARS-CoV-2 activity 24 hrs p.i., which

suggests additional MOA other than simple entry inhibition. Although we cannot conclude a

definitive and complete MOA, we show significant host cell modulation through increased

expression of several interferon-stimulated genes upon treatment with lactoferrin. Additionally,

lactoferrin has been previously shown to decrease the production of IL-6[40], which is one of the

key players of the "cytokine storm" produced by SARS-CoV-2 infection[41,42]. We found that

lactoferrin, either from bovine or human origin, retain activity in both the holo- and apo- forms, the

latter being the component of orally available lactoferrin supplements. Lactoferrin potential is

heightened by its ability to mitigate a high MOI SARS-CoV-2 infection in iAEC2 (Figure 5). Orally

available lactoferrin could be especially effective in resolving the gastrointestinal symptoms that

are present in COVID-19 patients[43]. The mechanisms may be similar to how lactoferrin reduces

human norovirus infection through induction of innate immune responses[44], especially as

lactoferrin gene polymorphisms are associated with increased susceptibility to infectious

diarrhea[45]. If lactoferrin reduces viral load in the GI tract, it could reduce fecal-oral transmission

of COVID-19[46].

Combination therapies are likely to be required for effectively treating SARS-CoV-2 infection, and

this approach has already shown promise. For example, combination therapy with interferon beta-

1b, lopinavir–ritonavir, and ribavirin showed efficacy against SARS-CoV-2 in a prospective, open-

label, randomized, phase 2 trial[47]. We show that lactoferrin potentiates the antiviral activity of both

remdesivir and hydroxychloroquine and could be used as a combination therapy with these drugs,

which are currently being used or studied for the treatment of COVID-19. Due to its wide

availability, limited cost, and lack of adverse effects, lactoferrin could be a rapidly deployable

option for both prophylaxis and the management of COVID-19. Likewise, ipratropium bromide, a

widely-used quaternary ammonium salt bronchodilator, holds promise as another agent for

combination therapies with potential to reduce bronchial viral burden. Although our findings are

promising, further studies are needed to confirm the efficacy of our lead antiviral compounds in

other representative *in vitro* cell lines and/or clinical studies.

High-content morphological cell profiling for drug repurposing screening enabled the identification

of both novel antivirals efficacious against SARS-CoV-2 and compounds that possibly exacerbate

SARS-CoV-2 infection. Confirmation in iAEC2s suggests high clinical translatability of these

compounds. This approach to preclinical testing has promise for identifying other anti-SARS-CoV-

2 drugs, rationally designing therapeutic combinations with multiple MOAs, and deployment of

optimized combinations in a rapid and systemic fashion.

Supplementary Information is available for this paper.

Correspondence and requests for materials should be addressed to jzsexton@umich.edu

327
328
329

ACKNOWLEDGEMENTS

Funding: University of Michigan Institute for Clinical and Health Research (MICHR) (NCATS - UL1TR002240) and its Center for Drug Repurposing. JZS is supported by the National Institute of Diabetes and Digestive and Kidney Diseases (R01DK120623). JWW is supported by the pharmacological sciences training program (PSTP) T32 training grant. CM is supported by Marie-Slodowska Curie individual fellowship (GA - 841247) and MICHR Postdoctoral Translational Scholars Program. KDA is supported by the I.M. Rosenzweig Junior Investigator Award from the Pulmonary Fibrosis Foundation. JRS is supported by the National Heart, Lung, and Blood Institute (NHLBI – R01HL119215), by the NIAID Novel Alternative Model Systems for Enteric Diseases (NAMSED) consortium (U19AI116482) and by grant number CZF2019-002440 from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation.

The authors would like to thank Matthew Chess for Amazon AWS support, Kevin Jan and Peyton Uhl at Yokogawa for imaging support, Nick Santoro at the University of Michigan Center for Chemical Genomics. We thank David Egan and Wienand Omta from Core Life Analytics for assisting high content data analytics as well as Philip Cheung and Brian Bolt at ReFactor Biosciences for assistance with HTS data registration. Finally, we thank Tracey Schultz and Dianne Jazdzyk for project management.

METHODS

Cells and virus. Vero E6, Caco-2 and Huh7 cells were maintained at 37°C with 5% CO_2 in Dulbecco's Modified Eagle's Medium (DMEM; Welgene), supplemented with 10% heat-

inactivated fetal bovine serum (FBS), HEPES, non-essential amino-acids, L-glutamine and 1X

Antibiotic-Antimycotic solution (Gibco). iPSC (SPC2 iPSC line, clone SPC2-ST-B2, Boston

University) derived alveolar epithelial type 2 cells (iAEC2s) were differentiated as previously

described and maintained as alveolospheres embedded in 3D Matrigel in "CK+DCI" media, as

previously described (Jacob et al. 2019). iAEC2s were passaged approximately every two weeks

by dissociation into single cells via the sequential application of dispase (2mg/ml, Thermo Fisher

Scientific, 17105-04) and 0.05% trypsin (Invitrogen, 25300054) and re-plated at a density of 400

cells/µl of Matrigel (Corning, 356231), as previously described (Jacob et al. 2019). SARS-CoV-2

WA1 strain was obtained by BEI resources and was propagated in Vero E6 cells. Viral titers were

determined by TCID50 assays in Vero E6 cells (Reed and Muench method) by microscopic

scoring. All experiments using SARS-CoV-2 were performed at the University of Michigan under

Biosafety Level 3 (BSL3) protocols in compliance with containment procedures in laboratories

approved for use by the University of Michigan Institutional Biosafety Committee (IBC) and

Environment, Health and Safety (EHS).

Viral titer determination. Vero E6, Caco-2 and Huh7 cells were seeded in a 48-well plate at

2x10^4 cells/well incubated overnight at 37°C with 5% CO2. Cells were then infected with SARS-

CoV-2 WA1 at a multiplicity of infection (MOI) of 0.2. One hour after infection, cells were harvested

(day 0 of infection) or kept at 37°C for 1, 2 and 3 days p.i. Viral titer determination was performed

by TCID50 assay on Vero E6 cells of the total virus (supernatant and intracellular fraction).

Alternatively, cells were harvested with Trizol and total cellular and viral RNA was extracted with

the ZymoGen Direct-zol RNA extraction kit. Viral RNA was quantified by RT-qPCR using the

2019-nCoV CDC qPCR Probe Assay and the probe set N1 (IDT technologies). IFNβ, viperin, MX1,

ISG15, IFITM3 and the housekeeping gene GAPDH mRNA levels were quantified by qPCR with

SsoAdvanced™ Universal SYBR® Green Supermix (Bio-Rad) with specific primers (IFNβ: F-

TTGACATCCCTGAGGAGATTAAGC, R- TCCCACGTACTCCAACTTCCA; MX1: F-

CCAGCTGCTGCATCCCACCC, R-AGGGGCGCACCTT CTCCTCA; ISG15: F-

TGGCGGGCAACGAATT, R- GGGTGATCTGCGCCTTCA; IFITM3: F-TCCCAC

GTACTCCAACTTCCA, R-AGCACCAGAAACACGTGCACT; GAPDH: F-

CTCTGCTCCTCCTGTTCGAC, R-GCGCCCCACCAAGCTCAAGA). Fold increase was

calculated by using the ΔΔCt method over non-infected untreated Huh7.

Viral infectivity assay. 384-well plates (Perkin Elmer, 6057300) were seeded with Huh7 cells at

3000 cells/well and allowed to adhere overnight. Compounds were then added to the cells and

incubated for 4 hours. The plates were then transferred to BSL3 containment and infected with

SARS-CoV-2 WA1 at a multiplicity of infection (MOI) of 0.2 in a 10 µL addition with shaking to

distribute virus. For the final dose-responses curves, porcine trypsin (Sigma-Aldrich, T0303) at a

final concentration of 2µg/ml was included during infection. After one hour of absorption, the virus

inoculum was removed, and media replaced with fresh compound. Uninfected cells and vehicle-

treated cells were included as positive and negative control, respectively. Two days post-infection,

cells were fixed with 4% PFA for 30 minutes at room temperature, permeabilized with 0.3% Triton

X-100 and blocked with antibody buffer (1.5% BSA, 1% goat serum and 0.0025% Tween 20). The

plates were then sealed, surface decontaminated, and transferred to BSL2 for staining with the

optimized fluorescent dye-set: anti-nucleocapsid protein (anti-NP) SARS-CoV-2 antibody

(Antibodies Online, Cat# ABIN6952432) overnight treatment at 4C followed by staining with

secondary antibody Alexa-647 (goat anti-mouse, Thermo Fisher, A21235), Hoechst-33342

pentahydrate (bis-benzimide) for nuclei staining (Thermo FIsher, H1398), HCS LipidTOX™ Green

Neutral Lipid Stain (Thermo Fisher, H34475), and HCS CellMask™ Orange for cell delineation

(Thermo Fisher H32713). iAEC2 maintained in 3D culture were dissociated to single cells and

seeded in collagen coated 384-well plates at a seeding density of 8000 cells/well in the presence

of 10 µM Y-27632 for the first 72 hours after plating (APExBIO, A3008 to grow to roughly 80%

confluence. Infection was performed at MOI of 10 in the presence of 2µg/ml of trypsin porcine

(Sigma-Aldrich, T0303). Staining protocol for the iAEC2s differed slightly with the addition of an

anti-acetylated tubulin primary antibody (Cell Signaling, 5335), instead of HCS CellMask Orange,

and the use of an additional secondary Alexa 488 antibody (donkey anti-rabbit, Jackson

ImmunoResearch, 711-545-152).

Viral Binding assay

Huh-7 cells were plated in 48-well plates at 100,000 cells per well and allowed to adhere

overnight. The following day, compounds were added at the indicated concentration in serum-

free DMEM and incubated for 1 hour at 4℃. Following compound incubation, cells were infected

with SARS-CoV-2 at an MOI of 10 for 1 hour at 4℃ to allow for viral binding. Cells were then

washed 3 times with ice cold PBS to remove unbound virus and RNA was extracted by using

the Direct-Zol RNA miniprep kit (Zymogen, R2052). Bound virus was then quantified by RT-

qPCR (see section Viral titer determination and host gene quantification) and percentages were

calculated over the Infected non-treated condition.

Multi-cycle cytopathogenic effect (CPE) reduction assay. Vero E6 were allowed to adhere

overnight in 96-well cell culture plates. A 1:2 10-point serial dilution of compounds (5000nM-5nM)

and SARS-CoV-2 at MOI of 0.002 were added. CPE was evaluated by microscopic scoring at

5dpi. The 50% inhibitory concentration (IC50) was calculated by logarithmic interpolation and is

defined as the concentration at which the virus-induced CPE is reduced by 50%.

Compound library. The compound library deployed for drug screening was created using the

FDA-Approved Drugs Screening Library (Item No. 23538) from Cayman Chemical Company. This

library of 875 compounds was supplemented with additional FDA approved drugs and rationally

included clinical candidates from other vendors including MedChemExpress, Sigma Aldrich, and

Tocris. Our library was formatted in five 384-well compound plates and was dissolved in DMSO

at 10 mM. Hololactoferrin (Sigma Aldrich, L4765), apolactoferrin (Jarrow Formulas, 121011),

native human lactoferrin (Creative BioMart, LFT-8196H) and transferrin (Sigma Aldrich, T2036)

were handled separately and added manually in cell culture media. Dilution plates were generated

for qHTS at concentrations of 2 mM, 1 mM, 500 μM, 250 μM and 50 μM and compounds were

dispensed at 1:1000 dilution.

qHTS primary screen and dose response confirmation. For the qHTS screen, compounds

were added to cells using a 50 nL pin tool Caliper Life Sciences Sciclone ALH 3000 Advanced

Liquid Handling system at the University of Michigan Center for Chemical Genomics (CCG).

Concentrations of 2 μM, 1 μM, 500 nM, 250 nM and 50 nM were included for the primary screen.

Post qHTS screen, all compounds were dispensed using an HP D300e Digital Compound

Dispenser and normalized to a final DMSO concentration of 0.1% DMSO. Confirmation dose

response was performed in triplicate and in 10-point:2-fold dilution.

Imaging. Stained cell plates were imaged on both Yokogawa CQ1 and Thermo Fisher CX5 high

content microscopes with a 20X/0.45NA LUCPlan FLN objective. Yokogawa CQ1 imaging was

performed with four excitation laser lines (405nm/488nm/561nm/640nm) with spinning disc

confocal and 100ms exposure times. Laser power was adjusted to yield optimal signal to noise

ratio for each channel. Maximum intensity projection images were collected from 5 confocal

planes with a 3-micron step size. Laser autofocus was performed and nine fields per well were

imaged covering approximately 80% of the well area. The Thermofisher CX5 with LED excitation

(386/23nm, 485/20nm, 560/25nm, 650/13nm) was also used and exposure times were optimized

to maximize signal/background. Nine fields were collected at a single Z-plane as determined by

image-based autofocus on the Hoechst channel. The primary qHTS screen was performed using

CX5 images and all dose-response plates were imaged using the CQ1.

Image segmentation and feature extraction. The open source CellProfiler software was used

in an Ubuntu Linux-based distributed Amazon AWS cloud implementation for segmentation,

feature extraction and results were written to an Amazon RDS relational database using MySQL.

A pipeline was developed to automatically identify the nuclei, cell, cytoplasm, nucleoli, neutral

lipid droplets and syncytia for feature extraction. Multiple intensity features and radial distributions

were measured for each object in each channel and cell size and shape features were measured.

Nuclei were segmented using the Hoechst-33342 image and the whole cell mask was generated

by expanding the nuclear mask to the edge of the Cell Mask Orange image.

Data pre-processing. Cell level data were pre-processed and analyzed in the open source

Knime analytics platform[48]. Cell-level data was imported into Knime from MySQL, drug treatment

metadata was joined, and features were centered and scaled. Features were pruned for low

variance (<5%) and high correlation (>95%) and resulted in 660 features per cell.

Statistical methods and hypothesis testing. Dose-response curves were fit and pairwise

differences between experimental conditions were tested using Prism (Graphpad Software,

San Diego, CA, USA). Other statistical tests were performed in the statistical programming

language and environment R.

Machine learning - infectivity score and field-level scoring. Multiple logistic regression as implemented in the statistical language and environment R was used to identify features characteristic of cells within infected wells. Models were fit to cells from infected and uninfected control wells in the first five plate-series of the quantitative high throughput screen. As an independent benchmark, these logistic regression models were validated against a manually selected set of individual infected and uninfected cells; features which degraded performance on the benchmark were excluded from the model. The final model included only virus channel intensity features in the cell and cytoplasm ROIs. As a threshold for initial classification, the minimum value from virus-infected cells in the benchmark was used; the final decision rule is given in Eq. 1.

(Eq.1) : A cell is infected if (Cells_Intensity_IntegratedIntensityEdge_Virus × 0.1487025 +

Cells_Intensity_MeanIntensityEdge_Virus × -38.40196 +

Cells_Intensity_MaxIntensityEdge_Virus × 42.70269 +

Cytoplasm_Intensity_StdIntensity_Virus × 42.54849) ≥ 1.525285

Then, individual field images from the infected control were categorized as confirmed-infected when the mean feature values, across all cells in the field, were above the threshold in Eq. 1. Using mean values for all 660 cell-profiler features in each field, a random forest classifier was trained to predict a probability of membership in the category of uninfected control fields vs confirmed-infected fields. The output of this random forest classifier is reported as "Probpos" (for the positive, uninfected control), throughout. Field level mean/median feature values were computed and a random forest model was fit between the positive control (32

uninfected wells) and the negative control (32 infected wells, 0.1% DMSO vehicle treated)

with 80/20 cross validation. The compound treated wells were scored with the RF model and

the efficacy score was normalized to the individual plate.

UMAP embedding. The embed_umap application of MPLearn (v0.1.0,

https://github.com/momeara/MPLearn) was used to generate UMAP embeddings. Briefly, each

for a set of cells, each feature was per-plate standardized and jointly orthogonalized using

sklearn.IncrementalPCA(n_components=379, batch_size=1000). Then features were embedded

into 2-dimensions using umap-learn (v0.4.1)[12] with umap. UMAP(n_components=2,

n_neighbors=15, min_dist=0, init='spectral', low_memory=True). Embeddings were visualized

using Holovies Datashader (v1.12.7)[49], using histogram equalization and the viridis color map.

Visualizing subsets was done in JMP Pro 14.

Data analytics. HC Stratominer (Core Life Analytics, Utrecht NL) was used as an independent

method for hit-calling and performs fully automated/streamlined cell-level data pre-processing and

score generation. IC Stratominer was also used to fit dose response curves for qHTS.

Compound registration and assay data registration were performed using the open source

ACAS platform (Refactor BioSciences github https://github.com/RefactorBio/acas).

Dose-response analysis and compound selection. In qHTS screening, a compound was

selected to be carried forward into full dose response confirmation when meeting one of the

following criteria: 1) Probpos greater than 0.75 for the median field in at least three concentrations,

with per-field cell counts at least 60% of the positive control, and without an observed standard

deviation in Probpos across-fields-in-the-well of 0.4 or greater, 2) a dose-response relationship

with Probpos was observed (by inspection) across the five concentrations tested, including

compounds with Propbos greater than 0.90 at the two highest concentrations, or 3) compounds

of interest not meeting this criteria were carried forward if reported positive in the literature or were

being evaluated in clinical trials for COVID-19.

Dose response analysis in the confirmation and combinatorial screening. Due to the spatial

inhomogeneity of infected cells across a single well, approximately half of the fields were

undersaturated, leading to a consistent distribution in Probpos that saturates in the top third of 27

rank-ordered fields (from 9 fields and triplicate wells) for each concentration tested. The Probpos

effect for a compound concentration was tabulated by averaging the top third of rank ordered

fields. Outlier fields with high Probpos values were visually inspected and eliminated if artifacts

(segmentation errors or debris) were observed. Cells treated with known fluorescence drugs

including Clofazimine, were confirmed to not have spectral interference. Dose response curves

were fit with Graphpad Prism using a semilog 4-parameter variable slope model.

544 **FIGURES AND LEGENDS**

545
546
Figure 1. Morphological profiling of SARS-CoV-2 infected Huh7 cells (MOI of 0.2 for 48 hrs). a)
representative field (left) with nuclei (cyan), neutral lipids (green), and SARS-CoV-2 NP
(magenta) and the NP image in the same area with "fire" false color LUT showing distinct
morphologies of infected cells showing small/round cells with a hollow center, cells with
protrusions, and large syncytia. Representative image was acquired on a Yokogawa CQ1 high-
content imager at with a 60X lens and visualized with Fiji ImageJ. b) 2 dimensional UMAP
embedding of 2.47 million infected cells using 379 morphological features. c) Cluster regions of

interest (ROI) in the UMAP are highlighted. d) UMAP projection colored by NP viral staining
intensity for four measurements where viral measurements with darker blue representing
greater value in those regions. e) For four ROIs, a representative cell is shown for the nuclear
(top), CellMask(middle), and SARS-CoV-2 NP channels (bottom).
558

Figure 2. a) Schematic representation of the anti-SARS-CoV-2 therapy discovery effort. 1) Compounds are administered to cells cultured on 384-well plates infected with SARS-CoV-2. Each plate contains 24 negative (infected) and 24 positive (non-infected) control wells to adjust for plate-to-plate variation. 2) Cells are fixed, stained, and imaged. Images are analyzed through a Cell Profiler-based pipeline which segments nuclei, cell boundaries, neutral lipid content and viral syncytia formation while extracting features of these cellular compartments. 3) Dose-response curves are calculated through multivariate-analysis to define per-image viral infectivity 4) Machine learning models are built around positive and negative control wells based on extracted features and applied to each drug condition. 5) Models inform on individual compound mode(s) of antiviral action through obtained features 6) confirmed antiviral hits; b) Dose-response curves of 16 hits of the drug screening. Graphs represent median SEM of 10-point 1:2 dilution series of selected compounds for N=3 biological replicates. IC50 were calculated based on normalization to the control and after fitting in GraphPad Prism.

575
Figure 3. Lactoferrin blocks SARS-CoV-2 replication at different stages of the viral cycle. a)
Huh7 cells were treated with lactoferrin (0 to 2.3 μM) and infected with SARS-CoV-2 (MOI of
0.2) in a 384-well plate. Plates were imaged using automated fluorescence microscopy and
processed using our image analysis pipeline to determine percent viral inhibition. Graph
indicates a dose-response (RED, IC50 = 308 μM). Cell viability is depicted in black. b) Huh7
were infected with SARS-CoV-2 (MOI of 1, 5 and 10; MOI of 0 indicates non-infected cells) and
treated with 2.3 μM of lactoferrin at 1 and 24 hrs p.i. Bars indicate the percentage of infected
cells in different conditions. Data is an average of eight replicates. Statistical significance
determined using multiple student's t-test with the Bonferroni-Dunn method, with alpha = 0.05.
Except for MOI of 0, all conditions (Untreated vs Lactoferrin, 1 hr or Untreated vs Lactoferrin, 24
hr) differ at P<0.0001. c-d) 2.5x104 Huh7 cells were infected with SARS-CoV-2 at MOI of 0.2.

48 hrs p.i., cells were harvested and RNA was extracted. Viral genome copies were calculated
with an absolute quantification method (standard curve) (c) and mRNA levels of cellular IFNβ,
MX1, ISG15 and IFITM3 (d) were calculated with ΔΔCt over non-infected Huh7. Data are
average, SD of N=2 biological replicates with n=3 technical replicates each. Statistical
significance determined using multiple student's t-test with the Bonferroni-Dunn method, with
alpha = 0.05. *P<0.001. e) Percentage of SARS-CoV-2 infected Huh7 cells upon treatment with
bovine apolactoferrin and hololactoferrin, native human lactoferrin and transferrin at a
concentration of 2.3 μM. f) 2-dimensional dose response heat maps of lactoferrin (0 to 2.3 μM)
in combination with remdesivir and hydroxychloroquine (0 to 30 nM and 0 to 10 μM,
respectively). Remdesivir combination was evaluated with a 0.2 MOI and HCQ was evaluated
with a MOI of 10 leading to a relative shift in lactoferrin
potency.

599
600   Figure 4. Antiviral activity of selected compounds was assessed in iAEC2 cells infected with
601   SARS-CoV-2 at MOI 10. Bovine and human lactoferrin exhibited IC50 of 44.9 and 466 nM
602   respectively. Remdesivir and S1RA exhibited IC50 of 18.4 nM and 1 μM respectively. Images of
603   nuclei (cyan), acetylated tubulin (green), and NP (magenta) from non-treated infected control,
604   IC50, and ICmax.

SUPPLEMENTARY INFORMATION

REFERENCES

1.   Xiao, F. *et al.* Evidence for Gastrointestinal Infection of SARS-CoV-2. *Gastroenterology* (2020) doi:10.1053/j.gastro.2020.02.055.

2.   Lin, L. *et al.* Gastrointestinal symptoms of 95 cases with SARS-CoV-2 infection. *Gut* (2020) doi:10.1136/gutjnl-2020-321013.

3.   Avula, A. *et al.* COVID-19 presenting as stroke. *Brain. Behav. Immun.* (2020) doi:10.1016/j.bbi.2020.04.077.

4.   Kochi, A. N., Tagliari, A. P., Forleo, G. B., Fassini, G. M. & Tondo, C. Cardiac and arrhythmic complications in patients with COVID-19. *Journal of Cardiovascular Electrophysiology* (2020) doi:10.1111/jce.14479.

5.   Mulangu, S. *et al.* A randomized, controlled trial of Ebola virus disease therapeutics. *N. Engl. J. Med.* (2019) doi:10.1056/NEJMoa1910993.

6.   Oprea, T. I. *et al.* Drug repurposing from an academic perspective. *Drug Discovery Today: Therapeutic Strategies* (2011) doi:10.1016/j.ddstr.2011.10.002.

7.   Chu, H. *et al.* Comparative tropism, replication kinetics, and cell damage profiling of SARS-CoV-2 and SARS-CoV with implications for clinical manifestations, transmissibility, and laboratory studies of COVID-19: an observational study. *The Lancet Microbe* (2020) doi:10.1016/s2666-5247(20)30004-5.

8.   Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is

Blocked by a Clinically Proven Protease Inhibitor. *Cell* (2020)

doi:10.1016/j.cell.2020.02.052.

9.   Gordon, D. E. *et al.* A SARS-CoV-2 protein interaction map reveals targets for drug

repurposing. *Nature* (2020) doi:10.1038/s41586-020-2286-9.

10.   McQuin, C. *et al.* CellProfiler 3.0: Next-generation image processing for biology. *PLoS*

Biol. (2018) doi:10.1371/journal.pbio.2005970.

11.   McInnes, L., Healy, J., Saul, N. & Großberger, L. UMAP: Uniform Manifold Approximation

and Projection. *J. Open Source Softw.* (2018) doi:10.21105/joss.00861.

12.   Riva, L. *et al.* A Large-scale Drug Repositioning Survey for SARS-CoV-2 Antivirals.

bioRxiv (2020) doi:10.1101/2020.04.16.044016.

13.   Katie Heiser *et al.* Identification of potential treatments for COVID-19 through artificial

intelligence enabled phenomic analysis of human cells infected with SARS-CoV-2.

bioRxiv (2020) doi:10.1101/2020.04.21.054387.

14.   Jeon, S. *et al.* Identification of antiviral drug candidates against SARS-CoV-2 from FDA-

approved drugs. *Antimicrob. Agents Chemother.* (2020) doi:10.1128/AAC.00819-20.

15.   Lang, J. *et al.* Inhibition of SARS pseudovirus cell entry by lactoferrin binding to heparan

sulfate proteoglycans. *PLoS One* (2011) doi:10.1371/journal.pone.0023710.

16.   Kell, D. B., Heyden, E. L. & Pretorius, E. The Biology of Lactoferrin, an Iron-Binding

Protein That Can Help Defend Against Viruses and Bacteria. *Front. Immunol.* (2020)

doi:10.3389/fimmu.2020.01221.

17.   Siqueiros-Cendón, T. *et al.* Immunomodulatory effects of lactoferrin. *Acta*

Pharmacologica Sinica (2014) doi:10.1038/aps.2013.200.

18.   Yeni, P. Update on HAART in HIV. in *Journal of Hepatology* (2006).

doi:10.1016/j.jhep.2005.11.021.

19.   Hurley, K. *et al.* Reconstructed Single-Cell Fate Trajectories Define Lineage Plasticity

Windows during Differentiation of Human PSC-Derived Distal Lung Progenitors. *Cell*

Stem Cell (2020) doi:10.1016/j.stem.2019.12.009.

20.   Jacob, A. *et al.* Derivation of self-renewing lung alveolar epithelial type II cells from

663   human pluripotent stem cells. *Nat. Protoc.* (2019) doi:10.1038/s41596-019-0220-0.

664   21.   Mason, R. J. Pathogenesis of COVID-19 from a cell biology perspective. *European*

665   *Respiratory Journal* (2020) doi:10.1183/13993003.00607-2020.

666   22.   Hibbs, A. M. & Lorch, S. A. Metoclopramide for the treatment of gastroesophageal reflux

667   disease in infants: A systematic review. *Pediatrics* (2006) doi:10.1542/peds.2005-2664.

668   23.   Hajifathalian, K. *et al.* SARS-COV-2 infection (coronavirus disease 2019) for the

669   gastrointestinal consultant. *World J. Gastroenterol.* (2020) doi:10.3748/wjg.v26.i14.1546.

670   24.   Roscow, O., Ganassin, R., Garver, K. & Polinski, M. Z-FA-FMK demonstrates differential

671   inhibition of aquatic orthoreovirus (PRV), aquareovirus (CSRV), and rhabdovirus (IHNV)

672   replication. *Virus Res.* (2018) doi:10.1016/j.virusres.2017.11.024.

673   25.   Ou, X. *et al.* Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its

674   immune cross-reactivity with SARS-CoV. *Nat. Commun.* (2020) doi:10.1038/s41467-020-

675   15562-9.

676   26.   Pardanani, A. *et al.* TG101209, a small molecule JAK2-selective kinase inhibitor potently

677   inhibits myeloproliferative disorder-associated JAK2V617F and MPLW515L/K mutations.

678   *Leukemia* (2007) doi:10.1038/sj.leu.2404750.

679   27.   Wu, D. & Yang, X. O. TH17 responses in cytokine storm of COVID-19: An emerging

680   target of JAK2 inhibitor Fedratinib. *J. Microbiol. Immunol. Infect.* (2020)

681   doi:10.1016/j.jmii.2020.03.005.

682   28.   Zhang, W. *et al.* The use of anti-inflammatory drugs in the treatment of people with

683   severe coronavirus disease 2019 (COVID-19): The experience of clinical immunologists

684   from China. *Clinical Immunology* (2020) doi:10.1016/j.clim.2020.108393.

685   29.   Stebbing, J. *et al.* COVID-19: combining antiviral and anti-inflammatory treatments. *The*

686   *Lancet Infectious Diseases* (2020) doi:10.1016/S1473-3099(20)30132-8.

30.   Treatment of Moderate to Severe Coronavirus Disease (COVID-19) in Hospitalized

Patients. (2020).

31.   Delprat, B., Crouzier, L., Su, T. P. & Maurice, T. At the Crossing of ER Stress and MAMs:

A Key Role of Sigma-1 Receptor? in *Advances in Experimental Medicine and Biology*

(2020). doi:10.1007/978-3-030-12457-1_28.

32.   Friesland, M., Mingorance, L., Chung, J., Chisari, F. V. & Gastaminza, P. Sigma-1

Receptor Regulates Early Steps of Viral RNA Replication at the Onset of Hepatitis C

Virus Infection. *J. Virol.* (2013) doi:10.1128/jvi.03557-12.

33.   Fung, T. S. & Liu, D. X. Coronavirus infection, ER stress, apoptosis and innate immunity.

Frontiers in Microbiology (2014) doi:10.3389/fmicb.2014.00296.

34.   Moebius, F. F., Reiter, R. J., Hanner, M. & Glossmann, H. High affinity of sigma1-binding

sites for sterol isomerization inhibitors: Evidence for a pharmacological relationship with

the yeast sterol C8-C7 isomerase. *Br. J. Pharmacol.* (1997) doi:10.1038/sj.bjp.0701079.

35.   Díaz, J. L. *et al.* Synthesis and biological evaluation of the 1-arylpyrazole class of σ1

receptor antagonists: Identification of 4-{2-[5-methyl-1- (naphthalen-2-yl)-1H-pyrazol-3-

yloxy]ethyl}morpholine (S1RA, E-52862). *J. Med. Chem.* (2012) doi:10.1021/jm3007323.

36.   Torres, V. *et al.* QT prolongation and the antiarrhythmic efficacy of amiodarone. *J. Am.*

Coll. Cardiol. (1986) doi:10.1016/S0735-1097(86)80272-8.

37.   Vidal-Torres, A. *et al.* Effects of the selective sigma-1 receptor antagonist S1RA on

formalin-induced pain behavior and neurotransmitter release in the spinal cord in rats. *J.*

Neurochem. (2014) doi:10.1111/jnc.12648.

38.   Gris, G. *et al.* The selective sigma-1 receptor antagonist E-52862 attenuates neuropathic

pain of different aetiology in rats. *Sci. Rep.* (2016) doi:10.1038/srep24591.

39.   Reghunathan, R. *et al.* Expression profile of immune response genes in patients with

severe acute respiratory syndrome. *BMC Immunol.* (2005) doi:10.1186/1471-2172-6-2.

40.   Cutone, A. *et al.* Lactoferrin prevents LPS-induced decrease of the iron exporter

ferroportin in human monocytes/macrophages. *BioMetals* (2014) doi:10.1007/s10534-

014-9742-7.

41. Conti, P. *et al.* Induction of pro-inflammatory cytokines (IL-1 and IL-6) and lung

inflammation by COVID-19: anti-inflammatory strategies. *Journal of biological regulators*

and homeostatic agents (2020) doi:10.23812/CONTI-E.

42. Lagunas-Rangel, F. A. & Chávez-Valencia, V. High IL-6/IFN-γ ratio could be associated

with severe disease in COVID-19 patients. *Journal of Medical Virology* (2020)

doi:10.1002/jmv.25900.

43. Han, C. *et al.* Digestive Symptoms in COVID-19 Patients With Mild Disease Severity:

Clinical Presentation, Stool Viral RNA Testing, and Outcomes. *Am. J. Gastroenterol.*

(2020) doi:10.14309/ajg.0000000000000664.

44. Oda, H. *et al.* Antiviral Effects of Bovine Lactoferrin on Human Norovirus. *Biochem. Cell*

Biol. (2020) doi:10.1139/bcb-2020-0035.

45. Mohamed, J. A. *et al.* A Novel Single-Nucleotide Polymorphism in the Lactoferrin Gene Is

Associated with Susceptibility to Diarrhea in North American Travelers to Mexico. *Clin.*

Infect. Dis. (2007) doi:10.1086/512199.

46. Gu, J., Han, B. & Wang, J. COVID-19: Gastrointestinal Manifestations and Potential

Fecal–Oral Transmission. *Gastroenterology* (2020) doi:10.1053/j.gastro.2020.02.054.

47. Hung, I. F.-N. *et al.* Triple combination of interferon beta-1b, lopinavir–ritonavir, and

ribavirin in the treatment of patients admitted to hospital with COVID-19: an open-label,

randomised, phase 2 trial. *Lancet* (2020) doi:10.1016/s0140-6736(20)31042-4.

48. Berthold, M. R. *et al.* KNIME: The konstanz information miner. in *4th International*

Industrial Simulation Conference 2006, ISC 2006 (2006). doi:10.1007/978-3-540-78246-

9_38.

49. Stevens, J.-L., Rudiger, P. & Bednar, J. HoloViews: Building Complex Visualizations

Easily for Reproducible Science. in *Proceedings of the 14th Python in Science*

Conference (2015). doi:10.25080/majora-7b98e3ed-00a.

740