1   **Genetic architecture and lifetime dynamics of inbreeding depression in a wild mammal**

4   Authors names and addresses:

6   Stoffel, M.A.[1]*, Johnston, S.E.[1], Pilkington, J.G.[1], Pemberton, J.M[1]

7
8   [1]Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh,

9   Edinburgh, EH9 3FL, United Kingdom

21   * Corresponding author:

22   Martin A. Stoffel

23   Postal address: Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, EH9 3FL, UK

24   E-mail: martin.stoffel@ed.ac.uk

Abstract

Inbreeding depression is a phenomenon of long-standing importance, but we know surprisingly little about its genetic architecture, precise effects and life-history dynamics in wild populations. Here, we combined 417K imputed SNP genotypes for 5952 wild Soay sheep with detailed long-term life-history data to explore inbreeding depression on a key fitness component, annual survival. Inbreeding manifests in long runs of homozygosity (ROH) and these are abundant in Soay sheep, covering on average 24% of the autosomal genome and up to 50% in the most inbred individuals. The ROH landscape is shaped by recombination rate variation and differs widely across the genome, including islands where up to 87% of the population have an ROH and deserts where the ROH prevalence is as low as 4%. We next quantified individual inbreeding as the proportion of the autosomal genome in ROH (F_{ROH}) and estimated its effect on annual survival. The consequences of inbreeding are severe; a 10% increase in F_{ROH} was associated with a 68% [95% CI 55-78%] decrease in the odds of survival. However, the strength of inbreeding depression is dynamic across the lifespan. We estimate depression to peak in young adults, to decrease into older ages and to be weaker in lambs, where inbreeding effects are possibly buffered by maternal care. Finally, using a genome-wide association scan on ROH, we show that inbreeding causes depression predominantly through many loci with small effects, but we also find three regions in the genome with putatively strongly deleterious mutations. Our study reveals population and genome-wide patterns of homozygosity caused by inbreeding and sheds light on the strength, dynamics and genetic architecture of inbreeding depression in a wild mammal population.

Introduction

Inbreeding depression, the reduced fitness of offspring from related parents, has been a core theme in evolutionary and conservation biology since Darwin [1]. The detrimental effects of inbreeding on a broad range of traits, individual fitness and population viability have now been recognized all across the animal and plant kingdoms [1-9]. However, despite the ubiquity of inbreeding depression, we still know very little about its most fundamental features in wild populations [7,10]: What are the genomic patterns of homozygosity caused by inbreeding? How strong is inbreeding depression in fitness and how do its effects change across different life stages? What is the distribution of effect sizes at loci causing inbreeding depression?

Inbreeding decreases fitness because it increases the fraction of the genome which is homozygous and identical-by-descent (IBD). This unmasks the effects of (partially-) recessive deleterious alleles or in rarer cases may decrease fitness at loci with heterozygote advantage [4,11]. While the probability of IBD at a genetic locus was traditionally estimated as the expected inbreeding coefficient based on a pedigree [12,13], modern genomic approaches enable us to gain a much more detailed picture. Genome-wide markers or whole genome sequences are now unraveling the genomic mosaic of homo- and heterozygosity and, unlike pedigree-based approaches, capture individual variation in homozygosity due to the stochastic effects of Mendelian segregation and recombination [10,14]. This makes it possible to quantify realized rather than expected individual inbreeding, and can provide insights into how inbreeding causes IBD along the genome [15,16].

An intuitive and powerful way of measuring IBD is through runs of homozygosity (ROH), which are long stretches of homozygous genotypes [17]. ROH arise when two IBD haplotypes come together, which happens more frequently with increasing parental relatedness. However, ROH do not only originate from recent inbreeding, but can also emerge when shorter IBD haplotypes are inherited from apparently unrelated individuals due to background relatedness in the population. Consequently, ROH are ubiquitous even in outbred human populations [18], and in cases of strong inbreeding can stretch along whole chromosomes, as observed in some Scandinavian wolves [15]. The proportion of the autosomal genome in ROH (F_{ROH}) is an estimate of realized individual inbreeding F [19] and has been used to uncover inbreeding depression in a wide range of traits in humans and farm animals [5,20,21]. While F_{ROH} condenses the information about an individual's IBD into a single number, the genomic location of ROH reflects where the genome is homozygous due to inbreeding and can therefore help to elucidate the genetic basis of inbreeding depression [10,22].

A major challenge towards a better understanding of inbreeding depression is in mapping the

underlying loci and estimating their effect sizes [10]. While simple recessive disease loci are relatively

straightforward to identify through homozygosity mapping [23,24], quantifying the genetic basis of

inbreeding depression in fitness and complex traits requires large samples in addition to dense

genomic data [22]. As a consequence, the genetic architecture of inbreeding depression has mostly

been studied in humans and livestock [25,26]. However, individual fitness will be different under natural

conditions and consequently there is a need to study inbreeding depression in wild populations to

understand its genetic basis in an evolutionary and ecological context. To date, only a handful of

studies have estimated inbreeding depression using genomic data in the wild [27-31]. While these

studies show that inbreeding depression in wild populations is more prevalent and more severe than

previously thought, all of them used genome-wide inbreeding coefficients and did not explore the

underlying genetic basis of depression.

Here, we combined long-term life-history data for 5952 free-living Soay sheep from the St. Kilda

archipelago, Scotland, with over 417K partially imputed genome-wide SNP markers for a detailed

genomic analysis of inbreeding depression. First, we quantified the genomic consequences of

inbreeding through patterns of ROH among individuals and across the genome. We then calculated

individual genomic inbreeding coefficients F_{ROH} to model inbreeding depression in annual survival

and estimate its strength and dynamics across the lifetime. Finally, we explored the genetic

architecture of inbreeding depression using a mixed-model based genome-wide association scan

on ROH to shed light on whether depression is caused by many loci with small effects, few loci with

large effects or a mixture of both.

Results

Study population. Soay sheep are descendants of early Bronze Age sheep and have lived on the

island of Soay in the St Kilda Archipelago for thousands of years [32]. Although the Soays have been

largely unmanaged, there is written and genomic evidence of an admixture event with the now

extinct Dunface breed approximately 150 years or 32 generations ago [33]. Since 1985, the sheep

inhabiting the Village Bay area of Hirta have been part of a long-term individual-based study [32].

Annual survival is assessed through routine mortality checks which are conducted throughout the

year, and over 80% of sheep in the study area are found after their death [27]. For all following analyses

we focused on a subset of 5952 sheep for which complete annual survival data and genomic data

were available. While many Soay sheep die early, some individuals live up to 15 years, resulting in a

total of 15889 observations for annual survival (Supplementary Figure 1).

Genotyping and imputation. All study individuals have been genotyped on on the Illumina Ovine

SNP50 BeadChip containing 51,135 SNP markers. In addition, 189 individuals have been genotyped

on the Ovine Infinium High-Density chip containing 606,066 SNPs. To increase the genomic

resolution for our analyses, we combined autosomal genotypes from both SNP chips with pedigree

information to impute missing SNPs in individuals genotyped at lower marker density using

AlphaImpute [34]. Cross-validation showed that imputation was successful, with a median of 99.3%

correctly imputed genotypes per individual (Supplementary Table 1). Moreover, the inferred

inbreeding coefficients F_{ROH} were very similar when comparing individuals genotyped on the high-

density chip (median F_{ROH} = 0.239) and individuals with imputed SNPs (median F_{ROH} = 0.241),

indicating no obvious bias in the abundance of inferred ROH based on imputed data

(Supplementary Figure 2). After quality control, the genomic dataset contained 417,373

polymorphic and autosomal SNPs with a mean minor allele frequency (MAF) of 23% and a mean call

rate of 99.5% across individuals.

Patterns of inbreeding in the genome. We first explored how inbreeding and a long-term small

population size (estimated N_e = 194, [35]) shaped patterns of ROH in Soay sheep (Figure 1). Individuals

had a mean of 194 ROH (sd = 11.6) longer than 1.2 Mb, which on average made up 24% of the

autosomal genome (i.e. mean F_{ROH} = 0.24, range = 0.18-0.50, Supplementary Figure 3). IBD in the

1% most inbred individuals compared to the 1% least inbred individuals was characterized through

longer ROH (mean: 5.27 Mb vs. 2.76 Mb), and slightly more abundant ROH (mean: 183 vs. 174).

Figure 1A contrasts ROH longer than 5 Mb between the seven most and least inbred individuals,

illustrating large differences in ROH length and coverage. The frequency of ROH in the population

also varied widely across the genome (Figure 1B). We scanned ROH in the population in non-

overlapping 500Kb windows, revealing regions with high ROH prevalence (ROH islands) and low

ROH prevalence (ROH deserts) on every chromosome (Figure 1B). The top ROH island on

chromosome 1 (227-227.5Mb) contained ROH in 87% of individuals, while only 4.4% of individuals

had an ROH in the top ROH desert on chromosome 11 (58.5-59Mb, see Supplementary Table 2 for

a list of the top ROH deserts and islands). Patterns of ROH on most chromosomes were shaped by

recombination rate variation (Supplementary Figure 4). On average across the genome, a

standardized (z-transformed) unit increase in recombination rate (measured as the summed fraction

of recombinants *r* in a window) was associated with a 2.3% decrease in ROH (β [95% confidence interval] = -0.023 [-0.026, -0.019], Supplementary Table 3 for model details).

Figure 1: ROH variation (A) among the seven most and least inbred individuals and (B) across the Soay sheep genome. Panel A shows ROH longer than 5 Mb in the seven individuals with the highest F_{ROH} in the seven top rows and the seven individuals with the lowest F_{ROH} in the seven bottom rows. Panel B shows the genome-wide ROH prevalence among all 5952 individuals in non-overlapping 500Kb windows. The colour gradient on Panel B has been scaled according to the ROH prevalence distribution, which is shown as a density plot in the figure legend.

Inbreeding depression in survival. Survival is a key fitness component. In Soay sheep, more than half of all individuals die over their first winter, minimizing their chances to reproduce (Supplementary Figure 1). The distribution of individual inbreeding coefficients F_{ROH} in different age classes revealed that highly inbred individuals rarely survive their early years of life and never reach old ages (Figure 2A). We modeled this using a binomial animal model with annual survival as a response, including 15889 observations for 5952 sheep over more than three decades (see Methods for details). The effect of inbreeding on survival was strong: A 10% increase in F_{ROH} was associated with a 68% reduction in the odds of survival (β [95% credible interval, CI] = 0.32 [0.22, 0.45]). Over the lifetime, inbreeding depression becomes weaker, with the slope of the interaction between F_{ROH} and age predicting an increase in the odds of survival (β [CI] = 1.19 [1.06, 1.32]). In addition, the

slope of the interaction between F_{ROH} and being a lamb (β [CI] = 1.94 [1.05, 3.34]) indicates that

inbreeding depression is weaker in lambs, though with considerable uncertainty (Figure 2B). As

model estimates for odds of survival are not easy to interpret, we plotted the predicted effects of

F_{ROH} on survival at different life stages (Figure 2C). Lambs (age 0) have a substantially lower survival

probability but the curve is on average less steep than for individuals at age one, reflecting stronger

depression in yearlings. Moreover, the three predicted survival curves for the age classes one, four

and seven differ in slopes, reflecting the estimated decrease of inbreeding depression across life. A

detailed list of the model estimates including the variable coding and standardization is shown in

Supplementary Table 4. As an alternative way to visualize the lifetime dynamics of inbreeding

depression, we also fitted separate models for each age class, which show essentially the same

pattern (Supplementary Figure 5). Finally we estimated the genetic load in Soay sheep through the

diploid number of lethal equivalents 2B. Lethal equivalents are a concept rooted in population

genetics, where one lethal equivalent is equal to a group of mutations which, if dispersed across

individuals, would on average cause one death [36]. We followed suggestions by Nietlisbach et al [37]

and refitted the survival model with a Poisson distribution and logarithmic link function using a

simplified model structure without interactions for a better comparability across studies. This gave

an estimate of 2B = 4.57 (95% CI 2.61-6.55) lethal equivalents for Soay sheep annual survival.

Figure 2: Inbreeding depression in annual survival. Panel A shows the distributions of inbreeding

coefficients F_{ROH} in Soay sheep age classes ranging from 0 to 10 years. Panel B shows the model estimates

(posterior mean and 95% CI) for individual inbreeding F_{ROH} and its interaction with age and being a lamb. F_{ROH}

has been centered around its mean and transformed so that a unit change and hence the model estimate

reflects the change in survival probability for a 0.1 increase in F_{ROH} from the mean. Panel C shows the predicted

survival probability and 95% CI over the range of inbreeding coefficients F_{ROH} for lambs (age 0), and 1, 4 and 7-

year old individuals respectively, while holding sex and twin constant at 1 (male) and 0 (no twin). The predictions

for age classes 1, 4, and 7 exceed the range of the data (panel A) but are shown across the full range for

comparability.

Genetic architecture of inbreeding depression. To quantify the survival consequences of being

IBD at each SNP location, we used a modified genome-wide association study (GWAS). Unlike in

traditional GWAS where p-values of additive SNP effects are of interest, we analysed a binary fixed

effect of ROH status, which indicates whether a focal SNP is in an ROH or not [25,26,38]. We fitted a

binomial mixed model of annual survival for each SNP, with individual ROH status at the focal SNP

position as fixed effect, and controlled for the additive SNP effect and mean individual inbreeding

F_{ROH} (based on all autosomes except for the focal chromosome) alongside a range of other individual

traits and environmental effects (see Methods for details). Under the null hypothesis that ROH status

does not have an effect on survival at any SNP position, we would expect a 50/50 distribution of

negative and positive GWAS effects, as all model estimates are due to chance. In contrast, we found

many more negative than positive effects of ROH status on survival across the genome than expected

by chance (Figure 3A, 3B; 247K neg. vs. 171K pos.; exact binomial test p = 2.2 * 10^{-16}). Using binomial

GLMs with effect direction as binary response variable, we also estimated an increasing proportion

of negative effects among larger estimates ($ß$ [95% CI] = 0.27 [0.26, 0.28]) and lower p-values ($ß$ [95%

CI] = 0.95 [0.94, 0.96]). Consequently, it is likely that many loci with small effects contribute to

inbreeding depression in survival. Moreover, ROH status reached genome-wide significance in three

regions on chromosomes 3, 10, and 23, revealing putative large effect mutations (Figure 3C,

Supplementary Table 5). While largely deleterious recessive variants might be expected to occur in

regions of elevated heterozygosity where they are rarely expressed in their homozygous state, we

did not observe clear common patterns of genetic diversity in these three regions, although

heterozygosity is slightly elevated and ROH frequency is lower around the top SNPs on

chromosomes 10 and 23 (Supplementary Figure 6).

Figure 3: GWAS of SNP-wise ROH status effect on annual survival. Regional inbreeding depression was
conceptualised and tested as a binary fixed effect of whether a SNP was part of an ROH (and hence IBD) or not.
Panel A and B show the distribution of effect sizes and p-values for this SNP-wise ROH effect. The yellow
histograms showing positive effects are superimposed on top of the purple histograms showing negative
effects to highlight that there was a substantially larger proportion of negative ROH status effects than expected
by chance. Panel C shows a Manhattan plot of the SNP-wise ROH status p-values across the genome. The dotted
line marks the genome-wide significance threshold for a Bonferroni correction which was based on the effective
number of tests when accounting for LD. All genome-wide significant p-values were associated with negative
effects on survival.

Discussion

The Soay sheep on St. Kilda have existed at a small population size and relative isolation for
thousands of years [32]. As a consequence, levels of IBD are high in the population and ROH make up
nearly a quarter of the average autosomal Soay sheep genome. This is more than three times as high
as the average F_{ROH} estimated across 78 mammal species based on genome-sequence data [39], but
still slightly lower than in some extremely inbred and very small populations such as mountain
gorillas [40], Scandinavian grey wolves [15] or Isle Royale wolves [41]. The prevalence of ROH varied
substantially across the genome and was broadly shaped by recombination rate variation, which is
known to impact ROH patterns alongside other factors such as gene density and selection [18,42,43].
We showed that most chromosomes contain both deserts where ROH are rare and islands where
ROH are common. In the most extreme regions as few as 4% and up to 87% of individuals had ROH
and this has potential implications for the distribution of deleterious mutations. ROH islands are likely
to contain few recessive deleterious alleles, as these are regularly exposed to selection when
homozygous and hence likely to be purged from the population. Expectations for deleterious
mutations in ROH deserts are more difficult to formulate because the (near-) absence of ROH could

be caused by several mechanisms, such as a high recombination rate or balancing selection.
However, a lack of ROH could potentially pinpoint regions harbouring lethal or semi-lethal alleles,
though this has only been demonstrated in farm animals using very large sample sizes (e.g. Jenko
et al., 2019).

We have only recently begun to understand the precise consequences of inbreeding for individual
fitness in natural populations. In Soay sheep, we found that the odds of survival decrease by 68%
with a mere 10% increase in F_{ROH}, adding to a small yet growing body of genomic studies showing
that inbreeding depression is stronger in wild populations than previously thought [27-29,31]. Other
recent examples include lifetime breeding success in red deer, which is reduced by up to 95% in
male offspring from half-sib matings [31] and lifetime reproductive success in helmeted honeyeaters,
which is up to 90% lower with a 9% increase in homozygosity [29]. The traditional way to compare
inbreeding depression among studies is to estimate the genetic load of a population using lethal
equivalents [36], although differences in methodology and inbreeding estimates often do not allow
direct comparisons [37]. We estimated the diploid number of lethal equivalents 2B for Soay sheep
annual survival at 4.57 (95% CI 2.61-6.55). This is a low to moderate inbreeding load compared to
other estimates from wild mammals obtained from appropriate statistical models [37], but
corroborates with theoretical expectations of lower genetic load in small populations [7] and a recent
comparative study estimating lower genetic load in smaller mammal populations [45].

Inbreeding depression is dynamic across life, and genomic measures are starting to unravel how
inbreeding depression affects fitness at different life-stages in wild populations [27,28,31]. Under the
mutation accumulation hypothesis [46], the adverse effects of deleterious mutations expressed late in
life should become stronger as selection becomes less efficient. This can theoretically increase the
effects of inbreeding depression later in life [47], although empirical evidence is sparse [48,49]. In contrast
to the idea of mutation accumulation, we showed that inbreeding depression in Soay sheep
generally becomes weaker over the lifetime. In addition, the sample for each successive age class
consists of increasingly outbred individuals (Figure 2C) due to a higher death rate among inbred
individuals earlier in life. This suggests that the effects of intragenerational purging [50] might
outweigh mutation accumulation in shaping the dynamics of inbreeding depression across the
lifetime. As a notable exception, we estimated inbreeding depression to be slightly weaker in lambs,
which may be because it is partially buffered by maternal care. Parental care has indeed been shown
to reduce inbreeding depression in burying beetles [51], but there is only limited evidence in mammals

to date [52,53]. Overall, we estimated inbreeding depression over a Soay sheep's life to start slightly

weaker in lambs, reaching its maximum in early adulthood and then decrease thereafter.

The effect size distribution of loci underpinning inbreeding depression has to our knowledge not

been studied in wild populations using fitness data, although deleterious mutations have been

predicted from sequencing data, for example in ibex and Isle Royale wolves [41,54]. Theoretical

predictions about the relative importance of weakly and strongly deleterious (partially-) recessive

alleles will depend on many factors, such as the distribution of dominance and selection coefficients

for mutations relative to the effective population size, and the frequency of inbreeding [55,56]. However,

we could expect that small populations purge largely deleterious recessive mutations more

efficiently as these are more frequently exposed to selection in the homozygous state [7,57,58], while

weakly deleterious mutations can more often drift to higher frequencies. We estimated the effect of

ROH status on Soay sheep survival at every SNP position within a GWAS framework. The effect size

distribution contained predominantly negative effects of ROH status on survival, particularly towards

larger effect sizes, showing that likely many alleles with weakly deleterious effects (or low

frequencies) contribute to inbreeding depression in survival.

Three regions in the genome also harboured putatively strongly deleterious alleles. This is

unexpected, as Soay sheep have a long-term small population size with an estimated N_e of 197 [35],

and highly deleterious recessive mutations should be rapidly purged. On the one hand however, it

is possible that genetic drift counteracted the effects of purifying selection and allowed deleterious

mutations to increase in frequency and be detected in a GWAS. On the other hand, a relatively recent

admixture event with the Dunface sheep breed around 150 years ago [33] could have introduced

deleterious variants into the population and recent selection has not been efficient enough to purge

them from the population yet. Identifying the loci carrying these strongly deleterious recessive

alleles will be challenging as ROH overlapping a given SNP vary in length among individuals, which

makes it difficult to pinpoint an exact effect location. Nevertheless, it is possible to identify the

haplotypes harbouring deleterious recessive alleles with large effects and monitor their frequency

changes in real time in natural populations, or select out individuals carrying them in conservation

breeding programs. To sum up, our study showed how genome-wide marker information for a large

sample of individuals with known fitness can deepen our understanding of the genetic architecture

and lifetime dynamics of inbreeding depression in the wild.

Methods

Study population, pedigree assembly and annual survival measurements. The Soay sheep is a primitive sheep breed descended from Bronze Age domestic sheep and has lived unmanaged on the St. Kilda archipelago, Scotland for thousands of years. When the last human inhabitants left St. Kilda in 1932, 107 Soays were transferred to the largest island, Hirta, and have roamed the island freely and unmanaged ever since. The population increased and fluctuates nowadays between 600 and 2200 individuals. A part of the population in the Village Bay area of Hirta (57 49'N, 8 34'W) has been the subject of a long-term individual based study since 1985 [32]. Most individuals born in the study area (95%) are ear-tagged and DNA samples are obtained from ear punches or blood sampling. Routine mortality checks are conducted throughout the year with peak mortality occurring at the end of winter and beginning of spring. Overall, around 80% of deceased animals are found [27]. For the analyses in this paper, survival was defined as dying (0) or surviving (1) from the 1st May of the previous year to the 30th April of that year, with measures available for 5952 individuals from 1979 to 2018. We focused on annual measures as this allowed us to incorporate the effects of age and environmental variation.

To assemble the pedigree, we inferred parentage for each individual using 438 unlinked SNP markers from the Ovine SNP50 BeadChip, on which most individuals since 1990 have been genotyped [59]. Based on these 438 markers, we inferred pedigree relationships using the R package Sequoia [60]. In the few cases where no SNP genotypes were available, we used either field observations (for mothers) or microsatellites [61]. All animal work was carried out according to UK Home Office procedures and was licensed under the UK Animals (Scientific Procedures) Act of 1986 (Project License no. PPL70/8818).

Genotyping. We genotyped a total of 7,700 Soay on the Illumina Ovine SNP50 BeadChip containing 51,135 SNP markers. To control for marker quality, we first filtered for SNPs with minor allele frequency (MAF) > 0.001, SNP locus genotyping success > 0.99 and individual sheep genotyping success > 0.95. We then used the check.marker function in GenABEL version 1.8-0 [62] with the same thresholds, including identity by state with another individual < 0.9. This resulted in a dataset containing 39,368 polymorphic SNPs in 7700 sheep. In addition, we genotyped 188 sheep on the Ovine Infinium HD SNP BeadChip containing 606,066 SNP loci. These sheep were specifically selected to maximise the genetic diversity represented in the full population as described in Johnston et al. [63], and underwent the same quality control as above, resulting in 430,702 polymorphic SNPs for 188 individuals. All genotype positions were based on the Oar_v3.1 sheep genome assembly (GenBank assembly ID GCA_000298735.1 [64])

Genotype imputation. In order to impute genotypes to high density, we merged the datasets from the 50K SNP chip and from the HD SNP chip using PLINK v1.90b6.12 with –bmerge [65]. This resulted in a dataset with 436,117 SNPs including 33,068 SNPs genotyped on both SNP chips. For genotype imputation, we discarded SNPs on the X chromosome and focused on the 419,281 SNPs located on autosomes. The merged dataset contained nearly complete genotype information for 188 individuals which have been genotyped on the HD chip, and genotypes at 38,130 SNPs for 7700 individuals which have been genotyped on the 50K chip. To impute the missing SNPs, we used AlphaImpute v1.98 [34], which combines information on shared haplotypes and pedigree relationships for phasing and genotype imputation. AlphaImpute works on a per-chromosome basis, and phasing and imputation are controlled using a parameter file (for the exact parameter file, see analysis code). Briefly, we phased individuals using core lengths ranging from 1% to 5% of the SNPs on a given chromosome over 10 iterations, resulting in a haplotype library. Based on the haplotype library, missing alleles were imputed using the heuristic method over five iterations which allowed us to use genotype information imputed in previous iterations. We only retained imputed genotypes for which all phased haplotypes matched and did not allow for errors. We also discarded SNPs with call rates below 95% after imputation. Overall, this resulted in a dataset with 7691 individuals, 417,373 SNPs and a mean genotyping rate per individual of 99.5 % (range 94.8%-100%).

To evaluate the accuracy of the imputation we used 10-fold leave-one-out cross-validation. In each iteration, we masked the genotypes unique to the high-density chip for one random individual that had been genotyped at high-density (HD) and imputed the masked genotypes. This allowed a direct comparison between the true and imputed genotypes. The imputation accuracy of the HD individuals should reflect of the average imputation accuracy across the whole population, because HD individuals were selected to be representative of the genetic variation observed across the pedigree (see Johnston et al., 2016 for details).

ROH calling and individual inbreeding coefficients. The final dataset contained genotypes at 417,373 SNPs autosomal SNPs for 5925 individuals for which annual survival data was available. We called runs of homozygosity (ROH) with a minimum length of 1200Kb and spanning at least 50 SNPs with the --homozyg function in Plink [65] and the following parameters: --homozyg-window-snp 50 --homozyg-snp 50 --homozyg-kb 1200 --homozyg-gap 300 --homozyg-density 200 --homozyg-window-missing 2 --homozyg-het 2 --homozyg-window-het 2. We chose 1200Kb as the minimum ROH length because between-individual variability in ROH abundance becomes very low for shorter ROH. Moreover, ROH of length 1200Kb extend well above the LD half decay in the population, thus

capturing variation in IBD due to more recent inbreeding rather than linkage disequilibrium

(Supplementary Figure 7). The minimum ROH length of 1200Kb also reflects the expected length

when the underlying IBD haplotypes had a most recent common ancestor haplotype 32 generations

ago, calculated as (100 / (2*g)) cM / 1.28 cM/Mb where g is 32 generations and 1.28 is the sex-

averaged genome-wide recombination rate in Soay sheep [63,66]. We then calculated individual

inbreeding coefficients F_{ROH} by summing up the total length of ROH for each individual and dividing

this by the total autosomal genome length (2,452Mb).

ROH landscape and recombination rate variation. To quantify variation in population-wide ROH

prevalence across the genome, we used a sliding window approach. We first calculated the number

of ROH overlapping each SNP position in the population using PLINK --homozyg. We then

calculated the mean number of ROH overlapping SNPs in 500 Kb non-overlapping sliding windows

in the population (Figure 1B). To estimate the top 0.5% ROH deserts and islands, we discarded

windows with less than 35 SNPs (the percentile of windows with the lowest SNP density). To estimate

the relationship between ROH prevalence and recombination rate we then quantified the

recombination rate in 500Kb window. We calculated recombination rate as the sum of the

recombination fractions *r* between consecutive loci in each window, using data from the Soay sheep

linkage map [63]. We then constructed a linear mixed model in lme4 [67] with population-wide ROH

prevalence per window as response, window recombination rate and chromosome size as fixed

effects and chromosome ID as random intercept (Supplementary Table 3). The fixed effects in the

model were standardised using z-transformation. To estimate confidence intervals, we used

parametric bootstrapping implemented in the tidy function of the broom.mixed package [68].

Modelling inbreeding depression in survival. We modelled the effects of inbreeding depression

in annual survival using a Bayesian animal model in INLA [69]. Annual survival data consists of a series

of 1s followed by a 0 in the year of a sheep's death, or only a 0 if it died as a lamb, and we

consequently used a binomial distribution with a logit link for the model. We used the following

model structure:

$\Pr(surv_i = 1) = logit^{-1}(\beta_0 + F_{ROH_i}\beta_1 + age_i\beta_2 + sex_i\beta_3 + twin_i\beta_4 + lamb_i\beta_5 + F_{ROH_i}age_i\beta_6 +$

$F_{ROH_i}lamb_i\beta_7 + \alpha_j^{capture\ year} + \alpha_k^{birth\ year} + \alpha_l^{id} + u_l^{ped})$

$\alpha_j^{capture\ year} \sim N(0, \sigma_{year}^2), \qquad for\ j = 1, ..., 40$

$\alpha_k^{birth\ year} \sim N(0, \sigma_{birth\ year}^2), \qquad for\ k = 1, ..., 40$

$$\alpha_l^{id} \qquad \sim N\left(0, \sigma_{id}^2\right), \qquad\qquad for\ l = 1, \dots, 5925$$

$$u_l^{ped} \qquad \sim N\left(0, A\sigma_A^2\right) \qquad\qquad for\ l = 1, \dots, 5925$$

Here, $\Pr(surv_i = 1)$ is the probability of survival for observation i, which depends on the intercept β_0, a series of fixed effects β_1 to β_7 , the random effects α which are assumed to be normally distributed with mean 0 and variance σ^2 and the breeding values u_l^{ped} which have a dependency structure corresponding to the pedigree, with a mean of 0 and a variance of $A\sigma_A^2$, where A is the relationship matrix and σ_A^2 is the additive genetic variance. As fixed effects, we fitted the individual inbreeding coefficient F_{ROH} (continuous), the *age* of the individual (continuous), *sex* (0 = female, 1 = male), a variable indicating whether an individual was born as a *twin* (0 = singleton, 1 = twin) and whether the individual is a *lamb* before its first year of age (0 = not a lamb, 1 = lamb). We also fitted an interaction term of F_{ROH} with *age* to estimate how inbreeding depression changes across the lifetime and an interaction term of F_{ROH} with *lamb* to estimate potential differences in inbreeding depression in lambs, as they receive maternal care which could ameliorate inbreeding depression. As random intercepts we fitted the birth year of an individual, the capture year to account for survival variation among years and the sheep identity to account for repeated measures. Finally, we controlled for additive genetic relatedness by including a covariance structure proportional to the pedigree relatedness matrix, as has previously been described for INLA models [70]. For all random effects, we used log-gamma priors with both shape and inverse-scale parameter values of 0.5.

To be able to interpret the slopes of the main effects of F_{ROH} and age despite their interactions at the mean value of the respective other variable (rather than at 0), we centered both variables [71,72]. Furthermore, we transformed F_{ROH} from its original range 0-1 to 0-10, which allowed us to directly interpret model estimates as resulting from a 10% increase in genome-wide IBD rather than the difference between a completely outbred and a completely inbred individual. Finally, we report model estimates on the response scale (as odds of survival) in the main paper, and on the link scale (as log-odds of survival) in the Supplementary Material.

Estimating lethal equivalents. The traditional way of comparing inbreeding depression among studies is to quantify the inbreeding load in terms of lethal equivalents, i.e. a group of genes which would cause on average one death if dispersed in different individuals and made homozygous [36]. However, differences in statistical methodology and inbreeding measures make it difficult to compare the strength of inbreeding depression in terms of lethal equivalents among studies [37]. Here, we used the approach suggested by Nietlisbach et al. (2019) and refitted the survival animal model with a Poisson distribution and a logarithmic link function. We were interested in lethal

equivalents for the overall strength of inbreeding depression rather than its lifetime dynamics, so we

fitted a slightly simplified animal model with F_{ROH}, age, age^2, twin and sex as fixed effects and birth

year, capture year, individual id and pedigree relatedness as random effects. The slope of F_{ROH}

estimates the decrease in survival due to a 10% increase in F_{ROH}, so we calculated the number of

diploid lethal equivalents 2B as $-(\beta_{FROH})/0.10 * 2$ where β_{FROH} is the Poisson model slope for F_{ROH}.

Mapping inbreeding depression. To map the effects of inbreeding depression in survival across

the genome, we used a modification of a genome-wide association study [25,26,38]. For each of the

~417K SNPs, we fitted a binomial mixed model with logit link in lme4 [67] with the following model

structure:

$$\Pr(surv_i = 1) = logit^{-1}(\beta_0 + SNP_{ROH_i}\beta_1 + SNP_{ADD_i}\beta_2 + F_{ROH_{mod_i}}\beta_3 + age_i\beta_4 + age_i^2\beta_5 + sex_i\beta_6 +$$

$$twin_i\beta_7 + \beta_{8-15}PC_{1-7} + \alpha_j^{capture\ year} + \alpha_k^{birth\ year} + \alpha_l^{id})$$

$$\alpha_j^{capture\ year} \sim N(0, \sigma_{capture\ year}^2), \quad for\ j = 1, \dots, 40$$

$$\alpha_k^{birth\ year} \sim N(0, \sigma_{birth\ year}^2), \quad for\ k = 1, \dots, 40$$

$$\alpha_l^{id} \sim N(0, \sigma_{id}^2), \quad for\ l = 1, \dots, 5925$$

Where the effect of interest is that of SNP_{ROH}, a binary variable indicating whether a SNP is in an ROH

(SNP_{ROH} = 1) or not (SNP_{ROH} = 0). SNP_{ADD} is the additive effect for the focal SNP (0,1,2 for

homozygous, heterozygous, homozygous for the alternative allele, respectively), and controls for the

possibility that a potential negative effect of ROH status is simply an additive effect. $F_{ROH_{mod}}$ is the

mean inbreeding coefficient of the individual based on all chromosomes except for the chromosome

where the focal SNP is located. We fitted F_{ROH} as we were interested in the effect of ROH status at a

certain locus (SNP_{ROH}), on top of the average individual inbreeding coefficient. Sex and twin are

again binary variables representing the sex of the individual and whether it is a twin. Age and age^2

control for the linear and quadratic effects of age, and are fitted as continuous covariates. Because

it was computationally impractical to fit 417K binomial animal models with our sample size and

because the additive genetic variance in survival is negligible (see Supplementary Table 4), we did

not fit an additive genetic effect. Instead, we used the top 7 principal components of the variance-

standardized additive relationship matrix (PC_{1-7}) as fixed effects [65]. Again, we added birth year,

capture year and individual id as random effects. For each fitted model, we extracted the estimated

slope of SNP_{ROH} and its p-value calculated based on a Wald-Z test. To determine a threshold for

genome-wide significance we used the 'simpleM' procedure [73]. The method uses composite linkage

disequilibrium to create a SNP correlation matrix and calculates the effective number of independent tests. We then used this value for a Bonferroni correction [74] of p-values resulting in a genome-wide significance threshold of $p < 1.28 * 10^{-6}$.

Data availability

Data will be deposited on Dryad upon acceptance.

Code availability

All code was written in R 3.6.1 [75] and all plots were made with ggplot2 [76]. The full analysis code is available on GitHub (https://github.com/mastoffel/sheep_ID).

Acknowledgements

We thank the National Trust for Scotland for permission to work on St. Kilda and QinetiQ, Eurest and Kilda Cruises for logistics and support. We thank I. Stevenson and many volunteers who have helped with fieldwork on the island and all those who have contributed to keeping the project going. SNP genotyping was conducted the Wellcome Trust Clinical Research Facility Genetic Core. This work has made extensive use of the Edinburgh Compute and Data Facility (http://www.ecdf.ed.ac.uk/). We thank John Hickey, Steve Thorn, Andrew Whalen and Martin Johnsson for help with the imputation. We are grateful for discussions with David Clark, Jon Slate, Jisca Huisman, Jarrod Hadfield, Peter Visscher, Anna Hewett, Deborah and Brian Charlesworth and the Wild Evolution Group. We also thank Emily Humble for comments on an earlier draft of the manuscript. The project was funded through an outgoing Postdoc fellowship from the German Science Foundation (DFG) awarded to MAS and a Leverhulme Grant (RPG-2019-072) awarded to JMP and SEJ. Field data collection has been supported by NERC over many years, and most of the SNP genotyping was supported by an ERC Advanced Grant to JMP.

References

1. Darwin, C. *The effect of cross and self fertilization in the vegetable kingdom*. (John Murray, 1876).

2. Angeloni, F., Ouborg, N. J. & Leimu, R. Meta-analysis on the association of population size and life history with inbreeding depression in plants. *Biological Conservation* **144**, 35–43 (2011).

3. Bozzuto, C., Biebach, I., Muff, S., Ives, A. R. & Keller, L. F. Inbreeding reduces long-term growth of Alpine ibex populations. *Nature ecology & evolution* **3**, 1359–1364 (2019).

4. Charlesworth, D. & Willis, J. H. The genetics of inbreeding depression. *Nature Reviews Genetics* (2009).

5. Clark, D. W. *et al.* Associations of autozygosity with a broad range of human phenotypes. *Nature Communications* **10**, 1–17 (2019).

6. Crnokrak, P. & Roff, D. A. Inbreeding depression in the wild. *Heredity* **83**, 260–270 (1999).

7. Hedrick, P. W. & Garcia-Dorado, A. Understanding inbreeding depression, purging, and genetic rescue. *Trends in ecology & evolution* **31**, 940–952 (2016).

8. Keller, L. Inbreeding effects in wild populations. *Trends in ecology & evolution* **17**, 230–241 (2002).

9. Ralls, K., Ballou, J. D. & Templeton, A. Estimates of lethal equivalents and the cost of inbreeding in mammals. *Conservation biology* **2**, 185–193 (1988).

10. Kardos, M., Taylor, H. R., Ellegren, H., Luikart, G. & Allendorf, F. W. Genomics advances the study of inbreeding depression in the wild. *Evolutionary applications* **9**, 1205–1218 (2016).

11. Hedrick, P. W. What is the evidence for heterozygote advantage selection? *Trends in ecology & evolution* **27**, 698–704 (2012).

12. Pemberton, J. Measuring inbreeding depression in the wild: the old ways are the best. *Trends in ecology & evolution* **19**, 613–615 (2004).

13. Wright, S. Coefficients of Inbreeding and Relationship. *The American Naturalist* **56**, 330–338 (1922).

14. Franklin, I. R. The distribution of the proportion of the genome which is homozygous by descent in inbred individuals. *Theoretical population biology* **11**, 60–80 (1977).

15. Kardos, M. *et al.* Genomic consequences of intensive inbreeding in an isolated wolf population. *Nature Ecology & Evolution* **2**, 124–131 (2018).

16. Kardos, M., Luikart, G. & Allendorf, F. W. Measuring individual inbreeding in the age of genomics: marker-based measures are better than pedigrees. *Heredity* **115**, 63 (2015).

17. Broman, K. W. & Weber, J. L. Long homozygous chromosomal segments in reference families
from the centre d'Etude du polymorphisme humain. *The American Journal of Human Genetics*
65, 1493–1500 (1999).

18. Gibson, J., Morton, N. E. & Collins, A. Extended tracts of homozygosity in outbred human
populations. *Hum Mol Genet* **15**, 789–795 (2006).

19. McQuillan, R. *et al.* Runs of homozygosity in European populations. *The American Journal of*
Human Genetics **83**, 359–372 (2008).

20. Curik, I., Ferenčaković, M. & Sölkner, J. Inbreeding and runs of homozygosity: A possible
solution to an old problem. *Livestock Science* **166**, 26–34 (2014).

21. Leroy, G. Inbreeding depression in livestock species: review and meta-analysis. *Animal*
Genetics **45**, 618–628 (2014).

22. Ceballos, F. C., Joshi, P. K., Clark, D. W., Ramsay, M. & Wilson, J. F. Runs of homozygosity:
windows into population history and trait architecture. *Nature Reviews Genetics* (2018).

23. Kijas, J. W. Detecting regions of homozygosity to map the cause of recessively inherited
disease. in *Genome-wide association studies and genomic prediction* 331–345 (Springer,
2013).

24. Lander, E. S. & Botstein, D. Homozygosity mapping: a way to map human recessive traits with
the DNA of inbred children. *Science* **236**, 1567–1570 (1987).

25. Ferenčaković, M., Sölkner, J., Kapš, M. & Curik, I. Genome-wide mapping and estimation of
inbreeding depression of semen quality traits in a cattle population. *Journal of Dairy Science*
100, 4721–4730 (2017).

26. Pryce, J. E., Haile-Mariam, M., Goddard, M. E. & Hayes, B. J. Identification of genomic regions
associated with inbreeding depression in Holstein and Jersey dairy cattle. *Genetics Selection*
Evolution **46**, (2014).

27. Bérénos, C., Ellis, P. A., Pilkington, J. G. & Pemberton, J. M. Genomic analysis reveals
depression due to both individual and maternal inbreeding in a free-living mammal
population. *Molecular ecology* **25**, 3152–3168 (2016).

28. Chen, N., Cosgrove, E. J., Bowman, R., Fitzpatrick, J. W. & Clark, A. G. Genomic consequences
of population decline in the endangered Florida scrub-jay. *Current Biology* **26**, 2974–2979
(2016).

29. Harrisson, K. A. *et al.* Lifetime Fitness Costs of Inbreeding and Being Inbred in a Critically
Endangered Bird. *Current Biology* **29**, 2711-2717.e4 (2019).

30. Hoffman, J. I. *et al.* High-throughput sequencing reveals inbreeding depression in a natural
population. *Proceedings of the National Academy of Sciences of the United States of America*
111, 3775–3780 (2014).

31. Huisman, J., Kruuk, L. E., Ellis, P. A., Clutton-Brock, T. & Pemberton, J. M. Inbreeding
depression across the lifespan in a wild mammal population. *Proceedings of the National*
Academy of Sciences **113**, 3585–3590 (2016).

32. Clutton-Brock, T. H. & Pemberton, J. M. *Soay sheep: dynamics and selection in an island*
population. (Cambridge University Press, 2004).

33. Feulner, P. G. D. *et al.* Introgression and the fate of domesticated genes in a wild mammal
population. *Molecular Ecology* **22**, 4210–4221 (2013).

34. Hickey, J. M., Kinghorn, B. P., Tier, B., van der Werf, J. H. & Cleveland, M. A. A phasing and
imputation method for pedigreed populations that results in a single-stage genomic
evaluation. *Genetics Selection Evolution* **44**, 9 (2012).

35. Kijas, J. W. *et al.* Genome-Wide Analysis of the World's Sheep Breeds Reveals High Levels of
Historic Mixture and Strong Recent Selection. *PLOS Biology* **10**, e1001258 (2012).

36. Morton, N. E., Crow, J. F. & Muller, H. J. An estimate of the mutational damage in man from
data on consanguineous marriages. *Proceedings of the National Academy of Sciences of the*
United States of America **42**, 855 (1956).

37. Nietlisbach, P., Muff, S., Reid, J. M., Whitlock, M. C. & Keller, L. F. Nonequivalent lethal
equivalents: Models and inbreeding metrics for unbiased estimation of inbreeding load.
Evolutionary applications **12**, 266–279 (2019).

38. Keller, M. C. *et al.* Runs of homozygosity implicate autozygosity as a schizophrenia risk factor.
PLoS genetics **8**, (2012).

39. Brüniche-Olsen, A., Kellner, K. F., Anderson, C. J. & DeWoody, J. A. Runs of homozygosity
have utility in mammalian conservation and evolutionary studies. *Conservation Genetics* **19**,
1295–1307 (2018).

40. Xue, Y. *et al.* Mountain gorilla genomes reveal the impact of long-term population decline and
inbreeding. *Science* **348**, 242–245 (2015).

41. Robinson, J. A. *et al.* Genomic signatures of extensive inbreeding in Isle Royale wolves, a
population on the threshold of extinction. *Science Advances* **5**, eaau0757 (2019).

42. Wang, H., Lin, C.-H., Chen, Y., Freimer, N. & Sabatti, C. Linkage disequilibrium and haplotype
homozygosity in population samples genotyped at a high marker density. *Human heredity* **62**,
175–189 (2006).

43. Pemberton, T. J. *et al.* Genomic patterns of homozygosity in worldwide human populations.
The American Journal of Human Genetics **91**, 275–292 (2012).

44. Jenko, J. *et al.* Analysis of a large dataset reveals haplotypes carrying putatively recessive lethal
and semi-lethal alleles with pleiotropic effects on economically important traits in beef cattle.
Genetics Selection Evolution **51**, 9 (2019).

45. van der Valk, T., de Manuel, M., Marques-Bonet, T. & Guschanski, K. Estimates of genetic load
in small populations suggest extensive purging of deleterious alleles. *bioRxiv* 696831 (2019).

46. Medawar, P. B. *An Unsolved Problem of Biology*. (H.K. Lewis, 1952).

47. Charlesworth, B. & Hughes, K. A. Age-specific inbreeding depression and components of
genetic variance in relation to the evolution of senescence. *Proceedings of the National*
Academy of Sciences **93**, 6140–6145 (1996).

48. Husband, B. C. & Schemske, D. W. Evolution of the magnitude and timing of inbreeding
depression in plants. *Evolution* **50**, 54–70 (1996).

49. Wilson, A. J. *et al.* Evidence for a genetic basis of aging in two wild vertebrate populations.
Current Biology **17**, 2136–2142 (2007).

50. Enders, L. S. & Nunney, L. Reduction in the cumulative effect of stress-induced inbreeding
depression due to intragenerational purging in Drosophila melanogaster. *Heredity* **116**, 304
(2016).

51. Pilakouta, N., Jamieson, S., Moorad, J. A. & Smiseth, P. T. Parental care buffers against
inbreeding depression in burying beetles. *Proceedings of the National Academy of Sciences*
112, 8031–8035 (2015).

52. Nielsen, J. F. *et al.* Inbreeding and inbreeding depression of early life traits in a cooperative
mammal. *Molecular Ecology* **21**, 2788–2804 (2012).

53. Wells, D. A., Cant, M. A., Nichols, H. J. & Hoffman, J. I. A high-quality pedigree and genetic
markers both reveal inbreeding depression for quality but not survival in a cooperative
mammal. *Molecular ecology* **27**, 2271–2288 (2018).

54. Grossen, C., Guillaume, F., Keller, L. F. & Croll, D. Purging of highly deleterious mutations
through severe bottlenecks in Alpine ibex. *Nature Communications* **11**, 1–12 (2020).

55. Bataillon, T. & Kirkpatrick, M. Inbreeding depression due to mildly deleterious mutations in
finite populations: size does matter. *Genetics Research* **75**, 75–81 (2000).

56. Glémin, S. How are deleterious mutations purged? Drift versus nonrandom mating. *Evolution*
57, 2678–2687 (2003).

57. Kyriazis, C. C., Wayne, R. K. & Lohmueller, K. E. High genetic diversity can contribute to
extinction in small populations. *bioRxiv* 678524 (2019).

58. Robinson, J. A., Brown, C., Kim, B. Y., Lohmueller, K. E. & Wayne, R. K. Purging of strongly
deleterious mutations explains long-term persistence and absence of inbreeding depression
in island foxes. *Current Biology* **28**, 3487–3494 (2018).

59. Bérénos, C., Ellis, P. A., Pilkington, J. G. & Pemberton, J. M. Estimating quantitative genetic
parameters in wild populations: a comparison of pedigree and genomic approaches.
Molecular ecology **23**, 3434–3451 (2014).

60. Huisman, J. Pedigree reconstruction from SNP data: parentage assignment, sibship clustering
and beyond. *Molecular ecology resources* **17**, 1009–1024 (2017).

61. Morrissey, M. B. *et al.* The prediction of adaptive evolution: empirical application of the
secondary theorem of selection and comparison to the breeder's equation. *Evolution:*
International Journal of Organic Evolution **66**, 2399–2410 (2012).

62. Aulchenko, Y. S., Ripke, S., Isaacs, A. & Van Duijn, C. M. GenABEL: an R library for genome-
wide association analysis. *Bioinformatics* **23**, 1294–1296 (2007).

63. Johnston, S. E., Bérénos, C., Slate, J. & Pemberton, J. M. Conserved Genetic Architecture
Underlying Individual Recombination Rate Variation in a Wild Population of Soay Sheep (Ovis
aries). *Genetics* **203**, 583–598 (2016).

64. Jiang, Y. *et al.* The sheep genome illuminates biology of the rumen and lipid metabolism.
Science **344**, 1168–1173 (2014).

65. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage
analyses. *The American Journal of Human Genetics* **81**, 559–575 (2007).

66. Thompson, E. A. Identity by descent: variation in meiosis, across genomes, and in populations.
Genetics **194**, 301–326 (2013).

67. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models using lme4.
Journal of Statistical Software **67**, (2015).

68. Bolker, B. & Robinson, D. *broom.mixed: Tidying Methods for Mixed Models. R package version*
0.2.4. (2019).

69. Rue, H., Martino, S. & Chopin, N. Approximate Bayesian inference for latent Gaussian models
by using integrated nested Laplace approximations. *Journal of the royal statistical society:*
Series b (statistical methodology) **71**, 319–392 (2009).

70. Holand, A. M., Steinsland, I., Martino, S. & Jensen, H. Animal models and integrated nested
Laplace approximations. *G3: Genes, Genomes, Genetics* **3**, 1241–1251 (2013).

71. Gelman, A. & Hill, J. *Data analysis using regression and multilevel/hierarchical models.*
(Cambridge university press, 2006).

72. Schielzeth, H. Simple means to improve the interpretability of regression coefficients. *Methods*
in Ecology and Evolution (2010).
73. Gao, X., Starmer, J. & Martin, E. R. A multiple testing correction method for genetic association
studies using correlated single nucleotide polymorphisms. *Genetic Epidemiology: The Official*
Publication of the International Genetic Epidemiology Society **32**, 361–369 (2008).
74. Holm, S. A simple sequentially rejective multiple test procedure. *Scandinavian journal of*
statistics 65–70 (1979).
75. R Core Team. *R: A language and environment for statistical computing. R Foundation for*
Statistical Computing. (2019).
76. Wickham, H. *ggplot2: elegant graphics for data analysis*. (Springer, 2016).
686