Functional reconstitution of a bacterial CO_2 concentrating mechanism in *E. coli*

Authors: Avi I. Flamholz[1], Eli Dugan[1], Cecilia Blikstad[1], Shmuel Gleizer[2], Roee Ben-Nissan[2], Shira Amram[2], Niv Antonovsky[2,†], Sumedha Ravishankar[1,‡], Elad Noor[2,§], Arren Bar-Even[3], Ron Milo[2,*] & David F. Savage[1,*]

Affiliations:

[1] Department of Molecular and Cell Biology, University of California, Berkeley, California 94720, United States

[2] Department of Plant and Environmental Sciences, Weizmann Institute of Science, Rehovot 76100, Israel

[3] Max Planck Institute of Molecular Plant Physiology, Am Mühlenberg 1, 14476 Potsdam, Germany

[†] Present address: Laboratory of Genetically Encoded Small Molecules, The Rockefeller University, 1230 York Avenue, New York, NY, 10065, USA

[‡] Present address: Division of Biological Sciences, Section of Molecular Biology, University of California, San Diego, La Jolla, California, 92093, United States.

[§] Present address: Institute of Molecular Systems Biology, Eidgenössische Technische Hochschule Zürich, Zürich CH-8093, Switzerland

*Correspondence to: ron.milo@weizmann.ac.il (R.M.) and savage@berkeley.edu (D.F.S.)

Abstract

Many photosynthetic organisms employ a CO_2 concentrating mechanism (CCM) to increase the rate of CO_2 fixation via the Calvin cycle. CCMs catalyze ≈50% of global photosynthesis, yet it remains unclear which genes and proteins are required to produce this complex adaptation. We describe the construction of a functional CCM in a non-native host, achieved by expressing genes from an autotrophic bacterium in an engineered *E. coli* strain. Expression of 20 CCM genes enabled *E. coli* to grow by fixing CO_2 from ambient air into biomass, with growth depending on CCM components. Bacterial CCMs are therefore genetically compact and readily transplanted, rationalizing their presence in diverse bacteria. Reconstitution enabled genetic experiments refining our understanding of the CCM, thereby laying the groundwork for deeper study and engineering of the cell biology supporting CO_2 assimilation in diverse organisms.

One Sentence Summary

A bacterial CO_2 concentrating mechanism enables *E. coli* to fix CO_2 from ambient air.

Introduction

Nearly all carbon in the biosphere enters by CO_2 fixation in the Calvin-Benson-Bassham cycle (Raven et al., 2017). Ribulose Bisphosphate Carboxylase/Oxygenase - commonly known as rubisco - is the CO_2 fixing enzyme in this cycle (Wildman, 2002) and likely the most abundant enzyme on Earth (Bar-On and Milo, 2019). As rubisco is abundant and central to biology, one might expect it to be an exceptional catalyst, but it is not. Photosynthetic rubiscos are modest enzymes, with carboxylation turnover numbers (k_{cat}) ranging from 1-10 s^{-1} (Flamholz et al., 2019; Iñiguez et al., 2020). Moreover, all known rubiscos catalyze a competing oxygenation of the five-carbon organic substrate, ribulose 1,5-bisphosphate (Bowes and Ogren, 1972; Cleland et al., 1998; Flamholz et al., 2019).

Rubisco arose > 2.5 billion years ago, when Earth's atmosphere contained little O_2 and abundant CO_2 (Fischer et al., 2016; Shih et al., 2016). In this environment, rubisco's eponymous oxygenase activity could not have hindered carbon fixation or the growth of CO_2-fixing organisms. Present-day atmosphere, however, poses a problem for plants and other autotrophs: their primary carbon source, CO_2, is relatively scarce ($\approx 0.04\%$) while a potent competing substrate, O_2, is abundant ($\approx 21\%$).

CO_2 concentrating mechanisms (CCMs) arose multiple times over the last 2 billion years (Flamholz and Shih, 2020; Raven et al., 2017) and overcome this problem by concentrating CO_2 near rubisco (Figure 1A). In elevated CO_2 environments most active sites are occupied with CO_2 and not O_2. As such, high CO_2 increases the rate of carboxylation and competitively inhibits oxygenation (Bowes and Ogren, 1972) thereby improving overall carbon assimilation (Figure 1B). Today, at least four varieties of CCMs are found in plants, algae and bacteria (Flamholz and Shih, 2020; Raven et al., 2017), organisms with CCMs are collectively responsible for $\approx 50\%$ of global net photosynthesis (Raven et al., 2017), and some of the most productive human crops (e.g. maize and sugarcane) rely on CCMs.

CCMs are particularly common among autotrophic bacteria: all Cyanobacteria and many Proteobacteria have CCM genes (Kerfeld and Melnicki, 2016; Rae et al., 2013). Bacterial CCMs rely on two crucial features: (i) energy-coupled inorganic carbon uptake at the cell membrane and (ii) a 200+ MDa protein organelle called the carboxysome that encapsulates rubisco with a carbonic anhydrase enzyme (Mangan et al., 2016; McGrath and Long, 2014). In the prevailing model of the carboxysome CCM, inorganic carbon uptake produces a high, above-equilibrium cytosolic HCO_3^- concentration (≈ 30 mM) that diffuses into the carboxysome, where carbonic anhydrase activity produces a high carboxysomal CO_2 concentration that promotes efficient carboxylation by rubisco (Figure 1A-B).

As CCMs accelerate CO_2 fixation, there is great interest in transplanting them into crops (Ermakova et al., 2020; McGrath and Long, 2014). Carboxysome-based CCMs are especially attractive because they natively function in single cells and appear to rely on a tractable number of genes (Lin et al., 2014; Long et al., 2018; Occhialini et al., 2016; Orr et al., 2020). Modeling suggests that introducing bacterial CCM components could improve plant photosynthesis

(McGrath and Long, 2014), especially if aspects of plant physiology can be modulated via genetic
engineering (Wu et al., 2019). However, expressing bacterial rubiscos and carboxysome
components has, so far, uniformly resulted in transgenic plants displaying impaired growth (Lin et
al., 2014; Long et al., 2018; Occhialini et al., 2016; Orr et al., 2020). More generally, as our
understanding of the genes and proteins participating in the carboxysome CCM rests mostly on
loss-of-function genetic experiments in native hosts (Cai et al., 2009; Desmarais et al., 2019;
Marcus et al., 1986; Price and Badger, 1989a), it is possible that some genetic, biochemical and
physiological aspects of CCM function remain unappreciated. We therefore sought to test whether
current understanding is sufficient to reconstitute the bacterial CCM in a non-native bacterial host,
namely *E. coli*.
88

89
Figure 1. Twenty genes form the basis of a bacterial CCM. *(**A**) The bacterial CCM consists of at least two essential*
components - energy-coupled carbon uptake and carboxysome structures that encapsulate rubisco with a carbonic
anhydrase (CA) enzyme (Mangan et al., 2016; McGrath and Long, 2014). Transport generates a large cytosolic HCO_3^-
*pool, which is rapidly converted to high carboxysomal CO_2 concentration by the carboxysomal CA. (**B**) Elevated CO_2*
increases the rubisco carboxylation rate (green) and suppresses oxygenation by competitive inhibition (grey). [O_2] was
*set to 270 μM for rate calculations. (**C**) H. neapolitanus CCM genes are mostly contained in a 20 gene cluster*

(Desmarais et al., 2019) expressing rubisco and its associated chaperones (green), carboxysome structural proteins
(purple), and an inorganic carbon transporter (orange).

Using a genome-wide screen in the CO_2-fixing proteobacterium *H. neapolitanus*, we recently
demonstrated that a 20-gene cluster encodes all activities required for the CCM, at least in
principle (Desmarais et al., 2019). These genes include rubisco large and small subunits, the
carboxysomal carbonic anhydrase, seven structural proteins of the α-carboxysome (Bonacci et
al., 2012), an energy-coupled inorganic carbon transporter (Desmarais et al., 2019; Scott et al.,
2019), three rubisco chaperones (Aigner et al., 2017; Mueller-Cajar, 2017; Wheatley et al., 2014),
and four genes of unknown function (Figure 1C). We aimed to test whether these genes are
sufficient to establish a functioning CCM in *E. coli*.

Figure 2. CCMB1 depends on rubisco
carboxylation for growth on glycerol. (A) Ribose-
5-phosphate (Ri5P) is required for nucleotide
biosynthesis. Deletion of ribose-phosphate
isomerase (Δrpi) in CCMB1 blocks ribulose-5-
phosphate (Ru5P) metabolism in the pentose
phosphate (PP) pathway. Expression of rubisco (H.
neapolitanus cbbLS) and phosphoribulokinase (S.
elongatus PCC7942 prk) on the p1A plasmid (B)
permits Ru5P metabolism, thus enabling growth on
M9 glycerol media in 10% CO_2 (C). Mutating the
rubisco active site (p1A cbbL⁻) abrogates growth, as
does mutating ATP-binding residues of prk (p1A
prk⁻). (D) CCMB1:p1A grows well under 10% CO_2,
but fails to grow in ambient air. Cells grown on M9
glycerol media throughout. The algorithmic design
of CCMB1 is described in figure supplement 1 and
the mechanism of rubisco-dependence is
diagrammed in figure supplement 2. Figure
supplement 3 shows CCMB1:p1A growth
phenotypes on various media and figure
supplement 4 demonstrates that rubisco
oxygenation is not required for growth by
demonstrating growth in the absence of O_2.
Acronyms: ribulose 1,5-bisphosphate (RuBP), 3-
phosphoglycerate (3PG).

Results

As *E. coli* is a heterotroph, consuming organic carbon molecules to produce energy and biomass,
it does not natively rely on rubisco. Therefore, in order to evaluate the effect of heterologous CCM
expression, we first designed an *E. coli* strain that depends on rubisco carboxylation for growth.
To grow on glycerol as the sole carbon source, *E. coli* must synthesize ribose 5-phosphate (Ri5P)
for nucleic acids. Synthesis of Ri5P via the pentose phosphate pathway forces co-production of
ribulose 5-phosphate (Ru5P). Deletion of ribose 5-phosphate isomerase (*rpiAB* genes, denoted
Δrpi), however, makes Ru5P a metabolic "dead-end" (Figure 2A). Expression of
phosphoribulokinase (*prk*) and rubisco creates a "detour" pathway converting Ru5P and CO_2 into

two units of the central metabolite 3-phosphoglycerate (3PG), enabling Ru5P metabolism and
growth (Figure 2A). Additionally, cytosolic carbonic anhydrase activity is incompatible with the
bacterial CCM (Price and Badger, 1989b). We therefore constructed a strain, named CCMB1 for
"**CCM B**ackground **1**", lacking *rpiAB* and all endogenous carbonic anhydrases (Methods).

As predicted, CCMB1 required rubisco and prk for growth on glycerol minimal media in 10% CO_2
(Figures 2B-C). When expressing rubisco and *prk* on the p1A plasmid (Figure 2B), CCMB1 also
grew reproducibly in an anoxic mix of 10:90 CO_2:N_2 (Figure 2 - figure supplement 4) implying that
rubisco carboxylation is sufficient for growth on glycerol media and rubisco-catalyzed oxygenation
of RuBP is not required. CCMB1:p1A failed to grow on glycerol media in ambient air, however,
presumably due to insufficient carboxylation at low CO_2 (Figure 2D). That is, CCMB1:p1A displays
the "high-CO_2 requiring" phenotype that is the hallmark of CCM mutants (Marcus et al., 1986;
Price and Badger, 1989a).

157
Figure 3. Expression of 20 CCM genes permits growth of CCMB1 in ambient air. Time course data give
representative growth curves from a bioreactor bubbling ambient air. CCMB1:pCB' + pCCM' grows well (purple, "full
CCM"), while rubisco and prk alone are insufficient for growth in ambient air (grey, CCMB1:p1A+vec). Inset: a plate
reader experiment in biological triplicate (different shades) gave the same result. Expressing the full complement of
CCM genes led to an increase in culture density (optical density at 600 nm) of ≈0.6 units after 80 hours of cultivation.
Bootstrapping was used to calculate a 99.9% confidence interval of 0.56-0.64 OD units for the effect of expressing the
full CCM during growth in ambient air. Figure supplement 1 shows triplicate growth curves and evaluates statistical
significance.

We expected that expressing a functional CO_2-concentrating mechanism would cure CCMB1 of
its high-CO_2 requirement and permit growth in ambient air. We therefore generated two plasmids,
pCB and pCCM, that together express all 20 genes from the *H. neapolitanus* CCM cluster (Figure
1C). pCB encodes ten carboxysome genes (Bonacci et al., 2012), including rubisco large and
small subunits, along with *prk*. The remaining *H. neapolitanus* genes, including putative rubisco
chaperones (Aigner et al., 2017; Mueller-Cajar, 2017; Wheatley et al., 2014) and an inorganic
carbon transporter (Desmarais et al., 2019; Scott et al., 2019), were cloned into the second
plasmid, pCCM.

CCMB1 co-transformed with pCB and pCCM initially failed to grow on glycerol media in ambient
air. We therefore conducted selection experiments, described fully in Figure S5, that resulted in

the isolation of mutant plasmids conferring growth in ambient air. Briefly, CCMB1:pCB + pCCM
cultures were grown to saturation in 10% CO_2. These cultures were washed and plated on glycerol
minimal media (Methods). Colonies became visible after 20 days of incubation in ambient air
(Figure S5). Deep-sequencing of plasmid DNA revealed mutations in regulatory sequences (e.g.
a promoter and transcriptional repressor) but none in sequences coding for CCM components
(Table S4). Individual post-selection plasmids pCB' and pCCM' were reconstructed by PCR,
resequenced, and transformed into naive CCMB1 (Methods). As shown in Figure 3, pCB' and
pCCM' together enabled reproducible growth of CCMB1 in ambient air, suggesting that the 20
genes expressed are sufficient to produce a heterologous CCM without any genomic mutations.
187

188
Figure 4. Growth in ambient air depends on the known components of the bacterial CCM. *We generated plasmid*
*variants carrying inactivating mutations to known components of the CCM. (**A**) Pre-cultures were grown in 10% CO_2*
and diluted into two tubes, one of which was cultured in 10% CO_2 and the other in ambient air (Methods). Strains were
*tested in biological quadruplicate and culture density was measured after four days. (**B**) Targeted mutations to CCM*
components ablated growth in ambient air while permitting growth in 10% CO_2, as expected. The left bar (darker color)
gives the mean endpoint density in 10% CO_2 for each strain. The right bar (lighter color) gives the mean in ambient air.
Error bars give the standard deviation. From left to right: a positive control (grey) grew in 10% CO_2 and ambient air,
while a negative control CCMB1 strain carrying catalytically inactive rubisco (CCMB1:pCB' cbbL⁻+pCCM') failed to grow
in either condition; CCMB1 expressing rubisco and prk but no CCM genes (green, CCMB1:p1A+vec) grew only in 10%
CO_2; CCMB1:pCB'+pCCM' grew in 10% CO_2 and ambient air, recapitulating results presented in Figure 3. The following
four pairs of maroon bars give growth data for strains carrying targeted mutations to CCM genes: an inactivating
mutation to carboxysomal carbonic anhydrase (CCMB1:pCB' CsoSCA⁻+pCCM'), deletion of the CsoS2 N-terminus
responsible for recruiting rubisco to the carboxysome (CCMB1:pCB' CsoS2 ΔNTD +pCCM'), deletion of pentameric
vertex proteins (CCMB1:pCB' ΔcsoS4AB + pCCM'), and inactivating mutations to the DAB carbon uptake system
(CCMB1:pCB' DabA1⁻ + pCCM'). All four CCM mutations abrogated growth in air while permitting growth in 10% CO_2.
The positive control is the CAfree strain expressing human carbonic anhydrase II (Methods). Figure supplement 1
describes additional controls, statistical analyses, and a longer timescale replicate experiment (12 days) that
additionally tests the contribution of rubisco chaperones to the CCM. Detailed description of all plasmid and mutation
abbreviations is given in Table S2.
208
To verify that growth in ambient air depends on the CCM, we generated plasmids carrying
targeted mutations to known CCM components (Figure 4). An inactivating mutation to the
carboxysomal rubisco (CbbL K194M) prohibited growth entirely. Mutations targeting the CCM,
rather than rubisco itself, should ablate growth in ambient air while permitting growth in high CO_2
(Desmarais et al., 2019; Mangan et al., 2016; Marcus et al., 1986; Price and Badger, 1989a; Rae
et al., 2013). Consistent with this understanding, an inactive mutant of the carboxysomal carbonic
anhydrase (CsoSCA C173S) required high-CO_2 for growth. Similarly, disruption of carboxysome
formation by removal of the pentameric shell proteins or the N-terminal domain of CsoS2 also

eliminated growth in ambient air. Removing the pentameric proteins CsoS4AB disrupts the
permeability barrier at the carboxysome shell (Cai et al., 2009), while truncating CsoS2 prohibits
carboxysome formation entirely (Oltrogge et al., 2020). Finally, an inactivating mutation to the
inorganic carbon transporter also eliminated growth in ambient air (Desmarais et al., 2019).
221
222 These experiments demonstrate that pCB' and pCCM' enable CCMB1 to grow in ambient air in a
223 manner that depends on the known components of the bacterial CCM. To confirm that these cells
224 produce carboxysome structures, we performed thin section electron microscopy. Regular
225 polyhedral inclusions of ≈100 nm diameter were visible in micrographs (Figure 5A), implying
226 production of morphologically-normal carboxysomes.
227

228
Figure 5. CCMB1:pCB'+pCCM' produces morphologically normal carboxysomes and fixes CO_2 from ambient
air into biomass. (A) Polyhedral bodies resembling carboxysomes are evident in electron micrographs of
CCMB1:pCB'+pCCM' cells grown in air. Figure supplement 1 shows images of control strains. (B) Biological replicate
cultures were grown in ambient air with 99% ^{13}C glycerol as the sole organic carbon source so that $^{12}CO_2$ in air is the
sole source of ^{12}C. As serine is a direct metabolic product of 3PG, we expect ^{12}C enrichment on serine when rubisco
is active. 3PG also derives from glycolytic metabolism of glycerol, so complete ^{12}C labeling of serine was not expected.
(C) The ^{12}C composition of serine from CCMB1:pCB' + pCCM' ("Experiment") is roughly threefold above the control.
Figure supplement 2 gives ^{12}C composition of all measured amino acids. (D) The fraction of 3PG production flux due
to rubisco was predicted via Flux Balance Analysis and estimated from isotopic labeling data (Methods). Estimates of
the rubisco flux fraction exceed 10% for all four biological replicates and the mean estimate accords well with a ≈20%
prediction. Figure supplement 3 details the flux calculation procedure.
240
We next conducted isotopic labeling experiments to determine whether CCMB1:pCB' + pCCM'
fixes CO_2 from ambient air into biomass. Cells were grown in minimal media with ^{13}C-labeled
glycerol as the sole organic carbon source, such that CO_2 from ambient air was the dominant
source of ^{12}C. The isotopic composition of amino acids in total biomass hydrolysate was analyzed
via mass spectrometry (Methods). Serine is a useful sentinel of rubisco activity because *E. coli*
produces serine from the rubisco product 3PG (Stauffer, 2004; Szyperski, 1995). 3PG is also an
intermediate of lower glycolysis (Bar-Even et al., 2012), and so the degree of ^{12}C labeling on
serine reports on the balance of fluxes through rubisco and lower glycolysis (Figure 5B). We
therefore expected excess ^{12}C labeling of serine when rubisco is active. Consistent with this
expectation, serine from CCMB1:pCB'+pCCM' cells contained roughly threefold more ^{12}C than
the rubisco-independent control (Figure 5C). We estimated the contribution of rubisco to 3PG

synthesis *in vivo* by comparing labeling patterns between the rubisco-dependent experimental
cultures and controls (Methods). Based on these estimates, rubisco carboxylation was
responsible for at least 10% of 3PG synthesis in all four biological replicates (Figure 5D, Methods),
confirming fixation of CO_2 from ambient air. As such, this work represents the first functional
reconstitution of any CCM.
257
Reconstitution in *E. coli* enabled us to investigate which *H. neapolitanus* genes are necessary for
CCM function in the absence of any regulation or genetic redundancy (i.e. genes with overlapping
function) present in the native host. We focused on genes involved in rubisco proteostasis and
generated plasmids lacking acRAF, a putative rubisco chaperone, or carrying targeted mutations
to CbbQ, an ATPase involved in activating rubisco catalysis (Aigner et al., 2017; Mueller-Cajar,
2017; Wheatley et al., 2014). Although *acRAF* deletion had a large negative effect in *H.*
neapolitanus (Desmarais et al., 2019), neither acRAF nor CbbQ were strictly required for CCMB1
to grow in ambient air. Consistent with our screen in the native host (Desmarais et al., 2019),
however, *acRAF* deletion produced a substantial growth defect (Figure 4 - figure supplement 1,
panel C), suggesting that the rate of rubisco complex assembly is an important determinant of
carboxysome biogenesis.

Discussion

Today, CCMs catalyze about half of global photosynthesis (Raven et al., 2017), but this was not
always so. Land plant CCMs, for example, arose only in the last 100 million years (Flamholz and
Shih, 2020; Raven et al., 2017; Sage et al., 2012). Though all contemporary Cyanobacteria have
CCM genes, these CCMs are found in two convergently-evolved varieties (Flamholz and Shih,
2020; Kerfeld and Melnicki, 2016; Rae et al., 2013), suggesting that the ancestor of present-day
Cyanobacteria and chloroplasts did not have a CCM (Rae et al., 2013). So how did carboxysome
CCMs come to dominate the cyanobacterial phylum?
277
Here we demonstrated that the α-carboxysome CCM from *H. neapolitanus* is readily transferred
between species and confers a large growth benefit, which can explain how these CCMs became
so widespread among bacteria (Kerfeld and Melnicki, 2016; Rae et al., 2013). We constructed a
CCM by expressing 20 genes in an engineered *E. coli* strain, CCMB1. In accordance with its role
in native autotrophic hosts (Desmarais et al., 2019; Long et al., 2018; Marcus et al., 1986; Price
and Badger, 1989a), the transplanted CCM required α-carboxysomes and inorganic carbon
uptake to enable CCMB1 to grow by fixing CO_2 from ambient air (Figures 3-5, S6-8). These results
conclusively demonstrate that at most 20 gene products are required to produce a bacterial CCM.
The α-carboxysome CCM is apparently genetically compact and "portable" between organisms.
It is possible, therefore, that expressing bacterial CCMs in non-native autotrophic hosts will
improve CO_2 assimilation and growth. This is a promising approach to improving plant growth
characteristics (Ermakova et al., 2020; Long et al., 2016; Wu et al., 2019) and also engineering
enhanced microbial production of fuel, food products and commodity chemicals from CO_2
(Claassens et al., 2016; Gleizer et al., 2019).
292

Reconstitution also enabled us to test, via simple genetic experiments, whether particular genes
play a role in the CCM (Figure 4 - figure supplement 1). These experiments demonstrated that
the rubisco chaperones are strictly dispensable for producing a functional bacterial CCM, though
removing the *acRAF* gene produced a substantial growth defect that warrants further
investigation. Further such can use our reconstituted CCM to delineate a minimal reconstitution
of the bacterial CCM suitable for plant expression (Du et al., 2014; Long et al., 2018, 2016;
Occhialini et al., 2016; Orr et al., 2020), test hypotheses about carboxysome biogenesis (Bonacci
et al., 2012; Oltrogge et al., 2020), and probe the relationship between CCMs and host physiology
(Mangan et al., 2016; Price and Badger, 1989b).
302
Our approach to studying CCMs by reconstitution in tractable non-native hosts can be applied to
other CCMs, including β-carboxysome CCMs, the algal pyrenoid, and plausible evolutionary
ancestors thereof (Flamholz and Shih, 2020). Historical trends in atmospheric CO_2 likely promoted
the evolution of CCMs (Fischer et al., 2016; Flamholz and Shih, 2020), so testing the growth of
plausible ancestors of bacterial CCMs (e.g. carboxysomes lacking carbonic anhydrase activity)
may provide insight into paths of CCM evolution and the composition of the ancient atmosphere
at the time bacterial CCMs arose. In response to these same pressures, diverse eukaryotic algae
evolved CCMs relying on micron-sized rubisco aggregates called the pyrenoids (Flamholz and
Shih, 2020; Wang and Jonikas, 2020). Pyrenoid CCMs are collectively responsible for perhaps
80% of oceanic photosynthesis (Mackinder et al., 2016), yet many fundamental questions remain
regarding the composition and operation of algal CCMs (Wang and Jonikas, 2020). Functional
reconstitution of a pyrenoid CCM is a worthy goal which, once achieved, will indicate enormous
progress in our collective understanding of the genetics, cell biology, biochemistry and physical
processes supporting the eukaryotic complement of oceanic photosynthesis. We hope such
studies will further our principled understanding of, and capacity to engineer, the cell biology
supporting CO_2 fixation in diverse organisms.
319
320

Materials and Methods

Growth conditions

Unless otherwise noted, cells were grown on M9 minimal media supplemented with 0.4% w/v glycerol, 0.5 ppm thiamin (10^4 dilution of 0.5% w/v stock) and a trace element mix. The trace element mix components and their final concentrations in M9 media are: 50 mg/L EDTA, 31 mM $FeCl_3$, 6.2 mM $ZnCl_2$, 0.76 mM $CuSO_4·5H_2O$, 0.42 mM $CoCl_2·6H_2O$, 1.62 mM H_3BO_3, 81 nM $MnCl_2·4H_2O$. 100 nM anhydrotetracycline (aTc) was used in induced cultures. For routine cloning, 25 mg/L chloramphenicol and 60 mg/L kanamycin selection were used as appropriate. Antibiotics were reduced to half concentration (12.5 and 30 mg/L, respectively) for CCMB1 growth experiments and kanamycin was omitted when evaluating rubisco-dependence of growth as pF plasmids carrying kanamycin resistance also express rubisco. Culture densities were measured at 600 nm in a table top spectrophotometer (Genesys 20, Thermo Scientific) and turbid cultures were measured in five or tenfold dilution as appropriate in order to reach the linear regime of the spectrophotometer.

Agar plates were incubated at 37 °C in defined CO_2 pressures in a CO_2 controlled incubator (S41i, New Brunswick). For experiments in which a frozen bacterial stock was used to inoculate the culture, cells were first streaked on agar plates and incubated at 10% CO_2 to facilitate fast growth. Pre-cultures derived from colonies were grown in 2-5 mL liquid M9 glycerol media under 10% CO_2 with a matching 1 mL control in ambient air. Negative control strains unable to grow in minimal media (i.e. active site mutants of rubisco) were streaked on and pre-cultured in LB media under 10% CO_2.

Growth curves were obtained using two complementary methods: an 8-chamber bioreactor for large-volume cultivation (MC1000, PSI), and 96-well plates in a gas controlled plate reader plate (Spark, Tecan). For the 96-well format, cells were pre-cultured in the appropriate permissive media, M9 glycerol under 10% CO_2 where possible. If rich media was used, e.g. for negative controls, stationary phase cells were washed in 2x the culture volume and resuspended in 1x culture volume of M9 with no carbon source. Cultures were diluted to an OD of 1.0 (600 nm) and 250 µl cultures were inoculated by adding 5 µl of cells to 245 µl media. A humidity cassette (Tecan) was refilled daily with distilled water to mitigate evaporation during multi-day cultivation at 37 °C. Evaporation nonetheless produced irregular growth curves (e.g. Figure 2 - figure supplement 3), which motivated larger volume cultivation in the bioreactor, which mixes by bubbling ambient air into each growth vessel. 80 ml bioreactor cultures were inoculated to a starting OD of 0.005 (600 nm) and grown at 37 °C to saturation. Optical density was monitored continuously at 680 nm.

Anaerobic cultivation of agar plates was accomplished using a BBL GasPak 150 jar (BD) flushed 6 times with an anoxic mix of 10% CO_2 and 90% N_2. Tenfold titers of biological duplicate cultures were plated on M9 glycerol media with and without 20 mM $NaNO_3$ supplementation. Because *E. coli* cannot ferment glycerol, NO_3^- was supplied as an alternative electron acceptor. Plates without

362 NO_3^- showed no growth (Figure 2 - figure supplement 4), confirming the presence of an anaerobic
363 atmosphere in the GasPak.

Computational design of rubisco-dependent strains

To computationally design mutant strains in which growth is coupled to rubisco carboxylation flux,
we used a variant of Flux Balance Analysis (Lewis et al., 2012) called "OptSlope" (Antonovsky et
al., 2016). Starting from a published model of *E. coli* central metabolism, the Core *Escherichia*
coli Metabolic Model (Orth et al., 2010), we considered all pairs of central metabolic knockouts
and ignored those that permit growth *in silico* in the absence of rubisco and phosphoribulokinase
(Prk) activities. For the remaining knockouts, we evaluated the degree of coupling between
rubisco flux and biomass production during growth in nine carbon sources: glucose, fructose,
gluconate, ribose, succinate, xylose, glycerate, acetate and glycerol. This approach highlighted
several candidate rubisco-dependent knockout strains, including Δ*rpiAB* Δ*edd*. OptSlope
predicted rubisco-dependent growth of Δ*rpiAB* Δ*edd* strains on glucose, fructose, succinate,
acetate, glycerate, xylose and gluconate. The OptSlope algorithm is outlined in Figure 2 - figure
supplement 1 and described fully in (Antonovsky et al., 2016). Proposed mechanisms of rubisco-
dependence are outlined in Figure 2 - figure supplement 2. OptSlope source code is available at
https://gitlab.com/elad.noor/optslope and calculations specific to CCMB1 can be found at
https://github.com/flamholz/carboxecoli.

Genomic modifications producing the CCMB1 strain

Strains used in this study are documented in Table S1. To produce CCMB1, we first constructed
a strain termed "Δ*rpiAB*" for short. This strain has the genotype Δ*rpiAB* Δ*edd* and was constructed
in the *E. coli* BW25113 background by repeated rounds of P1 transduction from the KEIO
collection followed by pCP20 curing of the kanamaycin selection marker (Baba et al., 2006;
Datsenko and Wanner, 2000). Deletion of *edd* removes the Entner-Doudoroff pathway (Peekhaus
and Conway, 1998), forcing rubisco-dependent metabolism of gluconate via the pentose
phosphate pathway (Figure 2 - figure supplement 2). CCMB1 has the genotype BW25113 Δ*rpiAB*
Δ*edd* Δ*cynT* Δ*can* and was constructed from Δ*rpiAB* by deleting both native carbonic anhydrases
using the same methods, first transducing the KEIO Δ*cynT* and then Δ*can* from EDCM636 (Merlin
and Masters, 2003), which was obtained from the Yale Coli Genetic Stock Center. Transformation
was performed by electroporation (ECM 630, Harvard Biosciences) and electrocompetent stocks
were prepared using standard protocols. Strain genotypes were verified by PCR, as described
below.
394
Plants, cyanobacteria and other autotrophs uniformly express "photorespiratory" pathways to
process the rubisco oxygenation product 2-phosphoglycolate, or 2PG (Eisenhut et al., 2008). The
E. coli genome encodes enzymes that could plausibly serve as a photorespiratory pathway
(Figure 2 - figure supplement 2). We attempted to delete the *gph* gene in CCMB1 as it encodes
the 2PG phosphatase that catalyzes the first step of this putative pathway. However, the Δ*gph*
knockout was challenging to transform by electroporation, consistent with a proposed role in DNA
repair (Pellicer et al., 2003). We reasoned that photorespiration might be required in CCMB1, as

photorespiratory genes are essential in cyanobacteria (Eisenhut et al., 2008) and
chemolithoautotrophic bacteria (Desmarais et al., 2019) even though both employ CCMs.

Recombinant expression of rubisco, prk, and CCM components

pFE21 and pFA31 are compatible vectors derived from pZE21 and pZA31 (Lutz and Bujard,
1997). These vectors use an anhydrotetracycline (aTc) inducible $P_{LtetO-1}$ promoter to regulate gene
expression. pF plasmids were modified from parent vectors to constitutively express the tet
repressor (TetR) under the P_{bla} promoter so that expression is repressed by default (Liang et al.,
1999). We found that an inducible system aids in cloning problematic genes like *prk* (Wilson et
al., 2018). We refer to these vectors as pFE and pFA respectively. The p1A plasmid (Figure 2A)
derives from pFE and expresses two additional genes: the Form IA rubisco from *H. neapolitanus*
and a *prk* gene from *Synechococcus elongatus PCC 7942*. The pCB plasmid is properly called
pFE-CB, while pCCM is pFA-CCM. The two CCM plasmids are diagrammed in Figure 3 - figure
supplement 1. Cloning was performed by Gibson and Golden-Gate approaches as appropriate.
Large plasmids (e.g. pCB, pCCM) were verified by Illumina resequencing (Harvard MGH DNA
Core plasmid sequencing service) and maps were updated manually after reviewing results
compiled by breseq resequencing software (Deatherage and Barrick, 2014). Plasmids used in
this study are described in Table S2 and available on Addgene at
https://www.addgene.org/David_Savage/.

Verifying the dependence of CCMB1 on rubisco carboxylation

To verify the dependence of CCMB1 on rubisco and Prk activities in minimal media, we
constructed the variants of p1A carrying inactive rubisco or *prk* genes. Rubisco was inactivated
by mutating the large subunit active site lysine to methionine, producing p1A CbbL K194M, or
p1A CbbL⁻ for short (Andersson et al., 1989; Cleland et al., 1998). Prk was inactivated by mutating
ATP-binding residues in the Walker A motif, producing p1A Prk K20M S21A, termed p1A Prk⁻ for
short (Cai et al., 2014; Higgins et al., 1986). CCMB1:p1A grew on glycerol and gluconate minimal
media when provided 10% CO_2 (Figure 2 - figure supplement 3). CCMB1:p1A CbbL⁻ and
CCMB1:p1A Prk⁻ both failed to grow on minimal media supplemented with glycerol or gluconate,
demonstrating a dependence on both enzymes. So long as high CO_2 was provided, neither
activity was required for growth in rich LB media, which contains abundant nucleic acids
precursors (Sezonov et al., 2007). Xylose minimal media was also tested but growth was
impractically slow (data not shown).
433
The high-CO_2 requirement of CCMB1:p1A growth was expected for two reasons: (i) bacterial
rubiscos typically display low net carboxylation rates in ambient air due to relatively low CO_2
(≈0.04%) and relatively high O_2 (≈21%), as shown in Figure 1B and discussed in (Flamholz et al.,
2019; Iñiguez et al., 2020), and (ii) CCMB1 entirely lacks carbonic anhydrase activity (Δ*cynT*
Δ*can*). Carbonic anhydrase knockouts of many microbes, including *E. coli* and *S. cerevisiae*, are
high-CO_2 requiring, likely due to cellular demand for HCO_3^- (Aguilera et al., 2005; Desmarais et
al., 2019; Du et al., 2014; Merlin and Masters, 2003).

To verify that CCMB1 growth depends specifically on rubisco carboxylation and not oxygenation,
we grew CCMB1:p1A on glycerol minimal medium in anoxic high-CO_2 conditions (10:90 CO_2:N_2,
Figure 2 - figure supplement 4). *E. coli* predominantly respires glycerol and, therefore, grows
extremely slowly on glycerol in anaerobic and low O_2 conditions (Stolper et al., 2010). We
therefore supplied 20 mM NO_3^- as an alternate terminal electron acceptor (Unden and Dünnwald,
2008) in anaerobic growth conditions (see "Growth conditions"). CCMB1:p1A grew on glycerol
media in anaerobic conditions when NO_3^- was provided. Growth is qualitatively weaker than a
wild-type control, but this is consistent with the growth differences observed in aerobic conditions
(Figure 2 - figure supplement 4). Anaerobic growth of CCMB1:p1A on glycerol minimal media
implies that growth can be supported by rubisco carboxylation alone and does not require the
rubisco-catalyzed oxygenation of RuBP.

453 Strain verification by PCR and phenotypic testing

As CCMB1 is a relatively slow-growing knockout strain, we occasionally observed contaminants
in growth experiments. We used two strategies to detect contamination by faster-growing
organisms (e.g. wild-type *E. coli*). As most strains grew poorly or not at all in ambient air, pre-
cultures grown in 10% CO_2 were accompanied by a matching 1 mL negative control in ambient
air. Pre-cultures showing growth in the negative control were discarded or verified by PCR
genotyping in cases where air-growth was plausible.

PCR genotyping was performed using primer sets documented in Table S3. Three primer pairs
were used to probe a control locus (*zwf*) and two target loci (*cynT* and *rpiA*). The *zwf* locus is
intact in all strains. *cynT* and *rpiA* probes test for the presence of the CCMB1 strain (genotype
BW25113 Δ*rpiAB* Δ*edd* Δ*cynT* Δ*can*). Notably, the CAfree strain (BW25113 Δ*cynT* Δ*can*) that we
previously used to test the activity of DAB-type transporters (Desmarais et al., 2019) is a *cynT*
knockout but has a wild-type *rpiA* locus, so this primer set can distinguish between wild-type,
CAfree and CCMB1. This was useful for some experiments where CAfree was used as a control
(e.g. Figures S7-8). Pooled colony PCRs were performed using Q5 polymerase (NEB), annealing
at 65 °C and with a 50 second extension time.

470 Selection for growth in novel conditions

CCMB1:pCB did not initially grow in glycerol minimal media, which was unexpected because pCB
carries rubisco and *prk* genes. We therefore performed a series of selection experiments (Herz
et al., 2017) to isolate plasmids conferring growth at elevated CO_2 and then in ambient air. We
first describe the methodology; the full series of experiments is diagrammed fully in Figure S5B
and described in paragraphs below. CCMB1 cultures carrying appropriate plasmids were first
grown to saturation in rich LB media in a 10% CO_2 incubator. Stationary phase cultures were
pelleted by centrifugation for 10 min at 4,000 x g, washed in 2x the culture volume, and
resuspended in 1x culture volume of M9 media with no carbon source. After resuspension,
multiple dilutions were plated on selective media (e.g. M9 glycerol media) and incubated in the
desired conditions (e.g. in ambient air) with a positive control in 10% CO_2 on appropriate media.

In later experiments, matching tenfold titers were plated in permissive conditions (e.g. in 10%
CO_2) to estimate the number of viable cells. When colonies formed in restrictive conditions, they
were picked into permissive media, grown to saturation, washed and tested for re-growth in
restrictive conditions by titer plating or streaking. Plasmid DNA was isolated from verified colonies
and transformed into naive CCMB1 cells to test whether plasmid mutations confer improved
growth (i.e. in the absence of genomic mutations).
487
We first selected for CCMB1:pCB growth on M9 glycerol media in 10% CO_2 and then in M9
gluconate media under 10% CO_2. Pre-cultures to stationary phase in rich media in 10% CO_2, and
then plated on selective media after washing. The resulting plasmid, pCB-gg for "gluconate
grower," was isolated and deep sequenced (Harvard MGH DNA Core plasmid sequencing
service). Plasmid maps were updated manually after running the breseq resequencing software
(Deatherage and Barrick, 2014). pCB-gg was found to carry two regulatory mutations: an amino
acid substitution to the tet repressor (TetR E37A) and a nucleotide substitution in the Tet operator
regulating the carboxysome operon (tetO$_2$ +8T, Table S4).
496
Following this first round of selection, CCMB1 was co-transformed with pCB-gg and pCCM. The
transformants grew in M9 glycerol media in 10% CO_2 but failed to grow on in ambient air. We
therefore performed another selection experiment, plating CCMB1:pCB-gg+pCCM on M9 glycerol
media in ambient CO_2. Parallel negative control selections were conducted on uninduced plates
(no aTc) and using CCMB1:p1A+pCCM, which lacks carboxysome genes.   Colonies formed on
induced CCMB1:pCB-gg+pCCM plates after 20 days, but not on control plates (Figure S5F).
503
Forty colonies were picked and tested for re-growth in ambient CO_2 by tenfold titer plating. 10 of
40 regrew (six examples are shown in Figure S5G). Pooled plasmid DNA was extracted from
verified colonies and electroporated into naive CCMB1 to test plasmid-linkage of growth. We
found that plasmid DNA from colony #4 produced the most robust growth in ambient air. This was
tested by picking 16 re-transformants and testing their growth in ambient air in liquid M9 glycerol
media. Re-transformant #13 regrew robustly in all 6 technical replicates. Pooled plasmid DNA
from colony #4 re-transformant #13 was resequenced by a combination deep sequencing (as
above) and targeted Sanger sequencing of the TetR locus and origins of replication, as these
regions share sequence between both parent plasmids. pCB carried the same mutations as pCB-
gg and pCCM had acquired the high-copy ColE1 origin of replication from pCB (Table S4). The
individual mutant plasmids, termed pCB' and pCCM', were reconstructed from pooled plasmid
extract by PCR and Gibson cloning.
516
517   These post-selection plasmids, termed pCB' and pCCM', were again verified by resequencing.
518   Naive CCMB1 was transformed with the reconstructed post-selection plasmids pCB' and pCCM'
519   and tested for growth in ambient air. We found that the post-selection plasmids confer
520   reproducible growth in ambient air in multiple growth conditions (Figure 3), implying that genomic
521   mutations that formed during selections were not required to produce growth in ambient air.

Design of mutant CCM plasmids

To verify that air-growth depends on the known components of the CCM, we generated variants of pCB' and pCCM' carrying known, targeted null mutations to the CCM. CCMB1 was co-transformed with two plasmids: a mutant plasmid (of either pCB' or pCCM') and its cognate, unmodified plasmid. Mutant plasmids are listed here along with expected growth phenotypes, with fuller detail in Table S2. pCB' CbbL K194M, or pCB'⁻, contains an inactivating mutation to the large subunit of the carboxysomal Form 1A rubisco (Andersson et al., 1989; Cleland et al., 1998). This mutation was expected to abrogate rubisco-dependent growth entirely.

Mutations targeting the CCM, rather than rubisco itself, are expected to ablate growth in ambient air but permit growth in high CO_2. The following plasmid mutations were designed to specifically target essential components of the CCM. pCB' CsoSCA C173S, or pCB' CsoSCA⁻, carries a mutation to an active site cysteine residue responsible for coordinating the catalytic Zn2+ ion in **β**-carbonic anhydrases (Sawaya et al., 2006). pCB' CsoS2 ΔNTD lacks the N-terminal domain of CsoS2, which is responsible for recruiting rubisco to the carboxysome during the biogenesis of the organelle (Oltrogge et al., 2020). Similarly, pCB' CbbL Y72R carries an arginine residue instead of the tyrosine responsible for mediating cation-π interactions between the rubisco large subunit and the N-termus of CsoS2. This mutation was shown to eliminate any binding interaction between the rubisco complex and the N-termus of CsoS2 (Oltrogge et al., 2020). pCB' Δ*csoS4AB* lacks both pentameric shell proteins, CsoS4AB, which was shown to disrupt the permeability barrier at the carboxysome shell (Cai et al., 2009). pCCM' DabA1 C462A, D464A, or pCCM' DabA1⁻, carries inactivating mutations to the putative active site of the inorganic carbon transporter component DabA1 (Desmarais et al., 2019).

Two more mutant plasmids were designed to test the roles of rubisco chaperones in producing a functional CCM. pCCM' CbbQ K46A, E107Q, denoted pCCM' CbbQ⁻, carries mutations that inactivate the ATPase activity of the CbbQ subunit of the CbbOQ rubisco activase complex (Tsai et al., 2015). pCCM' Δ*acRAF* lacks the putative rubisco chaperone acRAF. acRAF is homologous to a plant rubisco folding chaperone (Aigner et al., 2017) and likely involved in the folding of the *H. neapolitanus* Form IA rubisco (Wheatley et al., 2014). Experimental evaluation of growth phenotypes for the above-described mutants is detailed below and results are given in Figure 4 - figure supplement 1.

Phenotyping of matched cultures in 10% CO_2 and ambient air

To interrogate the phenotypic effects of mutations to the CCM, we tested the growth of matched biological replicate cultures of CCM mutants (e.g. disruption of carboxysome components or transporter function) in M9 glycerol medium in 10% CO_2 and ambient air (Figure 4A). For these experiments, individual colonies were picked into a round-bottom tube with 4 mL of M9 glycerol media with full strength antibiotic and 100 nM aTc. 1 mL of culture was then transferred to a second tube. The 3 mL pre-culture was incubated in 10% CO_2, while the 1 mL culture was incubated in ambient air as a negative control. Control strains unable to grow in minimal media (e.g. those expressing inactive rubisco mutants) were pre-cultured in LB media. High-CO_2 pre-

563   cultures were grown to saturation, after which optical density (OD600) was measured in five-fold
564   dilution.
565

Experimental cultures were inoculated with pre-culture to a starting OD600 of 0.01 in 3 mL of M9
glycerol media with 12.5 mg/L chloramphenicol and 100 nM aTc. Each pre-culture was used to
inoculate a matched pair of experimental cultures, one incubated in 10% CO_2 and another in
ambient air, as diagrammed in Figure 4A. After a defined period of growth (4 days in Figure 4B
and 12 days in Figure 4 - figure supplement 1, panel C) all culture densities were measured at
600 nm. All experiments were performed in biological quadruplicate, i.e. using four independent
pre-cultures deriving from distinct colonies to inoculate four pairs of matched cultures. A positive
control was included in all experiments to test the media composition. We used a complemented
double carbonic anhydrase knockout (CAfree:pFE-sfGFP+pFA-HCAII) for this purpose as its
growth in air depends on the expression of the human carbonic anhydrase II from the pFA-HCAII
plasmid (Desmarais et al., 2019).

577 Electron microscopy

CCMB1:pCB'+pCCM' was grown in ambient air in 3 ml of M9 glycerol medium and induced with
100 nM aTc. A carboxysome-negative control, CAfree:pFE-sfGFP+pFA-HCAII, was grown in the
same conditions. Sample preparation and sectioning were performed by the University of
California Berkeley Electron Microscope Laboratory. Cell pellets were fixed for 30 min at room
temperature in 2.5% glutaraldehyde in 0.1 M cacodylate buffer pH 7.4. Fixed cells were stabilized
in 1% very low melting-point agarose and cut into small cubes. Cubed sample was then rinsed
three times at room temperature for 10 min in 0.1 M sodium cacodylate buffer, pH 7.4 and then
immersed in 1% osmium tetroxide with 1.6% potassium ferricyanide in 0.1 M cacodylate buffer
for an hour in the dark on a rocker. Samples were later rinsed three times with a cacodylate buffer
and then subjected to an ascending series of acetone for 10 min each (35%, 50%, 75%, 80%,
90%, 100%, 100%). Samples were progressively infiltrated with Epon resin (EMS, Hatfield, PA,
USA) while rocking and later polymerized at 60 °C for 24 hours. 70 nm thin sections were cut
using an Ultracut E (Leica) and collected on 100 mesh formvar coated copper grids. The grids
were further stained for 5 min with 2% aqueous uranyl acetate and 4 min with Reynold's lead
citrate. The sections were imaged using a Tecnai 12 TEM at 120 KV (FEI) and images were
collected using UltraScan 1000 digital micrograph software (Gatan Inc.).

594 Sample preparation for LC-MS analysis

Protein-bound amino acids were analyzed in total biomass hydrolysate of 80 mL cultures grown
in minimal media with 99% ^{13}C glycerol (Cambridge Isotopes) as the sole organic carbon source.
These cultures were grown in 80 mL volumes in a bioreactor pumping ambient air (MC1000, PSI).
After harvesting biomass, samples were prepared and analyzed as described in (Antonovsky et
al., 2016). Briefly, the OD600 was recorded and 0.6 OD x mL of sample were pelleted by
centrifugation for 15 min at 4,000 x g. The pellet was resuspended in 1 mL of 6 N HCl and
incubated for 24 hours at 110 °C. The acid was subsequently evaporated under a nitrogen stream
using a custom-built gas manifold (Nevins et al., 2005), resulting in a dry hydrolysate. Dry

hydrolysates were resuspended in 0.6 mL of MilliQ water, centrifuged for 5 min at 14,000 x g, and
supernatant was analyzed by liquid chromatography-mass spectrometry (LC-MS). Hydrolyzed
amino acids were separated using ultra performance liquid chromatography (UPLC, Acquity,
Waters) on a C-8 column (Zorbax Eclipse XBD, Agilent) at a flow rate of 0.6 mL/min, and eluted
off the column using a hydrophobicity gradient. Buffers used were: A) H2O + 0.1% formic acid
and B) acetonitrile + 0.1% formic acid with the following gradient: 100% of A (0-3 min), 100% A
to 100% B (3-9 min), 100% B (9-13 min), 100% B to 100% A (13-14 min), 100% A (14-20 min).
The UPLC was coupled online to a triple quadrupole mass spectrometer (TQS, Waters). Data
were acquired using MassLynx v4.1 (Waters). Amino acids and metabolites used for analysis
were selected according to the following criteria: amino acids that had peaks at a distinct retention
time and m/z values for all isotopologues and also showed correct ^{13}C labeling fractions in control
samples that contained protein hydrolyzates of WT cells grown with known ratios of ^{13}C6-glucose
to ^{12}C-glucose.

Isotopic analysis composition of biomolecules

617   The total ^{13}C fraction of each metabolite was determined as the weighted average of the fractions
618   of all the isotopologues for that metabolite:
619

$$f_{13_C} = \frac{\sum_{i=0}^{N} f_i \times i}{N}$$

621
622   Here N is the number of carbons in the compound (e.g. N = 3 for serine) and f_i is the relative
623   fraction of the i-th isotopologue, i.e containing i ^{13}C carbon atoms. Each metabolite's total ^{12}C
624   fraction was calculated as $f_{12_C} = 1 - f_{13_C}$.

Estimating the effective intracellular ^{12}CO$_2$ fraction

E. coli cells grown in ^{13}C glycerol will simultaneously respire glycerol, producing intracellular
^{13}CO$_2$, and take up extracellular ^{12}CO$_2$ and H^{12}CO$_3^-$. The isotopic composition of the intracellular
inorganic carbon (C$_i$) pool will therefore reflect the balance of uptake and respiration. As rubisco
carboxylation draws from the intracellular CO$_2$ pool, we must estimate the isotopic composition of
the C$_i$ pool to evaluate the contribution of rubisco to metabolism. We used the carbamoyl-
phosphate moiety as a marker for the isotopic distribution of the intracellular C$_i$ pool, as described
in (Gleizer et al., 2019). Briefly, carbamoyl-phosphate is generated by phosphorylation of
bicarbonate, and is condensed with ornithine in the biosynthesis of L-arginine. We compared the
mass isotopologue distribution of L-arginine, which contains one carbon from carbamoyl-
phosphate, with the mass isotopologue distribution of L-glutamate as L-glutamate is an ornithine
precursor.
637
638   We estimated the effective ^{13}C labeling of intracellular inorganic carbon ($f_{13_{CO_2},effective}$) as follows:

$$f_{13_{CO_2},effective} = \sum_{i=0}^{6} f_{arg,i} - \sum_{i=0}^{5} f_{glu,i}$$

Here $f_{13_{CO_2},effective}$ is the relative fraction of $^{13}CO_2$ out of the total CO_2 pool (or, more formally,
the C_i), and $f_{arg,i}$ and $f_{glu,i}$ are the fraction of the i-th isotopologue of arginine and glutamate
respectively. We assumed fast equilibration of the intracellular C_i pool because the strains used
in labeling experiments express a carbonic anhydrase. An equivalent equation can be defined for
the arginine-proline comparison (Gleizer et al., 2019), however proline data were of insufficient
quality to use and so we report inferences based on the arginine-glutamate comparison. The
effective intracellular fraction of $^{12}CO_2$ was calculated as $f_{12_{CO_2},effective} = 1 - f_{13_{CO_2},effective}$.

For brevity, we refer to these fractions as $f_{12_{CO_2}}$ and $f_{13_{CO_2}}$, respectively.

Estimating the rubisco carboxylation flux *in vivo*

When CCMB1 cells are grown on 99% ^{13}C glycerol, 3-phosphoglycerate (3PG) can be produced
via two routes: (i) rubisco catalyzed carboxylation of RuBP and (ii) glycolytic metabolism of
glycerol via dihydroxyacetone phosphate, or DHAP (Booth, 2005). We denote these two fluxes
as $J_{rubisco}$ and J_{pgk}, where pgk (phosphoglycerate kinase) is the glycolytic enzyme producing 3PG
(Bar-Even et al., 2012). Serine is a direct metabolic product of 3PG (Stauffer, 2004; Szyperski,
1995) and was therefore assumed to have the same ^{12}C composition as 3PG. Rubisco-catalyzed
carboxylation of RuBP adds one CO_2 to the 5-carbon substrate, producing two 3PG molecules
containing a total of six carbon atoms. Therefore, ⅙ of carbon atoms on 3PG produced via rubisco
carboxylation must derive from an inorganic source (Figure 5 - figure supplement 3).
Carboxylation draws CO_2 from the intracellular inorganic carbon pool, whose ^{12}C composition
$f_{12_{CO_2}}$ was inferred as described above.

Based on these assumptions, the ^{12}C composition of 3PG, and therefore serine, equals a flux-
weighted sum of contributions from rubisco and pgk. As such, the relative 3PG production flux
that is due to rubisco, $J_{rubisco}/(J_{rubisco}+J_{pgk})$, can be inferred via the following calculation:
665

666
$$f_{ser,ctrl} = f_{3PG,ctrl} = 0 \times \frac{1}{6}(f_{12_{CO_2}} + 5 \times f_{RuBP,exp}) + 1 \times f_{DHAP,ctrl} = f_{DHAP,ctrl}$$

667
$$f_{ser,exp} = f_{3PG,ctrl} = \frac{J_{rubisco}}{J_{rubisco} + J_{pgk}} \times \frac{1}{6}(f_{12_{CO_2}} + 5 \times f_{RuBP,exp}) + \frac{J_{pgk}}{J_{rubisco} + J_{pgk}} \times f_{DHAP,exp}$$

669 Where the first equation is written for the control and the second for experimental cultures where
670 rubisco is active (CCMB1:pCB'+pCCM'). $f_{ser,ctrl}$ and $f_{ser,exp}$ denote the ^{12}C composition of serine
671 in the control and experiment respectively. Identical notation is used for RuBP and DHAP. As
672 there are only two routes of 3PG production, the above equations can be simplified to solve for
673 the relative flux through rubisco *in vivo*:
674

675
$$\frac{J_{pgk}}{J_{rubisco} + J_{pgk}} \equiv 1 - \frac{J_{rubisco}}{J_{rubisco} + J_{pgk}}$$

676
$$\frac{J_{rubisco}}{J_{rubisco} + J_{pgk}} = \frac{f_{ser,exp} - f_{DHAP,exp}}{\frac{1}{6}(f_{12_{CO_2}} + 5 \times f_{RUBP,exp}) - f_{DHAP,exp}}$$

To calculate the rubisco flux *in vivo* we must attach values to several parameters in the above
equation. $f_{12_{CO_2}}$ was inferred on a per-sample basis, with the mean values being $25\% \pm 4\%$ and
$67\% \pm 28\%$ for the control and experiment respectively (Figure 5 - figure supplement 3, panel C).
Because glycerol is converted into 3PG and serine via DHAP in wild-type *E. coli* (Booth, 2005),
we expect that $f_{ser,ctrl} = f_{DHAP,ctrl}$, as derived above. LC-MS measurements give $f_{ser,ctrl} =$
$0.6\% \pm 0.2\%$ and $f_{ser,exp} = 3.5\% \pm 2.2\%$ (Figure 5C). Valine is also a metabolic product of DHAP
(Szyperski, 1995) and was found to have a similar ^{12}C fraction $f_{val,ctrl} = 0.6\% \pm 0.1\%$ in control
cells (Figure 5 - figure supplement 2). Since glycerol is immediately converted to DHAP in *E. coli*,
we further assumed that $f_{DHAP,ctrl} = f_{DHAP,exp}$.

RuBP is produced in CCMB1 when rubisco and prk are expressed. Since glycerol is the sole
carbon source and there are no carboxylation reactions between DHAP and RuBP in CCMB1, we
assumed $f_{RuBP,exp} = f_{DHAP,ctrl}$. This assumption is supported by LC-MS measurements of
histidine in control cells. Like RuBP, histidine is synthesized from a pentose-phosphate pathway
intermediates (Szyperski, 1995; Winkler and Ramos-Montañez, 2009), and measured $f_{his,ctrl} =$
$0.7\% \pm 0.1\%$, which is very similar to $f_{ser,ctrl} = 0.6\% \pm 0.3\%$. Using mean values to illustrate the
calculation gives $\frac{J_{rubisco}}{J_{rubisco} + J_{pgk}} = \frac{3.5\% - 0.6\%}{\frac{1}{6}(67\% + 5 \times 0.6\%) - 0.6\%} = 0.26$, implying that 26% of 3PG production is
due to rubisco.

10^5 random samples were drawn from the experimentally determined parameter ranges to
estimate a 99% confidence interval on the rubisco flux fraction. As the ^{12}C composition of
inorganic carbon ($f_{12_{CO_2}}$) and serine are mechanistically linked via rubisco, these values were
assumed to co-vary. Distributions were estimated on a per-sample basis by assuming 0.1% error
in direct measurement of serine and 1% error in the inference of $f_{12_{CO_2}}$. These calculations gave
a median flux estimate of 19% with 99% of values falling between 4.0% and 47.3%. The sample
with the lowest inferred rubisco flux had a median estimate of 10.2% with 99% of values falling
between 2.5% and 17.9%, implying that rubisco is responsible for a nonzero fraction of 3PG
production in all samples. Applying a wider error range of 5% to $f_{12_{CO_2}}$ did not qualitatively change
results, giving an overall median flux fraction of 19.1% and a 99% confidence interval 3.9-50.7%.
This and above calculations can be found in the following Jupyter notebook:
https://github.com/flamholz/carboxecoli/blob/master/notebooks/00_LCMS_calcs.ipynb.

Predicting rubisco carboxylation flux via Flux Balance Analysis

A stoichiometric model of complemented CCMB1 was generated from the Core *Escherichia coli*
Metabolic Model (Orth et al., 2010) by adding rubisco and prk and then deleting the rpi and edd
reactions. Parsimonious Flux Balance Analysis (pFBA) was applied to the resulting model to
calculate intracellular metabolic metabolic fluxes that maximize the rate of biomass production.

As many distinct flux distributions can yield the same (maximal) rate of biomass production, pFBA
uses the minimum sum of fluxes objective to define a unique flux solution (Holzhütter, 2004). The
COBRApy implementation of pFBA introduces an additional free parameter, the permissible
fraction of the maximal biomass production rate f_{opt} (Ebrahim et al., 2013). When f_{opt} < 1.0, the
biomass production can be less-than-optimal if this would further decrease the sum of fluxes.
719
pFBA was run with f_{opt} ranging from 0.8 to 1.0 in increments of 0.01 to account for the fact that
CCMB1 has not undergone selection to maximize biomass production with rubisco expressed.
For each resulting flux distribution the fraction of 3PG production flux due to rubisco was
calculated as the fraction of 3PG molecules produced via rubisco carboxylation divided by the
total flux to 3PG. These calculations predict that 19.5%-21.5% of 3PG production is due to
rubisco. The model was rerun after removing all possibility for product secretion by deleting all
carbon-containing exchange reactions other than glycerol and CO_2 exchange. This modification
should give an upper bound on the fraction of 3PG production due to rubisco, as carbon cannot
be shunted away from biomass production to overflow products. The "no overflow" model
predicted that 23.9% of 3PG production is due to rubisco independent of f_{opt}. The range of
predictions from 19.5-23.9% is plotted in Figure 5D. All calculations were done using Python and
COBRApy (Ebrahim et al., 2013), and can be found in this Jupyter notebook:
https://github.com/flamholz/carboxecoli/blob/master/notebooks/01_FBA_rubisco_flux_prediction
.ipynb.
734
735

Acknowledgements: We thank Matt Davis for P1 transduction materials and advice, Hernan Garcia and Han Lim for pZ plasmids, Maggie Stoeva, Anna Engelbrektson, Anchal Mehra, Sophia Ewens and Tyler Barnum for help with anaerobic growth, Reena Zalpuri and Danielle Jorgens at the University of California Berkeley Electron Microscope Laboratory for advice and assistance with electron microscopy, and Rob Egbert and Adam Arkin for KEIO strains. We are grateful to Eric Estrin, Woody Fischer, Darcy McRose, Dipti Nayak, Sabeeha Merchant, Luke Oltrogge, and Naiya Phillips for detailed comments on the manuscript, and to Dan Arlow, Yinon Bar-On, Dan Davidi, Jack Desmarais, Hernan Garcia, Oliver Mueller-Cajar, Rob Nichols, Kris Niyogi, Dan Portnoy, Morgan Price, Noam Prywes, Jeremy Roop, Rachel Shipps, Patrick Shih, and Dan Tawfik, for support, advice and helpful discussions throughout.

Funding: This work was supported by a National Science Foundation Graduate Research Fellowship (to A.I.F.), grants from the US Department of Energy (no. DE-SC00016240) and Royal Dutch Shell (Energy Biosciences Institute project CW163755) to D.F.S., and from the European Research Council (Project NOVCARBFIX 646827) to R.M. R.M. is the Charles and Louise Gartner Professional Chair.

Author contributions: A.I.F. conceived of and designed all experiments with mentorship from R.M. and D.F.S. and support from all authors. S.A., N.A., E.N., A.B-E. and R.M. designed and constructed the Δ*rpiAB* strain from which A.I.F., E.J.D, and S.R. constructed CCMB1. A.I.F., E.J.D, and S.R. designed and constructed all other strains and plasmids. A.I.F and E.J.D. performed growth and selection experiments. C.B. performed electron microscopy. S.G. and R.B-N. performed LC-MS analysis on biomass hydrolysate prepared by A.I.F. and E.J.D. A.I.F., S.G., R.B-N., and E.N. analysed isotopic labeling data. A.I.F and E.N. designed and executed Flux Balance Analysis. A.I.F. wrote the manuscript with input from all authors.

Competing interests: D.F.S. is a co-founder of Scribe Therapeutics and a scientific advisory board member of Scribe Therapeutics and Mammoth Biosciences. A.B.-E. is co-founder of b.fab. These companies were not involved in this research in any way. All other authors declare no competing interests.

Data and materials availability: All data associated with this work is available in the main text, supplementary materials and the repository at https://github.com/flamholz/carboxecoli. Plasmids available on Addgene at https://www.addgene.org/David_Savage/, strains distributed on request.

References

Aguilera J, Van Dijken JP, De Winde JH, Pronk JT. 2005. Carbonic anhydrase (Nce103p): an essential biosynthetic enzyme for growth of Saccharomyces cerevisiae at atmospheric carbon dioxide pressure. *Biochem J* **391**:311–316.

Aigner H, Wilson RH, Bracher A, Calisse L, Bhat JY, Hartl FU, Hayer-Hartl M. 2017. Plant RuBisCo assembly in E. coli with five chloroplast chaperones including BSD2. *Science* **358**:1272–1278.

Andersson I, Knight S, Schneider G, Lindqvist Y, Lundqvist T, Brändén C-I, Lorimer GH. 1989. Crystal structure of the active site of ribulose-bisphosphate carboxylase. *Nature* **337**:229–234.

Antonovsky N, Gleizer S, Noor E, Zohar Y, Herz E, Barenholz U, Zelcbuch L, Amram S, Wides A, Tepper N, Davidi D, Bar-On Y, Bareia T, Wernick DG, Shani I, Malitsky S, Jona G, Bar-Even A, Milo R. 2016. Sugar Synthesis from CO2 in Escherichia coli. *Cell* **166**:115–125.

Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko K a., Tomita M, Wanner BL, Mori H. 2006. Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol Syst Biol* **2**:2006.0008.

Bar-Even A, Flamholz A, Noor E, Milo R. 2012. Rethinking glycolysis: on the biochemical logic of metabolic pathways. *Nat Chem Biol* **8**:509–517.

Bar-On YM, Milo R. 2019. The global mass and average rate of rubisco. *Proc Natl Acad Sci U S A* **116**:4738–4743.

Bonacci W, Teng PK, Afonso B, Niederholtmeyer H, Grob P, Silver P a., Savage DF. 2012. Modularity of a carbon-fixing protein organelle. *Proc Natl Acad Sci U S A* **109**:478–483.

Booth IR. 2005. Glycerol and Methylglyoxal Metabolism. *EcoSal Plus* **1**:1–8.

Bowes G, Ogren WL. 1972. Oxygen inhibition and other properties of soybean ribulose 1,5-diphosphate carboxylase. *J Biol Chem* **247**:2171–2176.

Bremer H, Dennis PP. 2008. Modulation of Chemical Composition and Other Parameters of the Cell at Different Exponential Growth Rates. *EcoSal Plus* **3**:1–49.

Cai F, Menon BB, Cannon GC, Curry KJ, Shively JM, Heinhorst S. 2009. The pentameric vertex proteins are necessary for the icosahedral carboxysome shell to function as a CO2 leakage barrier. *PLoS One* **4**:e7521.

Cai Z, Liu G, Zhang J, Li Y. 2014. Development of an activity-directed selection system enabled significant improvement of the carboxylation efficiency of Rubisco. *Protein Cell* 12–18.

Claassens NJ, Sousa DZ, Dos Santos VAPM, de Vos WM, van der Oost J. 2016. Harnessing the power of microbial autotrophy. *Nat Rev Microbiol* **14**:692–706.

Cleland WW, Andrews TJ, Gutteridge S, Hartman FC, Lorimer GH. 1998. Mechanism of Rubisco: The Carbamate as General Base. *Chem Rev* **98**:549–562.

Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**:6640–6645.

Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved microbes from next-generation sequencing data using breseq. *Methods Mol Biol* **1151**:165–188.

Desmarais JJ, Flamholz AI, Blikstad C, Dugan EJ, Laughlin TG, Oltrogge LM, Chen AW, Wetmore K, Diamond S, Wang JY, Savage DF. 2019. DABs are inorganic carbon pumps found throughout prokaryotic phyla. *Nat Microbiol* **4**:2204–2215.

Du J, Förster B, Rourke L, Howitt SM, Price GD. 2014. Characterisation of Cyanobacterial Bicarbonate Transporters in E. coli Shows that SbtA Homologs Are Functional in This Heterologous Expression System. *PLoS One* **9**:e115905.

Ebrahim A, Lerman JA, Palsson BO, Hyduke DR. 2013. COBRApy: COnstraints-Based Reconstruction and Analysis for Python. *BMC Syst Biol* **7**:74.

Eisenhut M, Ruth W, Haimovich M, Bauwe H, Kaplan A, Hagemann M. 2008. The

photorespiratory glycolate metabolism is essential for cyanobacteria and might have been
conveyed endosymbiontically to plants. *Proc Natl Acad Sci U S A* **105**:17199–17204.
Ermakova M, Danila FR, Furbank RT, von Caemmerer S. 2020. On the road to C4 rice:
advances and perspectives. *Plant J* **101**:940–950.
Fischer WW, Hemp J, Johnson JE. 2016. Evolution of Oxygenic Photosynthesis. *Annu Rev*
Earth Planet Sci **44**:647–683.
Flamholz AI, Prywes N, Moran U, Davidi D, Bar-On YM, Oltrogge LM, Alves R, Savage D, Milo
R. 2019. Revisiting Trade-offs between Rubisco Kinetic Parameters. *Biochemistry*
58:3365–3376.
Flamholz A, Shih PM. 2020. Cell biology of photosynthesis over geologic time. *Curr Biol*
30:R490–R494.
Gleizer S, Ben-Nissan R, Bar-On YM, Antonovsky N, Noor E, Zohar Y, Jona G, Krieger E,
Shamshoum M, Bar-Even A, Milo R. 2019. Conversion of Escherichia coli to Generate All
Biomass Carbon from CO2. *Cell* **179**:1255–1263.e12.
Herz E, Antonovsky N, Bar-On Y, Davidi D, Gleizer S, Prywes N, Noda-Garcia L, Frisch KL,
Zohar Y, Wernick DG, Others. 2017. The genetic basis for the adaptation of E. coli to sugar
synthesis from CO 2. *Nat Commun* **8**:1705.
Higgins CF, Hiles ID, Salmond GP, Gill DR, Downie JA, Evans IJ, Holland IB, Gray L, Buckel
SD, Bell AW. 1986. A family of related ATP-binding subunits coupled to many distinct
biological processes in bacteria. *Nature* **323**:448–450.
Holzhütter H-G. 2004. The principle of flux minimization and its application to estimate
stationary fluxes in metabolic networks. *Eur J Biochem* **271**:2905–2922.
Iñiguez C, Capó-Bauçà S, Niinemets Ü, Stoll H, Aguiló-Nicolau P, Galmés J. 2020. Evolutionary
trends in RuBisCO kinetics and their co-evolution with CO2 concentrating mechanisms.
Plant J **101**:897–918.
Kerfeld CA, Melnicki MR. 2016. Assembly, function and evolution of cyanobacterial
carboxysomes. *Curr Opin Plant Biol* **31**:66–75.
Lewis NE, Nagarajan H, Palsson BO. 2012. Constraining the metabolic genotype–phenotype
relationship using a phylogeny of in silico methods. *Nat Rev Microbiol* **10**:291–305.
Liang S, Bipatnath M, Xu Y, Chen S, Dennis P, Ehrenberg M, Bremer H. 1999. Activities of
constitutive promoters in Escherichia coli. *J Mol Biol* **292**:19–37.
Lin MT, Occhialini A, Andralojc PJ, Parry MAJ, Hanson MR. 2014. A faster Rubisco with
potential to increase photosynthesis in crops. *Nature* **513**:547–550.
Long BM, Hee WY, Sharwood RE, Rae BD, Kaines S, Lim Y-L, Nguyen ND, Massey B, Bala S,
von Caemmerer S, Badger MR, Price GD. 2018. Carboxysome encapsulation of the CO2-
fixing enzyme Rubisco in tobacco chloroplasts. *Nat Commun* **9**:3570.
Long BM, Rae BD, Rolland V, Förster B, Price GD. 2016. Cyanobacterial CO2-concentrating
mechanism components: function and prospects for plant metabolic engineering. *Curr Opin*
Plant Biol **31**:1–8.
Lutz R, Bujard H. 1997. Independent and tight regulation of transcriptional units in Escherichia
coli via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. *Nucleic Acids Res*
25:1203–1210.
Mackinder LCM, Meyer MT, Mettler-Altmann T, Chen VK, Mitchell MC, Caspari O, Freeman
Rosenzweig ES, Pallesen L, Reeves G, Itakura A, Roth R, Sommer F, Geimer S, Mühlhaus
T, Schroda M, Goodenough U, Stitt M, Griffiths H, Jonikas MC. 2016. A repeat protein links
Rubisco to form the eukaryotic carbon-concentrating organelle. *Proc Natl Acad Sci U S A*
113:5958–5963.
Mangan NM, Flamholz A, Hood RD, Milo R, Savage DF. 2016. pH determines the energetic
efficiency of the cyanobacterial CO2 concentrating mechanism. *Proc Natl Acad Sci U S A*
113:E5354–62.
Marcus Y, Schwarz R, Friedberg D, Kaplan A. 1986. High CO2 Requiring Mutant of Anacystis

nidulans R2. *Plant Physiol* **82**:610–612.

McGrath JM, Long SP. 2014. Can the cyanobacterial carbon-concentrating mechanism increase
photosynthesis in crop species? A theoretical analysis. *Plant Physiol* **164**:2247–2261.

Merlin C, Masters M. 2003. Why is carbonic anhydrase essential to Escherichia coli? *J Bacteriol*
185. doi:10.1128/JB.185.21.6415

Mueller-Cajar O. 2017. The Diverse AAA+ Machines that Repair Inhibited Rubisco Active Sites.
Front Mol Biosci **4**:31.

Mueller-Cajar O, Morell M, Whitney SM. 2007. Directed evolution of rubisco in Escherichia coli
reveals a specificity-determining hydrogen bond in the form II enzyme. *Biochemistry*
46:14067–14074.

Nevins CP, Vierck JL, Bogachus LD, Velotta NS, Castro-Munozledo F, Dodson MV. 2005. An
Inexpensive Method for Applying Nitrogen Evaporation to Hexane-containing 24- or 96-well
Plates. *Cytotechnology* **49**:71–75.

Occhialini A, Lin MT, Andralojc PJ, Hanson MR, Parry MAJ. 2016. Transgenic tobacco plants
with improved cyanobacterial Rubisco expression but no extra assembly factors grow at
near wild-type rates if provided with elevated CO2. *Plant J* **85**:148–160.

Oltrogge LM, Chaijarasphong T, Chen AW, Bolin ER, Marqusee S, Savage DF. 2020.
Multivalent interactions between CsoS2 and Rubisco mediate α-carboxysome formation.
Nat Struct Mol Biol **27**:281–287.

Orr DJ, Worrall D, Lin MT, Carmo-Silva E, Hanson MR, Parry MAJ. 2020. Hybrid
Cyanobacterial-Tobacco Rubisco Supports Autotrophic Growth and Procarboxysomal
Aggregation. *Plant Physiol* **182**:807–818.

Orth JD, Fleming RMT, Palsson BØ. 2010. Reconstruction and Use of Microbial Metabolic
Networks: the Core Escherichia coli Metabolic Model as an Educational Guide. *EcoSal Plus*
4:1–47.

Peekhaus N, Conway T. 1998. What's for dinner?: Entner-Doudoroff metabolism in Escherichia
coli. *J Bacteriol* **180**:3495.

Pellicer MT, Nuñez MF, Aguilar J, Badia J, Baldoma L. 2003. Role of 2-Phosphoglycolate
Phosphatase of Escherichia coli in Metabolism of the 2-Phosphoglycolate Formed in DNA
Repair. *J Bacteriol* **185**:5815–5821.

Price GD, Badger MR. 1989a. Isolation and characterization of high CO2-requiring-mutants of
the cyanobacterium Synechococcus PCC7942: two phenotypes that accumulate inorganic
carbon but are apparently unable to generate CO2 within the carboxysome. *Plant Physiol*
91:514–525.

Price GD, Badger MR. 1989b. Expression of Human Carbonic Anhydrase in the
Cyanobacterium Synechococcus PCC7942 Creates a High CO2-Requiring Phenotype
Evidence for a Central Role for Carboxysomes in the CO2 Concentrating Mechanism. *Plant*
Physiol **91**:505–513.

Rae BD, Long BM, Badger MR, Price GD. 2013. Functions, compositions, and evolution of the
two types of carboxysomes: polyhedral microcompartments that facilitate CO2 fixation in
cyanobacteria and some proteobacteria. *Microbiol Mol Biol Rev* **77**:357–379.

Raven JA, Beardall J, Sánchez-Baracaldo P. 2017. The possible evolution and future of CO2-
concentrating mechanisms. *J Exp Bot* **68**:3701–3716.

Sage RF, Sage TL, Kocacinar F. 2012. Photorespiration and the evolution of C4
photosynthesis. *Annu Rev Plant Biol* **63**:19–47.

Sawaya MR, Cannon GC, Heinhorst S, Tanaka S, Williams EB, Yeates TO, Kerfeld C a. 2006.
The structure of beta-carbonic anhydrase from the carboxysomal shell reveals a distinct
subclass with one active site for the price of two. *J Biol Chem* **281**:7546–7555.

Scott KM, Leonard JM, Boden R, Chaput D, Dennison C, Haller E, Harmer TL, Anderson A,
Arnold T, Budenstein S, Brown R, Brand J, Byers J, Calarco J, Campbell T, Carter E,
Chase M, Cole M, Dwyer D, Grasham J, Hanni C, Hazle A, Johnson C, Johnson R, Kirby B,

Lewis K, Neumann B, Nguyen T, Nino Charari J, Morakinyo O, Olsson B, Roundtree S, Skjerve E, Ubaldini A, Whittaker R. 2019. Diversity in CO2-Concentrating Mechanisms among Chemolithoautotrophs from the Genera Hydrogenovibrio, Thiomicrorhabdus, and Thiomicrospira, Ubiquitous in Sulfidic Habitats Worldwide. *Appl Environ Microbiol* **85**:1–19.

Sezonov G, Joseleau-Petit D, D'Ari R. 2007. Escherichia coli physiology in Luria-Bertani broth. *J Bacteriol* **189**:8746–8749.

Shih PM, Occhialini A, Cameron JC, Andralojc PJ, Parry MAJ, Kerfeld CA. 2016. Biochemical characterization of predicted Precambrian RuBisCO. *Nat Commun* **7**:10382.

Stauffer GV. 2004. Regulation of Serine, Glycine, and One-Carbon Biosynthesis. *EcoSal Plus* **1**:1–22.

Stolper DA, Revsbech NP, Canfield DE. 2010. Aerobic growth at nanomolar oxygen concentrations. *Proc Natl Acad Sci U S A* **107**:18755–18760.

Szyperski T. 1995. Biosynthetically directed fractional 13C-labeling of proteinogenic amino acids. An efficient analytical tool to investigate intermediary metabolism. *Eur J Biochem* **232**:433–448.

Taymaz-Nikerel H, Borujeni AE, Verheijen PJT, Heijnen JJ, van Gulik WM. 2010. Genome-derived minimal metabolic models for Escherichia coli MG1655 with estimated in vivo respiratory ATP stoichiometry. *Biotechnol Bioeng* **107**:369–381.

Tsai Y-CC, Lapina MC, Bhushan S, Mueller-Cajar O. 2015. Identification and characterization of multiple rubisco activases in chemoautotrophic bacteria. *Nat Commun* **6**:8883.

Unden G, Dünnwald P. 2008. The Aerobic and Anaerobic Respiratory Chain of Escherichia coli and Salmonella enterica: Enzymes and Energetics. *EcoSal Plus* **3**. doi:10.1128/ecosalplus.3.2.2

Wang L, Jonikas MC. 2020. The pyrenoid. *Curr Biol* **30**:R456–R458.

Wheatley NM, Sundberg CD, Gidaniyan SD, Cascio D, Yeates TO. 2014. Structure and identification of a pterin dehydratase-like protein as a ribulose-bisphosphate carboxylase/oxygenase (RuBisCO) assembly factor in the α-carboxysome. *J Biol Chem* **289**:7973–7981.

Wildman SG. 2002. Along the trail from Fraction I protein to Rubisco (ribulose bisphosphate carboxylase-oxygenase). *Photosynth Res* **73**:243–250.

Wilson RH, Martin-Avila E, Conlan C, Whitney SM. 2018. An improved Escherichia coli screen for Rubisco identifies a protein-protein interface that can enhance CO2-fixation kinetics. *J Biol Chem* **293**:18–27.

Winkler ME, Ramos-Montañez S. 2009. Biosynthesis of Histidine. *EcoSal Plus* **3**:1–33.

Wu A, Hammer GL, Doherty A, von Caemmerer S, Farquhar GD. 2019. Quantifying impacts of enhancing photosynthesis on crop yield. *Nat Plants* **5**:380–388.

 # Figures Supplements

 Figure 2 - figure supplement 1. The OptSlope algorithm for designing rubisco-coupled E. coli strains. *Optslope*
searches for metabolic knockout mutants in which biomass production is coupled to flux through a reaction of choice
*(e.g. rubisco) at all growth rates. (**A**) Shows the space of feasible biomass production and rubisco fluxes for wildtype*
(WT, grey) and a knockout mutant (green). For WT, biomass production and, therefore, growth rate, are independent
of rubisco at all feasible growth rates (i.e. within the grey polygon). The mutant is "rubisco-coupled" because maximal
biomass production requires non-zero rubisco carboxylation flux and increasing biomass production demands
*increased carboxylation. The slope of this relationship is the "coupling slope." (**B**) We computationally generated pairs*
of E. coli central metabolic knockouts and calculated the coupling slope on nine carbon sources: glucose (gluc), fructose
(fruc), gluconate (gnt), ribose (ribo), succinate (succ), xylose (xyl), glycerate (gly^{ate}), acetate (ace) and glycerol (gly^{ol}).
Each double knockout is summarized as a 3×3 matrix of coupling slopes. Black denotes a rubisco-independent mutant
and maroon a coupling slope of 0. The published mutant ΔgapA (Mueller-Cajar et al., 2007) has a coupling slope of 0
*(left), while the ΔrpiAB Δedd strain is rubisco-coupled on seven of the carbon sources (right). (**C**) Feasible phase space*
*diagram for the ΔgapA strain shows that biomass production is not coupled to rubisco flux. (**D**) ΔrpiAB Δedd has a*
positive coupling slope in glycerol, gluconate and xylose media.

982
Figure 2 - figure supplement 2. Proposed mechanisms of rubisco-dependent growth in CCMB1. (A) *CCMB1*
depends on rubisco and prk for growth in glycerol, gluconate, and xylose minimal media. The common mechanism is
an inability to metabolize ribulose-5-phosphate (Ru5P) due to the deletion of both ribose-phosphate isomerase genes
(ΔrpiAB). When gluconate or xylose is the growth substrate, Ru5P must be produced in order to metabolize the carbon
source. Though wild type E. coli can metabolize gluconate via the ED pathway, the ED dehydratase knockout (Δedd)
in CCMB1 blocks this route and forces 1:1 production of Ru5P from gluconate. Expression of prk and rubisco opens a
new route of Ru5P metabolism, thus enabling CCMB1 to grow in gluconate or xylose media. Since extracellular glycerol
is converted to glyceraldehyde 3-phosphate (GAP), it can be metabolized through lower glycolysis or through
gluconeogenesis. The gluconeogenesis route produces hexoses that enter the pentose phosphate pathway, which is
required to synthesize ribose 5-phosphate (Ri5P) for nucleotide and histidine biosynthesis. Depending on the growth
rate, products of Ri5P make up 5-25% of E. coli biomass (Bremer and Dennis, 2008; Taymaz-Nikerel et al., 2010). As
*shown in (**B**), the pentose phosphate pathway forces co-production of Ri5P, Ru5P and xylulose 5-phosphate (Xu5P).*
In the absence of rpi activity, there is no pathway for metabolism of Xu5P or Ru5P. This defect is complemented by the
*expression of rubisco and prk. Notably, rubisco can also oxygenate RuBP, as shown in (**C**). E. coli can, in principle,*
recycle the oxygenation product 2-phosphoglycolate (2PG) though an ersatz photorespiratory pathway via tartronate
semialdehyde. This pathway is not the dominant mechanism of rubisco complementation because CCMB1:p1A cannot
*grow in ambient air, where O₂ is abundant (Figure 2D). Panel (**D**) describes the initial metabolism of extracellular*
glycerol, gluconate and xylose in E. coli. Extracellular carbon sources are marked with a grey background throughout.
Abbreviations: 3-phosphoglycerate (3PG), 2-phosphoglycolate (2PG), glyceraldehyde 2-phosphate (GAP),
dihydroxyacetone phosphate (DHAP), ribose 5-phosphate (Ri5P), ribulose 5-phosphate (Ru5P), xylulose 5-phosphate
(Xu5P), ribulose 1,5-bisphosphate (RuBP), 2-keto-3-deoxy-6-phosphogluconate (KDGP), fructose 6-phosphate (F6P),
fructose 1,6-bisphosphate (F-1,6-BP), phosphoenolpyruvate (PEP).
1005

1006
Figure 2 - figure supplement 3. CCMB1 depends on rubisco and prk for growth in minimal media. (A) *Expression*
of rubisco and prk complements CCMB1 growth on M9 glycerol and gluconate media under 10% CO_2, but not in
ambient conditions (100 nM aTc induction in M9 plates). Mutations ablating rubisco (cbbL⁻) or prk (prk⁻) activity abrogate
growth in selective media but not in LB under 10% CO_2. Growth in LB is rubisco-independent in 10% CO_2, but CCMB1
does not grow in ambient air even when supplied rich media because it lacks CA genes (Merlin and Masters, 2003).
*Growth curves in (**B**) show the rubisco-dependence of CCMB1:p1A growth in glycerol (green) and gluconate (blue)*
media under 5% CO_2 in a gas controlled plate reader (Tecan Spark, Methods). Negative controls (CCMB1:p1A cbbL⁻
in glycerol or gluconate media) and uninduced cultures failed to grow in these conditions (dashed grey lines). Though
three curves are plotted for each condition in (B), experiments were conducted in technical sextuplicate. Replicates
were all consistent.
1017
1018
1019

1020
Figure 2 - figure supplement 4. CCMB1 does not require oxygen for growth in minimal media. *(**A**) Titer plating*
assays were used to measure the viability of CCMB1:p1A grown on glycerol media under ambient air (≈0.04% CO_2,
21% O_2), 10% CO_2 (balance air), and an anoxic mix of 10% CO_2 and 90% N_2 ("No O_2"). Since E. coli cannot ferment
*glycerol, 20 mM NO_3^- was provided as an alternate electron acceptor as marked. (**B**) CCMB1:p1A grows on glycerol*
media in the absence of O_2 so long as nitrate is provided. While CCMB1:p1A colonies are noticeably smaller than WT
in panel (A), the colony count is indistinguishable, as quantified in panel (B). Experiments were conducted in biological
duplicate (i.e. pre-cultures from distinct colonies) with at least two technical replicates (repeated spotting from the same
preculture).
1029

1030

Figure 3 - figure supplement 1. A series of selection experiments produced mutant plasmids that permit
rubisco-dependent growth in ambient air. (A) *pCB and pCCM plasmids together encode 20 H. neapolitanus genes*
including 12 confirmed CCM components. pCB carries kanamycin resistance and has two transcriptional units
expressed under an aTc-inducible $P_{LtetO-1}$ promoter (Lutz and Bujard, 1997). The first derives from pHnCB10 (Bonacci
et al., 2012) and expresses 10 carboxysome proteins. The second expresses phosphoribulokinase (prk). pCCM carries
chloramphenicol resistance and expresses an 11 gene operon from H. neapolitanus that contains both putative and
confirmed CCM genes (Desmarais et al., 2019). Although pCB expresses both rubisco and prk, CCMB1:pCB did not
initially grow in M9 media under 10% CO_2 (not shown) and so we undertook a series of selections, described in panels
(B-D) that ultimately led to isolation of pCB' and pCCM' plasmids that together enable CCMB1 to grow in ambient air.
(B) *We first selected CCMB1:pCB for growth on minimal media by screening for mutants able to grow on M9 glycerol*
and then M9 gluconate media. Gluconate growing mutant #9 (gg.9) was used for subsequent experiments as this
mutant was found to grow best on gluconate (as shown in E). (C) Plasmid extracted from gg.9 was deep sequenced
and electroporated into naive CCMB1 to test for plasmid linkage of growth on minimal. (D) Selection for rubisco-
dependent growth in ambient air. A turbid pre-culture of the CCMB1:pCB gg.9+pCCM double transformant was washed
and plated on M9 glycerol media under ambient air. Colonies formed after ≈20 days (as shown in F). 40 colonies (s.1-
40) were picked into rich media, grown to saturation, washed and plated on M9 glycerol media to verify growth under
ambient air. Roughly ¼ of chosen colonies regrew under ambient air to varying degrees (s.1-6 are shown in G). Plasmid
extracted from several strains was deep-sequenced and electroporated into naive CCMB1 to test plasmid-linkage of
growth on glycerol minimal media in ambient air. Pooled plasmid extracted from s.4 was found to confer replicable
growth in ambient air (as shown in H). PCR and Gibson cloning were used to reconstruct the individual pCB and pCCM
plasmids from this pool. We termed these reconstructed plasmids pCB' and pCCM'. (E) Restreaking of gluconate-
growing mutants gg.8-12 described in panel B shows that gg.9 grew best on gluconate. (F) CCMB1:pCB gg.9+pCCM
double transformants were plated for mutants on M9 glycerol media under ambient air. A negative control lacking
carboxysome genes (CCMB1:p1A+pCCM) was plated at the same time. Colonies formed after 20 days (bottom right)
only on induced plates (100 nM) and only when all CCM genes were provided (i.e. pCB gg.9 and pCCM). (G) Several
of the chosen colonies regrew in ambient air. Growth characteristics varied from colony to colony, suggesting genetic
variation. (H) Pooled plasmid extracted from s.4 was found to permit naive CCMB1 to grow in ambient air. For
comparison, plasmid from s.6 produced less reproducible air growth.

1059
Figure 3 - figure supplement 2. pCB' and pCCM' permit CCMB1 to grow in ambient air. (A) *Biological triplicate*
growth curves from a bioreactor bubbling ambient air. CCMB1 co-transformed with post-selection plasmids pCB' and
pCCM' (CCMB1:pCB' + pCCM') grows well (purple, "full CCM"), while rubisco and prk alone are insufficient for growth
in air (green, "rubisco+prk"). Maximal growth rates for the "full CCM" cultures ranged from 0.03-0.06 hr^{-1}, corresponding
to doubling times of 12-25 hours. As these are biological replicate cultures, heterogeneity in growth kinetics could be
due to genetic effects (e.g. point mutations in founding colonies) or non-genetic differences (e.g. varying degree of
carboxysome production during pre-culturing). (B) Data for the same strains grown in a 96 well plate in ambient air in
a shaking plate reader. Different shades mark biological replicates (pre-cultures deriving from three distinct colonies).
Additionally, each preculture was used to inoculate at least 12 technical replicates. (C) Quantification of the experiment
in panel (B) using endpoint data at 80 hours for biological and technical replicates. Panel (C) uses the same colors as
*(A) and (B) with the addition of a rubisco active site mutant as a negative control (grey, CCMB1:p1A⁻ + vec). '****'*
indicates $P < 10^{-10}$. P-values were calculated with a Bonferroni-corrected two-sided Mann-Whitney-Wilcoxon test. 10^{4}-
fold bootstrapping was used to compare "full CCM" data to "rubisco + prk" and estimate a confidence interval for the
effect of expressing a full CCM on growth in ambient air, which gave a 99.9% confidence interval of 0.56-0.64 OD units.
1074

1075
Figure 4 - figure supplement 1. Targeted mutations to the CCM eliminate growth in ambient air. *Pre-cultures*
were grown to saturation in 10% CO_2 and then diluted to an optical density of 0.01 (600 nm) into two tubes (Methods).
*One tube was grown in 10% CO_2 and the other in ambient air, as diagrammed in (**B**). Cells were incubated for 4 days*
*before measuring optical density in (**A**) and 12 days in (**C**). The left bar (darker color) gives the mean endpoint density*
of biological quadruplicate cultures in 10% CO_2 and the right bar (lighter color) gives the mean in ambient air. Error
bars give a 95% confidence interval of measurements. (A) and (C) share the leftmost 11 strains. From left to right: a
positive control (grey, grows in both conditions), two negative controls carrying active site mutants of rubisco
(CCMB1:p1A^-+vec and CCMB1:pCB^+^+pCCM'), CCMB1 expressing rubisco and prk but no CCM genes (green,
CCMB1:p1A+vec) or an incomplete set of CCM genes (green, CCMB1:p1A+pCCM'), CCMB1:pCB+pCCM which
carries the pre-selection CCM plasmids (purple), and CCMB1:pCB'+pCCM' which carries the post-selection plasmids.
"vec" denotes an appropriate vector control (pFA-sfGFP). The following pairs of maroon bars describe strains carrying
plasmids with targeted CCM mutations: CCMB1:pCB' CsoSCA^-+pCCM' which carries an inactivating mutation to
carboxysomal carbonic anhydrase, CCMB1:pCB' CsoS2 ΔNTD +pCCM' harboring a deletion of the N-terminal domain
of CsoS2 responsible for recruiting rubisco to the carboxysome, CCMB1:pCB' ΔcsoS4AB + pCCM' lacking both genes
pentameric vertex proteins, and CCMB1:pCB' DabA1^- + pCCM' carrying an inactivated DAB carbon uptake system.
*(**A**) CCMB1 grows well in ambient air only when given a full complement of CCM genes on the post-selection plasmids.*
*All mutations to the CCM abrogate growth in air (maroon). Panel (**C**) shows consistent results over a 12-day time period.*
(C) describes three additional mutants: CCMB1:pCB' CbbL Y72R + pCCM' carrying a mutation to the rubisco large
subunit that eliminates rubisco-CsoS2 binding, CCMB1:pCB' + pCCM' CbbQ^- harboring inactivating mutation to the
CbbQ subunit of the rubisco activase complex, and CCMB1:pCB' + pCCM' ΔacRAF lacking the putative rubisco
chaperone acRAF (CCMB1:pCB' + pCCM' ΔacRAF). Ablation of rubisco-CsoS2 interaction should eliminate
recruitment of rubisco to the carboxyome (Oltrogge et al., 2020). Accordingly, the Y72R mutation eliminated growth in
air. Chaperone mutants (CbbQ or acRAF) were both viable in air, though removal of acRAF produced a substantial
growth defect (2.5 fold in mean and 8.5 fold in median final density). The positive control strain is the CAfree strain
expressing human carbonic anhydrase II (Methods). P-values calculated by a one-sided Mann-Whitney-Wilcoxon test.
'' denotes a $P < 0.05$. Detailed description of all plasmid abbreviations is given in Table S2.*

1102
Figure 5 - figure supplement 1. CCMB1:pCB' + pCCM' produces polyhedral bodies resembling carboxysomes
when grown in ambient air. *Transmission electron micrographs of air-grown CCMB1:pCB'+pCCM' (images on the*
right) show morphological carboxysomes inside cells (white arrows). The negative control for carboxysome expression
is CAfree:pFE-sfGFP + pFA-HCAII (top left). WT:pHnCB10 is the parent strain transformed with a plasmid expressing
10 carboxysome genes and previously shown to enable purification of carboxysome structures from E. coli (Bonacci et
al., 2012). This was intended as a positive control, but we did not observe carboxysome structures in electron
micrographs of this strain, perhaps because of excessive IPTG induction (500 mM) as previously reported. Expression
of carboxysome genes was associated with production of black staining stress granules in both the experiment and
pHnCB10 control. These granules were not observed in images of the negative control.
1112

1113
Figure 5 - figure supplement 2. Isotopic composition of amino acids from total biomass hydrolysate. *Cells were*
grown under ambient air in M9 media containing 99% ^{13}C labeled glycerol (0.4% v/v) so that nearly all ^{12}C in biomass
must derive from inorganic carbon. The isotopic composition of amino acids in total biomass hydrolysate of
CCMB1:pCB' + pCCM' and an appropriate rubisco-independent control were measured via LC-MS (Methods). The
control strain is CAfree complemented with the human carbonic anhydrase II, which does not express rubisco
(Methods). Serine and valine, which are marked in green, are downstream of the rubisco product 3PG in E. coli central
metabolism and, accordingly, show significantly greater ^{12}C incorporation in CCMB1:pCB' + pCCM' than the control.
Histidine, threonine, proline and glutamate are synthesized from precursors deriving from the TCA cycle and pentose
phosphate pathways, and thus their carbon atoms do not derive from 3PG (Szyperski, 1995). Arginine is synthesized
via a rubisco-independent carboxylation of glutamate (by the addition of carboxyphosphate, (Gleizer et al., 2019)), and
so the difference between arginine and glutamate labeling is used to calculate the isotopic composition of intracellular
inorganic carbon (C_i, Methods). Notably, intracellular C_i derives both from extracellular C_i (predominantly ^{12}C) and
decarboxylation of the 99% ^{13}C glycerol carbon source. As such, the composition will depend on C_i uptake as well as
the rate of glycerol metabolism. Control cells grew faster than CCMB1:pCB'+pCCM, which can explain why arginine
from these cells contains significantly less ^{12}C and more ^{13}C (from rapid glycerol decarboxylation).
1129

1130
Figure 5 - figure supplement 1. ^{12}C enrichment on serine is consistent with in vivo CO_2 fixation. Cells were grown
under ambient air in M9 media containing 99% ^{13}C labeled glycerol (0.4% v/v) so that nearly all ^{12}C in biomass must
derive from inorganic carbon. In (**A**) ^{13}C atoms are depicted as open circles and fractional ^{12}C labeling by a partial green
fill color. In CCMB1, 3-phosphoglycerate (3PG) can be produced either through glycolytic metabolism of glycerol (via
dihydroxyacetone-phosphate, DHAP) or through rubisco-catalyzed carboxylation of RuBP. At most ⅙ of the carbon
atoms on 3PG will be ^{12}C when rubisco is active in vivo. In practice this fraction will be less than ⅙ because some of
the intracellular inorganic carbon pool (C_i) derives from decarboxylation of ^{13}C labeled glycerol and also because a
large fraction of intracellular 3PG is produced through glycolysis (Methods). Serine is a direct metabolic product of 3PG
and so reports on the labeling of 3PG. As such, we measured the ^{12}C composition of amino acids in total protein
hydrolysate via LC-MS (Methods). (**B**) Serine from CCM-expressing CCMB1 cells ('Experiment') displayed roughly
threefold higher ^{12}C labeling than controls, which grow in a rubisco-independent manner (Methods). (**C**) Rubisco
carboxylation draws from the intracellular inorganic carbon pool, whose ^{12}C composition can be inferred for each
sample by comparing the labeling of L-arginine and L-glutamate (Methods). The mean ^{12}C fraction of intracellular C_i
was estimated to be $25\% \pm 4\%$ and $67\% \pm 28\%$ for the control and experiment respectively. (**D**) These values were
integrated to estimate the percent of 3PG production flux that is due to carboxylation by rubisco (Methods), which was
inferred to be $24\% \pm 15\%$. These values compare favorably with predictions made via Flux Balance Analysis (19.5-
24%, Methods). A sampling method was used to estimate the uncertainty in these rubisco flux inferences (Methods).
99% confidence intervals on the rubisco flux fraction were strictly positive for each biological replicate, with 99% of all
posterior estimates between 4% and 51% across all four replicates.