Article

Human Endogenous Retrovirus K Rec forms a regulatory loop with MITF that opposes the progression of melanoma to an invasive stage

Manvendra Singh[1,2#], Huiqiang Cai[1#], Mario Bunse[1], Cedric Feschotte[2] and Zsuzsanna Izsvák*[1]

[1] Max-Delbrück-Center for Molecular Medicine (MDC), in the Helmholtz Society Robert-Rössle-Strasse 10, 13125 Berlin, Germany

[2] Department of Molecular Biology & Genetics, 526 Campus Road, Cornell University, Ithaca, NY 14853

[#] Equal Contributions

* Correspondence:

Abstract: In the human genome, HERV-K(HML2) is the most recently endogenized retrovirus (ERV). While HERV-K(HML2) transcription is observed in healthy tissues, various cancers showed the upregulation of retroviral derived endogenized accessory products (e.g., envelope (Env), Np9 and Rec). Still, it is not clear whether the different HERV-K-derived genes contribute to a disease, or they are mere by-products. Here, we focus on the potential role of Rec in melanoma. Our in vitro model and high throughput data mining, including single-cell transcriptome analyses of patent's material, reveal that Rec expression marks the proliferative (still controllable) stage of melanoma, and is involved in maintaining a delicate balance between cell proliferation and invasion. Thus, similar to melanocyte-inducing transcription factor (MITF), Rec is a sensitive marker of melanoma progression. Our Rec-knockdown in vitro system can faithfully model a subpopulation (MITF malignancy) of melanoma cells in human patients. Like Env, Rec modulates an endothelial-mesenchymal transition (EMT)-like process of cancer progression; however, they seem to affect the phenotype switch inversely. Rec inhibits the transition to the invasive state by altering the expression level of some key determinants of the EMT-like process, including MITF that directly binds the LTR5 _Hs of HERV-K. The Hominoid-specific HERV-K products might explain certain species-specific features of melanoma progression, and pinpoint to the limitation of using animal models in melanoma studies.

Keywords: HERV-K, Rec, Melanoma, Metastasis, and MITF

1. Introduction

Endogenous retroviruses (ERVs) are remnants of retroviruses that once infected the germline and became vertically inherited as part of the host genome. Sequences derived from various ERVs account for 8% of the human genome [1], reflecting multiple waves of retroviral invasion in the human lineage. In the human genome, HERV-K(HML2) is the most recently endogenized HERV, and several copies of this subgroup still express retroviral open reading frames (e.g., Gag, Pro-Protease, Pol-Polymerase, Env-Envelope), and are even capable of producing (non-infective) viral particles [3]. The canonical function of HERV-K Env is to enable the retrovirus to traffic through the endosomal membrane of the host cells or directly enter cells by fusing with their membranes. The endogenization process of HERV-K(HML2) (HERV-K thereafter) included the emergence of the accessory proteins, Np9 and Rec, translated from splicing products of the Env mRNA [3, 4]. Env, Np9, and Rec interact with different host-encoded cellular factors and have distinct functional activities. Rec is a small RNA-binding protein considered to be a functional homolog of the HIV Rev accessory protein [5]. Through its interaction with the CRM1 nuclear export factor [4, 6], Rec is involved in exporting un-spliced HERV-K RNA from the nucleus [3, 7]. Also, Rec can directly modulate cell signaling via binding to the promyelocytic leukemia zinc finger (PLZF) protein, a transcriptional repressor of the c-MYC proto-oncogene [8]. Rec also binds the TZFP (testicular zinc-finger protein) and the hSGT (human small glutamine-rich tetratricopeptide repeat protein), which are transcriptional repressors

of the androgen receptor [9, 10]. While it is evident that Rec must have first evolved to facilitate HERV-K replication, there is speculation that the protein may have been coopted for physiological function, such as spermatogenesis [6, 11] and defense against viral infections in the early human embryo [12].

It has been reported that at least 18 HERV-K genomic loci can be transcribed in healthy human tissues and express potentially-coding Rec or Np9 mRNAs (alone or in combination with retroviral genes). However, the landscape of transcribed HERV-K loci differs considerably between cell/tissue types [13]. Rec RNA/protein expression has been observed in human embryonic tissues, placenta, and retina [14, 15]. Interestingly, HERV-K is expressed in both germinal- and pluripotent stem cells, providing a particular link between these cell types [16]. While HERV-K transcription has been observed at low levels in various healthy tissues, it is strongly induced upon environmental stress, such as ultraviolet irradiation (e.g., UVB and UVC), starvation, or viral infection that may manifest the global DNA hypomethylation [17-21]. Furthermore, many studies have reported robust mRNA upregulation of several genomic loci of HERV-K in various disease states, including certain autoimmune diseases and several cancers, especially lung, breast cancers, germ cell tumors, and melanoma [18, 22, 23] and reviewed in [24, 25]. Moreover, transcriptional activation and translation of various HERV-K products [22, 26] and even virus-like particles have been observed in specific cancer cells such as teratoma, embryonic carcinoma, and melanomas [3, 27-29].

Whether and how the overexpression of HERV-K products contributes to the progression of these diseases is the subject of many studies, debates, and speculations. Several mechanisms have been proposed by which HERV-K might contribute to disease. The activated LTRs could act as alternative promoters and de-regulate tumor suppressor genes or proto-oncogenes (reviewed in [35, 36]). Furthermore, the Env protein of HERV-K, via its fusogenic property, is capable of inducing cell-cell fusion and could, therefore, contribute to tumor invasiveness [18, 37, 38]. Env could also have oncogenic properties through direct interference with cellular signaling pathways (e.g., RSS/MEK/ERK) [39, 40]. Indeed, the overexpression of the HERV-K Env protein has been reported to induce an epithelial to mesenchymal transition (EMT)-like process [39, 41], a crucial event in oncogenesis leading to a more malignant phenotype. In addition, Env was suggested to promote tumorigenesis via modulation of the immune response [42-44]. Confusingly, HERV-K may have either a positive or negative effect on the immune system [45, 46]. Similar to Env, the expression of the accessory genes, Rec and Np9 were readily increased in pathological conditions [47], including germ cell tumors and melanoma [4, 6, 22, 27, 28, 48-52]. Still again, it was not clear whether and how these factors could contribute to tumorigenesis and other disease states.

To begin answering this question, we investigate the oncogenic properties of HERV-K Rec (Rec thereafter) in melanoma. We used in vitro models as well as high throughput data mining, including single-cell transcriptome analyses of patient samples, to examine the impact of Rec expression in the spatiotemporal progression of melanoma. We found that Rec marks the proliferative state of melanoma, and similarly to Env [39], modulates the EMT-like process of cell transformation. However, surprisingly, in contrast to Env [39], Rec is inhibiting and not activating the transition from proliferative to invasive state and thus might be, in fact, a protective factor in melanoma.

2. Materials and Methods

2.1. Construction of the KD constructs

All 3 KD targeting the ERVK6(HML2.HOM) locus. KD1 targets 3'UTR, KD2 3'UTR, and KD3 the gene body of Rec.

KD1 construct 1 (3'UTR_I): (as) 5'-'ATCCATTCAACTCTGAGTGGA-3'

KD2 construct 2 (5'UTR_I): (as) 5'-TAAGGCTGACTGTAGATGTAC-3'

KD3 construct 4 (Rec): (as) 5'-CAACGGTGCTCGATTGCGATG-3'

2.2. Cell Culture

The melanoma cell line, A375 (ATCC CRL-1619) was cultivated in RPMI1640 (Thermo Fisher Scientific) supplemented with 10% fetal bovine serum (FBS; PAA Laboratories, Pasching, Austria), 80 U/mL penicillin and streptomycin (Lonza, Basel, Switzerland) at 37°C in a humidified atmosphere

99 of 95% air and 5% CO_2. The melanoma cell line, SKMel-28 (ATCC HTB-72) was maintained in
100 Dulbecco's modified Eagle's medium (DMEM, Life Technologies, England) containing GlutaMAX™-
101 I supplement (Gibco, catalog number: 31966-021), with 10% FBS and penicillin and streptomycin in
102 the same condition as above.

2.3. Transfection of melanoma cells and cell sorting

To knock down Rec, melanoma cells were transfected with pT2-MP71-KD1-Rec-EGFP, pT2-
MP71-KD2-Rec-EGFP, pT2-MP71-KD3-Rec-EGFP plasmids in the presence of pT2-CAGGS-SB100X,
using the Neon transfection system (Life technologies). A piggyBac transposon-based vector,
targeting GFP, was used as a control. 10μl Kit with the electroporation setting, 1350 V (pulse voltage),
20ms (pulse width), two pulses. 2×10^5 cells in 11 μL resuspension buffer R were combined with 2 μL
of purified plasmid mixture (50 ng transposase and 500 ng transposon). Transfected cells were
transferred to 6-well tissue culture plates containing 2 mL culture medium. EGFP+ cells were sorted
at weeks 1 and 3 post-transfection by FACSAria II cell sorter (Becton).

2.4. Quantitative Real-Time PCR and semi-quantitative PCR

Total RNA was extracted using the Direct-zol RNA MiniPrep kit (Zymo Research) according to
the protocol of the manufacturer. Reverse transcription was performed from 1μg total RNA with the
High Capacity RNA-to-cDNA kit (Applied Biosystems). The isolated RNA would be treated with
additional DNaseI (Invitrogen), to get rid of potential DNA contamination. Quantitative real-time
PCR (qRT-PCR) was carried out on ABI7900 with PowerUp SYBR Green Master Mix kit (Applied
Biosystems), according to the recommendations of the manufacturer. For quantification of mRNA
expression levels, real-time PCR was run in triplicates for each cDNA sample, using GAPDH and 18s
RNA as the internal controls for SKMel-28 and A375 cells, respectively, for primers and annealing
temperatures see Table 1. Data were analyzed using the comparative CT (2-ΔΔCT) method, which
describes relative gene expression.

Table 1. The primers/oligos used in this study.

Name	Application	Sequence (5' – 3')	Tm(°C)
HML-2-Env	Real-Time qPCR	F: GCTGCCCTGCCAAACCTGAG R: CCTGAGTGACATCCCGCTTACC	60
HML-2-Gag	Real-Time qPCR	F: AGCAGGTCAGGTGCCTGTAACATT R: TGGTGCCGTAGGATTAAGTCTCCT	60
HML-2-Np9	Real-Time qPCR	F: AGATGTCTGCAGGTGTACCCA R: CTCTTGCTTTTCCCCACATTTC	60
HML-2-Rec	Real-Time qPCR	F: ATCGAGCACCGTTGACTCACAAGA R: GGTACACCTGCAGACACCATTGAT	60
MITF-M	Real-Time qPCR	F:ATGCTGGAAATGCTAGAATATAATCACT R: GAATGTGTGTTCATGCCTGG	60
SPANXB1/B2	Real-Time qPCR	F: AGGCCAATGAGGCCAACAAGAC R: TCCTCCTGTAGCGAACCACTAG	60

Singh et al. (2020)

18S RNA	Real-Time qPCR	F: GTAACCCGTTGAACCCCATT R: CCATCCAATCGGTAGTAGCG	60
GAPDH	Real-Time qPCR	F: CAATGACCCCTTCATTGACCTC R: AGCATCGCCCCACTTGATT	60

2.5. RNA sequencing and data analysis

The concentration of RNA was measured on NanoDrop Spectrophotometer ND-1000, and the quality of RNA was analyzed using Agilent RNA 6000 Nano Kit on Aligent 2100 Bioanalyzer machine. The RNA sequencing library was prepared from 550 ng of RNA, using Illumina TruSeq Stranded mRNA LT Set A kit (Cat. no. RS-122-2101), according to TruSeq Stranded mRNA Sample Prep LS Protocol. Sequencing was performed on an Illumina HiSeq 2000 platform as 100 bp first strand-specific paired-end reads.

Sample-specific barcoded sequencing reads were de-multiplexed from multiplexed flow cells and by using CASAVA. 1.8.2 BCL files were converted to FASTQ format files. The quality of the raw sequence reads was determined by using the FastQC. Reads with a quality score below 30 were removed. We removed the highly variable two nt from the ends of the remaining sequencing reads and mapped them over the reference genome (Human hg19/GRCh37) and transcriptome model (hg19.refseq.gtf). hg19/GRCh37 and hg19.refseq.gtf that were downloaded from USCS tables (http://hgdownload.cse.ucsc.edu/goldenPath/hg19/bigZips/) by using Bowtie 2.0.5.0, Samtools 0.1.17.0 and TopHat v2.0.8 with parameters: "tophat2 -p 8 –r 150--mate-std-dev 140 --library-type fr-first-strand". Transcript assembly for each sample was conducted with Cufflinks v2.08, generating FPKM values. In the cross-study or/and cross-platform comparisons, we normalized the batch effect using the 'combat' package from R. Differentially expressed genes (DEGs) were calculated as counts per million (CPM) using featureCount and algorithms from "DESeq2". This strategy provided both quantification and statistical inference of systematic changes between conditions (with at least three replicates). We had generated various transcriptomes from A375 cells (e.g., A375-untransfected, A375-scambled transfection control, A375-knockdowns targeting HERV-K(HML2) Rec using three distinct RNAi constructs, Rec-KD1,2,3). We performed similar studies in SK-MEL28 cell lines. As an additional control, and to increase statistical power for DEGs identified in our transcriptomes upon Rec-KD, we also used publicly available A375 RNA-seq datasets (GSE110948). To analyze single-replicate RNA-seq datasets, we utilized the "GFOLD" algorithm for normalization, which calculates a variance of fold-changes from unreplicated RNA-seq data. Canonical pathways and biological function of the differentially expressed genes (DEGs) were further subjected to KEGG and Gene Set Enrichment Analyses (GSEA).

2.6. Single-cell RNA-seq data processing

We got the single-cell count matrix from GSE72056. We calculated the activity of genes in every cell at TPM expression levels. We considered samples expressing over 1000 genes with expression levels exceeding the defined threshold (Log2 TPM > 1). We used Seurat_3.1.1, a package from R to normalize the datasets at the logarithmic scale using "scale.factor = 10000". After normalization, we calculated scaled expression (z-scores for each gene) for downstream dimension reduction. We used the original annotations of datasets to classify the malignant, non-malignant, and heterogenous cell-types. To define cell population clusters, we employed the FindClusters function of "Seurat" using "PCA" as a reduction method. The specific markers for each cluster identified by "Seurat" were determined by the "FindAllMarkers" function, using "roc" as a test for significance. This provided us two lists of gene sets, 1) Malignant vs. Non-Malignant differentially expressed genes and 2) genes differentially expressed between MITF-High and MITF-low tumors, which we applied for comparing the DEGs from Rec-KD RNA-seq. Feature plots, violin plots, and heatmaps were constructed using default functions, except for the color scale that was set manually. The annotated cells were re-clustered using the methodologies described above and visualized on the UMAP coordinates.

Singh et al. (2020)

2.7. ATAC-seq and ChIP-seq data analyses

ATAC-seq and ChIP-seq raw datasets in sra format were downloaded from the listed studies and converted to fastq format using sratools function fastq-dump –split-3. Fastq reads were mapped against the hg19 reference genome with the bowtie2 parameters: --very-sensitive-local. All unmapped reads with MAPQ < 10 and PCR duplicates were removed using Picard and samtools. All the ATAC-seq peaks were called by MACS2 with the parameters –nomodel -q 0.01 -B. Blacklisted regions were excluded from called peaks (https://www.encodeproject.org/annotations/ENCSR636HFF/). To generate a set of unique peaks, we merged ATAC-seq peaks using the mergeBed function from bedtools, where the distance between peaks was less than 50 base pairs. We then intersected these peak sets with repeat elements from hg19 repeat-masked coordinates using bedtools intersectBed with a 50% overlap. To calculate the enrichment over the given repeat elements, we first extended 5KB upstream and 5KB downstream coordinates from the left boundary (TSS) of respective elements in a strand-specific manner. These 10KB windows were further divided into 100 bps bins, and tags (bedGraph output from MACS2) were counted in each bin. Tag counts in each bin were normalized by the total number of tags per million in given samples and presented as Counts Per Million (CPM) per 100 bps. We averaged CPM values for each bin between replicates before plotting the figures.   To find the MITF binding motifs, 25 bps sequences were extended from either side of ChIP-seq peak summits. The extended sequences were analyzed by the RSAT tool (http://rsat.sb-roscoff.fr/). The TF binding motifs were calculated from JASPAR libraries of human TF motifs.

2.8. Cell invasion assay

The cell invasion assays were performed using transwell chambers (8 μm pore size; Corning Costar) according to the vendor's instructions. Briefly, the insert of the wells was first coated with 50 μl BD matrigel. A375 cells were resuspended with RPMI1640 medium containing 0.5% FCS to reach a concentration of 5×10^5 cells/ml. Then 100 μl cells from each group were seeded into the upper chamber of the insert, with adding DMEM containing 10% FCS to the lower chamber. After incubation at 37 °C in a humidified atmosphere of 5% CO_2 for 24 h, non-invaded cells were removed with gentle swabbing while the invaded ones were stained. Then images were taken with microscopy. Cell numbers were calculated with ImageJ.

2.9. Accession of datasets used in this study:

Encode chromHMM datasets:
ftp://hgdownload.cse.ucsc.edu/goldenPath/hg19/encodeDCC/wgEncodeBroadHmm
Encode TF ChiP-seq datasets:
ftp://hgdownload.cse.ucsc.edu/goldenPath/hg19/encodeDCC/wgEncodeSydhTfbs
GSE60663: H3K27Ac Primary Melanoma
GSE60663: H3K4Me3 Primary Melanoma
GSE60663: MITF ChIP-seq Primary Melanoma
GSE60664: RNA-seq Primary Melanoma
GSE110948: A375 RNA-seq
GSE46817: RNA-seq of 7 distinct melanoma lines
GSE46805: RNA-seq Melanocyte BRAF overexpression
GSE50681: MITF ChIP-seq Melanoma and Melanocyte post-treatment
GSE82330: ATAC-seq Melanoma A375
GSE72056: Patient's Single cell RNA-seq data

3. Results

3.1. LTR5_Hs loci form active chromatin in cancer and pluripotent cell lines

The regulatory region of the youngest subfamily of HERV-K is the human-specific long terminal repeat 5 (LTR5_Hs). To dissect the regulation of HERV-K transcription in different cell types, we adopted a systematic genome-wide approach using publicly available data. For this, we investigated

Singh et al. (2020)

the chromatin state at ~600 full-length LTR5_Hs sequences using the uniquely mappable reads in 9
different cell types from the ENCODE project [50]. Analysis of the chromatin states suggested that a
substantial fraction (n = 227) of LTR5_Hs sequences overlap with active or permissive chromatin
states (Figure 1A). When compared to differentiated somatic cells, these LTR5_Hs genomic loci
showed features of active ('open') chromatin state in human embryonic stem cells (H1_ESCs) and in
cancer cell lines (Figure 1A).

Various transcription factors (TFs) have been reported to bind LTR5_Hs/HERV-K, including the
pluripotency factors such as OCT4 [12, 51] or the melanocyte inducing TF (MITF) [52]. In a systematic
approach, we mapped the peaks from 142 ChIP-seq TF datasets from 7 cell lines (ENCODE) over
LTR5_Hs genomic loci (n ~ 600). Our integrative analysis of various TF occupancy revealed that
LTR5_Hs recruits significantly more TFs in H1 embryonic stem cells (H1_ESCs) and leukemia K562
cells, compared with HepG2, GM12878 or HUVEC cells (Figure 1B), which may indicate elevated
levels of HERV-K transcription in embryonic stem cells and cancer cells. While in H1_ESCs, LTR5_Hs
was bound prominently by pluripotency specific TFs (e.g., NANOG) (Figure 1B-C), the cancer line
(K562) displays the binding of multiple TFs that control the cell cycle and proliferation (e.g., c-MYC,
MAX, COREST, P300, PU1, GATA2, and JUND) (Figure 1B-C and Table S1). Intriguingly, NANOG
binding over LTR5_Hs is overrepresented in 3iL naïve conditions (Figure 1C). The observed HERV-
K virion particles in the human blastocyst might be the products of HERVK driven by NANOG [12].
The active chromatin states and TF bindings over LTR5_Hs in cancer lines might be associated with
the oncogenic phenotype.

3.2. MITF regulated LTR5_Hs/HERV-K expression is a hallmark of the 'proliferative' type of melanoma

It has been previously reported that in melanoma, the melanocyte/melanoma-specific isoform,
MITF-M may bind LTR5_Hs, and promotes the expression of HERV-K transcripts [52]. MITF(-M) is
a critical transcription factor associated with melanoma progression [53, 54], and is a sensitive marker
of melanoma invasiveness; highly expressed in the melanocytes and the proliferative state of
melanoma (e.g., MITF-high) but lowly expressed in the invasive state (e.g., MITF-low) [55].

To investigate how HERV-K is controlled in melanoma, we mined ChIP-seq and RNA-seq data
from melanocytes, various melanoma cell lines, and primary melanoma cultures (Methods).
Importantly, characterizing primary melanoma cultures according to their MITF levels (Figure 1D)
revealed a positive correlation between MITF-regulated LTR5_Hs/HERV-K transcription and
invasiveness (Figure 2A). To see if additional transposable elements (TEs) respond to melanoma
proliferative/invasive status, we extended the systematic analysis over all of the TE families. This
analysis revealed that the expression of certain ERVs, including LTR5_Hs/HERV-K, LTR2C, and
LTR13, was correlated with proliferative melanoma associated genes (Figure S1A). Nevertheless,
when compared to the overall expression of other TE families, LTR5_Hs/HERVK was the most
abundantly expressed TE family and highly correlated with MITF level (rho = 0.43, corrected p-value
< 0.2e-9) in the melanoma primary cultures (N=10) (Figure 2B). These data suggest that
LTR5_Hs/HERV-K expression can be used as a marker that correlates negatively with melanoma
invasiveness. Next, we characterized LTR5_Hs loci activated in the proliferative type of melanoma
cells by layering ChIP-seq occupancy signals of H3K4Me3, H3K27Ac, and MITF (Figure 2C and S1B-
C). We found that MITF binding at LTR5_Hs loci coincides with the presence of the canonical E-box
motif (CA(C/T)GTG), which is recognized by MITF as well as c-MYC, and MAX TFs [56] (Figure 2C).

We also investigated the regulation of HERV-K transcription in the BRAF^{V600E} mutant, an
invasive type of melanoma. PLX4032, a commonly used BRAF^{V600E} inhibitor in the clinic, enhances
the expression of MITF by several folds that revert melanoma from an invasive to the proliferative
state [59]. To examine how PLX4032 treatment affects HERV-K expression, we reanalyzed the
publicly available MITF ChIP-seq datasets of primary melanocytes and a BRAF^{V600E} mutant
melanoma cell line (COLO829) with and without PLX4032 treatment (GSE50681). We observed that
PLX4032 treatment results in a substantial enrichment of MITF ChIP-seq peaks over LTR5_Hs
elements (123 unique loci were bound) (Figure 2D). When compared to other TE families, the gain of
MITF binding occurred specifically over LTR5_Hs (Figure 2E). Importantly, while MITF is expressed
in healthy pigmented cells, we found no elevated MITF ChIP-seq signal over LTR5_Hs elements upon

Singh et al. (2020)

270 PLX4032 treatment in normal, pigmented melanocytes (Figure 2D). Thus, MITF binding to LTR5_Hs
271 is specific to the proliferative state of melanoma.
272   Collectively, the MITF binding may lead to LTR5_Hs activation, not typical in healthy
273 melanocytes, but in the proliferative type of melanoma, thus can be considered as a sensitive marker
274 of the proliferative vs. invasive phenotype of melanoma.

3.3. Depleting HERV-K-Rec induces an EMT-like process in A375 melanoma cells

The ERVK6 locus (located on the Chr7 p22.1) produces both retroviral ORFs (Gag and Env) and
the accessory HERV-K product, Rec [13, 28]. To see how this specific locus responds to MITF in
proliferative (MITF-High), invasive (MITF-Low) primary melanoma lines, we analyzed a published
dataset for H3K27ac and MITF ChiP-seq (GSE60663), and also included the proliferative type A375
melanoma cell line in the analysis (GSE82330). This locus showed a higher H3K27Ac signal in both
proliferative MITF-High and A375 cells relative to MITF-Low cells, and also a significant ATAC-seq
signal in A375 (Figure 3A), indicating that this locus is preferentially active in a proliferative type of
melanoma. Moreover, ATAC-seq and active histone mark analysis (e.g., H3K27ac and H3K4Me1)
support the accessibility of multiple LTR5_Hs in A375 (Figure 3B-C), suggesting that A375 cell line
might be an adequate model for this type of melanoma.
To decipher the underlying difference of the invasiveness of various melanoma cells and the
potential role of Rec in melanoma, we first used an RNAi approach to deplete Rec expression in two
melanoma cell lines (e.g., A375 and SKMel-28) of different invasiveness. A375 and SKMEL-28, both
wild type for the tumor suppressor PTEN and carry the BRAF^{V600E} mutation, but A375 have a higher
proliferation rate, whereas SKMEL-28 is more invasive [62]. To knockdown (KD) Rec, we designed
three RNAi constructs. These constructs were delivered using Sleeping Beauty as a stable vector [63],
and also expressed a fluorescence reporter (GFP). To generate stable KD cell lines for both A375 and
SKMel-28, the KD constructs were transfected together with the SB100X transposase [64]. The stable
KD-Rec lines were selected in two-rounds by FACS sorting for the GFP-reporter signal. Real-time
qRT-PCR showed that Rec transcript levels were strongly depleted in each of the three independent
KD cell lines isolated for both A375 and SKMel-28 (Figure 3D). Notably, as a side effect, knocking
down Rec also led to an increased level of alternative retroviral HERV-K transcripts, Gag and Env in
SKMel-28, and A375, respectively (Figure 3E).
To detect global transcriptional changes in an unbiased way, we used RNA-seq analysis of the
stable KD-Rec cell lines. The study of individual HERV-K genomic loci revealed that our KD strategy
(using three biological replicates) significantly affected seven Rec expressing loci in KD-Rec_A375
cells (Figure 3F). SKMel-28, by contrast, has only three HERV-K loci expressing Rec (not shown) [28].
Having validated our Rec KD lines, we performed a global analysis of differentially expressed
genes (DEGs) (|2|-fold and FDR < 0.05, (Figure S1E, Table S1, see Methods) between the KD lines
and the parental cell lines. The list of the DEGs was different between the A375 and SKMel-28 knock-
downs, which could be in part explained by the distinct patterns of active HERV-K loci in the two
melanoma lines. In KD-Rec_SKMel-28, DEGs (Table S1) did not include MITF and were not
significantly enriched for any Gene Ontology (GO) categories, and formed no discernible pattern,
suggesting that knocking-down Rec in this particular (invasive) melanoma line was not instructive
of biological process alterations. The smaller number of Rec expressing HERV-K loci and the
antagonistic elevation of the alternative retroviral product (e.g., Rec) in KD-Rec_SKMel-28 might, at
least partially, explain the lack of distinct pattern.
In contrast, GO analysis of 1120 DEGs in KD-Rec_A375 revealed that most of the dysregulated
genes were associated with cell communication, followed by the cell cycle, cell proliferation (Figure
4A). Narrowing down the list of DEGs by excluding genes of the category of system development,
the remaining genes were enriched in clusters of cell differentiation, stem cell differentiation, cell
death, and system development. Genes in the stem cell differentiation and system development
categories could be linked to the WNT signaling pathway. Among DEGs in KD-Rec_A375, we
identified genes involved in an epithelial-mesenchymal transition (EMT)-like process, which has
been observed to play a critical role in phenotype switching in the proliferative to invasive state [33,
66, 67]. Intriguingly, this set of DEGs included matrix metallopeptidase 2 (MMP2), a classical marker
of cell invasion and additional four genes from the list of the 16 critical markers of the canonical EMT

process [65] (e.g. N-Cadherin (CDH2), zinc finger protein (SNAI2), forkhead box C2 (FOXC2) and
goosecoid Homeobox (GSC) (Figure 4B-C).
Nevertheless, as the melanocytes and melanoma cells differ from epithelial and mesenchymal
cells, respectively, the EMT-like process in melanoma phenotype switching is characteristically
different from the canonical EMT process [33]. In melanoma metastasis, the TF network switches to
the melanoma-specific $SNAI2^{low}/ZEB2^{low}/ZEB1^{high}$ phenotype [69-71], that we also observed in our
KD-Rec_A375 cells (Figure 4B-C), suggesting that Rec modulates the melanoma-specific EMT-like
phenotype switching. The elevated level of EMT markers in KD-Rec_A375 cells suggests that Rec
inhibits the EMT-like process.
Beside canonical EMT markers, we also observed the upregulation of further factors in KD-
Rec_A375 cells, known to act as inducers of the EMT-like process. For example, non-canonical WNT
signaling via WNT5A and GSK3B, which is markedly upregulated in KD-Rec_A375 cells (Figure 4D),
is critical for the switch to an EMT-like invasive phenotype when melanoma gets relapsed [72, 73].
Furthermore, the most extensively studied inducer of EMT, TGF-β [74, 75], reported to control SNAI2
[76], was also upregulated in our KD-Rec_A375 transcriptomes (Table S1).
In addition, among the top DEGs, we observe a strong upregulation of several members of the
SPANXC (sperm protein associated with the nucleus in the X chromosome) family of genes (Figure
4D-E), which are typically expressed in the germline. The upregulation of SPANXB1/2, clinically
relevant cancer-testis antigens, has been previously reported in melanoma [75] and was also
validated by qRT-PCR in our KD-Rec_A375 samples (Figure 4E). Simultaneous upregulation of
multiple genes physically located within the SPANX (B/C) chromosomal cluster suggests an
epigenetic disturbance affecting the entire locus. Indeed, the epigenetically sensitive Polycomb
repressor complex 2 (PRC2) has been implicated in maintaining SPANXB in a transcriptionally silent
state in differentiated cells [76]. Our data suggest that Rec may exert directly or indirectly an effect
on the epigenome of melanoma, an aspect that we did not follow up in this study.
Importantly, in our KD-Rec_A375 cells, we detected a robust transcriptional decrease in the level
of MITF-M, the governor of the melanocyte differentiation program (Figure 5A), confirmed by qPCR.
In addition to MITF, we observed a decreased expression of further melanocyte differentiation
markers (e.g., TYR, PMEL) [77] (Figure 5B).
Collectively, our transcriptome analysis in KD-Rec_A375 cells suggests that Rec might function
as a cellular factor inhibiting the melanoma phenotype switching to the invasive state.
354

3.4. Rec is a suppressor of cell invasion in melanoma

Our $SNAI2^{low}/ZEB2^{low}/ZEB1^{high}$ phenotype, attributed to aggressive melanoma, suggested that
knocking down Rec, via induction of an EMT-like process, might affect the invasiveness of A375 cells.
Further on this line, the decreased level of MITF, and the transcriptome-wide DEGs of invasive vs.
proliferative melanoma markers also predicted enhanced invasiveness of KD-Rec_A375 cells. To test
these predictions, we performed a trans-well invasion assay to compare the invasiveness of KD-
Rec_A375 cells to their parental line. The results showed that all three KD-Rec_A375 lines exhibited
significantly elevated invasiveness (Figure 5C-D). The enhanced metastatic potential of the
melanoma upon Rec KD supported the idea that Rec functions as a suppressor of the EMT-like
transition process, which modulates the invasiveness of melanoma.

3.5. The KD-Rec_A375 in vitro model mimics the MITF type malignancy in patients

Next, we sought to examine whether some of the genes dysregulated upon Rec KD in A375 cells
were also differentially regulated in malignant cell types. To assess this, and to monitor the dynamics
of Rec-regulated genes in the progression of tumors, we performed a single comprehensive cell
(sc)RNA-seq analysis (n=4645) derived from melanoma patients (n=19) [78]. Congruent with the
original study [78], our analysis with default parameters from the 'Seurat' algorithm distinguishes
clusters of non-malignant and malignant cell types (Figure 6A). The five distinct malignant cell types
were marked by high expression levels of MAGEC2, MITF, APOE, VIM, and SPP1, respectively
(Figure 6A-B).

Singh et al. (2020)

To see how faithfully our KD-Rec_A375 model mimics in vivo processes observed in melanoma
patients, we first compared the ~1500 DEGs identified between non-malignant and malignant cell
types (all five), to those identified in our KD-Rec_A375 experiments (Figure 7A). We found that ~200
genes overlapped between the two analyses in the opposite direction (Figure 7B), a significant
enrichment over the random expectation (p-value <0.00001, see Methods).

Importantly, the expression levels of MITF, APOE, and SPP1 (but not MAGEC2, and only
modestly VIM) were also significantly depleted upon Rec knockdown in A375 cells, suggesting that
the KD-Rec_A375 might recapitulate certain aspects of the melanoma process in patients (Figure 7C-
D). The affected MITF, APOE (Apolipoprotein E), and SPP1 (osteopontin, cytokine) highlight
different aspects of melanoma progression, crosstalk of differentiation, angiogenesis [79] and
inflammation [80], respectively.

As MITF directly regulates HERV-K (via binding LTR5_Hs), we next focused on the MITF
malignancy sub-cluster of the scRNA-seq dataset [78] (Figure 6B). We took the top 100 MITF-
correlated genes and determined their expression in our Rec-KD_A375 data. This approach revealed
a highly significant (Wilcoxon-test, p-value < 2e16) reverse differential expression pattern between
MITF-type malignancy and Rec-KD_A375 (Figure 7E), indicating an anti-correlation between the
expression of MITF-target gene expression and Rec depletion. Thus, our KD-Rec_A375 model
appears to recapitulate MITF-dependent aspects of melanoma in patients.

4. Discussion

HERV-K resembles "complex" retroviruses such as HIV-1 by its ability to encode multiple
accessory proteins, including Rec, through alternative splicing of its Env transcript. While Env has
the typical structure of a transmembrane protein (e.g., extracellular, transmembrane, cytoplasmic
domains), Rec lacks the majority of the domains required for a membrane protein function (Figure
8A). Still, it carries nuclear localization/export signals [3, 7]. Although Env and Rec are utterly
different in sequence and function, our results suggest that Rec, like Env [39], exerts a modulatory
effect on the EMT-like process in cancer progression. However, while Env affects signaling via
ERK1/2 activation, through the cytoplasmic tail of the transmembrane Env protein [39], this domain
is missing from Rec (Figure 8A). Rec, the splice variant of Env, inhibits the EMT-like process by
affecting the expression levels of critical determinants of the EMT-like process, and certain upstream
regulators. Thus, the two factors appear to act in the opposite direction. While Env overexpression
(both RNA and protein) may promote tumor progression [39], depletion of Rec seems to result in a
comparable phenotype. Curiously, similar to Env, the retroviral gene product, Gag also increases the
tumorigenic potential of melanoma cells [38], raising the possibility that retroviral and accessory
genes might generally act inversely, deciding on a long-lasting debate on the potential oncogenicity
of a retroviral versus accessory genes of HERV-K.

In about 50% of melanomas, the protein kinase BRAF is mutated (BRAF^{V600E}) [84]. MITF is
considered as a crucial regulator of melanoma invasiveness [85], and plays a critical role in controlling
BRAF^{V600E} melanoma [86, 87]. Accordingly, the expression of MITF has been widely used as a marker
to distinguish between the proliferative and invasive types of melanoma cells [61, 85, 86, 88]. Reduced
MITF expression generates invasive melanomas, with tumor-initiating properties [86]. Importantly,
our KD-Rec_A375 in vitro system appears to recapitulate a subpopulation of melanoma cells (MITF
malignancy) in human patients. Our data suggest that MITF activates LTR5_Hs only in melanoma
(not in melanocytes), but Rec expression specifically marks the proliferative type of melanoma cells.
Importantly, the proliferative phase of melanoma is still a better controllable stage of melanoma
progression, by contrast to the invasive stage that gets frequently relapsed. If Rec expression is
involved in maintaining a delicate balance between cell proliferation and invasion, as this study
suggests, then Rec activation might be used as a sensitive marker to distinguish between the
proliferative vs. invasive phenotypes of melanoma, not expressed in normal melanocytes.

To control cell plasticity during melanomagenesis, the EMT-like network operates upstream of
MITF. In cells harboring the SNAI2^{low}/ZEB2^{low}/ZEB1^{high} phenotype, the depleted SNAI2 and ZEB2,
expressed in normal melanocytes and function as tumor-suppressors, would fail to activate the MITF-
dependent melanocyte differentiation program [34, 87]. In parallel, the upregulated ZEB1 induces an
invasion-associated cascade that operates in concert with downregulating MITF.

Singh et al. (2020)

Our study reveals that depletion of Rec in a BRAF^{V600E} mutant melanoma cells (A375) results in
a decline in the MITF level. Thus, our KD-Rec_A375 cells suggest a model whereby a MITF-
dependent regulatory loop is disrupted through (i) decreased MITF binding to LTR5_Hs, which leads
to (ii) reduced Rec levels and (iii) a further decline in MITF level, contributing to (iv) escalated tumor
aggressiveness (Figure 8B). The observed cross-regulation between Rec and MITF reveals an
inhibitory effect of Rec on the invasive transformation of melanoma. Our data suggest that a high
level of Rec expression marks the proliferative (still controllable) state of melanoma cells, and opposes
the molecular switch to the invasive state. Future research is required to decipher the exact
mechanism of Rec control over MITF, and to test whether therapeutically induced Rec transcription,
similar to enhanced MITF expression (e.g., PLX4032 treatment), results in a reversion of the invasive
melanoma phenotype.

Under healthy physiological conditions, only a limited number (n=18, or often less) of HERV-K
loci appear to be active [13], whereas, in melanoma patients/cell lines, the number of transcribed
HERV-K loci is higher (n~24) [28], indicating that specific HERV-K loci are transcribed only under
pathologic conditions. Several loci express multiple HERV-K-derived products (both core and
accessory replication factors). Although forced expression of MITF-M in non-malignant cells has been
shown to enhance the levels of numerous HERV-K derived transcripts (e.g., Gag, Env, and Rec) [52],
the activated HERV-K loci might differ in both intensities of expression and the composition of their
HERV-K-derived transcripts. In line with this, HERV-K loci are variably activated and in various
combinations in melanomas [28, 89], suggesting that variability in the amounts of expressed HERV-
K products might contribute to melanoma heterogeneity [22, 27], and possibly the progression and
outcome of the disease.

It has been widely discussed how precisely melanoma can be recapitulated in animal models.
Interestingly, our analysis of single-cell transcriptome data from multiple human patients suggests a
distinct separation of proliferative and invasive types of cells within a patient. In contrast, these two
phases co-occur within the same melanoma cell in mice [88]   As Rec is hominid-specific, our study
also has an evolutionary perspective that might explain certain species-specific features of melanoma
progression in human patients, and help us to understand the limitation(s) of using animal models
in melanoma studies.

Singh et al. (2020)

Figure legends

Figure 1. (A) Stacked barplots show the chromatin state (chromHMM) of the LTR5_Hs
genomic loci (human Refseq genome) across 9 cell types. (B) Heatmap summarizing the
transcription factor (TF) occupancy (ChIP-seq peaks, ENCODE) over individual LTR5_Hs
loci in five human cell lines. This plot includes 234 out of 615 LTR5_Hs loci annotated in the
human Refseq genome, which is occupied by at least one ChIP-seq peak. Each row
represents an individual locus of LTR5_Hs. Each column shows TF ChIP-seq peak
occupancy of a different cell line. Grey denotes the absence, whereas black is the presence
of peaks in a given locus. Various TFs encompassing LTR5_Hs loci are manually annotated
in a given cell line. The cluster of TFs binding over a subset of LTR5_Hs loci is shown in the
red box. C. Line plots showing the distribution of normalized ChIP-seq tag counts of cMYC
and NANOG in K562 and naive ESCs, respectively. The ChIP-seq tag counts are calculated
relative to primed H1_ESCs, in a 3kb genomic window at the left boundaries of LTR5_Hs.
(D) Characterizing primary melanoma cultures according to their MITF expression levels.
Barplot is showing the level of MITF expression (Log2 TPM) in primary melanoma cultures
(n=10). According to their MITF expression level, cultures are defined as proliferative
melanoma (MITF-High) and invasive melanoma (MITF-Low) in vitro models.

Singh et al. (2020)

Figure 2. (A) Barplot is showing the LTR5_Hs/HERV-K-int expression (Log2 CPM) levels in proliferative melanoma (MITF-High) and invasive melanoma (MITF-Low) cell lines (GSE60664). (B) A Scatter plot visualizes the average expression (Log2 CPM) (X-axis) and standard deviation from the mean (Y-axis) of transposable element (TE) families in various primary melanoma cultures (n=10). (C) Line plot showing the MITF ChIP-seq signal occupancy in proliferative melanoma (MITF-High) and invasive melanoma (MITF-Low) cell lines averaged over all of the LTR5_Hs loci. The identified E-Box motif in MITF bound LTR5_Hs in A375 cells is shown. Note that the E-box motif was found using the sequences of LTR5_Hs loci with MITF peaks. (D) The effect of PLX4032 treatment on MITF binding at LTR5_Hs loci. Upper panel: Line plot showing the MITF ChIP-seq signal averaged over LTR5_Hs loci in primary melanocytes and melanoma cells with and without the treatment of PLX4032. Lower panel: Heatmap of MITF ChIP-seq signals. Note: MITF occupies LTR5_Hs genomic loci upon PLX4032 treatment, specifically in melanoma, but not in melanocytes (over 100 loci). (E) Barplot is showing the number of MITF peaks detected over the genomic loci of various TE families in melanoma cells treated with PLX4032. This plot indicates only those families which have at least 5 peaks over the individual locus.

Singh et al. (2020)

Figure 3. (A) Genome browser tracks showing the H3K27Ac, ATAC-seq, and MITF signals at a specific LTR5_Hs-HERVK locus (ERVK6, known to produce Rec) in MITF-High primary melanoma cultures and A375 cell line (proliferative) and MITF-low (invasive) melanoma primary cultures. (B) Normalized ATAC-seq (nucleosome-associated) signal averaged at LTR5_Hs genomic loci in A375 cells. (C) Normalized ChIP-seq signal of H3K4Me1 and H3K27Ac histone marks (corresponding to active chromatin states) averaged over LTR5_Hs genomic loci in A375 cells. (D) Knocking down (KD) Rec in A375 and SKMEL-28 cells using RNA interference. Stable KD lines were selected in two-rounds by FACS sorting for the GFP-reporter signal. Barplots are showing the normalized RT-qPCR quantification of Rec mRNA levels with three KD-Rec constructs. (E) Knocking down Rec affects the transcription of alternative HERV-K products. Barplot is showing the effect of knocking down Rec (KD-Rec) on the expression of alternative HERVK products, determined by real-time qPCR. Note: KD-Rec led to an increased level of Gag in SKMel-28 (all the three biological replicates), whereas Env in A375 (KD1 only). (F) Knocking down Rec in A375 affects at least seven genomic HERV-K loci. Combined barplots showing the specificity of the RNAi strategy against the Rec subunit. Relative quantification of HERV-K Rec genomic loci compared with negative and scrambled controls indicates that all three KD constructs significantly affected at least 7 Rec expressing loci in A375 cells.

Singh et al. (2020)

Figure 4. (A) Barplots of significant Gene Ontology (GO) Biological Process terms enrichment analysis of DEGs in KD-Rec_A375. Values are negative logarithms of the corrected p-values (Benjamini Hochberg FDR). (B) Barplots representing means ± SEM from RNA-seq expression values (Log2 TPM) for selected DEGs in KD-Rec_A375 (e.g., Matrix metallopeptidase 2 (MMP2), a classical marker of invasion; N-Cadherin (CDH2), Zinc finger protein SNAI2 (SNAI2), Forkhead box C2 (FOXC2) and Goosecoid Homeobox (GSC) are among the vital gene regulators of the canonical EMT transition. (C) Barplots representing means ± SEM from RNA-seq expression values (Log2 TPM) for DEGs in KD-Rec_A375 vs. control, involved in the EMT-like process in melanoma [34] (e.g., SNAI2, ZEB2, and ZEB1). (D) Barplots representing means ± SEM from RNA-seq expression values (Log2 TPM) for selected DEGs in KD-Rec_A375 vs. control, known as inducers of the EMT-like process (e.g., WNT5A and GSK3B) and SPANXC (sperm protein associated with the nucleus in the X chromosome). (E) Barplots are showing the effect of knocking down Rec in A375 vs. control on SPANXB1/B2 genes, detected as DEG in RNA-seq experiments, and validated using real-time qPCR.

Singh et al. (2020)

Figure 5. (A) Quantifications transcriptional variation of MITF-M by KD-Rec in A375 cells using RNA-seq and by real-time qPCR. (B) Barplot representing means ± SEM from RNA-seq expression values (Log2 TPM) for melanocyte markers (e.g., TYR and PMEL) upon Rec-KD in A375. (C) Representative inverted light microscope images of trans-well invasion assay performed in A375 melanoma cells upon knocking-down Rec by three KD constructs (KD1-3). UT, untransfected control. Scale bars, 400 μm. (D) Quantification of cells invading in the trans-well invasion assay (C). ImageJ counted the number of invaded cells. Columns indicate the average number of invaded cells per field from two independent experiments. Error bars represent the means±s.d. of two independent experiments. Mann Whitney test; **P ≤ 0.01, ****p ≤ 0.0001 (See methods for details).

Singh et al. (2020)

Figure 6. (A) Various plots with a maximum of thirteen distinct clusters are visualized on UMAP coordinates using 10XGenomics scRNA-seq [78] and their original annotations. The left panel shows the Malignant and Non-Malignant cell populations, whereas the right panel is displaying all the clusters obtained that were classified using their original annotations. (B) Feature plots based on the UMAP plot shown on (A) visualizing the expression of selected markers used to identify the distinct cell-types in the melanoma population. The color intensity gradient indicates the expression of the depicted marker gene. Each dot represents an individual cell. Dots in green denote lower, whereas in dark red, a higher level of gene expression in a given single cell. This resulted in five distinct malignant cell types, marked by MAGEC2, MITF, APOE, and Ki67, respectively (See Fig. 7A for SPP1 expression).

Singh et al. (2020)

Figure 7. (A) Heatmap visualization of scaled expression [log TPM (transcripts per million)] values of a distinctive set of ~ 1500 genes (log2 fold change > one and > 50% of cells expressing either in healthy or malignant cells), which are differentially expressed in scRNA-seq data. Colour scheme is based on Z-score distribution from -2.5 (gold) to 2.5 (purple). Left margin color bars highlight representative gene sets specific to the respective clusters. (B) The Violin plot visualizes the density and distribution of expression (Log2 TPM values) of ~ 200 DEGs upon Rec-KD in A375 cells, which are common in the set of genes shown in the figure above (C). (p-value calculated by Wilcoxon test). (C) Violin plots visualize the density and distribution of expression of distinct marker genes for malignant melanoma clusters. (D) Barplots with +/- SEM show the down-regulation of MITF, SPP1, and APOE upon knockdown of Rec in A375 cells. (E) Violin plot showing the comparison of previously identified gene set as "MITF-high," consisting of MITF itself and the top 100 genes correlated with MITF expression between malignant and non-malignant cells (blue) and upon Rec-KD in A375 by individual shRNA constructs (gold). (p-value calculated by Wilcoxon test).

Singh et al. (2020)

Figure 8. (A) Schematic structure of the HERV-K-derived alternative splice products, Env and Rec, and their effect on the endothelial-mesenchymal transition (EMT)-like process of cancer progression (EMT(L). The signal peptide (SP); transmembrane subunit (TM); Splice acceptor (Sa); Splice donor (Sd). (B) Modeling the interaction loop between MITF and LTR5_Hs and Rec in melanoma. MITF binds to LTR5_Hs and drives Rec expression. Rec marks the proliferative stage melanoma cells and, similarly to MITF, inhibits the endothelial-mesenchymal transition (EMT)-like process (EMT(L). Depletion of Rec results in MITF downregulation and the failure to maintain the balance between cell proliferation and invasion.

Singh et al. (2020)

Figure S1. (A) (Left panel) Ranked correlation matrix of genes and 42 transposable elements (TE) families, which are differentially expressed between proliferative and invasive primary melanoma cultures. Genes and TE families are manually annotated. Color codes on the left side represent the unsupervised clusters of most variable genes and TEs across the samples. On the right panel, selected genes are listed that are positively or negatively correlated with LTR5_Hs expression across the melanoma primary cultures. (B) Distribution of ChIP-seq signal of the H3K4Me3 histone mark occupancy, corresponding to active promoters averaged over LTR5_Hs loci in MITF-High (proliferative) vs. MITF-Low (invasive) state of primary melanoma cultures. (C) Distribution of ChIP-seq signal of the H3K27Ac histone mark occupancy (lower panel) averaged over LTR5_Hs loci in MITF-High (proliferative) vs. MITF-Low (invasive) state of primary melanoma cultures. (D) Heatmap displays the DEGs upon KD-Rec in A375 cells.

Author Contributions: Conceptualization, M.S, H.C., and Z.I.; methodology, H.C.; analyses, M.S.; writing, original draft preparation, Z.I., M.S, CF; writing—review and editing, all authors; visualization, A.K.; supervision, Z.I. All authors have read and agreed to the published version of the manuscript.

Funding:

Acknowledgments: H.C. was supported by the China Scholarship Council (CSC). M.S. was supported by a presidential postdoctoral fellowship from Cornell University. Z.I. was funded by European Research Council, ERC Advanced [ERC-2011-ADG 294742]. CF laboratory R35 GM122550 grant from the National Institutes of Health

Conflicts of Interest: Authors declare no conflict of interests

Singh et al. (2020)

References:

1. Lander ES, Linton LM, Birren B, Nusbaum C, et al., 2001. Initial sequencing and analysis of the human genome. *Nature,* 409: 860-921.

2. Weiss RA, 2006. The discovery of endogenous retroviruses. *Retrovirology,* 3: 67.

3. Lower R, Tonjes RR, Korbmacher C, Kurth R, et al., 1995. Identification of a Rev-related protein by analysis of spliced transcripts of the human endogenous retroviruses HTDV/HERV-K. *J Virol,* 69: 141-9.

4. Armbruester V, Sauter M, Krautkraemer E, Meese E, et al., 2002. A novel gene from the human endogenous retrovirus K expressed in transformed cells. *Clin Cancer Res,* 8: 1800-7.

5. Magin C, Lower R, and Lower J, 1999. cORF and RcRE, the Rev/Rex and RRE/RxRE homologues of the human endogenous retrovirus family HTDV/HERV-K. *J Virol,* 73: 9496-507.

6. Boese A, Sauter M, Galli U, Best B, et al., 2000. Human endogenous retrovirus protein cORF supports cell transformation and associates with the promyelocytic leukemia zinc finger protein. *Oncogene,* 19: 4328-36.

7. Yang J, Bogerd HP, Peng S, Wiegand H, et al., 1999. An ancient family of human endogenous retroviruses encodes a functional homolog of the HIV-1 Rev protein. *Proc Natl Acad Sci U S A,* 96: 13404-8.

8. Denne M, Sauter M, Armbruester V, Licht JD, et al., 2007. Physical and functional interactions of human endogenous retrovirus proteins Np9 and Rec with the promyelocytic leukemia zinc finger protein. *J Virol,* 81: 5607-16.

9. Kaufmann S, Sauter M, Schmitt M, Baumert B, et al., 2010. Human endogenous retrovirus protein Rec interacts with the testicular zinc-finger protein and androgen receptor. *J Gen Virol,* 91: 1494-502.

10. Hanke K, Hohn O, Liedgens L, Fiddeke K, et al., 2013. Staufen-1 interacts with the human endogenous retrovirus family HERV-K(HML-2) rec and gag proteins and increases virion production. *J Virol,* 87: 11019-30.

11. Costoya JA, Hobbs RM, Barna M, Cattoretti G, et al., 2004. Essential role of Plzf in maintenance of spermatogonial stem cells. *Nat Genet,* 36: 653-9.

12. Grow EJ, Flynn RA, Chavez SL, Bayless NL, et al., 2015. Intrinsic retroviral reactivation in human preimplantation embryos and pluripotent cells. *Nature,* 522: 221-5.

13. Schmitt K, Heyne K, Roemer K, Meese E, et al., 2015. HERV-K(HML-2) rec and np9 transcripts not restricted to disease but present in many normal human tissues. *Mob DNA,* 6: 4.

14. Andersson AC, Venables PJ, Tonjes RR, Scherer J, et al., 2002. Developmental expression of HERV-R (ERV3) and HERV-K in human tissue. *Virology,* 297: 220-5.

15. Mayer J, Ehlhardt S, Seifert M, Sauter M, et al., 2004. Human endogenous retrovirus HERV-K(HML-2) proviruses with Rec protein coding capacity and transcriptional activity. *Virology,* 322: 190-8.

16. Izsvak Z, Wang J, Singh M, Mager DL, et al., 2016. Pluripotency and the endogenous retrovirus HERVH: Conflict or serendipity? *Bioessays,* 38: 109-17.

17. Kreimer U, Schulz WA, Koch A, Niegisch G, et al., 2013. HERV-K and LINE-1 DNA Methylation and Reexpression in Urothelial Carcinoma. *Front Oncol,* 3: 255.

18. Serafino A, Balestrieri E, Pierimarchi P, Matteucci C, et al., 2009. The activation of human endogenous retrovirus K (HERV-K) is implicated in melanoma cell malignant transformation. *Exp Cell Res,* 315: 849-62.

19. Balestrieri E, Argaw-Denboba A, Gambacurta A, Cipriani C, et al., 2018. Human Endogenous Retrovirus K in the Crosstalk Between Cancer Cells Microenvironment and Plasticity: A New Perspective for Combination Therapy. *Front Microbiol,* 9: 1448.

20. Reiche J, Pauli G, and Ellerbrok H, 2010. Differential expression of human endogenous retrovirus K transcripts in primary human melanocytes and melanoma cell lines after UV irradiation. *Melanoma Res,* 20: 435-40.

21. Schanab O, Humer J, Gleiss A, Mikula M, et al., 2011. Expression of human endogenous retrovirus K is stimulated by ultraviolet radiation in melanoma. *Pigment Cell Melanoma Res,* 24: 656-65.

22. Muster T, Waltenberger A, Grassauer A, Hirschl S, et al., 2003. An endogenous retrovirus derived from human melanoma cells. *Cancer Res,* 63: 8735-41.

23. Stengel S, Fiebig U, Kurth R, and Denner J, 2010. Regulation of human endogenous retrovirus-K expression in melanomas by CpG methylation. *Genes Chromosomes Cancer,* 49: 401-11.

24. Clausen J, 2003. Endogenous retroviruses and MS: using ERVs as disease markers. *Int MS J,* 10: 22-8.

25. Matteucci C, Balestrieri E, Argaw-Denboba A, and Sinibaldi-Vallebona P, 2018. Human endogenous
retroviruses role in cancer cell stemness. *Semin Cancer Biol*, 53: 17-30.
26. Hahn S, Ugurel S, Hanschmann KM, Strobel H, et al., 2008. Serological response to human endogenous
retrovirus K in melanoma patients correlates with survival probability. *AIDS Res Hum Retroviruses*, 24:
717-23.
27. Buscher K, Trefzer U, Hofmann M, Sterry W, et al., 2005. Expression of human endogenous retrovirus K in
melanomas and melanoma cell lines. *Cancer Res*, 65: 4172-80.
28. Schmitt K, Reichrath J, Roesch A, Meese E, et al., 2013. Transcriptional profiling of human endogenous
retrovirus group HERV-K(HML-2) loci in melanoma. *Genome Biol Evol*, 5: 307-28.
29. Contreras-Galindo R, Kaplan MH, Dube D, Gonzalez-Hernandez MJ, et al., 2015. Human Endogenous
Retrovirus Type K (HERV-K) Particles Package and Transmit HERV-K-Related Sequences. *J Virol*, 89:
7187-201.
30. Verfaillie A, Imrichova H, Atak ZK, Dewaele M, et al., 2015. Decoding the regulatory landscape of
melanoma reveals TEADS as regulators of the invasive cell state. *Nat Commun*, 6: 6683.
31. Bradish JR and Cheng L, 2014. Molecular pathology of malignant melanoma: changing the clinical practice
paradigm toward a personalized approach. *Hum Pathol*, 45: 1315-26.
32. Bollag G, Hirth P, Tsai J, Zhang J, et al., 2010. Clinical efficacy of a RAF inhibitor needs broad target
blockade in BRAF-mutant melanoma. *Nature*, 467: 596-9.
33. Li FZ, Dhillon AS, Anderson RL, McArthur G, et al., 2015. Phenotype switching in melanoma: implications
for progression and therapy. *Front Oncol*, 5: 31.
34. Caramel J, Papadogeorgakis E, Hill L, Browne GJ, et al., 2013. A switch in the expression of embryonic
EMT-inducers drives the development of malignant melanoma. *Cancer Cell*, 24: 466-80.
35. Babaian A and Mager DL, 2016. Endogenous retroviral promoter exaptation in human cancer. *Mob DNA*,
7: 24.
36. Tomlins SA, Laxman B, Dhanasekaran SM, Helgeson BE, et al., 2007. Distinct classes of chromosomal
rearrangements create oncogenic ETS gene fusions in prostate cancer. *Nature*, 448: 595-9.
37. Huang FW, Hodis E, Xu MJ, Kryukov GV, et al., 2013. Highly recurrent TERT promoter mutations in
human melanoma. *Science*, 339: 957-9.
38. Oricchio E, Sciamanna I, Beraldi R, Tolstonog GV, et al., 2007. Distinct roles for LINE-1 and HERV-K
retroelements in cell proliferation, differentiation and tumor progression. *Oncogene*, 26: 4226-33.
39. Lemaitre C, Tsang J, Bireau C, Heidmann T, et al., 2017. A human endogenous retrovirus-derived gene that
can contribute to oncogenesis by activating the ERK pathway and inducing migration and invasion. *PLoS
Pathog*, 13: e1006451.
40. Li M, Radvanyi L, Yin B, Li J, et al., 2017. Downregulation of Human Endogenous Retrovirus Type K
(HERV-K) Viral env RNA in Pancreatic Cancer Cells Decreases Cell Proliferation and Tumor Growth. *Clin
Cancer Res*, 23: 5892-5911.
41. Zhou F, Li M, Wei Y, Lin K, et al., 2016. Activation of HERV-K Env protein is essential for tumorigenesis
and metastasis of breast cancer cells. *Oncotarget*, 7: 84093-84117.
42. Ishida T, Obata Y, Ohara N, Matsushita H, et al., 2008. Identification of the HERV-K gag antigen in prostate
cancer by SEREX using autologous patient serum and its immunogenicity. *Cancer Immun*, 8: 15.
43. Riker AI, Enkemann SA, Fodstad O, Liu S, et al., 2008. The gene expression profiles of primary and
metastatic melanoma yields a transition point of tumor progression and metastasis. *BMC Med Genomics*,
1: 13.
44. Wallace TA, Downey RF, Seufert CJ, Schetter A, et al., 2014. Elevated HERV-K mRNA expression in PBMC
is associated with a prostate cancer diagnosis particularly in older men and smokers. *Carcinogenesis*, 35:
2074-83.
45. Dong J, Sedrak M, Rincon L, Sun H, et al., 2014. K-type human endogenous retroviral elements in human
melanoma. *Advances in Genomics and Genetics*: 153.
46. Chiappinelli KB, Strissel PL, Desrichard A, Li H, et al., 2015. Inhibiting DNA Methylation Causes an
Interferon Response in Cancer via dsRNA Including Endogenous Retroviruses. *Cell*, 162: 974-86.
47. Nelson PN, Lever AM, Smith S, Pitman R, et al., 1999. Molecular investigations implicate human
endogenous retroviruses as mediators of anti-retroviral antibodies in autoimmune rheumatic disease.
Immunol Invest, 28: 277-89.

Singh et al. (2020)

48.   Hirschl S, Schanab O, Seppele H, Waltenberger A, et al., 2007. Sequence variability of retroviral particles
derived from human melanoma cells melanoma-associated retrovirus. *Virus Res,* 123: 211-5.

49.   Ruprecht K, Mayer J, Sauter M, Roemer K, et al., 2008. Endogenous retroviruses and cancer. *Cell Mol Life*
Sci, 65: 3366-82.

50.   Hohn O, Hanke K, and Bannert N, 2013. HERV-K(HML-2), the Best Preserved Family of HERVs:
Endogenization, Expression, and Implications in Health and Disease. *Front Oncol,* 3: 246.

51.   Galli UM, Sauter M, Lecher B, Maurer S, et al., 2005. Human endogenous retrovirus rec interferes with
germ cell development in mice and may cause carcinoma in situ, the predecessor lesion of germ cell tumors.
Oncogene, 24: 3223-8.

52.   Singh S, Kaye S, Francis N, Peston D, et al., 2013. Human endogenous retrovirus K (HERV-K) rec mRNA
is expressed in primary melanoma but not in benign naevi or normal skin. *Pigment Cell Melanoma Res,* 26:
426-8.

53.   Ernst J, Kheradpour P, Mikkelsen TS, Shoresh N, et al., 2011. Mapping and analysis of chromatin state
dynamics in nine human cell types. *Nature,* 473: 43-9.

54.   Sun Q, Yang J, Xing G, Sun Q, et al., 2008. Expression of GSDML Associates with Tumor Progression in
Uterine Cervix Cancer. *Translational Oncology,* 1: 73-IN1.

55.   Katoh I, Mirova A, Kurata S, Murakami Y, et al., 2011. Activation of the long terminal repeat of human
endogenous retrovirus K by melanoma-specific transcription factor MITF-M. *Neoplasia,* 13: 1081-92.

56.   Levy C, Khaled M, and Fisher DE, 2006. MITF: master regulator of melanocyte development and melanoma
oncogene. *Trends Mol Med,* 12: 406-14.

57.   Steingrimsson E, Copeland NG, and Jenkins NA, 2004. Melanocytes and the microphthalmia transcription
factor network. *Annu Rev Genet,* 38: 365-411.

58.   Blackwood EM and Eisenman RN, 1991. Max: a helix-loop-helix zipper protein that forms a sequence-
specific DNA-binding complex with Myc. *Science,* 251: 1211-7.

59.   Webster DE, Barajas B, Bussat RT, Yan KJ, et al., 2014. Enhancer-targeted genome editing selectively blocks
innate resistance to oncokinase inhibition. *Genome Res,* 24: 751-60.

60.   Muller J, Krijgsman O, Tsoi J, Robert L, et al., 2014. Low MITF/AXL ratio predicts early resistance to
multiple targeted drugs in melanoma. *Nat Commun,* 5: 5712.

61.   Hoek KS, Eichhoff OM, Schlegel NC, Dobbeling U, et al., 2008. In vivo switching of human melanoma cells
between proliferative and invasive states. *Cancer Res,* 68: 650-6.

62.   Kim HY, Lee H, Kim SH, Jin H, et al., 2017. Discovery of potential biomarkers in human melanoma cells
with different metastatic potential by metabolic and lipidomic profiling. *Sci Rep,* 7: 8864.

63.   Clauss J, Obenaus M, Miskey C, Ivics Z, et al., 2018. Efficient Non-Viral T-Cell Engineering by Sleeping
Beauty Minicircles Diminishing DNA Toxicity and miRNAs Silencing the Endogenous T-Cell Receptors.
Hum Gene Ther, 29: 569-584.

64.   Mates L, Chuah MK, Belay E, Jerchow B, et al., 2009. Molecular evolution of a novel hyperactive Sleeping
Beauty transposase enables robust stable gene transfer in vertebrates. *Nat Genet,* 41: 753-61.

65.   Chae YK, Chang S, Ko T, Anker J, et al., 2018. Epithelial-mesenchymal transition (EMT) signature is
inversely associated with T-cell infiltration in non-small cell lung cancer (NSCLC). *Sci Rep,* 8: 2918.

66.   Alonso SR, Tracey L, Ortiz P, Perez-Gomez B, et al., 2007. A high-throughput study in melanoma identifies
epithelial-mesenchymal transition as a major determinant of metastasis. *Cancer Res,* 67: 3450-60.

67.   Nieto MA, Huang RY, Jackson RA, and Thiery JP, 2016. Emt: 2016. *Cell,* 166: 21-45.

68.   Lin K, Baritaki S, Militello L, Malaponte G, et al., 2010. The Role of B-RAF Mutations in Melanoma and the
Induction of EMT via Dysregulation of the NF-kappaB/Snail/RKIP/PTEN Circuit. *Genes Cancer,* 1: 409-
420.

69.   Weiss MB, Abel EV, Mayberry MM, Basile KJ, et al., 2012. TWIST1 is an ERK1/2 effector that promotes
invasion and regulates MMP-1 expression in human melanoma cells. *Cancer Res,* 72: 6382-92.

70.   Dissanayake SK, Wade M, Johnson CE, O'Connell MP, et al., 2007. The Wnt5A/protein kinase C pathway
mediates motility in melanoma cells via the inhibition of metastasis suppressors and initiation of an
epithelial to mesenchymal transition. *J Biol Chem,* 282: 17259-71.

71.   O'Connell MP, Marchbank K, Webster MR, Valiga AA, et al., 2013. Hypoxia induces phenotypic plasticity
and therapy resistance in melanoma via the tyrosine kinase receptors ROR1 and ROR2. *Cancer Discov,* 3:
1378-93.

72.   Derynck R and Zhang YE, 2003. Smad-dependent and Smad-independent pathways in TGF-beta family
signalling. *Nature*, 425: 577-84.

73.   Zavadil J and Bottinger EP, 2005. TGF-beta and epithelial-to-mesenchymal transitions. *Oncogene*, 24: 5764-
74.

74.   Morita T, Mayanagi T, and Sobue K, 2007. Dual roles of myocardin-related transcription factors in epithelial
mesenchymal transition via slug induction and actin remodeling. *J Cell Biol*, 179: 1027-42.

75.   Almanzar G, Olkhanud PB, Bodogai M, Dell' agnola C, et al., 2009. Sperm-derived SPANX-B is a clinically
relevant tumor antigen that is expressed in human tumors and readily recognized by human CD4+ and
CD8+ T cells. *Clin Cancer Res*, 15: 1954-63.

76.   Yilmaz-Ozcan S, Sade A, Kucukkaraduman B, Kaygusuz Y, et al., 2014. Epigenetic mechanisms underlying
the dynamic expression of cancer-testis genes, PAGE2, -2B and SPANX-B, during mesenchymal-to-
epithelial transition. *PLoS One*, 9: e107905.

77.   Woods K, Pasam A, Jayachandran A, Andrews MC, et al., 2014. Effects of epithelial to mesenchymal
transition on T cell targeting of melanoma cells. *Front Oncol*, 4: 367.

78.   Tirosh I, Izar B, Prakadan SM, Wadsworth MH, 2nd, et al., 2016. Dissecting the multicellular ecosystem of
metastatic melanoma by single-cell RNA-seq. *Science*, 352: 189-96.

79.   Pencheva N, Tran H, Buss C, Huh D, et al., 2012. Convergent multi-miRNA targeting of ApoE drives
LRP1/LRP8-dependent melanoma metastasis and angiogenesis. *Cell*, 151: 1068-82.

80.   Castello LM, Raineri D, Salmi L, Clemente N, et al., 2017. Osteopontin at the Crossroads of Inflammation
and Tumor Progression. *Mediators Inflamm*, 2017: 4049098.

81.   Cheng L, Lopez-Beltran A, Massari F, MacLennan GT, et al., 2018. Molecular testing for BRAF mutations
to inform melanoma treatment decisions: a move toward precision medicine. *Mod Pathol*, 31: 24-38.

82.   Goding CR, 2011. Commentary. A picture of Mitf in melanoma immortality. *Oncogene*, 30: 2304-6.

83.   Hoek KS and Goding CR, 2010. Cancer stem cells versus phenotype-switching in melanoma. *Pigment Cell*
Melanoma Res, 23: 746-59.

84.   Vachtenheim J and Ondrusova L, 2015. Microphthalmia-associated transcription factor expression levels
in melanoma cells contribute to cell invasion and proliferation. *Exp Dermatol*, 24: 481-4.

85.   Carreira S, Goodall J, Denat L, Rodriguez M, et al., 2006. Mitf regulation of Dia1 controls melanoma
proliferation and invasiveness. *Genes Dev*, 20: 3426-39.

86.   Litvin O, Schwartz S, Wan Z, Schild T, et al., 2015. Interferon alpha/beta Enhances the Cytotoxic Response
of MEK Inhibition in Melanoma. *Mol Cell*, 57: 784-96.

87.   Shirley SH, Greene VR, Duncan LM, Torres Cabala CA, et al., 2012. Slug expression during melanoma
progression. *Am J Pathol*, 180: 2479-89.

88.   Tuncer E, Calcada RR, Zingg D, Varum S, et al., 2019. SMAD signaling promotes melanoma metastasis
independently of phenotype switching. *J Clin Invest*, 129: 2702-2716.