Expansion of SARS-CoV-2-specific Antibody-secreting Cells and Generation of

Neutralizing Antibodies in Hospitalized COVID-19 Patients

Running title: B cell and antibody responses in COVID-19 patients.

Authors

Renata Varnaitė[*], Marina García[*], Hedvig Glans[†,‡,1], Kimia T. Maleki[*,1], John Tyler

Sandberg[*,1], Janne Tynell[*], Wanda Christ[*], Nina Lagerqvist[§], Hilmir Asgeirsson[†,¶], Hans-

Gustaf Ljunggren[*,‡], Gustaf Ahlén[#], Lars Frelin[#], Matti Sällberg[#], Kim Blom[*], Jonas

Klingström[*,§], Sara Gredmark-Russ[*,†]

[1] H.G., K.T.M. and J.T.S. contributed equally to this work.

Affiliations

[*] Center for Infectious Medicine, ANA Futura, Department of Medicine Huddinge,

Karolinska Institutet, Stockholm, Sweden.

[†] Department of Infectious Diseases, Karolinska University Hospital, Stockholm, Sweden.

[‡] Department of Medicine Solna, Karolinska Institutet, Stockholm, Sweden.

[§] Public Health Agency of Sweden, Solna, Sweden.

[¶] Division of Infectious Diseases, Department of Medicine Huddinge, Karolinska Institutet,

Stockholm, Sweden.

[#] Department of Laboratory Medicine, Division of Clinical Microbiology, ANA Futura,

Karolinska Institutet, Stockholm, Sweden.

Abstract

Coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome

coronavirus 2 (SARS-CoV-2), emerged in late 2019 and has since become a global

pandemic. Pathogen-specific antibodies are typically a major predictor of protective

immunity, yet B cell and antibody responses during COVID-19 are not fully understood.

Here, we analyzed antibody-secreting cell (ASC) and antibody responses in twenty

hospitalized COVID-19 patients. The patients exhibited typical symptoms of COVID-19, and

presented with reduced lymphocyte numbers and increased T cell and B cell activation.

Importantly, we detected an expansion of SARS-CoV-2 nucleocapsid protein-specific ASCs

in all twenty COVID-19 patients using a multicolor FluoroSpot assay. Out of the 20 patients,

16 had developed SARS-CoV-2-neutralizing antibodies by the time of inclusion in the study.

SARS-CoV-2-specific IgA, IgG and IgM antibody levels positively correlated with SARS-

CoV-2-neutralizing antibody titers, suggesting that SARS-CoV-2-specific antibody levels

may reflect the titers of neutralizing antibodies in COVID-19 patients during the acute phase

of infection. Lastly, we showed that interleukin 6 (IL-6) and C-reactive protein (CRP)

concentrations were higher in serum of patients who were hospitalized for longer, supporting

the recent observations that IL-6 and CRP could be used to predict COVID-19 severity.

Altogether, this study constitutes a detailed description of clinical and immunological

parameters in twenty COVID-19 patients, with a focus on B cell and antibody responses, and

provides tools to study immune responses to SARS-CoV-2 infection and vaccination.

Keywords: SARS-CoV-2, COVID-19, antibody-secreting cells, B cells, antibodies, T cells,

interleukin 6

Introduction

Coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), emerged in late 2019 and has since become a global pandemic (1). SARS-CoV-2 primarily affects the respiratory tract and can cause severe respiratory illness, which sometimes requires mechanical ventilation (2). The world is racing to understand the immune responses to SARS-CoV-2 in order to identify correlates of protection, and to aid treatment and vaccine development. In the majority of infections, pathogen-specific antibodies are one of the main contributors to protective immunity, yet B cell and antibody responses during COVID-19 are currently not fully understood.

B cells are one of the key players of the adaptive immune system and can provide sterilizing immunity against pathogens through the secretion of pathogen-specific antibodies. B cells carry immunoglobulin (Ig) surface receptors with a high range of specificity that can directly recognize different antigens. Early during acute infections upon the encounter with the antigen, B cells differentiate into antibody-secreting cells (ASCs), including plasmablasts and plasma cells, which produce large quantities of pathogen-specific antibodies (3).

ASCs can expand to very high levels during some acute human infections (4-7). For example, during human dengue virus infection, ASCs expand to constitute an average of 47% of all circulating B cells (5). Meanwhile, at steady state conditions in healthy individuals, ASCs comprise only up to around 1% of all circulating B cells (5). The ASC response is typically transient, peaking at around day 7 of the febrile stage of infection (8). However, during acute human respiratory syncytial virus (RSV) infection, ASC response could be detected for as long as 22-45 days after the onset of symptoms, which was found to be associated with

prolonged shedding of RSV in the airways (9). Thus, the kinetics of ASC response may be

dependent on pathogen persistence.

In humans, ASCs can be identified using flow cytometry as $CD19^+CD20^{low/-}$ isotype-

switched B cells with high cell surface expression of CD38 and CD27 molecules, as well as

intracellular expression of proliferation marker Ki-67 (10, 11). Although flow cytometry is a

valuable tool to measure the frequencies of total ASCs in peripheral blood, it only allows

phenotypic characterization. Functional antibody secretion capacity and antigen-specificity of

ASCs are typically described using the gold standard enzyme-linked immunospot (ELISpot)

assay (12). ELISpot is a highly sensitive method allowing for detection of ASCs producing

different antibody isotypes at single-cell level, including IgA, IgG and IgM, against different

antigens. Flow cytometry and ELISpot can therefore be used in combination to

comprehensively describe the magnitude and specificity of B cell responses to infectious

agents.

Antigen-specific antibodies can contribute to pathogen control either by (i) directly

interfering with the entry of a pathogen into the target cells (referred to as neutralization); or

by (ii) assisting the effector cells in recognizing and eliminating infected target cells (known

as antibody-dependent cellular cytotoxicity) (13). Neutralizing antibodies are typically

considered to be the strongest predictor of protective immunity following natural infection or

vaccination. Therefore, it is of great importance to evaluate the neutralizing capacity and

persistence of specific antibodies in acute viral infections such as SARS-CoV-2.

Here, we provide a detailed description of early B cell and antibody responses to SARS-CoV-

2 infection in a cohort of twenty hospitalized patients with severe COVID-19. We found a

substantial level of ASCs expansion in all patients, and that not all patients developed

detectable levels of SARS-CoV-2-specific and neutralizing antibodies at the time of

sampling. We also show increased levels of adaptive immune system activation and increased

serum concentrations of inflammatory markers interleukin 6 (IL-6) and C-reactive protein

(CRP), the latter two correlating with the duration of hospitalization in the present COVID-

19 patient cohort. This study constitutes a detailed description of clinical and immunological

B cell-related parameters in COVID-19 patients, and describes tools to study immune

responses to SARS-CoV-2 infection and vaccination.

Materials and Methods

Ethics statement

The study was approved by the Regional Ethical Review Board in Stockholm, Sweden and by the Swedish Ethical Review Authority. All COVID-19 patients and healthy controls included in this study provided a written informed consent for participation.

Study subjects and sampling of peripheral blood

Peripheral blood samples were collected from 20 adult COVID-19 patients hospitalized in April 2020 at the Karolinska University Hospital in Stockholm, Sweden (5 females and 15 males; age range between 34 and 67 years; median age 53 years) (Table I). Patients were diagnosed with COVID-19 by RT-PCR for SARS-CoV-2 in either nasopharyngeal swabs (18/20 patients) or sputum (2/20 patients). Diagnostics were performed at the diagnostic laboratory at the Karolinska University Hospital, Stockholm, Sweden. Peripheral blood samples from patients were taken at a median of 15 days after self-reported onset of symptoms (range 7-19 days). Peripheral blood samples of 7 healthy controls were collected in parallel (2 females and 5 males; age range between 26 and 53 years; median age 31 years).

Peripheral blood mononuclear cells (PBMCs) were isolated from heparinized anti-coagulated blood using density gradient Lymphoprep medium (Stemcell Technologies) following the manufacturer's instructions, and immediately used for flow cytometry and FluoroSpot assays. Serum was collected from COVID-19 patients and healthy controls in BD Vacutainer serum tubes with spray-coated silica (BD Biosciences). After coagulation for up to 2 hours at RT, serum was isolated by centrifugation at 2000 g for 10 min and immediately stored at -80 ºC for later analysis.

Absolute counts of lymphocytes in peripheral blood

Absolute numbers of $CD45^+$, $CD3^+$, $CD4^+$, $CD8^+$, and $CD19^+$ cells in peripheral blood were

measured using BD Trucount Tubes (BD Biosciences). 50 µL of anti-coagulated whole blood

were added into Trucount Tubes within 3 hours after blood extraction and stained with either

anti-CD45-PerCP (2D1), anti-CD3-FITC (SK7), anti-CD4-APC (SK3) and anti-CD8-PE

(SK1), or anti-CD45-PerCP (2D1) and anti-CD19-AF488 (HIB19) (all from BioLegend).

After 15 minutes of incubation at RT, stained whole blood was fixed and red blood cells

lysed with 2X BD FACS Lysing Solution (BD Biosciences). Samples were acquired on a BD

Accuri C6 Plus flow cytometer. Bead number recorded was used to quantify absolute $CD45^+$,

$CD3^+$, $CD4^+$, $CD8^+$, and $CD19^+$ cell counts per microliter of blood. ASC numbers per

microliter of blood were calculated based on $CD19^+$ B cell numbers measured by absolute

cell counting and on frequencies of ASCs within $CD19^+$ cells measured by flow cytometry.

Flow cytometry

Staining with fluorescently-labelled antibodies was performed on freshly isolated PBMCs.

Briefly, cells were incubated with surface staining antibodies diluted in PBS for 30 min at

4°C in the dark, followed by 3 washes with flow cytometry buffer (2% FCS and 2 mM

EDTA in PBS). Cells were then fixed and permeabilized using eBioscience

Foxp3/Transcription Factor Staining Buffer Set (Thermo Fisher Scientific) and later

incubated with antibodies diluted in PBS for intracellular staining for 30 min at 4°C in the

dark. Finally, samples were incubated in a 2% formaldehyde solution (Polysciences) for 2 h,

washed and resuspended in flow cytometry buffer, and data subsequently acquired on a BD

LSRFortessa flow cytometer equipped with 355, 405, 488, 561, and 639 nm lasers and BD

FACSDiva Software (BD Biosciences). For a detailed gating strategy see Supplementary Fig.

1.

The following monoclonal antibody conjugates were used for cell surface staining: anti-CD8-

Qdot605 (3B5) (Thermo Fisher Scientific), anti-CD19-BUV395 (SJ25C1), anti-CD14-V500

(MφP9), anti-CD4 -BUV737 (RPA-T4) (all from BD Biosciences), anti-CD123-BV510

(6H6), anti-CD27-BV650 (O323), anti-CD20-FITC (2H7), anti-CD38-BV421 (HB-7), anti-

IgD-PE-Cy7 (IA6-2), anti-IgM-BV785 (MHM-88) (all from BioLegend), anti-CD3-PE-Cy5

(UCHT1), anti-CD56-ECD (N901) (all from Beckman Coulter), and anti-IgA-APC

(REA1014) (Miltenyi). LIVE/DEAD Fixable Near-IR Dead Cell Stain Kit (Thermo Fisher

Scientific) was used as a viability marker. The following monoclonal antibody conjugates

were used for intracellular staining: anti-IgG-PE (HP6017) (BioLegend) and anti-Ki-67-

AF700 (B56) (BD Biosciences).

FluoroSpot assay for antibody-secreting cells

The number of SARS-CoV-2 nucleocapsid (N) protein-specific IgA, IgG and IgM antibody-

secreting cells (ASCs), as well as the total number of IgA-, IgG- and IgM-ASCs in freshly

isolated PBMCs were measured using a multicolor B cell FluoroSpot kit with modifications

(Mabtech). Briefly, ethanol-activated IPFL membrane plates were coated overnight with

either: (i) anti-IgG, anti-IgA, and anti-IgM capture antibodies (15μg/mL of each) for the

detection of all ASCs, or (ii) SARS-CoV-2 N protein (10 μg/mL) for the detection of SARS-

CoV-2-specific ASCs. The plates were washed with PBS and blocked with R10 media

(RPMI-1640 with 10% FCS, 1% Pen/Strep, 2mM L-Glutamine (all from Thermo Fisher

Scientific)) for 30 minutes at RT before the addition of freshly isolated PBMCs. Plates were

then incubated at 37°C in 5% CO_2 for 20 hours and then developed with anti-human IgG-550

(yellow fluorescence), anti-human IgA-490 (green fluorescence) and anti-human IgM-640

(red fluorescence) secondary detection antibodies (diluted 1:500 each) (all antibodies from

Mabtech). Fluorescent spots indicating a single ASC were detected with an IRIS FluoroSpot

reader and counted with Apex software (Mabtech).

Recombinant SARS-CoV-2 nucleocapsid protein

A full-length nucleocapsid (N) phosphoprotein nucleotide sequence (1293 base-pairs) of the

SARS-CoV-2 virus was optimized and synthesized (Genscript). The synthesized sequence

was cloned into a PET-30a(+) vector with a carboxyterminal His tag for detection of protein

expression in *E.coli*. The *E. coli* strain BL21 Star (DE3) was transformed with the

recombinant plasmid and a single colony was inoculated into TB medium containing

antibiotic and cultured at 37°C at 200 rpm and then induced with IPTG. Protein purity and

molecular weight were determined by SDS-PAGE and Western blot according to standard

procedures (Genscript).

SARS-CoV-2 isolation from serum

50 µL of serum were mixed with 150 µL of EMEM (Gibco) and added to confluent Vero E6

cells seeded in 24-well plates. Cells were incubated with diluted serum for 1 hour, and 1 mL

of Vero E6 medium was added after the incubation. Cells were subsequently incubated

further for 10 days at 37 °C and 5% CO_2 and monitored for cytopathic effect (CPE) by

optical microscopy.

Immunofluorescence assay (IFA) for IgG against SARS-CoV-2

Vero E6 cells were infected with SARS-CoV-2 (isolate SARS-CoV-2/human/SWE/01/2020,

accession number MT093571) for 24 hours, trypsinized and mixed with uninfected Vero E6

cells, and then seeded on microscope slides. Twelve hours later, slides were fixed in acetone

and stored at -80˚C until further use. Serum samples were heat-inactivated at 56˚C for 30

minutes prior to analysis. For analysis of total SARS-CoV-2 IgG antibody titers, serum

samples were serially diluted from 1:20 to 1:5120. 25 μL of diluted serum was then added to

fixed cells and incubated at 37˚C for 30 min, after which the slides were washed in NaCl for

30 min. Bound SARS-CoV-2 IgG antibodies were then detected by incubating for 30 min at

37˚C with a secondary AF488-conjugated AffiniPure goat anti-human IgG antibody (Jackson

Immunoresearch), diluted 1:200 in 0.1% Evan's Blue. SARS-CoV-2 IgG positive cells were

visualized using a Nikon Eclipse Ni fluorescence microscope (x40 magnification). The titer

of IgG in each serum sample was determined by the inverted dilution factor value for the

highest dilution with positive staining.

ELISAs

SARS-CoV-2 specific IgG and IgA antibodies in serum were detected using anti-SARS-

CoV-2 ELISA kits (both from Euroimmun), according to the manufacturer's instructions.

SARS-CoV-2 specific IgM antibodies were detected using EDI Novel Coronavirus COVID-

19 IgM ELISA kit (Epitope Diagnostics), according to the manufacturer's instructions. Serum

samples were heat-inactivated at 56˚C for 30 minutes prior to analysis.

IL-6 levels in serum from patients and healthy controls were measured in freshly thawed

serum using human IL-6 ELISA development kit (Mabtech), according to the manufacturer's

instructions. Serum samples were diluted 1:2 in ready-to-use ELISA diluent (Mabtech) prior

to performing the IL-6 ELISA assay.

Micro-neutralization assay

Two-fold dilution series from 1:10 to 1:10240 in EMEM (Gibco) + 5% FCS (Thermo Fisher

Scientific) were performed on the serum samples, which were previously heat inactivated at

56˚C for 30 minutes. Each dilution was subsequently mixed with equal volume of 4000

$TCID_{50}$/ml SARS-CoV-2 (50 μl serum plus 50 μl virus) and incubated for 1 hour at 37 °C

and 5% CO_2. Each sample was prepared in duplicates. After incubation, the mixtures were

added on confluent Vero E6 cells seeded on 96-well plates and incubated at 37 °C 5% CO_2.

Four days later the cells were inspected for signs of cytopathic effect (CPE) by optical

microscopy. Each well was scored as either 'neutralizing' if less than 50% of the cell layer

showed signs of CPE, or 'non-neutralizing' if ≥50% CPE was observed. Results are shown as

the arithmetic mean of the reciprocals of the highest neutralizing dilutions from the two

duplicates for each sample.

Real-time RT-PCR

RNA was extracted from freshly thawed serum samples using the MagDEA Dx SV reagent

kit and the magLEAD instrument (Precision System Science). The assay used to detect

SARS-CoV-2 RNA was modified from (14): forward primer 5'-

CATGTGTGGCGGTTCACTATATGT-3', reverse primer 5'-

TGTTAAARACACTATTAGCATAWGCAGT-3', and RdRp_SARSr-P2 probe. The assay

was carried out in 25 μL reaction mixtures containing 5μL RNA template, TaqMan Fast

Virus 1-Step Master Mix, 0.6 μM forward primer, 0.8 μM reverse primer, and 0.2 μM probe

(Applied Biosystems). Thermal cycling was performed at 50°C for 5 min, 95°C for 20 sec,

followed by 40 cycles of 95°C for 3 sec, and 60°C for 30 sec in a StepOne Plus real-time

PCR (Applied Biosystems).

Statistics and data analysis

Statistical analyses were performed using GraphPad Prism software 7.0 for MacOSX

(GraphPad Software). Correlation analyses were performed using Spearman's correlation

test. Statistical significance for differences between COVID-19 patients and healthy controls

was determined by two-sided Mann-Whitney *U* test. P values of < 0.05 were considered

statistically significant. FlowJo software version 10.5.3 (Tree Star) was used to analyze all

flow cytometry data. FluoroSpot data was analyzed with Apex software (Mabtech).

Results

Twenty COVID-19 patients were enrolled in this study during hospitalization at Karolinska

University Hospital, Sweden. All twenty patients exhibited typical COVID-19 symptoms of

fever, cough and breathing difficulties, and a few also experienced chest pain, myalgia and

diarrhea (Table I). Some of the patients presented with common co-morbidities such as

overweightness, hypertension, asthma, cardiovascular disease and diabetes mellitus type II.

According to the guidelines for diagnosis and treatment of COVID-19 (Version 7) released

by the National Health Commission and State Administration of Traditional Chinese

Medicine, 19 out of the 20 patients were classified as severe, and 1 out of the 20 as critically

ill (15). While all patients enrolled in this study were hospitalized at the infectious diseases

wards at the time of sampling, two patients were treated at the intensive care unit (ICU)

before inclusion in this study, while one patient required ICU treatment after the inclusion.

Peripheral blood samples from the twenty COVID-19 patients were collected at a median of

15 days after the self-reported onset of symptoms, and at a median of 3 days after

hospitalization (Table I, Fig. 1A and B). In parallel, seven donors without an ongoing

respiratory disease or signs of inflammation were included in the study as healthy controls.

Clinical chemistry parameters in COVID-19 patients

Blood cell counts and clinical chemistry parameters in peripheral blood of COVID-19

patients were measured at the Diagnostics Laboratory of the Karolinska University Hospital

on the same day as the inclusion in this study (+/− 24 hours). Clinical chemistry tests showed

increased levels of inflammatory markers, including C-reactive protein (CRP), lactate

dehydrogenase (LDH) and ferritin (Fig. 1C). Some patients also presented with increased

levels of aspartate transaminase (AST), alanine transaminase (ALT) and D-dimer, while

hemoglobin, procalcitonin (PCT) and troponin T levels fell within the range of reference

values in the majority of patients (Fig. 1C). Numbers of total leukocytes, neutrophils,

monocytes and platelets in peripheral blood were within the normal range in the majority of

COVID-19 patients at the time of sampling (Fig. 1C). In contrast, we observed reduced

lymphocyte numbers, with the median lymphocyte numbers falling below or within the lower

range of reference values (Fig. 1C). Reduced lymphocyte numbers in peripheral blood of

COVID-19 patients were confirmed by flow cytometry analysis of absolute cell counts. We

found significantly lower $CD45^+$ lymphocyte numbers in the patients compared to healthy

controls (Fig. 1D). $CD3^+$ T cell and $CD3^+CD8^+$ T cell numbers were also significantly lower

in patients compared to the controls, while no significant difference in the numbers of

$CD3^+CD4^+$ T cells and $CD19^+$ B cells was observed between COVID-19 patients and healthy

controls (Fig. 1D).

T cells and B cells are activated in COVID-19 patients

We next assessed the overall activation level of the adaptive immune system in COVID-19

patients. T cell and B cell activation levels were measured based on the expression of the

activation marker CD38 and the proliferation marker Ki-67 using multicolor flow cytometry

on freshly isolated peripheral blood mononuclear cells (PBMCs) (Fig. 2A). We observed

significantly higher frequencies of activated $CD4^+$ T cells, $CD8^+$ T cells and $CD19^+$ B cells in

COVID-19 patients compared to healthy controls (Fig. 2B). Additionally, there was a strong

positive correlation between the frequencies of activated $CD8^+$ T cells and $CD4^+$ T cells in

COVID-19 patients ($r_s = 0.699$, $P < 0.001$) (Supplementary Fig. 2A).

Significant expansion of antibody-secreting cells in COVID-19 patients

Flow cytometric analyses revealed increased frequencies of activated $CD38^+Ki-67^+$ B cells in

the COVID-19 patients (Fig. 2A and B). Therefore, we further analyzed if antibody-secreting

cell (ASC) frequencies were also increased in peripheral blood. ASCs were identified as isotype-switched B cells expressing high levels of surface CD38 and CD27 ($CD19^+CD20^{low/-}$ $IgD^-CD38^{high}CD27^{high}$) (Fig. 3A). As expected, low frequencies and numbers of ASCs were detected in peripheral blood of healthy controls (Fig. 3B and C). However, significantly higher ASC frequencies and numbers were observed in COVID-19 patients compared to healthy controls, constituting up to 31% of all circulating B cells (Fig. 3B and C).

Next, we assessed the expression of immunoglobulins (Ig) by the ASCs in COVID-19 patients and healthy controls (Fig. 3A). IgA has previously been shown to be expressed by 80% of all ASCs at steady state (16). In agreement with this, the largest proportion within the ASC compartment in healthy controls was IgA^+ ASCs, whereas the majority of ASCs in COVID-19 patients expressed IgG (Fig. 3D). IgM^+ ASCs were present at comparable frequencies in both COVID-19 patients and healthy controls (Fig. 3D). Noteworthy, a substantial ASC expansion was detected as early as 7 days, and as late as 19 days after the onset of symptoms (Supplementary Fig. 2B).

Detection of SARS-CoV-2-specific antibody-secreting cells in COVID-19 patients

The expansion of ASCs is usually a result of an ongoing infection and is characterized by high specificity towards the infectious agent (17). To investigate if the ASCs identified by flow cytometry were SARS-CoV-2-specific, we developed a FluoroSpot assay - a fluorescence-based variant of the enzyme-linked immunospot (ELISpot) assay, which allows for simultaneous detection of total and antigen-specific IgA-, IgG- and IgM-ASCs. To detect SARS-CoV-2-specific IgA-, IgG- and IgM-ASCs, we generated recombinant SARS-CoV-2 nucleocapsid protein (N protein).

The FluoroSpot assay confirmed the ASC expansion detected by flow cytometry, as the total

numbers of IgA-, IgG-, and IgM-ASCs in COVID-19 patients were significantly higher than

in healthy controls (Fig. 3E and G). Moreover, the total ASC frequencies and numbers

detected by flow cytometry positively correlated with total numbers of ASCs measured by

FluoroSpot (r_s = 0.537, P = 0.01; and r_s = 0.636, P = 0.003, respectively). Consistent with

flow cytometry data, IgG-ASCs constituted the predominant subset of all ASCs in COVID-

19 patients, followed by IgA-ASCs (Fig. 3D and G).

Importantly, we detected SARS-CoV-2 N protein-specific ASCs in all twenty COVID-19

patients, but not in controls, suggesting an active SARS-CoV-2-specific B cell response in

acute COVID-19 (Fig. 3F and H). We detected IgA-, IgG- and IgM-ASCs that are specific

for SARS-CoV-2 N-protein, yet at highly variable numbers among the patients. Overall, N-

protein-specific ASCs constituted a small fraction of total ASCs [median (IQR) = 3.47%

(2.5-4.9)]. Nonetheless, total ASC frequencies determined by flow cytometry positively

correlated with N protein-specific ASC numbers (r_s = 0.574, P = 0.008), suggesting that the

ASC expansion detected by flow cytometry may reflect the magnitude of SARS-CoV-2 N

protein-specific ASC response.

Generation of SARS-CoV-2-specific and neutralizing antibodies in COVID-19 patients

The expansion of SARS-CoV-2-specific ASCs in all of the patients in our cohort suggested

that the patients had developed SARS-CoV-2-specific antibodies in response to the infection.

To investigate this in detail, we next analyzed SARS-CoV-2-specific antibody levels.

First, we measured SARS-CoV-2 spike S1-specific IgA and IgG, as well as N-protein-

specific IgM antibody levels by ELISAs. We found detectable SARS-CoV-2-specific IgA

(15/20 patients), IgG (15/20 patients) and IgM (16/20 patients) antibody levels in most of the

COVID-19 patients (Fig. 4A and B). Next, we determined total anti-SARS-CoV-2 IgG

antibody levels measured towards SARS-CoV-2-infected cells using an immunofluorescence

assay (IFA) (Fig. 4C). We found that 16 out of the 20 patients were positive in this assay,

with titers ranging from 40 to 5120 (Fig. 4C). As expected, none of the healthy controls were

positive in any of the antibody assays (Fig. 4A). Higher IgG levels were detected in patients

who were sampled late compared to early after the onset of symptoms, as total and spike S1-

specific SARS-CoV-2 IgG antibody levels positively correlated with the number of days

since symptom onset ($r_s = 0.577$, P = 0.01, and $r_s = 0.603$, P = 0.005, respectively)

(Supplementary Fig. 2C and D).

Next, using a micro-neutralization assay we assessed if the COVID-19 patients had generated

SARS-CoV-2-neutralizing antibodies (18). This assay measures the ability of COVID-19

patient serum to reduce the cytopathic effect (CPE) caused by SARS-CoV-2 infection of

susceptible cells *in vitro*. Neutralizing antibodies were detected in most of the patients

(16/20), but at heterogeneous titers ranging from 10 to 1920 (Fig. 4D). Three of the four

patients with undetectable levels of SARS-CoV-2-neutralizing antibodies were also below

the level of detection for SARS-CoV-2-specific antibodies in IFA and all ELISAs (Fig. 4A).

Interestingly, we observed a strong positive correlation between total SARS-CoV-2-specific

IgG antibody levels and neutralizing antibody titers in COVID-19 patients (Fig. 4E). Both

S1-specific IgA and IgG, as well as N-specific IgM levels also correlated with SARS-CoV-2-

neutralizing antibody titers suggesting that SARS-CoV-2-specific antibody titers may reflect

the levels of neutralizing antibodies during the acute phase of COVID-19 (Fig. 4E).

Next, we assessed if the frequencies of activated T cells and B cells correlate with SARS-

CoV-2-neutralizing antibody responses. We found that $CD8^+$ T cell activation level, but not

$CD4^+$ T cell or $CD19^+$ B cell activation level, positively correlated with neutralizing antibody

titers ($r_s = 0.544$, $P = 0.01$; $r_s = 0.271$, $P = 0.25$, and $r_s = -0.136$, $P = 0.57$, respectively)

(Supplementary Fig. 2E - G).

Detection of SARS-CoV-2 RNA in serum of COVID-19 patients

An important role of neutralizing antibodies is to limit the spread of the virus. We screened

all serum samples by real time RT-PCR for SARS-CoV-2 RNA (14). Three COVID-19

patients were positive for SARS-CoV-2 RNA in serum (Fig. 4A). Notably, 2 out of the 3

SARS-CoV-2 RT-PCR-positive patients lacked detectable levels of neutralizing antibodies,

which might allow for a more efficient spread of SARS-CoV-2 in those patients (Fig. 4A).

However, our attempts to isolate live SARS-CoV-2 from patient serum on Vero E6 cells were

unsuccessful (data not shown), suggesting absence or low levels of live SARS-CoV-2 in

serum of COVID-19 patients.

Inflammatory makers, neutralizing antibodies and disease severity in COVID-19

The pro-inflammatory cytokine IL-6 and CRP serum levels have been shown to correlate

with disease severity in COVID-19 patients (19, 20). We measured IL-6 serum concentration

in all COVID-19 patients at the time of inclusion in this study using ELISA. We found

heterogenous, yet significantly higher median concentration of IL-6 in COVID-19 patients

compared to the controls (Fig. 5A).

There was a strong positive correlation between the IL-6 and CRP levels at the time of

inclusion in this study (Fig. 5B). Higher serum concentrations of both IL-6 and CRP were

higher in patients with longer duration of hospitalization (Fig. 5B). Notably, IL-6 and CRP

serum concentrations negatively correlated with neutralizing antibody titers in COVID-19

patients, indicating a possible link between inflammation and humoral responses in COVID-

19 (Fig. 5B).

Discussion

Here, we demonstrated that COVID-19 patients elicit a SARS-CoV-2-specific B cell response, indicated by the expansion of SARS-CoV-2-specific ASCs. Although not all patients in the present cohort had detectable levels of SARS-CoV-2-specific antibodies at the time of sampling, SARS-CoV-2 nucleocapsid protein-specific ASCs could be detected in all patients using the FluoroSpot assay. In addition, we showed a clear relationship between the levels of SARS-CoV-2-specific antibodies and total SARS-CoV-2-neutralizing antibodies. This suggests that standard serological assays may inform about the levels of neutralizing SARS-CoV-2-specific antibodies, which may offer protection from a re-infection. Additionally, tools employed in this study may be of relevance in the assessment of long-lasting immunity after SARS-CoV-2 infection and vaccination.

The expansion of ASCs in the present cohort of COVID-19 patients highlights an active B cell response to SARS-CoV-2 infection. The frequencies of ASCs within the total circulating B cell population were highly variable between COVID-19 patients, ranging from 3 to 31%. Although the patients in this cohort were sampled at different timepoints following the onset of symptoms, a high variability in ASC frequencies have also been observed among patients infected with influenza or dengue virus sampled at normalized timepoints (e.g., 7 days after febrile illness) (4, 5). Although ASCs have been shown to expand to very high levels in some viral infections, constituting up to 50-80% of all peripheral B cells, the highest ASCs frequency in this cohort of COVID-19 patients was 31% (5, 7, 21). Healthy controls in this study had very low frequencies of ASCs in the circulation, a finding consistent with previous observations (5, 6, 16).

The ASC response in peripheral blood is usually transient and peak at around 7 days after the

onset of febrile illness, dropping to steady-state levels 2-3 weeks later (8). Interestingly, we

observed a substantial ASC expansion as late as 19 days following symptom debut. This

finding highlights the similarity between COVID-19 and respiratory syncytial virus (RSV)

infection, where a significant ASC expansion could be detected as late as 22-45 days after the

onset of symptoms (9). The prolonged ASC expansion in RSV infection was associated with

longer RSV shedding in the airways. Thus, the kinetics of ASC response may be dependent

on pathogen persistence. In support of our findings, a recent report on longitudinal immune

responses in a single COVID-19 patient with mild disease showed the peak of ASC

expansion at day 8 after symptom onset, but a substantial ASCs response could still be

detected by day 20 (22). Thus, it remains to be investigated whether SARS-CoV-2

persistence in the airways may lead to a prolonged ASC response in circulation.

The expansion of ASCs is usually characterized by high specificity towards the infectious

agent (17). Here, we successfully demonstrated that COVID-19 patients generate SARS-

CoV-2-specific ASC responses using a FluoroSpot assay. We were able to successfully

identify SARS-CoV-2-specific IgA-, IgG- and IgM-ASCs in all COVID-19 patients. The

frequency of SARS-CoV-2-specific ASCs within the total ASC population in this patient

cohort was relatively low. However, since only the SARS-CoV-2 N protein was used to

detect specific ASCs in this study, the numbers of N protein-specific ASCs do not reflect the

overall magnitude of the specific ASC response. The use of SARS-CoV-2 viral lysate or

recombinant SARS-CoV-2 proteins in combination would allow for the evaluation of the

overall ASC response. Furthermore, the FluoroSpot assay described in this study could be

adapted to evaluate memory B cell responses to various SARS-CoV-2 proteins in

convalescent COVID-19 patient samples.

The protective level of SARS-CoV-2-neutralizing antibodies in COVID-19 has not yet been

determined. Monoclonal antibodies isolated from convalescent COVID-19 patients showed a

strong SARS-CoV-2-neutralizing capacity *in vitro* and *in vivo* in transgenic mice expressing

human SARS-CoV-2 receptor ACE2 (23). However, extensive longitudinal studies in

humans are required to investigate if recovered COVID-19 patients are protected from re-

infection with SARS-CoV-2. Seroconversion in COVID-19 patients, measured by detectable

SARS-CoV-2-specific IgG levels, has been recently shown to take place within nineteen days

after the onset of symptoms (24). The four patients in the present COVID-19 cohort without

detectable neutralizing antibody titers were sampled at days 7, 8, 9, and 15 after symptom

debut. Although these four patients did not have detectable neutralizing antibody titers,

SARS-CoV-2 specific ASCs were detected. This indicates that the FluoroSpot assay may be

a more sensitive method to detect early B cell responses to SARS-CoV-2, compared to

standard serological assays.

In this cohort of patients, 2 out of the 4 patients with undetectable neutralizing antibody

levels had SARS-CoV-2 RNA in serum. Currently, the extent to which neutralizing

antibodies contribute to the control of viremia in COVID-19 is unclear. Our results seem to

suggest that neutralizing antibodies may not be the only player in SARS-CoV-2 control in

circulation, as one patient with substantial levels of SARS-CoV-2-neutralizing antibodies in

this cohort had detectable serum levels of SARS-CoV-2 RNA. A pilot study assessing the

effectiveness of COVID-19 patient treatment with convalescent plasma containing SARS-

CoV-2-neutralizing antibodies found that the treatment reduced viral load in patients within a

few days after administration, suggesting that neutralizing antibodies have a role in the

management of viremia in COVID-19 (25).

Rapid serology tests are essential for COVID-19 diagnostics and for national serosurveillance

programs. Neutralizing antibody tests, however, are usually laborious and time-consuming.

Therefore, diagnostic tests often rely on other methods such as ELISA. Antibody levels

detected by ELISA, however, do not necessarily reflect the neutralization capacity of patient

serum. Here, we demonstrated a strong correlation between SARS-CoV-2-specific and

SARS-CoV-2-neutralizing antibody titers. These data suggest that the levels of SARS-CoV-

2-specific antibodies measured with ELISA may reflect the capacity of serum from COVID-

19 patients to neutralize SARS-CoV-2.

COVID-19 patients generally present with decreased lymphocyte numbers in peripheral

blood, which is particularly pronounced in severe cases (19, 26, 27). T cells appear to be the

most affected lymphocyte subset. In agreement with these observations, we found decreased

numbers of lymphocytes, and significantly lower T cell numbers in the present cohort of

COVID-19 patients.

In addition to lymphopenia, COVID-19 patients typically display increased concentrations of

inflammatory cytokines in serum, especially in severe cases (19, 27). The levels of

inflammatory markers IL-6 and CRP, in particular, have been shown to correlate with disease

severity in COVID-19 patients (19, 20). In this cohort of COVID-19 patients we also

observed increased levels of IL-6 and CRP, with comparable median values to those

measured by Liu *et al.* in severe COVID-19 patients (20). We show that higher IL-6 and CRP

levels were observed in patients who were hospitalized for longer, suggesting that these

markers may be of use in predicting the duration of COVID-19.

Lastly, we observed an inverse correlation between IL-6 levels and SARS-CoV-2-

neutralizing antibody titers in COVID-19 patients. This relationship may indicate that the

lack of SARS-CoV-2-neutralizing antibodies leads to higher inflammation, or that high levels

of inflammation impair or delay antibody responses. Further studies where a longitudinal

patient sampling at controlled timepoints is performed are required to describe the

relationship between inflammation and humoral responses in COVID-19.

Taken together, we provided a detailed description of clinical and immunological parameters

in twenty COVID-19 patients, with a focus on B cell and antibody responses. We

demonstrated that COVID-19 patients elicit a SARS-CoV-2-specific B cell response,

indicated by the expansion of SARS-CoV-2-specific ASCs. In addition, we showed a clear

relationship between the levels of SARS-CoV-2-specific antibodies and total SARS-CoV-2-

neutralizing antibodies. The tools employed in this study may be of relevance in the

assessment of long-lasting immunity after SARS-CoV-2 infection and vaccination.

Conflict of interest

Authors declare no conflicts of interest.

Acknowledgments

We would like to thank the patients and healthy controls for participating in the study. We

thank the nurses at the Karolinska University Hospital, Stockholm, Sweden for blood

sampling of patients, T. Aktas and M. Olausson at the Public Health Agency of Sweden for

technical assistance, and P. Schierloh at IBB-UNER-CONICET (Argentina) for providing

input on statistical analysis.

References

1. Zhou, P., X.-L. Yang, X.-G. Wang, Ben Hu, L. Zhang, W. Zhang, H.-R. Si, Y. Zhu, B. Li, C.-L. Huang, H.-D. Chen, J. Chen, Y. Luo, H. Guo, R.-D. Jiang, M.-Q. Liu, Y. Chen, X.-R. Shen, X. Wang, X.-S. Zheng, K. Zhao, Q.-J. Chen, F. Deng, L.-L. Liu, B. Yan, F.-X. Zhan, Y.-Y. Wang, G.-F. Xiao, and Z.-L. Shi. 2020. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 579: 270–273.

2. Huang, C., Y. Wang, X. Li, L. Ren, J. Zhao, Y. Hu, L. Zhang, G. Fan, J. Xu, X. Gu, Z. Cheng, T. Yu, J. Xia, Y. Wei, W. Wu, X. Xie, W. Yin, H. Li, M. Liu, Y. Xiao, H. Gao, L. Guo, J. Xie, G. Wang, R. Jiang, Z. Gao, Q. Jin, J. Wang, and B. Cao. 2020. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 395: 497–506.

3. Nutt, S. L., P. D. Hodgkin, D. M. Tarlinton, and L. M. Corcoran. 2015. The generation of antibody-secreting plasma cells. *Nat Immunol* 15: 160–171.

4. Huang, K.-Y. A., C. K.-F. Li, E. Clutterbuck, C. Chui, T. Wilkinson, A. Gilbert, J. Oxford, R. Lambkin-Williams, T.-Y. Lin, A. J. McMichael, and X.-N. Xu. 2014. Virus-Specific Antibody Secreting Cell, Memory B-cell, and Sero-Antibody Responses in the Human Influenza Challenge Model. *J. Infect. Dis.* 209: 1354–1361.

5. Wrammert, J., N. Onlamoon, R. S. Akondy, G. C. Perng, K. Polsrila, A. Chandele, M. Kwissa, B. Pulendran, P. C. Wilson, O. Wittawatmongkol, S. Yoksan, N. Angkasekwinai, K. Pattanapanyasat, K. Chokephaibulkit, and R. Ahmed. 2012. Rapid and massive virus-specific plasmablast responses during acute dengue virus infection in humans. *J. Virol.* 86: 2911–2918.

6. García, M., A. Iglesias, V. I. Landoni, C. Bellomo, A. Bruno, M. T. Córdoba, L. Balboa, G. C. Fernández, M. D. C. Sasiain, V. P. Martínez, and P. Schierloh. 2017. Massive plasmablast response elicited in the acute phase of hantavirus pulmonary syndrome. *Immunology* 151: 122–135.

7. McElroy, A. K., R. S. Akondy, C. W. Davis, A. H. Ellebedy, A. K. Mehta, C. S. Kraft, G.

M. Lyon, B. S. Ribner, J. Varkey, J. Sidney, A. Sette, S. Campbell, U. Ströher, I. Damon, S.

T. Nichol, C. F. Spiropoulou, and R. Ahmed. 2015. Human Ebola virus infection results in

substantial immune activation. *Proc. Natl. Acad. Sci. U.S.A.* 112: 4719–4724.

8. Fink, K. 2012. Origin and function of circulating plasmablasts during acute viral

infections. *Front Immunol* 3: 1–5.

9. Lee, F. E.-H., A. R. Falsey, J. L. Halliley, I. Sanz, and E. E. Walsh. 2010. Circulating

antibody-secreting cells during acute respiratory syncytial virus infection in adults. *J. Infect.*

Dis. 202: 1659–1666.

10. Wrammert, J., K. Smith, J. Miller, W. A. Langley, K. Kokko, C. Larsen, N.-Y. Zheng, I.

Mays, L. Garman, C. Helms, J. James, G. M. Air, J. D. Capra, R. Ahmed, and P. C. Wilson.

2008. Rapid cloning of high-affinity human monoclonal antibodies against influenza virus.

Nature 453: 667–671.

11. Qian, Y., C. Wei, F. Eun-Hyung Lee, J. Campbell, J. Halliley, J. A. Lee, J. Cai, Y. M.

Kong, E. Sadat, E. Thomson, P. Dunn, A. C. Seegmiller, N. J. Karandikar, C. M. Tipton, T.

Mosmann, I. Sanz, and R. H. Scheuermann. 2010. Elucidation of seventeen human peripheral

blood B-cell subsets and quantification of the tetanus response using a density-based method

for the automated identification of cell populations in multidimensional flow cytometry data.

Cytometry B Clin Cytom 78: S69–82.

12. Crotty, S., R. D. Aubert, J. Glidewell, and R. Ahmed. 2004. Tracking human antigen-

specific memory B cells: a sensitive and generalized ELISPOT system. *J. Immunol. Methods*

286: 111–122.

13. Murin, C. D., I. A. Wilson, and A. B. Ward. 2019. Antibody responses to viral infections:

a structural perspective across three different enveloped viruses. *Nature Microbiology* 4:

734–747.

14. Corman, V. M., O. Landt, M. Kaiser, R. Molenkamp, A. Meijer, D. K. Chu, T. Bleicker,

S. Brünink, J. Schneider, M. L. Schmidt, D. G. Mulders, B. L. Haagmans, B. van der Veer, S.

van den Brink, L. Wijsman, G. Goderski, J.-L. Romette, J. Ellis, M. Zambon, M. Peiris, H.

Goossens, C. Reusken, M. P. Koopmans, and C. Drosten. 2020. Detection of 2019 novel

coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill.* 25: 2431.

15. National Health Commission State Administration of Traditional Chinese Medicine.

2020. Diagnosis and Treatment Protocol for Novel Coronavirus Pneumonia (Trial Version 7).

16. Mei, H. E., T. Yoshida, W. Sime, F. Hiepe, K. Thiele, R. A. Manz, A. Radbruch, and T.

Doerner. 2009. Blood-borne human plasma cells in steady state are derived from mucosal

immune responses. *Blood* 113: 2461–2469.

17. Lee, F. E. H., J. L. Halliley, E. E. Walsh, A. P. Moscatiello, B. L. Kmush, A. R. Falsey,

T. D. Randall, D. A. Kaminiski, R. K. Miller, and I. Sanz. 2011. Circulating Human

Antibody-Secreting Cells during Vaccinations and Respiratory Viral Infections Are

Characterized by High Specificity and Lack of Bystander Effect. *J. Immunol.* 186: 5514–

5521.

18. Manenti, A., M. Maggetti, E. Casa, D. Martinuzzi, A. Torelli, C. M. Trombetta, S.

Marchi, and E. Montomoli. 2020. Evaluation of SARS-CoV-2 neutralizing antibodies using a

CPE-based colorimetric live virus micro-neutralization assay in human serum samples. *J.*

Med. Virol. 38: 1.

19. Zhang, X., Y. Tan, Y. Ling, G. Lu, F. Liu, Z. Yi, X. Jia, M. Wu, B. Shi, S. Xu, J. Chen,

W. Wang, B. Chen, L. Jiang, S. Yu, J. Lu, J. Wang, M. Xu, Z. Yuan, Q. Zhang, X. Zhang, G.

Zhao, S. Wang, S. Chen, and H. Lu. 2020. Viral and host factors related to the clinical

outcome of COVID-19. *Nature* 1–7.

20. Liu, F., L. Li, M. Xu, J. Wu, D. Luo, Y. Zhu, B. Li, X. Song, and X. Zhou. 2020.

Prognostic value of interleukin-6, C-reactive protein, and procalcitonin in patients with

COVID-19. *J. Clin. Virol.* 127: 104370.

21. Garcia-Bates, T. M., M. T. Cordeiro, E. J. M. Nascimento, A. P. Smith, K. M. Soares de

Melo, S. P. McBurney, J. D. Evans, E. T. A. Marques, and S. M. Barratt-Boyes. 2013.

Association between magnitude of the virus-specific plasmablast response and disease

severity in dengue patients. *J. Immunol.* 190: 80–87.

22. Thevarajan, I., T. H. O. Nguyen, M. Koutsakos, J. Druce, L. Caly, C. E. Sandt, X. Jia, S.

Nicholson, M. Catton, B. Cowie, S. Y. C. Tong, S. R. Lewin, and K. Kedzierska. 2020.

Breadth of concomitant immune responses prior to patient recovery: a case report of non-

severe COVID-19. *Nat. Med.* 1–10.

23. Cao, Y., B. Su, X. Guo, W. Sun, Y. Deng, L. Bao, Q. Zhu, X. Zhang, Y. Zheng, C. Geng,

X. Chai, R. He, X. Li, Q. Lv, H. Zhu, W. Deng, Y. Xu, Y. Wang, L. Qiao, Y. Tan, L. Song,

G. Wang, X. Du, N. Gao, J. Liu, J. Xiao, X.-D. Su, Z. Du, Y. Feng, C. Qin, C. Qin, R. Jin,

and X. S. Xie. 2020. Potent neutralizing antibodies against SARS-CoV-2 identified by high-

throughput single-cell sequencing of convalescent patients' B cells. *Cell* 183: 1–12.

24. Long, Q.-X., B.-Z. Liu, H.-J. Deng, G.-C. Wu, K. Deng, Y.-K. Chen, P. Liao, J.-F. Qiu,

Y. Lin, X.-F. Cai, D.-Q. Wang, Y. Hu, J.-H. Ren, N. Tang, Y.-Y. Xu, L.-H. Yu, Z. Mo, F.

Gong, X.-L. Zhang, W.-G. Tian, L. Hu, X.-X. Zhang, J.-L. Xiang, H.-X. Du, H.-W. Liu, C.-

H. Lang, X.-H. Luo, S.-B. Wu, X.-P. Cui, Z. Zhou, M.-M. Zhu, J. Wang, C.-J. Xue, X.-F. Li,

L. Wang, Z.-J. Li, K. Wang, C.-C. Niu, Q.-J. Yang, X.-J. Tang, Y. Zhang, X.-M. Liu, J.-J. Li,

D.-C. Zhang, F. Zhang, P. Liu, J. Yuan, Q. Li, J.-L. Hu, J. Chen, and A.-L. Huang. 2020.

Antibody responses to SARS-CoV-2 in patients with COVID-19. *Nat. Med.* 1–15.

25. Duan, K., B. Liu, C. Li, H. Zhang, T. Yu, J. Qu, M. Zhou, L. Chen, S. Meng, Y. Hu, C.

Peng, M. Yuan, J. Huang, Z. Wang, J. Yu, X. Gao, D. Wang, X. Yu, L. Li, J. Zhang, X. Wu,

B. Li, Y. Xu, W. Chen, Y. Peng, Y. Hu, L. Lin, X. Liu, S. Huang, Z. Zhou, L. Zhang, Y.

Wang, Z. Zhang, K. Deng, Z. Xia, Q. Gong, W. Zhang, X. Zheng, Y. Liu, H. Yang, D. Zhou,

D. Yu, J. Hou, Z. Shi, S. Chen, Z. Chen, X. Zhang, and X. Yang. 2020. Effectiveness of

convalescent plasma therapy in severe COVID-19 patients. *Proc. Natl. Acad. Sci. U.S.A.* 117:

9490–9496.

26. Wang, F., J. Nie, H. Wang, Q. Zhao, and Y. X. 2020. Characteristics of Peripheral

Lymphocyte Subset Alteration in COVID-19 Pneumonia. *J. Infect. Dis.* 221: 1762–1769.

27. Qin, C., L. Zhou, Z. Hu, S. Zhang, S. Yang, Y. Tao, C. Xie, K. Ma, K. Shang, W. Wang,

and D.-S. Tian. 2020. Dysregulation of immune response in patients with COVID-19 in

Wuhan, China. *Clin. Infect. Dis.* 1-7

Footnotes

1. Grant support

This work was supported by the Marianne and Marcus Wallenberg Foundation (SGR), and

the KID PhD student funding program from Karolinska Institutet (SGR), the Swedish

Research Council (JK), and the SciLifeLab national COVID-19 research program (JK).

2. Abbreviations

ASC – antibody-secreting cell

CPE – cytopathic effect

COVID-19 – coronavirus disease 2019

N protein – nucleocapsid protein

SARS-CoV-2 – severe acute respiratory syndrome coronavirus 2

S1 protein – subunit 1 of SARS-CoV-2 spike protein

3. Corresponding author

Sara Gredmark-Russ, Center for Infectious Medicine, ANA Futura, Department of Medicine,

Karolinska Institutet, Karolinska University Hospital Huddinge, 141 52 Stockholm, Sweden,

Telephone: +46 8 58589688, Fax: +46 8 7467637, E-mail: sara.gredmark.russ@ki.se

Figure legends

Figure 1. Clinical chemistry parameters in peripheral blood of COVID-19 patients.

(A) Number of days since the onset of COVID-19 symptoms that peripheral blood samples

were taken. (B) Number of days since hospitalization that peripheral blood samples were

taken. (C) Blood cell counts and clinical chemistry parameters in peripheral blood of

COVID-19 patients measured on the day of the inclusion in this study (+/− 24 hours). Gray

boxes indicate the range for reference values. (D) Absolute numbers of $CD45^+$, $CD3^+$, $CD4^+$,

$CD8^+$, and $CD19^+$ cells in peripheral blood of COVID-19 patients (n = 20) and healthy

controls (HC) (n = 7) measured by flow cytometry. Statistical significance was determined

using Mann-Whitney *U* test (D). Graphs display median and IQR. n = number of patients.

CRP – C-reactive protein; LDH – lactate dehydrogenase; AST – aspartate transaminase; ALT

– alanine transaminase; PCT – procalcitonin; DL – detection limit. *, P < 0.05; **, P < 0.01;

***, P < 0.001; ns – not significant.

Figure 2. T cells and B cells are activated in peripheral blood of COVID-19 patients.

(A) Representative flow cytometry plots of CD38 and Ki-67 co-expression in $CD4^+$ and

$CD8^+$ T cells, as well as $CD19^+$ B cells in one representative COVID-19 patient (16 days

after symptom onset) and one healthy control (HC). (B) Frequencies of CD38 and Ki-67 co-

expressing $CD4^+$ T cells, $CD8^+$ T cells and $CD19^+$ B cells in COVID-19 patients (n = 20) and

healthy donors (n = 7). Bar graphs display median and IQR. Statistical significance was

determined using Mann-Whitney *U* test. *, P < 0.05; ***, P < 0.001.

Figure 3. SARS-CoV-2 nucleocapsid protein-specific antibody-secreting cells (ASCs)

expand in COVID-19 patients. (A) Flow cytometry gating strategy used to identify ASCs

and the IgA-, IgG-, and IgM-ASC subsets. B cells are gated on live $CD14^-CD123^-CD3^-CD4^-$

cells. (B) Frequencies of ASCs within all B cells in COVID-19 patients and in HCs. ASCs

were defined as $CD19^+CD20^{low/-}IgD^-CD38^{high}CD27^{high}$. (C) Numbers of ASCs per microliter

of whole blood, calculated using absolute B cell numbers and frequencies of ASCs measured

by flow cytometry. (D) Frequencies of IgA-, IgG-, and IgM-ASCs within the total ASC

population measured by flow cytometry. (E and F) Representative images of wells from a

FluoroSpot assay showing total IgA-, IgG-, and IgM-ASCs (E), and SARS-CoV-2

nucleocapsid protein-specific ASCs (F) from one COVID-19 patient (13 days after symptom

onset) and one healthy control. IgM fluorescence is originally red, but replaced with white in

this figure for visualization purpose. (G) Numbers of total IgA-, IgG-, and IgM-ASCs per

million PBMCs, as measured by FluoroSpot assay. (H) Numbers of SARS-CoV-2

nucleocapsid protein-specific IgA-, IgG-, and IgM-ASCs per million PBMCs. Experiments

were performed on all COVID-19 patients (n = 20) and healthy controls (n = 7). Statistical

significance was determined using Mann-Whitney U test (B, C, D and G). Bar graphs display

median and IQR. **, P < 0.01; ***, P < 0.001; ns – not significant.

Figure 4. SARS-CoV-2-specific and neutralizing antibody levels in COVID-19 patients.

(A) Individual antibody responses to SARS-CoV-2 in COVID-19 patients (n = 20) and

healthy controls (HC) (n = 7). Positivity for IgA against S1-protein, IgG against S1-protein,

IgM against N protein, total IgG antibodies against whole SARS-CoV-2, and SARS-CoV-2-

neutralizing antibody titers are presented in the heatmap. Blue color indicates a positive

response and the color scale is adjusted for the minimum positive assay value and the highest

value recorded within the patient cohort for each assay. White boxes indicate values below

positive threshold or below detection level for each assay. Red asterisks represent patients

with detectable levels of SARS-CoV-2 RNA in serum. (B) IgA, IgG, and IgM antibody levels

in COVID-19 patients and controls, analyzed by ELISAs. Dotted horizontal line indicates the

threshold for positive result. OD, optical density. OD ratio = OD of the sample divided by

OD of the calibrator. (C) Total SARS-CoV-2 IgG antibody titers determined by

immunofluorescence assay. Patients with titers < 20 were assigned a value of 1. (D) SARS-

CoV-2-neutralizing antibody titers determined by microneutralization assay. Patients with

titers < 10 were assigned a value of 1. (E) Correlation between SARS-CoV-2-specific IgA,

IgG and IgM antibody titers and SARS-CoV-2-neutralizing antibody titers, tested with

Spearman's correlation test. r_s: Spearman's rank correlation coefficient. P < 0.05 was

considered statistically significant. Four data points from COVID-19 patients with identical

values are highlighted (5, 7, 10 and 13). Bar graphs display median and IQR.

Figure 5. Correlations between the levels of IL-6, CRP, neutralizing antibodies and the

duration of COVID-19. (A) Serum concentrations of IL-6 in COVID-19 patients and

healthy controls. Bar graph displays median and IQR. Statistical significance was determined

using Mann-Whitney *U* test. (B) Correlation matrix for serum IL-6, CRP, neutralizing

antibody titers, duration of hospitalization and the number of days between symptom onset to

discharge from hospital. Correlations were examined by Spearman's correlation test. The

color scale represents the range values for Spearman's rank correlation coefficient (r_s). *, P <

0.05; **, P < 0.01; ***, P < 0.001.

Table I. Clinical characteristics of 20 hospitalized COVID-19 patients.

Cohort characteristics	
Age, years, median (range)	53 (34-67)
Male, n (%)	15 (75)
Symptom onset to sampling, days, median (range)	15 (7-19)
Hospitalization to sampling, days, median (range)	3 (2-10)
Symptom onset to discharge, days, median (range)	19.5 (9-30)
Duration of hospitalization, days, median (range)	7.5 (3-25)
Intensive care unit (ICU) treatment[a], n (%)	3 (15)
Fatal outcome, n (%)	1 (5)
Comorbidities	**n (%)**
Overweight (BMI >25), n/N (%)	14/14 (100)[b]
Diabetes mellitus type II	4 (20)
Hypertension	3 (15)
Asthma	2 (10)
Cardiovascular disease	1 (5)
Symptoms	**n (%)**
Fever (>38°C)	20 (100)
Cough	20 (100)
Respiratory difficulties (SpO$_2$ ≤ 93% on room air)	20 (100)
Myalgia	9 (45)
Chest pain	6 (30)
Diarrhea	4 (20)
Treatment	**n (%)**
Supplemental oxygen[c]	20 (100)
High flow nasal oxygen	4 (20)
Inhaled bronchodilators	2 (10)
Low molecular weight heparin	17 (85)
Antibiotics	14 (70)
Immunomodulatory drugs[d] (given before sampling)	3 (15)

Abbreviations: n: number of patients; BMI: body mass index. SpO$_2$: percutaneous oxygen saturation.

[a] ICU treatment before study sampling (n=2), ICU treatment after study sampling (n=1).
[b] n/N (%), where N is the total number of patients with available data.
[c] Oxygen flow rate – minimum 2.5 L/min. 1 patient in mechanical ventilation.
[d] Chloroquine phosphate (n=2) or Anakinra (n=1).

Figure 1 Varnaite et al.

Figure 2 Varnaite et al.

Figure 3 Varnaite et al.

Figure 4 Varnaite et al.

Figure 5 Varnaite et al.

(A) Flow cytometry gating strategy for antibody-secreting cells (ASCs), the IgA, IgG and IgM ASC subsets, as well as activated B cells. (B) Flow cytometry gating strategy for CD4[+] and CD8[+] T cell activation, defined as the co-expression of CD38 and Ki-67.

Supplementary Figure 2

(A) Correlation between CD4+ and CD8+ T cell activation levels, as measured by CD38 and Ki-67 co-expression using flow cytometry. (B) Graph displaying the antibody-secreting cell (ASC) frequencies within the total pool of CD19+ B cells and the number of days since symptom onset of COVID-19. (C) Correlation between total IgG titers against SARS-CoV-2 (measured by IFA) and the number of days since symptom onset of COVID-19. (D) Correlation between levels of SARS-CoV-2 S1 protein-specific IgG levels and the number of days since symptom onset of COVID-19. (E-G) Correlations between activated T cells and B cells with SARS-CoV-2 neutralizing antibody titers. Correlations were examined by Spearman's correlation test. r_s: Spearman's rank correlation coefficient. $P < 0.05$ was considered statistically significant.