Running Title: mEos3.2 photophysics in cells

Sample preparation and imaging conditions affect mEos3.2 photophysics in fission yeast cells

Mengyuan Sun, Kevin Hu, Joerg Bewersdorf[*] and Thomas D. Pollard[*]

[1]Departments of Molecular Biophysics and Biochemistry

[2]Departments of Molecular Cellular and Developmental Biology

[3]Department of Cell Biology

[4]Department of Biomedical Engineering

Yale University, PO Box 208103

New Haven, CT 06520-8103 USA

[*]Co-corresponding authors

Abstract

Photoconvertible fluorescent proteins (PCFPs) are widely used in super-resolution microscopy and studies of cellular dynamics. However, our understanding of their photophysics is still limited, hampering their quantitative application. For example, we do not know the optimal sample preparation methods or imaging conditions to count protein molecules fused to PCFPs by single-molecule localization microscopy in live and fixed cells. We also do not know how the behavior of PCFPs in live cells compares with fixed cells. Therefore, we investigated how formaldehyde fixation influences the photophysical properties of the popular green-to-red PCFP mEos3.2 in fission yeast cells under a wide range of imaging conditions. We estimated photophysical parameters by fitting a 3-state model of photoconversion and photobleaching to the time course of fluorescence signal per yeast cell expressing mEos3.2. We discovered that formaldehyde fixation makes the fluorescence signal, photoconversion rate and photobleaching rate of mEos3.2 sensitive to the buffer conditions by permeabilizing the yeast cell membrane. Under some imaging conditions, the time-integrated mEos3.2 signal per yeast cell is similar in live cells and fixed cells imaged in buffer at pH 8.5 with 1 mM DTT, indicating that light chemical fixation does not destroy mEos3.2 molecules. We also discovered that some red-state mEos3.2 molecules entered an intermediate dark state that is converted back to the red fluorescent state by 561-nm illumination. Our findings provide a guide to compare quantitatively conditions for imaging and counting of mEos3.2-tagged molecules in yeast cells. Our imaging assay and mathematical model are easy to implement and provide a simple quantitative approach to measure the time-integrated signal and the photoconversion and photobleaching rates of fluorescent proteins in cells.

STATEMENT OF SIGNIFICANCE

Making quantitative measurements with single-molecule localization microscopy (SMLM) has been impeded by limited understanding of the photophysics of the fluorophores, which is very sensitive to the sample preparation and imaging conditions. We characterized the photophysics of the green-to-red photoconvertible fluorescent protein mEos3.2, which is widely used in SMLM. We combined quantitative fluorescence microscopy and mathematical modeling to measure the fluorescence signal and rate constants for photoconversion and photobleaching of mEos3.2 in live and fixed cells under a wide range of illumination intensities. Our findings

46   provide a guide to compare conditions for imaging and counting mEos3.2-tagged proteins in

47   cells. The presented approach is generally applicable to characterize other fluorescent proteins or

48   dyes in cells.

INTRODUCTION

Photoactivatable or photoconvertible fluorescent proteins (PAFPs or PCFPs) have enabled super-resolution imaging by temporally separating closely-spaced molecules (1-3). The fluorescent protein EosFP (4) and its derivatives (5-7) have been widely used in SMLM for both live (8-10) and fixed biological samples (11-13). The fluorescent protein is fused to the coding sequence of a protein of interest in the genome for endogenous expression or expressed exogenously and transiently in cells. Irradiation at 405 nm photoconverts EosFPs from their native green state with an emission peak at 516 nm to their red state with an emission peak at 580 nm (4, 14, 15). In SMLM, sparsely distributed photoconverted red EosFPs are excited at 561 nm and then localized (13). The rationally designed, monomeric derivative of EosFP mEos3.2 is favored by many due to its monomeric property, high brightness, photostability, and compatibility with live cells (6).

Counting fluorescently-tagged fusion proteins is a potential strength of SMLM, as the images are assembled from discrete localizations of individual molecules (16, 17). The total number of localizations in the SMLM images is closely correlated to the total number of fusion proteins, which allows the measurement of this important quantity even in a diffraction-limited subcellular structure (9, 11, 12, 18-23). Genetically encoded tagging with PAFPs or PCFPs can ensure 1:1 labeling stoichiometry (16), without the uncertainties associated with extrinsic labeling techniques (22, 24-28). However, even with genetically encoded tags, quantitative SMLM still faces several challenges that can lead to undercounting or overcounting the molecules. Fluorescent proteins mature slowly, so an unknown fraction of the FPs is fluorescent at the time of imaging (29). Some of the PAFPs or PCFPs might never be photoconverted or photoactivated to the active state for SMLM imaging (30). Moreover, activated PAFPs or PCFPs may blink by entering a transient dark state and return to the fluorescent state an unknown number of times, which can lead to overcounting (18, 20). A promising way of doing quantitative SMLM is to obtain numbers of the tagged molecules relative to internal calibration standards of known number (9, 11, 12, 21, 23). The sources of error mentioned above can be accounted for if the target and calibration standards are prepared, imaged, and analyzed consistently in the same way (11, 12, 16, 31).

However, any inconsistency in the process of sample preparation, SMLM imaging, or data analysis can introduce errors. Diffusion and other movements of the fluorescent protein can

further complicate the quantification process, so light chemical fixation can be used to preserve

the targeted structure and eliminate movements for quantitative SMLM of cells expressing

proteins tagged with mEos2 (11, 12). However, fixation can introduce measurement errors. For

example, fixation might destroy some FPs or change their photophysical properties (32), which

can change the average number of localizations for the FP. Therefore, one must understand how

fixation and sample preparation affect the mEos3.2 photophysical parameters related to single-

molecule imaging and counting. Photoconversion and photobleaching rates determine the density

of active fluorophores at each frame, which is essential for separating closely-spaced molecules.

The fluorescence signal of fluorescence fusion proteins in the structure of interest contains

information about the brightness of individual molecules and the number of molecules able to

fluoresce, both aspects being important for molecule counting with diffraction-limited (31) and

super-resolution imaging (17).

In this study, we measured the photophysical properties of mEos3.2 in fission yeast cells

by fitting a 3-state model of photoconversion and photobleaching to the time course of the

mEos3.2 fluorescence signal per cell measured by quantitative fluorescence microscopy. We

found that formaldehyde fixation permeabilized the *S. pombe* cells for small molecules, making

the photophysical properties of mEos3.2 sensitive to the buffer conditions. To find conditions

where the mEos3.2 photophysical parameters are comparable in live and fixed yeast cells, we

investigated the effect of fixation and imaging buffer under a wide range of imaging conditions

with point-scanning and widefield illumination. We also discovered that a subpopulation of red-

state mEos3.2 molecules entered an intermediate state that is converted back to the red

fluorescent state by illumination at 561-nm but not 405-nm. Our data provide information on

sample preparation for imaging and counting mEos3.2 in live and fixed yeast cells. Our

quantitative imaging assay combined with the 3-state model can be applied to study the

photophysical properties of other PAFPs and PCFPs quantitatively without doing single-

molecule localization.

MATERIALS AND METHODS

Plasmids and *S. pombe* Strains

The open reading frame encoding mEos3.2 was cloned into the pJK148-pAct1-nmt1Term

plasmid with PCR and NEB HiFi Builder Assembly. Both the newly constructed plasmid and

chromosomal insertion were verified by sequencing.

Preparation of *S. pombe* Cells for Imaging

S. pombe cells expressing mEos3.2 were grown in exponential phase at 25 °C in YE5S-rich

liquid medium in 50-mL flasks in the dark before switching to EMM5S-synthetic medium ~12-

18 hours before imaging to reduce the cellular autofluorescence background. Live cells were

concentrated 10- to 20-fold by centrifugation at 3,000 rpm for 30 s and resuspended in EMM5S

for imaging.

Cells were fixed by mixing an equal volume of fresh, room temperature 4%

formaldehyde aqueous solution (EMS) with the cell culture and shaking at 150 rpm at 25° C for

15 min or 30 min. The cells were pelleted by centrifugation at 3,000 rpm for 30 s and washed by

pelleting in EMM5S or other buffers 3 times, and then resuspended in EMM5S or other buffers.

Concentrated cells in 5 µL were mounted on a thin layer formed from 35 μL of 25% gelatin

(Sigma-Aldrich; G-2500) in EMM5S or other buffers (without the DTT). To assess how pH and

reducing agent affect the photophysical properties of mEos3.2, we fixed cells for 30 min and

then washed and resuspended the cells for imaging in one of the following buffers: 50 mM MES

(pH 5.5), 50 mM MES (pH 6.5), 50 mM Tris-HCl (pH 7.5), 50 mM Tris-HCl (pH 8.5), 50 mM

Tris-HCl (pH 7.5) with 1 mM DTT, 50 mM Tris-HCl (pH 8.5) with 1 mM DTT, or 50 mM Tris-

HCl (pH 8.5) with 10 mM DTT.

Point-scanning Confocal Imaging Conditions

Time lapse videos were acquired on a Zeiss Laser Scanning Microscope (LSM 880) using an

alphaPlan-Apochromat 100x/NA 1.46 oil-immersion objective and an emission band path filter

collecting fluorescence in the 566 - 719 nm wavelength range. Samples were illuminated by

scanning a field of view (FOV) of 85 x 85 µm (512 x 512 pixels; 160 nm pixel size) with both

the 405 nm and 561 nm lasers at constant intensities. To test different imaging conditions, we set

the 405 nm laser power at the sample constant ranging from 16 to 56 µW and the 561 nm laser

power constant ranging from 11 to 37 µW. To compare with widefield imaging conditions, we

estimated the average intensity and the peak intensity of point-scanning illumination. The

average intensities in the FOV (power at the sample divided by the FOV area) were 0.22 to 0.78

W/cm^2 at 405 nm and 0.15 to 0.51 W/cm^2 at 561 nm. The peak intensities (power at the sample

divided by the size of the point spread function) were ~80 to 240 kW/cm^2 at 405 nm and ~20 to

80 kW/cm^2 at 561 nm. A Z-stack of 19 slices spaced at 600-nm intervals was acquired with a

pixel dwell time of 0.85 μs. The total exposure time for each Z-stack was 4.23 s (0.85 μs x 512 x

512 x 19). An entire time lapse data set consisted of 50 or 100 Z-stacks at an acquisition rate of

approximately 4 Z-stacks per minute (due to overhead in the scan process).

For experiments with alternating 405 and 561-nm laser illumination, the 561-nm laser

scanned the FOV for 10 cycles followed by 405-nm laser illumination for 5 cycles with either no

break or a 2 min break between the 405-nm period and the following 561-nm illumination period.

The laser powers at the sample were 56 μW at 405 nm and 37 μW at 561 nm. This procedure

was repeated 7 times. We increased the temporal resolution by reducing the pixel number in the

85 x 85 μm FOV to 128 x 128 (640 nm pixel size) with a pixel dwell time of 3.39 μs. This

approach reduced the total exposure time for each Z-stack to 1.05 s (3.39 μs x 128 x 128 x 19).

These 3D stacks were collected with 40 cycles of 561-nm illumination followed by 20 cycles of

405-nm irradiation without breaks between the 405-nm and 561-nm illumination periods. The Z-

stacks were acquired at a rate of approximately 5 Z-stacks per minute due to the scanning

overhead. This procedure was repeated 7 times.

For GFP photobleaching measurements, wildtype cells and cells expressing Fim1-GFP

(TP347) were excited at 488 nm (~60 μW at the sample) and emission fluorescence in the range

of 505-735 nm was collected. A 19-slice Z stack with 600-nm Z-step intervals covering a FOV

of 82 x 82 μm (512 x 512 pixels; 160 nm pixel size) was imaged at each time point with a pixel

dwell time of 0.85 μs. The entire time-lapse data set consisted of 100 Z-stacks.

Widefield Fluorescence Imaging Conditions

Time lapse videos were acquired with a custom-built single-molecule localization microscope

(SMLM) based on a Leica DMi8 stand with widefield illumination, a 63x/1.47 NA oil-

immersion objective, and a band pass filter to collect emission fluorescence in the 584-676 nm

wavelength range. Samples were illuminated at both 405 nm and 561 nm and imaged with an

sCMOS camera (Hamamatsu ORCA-Flash4.0 V2) at 50 frames per second (fps) for 15,000

frames. To test different imaging conditions, the 405-nm laser intensity was set constant in a

range from 0.5 to 2 W/cm^2, and the 561-nm laser intensity from 1 W/cm^2 to 1 kW/cm^2. A

custom-written LABVIEW program was used to control the lasers for alternating 405-nm and

561-nm illumination.

Image Analysis

Images recorded by confocal and widefield fluorescence microscopes were viewed and analyzed

in Fiji (Fiji is Just ImageJ) (33). We made a sum projection of the 19-slice Z-stacks of the time-

lapse confocal images. We manually selected a region of interest (ROI) 1 (containing typically

~50-100 cells for the confocal images and ~ 25 cells for the widefield images) with the polygon

tool and selected the background ROI 2 with the square tool (Fig. S1). The area and mean signal

per pixel (MSPP) of both ROIs were measured and the fluorescence signal per cell at each time

point was calculated based on: [Area 1 * (MSPP 1 –MSPP 2)] / number of cells in ROI 1. We

calculated the weighted mean and standard deviation of the fluorescence signal per cell from all

the FOVs included for each condition, weighted by the number of cells in each FOV. To correct

for autofluorescence background, we subtracted the autofluorescence signal per wildtype cell at

each time point from the fluorescence signal per cell expressing mEos3.2.

Analytical Model of the Time Course of Red mEos3.2 Fluorescence Signal per Cell

Our three-state model (Fig. 1C) considers mEos3.2 molecules to have 3 different states: a non-

activated green (G) state, an activated red (R) state, and a bleached (B) state. Photoconversion

converts molecules from the G- to the R-state by an irreversible first-order reaction with a rate

constant of k_{act} ($k_{activation}$). Molecules in the R-state emit red photons until is photobleached to the

B state by an irreversible first-order reaction with a rate constant of k_{bl} ($k_{bleaching}$). With

illumination at 405 nm and 561 nm, the rates of change in the numbers (n) of G-, R-, and B-state

mEos3.2 molecules are described by the following differential equations (Fig. S2A):

191
$$\frac{dG_n(t)}{dt} = -k_{act}G_n(t) \qquad (1)$$

192
$$\frac{dR_n(t)}{dt} = k_{act}G_n(t) - k_{bl}R_n(t) \qquad (2)$$

193
$$\frac{dB_n(t)}{dt} = k_{bl}R_n(t) \qquad (3)$$

194   We defined the total number of mEos3.2 molecules in a cell as M_n, and assumed that all

195 mEos3.2 molecules were in the green state at the start of the experiment, so $G_n(t = 0) = M_n$, $R_n(t$

196 $= 0) = 0$, and $B_n(t = 0) = 0$. Solving the system of differential equations analytically resulted in

197 the following equations for the number of G-, R-, and B-state molecules in a cell changing over

198 time:

199
$$G_n(t) = M_n e^{-k_{act}t} \qquad (4)$$

200
$$R_n(t) = \frac{M_n k_{act}}{k_{bl}-k_{act}} (e^{-k_{act}t} - e^{-k_{bl}t}) \qquad (5)$$

201
$$B_n(t) = \frac{M_n}{k_{bl}-k_{act}}[k_{bl}(1-e^{-k_{act}t}) - k_{act}(1-e^{-k_{bl}t})] \quad (6)$$

Eq. 5 describes how the number of R-state mEos3.2 molecules in a cell (R_n) changes with

continuous photoconversion and photobleaching. Assuming that the signal of an R-state

molecule per frame is ε_f, the fluorescence signal from the red mEos3.2 molecules per cell (R_s) at

each frame recorded at a given time t is $R_s(t) = R_n(t) \times \varepsilon_f$. Multiplying both sides of Eq. 5 by ε_f

gives Eq. 7 that describes how the fluorescence signal per cell in each frame $R_s(t)$ changes over

time with continuous photoconversion and photobleaching:

208
$$R_s(t) = \frac{M_n \varepsilon_f k_{act}}{k_{bl}-k_{act}}(e^{-k_{act}t} - e^{-k_{bl}t}) \quad (7)$$

We estimated $M_n \times \varepsilon_f$, k_{act}, and k_{bl} using Levenberg-Marquardt nonlinear least squares

regression to fit Eq. 7 to the time course of fluorescence signal per cell $R_s(t)$. We calculated the

95% confidence intervals (CI) of the fitted parameters from the covariance argument of the fit.

For the confocal experiments, we fit Eq. 7 to the weighted average time course of fluorescence

signal per cell from all FOVs and report the 95% CI of the fitted parameters. For the widefield

illumination experiments, we fit Eq. 7 individually to the time course of fluorescence signal per

cell (after autofluorescence background subtraction) from each FOV. We calculated the mean

fitted parameters and standard deviation weighted by the number of cells in each FOV.

Since $R_s(t)$ is the fluorescence signal from red mEos3.2 molecules per cell in each frame,

the fluorescence signal from red mEos3.2 molecules per cell per second is $R_s(t) \times f$ (frame rate,

fps). Integrating the function $R_s(t) \times f$ with respect to time (t, second) over the interval of [0, ∞]

gives the time-integrated signal $\overline{R_s}$ of mEos3.2 per cell:

221
$$\overline{R_s} = \int_0^\infty [f \cdot R_s(t)]dt = \int_0^\infty \frac{fM_n \varepsilon_f k_{act}}{k_{bl}-k_{act}}(e^{-k_{act}t} - e^{-k_{bl}t})dt = \frac{fM_n \varepsilon_f}{k_{bl}} \quad (8)$$

We used Eq. 8 to calculate $\overline{R_s}$ using the parameters $M_n \times \varepsilon_f$ and k_{bl} from the previous fit

of Eq. 7. We estimated the 95% CI or standard deviation through error propagation.

Experiments with alternating illumination at 405 nm and 561 nm revealed a fourth

intermediate (I) state of mEos3.2 molecules (Fig. 6). Illumination at 405 nm converts mEos3.2

molecules in the G-state into the R-state and molecules in the G- and/or R-state into the I-state.

Irradiation at 561 nm converts the mEos3.2 molecules from the I-state to R-state with an

activation rate constant of $k_{act,561}$. R-state molecules are photobleached with a bleaching rate

constant of k_{bl}. During a 561-nm illumination period after a previous 405-nm illumination period,

we assumed that no mEos3.2 molecules in the G- and/or R-state converted to the I-state and I-

states molecules were not photobleached. The rates of change in the numbers (n) of I-, R-, and B-
state mEos3.2 molecules are described by the following differential equations:

$$\frac{d I_n(t)}{dt} = -k_{act,561} I_n(t) \qquad (9)$$

$$\frac{d R_n(t)}{dt} = k_{act,561} I_n(t) - k_{bl} R_n(t) \qquad (10)$$

$$\frac{d B_n(t)}{dt} = k_{bl} R_n(t) \qquad (3)$$

We defined the total number of mEos3.2 molecules, i.e. the sum of molecules in the I-,
R- and B-states in a cell, after the previous 405-nm irradiation as S_n. We further defined $t = 0$ as
the time at which 561-nm illumination starts and assumed that S_n is constant during the 561-nm
illumination period, since the 405-nm laser was off and conversion of molecules to the I-state by
561-nm light is negligible. We further assume that the number of molecules in the different states
at the beginning of the 561-nm illumination period is $R_n(t = 0) = R_{n,0}$, $I_n(t = 0) = S_n - R_{n,0}$, and
$B_n(t = 0) = 0$. Solving the system of differential equations analytically resulted in the following
equations for the number of I-, R-, B-state molecules changing over time:

$$I_n(t) = \left(S_n - R_{n,0}\right)e^{-k_{act,561}t} \qquad (11)$$

$$R_n(t) = \frac{1}{k_{bl} - k_{act,561}}\left[\left(S_n - R_{n,0}\right)k_{act,561}e^{-k_{act,561}t} - \left(S_n k_{act,561} - R_{n,0}k_{bl}\right)e^{-k_{bl}t}\right] \qquad (12)$$

$$B_n(t) = \frac{S_n}{k_{bl} - k_{act,561}}\left[k_{bl}\left(1 - e^{-k_{act,561}t}\right) - k_{act,561}\left(1 - e^{-k_{bl}t}\right)\right]$$

$$+ \frac{R_{n,0}k_{bl}}{k_{bl} - k_{act,561}}\left(e^{-k_{act,561}t} - e^{k_{bl}t}\right) \qquad (13)$$

Eq. 12 describes how the number of red mEos3.2 molecules in a cell (R_n) changes during
the period of 561-nm illumination. Multiplying both sides of Eq. 12 by ε_f gives equation (14) that
describes how the fluorescence signal per cell changes over time during this period:

$$R_s(t) = \frac{1}{k_{bl} - k_{act,561}}\left[\left(S_n \varepsilon_f - R_{n,0}\varepsilon_f\right)k_{act,561}e^{-k_{act,561}t} - \left(S_n \varepsilon_f k_{act,561} - R_{n,0}\varepsilon_f k_{bl}\right)e^{-k_{bl}t}\right] \qquad (14)$$

We estimated $S_n \times \varepsilon_f$, $R_{n,0} \times \varepsilon_f$, $k_{act,561}$, and k_{bl} using Levenberg-Marquardt nonlinear least
squares regression to fit Eq. 14 to the time courses of the fluorescence signal per cell during the
561-nm illumination period. We calculated the mean and standard deviation of the activation rate
constant ($k_{act,561}$) for converting mEos3.2 molecules from I- to R-state by 561-nm irradiation
from averaging the 7 periods of 561-nm irradiation for each condition.

Single-molecule Localization

We acquired data on single R-state mEos3.2 molecules with our custom-built SMLM using a

405-nm laser intensity of 1 W/cm^2, a 561-nm laser intensity of 1 kW/cm^2, and a frame rate of 50

fps. We localized single molecules with the Python Microscopy Environment (PYME) package

(34), using a threshold of 0.6 for event detection computed from the estimated pixel signal-to-

noise ratio (SNR).We corrected pixel-dependent noise with maps generated from dark camera

frames. We measured the number of photons from single R-state mEos3.2 molecules in each 20-

ms frame between frames 5,000 and 10,000 only, since the R-state molecules before frame 5,000

were too dense for localization and most were bleached after frame 10,000.

Cell Permeability Experiments

Wildtype *S. pombe* cells were fixed for 15 min or 30 min as above. Fixed and live cells were

mounted on coverslips coated with 0.5 mg/mL peanut-lectin (Sigma-Aldrich; L0881-5MG) for

30 min. The cells were then incubated for 30 min in EMM5S medium containing 20 µM

fluorescein (Sigma-Aldrich 46960-25G-F) or fluorescein-dextran 3,000 MW (ThermoFisher

D3305). The live cells mixed with the dyes were imaged at room temperature (~23° C)

immediately to minimize endocytosis of the dyes. Wildtype cells in EMM5S medium alone were

prepared and imaged as negative controls for autofluorescence subtraction. Brightfield and

confocal fluorescence images of the mid-sections of cells were acquired with LSM880. The

samples were excited at 488 nm (~22 µW at the sample) and the emitted fluorescence was

collected using an emission band path filter with a 519-601 nm wavelength range. We analyzed

the images in Fiji (33) by manually selecting ROIs of 5x5-pixel squares in the cells (~30-40 cells

for each condition; 1 ROI per cell) and the extracellular environment (6~12 ROIs for each

condition) (Fig. S3A). We measured the MSPP of all ROIs including the negative controls (for

autofluorescence subtraction), quantified the cell permeability in each condition by calculating

($\text{MSPP}_{\text{cytoplasmic}} - \text{MSPP}_{\text{negative control}}$)/ $\text{MSPP}_{\text{extracellular}}$, and reported the mean and standard

deviation.

RESULTS

Quantitative assessment of mEos3.2 photophysics in yeast cells by fitting a 3-state model to

fluorescence microscopy data

We combined quantitative fluorescence microscopy with mathematical modeling to estimate the

time-integrated signal per cell detected in the red channel ($\overline{R_s}$), and the rate constants for

photoconversion and photobleaching (k_{act} and k_{bl}) of mEos3.2 in the cytoplasm of fission yeast

cells (Fig. 1). Our model assumes that illumination at 405 nm photoconverts mEos3.2 molecules

irreversibly from their green (G) to their red (R) state with an activation rate constant of k_{act} and

that the 561-nm light excites the red-form mEos3.2 with a peak emission at ~580 nm.

Illumination at either 405 nm or 561 nm converts R-state mEos3.2 molecules to the bleached (B)

state by an irreversible first-order reaction with a bleaching rate constant of k_{bl} (Fig. 1C). Our 3-

state model did not consider mEos3.2 photoswitching or "blinking" in its G- or R-state, where

the protein enters a transient off state and can be converted back to the fluorescent state.

Fission yeast cells expressing mEos3.2 from the actin promoter in the *leu1* locus ensured

a relatively high and homogenous cytoplasmic mEos3.2 expression level (Fig. 1A). We used

point-scanning confocal microscopy to illuminate the cells at both 405 nm and 561 nm and

collect time-lapse images in the red wavelength range of 566-719 nm (Fig. 1A). The time course

of the fluorescence signal per cell first rose as the large pool of molecules in the G-state was

photoconverted to the R-state, from which we detected red photons as signal, and then declined

as the pool of molecules in the G-state was depleted and photobleaching depleted the R-state

pool (Fig. 1B).

The equation (Eq. 7) of our 3-state model (Fig. 1C) fit the time courses of fluorescence

signal per cell very closely (Fig. 1B). The best fits yielded estimates of the product of total

number of molecules per cell and detected signal per R-state molecule per frame (M_n x ε_f), and

the rate constants for photoconversion (k_{act}) and photobleaching (k_{bl}) (Table S1). We then used

Eq. 8 to calculate the time-integrated signal per cell ($\overline{R_s}$) using fitted parameters M_n x ε_f and k_{bl}

(Table S1). We used this approach to measure how sample preparation and imaging conditions

influence these photophysical properties.

Effects of formaldehyde fixation and imaging buffer on photophysical properties of

mEos3.2 in fission yeast cells

We investigated how fixation affects the photophysical properties of mEos3.2, so we could

compare experiments on live and fixed yeast cells (Fig. 2). Yeast cells fixed with formaldehyde

in EMM5S-synthetic growth medium emitted fewer red photons than live cells imaged under the

same conditions (Fig. 2C). Fitting Eq. 7 of the 3-state model (Fig. 1C) to the time courses of

fluorescence signal per yeast cell showed that mEos3.2 in fixed cells had lower M_n x ε_f (Fig. 2D)

and $\overline{R_s}$ (Fig. 2G) and higher k_{act} (Fig. 2E) and k_{bl} (Fig. 2F) than live cells. Yeast cells fixed for 30

min had even lower fluorescence signals (Fig. 2D, G) and higher rate constants than cells fixed

for 15 min (Fig. 2E, F).

We used fluorescein to test our hypothesis that formaldehyde fixation affects mEos3.2

photophysics by permeabilizing the yeast cell membrane for small molecules (Fig. S3). Live

yeast cells excluded fluorescein (332 g/mol), but fixation with formaldehyde (without detergents

or organic solvents) partially permeabilized the cell membrane, allowing the entry of fluorescein

(Fig. S3D). Moreover, more fluorescein entered the cells fixed for 30 min than the cells fixed for

15 min (Fig. S3D). Thus, ions and small molecules in the imaging buffer, such as thiol DTT (154

g/mol), equilibrated with the interior of the fixed yeast cells and affected the photophysical

properties of mEos3.2.

Knowing that fixed *S. pombe* cells are permeable for small molecules, we tested how the

composition of the imaging buffer influenced mEos3.2 photophysics (Fig. 3). We hypothesized

that the photophysical changes of mEos3.2 in the fixed cells were due to the low pH (~5.5) and

oxidizing environment of EMM5S relative to the live-cell cytoplasmic environment. We found

that $\overline{R_s}$ was higher (Fig. 3B), and k_{act} (Fig. 3D) and k_{bl} (Fig. 3E) were lower in imaging buffers

with higher pH. Adding the reducing agent DTT to the imaging buffer further increased $\overline{R_s}$ (Fig.

3B) and decreased k_{bl} in the pH range we tested (Fig. 3E). A concentration of 1 mM DTT was

more effective than 10 mM DTT at increasing $\overline{R_s}$ (Fig. 3B). Additionally, the value of $M_n \times \varepsilon_f$

from fixed cells in 50 mM Tris-HCl (pH 8.5) with 1 mM DTT (9187 A.U., 95% CI: 9024 – 9351,

Table S2) was similar to that from live cells (9489 A.U., 95% CI: 9291 - 9687, Table S1). Values

of $\overline{R_s}$ were also similar in live cells and fixed cells at pH 8.5 with 1 mM DTT (Table S1, S2).

Thus, mEos3.2 molecules survived fixation and the total number of fluorescence-competent

molecules per cell (M_n) did not change, but the extracellular imaging buffer influenced the

intracellular mEos3.2 signal per frame (ε_f) and other photophysical properties as

photoconversion and photobleaching rates. In the following, we therefore used the imaging

buffer at pH 8.5 with 1 mM DTT (called 'Tris8.5-DTT buffer') for many fixed cell imaging

conditions.

Effects of the 405-nm and 561-nm laser intensities on photophysical properties of mEos3.2

in fission yeast cells

We used laser-scanning confocal microscopy and widefield fluorescence microscopy to test the

effects of a wide range of laser intensities on mEos3.2 photophysics in fixed *S. pombe* cells in

Tris8.5-DTT buffer (Fig. 4). The products of total mEos3.2 molecules per cell and signal per R-

state molecule per frame ($M_n \times \varepsilon_f$) were similar in live cells and fixed cells in Tris8.5-DTT buffer

(Fig. 4A-D). This was true for low-power laser-scanning confocal microscopy conditions as well

as widefield SMLM imaging conditions with a 405-nm laser intensity of 0.5-2 W/cm^2 and a 561-

nm laser intensity of 1 kW/cm^2 (Fig. 4C). However, at widefield 561-nm laser intensities of 10

and 100 W/cm^2, the time-integrated signal per cell ($\overline{R_s}$) differed in live cells and fixed cells in

Tris8.5-DTT buffer (Fig. 4P). Fixation and the imaging buffer had different effects on $M_n \times \varepsilon_f$

(Fig. 4D), k_{act} (Fig. 4H), k_{bl} (Fig. 4L) and $\overline{R_s}$ (Fig. 4P) of mEos3.2 depending on the 561-nm

laser intensities.

To compare imaging conditions quantitatively, we explored the effects of laser intensities

on mEos3.2 photophysics by point-scanning illumination (Fig. 4, left 2 columns, Table S3). $\overline{R_s}$

decreased (Fig. 4M), and both rate constants (Fig. 4E, I) increased with higher 405-nm laser

intensity. $\overline{R_s}$ increased with higher 561-nm laser intensity (Fig. 4N), but the 561-nm laser

intensity had only modest effects on both rate constants (Fig. 4F, J) in the range we tested.

We used widefield illumination to explore the effects of a wider range of 561-nm laser

intensities on mEos3.2 photophysics, including the high 561-nm laser intensity of ~1 kW/cm^2

often used in SMLM (Fig. 4, right 2 columns, Table S4). Values of k_{bl} increased with higher

405-nm laser intensity (Fig. 4K), as observed with point-scanning illumination (Fig. 4I), but the

405-nm laser intensity had remarkably little impact on k_{act} (Fig. 4G) under SMLM conditions,

which was likely caused by the high 561-nm laser intensity of 1 kW/cm^2 driving some

photoconversion. The time-integrated signal per cell $\overline{R_s}$ increased with higher 561-nm laser

intensity and then plateaued and dropped (Fig. 4P). Both k_{act} and k_{bl} increased dramatically with

561-nm laser intensities (Fig. 4H, L) above the intensities used for point-scanning illumination

(Fig. 4F, J). The discussion section considers these differences between the confocal and

widefield conditions.

Photon counts from single red-state mEos3.2 molecules in live and fixed *S. pombe* cells

To assess the effects of fixation and the imaging buffer on mEos3.2 photophysics under SMLM

conditions, we compared the photon counts of single R-state mEos3.2 molecules per frame in

live and fixed yeast cells (Fig. 5) under SMLM imaging conditions with a frame rate of 50 fps

and a 561-nm laser intensity of 1 kW/cm^2. We localized the single R-state molecules in each

frame and measured their photon counts when the density of the R-state molecules was sparse

enough for localization.

The mean photon counts per frame from single R-state mEos3.2 molecules in live cells

and fixed cells imaged in the Tris8.5-DTT buffer were identical (P = 1.000) and higher than the

counts in fixed cells imaged in the EMM5S medium at pH 5.5 (P = 0.007, Fig.5, inset).

Therefore, the higher brightness per frame of R-state mEos3.2 molecule in the fixed cells in

Tris8.5-DTT buffer can improve localization precision (35). Besides, the oxidizing environment

in fixed cells in EMM5S medium may prolong the total on-time (the total time a mEos3.2

molecule is in the red fluorescent state until bleached) of single R-state mEos3.2 molecules,

suggested by the time-integrated signal per cell $\overline{R_s}$ (the product of brightness per frame and total

on-time of single R-state mEos3.2, and the total number of molecules per cell) being higher in

the fixed cells in EMM5S medium than in the other 2 conditions (Fig. 4O).

Intermediate state of mEos3.2 that converts to the red fluorescent state upon 561-nm

illumination

To investigate the effects of 405-nm and 561-nm illumination separately, we alternated periods

of multiple cycles of 405-nm and 561-nm illumination. Each period of 405-nm illumination

increased the fluorescence emission during 561-nm illumination; surprisingly, the signal rose

transiently above the initial value despite the 405-nm illumination being switched off (Fig. 6A).

After peaking during the fifth cycle of each 561 nm illumination period, the signal decreased due

to photobleaching (Fig. 6A, B). The signal during each 561-nm illumination period (except for

the first one without prior 405-nm illumination) followed similar time courses (Fig. 6B). These

transient increases in the fluorescence signal also occurred using alternating widefield

illumination conditions (Fig. S4).

We considered four hypotheses to explain these transient increases in the fluorescence

signal. First, illumination at 561 nm might photoconvert G-state mEos3.2 to the R-state directly,

but we observed no comparable activation with 561-nm irradiation alone (Fig. S5B, and first 40

cycles of 561-nm irradiation in Fig. 6A). A second hypothesis is that the first-order

photoconversion reaction is slow after absorption of 405-nm photons, delaying accumulation of

R-state molecules. We ruled out this mechanism by adding a 2-min pause between each 405-nm

illumination period and the following 561-nm illumination period. The transient increase in the

fluorescence signal was still observed, ruling out this hypothesis as a dominant effect (Fig. S5A).

Similarly, we could rule out the third hypothesis that the observed increase in the fluorescence

signal over the course of 561-nm illumination was related to protein maturation or similar live-

cell phenomena by observing the same effect in fixed cells (Fig. S5C).

The fourth hypothesis is that 405-nm illumination leaves a subpopulation of mEos3.2

molecules in an intermediate (I) state that requires 561-nm illumination to convert to the

fluorescent R-state. To test this hypothesis, we added a fourth I-state to the model (Fig. 6D).

Whether this I-state is populated from the R-state or G-state mEos3.2 molecules does not affect

Eqs. 9-14, which only consider the period of 561-nm illumination when conversion into the I-

state is negligible. Eq. 14 of our 4-state model fits closely the time courses of mEos3.2

fluorescence signal during each 561-nm illumination period (Fig. 6B). The best fits gave an

average activation rate constant for conversion from the I-state to R-state by 561-nm irradiation

$k_{act,561} = 0.34$ s^{-1} (SD: 0.02 s^{-1}) (Fig. 6B), which is 4-fold higher than the rate constant for

photoconversion from the G- to R-state, $k_{act} = 0.050$ s^{-1} (95% CI: 0.048- 0.051 s^{-1}), as measured

with simultaneous 405-nm and 561-nm illumination (Fig. S5D).

We compared the number of I-state molecules changing over time with the respective G-

and R-state populations (Fig. 6C). The normalized time course of the I-state molecules was

similar to R-state molecules and distinctly different from the G-state molecules, strongly

suggesting that the I-state is populated from the R-state molecules. Overall, these experiments

revealed that a subpopulation of the R-state molecules is converted to the I-state but can be

converted back to the R-state by 561-nm illumination (Fig. 6D). We did not detect any

spontaneous conversion from I- to R-state over 2 minutes (Fig. S5A). However, for experiments

with simultaneous illumination at both 405 nm and 561 nm, Eq. 7 of our simplified 3-state model

(without considering the I-state) fully accounted for the time courses of fluorescence signal per

cell under all the conditions (Fig. 1B, 2C, 3A,S8, S9).

DISCUSSION

Characterizing the photophysics of mEos3.2 by fitting our 3-state model to quantitative

time-lapse fluorescence microscopy data

We show how to measure the product of total number of molecules per cell and the signal per R-

state molecule per frame (M_n x ε_f) and rate constants for photoconversion (k_{act}) and

photobleaching (k_{bl}) in cells by fitting a simple 3-state model to the time course of measured

fluorescence signals in the red channel (Fig. 1). Eq. 7 of our 3-state model fit the time courses of

the fluorescence signal per cell very closely (Fig. 2C, 3A) and revealed how different conditions

changed the three parameters separately. Furthermore, simulations of the 3-state model showed

how the values of the three parameters influenced the time courses of the number of R-state

molecules per cell (Fig. S2B – D). We also used Eq. 8 to calculate the time-integrated signal per

cell ($\overline{R_s}$) using M_n x ε_f and k_{bl}.

Our approach has several advantages over the alternative approach of characterizing

single fluorescent molecules *in vitro* (18, 37, 38). (1) It avoids potential artifacts caused by using

arbitrary photon number or localization uncertainty thresholds in single-molecule localization

algorithms to separate molecules from noise. (2) It is easy to implement with conventional

microscopes and whole cells. (3) Large numbers of cells can be imaged in just hours to test

different sample preparation and imaging conditions, including a wide range of laser intensities.

(4) It extracts photoconversion or photoactivation rate constants from PAFPs or PCFPs in cells

more easily than single-molecule methods, as photobleaching is hard to account for in single-

molecule data (30). These rate constants are useful for optimizing SMLM imaging conditions

and simulating SMLM raw data.

On the other hand, our approach cannot measure the single-molecule blinking kinetics of

mEos variants in either their green (39, 40) or red states (41, 42). Blinking in the green state of

mEos variants could affect photoconversion to the red state (39, 40). The interplay between

blinking and photoconversion has also been described in other photoconvertible fluorescent

proteins, such as SAASoti (43) and LEA (44). Thus, combining our approach with single-

molecule measurements will offer a more complete and quantitative understanding of the

photophysics of PAFPs or PCFPs.

Sample preparation for imaging mEos3.2 in fixed fission yeast cells

Preserving the fluorescence signal and structures of interest is crucial when fixing cells with

fluorescent protein labels. We show that formaldehyde fixation does not destroy mEos3.2

molecules but permeabilizes yeast cells for small molecules, making the photophysical properties

of mEos3.2 sensitive to the composition of the imaging buffer. The low pH of 5.5 and lack of

oxygen scavenging system in the EMM5S synthetic medium affected the photophysical

properties of mEos3.2 in fixed cells as expected from previous works showing that EosFP

photoconverts faster (4) and mEos3.2 emits fewer red photons at acidic pH (6). Photooxidation

can increase the photobleaching rate (45). Oxygen in the solution can also affect the fluorescence

signal by promoting intersystem crossing (46) and convert the excited molecules to the non-

fluorescent triplet state.

Our experiments also show that fixation conditions should be tested and optimized for

each fluorescent protein. For example, mEos3.2 had even lower $\overline{R_s}$ and higher k_{act} and k_{bl} in cells

fixed for 30 min rather than 15 min under the tested imaging condition (Fig. 2E-G). On the other

hand, GFP was far less sensitive to fixation, as formaldehyde treatment had little effect on its

fluorescence signal and photobleaching rate (Fig. S6).

The photophysics of mEos3.2 was similar in live cells and fixed cells in the Tris8.5-DTT

buffer (Fig. 3). Imaging buffers with a pH equal to or slightly higher than the cytoplasmic pH of

fission yeast at ~7.3 (47) increased $\overline{R_s}$ and reduced k_{act} and k_{bl} (Fig. 3B, D, E). Adding a reducing

agent DTT to the imaging buffer further increased $\overline{R_s}$ (Fig. 3B). The Tris8.5-DTT buffer not only

maintained the photon counts of single R-state mEos3.2 molecules in each frame (Fig. 5) but

also reduced the cellular autofluorescence compared with fixed cells in EMM5S medium (Fig.

S8E, S8H). This is crucial for SMLM imaging, where the signal-to-noise ratio is important for

obtaining high localization precision and consequently resolution (37).

Comparison of point-scanning and widefield illumination for mEos3.2 photophysics

We compared the photophysical properties of mEos3.2 under point-scanning and widefield

illumination. Under point-scanning illumination, each area of the sample was illuminated for a

very short time at high peak intensity (e.g. ~80 kW/cm^2, Fig. 4), while the other pixels were kept

in the dark. Thus, the average intensity of the laser power over the entire field of view was ~10^4

times lower than the peak intensity (e.g. ~0.5 W/cm^2, Fig. 4).

Despite huge differences in the instantaneous peak intensities in point-scanning and

widefield microscopy, the rate constants for photoconversion and photobleaching in live cells

were similar at comparable average intensities (Table S3, S4). For example, with average

intensities of ~ 0.5 to 1.1 W/cm^2 for both lasers, confocal and widefield imaging of live cells

gave similar values for k_{act} (~1.5×10^{-2} s^{-1} vs. ~5×10^{-2} s^{-1}) and k_{bl} (~2×10^{-3} s^{-1} vs. ~1×10^{-3} s^{-1}).

Moreover, the photophysical parameters of mEos3.2 trended similarly with illumination

intensities by both point-scanning confocal and widefield microscopy. For example, k_{bl} increased

with higher 405-nm intensity (Fig. 4I, K) and M_n x ε_f increased with higher 561-nm laser

intensity (Fig. 4B, D). However, these trends diverged at higher 561-nm laser intensities of 100

and 1000 W/cm^2 above the range of the confocal microscope. For example, k_{act} increased

dramatically with 561-nm laser intensity (Fig. 4H), but only increased slightly with 405-nm laser

intensity (Fig. 4G, 561-nm laser intensity = 1 kW/cm^2). We conclude that the 561-nm laser

contributed strongly to photoconversion at high intensities.

The effects of the fixation and imaging buffer on mEos3.2 photophysics is also consistent

under different illumination methods. For example, under comparable, low average 405-nm and

561-nm laser intensities (~1 W/cm^2), the time-integrated signal per cell $\overline{R_s}$ (Fig. 4N, P) was

lower and photoconversion (Fig. 4F, H) and photobleaching rates constants (Fig. 4J, L) were

higher in fixed cells in EMM5S than in live cells and fixed cells in the Tris8.5-DTT buffer.

However, fixation and imaging buffer affect mEos3.2 photophysics differently at higher 561-nm

laser intensities of 100 and 1000 W/cm^2, above the range of confocal microscope. For example,

$\overline{R_s}$ was lower in fixed cells in EMM5S (pH 5.5) than in live cells under low 561-nm laser

intensity in the range of 1-10 W/cm^2, while for a high 561-nm laser intensity of 1 kW/cm^2 the

trend was the opposite (Fig. 4P). The $\overline{R_s}$ in the fixed cells was only higher in the Tris8.5-DTT

buffer than in EMM5S under low 561-nm laser intensities of 1-10 W/cm^2 (Fig. 4P). Moreover,

under low 561-nm laser intensities both k_{act} (Fig. 4E, H) and k_{bl} (Fig. 4I, L) in fixed cells in

EMM5S were higher than in live cells and fixed cells in Tris8.5-DTT buffer, while under higher

561-nm laser intensities some measurements of the rates were lower than in the Tris8.5-DTT

buffer (Fig. 4H, L). Oxygen in the environment could promote intersystem crossing and convert

the excited fluorophores to the non-fluorescent triplet state, where the molecules could return to

the fluorescent state through laser excitation. The laser intensity could potentially affect the

probability that the triple state molecules return to the fluorescent state. Therefore, redox

environment in the imaging buffer for fixed cells may affect mEos3.2 photophysics differently

depending on the 561-nm laser intensity.

Comparing laser intensities for imaging mEos3.2 in live and fixed fission yeast cells

Our quantitative measurements provide information for selecting laser intensities to image

proteins tagged with mEos3.2. Maximizing the red fluorescence signal of mEos3.2 while

maintaining a relatively low level of autofluorescence background is the key to optimize imaging

quality. Higher signal-noise-ratios can increase localization precision and thus the resolution in

SMLM (35). For SMLM, one controls the rates of photoconversion and photobleaching to

achieve a density of active fluorophores sparse enough for localization but also dense enough to

image quickly. High laser intensities are usually used for fast SMLM imaging in fixed samples

(37).

Higher 405-nm illumination intensity had four effects relevant for SMLM image quality:

(1) it decreased $\overline{R_s}$ under a low average 561-nm laser intensity of ~ 0.3 W/cm^2 (Fig. 4M), but the

effect was not obvious under high 561-nm laser intensity of 1 kW/cm^2 (Fig. 4O); (2) it increased

background autofluorescence (Fig. S8B, E, H), especially in live cells (Fig. S8B); (3) it increased

k_{act} (Fig. 4E); and (4) it increased k_{bl} (Fig. 4I, K). One may ramp up the 405-nm laser intensity

while imaging a FOV to increase k_{act} and compensate for the loss of bleached molecules.

However, high 405-nm laser intensities can potentially decrease SMLM imaging quality in two

ways: increased autofluorescence compromises accurate localization; and rapid photobleaching

decreases $\overline{R_s}$, likely decreasing the total number of localizations.

The time-integrated signal $\overline{R_s}$ increased over the low range of 561-nm laser intensities

(Fig. 4N) and peaked in the middle of the high range of 561-nm laser intensities (Fig. 4P), so

very high 561-nm laser intensities could compromise image quality. Moreover, both the

photoconversion (Fig. 4H) and photobleaching rates (Fig. 4L) increased at high 561-nm laser

intensities. The photoconversion rate likely increases, because 561-nm laser is the predominant

source of photoconversion at high intensities (Fig. 6D). Our results differ from Thedie *et al.* (39),

who reported that high 561-nm laser intensities (1.2 – 4.8 kW/cm^2) slow the photoconversion of

single mEos2 molecules embedded in PVA under weak 405-nm illumination (0.03 W/cm^2).

Their interpretation was that 561-nm illumination pushes green-state mEos2 molecules into a

transient off state that cannot be photoconverted. Both the sample conditions and illumination

intensities differ in the 2 experiments. Their mEos2 molecules were immobilized in PVA with

restricted access to oxygen rather than being in a physiological environment in cells.

Additionally, we used higher 405-nm illumination (1 W/cm^2) and lower 561-nm laser intensity

(0.001 – 1 kW/cm^2) where green molecules in the transient off state may convert back to the

fluorescent state, thus promoting photoconversion to the red-state.

Based on our data, we recommend the following for SMLM imaging of mEos3.2 in live

and fixed fission yeast cells. (1) Use high but not maximal 561-nm laser intensities to avoid

decreasing $\overline{R_s}$ (Fig. 4P) and degrading the SMLM image quality. (2) Use low 405-nm laser

intensities to achieve sparse activation for temporal separation of molecules and to compensate

for the high photoconversion rate from strong 561-nm laser illumination.

Application of our findings to quantitative SMLM with mEos3.2 in live and fixed yeast cells

Several variants of EosFP fluorescent proteins are used for SMLM of live and fixed cells (9, 11,

12). Both sample preparation and imaging conditions can affect quantitative measurement. The

time-integrated signal ($\overline{R_s}$) contains information of both the brightness per single fluorescent

molecule and the total number of molecules able to fluoresce in the structure of interest. The

single-molecule brightness also affects the number of mEos3.2 localizations, as only molecules

emitting more than a threshold number of photons in each frame are counted (34). Under SMLM

imaging conditions (Fig. 4O), $\overline{R_s}$ measurements differed in live cells and fixed cells in the

Tris8.5-DTT buffer, so they cannot be used for direct comparisons. However, the photon counts

from single mEos3.2 emission bursts in each SMLM frame were similar in live cells and fixed

cells (Fig. 5), so the same analysis thresholds can be used for single-molecule localizations.

Combining the time-integrated signal $\overline{R_s}$ (Fig. 4O) and photon counts (Fig. 5, inset) suggests that

fixation and the imaging buffer affect the total on-time of single R-state mEos3.2, likely due to

the oxidation-reduction environment. Similarly, adding reducing thiol MEA and removing

oxygen from the imaging buffer increase the number of blinking cycles of R-state mEos2 in

fixed mammalian cells (41) and of purified mEos3.2 (48).

Illumination at 561-nm converts intermediate-state mEos3.2 to the red fluorescent state

Our alternating illumination experiments (Fig. 6) revealed that some R-state molecules enter the

I-state but can be reconverted back to the R-state by 561-nm illumination. De Zitter *et al.* has

reported a similar observation, that mEos4b – an mEos3.2 derivative – recovered from a long-

lived dark state to the red state in response to 561-nm illumination (42). The long-lived dark state

recovery rate increased with 561-nm laser intensity and could be recovered to the red state by

405-nm irradiation. Our I-state converts to R-state upon 561-nm illumination, even at as low as 1

W/cm^2 (Fig. S4A). However, our I-state molecules did not seem to respond to 405-nm

illumination. The period of 561-nm illumination would not increase the fluorescence signal

transiently (Fig. 6A), if 405-nm irradiation could convert the I-state to R-state. Therefore, the I-

state we observed for mEos3.2 is unlikely to be the long-lived dark state as described for

mEos4b (42). Further structural studies are needed to characterize the nature of this intermediate

state.

CONCLUSION

We measured the time-integrated signal per cell and the rate constants for photoconverting and

photobleaching of mEos3.2 in fission yeast cells by fitting a 3-state model to time-lapse

quantitative fluorescence microscopy data. Formaldehyde fixation partially permeabilized the

yeast cell membrane for small molecules, allowing the reducing agent and ions in the imaging

buffer to enter fixed yeast cells and affect mEos3.2 photophysics. Using an imaging buffer at pH

8.5 with 1 mM DTT resulted in a time-integrated signal comparable to live cells under certain

imaging condition, suggesting that mEos3.2 molecules survived the light chemical fixation. We

investigated the effects of fixation and imaging buffer on mEos3.2 photophysics in yeast cells

over a wide range of laser intensities by point-scanning and widefield microscopy. We

discovered an intermediate state of mEos3.2, populated by red-state molecules, that can be

convert to the red fluorescent state by 561-nm illumination. Our results provide information for

sample preparation and laser intensities for imaging and counting mEos3.2-tagged molecules in

fission yeast cells. Our imaging assay and 3-state model can also be applied to study the

photophysical properties of other photoactivatable or photoconvertible fluorescent proteins.

SUPPORTING MATERIAL

Supporting Material can be found online at ***.

AUTHOR CONTRIBUTIONS

HS, JB and TDP designed research; HS carried out the experiments; KH built the single-

molecule localization microscope and assisted in imaging and data analysis; all authors analyzed

the data and wrote the paper.

ACKNOWLEDGMENTS

Research reported in this publication was supported by National Institute of General Medical

Sciences of the National Institutes of Health under award number R01GM026132 and R01

GM118486, and the Wellcome Trust (203285/B/16/Z). MS and KH were supported by NIH

Grant No. T32EB019941. The content is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institutes of Health. The authors thank

Bennett Rollins and Zach Marin for help with single-molecule localization imaging and data

analysis, Yongdeng Zhang for helpful discussion, and Phylicia Kidd for help with sample

preparation. We also acknowledge support from the Program in Physics, Engineering and

Biology. JB discloses a significant financial interest in Bruker Corp. and Hamamatsu Photonics.

References:

1.   Rust, M. J., M. Bates, and X. Zhuang. 2006. Sub-diffraction-limit imaging by stochastic
optical reconstruction microscopy (STORM). Nat Methods 3:793-795.
2.   Betzig, E., G. H. Patterson, R. Sougrat, O. W. Lindwasser, S. Olenych, J. S. Bonifacino,
M. W. Davidson, J. Lippincott-Schwartz, and H. F. Hess. 2006. Imaging intracellular
fluorescent proteins at nanometer resolution. Science 313:1642-1645.
3.   Hess, S. T., T. P. K. Girirajan, and M. D. Mason. 2006. Ultra-high resolution imaging by
fluorescence photoactivation localization microscopy. Biophysical Journal 91:4258-4272.
4.   Wiedenmann, J., S. Ivanchenko, F. Oswald, F. Schmitt, C. Rocker, A. Salih, K. D.
Spindler, and G. U. Nienhaus. 2004. EosFP, a fluorescent marker protein with UV-
inducible green-to-red fluorescence conversion. Proc Natl Acad Sci U S A 101:15905-
15910.
5.   McKinney, S. A., C. S. Murphy, K. L. Hazelwood, M. W. Davidson, and L. L. Looger.
2009. A bright and photostable photoconvertible fluorescent protein. Nat Methods 6:131-
133.
6.   Zhang, M., H. Chang, Y. Zhang, J. Yu, L. Wu, W. Ji, J. Chen, B. Liu, J. Lu, Y. Liu, J.
Zhang, P. Xu, and T. Xu. 2012. Rational design of true monomeric and bright
photoactivatable fluorescent proteins. Nat Methods 9:727-729.
7.   Paez-Segala, M. G., M. G. Sun, G. Shtengel, S. Viswanathan, M. A. Baird, J. J. Macklin,
R. Patel, J. R. Allen, E. S. Howe, G. Piszczek, H. F. Hess, M. W. Davidson, Y. Wang,
and L. L. Looger. 2015. Fixation-resistant photoactivatable fluorescent proteins for
CLEM. Nat Methods 12:215-218, 214 p following 218.
8.   Laplante, C., F. Huang, I. R. Tebbs, J. Bewersdorf, and T. D. Pollard. 2016. Molecular
organization of cytokinesis nodes and contractile rings by super-resolution fluorescence
microscopy of live fission yeast. Proc Natl Acad Sci U S A.
9.   Zhang, Y., M. Lara-Tejero, J. Bewersdorf, and J. E. Galan. 2017. Visualization and
characterization of individual type III protein secretion machines in live bacteria. Proc
Natl Acad Sci U S A 114:6098-6103.

10.   Akamatsu, M., Y. Lin, J. Bewersdorf, and T. D. Pollard. 2017. Analysis of interphase
670   node proteins in fission yeast by quantitative and superresolution fluorescence
671   microscopy. Mol Biol Cell 28:3203-3214.

672   11.   Puchner, E. M., J. M. Walter, R. Kasper, B. Huang, and W. A. Lim. 2013. Counting
673   molecules in single organelles with superresolution microscopy allows tracking of the
674   endosome maturation trajectory. Proc Natl Acad Sci U S A 110:16015-16020.

675   12.   Fricke, F., J. Beaudouin, R. Eils, and M. Heilemann. 2015. One, two or three? Probing
676   the stoichiometry of membrane proteins by single-molecule localization microscopy. Sci
677   Rep 5:14072.

678   13.   Shroff, H., C. G. Galbraith, J. A. Galbraith, H. White, J. Gillette, S. Olenych, M. W.
679   Davidson, and E. Betzig. 2007. Dual-color superresolution imaging of genetically
680   expressed probes within individual adhesion complexes. P Natl Acad Sci USA
681   104:20308-20313.

682   14.   Nienhaus, K., G. U. Nienhaus, J. Wiedenmann, and H. Nar. 2005. Structural basis for
683   photo-induced protein cleavage and green-to-red conversion of fluorescent protein EosFP.
684   Proc Natl Acad Sci U S A 102:9156-9159.

685   15.   Turkowyd, B., A. Balinovic, D. Virant, H. G. G. Carnero, F. Caldana, M. Endesfelder, D.
686   Bourgeois, and U. Endesfelder. 2017. A General Mechanism of Photoconversion of
687   Green-to-Red Fluorescent Proteins Based on Blue and Infrared Light Reduces
688   Phototoxicity in Live-Cell Single-Molecule Imaging. Angew Chem Int Edit 56:11634-
689   11639.

690   16.   Coffman, V. C., and J. Q. Wu. 2012. Counting protein molecules using quantitative
691   fluorescence microscopy. Trends Biochem Sci 37:499-506.

692   17.   Baddeley, D., and J. Bewersdorf. 2017. Biological Insight from Super-Resolution
693   Microscopy: What We Can Learn from Localization-Based Images. Annu Rev Biochem.

694   18.   Lee, S. H., J. Y. Shin, A. Lee, and C. Bustamante. 2012. Counting single photoactivatable
695   fluorescent molecules by photoactivated localization microscopy (PALM). Proc Natl
696   Acad Sci U S A 109:17436-17441.

697   19.   Rollins, G. C., J. Y. Shin, C. Bustamante, and S. Presse. 2015. Stochastic approach to the
698   molecular counting problem in superresolution microscopy. Proc Natl Acad Sci U S A
699   112:E110-118.

700   20.   Annibale, P., S. Vanni, M. Scarselli, U. Rothlisberger, and A. Radenovic. 2011.
701   Quantitative photo activated localization microscopy: unraveling the effects of
702   photoblinking. PLoS One 6:e22678.

703   21.   Hummer, G., F. Fricke, and M. Heilemann. 2016. Model-independent counting of
704   molecules in single-molecule localization microscopy. Mol Biol Cell 27:3637-3644.

705   22.   Ehmann, N., S. van de Linde, A. Alon, D. Ljaschenko, X. Z. Keung, T. Holm, A. Rings,
706   A. DiAntonio, S. Hallermann, U. Ashery, M. Heckmann, M. Sauer, and R. J. Kittel. 2014.
707   Quantitative super-resolution imaging of Bruchpilot distinguishes active zone states. Nat
708   Commun 5:4650.

709   23.   Karathanasis, C., J. Medler, F. Fricke, S. Smith, S. Malkusch, D. Widera, S. Fulda, H.
710   Wajant, S. J. L. van Wijk, I. Dikic, and M. Heilemann. 2020. Single-molecule imaging
711   reveals the oligomeric state of functional TNFalpha-induced plasma membrane TNFR1
712   clusters in cells. Sci Signal 13.

713   24.   Virant, D., B. Traenkle, J. Maier, P. D. Kaiser, M. Bodenhofer, C. Schmees, I. Vojnovic,
714   B. Pisak-Lukats, U. Endesfelder, and U. Rothbauer. 2018. A peptide tag-specific
715   nanobody enables high-quality labeling for dSTORM imaging. Nat Commun 9:930.
716   25.   Gautier, A., A. Juillerat, C. Heinis, I. R. Correa, Jr., M. Kindermann, F. Beaufils, and K.
717   Johnsson. 2008. An engineered protein tag for multiprotein labeling in living cells. Chem
718   Biol 15:128-136.
719   26.   Los, G., A. Darzins, N. Karassina, C. Zimprich, R. Learish, M. McDougall, L. Encell, R.
720   Friedman-Ohana, M. Wood, and G. Vidugiris. 2005. HaloTag™ Interchangeable labeling
721   technology for cell imaging and protein capture. Cell Notes 11:2-6.
722   27.   Jungmann, R., M. S. Avendano, M. Dai, J. B. Woehrstein, S. S. Agasti, Z. Feiger, A.
723   Rodal, and P. Yin. 2016. Quantitative super-resolution imaging with qPAINT. Nat
724   Methods 13:439-442.
725   28.   Thevathasan, J. V., M. Kahnwald, K. Cieslinski, P. Hoess, S. K. Peneti, M. Reitberger, D.
726   Heid, K. C. Kasuba, S. J. Hoerner, Y. Li, Y. L. Wu, M. Mund, U. Matti, P. M. Pereira, R.
727   Henriques, B. Nijmeijer, M. Kueblbeck, V. J. Sabinina, J. Ellenberg, and J. Ries. 2019.
728   Nuclear pores as versatile reference standards for quantitative superresolution
729   microscopy. Nat Methods 16:1045-1053.
730   29.   Wang, S., J. R. Moffitt, G. T. Dempsey, X. S. Xie, and X. Zhuang. 2014.
731   Characterization and development of photoactivatable fluorescent proteins for single-
732   molecule-based superresolution imaging. Proc Natl Acad Sci U S A 111:8452-8457.
733   30.   Durisic, N., L. Laparra-Cuervo, A. Sandoval-Alvarez, J. S. Borbely, and M. Lakadamyali.
734   2014. Single-molecule evaluation of fluorescent protein photoactivation efficiency using
735   an in vivo nanotemplate. Nat Methods 11:156-162.
736   31.   Wu, J. Q., and T. D. Pollard. 2005. Counting cytokinesis proteins globally and locally in
737   fission yeast. Science 310:310-314.
738   32.   Ganguly, S., A. H. Clayton, and A. Chattopadhyay. 2011. Fixation alters fluorescence
739   lifetime and anisotropy of cells expressing EYFP-tagged serotonin1A receptor. Biochem
740   Biophys Res Commun 405:234-237.
741   33.   Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S.
742   Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J. Y. Tinevez, D. J. White, V. Hartenstein,
743   K. Eliceiri, P. Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for
744   biological-image analysis. Nat Methods 9:676-682.
745   34.   Lin, R., A. H. Clowsley, I. D. Jayasinghe, D. Baddeley, and C. Soeller. 2017.
746   Algorithmic corrections for localization microscopy with sCMOS cameras -
747   characterisation of a computationally efficient localization approach. Opt Express
748   25:11701-11716.
749   35.   Deschout, H., F. Cella Zanacchi, M. Mlodzianoski, A. Diaspro, J. Bewersdorf, S. T. Hess,
750   and K. Braeckmans. 2014. Precisely and accurately localizing single emitters in
751   fluorescence microscopy. Nat Methods 11:253-266.
752   36.   Nienhaus, G. U., K. Nienhaus, A. Holzle, S. Ivanchenko, F. Renzi, F. Oswald, M. Wolff,
753   F. Schmitt, C. Rocker, B. Vallone, W. Weidemann, R. Heilker, H. Nar, and J.
754   Wiedenmann. 2006. Photoconvertible fluorescent protein EosFP: biophysical properties
755   and cell biology applications. Photochem Photobiol 82:351-358.
756   37.   Lin, Y., J. J. Long, F. Huang, W. C. Duim, S. Kirschbaum, Y. Zhang, L. K. Schroeder, A.
757   A. Rebane, M. G. Velasco, A. Virrueta, D. W. Moonan, J. Jiao, S. Y. Hernandez, Y.

Zhang, and J. Bewersdorf. 2015. Quantifying and optimizing single-molecule switching nanoscopy at high speeds. PLoS One 10:e0128135.

38. Zanacchi, F. C., C. Manzo, A. S. Alvarez, N. D. Derr, M. F. Garcia-Parajo, and M. Lakadamyali. 2017. A DNA origami platform for quantifying protein copy number in super-resolution. Nat Methods 14:789-792.

39. Thedie, D., R. Berardozzi, V. Adam, and D. Bourgeois. 2017. Photoswitching of Green mEos2 by Intense 561 nm Light Perturbs Efficient Green-to-Red Photoconversion in Localization Microscopy. J Phys Chem Lett 8:4424-4430.

40. De Zitter, E., J. Ridard, D. Thedie, V. Adam, B. Levy, M. Byrdin, G. Gotthard, L. Van Meervelt, P. Dedecker, I. Demachy, and D. Bourgeois. 2020. Mechanistic Investigations of Green mEos4b Reveal a Dynamic Long-Lived Dark State. J Am Chem Soc 142:10978-10988.

41. Endesfelder, U., S. Malkusch, B. Flottmann, J. Mondry, P. Liguzinski, P. J. Verveer, and M. Heilemann. 2011. Chemically induced photoswitching of fluorescent probes--a general concept for super-resolution microscopy. Molecules 16:3106-3118.

42. De Zitter, E., D. Thedie, V. Monkemoller, S. Hugelier, J. Beaudouin, V. Adam, M. Byrdin, L. Van Meervelt, P. Dedecker, and D. Bourgeois. 2019. Mechanistic investigation of mEos4b reveals a strategy to reduce track interruptions in sptPALM. Nat Methods 16:707-710.

43. Solovyev, I. D., A. V. Gavshina, and A. P. Savitsky. 2019. Novel Phototransformable Fluorescent Protein SAASoti with Unique Photochemical Properties. Int J Mol Sci 20.

44. Krueger, T. D., L. Tang, L. Zhu, I. L. Breen, R. M. Wachter, and C. Fang. 2020. Dual Illumination Enhances Transformation of an Engineered Green-to-Red Photoconvertible Fluorescent Protein. Angew Chem Int Ed Engl 59:1644-1652.

45. Greenbaum, L., C. Rothmann, R. Lavie, and Z. Malik. 2000. Green fluorescent protein photobleaching: a model for protein damage by endogenous and exogenous singlet oxygen. Biological chemistry 381:1251-1258.

46. Mclean, A. J., D. J. Mcgarvey, T. G. Truscott, C. R. Lambert, and E. J. Land. 1990. Effect of Oxygen-Enhanced Intersystem Crossing on the Observed Efficiency of Formation of Singlet Oxygen. J Chem Soc Faraday T 86:3075-3080.

47. Zhang, Y., H. Shen, H. Liu, H. Feng, Y. Liu, X. Zhu, and X. Liu. 2017. Arp2/3 complex controls T cell homeostasis by maintaining surface TCR levels via regulating TCR(+) endosome trafficking. Sci Rep 7:8952.

48. Baldering, T. N., M. S. Dietz, K. Gatterdam, C. Karathanasis, R. Wieneke, R. Tampe, and M. Heilemann. 2019. Synthetic and genetic dimers as quantification ruler for single-molecule counting with PALM. Mol Biol Cell 30:1369-1376.

<div align="center">**Figure 1**</div>

Figure 1: Photoconversion and photobleaching of mEos3.2. (**A**) Time series of fluorescence

micrographs of a field of *S. pombe* cells expressing cytoplasmic mEos3.2 at the 1[st], 31[st], 61[st], and

91[st] time cycles. At each of the 100 cycles, a point-scanning confocal microscope illuminated the

cells simultaneously at both 405 nm and 561 nm, 19-slices in a Z stack were imaged with a total

exposure time of 4.23 s and sum-projected with the same contrast. Scale bar = 10 µm. (**B**) Time

course of the fluorescence signal per cell at 566-719 nm (after autofluorescence subtraction).

Fitting Eq. 7 of the 3-state model in panel C (line) to the data (dots) gave a photoconversion rate

constant (k_{act}) of 1.2×10^{-2} s^{-1} (95% CI: 1.16 -1.24 x 10^{-2}) and a photobleaching rate constant (k_{bl})

of 1.6×10^{-3} s^{-1} (95% CI: 1.5-1.7 x10^{-3}). (**C**) Three state model for mEos3.2 photoconversion and

bleaching. Illumination at 405 nm photoconverts mature mEos3.2 molecules from the green (G)

state to the red (R) state with a photoconversion rate constant of k_{act}. Illumination at both

wavelengths photobleaches red mEos3.2 molecules with a rate constant of k_{bl}.

Figure 2

Figure 2: Effects of formaldehyde fixation on the fluorescence signal of mEos3.2 and rate

constants for photoconversion and photobleaching. (A) Time courses of the fluorescence

signal per *S. pombe* cell expressing cytoplasmic mEos3.2 illuminated at 405 nm (22 µW) and

561 nm (15 µW) by point-scanning confocal microscopy under 3 conditions: live cells (red dots)

or cells fixed with 2% formaldehyde for 15 min (blue dots) or 30 min (black dots) in EMM5S

medium and imaged in EMM5S. Nine fields of view (FOV) of 85 µm x 85 µm were taken over

time for each condition. Plots are weighted mean (dots) and standard deviations (shaded area) of

the fluorescence signal per cell. **(B)** Time courses of the total autofluorescence signal per

wildtype *S. pombe* cell under the same conditions as in panel A. Four FOVs of 85 µm x 85 µm

over time were taken for each condition. Plots show weighted mean (dots) and standard

deviations (shaded area) of the fluorescence signal per cell. **(C)** Time courses of the mEos3.2

fluorescence signal per cell at 566-719 nm after autofluorescence subtraction. Eq. 7 of the 3-state

model was fit to the experimental data (dots). The lines are theoretical curves using the

parameters that best fit the data. **(D-G)** Comparison of parameters of live cells and cells fixed for

15 or 30 min. The error bars are 95% confidence intervals of the parameters (Table S1). **(D)** The

product of total number of molecules per cell and signal of an R-state mEos3.2 molecule per

frame (M_n x ε_f) from the fit. **(E)** Photoconversion rate constant (k_{act}) from the fit. **(F)**

Photobleaching rate constant (k_{bl}) from the fit. **(G)** Time-integrated signal per cell ($\overline{R_s}$)

calculated using Eq. 8.

                                            Figure 3

Figure 3: Effects of pH and DTT on the fluorescence signal and rate constants for

photoconversion and photobleaching of mEos3.2 in fixed *S. pombe* cells. (A) Time courses of

the fluorescence signal per cell expressing cytoplasmic mEos3.2 (after autofluorescence

subtraction). The cells were fixed with 2% formaldehyde for 30 min and illuminated at 405 nm

(22 µW) and 561 nm (15 µW) by point-scanning confocal microscope under 9 different buffer

conditions: (red dot) 50 mM MES (pH 5.5); (red circle) 50 mM MES (pH 5.5) with 1 mM DTT;

(orange dot) 50 mM MES (pH 6.5); (orange circle) 50 mM MES (pH 6.5) with 1 mM DTT;

(green dot) 50 mM Tris-HCl (pH 7.5); (green circle) 50 mM Tris-HCl (pH 7.5) with 1 mM DTT;

(blue dot) 50 mM Tris-HCl (pH 8.5); (blue circle) 50 mM Tris-HCl (pH 8.5) with 1 mM DTT:

(blue triangle) 50 mM Tris-HCl (pH 8.5) with 10 mM DTT. The continuous lines are best fits of

Eq. 7 of the 3-state model to the time courses of the fluorescence signal per cell. Fig. S7 reports

the raw data. **(B-E)** Dependence of the parameters on pH and DTT. The error bars are 95%

confidence intervals for the parameters (Table S2). **(B)** Time-integrated signal per cell ($\overline{R_s}$)

calculated using Eq. 8. **(C)** The product of total number of molecules per cell and the signal of an

R-state mEos3.2 molecule per frame (M_n x ε_f) from the fit. **(D)** Photoconversion rate constant

(k_{act}) from the fit. **(E)** Photobleaching rate constant (k_{bl}) from the fit.

Figure 4

Figure 4: The effects of 405-nm and 561-nm laser intensities on the fluorescence signal and

rate constants for photoconversion and photobleaching of mEos3.2 in live and fixed *S.*

***pombe* cells.** Cells expressing cytoplasmic mEos3.2 were imaged live (red circles) or fixed with

2% formaldehyde for 30 min in EMMS5 medium and mounted in EMM5S medium (black

circles) or 50 mM Tris-HCl (pH 8.5) with 1 mM DTT (blue circles). Cells were imaged with 7

different laser intensities by confocal microscope and 6 different laser intensities by widefield

microscope. For confocal imaging, the laser powers and corresponding average intensities were

set as follows: Panels A, E, I and M, the 561 nm laser power was set at 15 µW (~ 0.3 W/cm^2)

and the 405 nm laser intensity ranged from ~ 0.2 - 0.8 W/cm^2; Panels B, F, J, and N, the 405 nm

laser power was set at 22 µW (~ 0.2 W/cm^2) and the 561 nm laser intensity ranged from ~ 0.15 -

0.5 W/cm^2. Four FOVs were taken for each condition with cells expressing mEos3.2. Two FOVs

were taken for each condition with wildtype cells. For widefield imaging, 561 nm laser intensity

was set at 1 kW/cm^2 for panels C, G, K, O; 405 nm laser intensity was set at 1 W/cm^2 for panels

D, H, L, P. Eight to ten FOVs were taken for each condition with cells expressing mEos3.2.

Three or four FOVs were taken for each condition with wildtype cells. Eq. 7 of the 3-state model

was fit to the time courses of fluorescence signal to determine the parameters giving the best fit:

(A-D) The product of total number of molecules per cell and the signal of an R-state mEos3.2

molecule per frame (M_n x ε_f); **(E-H)** photoconversion rate constant (k_{act}); and **(I-L)**

photobleaching rate constant (k_{bl}) (Table S3, S4). **(M-P)** Time-integrated signal per cell ($\overline{R_s}$)

calculated using Eq. 8 (Table S3, S4). The error bars in the left 2 columns are 95% confidence

intervals of the fit. The error bars in the right 2 columns are the weighted standard deviations

among different FOVs. Fig. S8 and S9 report the raw data.

Figure 5

Figure 5: The effect of fixation and imaging buffer on photon counts from mEos3.2 by

single molecule localization microscopy (SMLM). SMLM imaging of cytoplasmic mEos3.2 in

S. pombe cells with continuous illumination at 405 nm (1 W/cm

^{2}) and 561 nm (1 kW/cm^2) under 3 conditions: live cells (red), cells fixed with 2% formaldehyde

in EMM5S medium for 30 min and mounted in EMM5S synthetic medium (black) or mounted in

50 mM Tris-HCl (pH 8.5) with 1 mM DTT) (blue). Four FOVs of 40 µm x 40 µm were taken

over time at 50 fps for 15,000 frames for each condition. All emission bursts between frame

5,000 and 10,000 were localized to measure the photon counts from single red mEos3.2

molecules in each 20-ms frame, when the mEos3.2 molecules were sparse enough for Gaussian

center fitting. The curves show the cumulative probability distribution of the photon counts of

single mEos3.2 molecules in each 20-ms frame under all three conditions. Inset: The table

reports the mean number of photons (± standard deviation between the 4 FOVs) emitted by

single mEos3.2 molecules in each 20-ms frame under the three conditions. About 2 – 5 x 10^5

emission bursts were recorded for the histogram.

Figure 6

Figure 6: An intermediate state of mEos3.2 that requires 561-nm illumination to reach the

red-state (A) Time courses of the fluorescence signal per live *S. pombe* cell expressing mEos3.2

and subjected to alternating illumination by point-scanning confocal microscopy at 561 nm (37

µW) for 40 cycles followed by illumination at 405 nm (56 µW) for 20 cycles. Illumination at 561

nm increased the fluorescence signal beyond the start of the 561-nm illumination period after

each period of 405-nm illumination. **(B)** Comparisons of the time courses of the fluorescence

signals during 7 periods with 40 cycles of 561-nm illumination (not including the first 561-nm

illumination period before any 405-nm irradiation) from panel A. Eq. 14 of the 4-state model in

panel D (lines) was fit to these time courses (dots) to determine rate constants giving the best fit.

The mean activation rate constant ($k_{act,561}$) from the I-state to the R-state is 0.34 s^{-1} (SD: 0.02)..

(C) Normalized number of red (red), intermediate (black) and green molecules (green) at the

beginning of each 561-nm illumination period. The normalized number of red and intermediate

molecules were estimated from fitting the Eq. 14 of the 4-state model to the time courses of

fluorescence signal during the 7 periods of 561-nm illumination in Panel B. The depletion of

green molecules was estimated from subtracting the fluorescence signal at the last cycle of the

previous 561-nm illumination period from the signal at the first cycle of the previous 561-nm

illumination period. **(D)** Four-state model for mEos3.2 photoconversion and bleaching.

Illumination at 405 nm photoconverts mature mEos3.2 molecules from the G-state to the R-state

with a photoconversion rate constant of k_{act}. A subpopulation of the R-state molecules enter an

intermediate state (gray circle), and irradiation at 561 nm activates the I-state mEos3.2 molecules

to the R-state with a rate constant of $k_{act,561}$. Illumination at both wavelengths photobleaches red

mEos3.2 molecules with a rate constant of k_{bl}. See Fig. S4 and S5 for supplementary information.

A

1st time cycle

31st time cycle

61st time cycle

91st time cycle

B

C