Lateral and longitudinal fish eDNA distribution in

dynamic riverine habitats

Running title: spatio-temporal eDNA distribution

Bettina Thalinger[1,2*], Dominik Kirschner[1,3], Yannick Pütz[1], Christian Moritz[4], Richard

Schwarzenberger[4], Josef Wanzenböck[5] & Michael Traugott[1,6]

[1] Department of Zoology, Universität Innsbruck, Technikerstr. 25, 6020 Innsbruck, Austria

[2] Centre for Biodiversity Genomics, University of Guelph, 50 Stone Road E, N1G2W1,

Guelph, Ontario, Canada

[3] Department of Aquatic Ecology, Eawag: Swiss Federal Institute of Aquatic Science and

Technology, Dübendorf, Switzerland

[4] ARGE Limnologie GesmbH, Hunoldstr. 14, 6020 Innsbruck, Austria

[5] Research Department for Limnology, Mondsee, Universität Innsbruck, Mondseestr. 9, 5310

Mondsee, Austria

[6] Sinsoma GmbH, Lannes 6, 6176 Voels, Austria

*Corresponding author:

Bettina Thalinger, bettina.thalinger@gmail.com

Centre for Biodiversity Genomics, University of Guelph, 50 Stone Road E, N1G 2W1,

Guelph, Ontario, Canada; phone: +1 519-824-4120 Ext. 53800

Keywords: lotic, environmental DNA, hydrology, sampling scheme, eDNA plume

Abstract

Assessing the status and distribution of fish populations in rivers is essential for management and conservation efforts in these dynamic habitats. Currently, methods detecting environmental DNA (eDNA) are being established as an alternative and/or complementary approach to the traditional monitoring of fish species. In lotic systems, a sound understanding of hydrological conditions and their influence on the local target detection probability and DNA quantity is key for the interpretation of eDNA-based results. However, the effect of seasonal and diurnal changes in discharge and the comparability of semi-quantitative results between species remain hardly addressed. We conducted a cage experiment with four fish species (three salmonid and one cyprinid species) in a glacier-fed, fish-free river in Tyrol (Austria) during summer, fall, and winter discharge situations (i.e. 25-fold increase from winter to summer). Each season, water samples were obtained on three consecutive days at 13 locations downstream of the cages including lateral sampling every 1-2 m across the wetted width. Fish eDNA was quantified by species-specific endpoint PCR followed by capillary electrophoresis. Close to the cages, lateral eDNA distribution was heterogenous and mirrored cage placement within the stream. In addition to the diluting effect of increased discharge, longitudinal signal changes within the first 20 m were weakest at high discharge. For downstream locations with laterally homogenous eDNA distribution, the signals decreased significantly with increasing distance and discharge. Generally, the eDNA of the larger-bodied salmonid species was less frequently detected, and signal strengths were lower compared to the cyprinid species. This study exemplifies the importance of hydrological conditions for the interpretation of eDNA-based data across seasons. To control for heterogenous eDNA distribution and enable comparisons over time, sampling schemes in lotic habitats need to incorporate hydrological conditions and species traits.

Keywords: lotic, environmental DNA, hydrology, sampling scheme**,** *Salmo trutta*, *Oncorhynchus mykiss*, *Salvelinus fontinalis*, *Phoxinus phoxinus***,**

Introduction

In times of rapid environmental changes there is a growing need for biomonitoring in both terrestrial and aquatic systems (Cardinale et al., 2012; Tickner et al., 2020). Reliable and cost-effective approaches for species detection are thus key for tracking species in time and space and informing conservation and management efforts (Jetz et al., 2019). Molecular methods, like the detection of environmental DNA (eDNA) released by organisms into their environment, have the capacity to accommodate this demand as they are non-invasive, sensitive, and enable the processing of large sample numbers (Barnes & Turner, 2016; Deiner et al., 2017; Thomsen & Willerslev, 2015). In aquatic habitats, eDNA can be used for the monitoring of taxa in both lotic and lentic systems with a focus on endangered or invasive species and flexible year-round application (Beng & Corlett, 2020; e.g. Harper, Griffiths, et al., 2019; Thomsen & Willerslev, 2015).

Rivers and streams contain, absorb, and transport eDNA from aquatic and terrestrial species and are thus ideal for cross-habitat species detection (Deiner, Fronhofer, Mächler, Walser, & Altermatt, 2016; Sales et al., 2020), although the dynamic nature of these ecosystems leads to constantly changing conditions during sampling (Shogren et al., 2017; Willett, McCoy, Taylor Perron, Goren, & Chen, 2014). Nevertheless, aspects such as a species' upstream distribution limits (Carim et al., 2019; Robinson, de Leaniz, & Consuegra, 2019) or local abundance and its change over time (Doi et al., 2017; Levi et al., 2019; Thalinger, Wolf, Traugott, & Wanzenböck, 2019) have been successfully examined in lotic systems via eDNA. Often, these efforts are combined with traditional monitoring techniques to confirm molecular results and facilitate their interpretation (Evans, Shirey, Wieringa, Mahon, & Lamberti, 2017; Wilcox et al., 2016) and there is generally a good consensus between molecular and non-molecular data.

However, changes in hydrological conditions have profound influence on the distribution and persistence of eDNA in the water column (reviewed by Harrison, Sunday, & Rogers, 2019), which makes the interpretation of local eDNA signals or comparisons between sampling campaigns challenging. The longitudinal eDNA detection probability in

rivers depends on dilution, transport, deposition, resuspension, and degradation (reviewed

by Harrison et al., 2019). These effects were examined for small streams (Fremier, Strickler,

Parzych, Powers, & Goldberg, 2019; Shogren et al., 2017; Wilcox et al., 2016) and large river

systems (Deiner & Altermatt, 2014; Pont et al., 2018), revealing the behavior of eDNA as

similar to that of fine particulate organic matter. Recently, studies also focused on lateral

eDNA distribution, describing a "plume" downstream of the source and gradual lateral

homogenization with increasing downstream distance (Laporte et al., 2020; Wood, Erdman,

York, Trial, & Kinnison, 2020). Albeit discharge-associated changes in longitudinal eDNA

distribution were previously examined for small streams (Jane et al., 2015), changes in the

size and shape of eDNA plumes have been neglected so far.

Alpine rivers exemplify the benefits of eDNA-based monitoring as well as the

challenges associated with sampling campaigns in such dynamic ecosystems. The prevailing

low water temperatures permit only humble population densities and a limited species

inventory at risk of biodiversity loss due to increased climate change effects (Settele et al.,

2015). Anthropogenic influences in the form of straightened riverbeds, hydropower plants,

and dams add to this strained situation and intensify the need for monitoring the remaining

natural fish populations (Faulks, Gilligan, & Beheregaray, 2011; Fette, Weber, Peter, &

Wehrli, 2007). The discharge in these rivers varies with changing seasons from winter

drought to high water levels in spring and summer (snow melt and glacial melt), with

intermediate conditions in fall. Additionally, the melting processes induce substantial diurnal

discharge changes in spring and summer (Bard et al., 2015). Therefore, traditional

monitoring via electrofishing is only possible outside of protected periods (e.g. spawning

seasons) and at low discharge and turbidity in fall and early winter. eDNA-based approaches

can potentially overcome these limitations, but only after accounting for the spatio-temporal

dynamics of Alpine rivers and by optimizing sampling schemes and carefully interpreting the

obtained results.

In this study, we examined changes in lateral and longitudinal eDNA distribution

between seasons in a glacier-fed Alpine river. The experiments were conducted with caged

fish and we used species-specific endpoint PCR combined with capillary electrophoresis

(celPCR) to investigate how longitudinal and lateral eDNA detection probability and signal

strength vary on a small scale (<20 m) between summer, fall, and winter, (i.e. high, medium,

and low water levels) with up to 25-fold variations in discharge. Additionally, the longitudinal

change in eDNA signal strength was examined at an intermediate scale (~1 km) for fish

species of different size, but constant total biomass.

Materials and Methods

Study site

The three caged-fish trials took place in the Melach, a fish-free, glacier-fed, Alpine river in

Lüsens, Tyrol (Latitude 47°7'0,32", Longitude 11°8'4,92", WGS1984) in August (22nd-26th)

2016, November (21st-25th) 2016, and September (26th-29th) 2017. The Melach shows

seasonal and daily fluctuations in discharge and sediment load, typical for glacially

influenced rivers in the Alps (Sertić Perić, Jolidon, Uehlinger, & Robinson, 2015). Following

the European river zonation, the sampling site is located in the epirhithral of the river at an

elevation of 1700 m a.s.l. Permanent fish populations cannot be established in this part of the

river due to extreme discharge situations. Additionally, a transverse structure at the

downstream end of the examined range prohibited potential upstream migration.

Experimental design

All handling of study animals conformed to Directive 2010/63/EU, and permission for field

work was granted by the fishing area manager H. Raffl. Prior to the cage experiment, the

absence of fish was confirmed via electrofishing, starting 1.3 km downstream at the

transverse structure. For each of the three trials, two steel cages 1 × 1 × 0.6 m with "mesh"

size of 10 mm and 5 mm, respectively, were installed at the same location in full current of

the river for the duration of the experimental run (four days), secured against drift, and

equipped with a few stones to provide natural structure and potential resting places. Four fish

species were used in the experiment: A cyprinid species (Eurasian minnow – *Phoxinus phoxinus* L. 1758) was placed in the cage with the narrow mesh size and three salmonid species (brown trout – *Salmo trutta* L. 1758, brook trout – *Salvelinus fontinalis* M. 1814, and rainbow trout – *Oncorhynchus mykiss* W. 1792) were placed in the other cage. On days two, three, and four of each trial, a minimum of 200 g fish per species (exception: *P. phoxinus* on day one in August (196 g) and in November (183 g)) were placed in the cages, corresponding to two to five individuals per salmonid species and 40 to 90 *P. phoxinus* individuals. At the end of each sampling day, fish were removed from the cages and exchanged for new individuals to generate true biological replicates, prevent adaption to the local conditions, and reduce stress-periods for the animals. The fish were kept in separate tanks afterwards to avoid the multiple use of individuals. Due to their life cycle and therefore a lack of catchable individuals, the availability of *P. phoxinus* was limited in November and the same individuals had to be used on all sampling days.

Two-liter water samples were used for the detection of fish eDNA. On the first day of each trial (prior to fish placement in cages), control samples were taken from at least five locations between the cage positions and the transverse structure. On days two, three, and four, water samples were taken daily at 13 locations downstream of the cages (1.3 km, 555 m, 423 m, 323 m, 223 m, 130 m, 65 m, 33 m, 16.8 m, 8 m, 4 m, 2.5 m, and 0.5 m) at approximately the same daytime and before fish were exchanged. Because of changes in the structure of the riverbed, the distance between the cages and the first seven transects sometimes varied by one to two meters between trials, which was accounted for during data analyses. At 0.5 m to 130 m distance, two to four water samples were taken in transects, depending on the width of the river (Fig. 1). Further downstream, one sample was taken per location. In November, only two to three transect samples were taken due to limited discharge, resulting in reduced wetted width, and sampling at transect 5 had to be omitted due to a slight position change of the cages. Each day, sampling was carried out from the most downstream location moving upstream towards the cages. In September, two such sampling runs were carried out per day.

The water samples were collected right below the stream surface using 2 L wide-neck

bottles, which had been treated with chlorine bleach (3.6 g sodium hypochlorite per 100 g

liquid) overnight and thoroughly washed using fish-DNA-free tap water. Filtration was carried

out on 500 ml filter towers with a peristaltic pump (Solinst; Model 410) and glass fibre filters

with 47 mm diameter (1.2 µm mesh width, Whatman GF/C), followed by storage at –80 °C

until further processing. In case of filter clogging (high turbidity; only in August), up to three

filters were utilized per sample. During all water processing steps, DNA-free gloves were

worn and frequently changed. All multi-use equipment was soaked in chlorine bleach for at

least ten minutes between samples and thoroughly rinsed using MilliQ water. Forceps for

filter handling were singed three times prior to each use. Every ten samples, 2 L of MilliQ

water were filtered as negative controls to check for cross-contamination during fieldwork,

whereas a sample from the fish transport container served as daily positive control.

Total discharge and its lateral differences were measured with a FlowTracker

(Sontek, USA) during the August trial, and with salt and a TQ Tracer (Sommer Messtechnik,

Austria) during the November and September trials. In August, discharge increased during

the sampling time (10 AM to 5 PM) due to the strong glacial influence, but remained constant

along the longitudinal gradient. Therefore, diurnal discharge was measured once and

supported by point-measurements at 3 PM. In September and November, the inflow of side

streams caused an increase along the 1.3 km sampling range, with hardly any changes in

the course of a day. Hence, measurements were carried out daily directly at the cages, at

223 m, 555 m, and 1,300 m downstream, and supported by a diurnal measurement per trial.

Discharge at each water sampling point (in space and time) was interpolated; no rainfall

occurred during any of the sampling days. For all three trials, turbidity was measured with a

turbidity meter using an infrared light source (AL250T-IR, Aqualytic, Germany). Water

temperature, pH, conductivity, and oxygen saturation were obtained with a multi-parameter

probe (WTW, Germany).

Laboratory processing

All molecular work was carried out in a clean-room laboratory at the University of Innsbruck

(Austria) compliant with ancient-DNA processing standards. First, filters were lysed using

190 µL TES (0.1 M TRIS, 10mM EDTA, 2% sodium dodecyl sulphate; pH 8) and 10 µL

Proteinase K (VWR, 20 mg/mL) each, followed by over-night incubation on a rocking platform

at 56°C. Then, filters were transferred to plastic inserts with a perforated bottom and

centrifuged for 10 min at 14,000 rpm. The entire resulting lysate was extracted on the

Biosprint 96 platform (QIAGEN) using a custom DNA-uptake protocol which combined the

DNA contained in up to 900 µL lysate, prior to extraction with the "BS 96 tissue protocol"

according to the manufacturer's instructions; except for elution in 100 µL TE. Lysates of

filters stemming from the same water sample were combined during the uptake process.

Each 96-well plate contained four extraction negative controls and all resulting eluates were

stored at -32°C until further processing.

Target DNA was amplified using species-specific primers and celPCR. These were

run on Nexus Mastercyclers (Eppendorf); each plate included at least one positive control

(DNA extract from target species) and one negative control (molecular grade water).

Previously published primers (Thalinger et al., 2016) and a newly developed primer pair for

S. trutta (Table 1) were utilized. During PCR optimization, primers were tested for specificity

against other fish species and aquatic invertebrates occurring in Central European

freshwaters (Thalinger et al., 2016). Additionally, we specified assay sensitivity following Sint

et al. (2012): reliable positive amplifications were possible for all primer pairs from 10 DNA

ds. Each 10 µL PCR master mix contained 1 × Multiplex reaction mix (QIAGEN), 0.5 µM of

each primer, 30 mM TMAC, 5 µg BSA (Bovine Serum Albumin) and 3.2 µL DNA extract.

Optimized thermocycling conditions were: denaturation at 95°C for 15 min, 35 cycles of 94°C

for 30 s, 64°C (66°C for *P. phoxinus*) for 3 min and 72°C for 60 s followed by final elongation

at 72°C for 10 min. Target DNA signal strength was determined via capillary electrophoresis

on the QIAxcel (Qiagen) with the associated software QIAxcel Screengel version 1.6.0.10

using the method AM320-30s. The fluorescent signal measured in Relative Fluorescence

Units (RFU) was used as a semi-quantitative measure of target DNA (Thalinger, Pütz, &

Traugott, 2020; Thalinger et al., 2019) and signals ≥0.08 RFU were deemed positive. To

verify the applicability of this approach, 40 samples positive for *S. trutta* were re-tested with

digital PCR (Supporting Information (SI) 1a). All extraction and PCR negative controls

resulted negative, however, in the August and the November trial six of 16 negative controls

showed low levels of contamination (≤0.2 RFU), hence, fluorescence values of potentially

affected field samples (10.9% of all obtained PCR results) were down-corrected by the

respective values.

Statistical analysis

All data were analyzed in R (R Core Team, 2020) and visualized with "ggplot2" (Hadley

Wickham, 2016), "ggpubr" (Kassambara, 2019), and "viridis" (Garnier, 2018). We generated

eDNA heatmaps per fish cage, i.e. for salmonids and *P. phoxinus,* for each of the three trials

up to 20 m and 65 m downstream distance. The lateral distance from the orographically left

shore (x), the longitudinal distance from the respective cage (y) and mean RFU (z) were

used for the linear interpolation of irregular gridded data with the "akima" package (Akima &

Gebhardt, 2016). eDNA signals were not extrapolated towards the edge of the water body

and RFU were interpolated on a 5 × 5 cm grid.

Differences in detection rates and eDNA signal strengths amongst salmonid species

up to 33 m downstream distance were examined per trial with Z-tests, Kruskal-Wallis tests,

and Holm-Sidak-corrected p-values. As no significant differences were detected (SI 1b),

salmonid data remained pooled for the subsequent analysis of small-scale lateral and

longitudinal eDNA distribution using the packages "reshape2" (H. Wickham, 2007), "ade4"

(Bougeard & Dray, 2018; Chessel, Dufour, & Thioulouse, 2004), and "data.table" (Dowle &

Srinivasan, 2020). Horizontal changes in interpolated eDNA signals were examined per cage

and trial at 2.5 m, 5 m, 10 m and 20 m by testing salmonid and *P. Phoxinus* data for

normality and calculating their Spearman Rank Correlation. To investigate the magnitude of

longitudinal eDNA signal changes between 2.5 m and 20 m, differences between eDNA

signals of adjacent interpolated datapoints were calculated 1 m from the left shore, the

stream center, and 1m from the right shore individually for each trial, and salmonids and

P. phoxinus, respectively. Differences were tested for normality and per position, Kruskal-

Wallis-tests followed by pairwise Wilcoxon Rank Sum Tests were used to compare between

trials for each position (left, center, right) and taxon (salmonids, *P. phoxinus*); all p-values

were Holm-Sidak-corrected.

To analyze the relationship between eDNA signals, distance from the cages, and

discharge on a larger scale, only transects with homogenous eDNA distribution at all trials

were considered. Therefore, salmonid RFU were tested for normal distribution with a

Shapiro-Wilk-test, followed by Kruskal-Wallis-tests per transect and trial. *P. phoxinus* data

were not considered as the dataset was too small for separate testing. Also, mirrored lateral

eDNA distributions between the two cages would potentially mask patterns, if considered

together. Based on these tests (SI 1c), only data with downstream distances ≥ 130 m were

used for the two subsequent analyses: once entering detections as binary variable (0/1), and

once using RUF of positive samples and considering non-detections as random results

induced by the sampling process (Hagenaars & McCutcheon, 2002). To account for the

binary nature and the right-skewed distribution of RFU, Generalized Linear Models (GLM)

were used with family "binomial" and a logit-link for detections (0/1) and family "Gamma" with

a log-link for RFU (Faraway, 2016). After initial data inspection, discharge, downstream

distance, their interaction and fish species were entered as independent variables in both

models, and *P. phoxinus* was set as base category (see SI 2 for the complete R code). For

both models, simulated, scaled residuals were calculated (package: "DHARMa" (Florian

Hartig, 2020); function: "simulateResiduals"; n = 1000) to confirm model fit and test for

outliers and overdispersion.

Results

Altogether, 306 water samples were analyzed: 84, 168, and 54 from the August, September,

and November trial respectively, and average discharge during sampling 0-20 m downstream

of the cages was 835 L/s in August, 174 L/s in September, and 61 L/s in November. Fifty

percent of the 1,224 PCRs resulted positive with the majority of detections occurring in

September and November. The environmental variables turbidity (0 – 4.6 NTU), oxygen

saturation (9.43 – 11.25 mg/L), water temperature, (4.2 – 10.5°C), conductivity (49.3 – 69.2

µS/cm), and pH (7.11 – 7.6) were near-constant between the three trials. Only in August

during sampling within 100 m of the cages (i.e. at a later daytime) turbidity increased (88.7

NTU ± 50.3 NTU DS) and conductivity decreased (36.5 µS/cm ± 9.70 µS/cm SD) because of

glacial run-off. Within the first 30 m downstream of the cages, RFU per trial did not differ

between the three salmonid species (SI 1b), leading to a combined analysis of salmonid

eDNA signals on a small scale: heatmaps of eDNA signals up to 20 m downstream of the

cages revealed eDNA plumes below the respective fish cage and much weaker or no signals

at small lateral distances of ~1 m (Fig. 2). In August, cages were placed in the center of the

stream and positive samples within 20 m distance were almost exclusively obtained from

sampling points with no lateral offset (Fig. 1 and Fig. 2). The interpolated eDNA signals from

the salmonids and *P. phoxinus* obtained in August showed significant positive correlations at

all examined distances (2.5 m, 5 m, 10 m, and 20 m; Table 2). In September, the main water

flow out of the salmonid cage was observed towards the orographically right edge of the

river; the main water flow from the *P. phoxinus* cage was on the opposite side. This situation

was mirrored by high target eDNA signals downstream of the respective cage, low eDNA

signals at the opposite sides, and significantly negative correlations between salmonid and

P. phoxinus RFU at 5 m, 10 m, and 20 m (Table 2). In November, eDNA signals were

highest for both salmonids and *P. phoxinus* and the eDNA distribution pattern was similar to

September with significant negative correlations at the same downstream distances (Table

2). The small-scale longitudinal changes of the interpolated eDNA signals at the left, center,

and right position differed significantly for both salmonids and *P. phoxinus* (Table 3). The

longitudinal changes were strongest in November, with e.g. *P. phoxinus* values decreasing

on average 0.036 RFU ± 0.05 RFU SD with each meter; the smallest changes were detected

for salmonids in August at the left and right position (Table 3, Fig. 2).

At 130 m downstream distance from the cages and beyond, salmonid signals were

laterally homogenous during all three trials (SI 1c) and thus used to examine large-scale

patterns of detection rates and RFU. In August, only few samples resulted positive with low

mean values (e.g. *P. phoxinus* 0.25 RFU ± 0.16 RFU SD; Fig. 3 and Fig. 4). In September,

the signals remained at a similar level, but 100% of *P. phoxinus* PCRs and 71% of salmonid

species PCRs resulted positive. In November, mean *P. phoxinus* eDNA signals were highest

(0.34 RFU ± 0.22 RFU SD), and salmonid signals remained similar to September (Fig. 3 and

Fig. 4). The longitudinal increase in discharge was similar in September and November with

a 2.1-fold increase from 201 L/s to 335 L/s and a 2.7-fold increase from 71 L/s to 167 L/s,

respectively (Fig. 3). Both GLMs (for binomial data and RFU) passed the diagnostics for

model fit, outliers, and overdispersion. They displayed similar patterns of parameter

estimates and significances (Table 4 and Table 5): salmonids were significantly less likely to

be detected and had significantly lower RFU in comparison to *P. phoxinus*. Discharge and

downstream distance were significantly correlated in both models and only the individual

effect of discharge was not significant ($p > 0.05$) in the Gamma GLM (Table 5).

Discussion

Our results demonstrate profound spatio-temporal changes in eDNA distribution induced by

seasonal discharge conditions. The lateral extent of the eDNA plumes downstream of the

caged fish mirrored their position inside the river, longitudinal changes in eDNA signal

strength were strongest at low discharge, and lateral mixing occurred further downstream at

higher discharge. Additionally, the small-sized *P. phoxinus* individuals emitted a significantly

stronger eDNA signal and had higher detection rates than the three larger-bodied salmonid

species, particularly in November. Along the 1.3 km flow path, discharge and distance both

had significantly negative effects on detection rates and eDNA signal strength.

The eDNA signals recorded at a small lateral and longitudinal scale (<20 m) coincide

with previous results describing a plume-shaped distribution of eDNA downstream of the

source (Laporte et al., 2020; Wilcox, McKelvey, Young, Lowe, & Schwartz, 2015; Wood et

al., 2020). Additionally, the shape of the salmonid and *P. phoxinus* plume mirrored the lateral

position of the respective cage when these were not aligned in flow direction in September

and November. This finding complicates the interpretation of field-derived eDNA results

when local conditions differ across the wetted width, especially as individual species may

prefer specific micro-habitats depending on hydrological conditions (e.g. seeking shelter from

high flow velocities) (Aarts & Nienhuis, 2003). Based on our experimental setup with constant

fish biomasses, it was not possible to confirm the results of Wood et al. (2020) who found a

correlation between increased eDNA quantities and detection probability at higher lateral

distance. However, our data show a comparable effect induced by lower discharge and

slower flow velocities (Wondzell, Gooseff, & McGlynn, 2007) at constant fish biomass. Future

studies examining lateral and longitudinal eDNA distribution should therefore take biomass-

induced variations and hydrological effects into account.

All three salmonid species (average individual mass 101 g) had lower detection

probabilities and lower eDNA signals than *P. phoxinus* (average individual mass 3.5 g) at

similar total biomass. eDNA shedding rates and biomass in most cases exhibit a positive

relationship in controlled and natural settings (Lacoursière-Roussel, Rosabal, & Bernatchez,

2016; Takahara, Minamoto, Yamanaka, Doi, & Kawabata, 2012; Thalinger et al., 2019), but

the influence of fish size is not well studied. Maruyama et al. (2014) found elevated levels of

eDNA release per gram bodyweight in juvenile fish, but attributed this effect to higher

metabolic activity (Vinberg, 1960). In the present study, several scenarios can explain the

difference in detection probability and signal strength: the *P. phoxinus* individuals, although

mostly not juvenile, could indeed have had higher metabolic rates, which are common for

smaller fish species (Clarke & Johnston, 1999). The increased surface to volume ratio of the

smaller fish is another potential explanation as more *P. phoxinus* surface was exposed to the

water flow. The eDNA release rates could generally differ between cyprinids and salmonids,

which also exhibit structural and physiological differences (Freyhof & Kottelat, 2007). Albeit

the daily exchange of fish individuals after eDNA sampling was another potential source of

variation (e.g. slight changes in total biomass or individual counts) we deemed this measure

appropriate to obtain true biological replicates, avoid adaption effects to the stream

environment, and minimize the exposure to stressful flow conditions. Generally, the

salmonids were better adapted to the high flow conditions in August, when some *P. phoxinus*

did not survive the exposure to 1,500 L/s discharge. This could explain the distinct detection

probability for *P. phoxinus*, as dead individuals can emit more eDNA into the water (Tillotson

et al., 2018). However, the difference in eDNA signal strength was most pronounced in

November at ~60 L/s, when all but one individual of *P. Phoxinus* survived. Though celPCR

assay optimization (Sint et al., 2012; Thalinger et al., 2016) can not fully account for

differences in amplification efficiency between primer pairs (Thalinger et al., 2020), primer

bias is an unlikely cause for the strong *P. phoxinus* signals, because detection patterns and

signal strength did not differ between the salmonids. Nevertheless, celPCR produces a semi-

quantitative measure of target copy numbers and without direct comparison to a fully

quantitative approach (i.e. digital PCR), it is not possible to infer absolute eDNA

concentrations from RFU (Thalinger et al., 2020). Most likely, the strong eDNA signals of

P. phoxinus can be attributed to a combination of the aforementioned physiological and

morphological differences. These effects could be even stronger when fish are compared to

other aquatic groups such as amphibians, mussels, or crayfish (Bedwell & Goldberg, 2020;

Robinson, Uren Webster, Cable, James, & Consuegra, 2018; Wacker et al., 2019).

As expected for glacier-fed Alpine rivers and streams, discharge changed substantially

(25-fold) between trials (Bard et al., 2015). The associated diluting effect was clearly visible

from eDNA signal strengths directly downstream of the cages (≤20 m): average *P. phoxinus*

signals at this location ranged from 0.30 RFU ± 0.30 RFU SD in August to 1.06 RFU ± 0.46

RFF SD in November. Few of the samples taken at high discharge >1,000 L/s in August

tested positive, but in this situation the 200 g fish mass per species were less than 0.2 ‰ of

the passing water mass. The summer situation was most challenging to examine, as

discharge and turbidity increased from morning (~270 L/s and clear water) to afternoon

(~1,100 L/s and high turbidity). The analysis of samples with high variations in turbidity

should include this variable in the modelling process to account for changes in extraction

efficiency and inhibition associated with increased turbidity (Harper, Buxton, et al., 2019). We

refrained from doing so as the samples taken in August at high turbidity were not part of the

dataset used for modelling. In contrast to classic fish monitoring via electrofishing, eDNA-

based methods are not restricted to low flow conditions outside of protected periods and

spawning seasons. Based on our results, spring and summer sampling in glacier-fed Alpine

rivers should, however, not take place at high discharge in the afternoon and evening and is

also not advisable during floods and after strong rainfalls. In September and November

discharge remained almost constant over time but increased more than two-fold (side stream

inflow) within the examined 1.3 km. So far, most eDNA studies only discuss the effect of

changing discharge (Harrison et al., 2019; Laramie, Pilliod, & Goldberg, 2015; Wood et al.,

2020) even though it generally increases along the course of running waters and significantly

influenced eDNA detection rates this study. Furthermore, the significance of the discharge-

distance interaction term in our analysis shows the effect of the distinct hydrological

conditions at each trial on eDNA detection rates and signal strengths.

DNA signals and detection rates of both *P. phoxinus* and the salmonids declined

significantly with increasing distance from their source, as expected from previous work on

eDNA deposition and degradation (Harrison et al., 2019). At constant longitudinal and

temporal discharge, average transport distance (S_P) and depositional velocity (v_{dep}) are used

to describe eDNA deposition (Pont et al., 2018; Shogren et al., 2017; Wilcox et al., 2016).

The calculation of v_{dep} relies on flow velocity and depth data, which can fluctuate

considerably in natural and semi-natural rivers and are directly influenced by discharge.

Therefore, we refrained from calculating this factor. S_P is commonly described as the slope

parameter of a first order exponential decline $S_P = 1/\text{-}k$ (but also see Wood et al., 2020). In

our case this would result in an average transport distance of ~ 3,000 m and ~850 m for

P. phoxinus eDNA in September and November, respectively (based on data from locations

with homogenous lateral eDNA distribution), and confirm the positive correlation between

river size in general and transport distance (Deiner & Altermatt, 2014; Jane et al., 2015; Pont

et al., 2018; Shogren et al., 2017). However, S_P is non-generalizable between streams

(Harrison et al., 2019) and thus of limited use outside of a systematic framework

incorporating the complex flow regime of Alpine rivers.

Our work aids to the understanding of how eDNA signals obtained from field-collected

samples can be interpreted for species monitoring and conservation. Sampling campaigns

carried out in dynamic habitats such as Alpine rivers and streams need to account for the

heterogenous lateral eDNA distribution, adapt the sampling scheme to habitat preferences of

the target species, and address the prevailing discharge situation. In a best-case scenario,

the target species has distinct habitat preferences, and discharge is low and constant during

the entire sampling period. Then, eDNA quantities measured directly downstream of suitable

habitats are likely to be directly correlated with local target species biomass (Hinlo,

Lintermans, Gleeson, Broadhurst, & Furlan, 2018). Otherwise, only discharge measurements

at each sampling location can prevent flawed inferences (Thalinger et al., 2019). The

detection of rare species, however, is best accomplished by determining a suitable distance

between sampling points and preliminary investigations of the respective eDNA shedding

rates (cp. Wood et al., 2020). The population densities and the position of individuals within

lotic systems should not be inferred from eDNA signals without any *a priori* knowledge on

local hydrology. Comparisons between species are also not advisable without previous tests

under controlled conditions. Therefore, we advocate for the reporting of the sampling

position, local discharge, time, and species biology during field sampling. In the future, these

data can be incorporated in hydrological models specifically designed for eDNA-based

species monitoring.

Data Availability Statement

All data on fish, discharge, eDNA signals, environmental conditions, sampling and

comparison between celPCR and dPCR have been uploaded to Figshare and are available

at https://doi.org/10.6084/m9.figshare.12380642.v1

Acknowledgements

This research was conducted within the eDNA-Alpfish project funded by the Austria
Research Promotion Agency (FFG); project number 853219. We thank F. Drewes for his
extraordinary patience, support and organisational skills during setting up the cages, fish
handling, and discharge measurements in the face of non-existent mobile reception. We are
grateful to H. Raffl for letting us conduct the experiments in his fishing area and M. Böcker for
supporting this work with her graphic skills. We thank two anonymous reviewers for their
constructive feedback on the original manuscript.

Conflict of interest

459 MT is the co-founder of Sinsoma GmbH, a for-profit company dedicated to DNA analyses in
460 environmental studies; CM and RS are co-founders of the ARGE Limnologie GesmbH, a for-
461 profit consultancy in aquatic ecology.

Author contributions

MT, JW, CM, and RS conceived the study; the experiment was designed by BT, MT, RS, and

CM. Data were acquired and analyzed by BT, DK, and YP. BT wrote the first draft of the

manuscript which was revised by DK, YP, JW, and MT.

References

Aarts, B. G. W., & Nienhuis, P. H. (2003). Fish zonations and guilds as the basis for

assessment of ecological integrity of large rivers. In *Aquatic Biodiversity* (pp. 157–178).

https://doi.org/10.1007/978-94-007-1084-9_11

Akima, H., & Gebhardt, A. (2016). *akima: Interpolation of Irregularly and Regularly Spaced*

Data. Retrieved from https://cran.r-project.org/package=akima

Bard, A., Renard, B., Lang, M., Giuntoli, I., Korck, J., Koboltschnig, G., … Volken, D. (2015).

Trends in the hydrologic regime of Alpine rivers. *Journal of Hydrology*, *529*, 1823–1837.

https://doi.org/10.1016/j.jhydrol.2015.07.052

Barnes, M. A., & Turner, C. R. (2016). The ecology of environmental DNA and implications

for conservation genetics. *Conservation Genetics*, Vol. 17, pp. 1–17.

https://doi.org/10.1007/s10592-015-0775-4

Bedwell, M. E., & Goldberg, C. S. (2020). Spatial and temporal patterns of environmental

DNA detection to inform sampling protocols in lentic and lotic systems. *Ecology and*

Evolution, *10*(3), 1602–1612. https://doi.org/10.1002/ece3.6014

Beng, K. C., & Corlett, R. T. (2020). Applications of environmental DNA (eDNA) in ecology

and conservation: opportunities, challenges and prospects. *Biodiversity and*

Conservation, Vol. 29, pp. 2089–2121. https://doi.org/10.1007/s10531-020-01980-0

Bougeard, S., & Dray, S. (2018). Supervised Multiblock Analysis in R with the ade4 Package.

Journal of Statistical Software, *86*(1), 1–17. https://doi.org/10.18637/jss.v086.i01

Cardinale, B. J., Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C., Venail, P., …

Naeem, S. (2012). Biodiversity loss and its impact on humanity. *Nature*, Vol. 486, pp.

59–67. https://doi.org/10.1038/nature11148

Carim, K. J., Caleb Dysthe, J., McLellan, H., Young, M. K., McKelvey, K. S., & Schwartz, M.

K. (2019). Using environmental DNA sampling to monitor the invasion of nonnative *Esox*

lucius (northern pike) in the Columbia River basin, USA. *Environmental DNA*, *1*(3), 215–

226. https://doi.org/10.1002/edn3.22

Chessel, D., Dufour, A., & Thioulouse, J. (2004). The ade4 Package – I: One-Table Methods.

R News, *4*(1), 5–10. Retrieved from https://cran.r-project.org/doc/Rnews/

Clarke, A., & Johnston, N. M. (1999). Scaling of metabolic rate with body mass and

temperature in teleost fish. *Journal of Animal Ecology*, *68*(5), 893–905.

https://doi.org/10.1046/j.1365-2656.1999.00337.x

Deiner, K., & Altermatt, F. (2014). Transport distance of invertebrate environmental DNA in a

natural river. *PLOS ONE*, *9*(2). https://doi.org/10.1371/journal.pone.0088786

Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., …

Bernatchez, L. (2017). Environmental DNA metabarcoding: Transforming how we

survey animal and plant communities. *Molecular Ecology*, *26*(21), 5872–5895.

https://doi.org/10.1111/mec.14350

Deiner, K., Fronhofer, E. A., Mächler, E., Walser, J. C., & Altermatt, F. (2016). Environmental

DNA reveals that rivers are conveyer belts of biodiversity information. *Nature*

Communications, *7*. https://doi.org/10.1038/ncomms12544

Doi, H., Inui, R., Akamatsu, Y., Kanno, K., Yamanaka, H., Takahara, T., & Minamoto, T.

(2017). Environmental DNA analysis for estimating the abundance and biomass of

stream fish. *Freshwater Biology*, *62*(1), 30–39. https://doi.org/10.1111/fwb.12846

Dowle, M., & Srinivasan, A. (2020). *data.table: Extension of `data.frame`*. Retrieved from

https://cran.r-project.org/package=data.table

Evans, N. T., Shirey, P. D., Wieringa, J. G., Mahon, A. R., & Lamberti, G. A. (2017).

Comparative Cost and Effort of Fish Distribution Detection via Environmental DNA

Analysis and Electrofishing. *Fisheries*, *42*(2), 90–99.

https://doi.org/10.1080/03632415.2017.1276329

Faraway, J. J. (2016). *Extending the Linear Model with R. Generalized Linear, Mixed Effects*

and Nonparametric Regression Models (2nd ed.).

https://doi.org/10.1201/9781315382722

Faulks, L. K., Gilligan, D. M., & Beheregaray, L. B. (2011). The role of anthropogenic vs.

natural in-stream structures in determining connectivity and genetic diversity in an

endangered freshwater fish, Macquarie perch (Macquaria australasica). *Evolutionary*

Applications, *4*(4), 589–601. https://doi.org/10.1111/j.1752-4571.2011.00183.x

Fette, M., Weber, C., Peter, A., & Wehrli, B. (2007). Hydropower production and river

rehabilitation: A case study on an alpine river. *Environmental Modeling & Assessment.*

Retrieved from https://doi.org/10.1007/s10666-006-9061-7

Fremier, A. K., Strickler, K. M., Parzych, J., Powers, S., & Goldberg, C. S. (2019). Stream

Transport and Retention of Environmental DNA Pulse Releases in Relation to

Hydrogeomorphic Scaling Factors. *Environmental Science and Technology*, *53*(12),

6640–6649. https://doi.org/10.1021/acs.est.8b06829

Freyhof, J., & Kottelat, M. (2007). Handbook of European freshwater fishes . In *Ichthyological*

Research (Vol. 55). https://doi.org/10.1007/s10228-007-0012-3

Garnier, S. (2018). *viridis: Default Color Maps from "matplotlib."* Retrieved from https://cran.r-

project.org/package=viridis

Hagenaars, J. A., & McCutcheon, A. L. (2002). Applied Latent Calss Analysis. In J. A.

Hagenaars & A. L. McCutcheon (Eds.), *Applied Latent Class Analysis.* Edinburgh:

Cambridge University Press.

Harper, L. R., Buxton, A. S., Rees, H. C., Bruce, K., Brys, R., Halfmaerten, D., … Hänfling, B.

(2019). Prospects and challenges of environmental DNA (eDNA) monitoring in

freshwater ponds. *Hydrobiologia*, Vol. 826, pp. 25–41. https://doi.org/10.1007/s10750-

018-3750-5

Harper, L. R., Griffiths, N. P., Lawson Handley, L., Sayer, C. D., Read, D. S., Harper, K. J.,

… Hänfling, B. (2019). Development and application of environmental DNA surveillance

for the threatened crucian carp (*Carassius carassius*). *Freshwater Biology*, *64*(1), 93–

107. https://doi.org/10.1111/fwb.13197

Harrison, J. B., Sunday, J. M., & Rogers, S. M. (2019). Predicting the fate of eDNA in the

environment and implications for studying biodiversity. *Proceedings of the Royal Society*

B: Biological Sciences, Vol. 286. https://doi.org/10.1098/rspb.2019.1409

Hinlo, R., Lintermans, M., Gleeson, D., Broadhurst, B., & Furlan, E. (2018). Performance of

eDNA assays to detect and quantify an elusive benthic fish in upland streams. *Biological*

Invasions, *20*(11), 3079–3093. https://doi.org/10.1007/s10530-018-1760-x

Jane, S. F., Wilcox, T. M., Mckelvey, K. S., Young, M. K., Schwartz, M. K., Lowe, W. H., …

Whiteley, A. R. (2015). Distance, flow and PCR inhibition: EDNA dynamics in two

headwater streams. *Molecular Ecology Resources*, *15*(1), 216–227.

https://doi.org/10.1111/1755-0998.12285

Jetz, W., McGeoch, M. A., Guralnick, R., Ferrier, S., Beck, J., Costello, M. J., … Turak, E.

(2019). Essential biodiversity variables for mapping and monitoring species populations.

Nature Ecology and Evolution, Vol. 3, pp. 539–551. https://doi.org/10.1038/s41559-019-

0826-1

Kassambara, A. (2019). *ggpubr: "ggplot2" Based Publication Ready Plots*. Retrieved from

https://cran.r-project.org/package=ggpubr

Lacoursière-Roussel, A., Rosabal, M., & Bernatchez, L. (2016). Estimating fish abundance

and biomass from eDNA concentrations: variability among capture methods and

environmental conditions. *Molecular Ecology Resources*, *16*(6), 1401–1414.

https://doi.org/10.1111/1755-0998.12522

Laporte, M., Bougas, B., Côté, G., Champoux, O., Paradis, Y., Morin, J., & Bernatchez, L.

(2020). Caged fish experiment and hydrodynamic bidimensional modeling highlight the

importance to consider 2D dispersion in fluvial environmental DNA studies.

Environmental DNA, edn3.88. https://doi.org/10.1002/edn3.88

Laramie, M. B., Pilliod, D. S., & Goldberg, C. S. (2015). Characterizing the distribution of an

endangered salmonid using environmental DNA analysis. *Biological Conservation*, *183*,

29–37. https://doi.org/10.1016/j.biocon.2014.11.025

Levi, T., Allen, J. M., Bell, D., Joyce, J., Russell, J. R., Tallmon, D. A., … Yu, D. W. (2019).

Environmental DNA for the enumeration and management of Pacific salmon. *Molecular*

Ecology Resources, *19*(3), 597–608. https://doi.org/10.1111/1755-0998.12987

Maruyama, A., Nakamura, K., Yamanaka, H., Kondoh, M., & Minamoto, T. (2014). The

release rate of environmental DNA from juvenile and adult fish. *PLoS ONE*, *9*(12).

https://doi.org/10.1371/journal.pone.0114639

Pont, D., Rocle, M., Valentini, A., Civade, R., Jean, P., Maire, A., … Dejean, T. (2018).

Environmental DNA reveals quantitative patterns of fish biodiversity in large rivers

despite its downstream transportation. *Scientific Reports*, *8*(1).

https://doi.org/10.1038/s41598-018-28424-8

R Core Team. (2020). *R: A Language and Environment for Statistical Computing.* Retrieved

from https://www.r-project.org/

Robinson, C. V., de Leaniz, C. G., & Consuegra, S. (2019). Effect of artificial barriers on the

distribution of the invasive signal crayfish and Chinese mitten crab. *Scientific Reports*,

9(1), 1–11. https://doi.org/10.1038/s41598-019-43570-3

Robinson, C. V., Uren Webster, T. M., Cable, J., James, J., & Consuegra, S. (2018).

Simultaneous detection of invasive signal crayfish, endangered white-clawed crayfish

and the crayfish plague pathogen using environmental DNA. *Biological Conservation*,

222, 241–252. https://doi.org/10.1016/j.biocon.2018.04.009

Sales, N. G., McKenzie, M. B., Drake, J., Harper, L. R., Browett, S. S., Coscia, I., …

McDevitt, A. D. (2020). Fishing for mammals: Landscape-level monitoring of terrestrial

and semi-aquatic communities using eDNA from riverine systems. *Journal of Applied*

Ecology, *57*(4), 707–716. https://doi.org/10.1111/1365-2664.13592

Sertić Perić, M., Jolidon, C., Uehlinger, U., & Robinson, C. T. (2015). Long-term ecological

patterns of alpine streams: An imprint of glacial legacies. *Limnology and Oceanography*,

60(3), 992–1007. https://doi.org/10.1002/lno.10069

Settele, J., Scholes, R., Betts, R. A., Bunn, S., Leadley, P., Nepstad, D., … Winter, M.

(2015). Terrestrial and Inland water systems. In *Climate Change 2014 Impacts,*

Adaptation and Vulnerability: Part A: Global and Sectoral Aspects (pp. 271–360).

https://doi.org/10.1017/CBO9781107415379.009

Shogren, A. J., Tank, J. L., Andruszkiewicz, E., Olds, B., Mahon, A. R., Jerde, C. L., &

Bolster, D. (2017). Controls on eDNA movement in streams: Transport, Retention, and

Resuspension /704/158/2464 /704/242 /45/77 article. *Scientific Reports*, *7*(1).

https://doi.org/10.1038/s41598-017-05223-1

Sint, D., Raso, L., & Traugott, M. (2012). Advances in multiplex PCR: Balancing primer

efficiencies and improving detection success. *Methods in Ecology and Evolution*, *3*(5),

898–905. https://doi.org/10.1111/j.2041-210X.2012.00215.x

Takahara, T., Minamoto, T., Yamanaka, H., Doi, H., & Kawabata, Z. (2012). Estimation of

Fish Biomass Using Environmental DNA. *PLoS ONE*, *7*(4), e35868.

https://doi.org/10.1371/journal.pone.0035868

Thalinger, B., Oehm, J., Mayr, H., Obwexer, A., Zeisler, C., & Traugott, M. (2016). Molecular

prey identification in Central European piscivores. *Molecular Ecology Resources*, *16*(1),

123–137. https://doi.org/10.1111/1755-0998.12436

Thalinger, B., Pütz, Y., & Traugott, M. (2020). Endpoint PCR coupled with capillary

electrophoresis (celPCR) provides sensitive and quantitative measures of environmental

DNA in singleplex and multiplex reactions. *BioRxiv*, 2020.10.24.353730.

https://doi.org/10.1101/2020.10.24.353730

Thalinger, B., Wolf, E., Traugott, M., & Wanzenböck, J. (2019). Monitoring spawning

migrations of potamodromous fish species via eDNA. *Scientific Reports*, *9*(1).

https://doi.org/10.1038/s41598-019-51398-0

Thomsen, P. F., & Willerslev, E. (2015). Environmental DNA - An emerging tool in

conservation for monitoring past and present biodiversity. *Biological Conservation*, *183*,

4–18. https://doi.org/10.1016/j.biocon.2014.11.019

Tickner, D., Opperman, J. J., Abell, R., Acreman, M., Arthington, A. H., Bunn, S. E., …

Young, L. (2020). Bending the Curve of Global Freshwater Biodiversity Loss: An

Emergency Recovery Plan. *BioScience*, *70*(4), 330–342.

https://doi.org/10.1093/biosci/biaa002

Tillotson, M. D., Kelly, R. P., Duda, J. J., Hoy, M., Kralj, J., & Quinn, T. P. (2018).

Concentrations of environmental DNA (eDNA) reflect spawning salmon abundance at

fine spatial and temporal scales. *Biological Conservation*, *220*, 1–11.

https://doi.org/10.1016/j.biocon.2018.01.030

Vinberg, G. G. (1960). *Rate of Metabolism and Food Requirements of Fishes*. Nanaimo,

B.C.: Distributed by the Fisheries Research Board of Canada, Biological Station

(1960).

Wacker, S., Fossøy, F., Larsen, B. M., Brandsegg, H., Sivertsgård, R., & Karlsson, S. (2019).

Downstream transport and seasonal variation in freshwater pearl mussel (*Margaritifera*

margaritifera) eDNA concentration. *Environmental DNA*, *1*(1), 64–73.

https://doi.org/10.1002/edn3.10

Wickham, H. (2007). Reshaping Data with the reshape Package. *Journal of Statistical*

Software, *21*(12), 1–20. Retrieved from http://www.jstatsoft.org/v21/i12/

Wickham, Hadley. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Retrieved from

https://ggplot2.tidyverse.org

Wilcox, T. M., McKelvey, K. S., Young, M. K., Lowe, W. H., & Schwartz, M. K. (2015).

Environmental DNA particle size distribution from Brook Trout (Salvelinus fontinalis).

Conservation Genetics Resources, *7*(3), 639–641. https://doi.org/10.1007/s12686-015-

0465-z

Wilcox, T. M., McKelvey, K. S., Young, M. K., Sepulveda, A. J., Shepard, B. B., Jane, S. F.,

… Schwartz, M. K. (2016). Understanding environmental DNA detection probabilities: A

case study using a stream-dwelling char Salvelinus fontinalis. *Biological Conservation*,

194, 209–216. https://doi.org/10.1016/j.biocon.2015.12.023

Willett, S. D., McCoy, S. W., Taylor Perron, J., Goren, L., & Chen, C. Y. (2014). Dynamic

reorganization of River Basins. *Science*, *343*(6175).

https://doi.org/10.1126/science.1248765

Wondzell, S. M., Gooseff, M. N., & McGlynn, B. L. (2007). Flow velocity and the hydrologic

behavior of streams during baseflow. *Geophysical Research Letters*, *34*(24), L24404.

https://doi.org/10.1029/2007GL031256

Wood, Z. T., Erdman, B. F., York, G., Trial, J. G., & Kinnison, M. T. (2020). Experimental

assessment of optimal lotic eDNA sampling and assay multiplexing for a critically

endangered fish. *Environmental DNA*, edn3.64. https://doi.org/10.1002/edn3.64

Tables and Figures

Table 1: Primer pairs used for the molecular analysis of the eDNA samples.

Species	Primer name	5' - 3'	Target gene	Fragment length (bp)	Reference
Phoxinus phoxinus	Pho-Pho-S639 Pho-Pho-A648	CGTGCAGAAGCGGATATAAATAC CCAACCGAAGGTAAAGTCTTATTG	16s	128	Thalinger et al. 2016
Salvelinus **sp.**	Sal-vel-S651 Sal-vel-A651	ATAGTCGGCACCGCCCTT TAACGAAGGCATGGGCTGTT	COI	112	Thalinger et al. 2016
Oncorhynchus mykiss	Onc-myk-S655 Onc-myk-A655	TCTCCCTTCATTTAGCTGGAATC GCTGGAGGTTTTATGTTAATAATGGTC	COI	82	Thalinger et al. 2016
Salmo trutta	Sal-tru-S1002 Sal-tru-A1002	TCTCTTGATTCGGGCAGAACTC CGAAGGCATGGGCTGTAACA	COI	89	

669 The respective target taxon, primer sequence, target gene, fragment length in base pairs and the source for previously published primers are

670 displayed. Please note that the *Salvelinus* sp. primer pair was designed to amplify both *S. fontinalis* and *Salvelinus umbla*.

672   **Table 2:** The correlation between interpolated eDNA signals from the two fish cages.

trial	longitudinal position	W salmonids	p-value	W *P. phoxinus*	p-value	Spearman's rho	p-value
August	2.5 m	0.89	<0.001	0.96	< 0.01	0.89	<0.001
	5 m	0.94	<0.001	0.92	<0.001	0.84	<0.001
	10 m	0.84	<0.001	0.93	<0.001	0.95	<0.001
	20 m	0.61	<0.001	0.97	<0.05	0.62	<0.001
September	2.5 m	0.97	0.12	0.85	<0.001	0.64	<0.001
	5 m	0.92	<0.001	0.87	<0.001	-0.35	<0.01
	10 m	0.92	<0.001	0.91	<0.001	-0.52	<0.001
	20 m	0.87	<0.001	0.95	<0.01	-0.31	<0.01
November	2.5 m	0.42	<0.001	0.96	0.30	0.30	0.11
	5 m	0.84	<0.001	0.91	<0.01	-0.87	<0.001
	10 m	0.82	<0.001	0.76	<0.001	-0.82	<0.001
	20 m	0.96	<0.05	0.95	<0.05	-0.26	<0.05

674   The test statistic (W) and corresponding p-values of Shapiro-Wilk tests for normality are provided per trial, taxon, and for four downstream

675   distances (2.5 m, 5 m, 10 m and 20 m). Spearman's rho and the corresponding p-value were calculated per distance and trial.

Table 3: The difference in longitudinal eDNA signal change for the left shore, center, and right shore of the river. The change was compared between the three trials per position and taxon.

taxon	lateral position	trial	mean RFU-change per m	SD	W	p-value	Chi²	p-value	September p-value	November p-value
salmonids	left	August	0.0004	0.0013	0.76	<0.001	1043	<0.001	<0.001	<0.01
		September	0.0034	0.0050	0.71	<0.001				<0.001
		November	0.0078	0.0294	0.66	<0.001				
	center	August	-0.0040	0.0079	0.53	<0.001	1049	<0.001	<0.001	<0.001
		September	0.0033	0.0130	0.61	<0.001				<0.001
		November	0.0080	0.0133	0.64	<0.001				
	right	August	-0.0011	0.0083	0.80	<0.001	954	<0.001	<0.05	<0.001
		September	0.0048	0.0156	0.77	<0.001				<0.001
		November	0.0063	0.0148	0.68	<0.001				
P. phoxinus	left	August	0.0033	0.0080	0.76	<0.001	1049	<0.001	0.49	<0.001
		September	0.0076	0.0337	0.76	<0.001				<0.001
		November	-0.0300	0.0358	0.72	<0.001				
	center	August	0.0098	0.0269	0.80	<0.001	1023	<0.001	<0.001	<0.001
		September	0.0069	0.0234	0.55	<0.001				<0.001
		November	-0.0360	0.0549	0.61	<0.001				
	right	August	0.0051	0.0122	0.83	<0.001	916	<0.001	<0.01	<0.001
		September	0.0117	0.0130	0.71	<0.001				<0.001
		November	-0.0334	0.0538	0.59	<0.001				

The test statistic (W) and p-value of Shapiro-Wilk tests for normality are provided together with the Chi² and p-values obtained from Kruskal-Wallis tests per position and taxon. The columns "September p-value" and "November p-value" denote for the results of pairwise Wilcoxon Rank Sum tests between individual trials. All displayed p-values were Holm-Sidak corrected.

Table 4: The Generalized Linear Model (Binomial family, logit-link) describing the relation between positive detections and discharge, downstream distance, their interaction, and fish species.

A)	Estimate	Std. Error	z-value	lower 95% CI	upper 95% CI	p-value
(Intercept)	4.64	0.79	5.91	3.17	6.26	<0.001
discharge	-0.01	0.00	-3.42	-0.02	0.00	<0.001
distance	0.00	0.00	-3.39	-0.01	0.00	<0.001
Oncorhynchus mykiss	-2.32	0.43	-5.45	-3.21	-1.52	<0.001
Salvelinus fontinalis	-1.86	0.43	-4.37	-2.75	-1.06	<0.001
Salmo trutta	-1.43	0.43	-3.31	-2.32	-0.61	<0.001
discharge*distance	0.00	0.00	3.32	0.00	0.00	<0.001

B)	Df	Deviance	Resid. Df	Resid. Dev
NULL			335	431.78
discharge	1	3.728	334	428.05
distance	1	0.433	333	427.62
fish species	3	39.212	330	388.4
discharge*distance	1	12.017	329	376.39

689 A) The columns show the test statistics (z-value), the lower and upper 95% confidence intervals, and p-values returned from the GLM for the input
690 variables. Table B) shows the degrees of freedom (Df) residuals and deviance for each variable.

692   **Table 5**: The Generalized Linear Model (Gamma-family, log-link) describing the relation between RFUs and discharge, downstream distance, their

693   interaction, and fish species.

A)	Estimate	Std. Error	t-value	lower 95% CI	upper 95% CI	p-value
(Intercept)	-0.42	0.22	-1.96	-0.82	-0.01	0.05
discharge	0.00	0.00	-1.74	0.00	0.00	0.08
distance	0.00	0.00	-4.21	0.00	0.00	<0.001
Oncorhynchus mykiss	-0.33	0.13	-2.52	-0.58	-0.07	<0.05
Salvelinus fontinalis	-0.38	0.12	-3.14	-0.62	-0.14	<0.01
Salmo trutta	-0.30	0.12	-2.61	-0.53	-0.07	<0.01
discharge*distance	0.00	0.00	3.25	0.00	0.00	<0.01

B)	Df	Deviance	Resid. Df	Resid. Dev
NULL			220	125.14
discharge	1	1.4257	219	123.72
distance	1	3.8624	218	119.85
fish species	3	4.9894	215	114.86
discharge*distance	1	5.0312	214	109.83

A) The columns show the test statistics (t-value), the lower and upper 95% confidence intervals, and p-values returned from the GLM for the input

variables. Table B) shows the degrees of freedom (Df) residuals and deviance for each variable.

Figure 1: The geographic location of the study region and the aerial view (source: Land Tirol –
data.gv.at) of the entire sampling reach is displayed along the schematic view of the cage
placement in the stream and the hydrological conditions prevailing during the August,
September, and November trials. Larger flowlines and eddies code for stronger currents. Due to
reduced discharge, the relative location and number of eDNA samples were not kept uniform
between trials. Note that additional sampling distances up to 1.3 km downstream of the cages
are not displayed.

Figure 2: Heatmaps showing the small-scale eDNA distribution ≤20 m downstream of the cages
during the three examined discharge situations. As eDNA levels differed between
Phoxinus phoxinus and the salmonids, different colour scales where used. However, coloring
per taxon remains constant for the three trials. Isotherms display interpolated differences of 0.05
RFU; the irregular shape in November reflects an increase in wetted width downstream of the
cages. Dashed lines to the left and the bottom of the figures show locations of lateral and
longitudinal transects used to compute small-scale differences in eDNA signals; these transects
differ from the actual water sampling locations.

Figure 3: The eDNA signal strengths of *P. phoxinus* and the salmonid species are displayed for
downstream distances with homogeneous lateral eDNA distribution (≥130 m) for August,
September, and November. Additionally, the discharge at each distance (three values each) is
displayed.

Figure 4: The pie charts display the proportion of PCR positives for *P. phoxinus* ("Pho pho") and
salmonid species ("Onc myk" = *Oncorhynchus mykiss*, "Sal fon" = *Salvelinus fontinalis*, "Sal tru"
= *Salmo trutta*) per trial; dark shading codes for successful detection; the number of positives

and total reactions are shown between pie charts and boxplots. The RFUs of positive samples

are displayed as boxplots; white circles code for mean values.

Figure 1

Fish cages

Maximum fish-free distance (1.3 km)

August   September   November

Water flow

Legend
- *Salvelinus fontinalis*
- *Oncorhynchus mykiss*
- *Salmo trutta*
- *Phoxinus phoxinus*
- Sampling point

Figure 2

Figure 3

Figure 4