1

Non-Invasive Single-Cell Morphometry in Living Bacterial Biofilms

Mingxing Zhang[1,§], Ji Zhang[1,§], Yibo Wang[1,§], Jie Wang[3], Alecia M. Achimovich[2],

Scott T. Acton[3], and Andreas Gahlmann[1,2,*]

[1]Department of Chemistry, University of Virginia, Charlottesville, Virginia, USA

[2]Department of Molecular Physiology & Biological Physics, University of Virginia School of

Medicine, Charlottesville, Virginia, USA

[3]Department of Electrical & Computer Engineering, University of Virginia, Charlottesville,

Virginia, USA

§These authors contribute equally to this paper

*Corresponding Author

Abstract

Fluorescence microscopy enables spatial and temporal measurements of live cells and cellular communities. However, this potential has not yet been fully realized for investigations of individual cell behaviors and phenotypic changes in dense, three-dimensional (3D) bacterial biofilms. Accurate cell detection and cellular shape measurement in densely packed biofilms are challenging because of the limited resolution and low signal to background ratio in fluorescence microscopy images. In this work, we present Bacterial Cell Morphometry 3D (*BCM3D*), an image analysis workflow that combines deep learning with mathematical image analysis to accurately segment and classify single bacterial cells in 3D fluorescence images. In *BCM3D*, deep convolutional neural networks (CNNs) are trained using simulated biofilm images with experimentally realistic signal-to-background ratios (SBRs), cell densities, labelling methods, and cell shapes. We systematically evaluated the segmentation accuracy of *BCM3D* using both simulated and experimental images. Compared to state-of-the-art bacterial cell segmentation approaches, *BCM3D* consistently achieves higher segmentation accuracy and further enables automated morphometric cell classifications in multi-population biofilms.

Introduction

Biofilms are multicellular communities of microorganisms that grow on biotic or abiotic surfaces[1-4]. In addition to cellular biomass, biofilms also contain an extracellular matrix (ECM) which is composed of polysaccharides, DNA, and proteins. Individual cells in biofilms interact with other cells, the ECM, or with the substrate surface, and the sum total of these interactions provide bacterial biofilms with emergent functional capabilities beyond those of individual cells. For example, biofilms are orders of magnitude more tolerant towards physical, chemical, and biological stressors, including antibiotic treatments and immune system clearance[1,2,5-8]. Understanding how such capabilities emerge from the coordination of individual cell behavior requires imaging technologies capable of resolving and simultaneous tracking of individual bacterial cells in 3D biofilms.

Live cell-compatible imaging technologies, such as light microscopy, can reveal the spatial and temporal context that affects cellular behaviors. However, conventional imaging modalities are not able to resolve individual cells within thick 3D biofilms over extended periods of time. For example, the diffraction-limited lateral x,y-resolution (~230 nm) of a confocal fluorescence microscope is barely sufficient to resolve bacterial cells positioned next to each other on flat glass coverslips. Even worse, the diffraction-limited axial z-resolution (570 nm) is comparable to the size of a single bacterial cell, so that densely-packed cells become unresolvable in the axial z-dimension[9,10]. Notable exceptions include loose biofilms (low cell density), spherical cell shapes[11,12], and mutant *Vibrio cholera* biofilms, in which cell-cell spacing is increased through the overproduction of ECM materials[13-15]. While single-cell resolved images have been obtained in such special situations, conventional optical microscopy modalities are not generally capable to accurately resolve and quantitatively track individual cells in dense 3D biofilms.

Super-resolution derivatives of confocal microscopy, known as Image Scanning

Microscopy[16], can improve spatial resolution. However, a perhaps more important limitation for

long-term live-cell imaging is photodamage to the specimen (phototoxicity) and to the

fluorophores used for labeling (photobleaching)[17-19]. In confocal microscopy-based approaches,

undesired out-of-focus fluorescence emission is filtered out by confocal pinholes to yield optically-

sectioned images with high contrast, i.e. high signal-to-background ratios (SBRs). However,

repeated illumination of out-of-focus regions during laser scanning and high light intensities at the

focal volume result in rapid photobleaching of fluorophores and unacceptable phototoxicity for

light sensitive specimens[17-19]. In fact, confocal fluorescence microscopy (as well as its super-

resolution derivatives) uses illumination light intensities that are two to three orders of magnitude

higher than the light intensities under which life has evolved[18]. The high rates of phototoxicity and

photobleaching make confocal-based microscopy unsuitable for high frame-rate time-lapse

imaging of living specimens over many hours and days[14,15,17,20,21].

In recent years, light sheet-based fluorescence excitation and imaging approaches have

been developed to overcome the drawbacks of confocal microscopy. Among these, lattice light

sheet microscopy (LLSM)[18,19] and field synthesis variants thereof[22], axially-swept light sheet

microscopy (ASLM)[23,24], and single-objective oblique plane light sheet microscopes[25-30] now

combine excellent 3D spatial resolution with fast temporal resolution and low phototoxicity at

levels that cannot be matched by confocal microscopy. Specifically, light sheet-based microscopy

approaches can operate at illumination intensities that are below the levels of cellular

phototoxicity, even for notoriously light sensitive specimens, and reduce fluorophore

photobleaching by 20-50 times compared to confocal microscopy, while maintaining comparable

spatial resolution and contrast/SBR[18,26].

An additional challenge in high-resolution biofilm imaging is data quantification. Even if

sufficient resolution and high SBRs can be achieved to visually discern, i.e. qualitatively resolve

individual cells, robust computational algorithms are still needed for automated cell segmentation

and quantitative cell tracking. Towards this goal, image processing approaches based on the

watershed technique and intensity thresholding have been developed over the years for single-cell

segmentation in bacterial biofilms[13-15,21]. The broad applicability of watershed- and threshold-

based image processing algorithms is limited, because these algorithms require manual

optimization of many user-selected parameters. Even with optimal parameters, watershed- and

threshold-based image processing methods produce sub-optimal segmentation results, especially

when cell densities are high, when SBRs are low, and when cellular fluorescence intensities are

not uniform across the cytosol or on the cell surface. To overcome the drawbacks of watershed-

and threshold-based image processing approaches, deep learning by CNNs have been used in

recent years with outstanding success for a wide range of problems in biomedical image analysis[31].

Here, we present Bacterial Cell Morphometry 3D (*BCM3D*)[32], a generally applicable

workflow for single-cell segmentation and shape determination in high-resolution 3D images of

bacterial biofilms. *BCM3D* uses CNNs in combination with mathematical image analysis to

achieve accurate single cell segmentation in 3D. The CNNs employed in *BCM3D* are based on the

3D U-Net architecture and training strategy, which has achieved excellent performance in

biomedical data analysis benchmark tests[31]. The mathematical image analysis modules of *BCM3D*

enable post-processing of the CNN results to further improve the segmentation accuracy. We

establish that experimental bacterial biofilms images, acquired by lattice light sheet microscopy,

can be successfully segmented using CNNs trained with computationally simulated biofilm

images, for which the ground-truth voxel-level annotation maps are known accurately and

precisely. By systematically evaluating the performance of *BCM3D* for a range of SBRs, cell

densities, and cell shapes, we find that voxel-level segmentation accuracies of >80%, as well as

cell counting accuracies of >90%, can be robustly achieved. *BCM3D* consistently outperforms

previously reported image segmentation approaches that rely exclusively on conventional image

processing approaches. We expect that *BCM3D*, and CNN-based single-cell segmentation

approaches in general, combined with non-invasive light sheet-based fluorescence microscopy

will enable accurate cell tracking over time in dense 3D biofilms. This capability will launch a

new era for bacterial biofilm research, in which the emergent properties of microbial populations

can be studied in terms of the fully-resolved behavioral phenotypes of individual cells.

Materials and Methods

Lattice Light Sheet Imaging of Bacterial Biofilms

Fluorescence images of bacterial biofilms were acquired on a home-built lattice light sheet

microscope (LLSM). LLSM enables specimen illumination with a thin light sheet derived from

2D optical lattice[33,34]. Here, a continuous illumination light sheet was produced by a time-averaged

(dithered), square lattice pattern[18], and the illumination intensity at the sample was <1 W/cm^2. The

submicrometer thickness of the excitation light sheet is maintained over long propagation distances

(~30 μm), which enables optical sectioning, and thus high resolution, high contrast imaging of 3D

specimens comparable to confocal microscopy. However, fluorophore excitation by a 2D light

sheet reduces phototoxicity, because each excitation photon has multiple opportunities to be

absorbed by fluorophores in the excitation plane and produce in-focus fluorescence. Widefield

fluorescence images corresponding to each illuminated specimen plane are recorded on a sCMOS

detector (Hamamatsu ORCA Flash v2). In this work, 3D biofilm images were acquired by

translating the specimen through the light sheet in 200 nm steps using a piezo nanopositioning

stage (Physik Instrumente, P-621.1CD).

E.coli K12 Biofilm Imaging: Ampicillin resistant *E.coli* K12, constitutively expressing

GFP[35], were cultured at 37 degrees overnight in LB medium with 100 μg/ml ampicillin. Overnight

cultures were diluted 100 times into the same culture medium, grown to an optical density at 600

nm (OD600) of 0.6 – 1.0, and then diluted by an additional factor of 10. Round glass coverslips

with the diameter of 5 mm were put into a 24-well plate (Falcon) and 400 μL of cell culture was

added to the well. Cells were allowed to settle to the bottom of the well and adhere to the coverslip

for 1 hour. The round coverslips were then mounted onto a sample holder and placed into the

LLSM sample-basin filled with M9 medium. GFP fluorescence was excited using 488 nm light

sheet excitation. Biofilm growth was imaged at room temperature every 30 min for a total of 20

time points. At each time point, a single 3D image stack contained 400 images, each acquired with

a 15 ms exposure time to avoid motion blur.

M. xanthus Biofilm Imaging: Strain LS3908 expressing tdTomato under the control of the

IPTG-inducible promoter[36] and DK1622 (WT) were cultured in the nutrient rich CYE media at 30

degrees Celsius until it reached an OD600 of 0.6 - 1.0. Media was supplemented with 1 mM IPTG

for tdTomato expressing cells. Chitosan (Thermo Fisher)-coated 5 mm round glass coverslips were

prepared by incubating coverslips with 1% (w/v) chitosan (1.5 % glacial acetic acid (v/v)) at room

temperature for 1 hour. Coverslips were then rinsed with water and placed into a 24-well plate

(Falcon) with 350-400 μL of undiluted cell culture. WT cells were stained directly in the 24 well

plate with 5 ng/ml FM4-64 (Thermo Fisher) dye. Cells were allowed to settle and adhere to the

coverslip for 2 hours. After the settling period, the coverslip was gently rinsed with CYE media to

flush away unattached cells. The rinsed coverslip was then mounted onto a sample holder and

placed into the LLSM sample-basin filled with MC7 starvation buffer. tdTomato and FM 4-64

fluorescence was excited using 561 nm light sheet. The 3D image stack contained 400 2D images.

Each 2D slice was acquired with an exposure time of 30 ms.

<u>Mixed Strain Biofilm Imaging</u>: Ampicillin resistant *E.coli* K12, constitutively expressing

GFP[35], and ampicillin resistant *E.coli* K12, expressing mScarlet (pBAD vector, arabinose induce)

were cultured separately at 37 degrees overnight in LB medium with 100 μg/ml ampicillin.

Overnight cultures were diluted 100 times into the same culture medium, grown to an optical

density at 600 nm (OD600) of 0.6 – 1.0, and then diluted to an OD of 0.1. After dilution, the two

strains were mixed together. Round glass coverslips with the diameter of 5 mm were put into a 24-

well plate (Falcon) and 500 μL of cell culture was added to the well. Cells were allowed to settle

to the bottom of the well and adhere to the coverslip for 1 hour. The cell culture medium was then

removed and replaced by 500 uL M9 medium containing 0.2% (w/v) arabinose. The co-culture

was incubated at 30 degrees overnight. 10 mins before imaging, the co-culture was stained with

5 ng/ml FM4-64 (Thermo Fisher) dye. 3D image stacks of 20 planes with 5 ms exposure time per

frame were acquired using 488 nm excitation.

Raw Data Processing

Raw 3D images were background subtracted and then deskewed and deconvolved as

described previously[18,19]. The background was estimated by averaging intensity values of dark

areas (devoid of cells) in the field of view. Deconvolution was performed using the Richardson-

Lucy algorithm with 10 iterations using experimentally measured point spread functions (PSFs) as

the deconvolution kernel. The experimentally measured PSFs were obtained separately for each

color channel using fluorescent beads (200 nm FluoSpheres®, Thermo Fisher) coated on a

coverslip[37]. 3D images were rendered using the 3D Viewer plugin in Fiji[38] or ChimeraX[39].

Generation of simulated biofilm images

176   To generate data for training of CNNs, we computationally simulated fluorescence images

of 3D biofilms, for which spatial arrangements among individual cells are known precisely and

accurately. Growth and division of individual rod-shaped cells in a population were simulated

using CellModeller, an individual-based computational model of biofilm growth (**Figure 1a**)[40]. In

individual-based biofilm growth models, cells are the basic modeling units. Each cell is

characterized by a set of parameters, including its 3D position, volume, and spatial orientation. All

the cells in the simulated biofilm are then allowed to evolve in time according to predefined

biological, chemical, and mechanical rules. For example, cells grow at a defined rate and then

divide after reaching a certain volume threshold. Cellular collisions that are due to cell growth are

alleviated by imposing a minimum distance criterion between cells at each time point. For our

simulations, we chose cell diameter and cell length ($d,\ l$) parameters consistent with a given

bacterial species, namely (1 μm, 3 μm) for *E. coli*[41], (0.7 μm, 6 μm) for *M. xanthus*[42], and (1 μm,

1 μm) for spherically symmetric *S.aureus*[43].

189   While the cell volume can be readily adjusted in CellModeller, the cellular volume density,

which is determined by the intercellular spacing, is not directly adjustable. We therefore adjusted

the cellular volume density after each simulation by scaling the cellular positions (cell centroids)

and thus the intercellular distances by a constant factor, while leaving cell sizes, shapes, and

orientations unchanged. This post-processing procedure enabled simulation of the exact same 3D

cell arrangements at adjustable cell volume densities.

We fluorescently labeled simulated cell volumes and surfaces according to two commonly

used labeling strategies in fluorescence microscopy. To simulate expression of intracellular

fluorescent proteins, the fluorescence emitters were placed at random positions within the cell

volume. To simulate membrane staining, the fluorescence emitters were placed at random

positions on the cell surface. Each cell contained between 500 - 1000 fluorophores to simulate

expression level variations between cells, which is often observed in experimental images. Once

the fluorophore spatial distributions were determined, a 3D fluorescence image (**Figure 1b**) was

computationally generated. Each fluorophore was treated as an isotropic point emitter, so that it

would produce a diffraction-limited point-spread-function (PSF) on the detector. Experimentally

measured 3D PSF shapes (see Raw Data Processing) were used as the convolution kernel. Next,

the fluorescence signal intensity was scaled by multiplying the image by a constant factor and then

a constant background intensity was added to the image at ~200 photons per pixel, as measured in

experimental data. This procedure enabled independent adjustments of the fluorescence signal and

background to obtain signal-to-background ratios (SBRs) consistent with experimental data. In a

final step, we introduced Poisson-distributed counting noise, based on the summed background

and signal intensities, as well as Gaussian-distributed camera read-out noise (experimentally

calibrated for our detector at 3.04 photons per pixel on average)[44]. This resulting image data

(**Figure 1c**) was then processed in the same manner as experimental data (see Raw Data

Processing). In contrast to experimental data, generation of the corresponding voxel-level

annotation maps is fast and error free, because the underlying ground truth cell arrangements are

known *a priori* (**Figure 1d**).

Differential labeling: To mimic imaging of reporter gene expression in a subset of cells,

we simulated biofilm images, in which all cells were stained at the cell surface (e.g. with a

membrane intercalating fluorescent dye) and a subset of cells additionally contained intracellular

fluorophores (e.g. through the expression of an intracellular fluorescent protein) (**Figure S1ab**).

The mixing ratios between membrane-labelled, and membrane + interior labelled cells were 10:90,

30:70, 50:50, 70:30 and 90:10. Ten different cell arrangements containing ~300 cells were

simulated for each ratio. To train the CNNs (see next section), five datasets were used, all with a

50:50 mixing ratio.

Mixed cell shapes: To mimic imaging of cells with different morphologies, we simulated

biofilms containing spherical and rod-shaped cells **(Figure S1cd)**. The spatial arrangements of the

mixed biofilm were generated by randomly positioning cells in a 3D volume. Collision among

cells was eliminated by adjusting cell positions and orientations. The size of the rod-shaped cells

is that of *E. coli* (~3 × 1 μm, length by diameter). The size of the spherical cells is that of *S. aureus*

(~1 μm in diameter)[45]. Both cell types were labelled by intracellular fluorophores, as described

above. The mixing ratios between rod-shaped and spherical cells were 10:90, 30:70, 50:50, 70:30

and 90:10. Ten different cell arrangements containing ~300 cells were simulated for each ratio. To

train the CNNs (see next section), we picked one image from each mixing ratio for a total of five

images.

Training the convolutional neural networks

We trained 3D U-Net CNNs for voxel-level classification tasks[46] within the NiftyNet platform[47]

(network architecture depth 4, convolution kernel size 3, ReLU activation function, 32 initial

feature maps, and random dropout of 0.5 during training). To achieve robust performance, we

trained these networks using five to ten simulated biofilm images with randomly selected cell

densities and signal-to-background ratios (see Generation of simulated biofilm images). The raw

data processing steps used for experimental data (see Raw Data Processing) were equally applied

to simulated data. 3D deconvolved simulated data and their corresponding voxel-level annotations

were used to train the CNNs. Each image used for training contained ~9 million voxels. We trained

CNNs by classifying each voxel as 'background', 'cellular interior' or as 'cellular boundary' based

on the underlying cell arrangements. For mixed-species biofilms, two additional classes, 'cellular

interior' and 'cellular boundary' of the second species, were used. This type of annotation scheme

has been shown to increase separation of bacterial cells in 2D[48]. For data augmentation, we applied

NiftyNet's built-in scaling, rotation, and elastic deformation functions. Instead of the original

cross-entropy loss function combined with uniform sampling, we used the Dice loss function and

'balanced sampler', so that every label has the same probability of occurrence in training. All

networks were trained for 2000 to 3600 iterations with a learning rate of 0.0001. Using these

parameters, it took approximately 24 hours to train the CNNs on a NVIDIA Tesla V100 GPU with

16 GB memory.

Thresholding of CNN-produced confidence maps

Voxel-level classification by CNNs generates different confidence maps (one confidence

map for each annotation class). The confidence values range between 0 and 1 and represent the

confidence of assigning individual voxels to a given class. After thresholding the 'cell interior'

confidence map to obtain a binary image (**Figure S2a-c**), connected voxel clusters can be isolated

and identified as single cell objects using 3D connected component labeling[49]. A conservative size-

exclusion filter was applied: small objects with a volume ~10 times less than the expected cell size

were considered background noise and filtered out using an area open operator[49]. Since the cell-

interior volumes do not contain the cell boundaries, we dilated each object by 1-2 voxels to

increase the cell volumes using standard morphological dilation[49]. The threshold value to segment

individual cell objects based on the 'cell interior' confidence map was determined by plotting the

overall voxel-level segmentation accuracy (see Performance Evaluation) versus the threshold

values (**Figure S2**). High segmentation accuracies were consistently obtained using confidence

thresholds between 0.88 and 0.94.

Post-processing of U-Net result using a refined LCuts algorithm

Thresholding of the 'cell interior' confidence map produces a binary segmentation result

(background = 0, cell interior =1), where groups of connected, non-zero voxels identify individual

cells in most cases (**Figure S2**). However, when cells are touching, they are often not segmented

as individuals, but remain part of the same voxel cluster (undersegmentation). On the other hand,

a single cell may be erroneously split into smaller sub-cellular objects (oversegmentation). Finally,

in datasets with low SBR, connected voxel clusters may be detected that do not correspond to cells

and thus produce false positive objects (**Figure 2a**). To address these errors and improve the

segmentation accuracy further, we included additional mathematical image analysis steps to post-

process the CNN results and reduce undersegmentation and oversegmentation errors.

<u>Step 1. Filtering:</u> False positive objects are identified by evaluating the coefficient of

variation[50,51] for each connected voxel cluster i:

$$CV_i = \frac{\sigma_i}{\mu_i}$$

where σ_i and μ_i denote the standard deviation and the mean of the intensity taken over all voxels

contained in connected voxel cluster i. If the coefficient of variation is larger than ρ, then the

current object will be classified as a false positive object and removed from the confidence map

by setting all its voxels to zero. The removed objects will then no longer be counted when

evaluating the cell counting accuracy. The value of ρ is selected based on the coefficient of

variation of the background. For the datasets analyzed here, this sample coefficient of variation

was determined to be $\rho = 1.1$. After CV-filtering, objects smaller than 25% of the expected

bacterial cell size are also removed by setting its voxels to zero. The remaining connected voxel

clusters are then considered for further processing (**Figure 2a**).

Step 2. Identification of point coordinates along central bacterial cell axes: To identify and

delineate individual cells in the connected voxel clusters identified in the previous step, we

implemented medial axis extraction using the method of inscribed spheres[52], with the constraint

that the sphere radii do not exceed the expected diameter of a single bacterial cell (e.g. $d = 0.8$ μm)

(**Figure 2b** left). The set of N inscribed spheres are tangent to the object's surface and

parameterized by $(x_i, y_i, z_i; r_i < d)$ for $i = 1, \ldots, N$. Determination of the $(x_i, y_i, z_i; r_i)$ coordinates is

achieved using the Euclidean distance transform of the objects' boundary[53], so that the points with

coordinates (x_i, y_i, z_i) reliably trace out the central cell axes of individual bacterial cells (**Figure 2b**

right).

Step 3. Identification of individual cell axes: To separate different linear segments after

cell axis extraction (**Figure 2c**), we used a refined version of the linear cuts (*LCuts*) algorithm[54,55].

LCuts is a graph-based data clustering method designed to detect linearly oriented groups of points

with certain properties. The fundamental elements of a weighted mathematical graph are nodes,

edges, and edge weights. Here, the points with coordinates (x_i, y_i, z_i) represent the graph nodes.

Edges are the connections among nodes. Edges are assigned weights, for example, to reflect the

confidence that two nodes belong to the same group. *LCuts* achieves grouping by assigning

weights to edges in the fully connected graph to reflect the similarity between two nodes. The

features of each node include its location and direction, where the location of each node is simply

its Cartesian coordinates. The direction of each node is found by first determining its 5-hop

neighborhood, removing nodes at large relative angles, and evaluating the major direction of the

outlier removed neighborhood (**Figure S3**).

The algorithm to separate the nodes into different groups is a recursive graph cutting

method[54]. Graph cuts (*e.g.* nCut[56]) disconnect the edges between two groups of nodes when the

combined weights of these edges are minimized. The weights, between node i and node j, are

calculated as follows:

317
$$w_{ij} = w_D \cdot w_T$$

318   where

319
$$w_D = \begin{cases} e^{-D_{ij}^2/\sigma_D^2} & \text{if } D_{ij}^2 \leq r \\ 0 & \text{if } D_{ij}^2 > r \end{cases}$$

320
$$w_T = e^{-(\cos(\theta_{ij})-1)^2/\sigma_T^2}$$

w_D weighs the distance between two nodes and w_T weighs difference between node directions.

D_{ij} is the Euclidean distance between node i and node j, and r is set to eliminate edges between

two far away nodes. θ_{ij} is the relative angle between the directions of nodes i and j. σ_D and σ_T are

adjustable parameters that control the rate of exponential decay. *LCuts* continues to separate groups

of nodes until each group satisfies a stopping criterion. The stopping criterion is biologically

inspired based on the expected length of a single cell and a group's linearity after each recursion.

LCuts yields linearly oriented groups of points that trace out the central axes of individual cells

(**Figure 2c**). Importantly, cell separation is achieved without having to specify the number of cells

in the biofilm in advance. With this step, the clustering provided by the *LCuts* algorithm can fix

under- and over-segmentation problems from the CNN-based segmentation.

<u>Step 4. Cell reconstruction:</u> The final output of linear clustering can provide length,

location and orientation of each cell. Based on these linear clusters, the cellular architecture of the

biofilms can be reconstructed by placing geometrical models of cells in space as shown in **Figure**

2d. For fast computation, spherocylinders are used as the geometrical model using a radius

consistent with the known sizes of bacterial cells. To further refine the cell surfaces to better align

with the CNN-segmented volumes, we enclosed the inscribed spheres found in Step 2 in a convex

hull (**Figure 2d**).

Performance Evaluation

We quantified segmentation accuracy both at the cell-level (object counting) and at the

voxel-level (cell shape estimation). To quantify the cell-level segmentation accuracy, we

designated segmented objects as true positive (*TP*) if their voxel overlap with the ground truth or

the manual annotation resulted in a Jaccard index[57] larger than 0.5. This criterion ensures one-to-

one matching. If the segmented cell object could not be matched to a ground truth/manually

annotated cell volume, then it was counted as a false positive (*FP*) and the Jaccard index of that

segmented object was set to zero. If a ground truth/manually annotated cell volume was not

identified in the image, then it was counted as false negative (*FN*). The cell counting accuracy was

defined as *TP*/(*TP+FP+FN*). The voxel-level segmentation accuracy quantifies the accuracy of

cell shape estimation and is calculated as the average Jaccard index over all segmented objects in

the image.

To evaluate the accuracy of cell segmentation on experimental data, three researchers

separately traced the cell contours on experimental 2D slices by using freehand selections in Fiji

ROI Manger[58]. Because human annotation is very time consuming (about 50 hours for a complete

3D dataset containing ~300 cells in a 22 x 32 x 12 um^3 volume), one to three single 2D slices were

selected for each dataset. One exception is the 3D *M. xanthus*, for which the cell outlines in all

available x, y and z slices were traced manually. For straight, rod-shaped cells, the centroids of the

resulting 2D cell contours all fall within the cell interior volume. To group together the contours

belonging to the same cells, the centroid of each contour was projected along the x, y and z

dimension. If the projected centroid was enclosed by any other contour in a different slice, then

the centroid of that contour was projected onto the plane of the initial contour. Two contours were

labeled as related if they contained each other's projected centroids. This process is repeated for

all possible contour pairs and their relationship is recorded in an adjacency matrix. Next, related

contours were assigned to individual cells. To separate incorrectly grouped contours, we

additionally identified clusters of centroids using the DBSCAN point clustering algorithm[59]. In a

final step, we manually removed incorrectly traced contours. This procedure determined the

approximate positions, shapes, and orientations of individual cells in the 3D biofilm.

To estimate the SBRs of both simulated and experimental images, we manually selected

and determined the intensities of approximately ten 'signal' and ten 'background' regions in the

images. We computed the SBR as the mean signal intensity divided by the mean background

intensity. To estimate the local density of a biofilm cluster of simulated images, we sliced the

considered several 3D tiles within the biofilm of size 64 by 64 by 8 voxels. We then estimated the

local density as total cell volume contained in each tile divided by the tile volume. We calculated

the mean density of the 10 densest tiles to define the 'local density' metric reported for each dataset

in the paper. To estimate the cell density in an experimentally acquired biofilm image, the same

calculations were performed on either 3D manual annotations (if available) or binary masks

obtained by CNN-processing.

Results and Discussion

Cell segmentation by thresholding CNN confidence maps

CNNs have been shown to perform well on pixel-level classification tasks for both 2D and 3D data[60,61]. Bacterial biofilms, however, present a unique challenge in this context. The cell shapes to be segmented are densely packed and barely resolvable even with the highest resolution optical microscopes. Additionally, living biofilms in fluorescence microscopes can only be excited with low laser intensities to ameliorate phototoxicity and photobleaching concerns. Low intensity excitation unfortunately also reduces the SBR in the acquired images. It remains unclear to what extent single-cell segmentation approaches can accurately identify and delineate cell shapes in bacterial biofilm images obtained under low intensity illumination conditions. To address this question, we implemented an *in silico* CNN training strategy for *BCM3D* (see Materials and Methods) and systematically evaluated its voxel-level classification (cell morphometry) and cell counting accuracies using simulated biofilm images with varying cell densities and SBRs similar to those encountered in experimental data.

We compared two commonly used cell labeling approaches, namely genetic labeling through the expression of cell-internal fluorescent proteins (**Figure 3a-c**) and staining of the cell membranes using fluorescent dyes (**Figure 3d-f**). For both labeling approaches, voxel-level segmentation and cell counting accuracies, obtained by thresholding CNN confidence maps (see Materials and Methods), depend mostly on cell density, whereas the SBR plays a less important role. For cell-internal labeling, SBRs of >1.7 and cell densities of <60% consistently produce voxel-level classification accuracies of >80% and cell counting accuracies of >95%. On the other hand, SBRs of <1.7 and cell densities of >60% lead to lower segmentation accuracies. While lower segmentation accuracies are expected for higher cell densities and lower SBRs, the sharp drop-

offs observed here may indicate a fundamental performance limitation of the CNNs employed.

Still, the voxel-level classification and cell counting accuracies consistently surpass previous

approaches for bacterial cell segmentation for commonly encountered cell densities and SBRs.

Specifically, the cell counting accuracies obtained by *BiofilmQ*[62], *Seg3D*[63], and Yan *et al.*[13] quickly

drop to zero as a function of increasing the Jaccard index threshold (a quantitative measure of cell

shape similarity relative to the ground truth, see Materials and Methods), indicating that cell shapes

are not accurately estimated by conventional image processing approaches (**Figure 3g-i**). By

contrast, the cell counting accuracy obtained by thresholding CNN confidence maps remain high

even for Jaccard index thresholds larger than 0.5, indicating that cell shapes are better estimated

by the CNNs.

The accuracies of single-cell shape estimation and cell counting are predominantly affected

by cell density. The variation is more pronounced for membrane-stained cells, because *inter-*

cellular fluorescence intensity minima are less pronounced when cell membranes are labeled and

cells physically contact each other (red arrow in **Figures 3c** and **3f**). By contrast, intracellular

fluorophores produce the highest intensities at the cell center, so that the gaps between cells are

more readily resolvable. Also noteworthy is the sharp drop-off in segmentation accuracies for

SBRs of <1.7 for all cases. In such low SBR regimes, fluorescence signals of the cells become too

difficult to be distinguished from the background. As a result, the CNNs falsely identify random

noisy patterns in the background as cells. Additionally, thresholding of the CNN confidence maps

often yields connected voxel clusters that contain multiple bacterial cells. False identification and

incomplete delineation of cells cause the pronounced decrease in segmentation accuracy for SBRs

of <1.7.

Post-processing of CNN confidence maps

To better identify individual cells in low SBR and high cell density datasets, we developed a graph-based post-processing module (see Materials and Methods) that takes advantage of highly conserved bacterial cell shape of a given species. Briefly, we transformed the CNN 'cell interior' confidence maps into 3D point cloud data that trace out the central axes of individual cells. This transformation was achieved by medial axis extraction using size-constrained inscribed spheres[52] (**Figure 2**). Single-cell axes are then identified as linearly clustered data points by *LCuts* – a graph-based data clustering method designed to detect linearly oriented groups of points[54]. The so-identified single-cell axes are then mapped back onto the original segmentation volumes to obtain estimates of the 3D positions, shapes, and orientations of the now separated cells.

Post-processing with *LCuts* takes advantage of *a priori* knowledge about expected bacterial cell sizes (**Figure S4**) by removing erroneously segmented volumes that are significantly smaller than the expected value and by splitting incompletely segmented volumes representing fused cells. Improvements of up to 15% and 36% are observed for cells labeled with cytosolic fluorophores (**Figure 4a-c**) and membrane-localized fluorophores (**Figure 4d-f**), respectively. The more substantial improvement for membrane-stained cells is due to fact that CNNs trained on membrane-stained cells are more prone to erroneously identifying speckled background noise as fluorescence signals in low SBR images. In addition, membrane-intercalating fluorophores of two adjacent cells are in close proximity, making it difficult to resolve fluorescence signals from two separate cells due to spatial signal overlap (see the red arrow, **Figure 3c** and **3f**).

Segmentation of experimental biofilm images

To test the performance of *BCM3D* on experimentally acquired biofilm images, we acquired time-lapse images of GFP expressing *E. coli* biofilms every thirty minutes for ten hours. We then manually annotated one 2D slice in the 3D images at the t = 5, 6, and 10-hour time points. When referenced to these manual segmentation results, both the CNN outputs and the *LCuts*-processed CNN outputs consistently achieved better cell counting accuracies than conventional segmentation methods (**Figure 5, Figure S5**). We note however that the cell counting accuracy is much lower (45% at a 0.5 Jaccard index threshold) when referencing manual annotation. The more rapid drop-off of the cell counting accuracy as a function of increasing Jaccard index threshold can be attributed to the following factors. First, human annotation of experimentally acquired biofilm images is not as consistent as simulated data, and therefore, it is less likely to produce good matches between segmented and annotated cell shapes. Second, the experimentally acquired biofilm images differ from the simulated training and annotation datasets in several ways. Real cells may appear blurred due to partially immobilized/wiggling cells. Some cells are in the process of cell division, so that different human annotators may count these cells either as one cell or two cells. Finally, optical aberrations and scattering effects were not included in training data simulations. We note, however that, at a lower Jaccard index threshold, the cell counting accuracies is in the range of 60% to 70%, which is substantially higher than the current state-of-the-art in biofilm imaging (**Figure 5**).

To demonstrate that *BCM3D* can achieve similarly high segmentation accuracies for membrane-stained cells, we analyzed a small patch of a *M. xanthus* biofilm, which was stained with the membrane intercalating dye FM4-64 (**Figure 6a**). For this dataset, we manually annotated each *xy*, *xz*, and *yz* slice through a single 3D image stack (**Figure 6b**). When referenced to this 3D

466 manual segmentation result, *BCM3D* (**Figure 6c**) again produced cell counting accuracies in the

467 range of 60% to 70% at low Jaccard index thresholds (**Figure 6d**). We note that comparison with

468 the state-of-the-art cannot be made here, because previous biofilm segmentation approaches were

469 not designed to work with membrane-stained cells.

Morphological separation of mixed cell populations

Given the much-improved segmentation results obtained using *BCM3D*, we reasoned that

the same CNNs may also be able to assign segmented objects to different cell types based on subtle

morphological differences in the acquired images. Differences in the imaged cell morphologies

arise due to physical differences in cell shapes (e.g. spherical vs. rod-shaped cells) or due to

differences in the fluorescent labeling protocols (e.g. intracellular vs. cell membrane labeling),

because fluorescence microscopes simply measure the spatial distributions of fluorophores in the

sample. The ability to separate different cell morphologies is important for the study of

multispecies biofilms, where interspecies cooperation and competition dictate population-level

outcomes[3,64-71]. Separation of differentially labeled cells is also important for the study of gene

activation in response to cell-to-cell signaling[72]. Expression of cytosolic fluorescent proteins by

transcriptional reporter strains is a widely-used technique to visualize activation of a specific gene

or genetic pathway in living cells. Such genetic labeling approaches can be complemented by

chemical labeling approaches, e.g. using membrane intercalating chemical dyes that help visualize

cells non-specifically or environmentally-sensitive membrane dyes that provide physiological

information, including membrane composition[73,74], membrane organization/integrity[75-77], and

membrane potential[78,79]. Chemical and genetic labeling approaches are traditionally implemented

in two different color channels. However, there are important drawbacks to using multiple colors.

First and foremost, the amount of excitation light delivered is increased to excite differently

colored fluorophores, raising phototoxicity and photobleaching concerns. Second, it takes N times

as along to acquire N-color images (unless different color channels can be acquired

simultaneously), making it challenging to achieve high temporal sampling in time-lapse

acquisition. For these reasons, methods that extract complementary physiological information

from a single-color image are preferable.

We evaluated the ability of *BCM3D* to automatically segment and separate rod-shaped and

spherical bacterial cells consistent with sizes of *E. coli* and *S. aureus*, respectively, in simulated

images (**Figure S6**). Similar to a single-population segmentation problem, we trained CNNs that

classify pixels into five different classes: 'background', 'cellular interior of population 1', 'cellular

boundary of population 1', 'cellular interior of population 2' and 'cellular boundary of population

2'. Thresholding the CNNs confidence maps can achieve cell counting accuracies larger than 90%

for both cell types, independent of their population fractions (**Figure 7a**). Post-processing of this

result using *LCuts* improved the cell counting accuracy by less than 0.5% on average, indicating

that under-segmented cell clusters are not prevalent in this dataset.

We next evaluated the ability of *BCM3D* to automatically segment and separate membrane-

stained cells that express cytosolic fluorescent proteins from those that do not (**Figure S7**). Again,

the cell counting accuracy is consistently above 80% for all tested mixing ratios (**Figure 7b**).

Finally, we applied *BCM3D* to experimentally acquired biofilm images of two different *E. coli*

strains. Both strains were stained by the membrane intercalating dye FM4-64, but the second strain

additionally expressed GFP (**Figure S8**). The cells were homogeneously mixed prior to mounting

to randomize the spatial distribution of different cell types in the biofilm (see Materials and

Methods). Multiple 2D slices from the 3D image stack were manually annotated and compared

with the results obtained by *BCM3D*. Consistent with the single-species experimental data, a cell

counting accuracy of 50% is achieved for each cell type at a 0.5 Jaccard index threshold and, at a

lower Jaccard index thresholds, the counting accuracies increased to 60% to 70%, (**Figure 7cd**).

Thus, *BCM3D* enables automated and accurate cell type assignments based on subtle differences

in cell morphologies in mixed population biofilms – a capability not available using conventional

image processing methods.

Conclusions

CNNs have been successful applied to many different problems in biological image

analysis, but their ability to segment individual cells in 3D and time-lapse 3D bacterial biofilm

images has not yet been fully explored. Here, we demonstrated a new CNN-based image analysis

workflow, termed *BCM3D*, for single-cell segmentation and shape classification (morphometry)

in 3D images of bacterial biofilms. We applied *BCM3D* to 3D images acquired by lattice light

sheet microscopy. However, *BCM3D* can also be applied to 3D images acquired by confocal

microscopy or more advanced super-resolution microscopy modalities, provided that realistic

image formation models are used to simulate the training datasets. The use of simulated training

data is a major advantage of *BCM3D*, because it overcomes inconsistencies inherent in manual

dataset annotation and provides needed flexibility not only in terms of the microscope platform

used, but also in terms of the bacterial cell shapes that are to be segmented.

We systematically investigated the advantages and limitations of *BCM3D* by evaluating its

voxel- and cell-level segmentation accuracy using simulated and experimental datasets of different

cell densities and SBRs. CNN-based image analysis enabled accurate segmentation of individual

cells in crowded environments and automatic assignments of individual cells to specific cell

populations for most of the tested parameter space. Such capabilities are not readily available when using previously established segmentation methods that rely exclusively on conventional image and signal processing algorithms.

While *BCM3D* surpasses the performance of previous approaches, we stress that further improvements are still possible and, for certain experiments, absolutely needed. Our systematic analysis revealed that high cell density and low SBR datasets are particularly challenging for the CNNs used in this work. Future work will therefore focus on increasing the contrast and resolution in bacterial biofilm images. The use of optical super-resolution modalities can certainly provide higher spatial resolution, but such resolution improvements often come at a cost of reduced image contrast and faster photobleaching/phototoxicity, especially for fast frame-rate time-lapse imaging. Software solutions will therefore play an increasingly important role in biological imaging. Indeed, CNN-based processing modules developed for contrast enhancement and image denoising have also surpassed the performance of conventional methods based on mathematical signal processing[80-83]. Incorporation of these tools into *BCM3D* may yield further improvements in the single-cell segmentation accuracies. We anticipate that the ability to accurately identify and delineate individual cells in dense 3D biofilms will in turn enable accurate cell tracking over long periods of time. Detailed measurements of behavioral single-cell phenotypes in larger bacterial communities will help determine how macroscopic biofilm properties, such as its mechanical cohesion/adhesion and its biochemical metabolism, emerge from the collective actions of *individual* bacteria.

Author Contributions

M.Z., J.Z., Y.W., J.W., S.T.A., and A. G. designed research;

M.Z., J.Z., Y.W., J.W., and A.M.A. performed research;

M.Z., J.Z., Y.W., J.W., S.T.A., and A. G. analyzed data;

and M.Z., J.Z., Y.W., J.W., S.T.A., and A. G. wrote the paper.

Acknowledgements

This work was supported by a Jeffress Memorial Trust Awards in Interdisciplinary Research (to A.G and S.T.A.). M.Z. and J.W. were supported by a University of Virginia Presidential Fellowship in Data Science. Y.W. was supported in part by a Dean's M.S.-Ph.D. Fellowship in Data Science. We thank Lotte Søgaard-Andersen, Larry Shimkets, Huiwang Ai and Ingmar Riedel-Kruse for providing the bacterial strains used in this work. We thank Karsten Siller, computational research consultant at the University of Virginia, for help with utilizing the Universities super-computing resources.

Figure 1.

Figure 1. Simulation of fluorescent biofilms images and annotation maps. **(a)** Cell arrangements

obtained by CellModeller. **(b)** Simulated 3D fluorescence image based on the cell arrangements

in **a**. **(c)** XY slice through the 3D simulated fluorescence image in **b** (upper panel shows cells

expressing cytosolic fluorescent proteins, lower panel shows cells stained with membrane-

intercalating dyes). **(d)** Ground truth cell arrangements giving rise to the image shown in **c**. Voxels

are displayed as black (background), or in different colors (indicating different cells).

Figure 2.

Figure 2. Post-processing of CNN-produced confidence maps using a refined *LCuts* processing

pipeline. (**a**) False positive objects are detected and removed by CV- and size- filtering. Under-

segmented clusters that are larger than single cells are selected for further splitting. (**b**) Illustration

of modified medial axis (red dashed lines) extraction to generate point cloud data from fused

clusters of rod-shaped cells using the method of inscribed spheres. When cells are touching, the

traditional medial axis extraction process fails to align with the actual cell central axis (left). To

overcome this drawback, we limited the size of the inscribed spheres based on prior knowledge of

bacterial cell diameters (right). **(c)** The set of inscribed sphere centers are then treated as a fully-

connected, undirected graph in 3D with two node features: location and direction (see text and

Figure S3 for details). The graph (blue nodes) is then iteratively cut into smaller graphs (red nodes)

until the stopping criteria are reached (see text for details). **(d)** Post-processed graphs represented

in different color denoting different cells. The 3D surface of individual cells can be determined

using a geometrical cell shape model (e.g. a spherocylinder for rod shaped bacteria) or by

calculating the convex hull around the inscribed spheres found in step 2.

Figure 3.

Figure 3. Performance of *BCM3D* using *in silico*-trained CNNs only on previously unseen simulated biofilm images. **(a)** The Jaccard index quantifies whether each voxel has been assigned to the correct class ('cell interior', cell surface', or 'background'). Solid circles represent the maximum local density and average SBRs encountered in experimental datasets (red, orange and blue: *E. coli* expressing GFP). **(b)** The cell counting accuracy using a Jaccard Index threshold of 0.5 for each segmented object (Materials and Methods) averaged over n=10 replicate datasets for cells labeled with cytosolic fluorophores. **(c)** Example image of cells labeled with cytosolic

fluorophores (Cell density = 60.0%, SBR = 1.34, indicated by white rectangle in panels a and b).

(d) Jaccard index and **(e)** Cell counting accuracy averaged over n=10 replicate datasets for cells

labeled with membrane-localized fluorophores. Purple solid circles represent experimental

datasets of *M. xanthus* stained with the membrane intercalating dye FM4-64. **(f)** Example image

of cells labeled with membrane-localized fluorophores (Cell density = 60.0%, SBR = 1.34,

indicated by white rectangles in panels d and e). The red arrows indicate a close cell-to-cell contact

point. **(g), (h) and (i)** Comparison of segmentation accuracies achieved by conventional (*BiofilmQ*,

Seg3D, and Yan *et al.*) and *BCM3D* using *in silico*-trained CNNs only. Three simulated datasets

(cytosolic fluorophores) with different SBRs and cell densities are shown. Segmentation accuracy

is parameterized in terms of cell counting accuracy (*y* axis) and accuracy of single-cell shape

estimation (*x* axis, quantified as the single-cell Jaccard index). Curves approaching the upper right-

hand corner indicate higher overall segmentation accuracy.

Figure 4.

Figure 4. Performance of *BCM3D* (*in silico*-trained CNNs *and* additional post-processing by *LCuts*) on previously unseen simulated data. **(a)** Jaccard index and **(b)** The cell counting accuracy using a Jaccard Index threshold of 0.5 for each segmented object averaged over n=10 replicate datasets for cells labeled with cytosolic fluorophores. **(c)** Improvement relative to *silico*-trained convolutional neural networks without post-processing. **(d)** Jaccard index and **(e)** Cell counting accuracy averaged over n=10 replicate datasets for cells labeled with membrane-localized fluorophores. **(f)** Improvements relative to *silico*-trained convolutional neural networks without post-processing.

Figure 5.

Figure 5. Comparison of segmentation accuracies achieved by conventional (*BiofilmQ*, *Seg3D*,

and Yan *et al.*), and *BCM3D* (this work). The estimated SBRs are 2.2, 1.8, and 1.3, respectively.

The estimated cell densities are 54.8%, 59.0%, and 64.6%, respectively. **(a-c)** Three experimental

E. coli datasets (cytosolic expression of GFP) acquired at different time points after inoculation of

cells. Segmentation accuracy is parameterized in terms of cell counting accuracy (y axis) and

accuracy of single-cell shape estimation (x axis, quantified as the single-cell Jaccard index). Curves

approaching the upper right-hand corner indicate higher overall segmentation accuracy. **(d)**

Comparison of segmentation results achieved at the t= 600 minutes time point by manual

annotation, and by BCM3D using *in silico*-trained CNNs only and after further refinement of CNN

outputs using *LCuts*. Segmentation results of the other methods are shown in **Figure S5**.

Figure 6.

Figure 6. 3D Segmentation accuracy evaluation using *M. xanthus* biofilm images (cell density =

36.2%, and SBR = 1.58) using *in silico*-trained CNN processing. **(a)** Maximum intensity projection

of a 3D *M. xanthus* fluorescence image. Cells were labeled with membrane-intercalating dye,

FM4-64. **(b)** Maximum intensity projection of the manually obtained 3D segmentation result. **(c)**

Maximum intensity projection of a CNN-based 3D segmentation result after *LCuts* post-

processing. Cells that can be matched with the GT are displayed in the same colors as GT or

otherwise colored in white. **(d)** Segmentation accuracy parameterized in terms of cell counting

accuracy (*y* axis) and accuracy of single-cell shape estimation (*x* axis, quantified as the single-cell

Jaccard index).

Figure 7.

Figure 7. **(a)** Cell counting accuracy of *BCM3D* on simulated images containing different ratios

of rod-shaped and spherical cells. Black diamonds represent the counting accuracy for each

independent dataset. **(b)** Cell counting accuracy of *BCM3D* on simulated images with different

ratios of membrane-labeled, and membrane-labeled and interior fluorescent protein expressing

cells. Black diamonds represent the counting accuracy for 10 independent datasets. **(c** and **d)** Cell

counting accuracy of *BCM3D* on experimental images of (c) membrane-labeled, and (d)

membrane-labeled and interior fluorescent protein expressing *E. coli* cells (mixing ratio ~ 1:1).

References

1. Hall-Stoodley, L., Costerton, J.W. & Stoodley, P. Bacterial biofilms: from the natural environment to infectious diseases. *Nat Rev Microbiol* **2**, 95-108 (2004).

2. O'Toole, G., Kaplan, H.B. & Kolter, R. Biofilm formation as microbial development. *Annual Review of Microbiology* **54**, 49-79 (2000).

3. Nadell, C.D., Drescher, K. & Foster, K.R. Spatial structure, cooperation and competition in biofilms. *Nat Rev Microbiol* **14**, 589-600 (2016).

4. Persat, A., Inclan, Y.F., Engel, J.N., Stone, H.A. & Gitai, Z. Type IV pili mechanochemically regulate virulence factors in Pseudomonas aeruginosa. *Proc Natl Acad Sci U S A* **112**, 7563-7568 (2015).

5. Zhang, Q., Lambert, G., Liao, D., Kim, H., Robin, K., Tung, C.-k., Pourmand, N. & Austin, R.H. Acceleration of Emergence of Bacterial Antibiotic Resistance in Connected Microenvironments. *Science* **333**, 1764-1767 (2011).

6. Donlan, R.M. Biofilms: microbial life on surfaces. *Emerg Infect Dis* **8**, 881-890 (2002).

7. Yan, J. & Bassler, B.L. Surviving as a Community: Antibiotic Tolerance and Persistence in Bacterial Biofilms. *Cell host & microbe* **26**, 15-21 (2019).

8. Kovach, K., Davis-Fields, M., Irie, Y., Jain, K., Doorwar, S., Vuong, K., Dhamani, N., Mohanty, K., Touhami, A. & Gordon, V.D. Evolutionary adaptations of biofilms infecting cystic fibrosis lungs promote mechanical toughness by adjusting polysaccharide production. *NPJ Biofilms Microbiomes* **3**, 1 (2017).

9. Berk, V., Fong, J.C., Dempsey, G.T., Develioglu, O.N., Zhuang, X., Liphardt, J., Yildiz, F.H. & Chu, S. Molecular architecture and assembly principles of *Vibrio cholerae* biofilms. *Science* **337**, 236-239 (2012).

10. Vidakovic, L., Singh, P.K., Hartmann, R., Nadell, C.D. & Drescher, K. Dynamic biofilm architecture confers individual and collective mechanisms of viral protection. *Nat Microbiol* **3**, 26-31 (2018).

11. Stewart, E.J., Ganesan, M., Younger, J.G. & Solomon, M.J. Artificial biofilms establish the role of matrix interactions in staphylococcal biofilm assembly and disassembly. *Sci Rep* **5**, 13081 (2015).

12. Stewart, E.J., Satorius, A.E., Younger, J.G. & Solomon, M.J. Role of environmental and antibiotic stress on Staphylococcus epidermidis biofilm microstructure. *Langmuir* **29**, 7017-7024 (2013).

13. Yan, J., Sharo, A.G., Stone, H.A., Wingreen, N.S. & Bassler, B.L. Vibrio cholerae biofilm growth program and architecture revealed by single-cell live imaging. *Proc Natl Acad Sci U S A* **113**, E5337-5343 (2016).

14. Drescher, K., Dunkel, J., Nadell, C.D., van Teeffelen, S., Grnja, I., Wingreen, N.S., Stone,
H.A. & Bassler, B.L. Architectural transitions in Vibrio cholerae biofilms at single-cell
resolution. *Proceedings of the National Academy of Sciences* **113**, E2066-E2072 (2016).

15. Hartmann, R., Singh, P.K., Pearce, P., Mok, R., Song, B., Díaz-Pascual, F., Dunkel, J. &
Drescher, K. Emergence of three-dimensional order and structure in growing biofilms.
Nature Physics **15**, 251-256 (2019).

16. Gregor, I. & Enderlein, J. Image scanning microscopy. *Curr Opin Chem Biol* **51**, 74-83
(2019).

17. Waldchen, S., Lehmann, J., Klein, T., van de Linde, S. & Sauer, M. Light-induced cell
damage in live-cell super-resolution microscopy. *Sci Rep* **5**, 15348 (2015).

18. Chen, B.C., Legant, W.R., Wang, K., Shao, L., Milkie, D.E., Davidson, M.W., Janetopoulos,
C., Wu, X.S., Hammer, J.A., 3rd, Liu, Z., English, B.P., Mimori-Kiyosue, Y., Romero, D.P.,
Ritter, A.T., Lippincott-Schwartz, J., Fritz-Laylin, L., Mullins, R.D., Mitchell, D.M.,
Bembenek, J.N., Reymann, A.C., Bohme, R., Grill, S.W., Wang, J.T., Seydoux, G., Tulu,
U.S., Kiehart, D.P. & Betzig, E. Lattice light-sheet microscopy: imaging molecules to
embryos at high spatiotemporal resolution. *Science* **346**, 1257998 (2014).

19. Li, D., Shao, L., Chen, B.C., Zhang, X., Zhang, M., Moses, B., Milkie, D.E., Beach, J.R.,
Hammer, J.A., 3rd, Pasham, M., Kirchhausen, T., Baird, M.A., Davidson, M.W., Xu, P. &
Betzig, E. Extended-resolution structured illumination imaging of endocytic and cytoskeletal
dynamics. *Science* **349**, aab3500 (2015).

20. Nadell, C.D., Ricaurte, D., Yan, J., Drescher, K. & Bassler, B.L. Flow environment and
matrix structure interact to determine spatial competition in Pseudomonas aeruginosa
biofilms. *Elife* **6**(2017).

21. Diaz-Pascual, F., Hartmann, R., Lempp, M., Vidakovic, L., Song, B., Jeckel, H., Thormann,
K.M., Yildiz, F.H., Dunkel, J., Link, H., Nadell, C.D. & Drescher, K. Breakdown of Vibrio
cholerae biofilm architecture induced by antibiotics disrupts community barrier function. *Nat
Microbiol* (2019).

22. Chang, B.J., Kittisopikul, M., Dean, K.M., Roudot, P., Welf, E.S. & Fiolka, R. Universal
light-sheet generation with field synthesis. *Nat Methods* **16**, 235-238 (2019).

23. Dean, K.M., Roudot, P., Welf, E.S., Danuser, G. & Fiolka, R. Deconvolution-free
Subcellular Imaging with Axially Swept Light Sheet Microscopy. *Biophysical journal* **108**,
2807-2815 (2015).

24. Chakraborty, T., Driscoll, M.K., Jeffery, E., Murphy, M.M., Roudot, P., Chang, B.J., Vora,
S., Wong, W.M., Nielson, C.D., Zhang, H., Zhemkov, V., Hiremath, C., De La Cruz, E.D.,
Yi, Y., Bezprozvanny, I., Zhao, H., Tomer, R., Heintzmann, R., Meeks, J.P., Marciano, D.K.,
Morrison, S.J., Danuser, G., Dean, K.M. & Fiolka, R. Light-sheet microscopy of cleared
tissues with isotropic, subcellular resolution. *Nat Methods* **16**, 1109-1113 (2019).

25. Dunsby, C. Optically sectioned imaging by oblique plane microscopy. *Optics Express* **16**, 20306-20316 (2008).

26. Voleti, V., Patel, K.B., Li, W., Perez Campos, C., Bharadwaj, S., Yu, H., Ford, C., Casper, M.J., Yan, R.W., Liang, W., Wen, C., Kimura, K.D., Targoff, K.L. & Hillman, E.M.C. Real-time volumetric microscopy of in vivo dynamics and large-scale samples with SCAPE 2.0. *Nat Methods* **16**, 1054-1062 (2019).

27. Flemming, H.C. & Wuertz, S. Bacteria and archaea on Earth and their abundance in biofilms. *Nat Rev Microbiol* **17**, 247-260 (2019).

28. York, A. High NA single-objective light-sheet. in *https://andrewgyork.github.io/high_na_single_objective_lightsheet/* (Github, 2020).

29. Voleti, V., Patel, K.B., Li, W., Campos, C.P., Bharadwaj, S., Yu, H., Ford, C., Casper, M.J., Yan, R.W. & Liang, W. Real-time volumetric microscopy of in vivo dynamics and large-scale samples with SCAPE 2.0. *Nature methods* **16**, 1054-1062 (2019).

30. Bouchard, M.B., Voleti, V., Mendes, C.S., Lacefield, C., Grueber, W.B., Mann, R.S., Bruno, R.M. & Hillman, E.M. Swept confocally-aligned planar excitation (SCAPE) microscopy for high-speed volumetric imaging of behaving organisms. *Nature photonics* **9**, 113 (2015).

31. Ulman, V., Maska, M., Magnusson, K.E.G., Ronneberger, O., Haubold, C., Harder, N., Matula, P., Matula, P., Svoboda, D., Radojevic, M., Smal, I., Rohr, K., Jalden, J., Blau, H.M., Dzyubachyk, O., Lelieveldt, B., Xiao, P., Li, Y., Cho, S.Y., Dufour, A.C., Olivo-Marin, J.C., Reyes-Aldasoro, C.C., Solis-Lemus, J.A., Bensch, R., Brox, T., Stegmaier, J., Mikut, R., Wolf, S., Hamprecht, F.A., Esteves, T., Quelhas, P., Demirel, O., Malmstrom, L., Jug, F., Tomancak, P., Meijering, E., Munoz-Barrutia, A., Kozubek, M. & Ortiz-de-Solorzano, C. An objective comparison of cell-tracking algorithms. *Nat Methods* **14**, 1141-1152 (2017).

32. Source code available at: https://github.com/GahlmannLab/BCM3D.git.

33. Chen, B.C., Legant, W.R., Wang, K., Shao, L., Milkie, D.E., Davidson, M.W., Janetopoulos, C., Wu, X.S., Hammer, J.A. & Liu, Z. Lattice light-sheet microscopy: imaging molecules to embryos at high spatiotemporal resolution. *Science* **346**, 1257998 (2014).

34. Valm, A.M., Cohen, S., Legant, W.R., Melunis, J., Hershberg, U., Wait, E., Cohen, A.R., Davidson, M.W., Betzig, E. & Lippincott-Schwartz, J. Applying systems-level spectral imaging and analysis to reveal the organelle interactome. *Nature* **546**, 162 (2017).

35. Glass, D.S. & Riedel-Kruse, I.H. A synthetic bacterial cell-cell adhesion toolbox for programming multicellular morphologies and patterns. *Cell* **174**, 649-658. e616 (2018).

36. Cotter, C.R., Schuttler, H.B., Igoshin, O.A. & Shimkets, L.J. Data-driven modeling reveals cell behaviors controlling self-organization during Myxococcus xanthus development. *Proc Natl Acad Sci U S A* **114**, E4592-E4601 (2017).

37. Gao, R., Asano, S.M., Upadhyayula, S., Pisarev, I., Milkie, D.E., Liu, T.-L., Singh, V.,
Graves, A., Huynh, G.H. & Zhao, Y. Cortical column and whole-brain imaging with
molecular contrast and nanoscale resolution. *Science* **363**, eaau8302 (2019).

38. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.Y., White, D.J., Hartenstein,
V., Eliceiri, K., Tomancak, P. & Cardona, A. Fiji: an open-source platform for biological-
image analysis. *Nature Methods* **9**, 676-682 (2012).

39. Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris, J.H. & Ferrin,
T.E. UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Sci*
27, 14-25 (2018).

40. Rudge, T.J., Steiner, P.J., Phillips, A. & Haseloff, J. Computational modeling of synthetic
microbial biofilms. *ACS Synth Biol* **1**, 345-352 (2012).

41. Reshes, G., Vanounou, S., Fishov, I. & Feingold, M. Cell shape dynamics in Escherichia
coli. *Biophysical journal* **94**, 251-264 (2008).

42. Starruß, J., Peruani, F., Jakovljevic, V., Søgaard-Andersen, L., Deutsch, A. & Bär, M.
Pattern-formation mechanisms in motility mutants of Myxococcus xanthus. *Interface focus* **2**,
774-785 (2012).

43. Harris, L.G., Foster, S. & Richards, R.G. An introduction to Staphylococcus aureus, and
techniques for identifying and quantifying S. aureus adhesins in relation to adhesion to
biomaterials: review. *Eur Cell Mater* **4**, 39-60 (2002).

44. Yan, T., Richardson, C.J., Zhang, M. & Gahlmann, A. Computational correction of spatially
variant optical aberrations in 3D single-molecule localization microscopy. *Optics Express* **27**,
12582-12599 (2019).

45. Muller, F.D., Schink, C.W., Hoiczyk, E., Cserti, E. & Higgs, P.I. Spore formation in
Myxococcus xanthus is tied to cytoskeleton functions and polysaccharide spore coat
deposition. *Mol Microbiol* **83**, 486-505 (2012).

46. Çiçek, Ö., Abdulkadir, A., Lienkamp, S.S., Brox, T. & Ronneberger, O. 3D U-Net: Learning
Dense Volumetric Segmentation from Sparse Annotation. 424-432 (Springer International
Publishing, Cham, 2016).

47. Gibson, E., Li, W., Sudre, C., Fidon, L., Shakir, D.I., Wang, G., Eaton-Rosen, Z., Gray, R.,
Doel, T., Hu, Y., Whyntie, T., Nachev, P., Modat, M., Barratt, D.C., Ourselin, S., Cardoso,
M.J. & Vercauteren, T. NiftyNet: a deep-learning platform for medical imaging. *Computer
Methods and Programs in Biomedicine* **158**, 113-122 (2018).

48. Van Valen, D.A., Kudo, T., Lane, K.M., Macklin, D.N., Quach, N.T., DeFelice, M.M.,
Maayan, I., Tanouchi, Y., Ashley, E.A. & Covert, M.W. Deep Learning Automates the
Quantitative Analysis of Individual Cells in Live-Cell Imaging Experiments. *PLoS Comput
Biol* **12**, e1005177 (2016).

49. Acton, S.T. Fast Algorithms for Area Morphology. *Digit Signal Process* **11**, 187-203 (2001).

50. Yu, Y. & Acton, S.T. Speckle reducing anisotropic diffusion. *Trans. Img. Proc.* **11**, 1260-
1270 (2002).

51. Tabassum, N., Vaccari, A. & Acton, S. Speckle removal and change preservation by
distance-driven anisotropic diffusion of synthetic aperture radar temporal stacks. *Digit Signal*
Process **74**, 43-55 (2018).

52. Choi, H.I., Choi, S.W. & Moon, H.P. Mathematical theory of medial axis transform. *Pacific*
Journal of Mathematics **181**, 57-88 (1997).

53. Yan, T.Q. & Zhou, C.X. A Continuous Skeletonization Method Based on Distance
Transform. *Comm Com Inf Sc* **304**, 251-258 (2012).

54. Wang, J., Batabyal, T., Zhang, M.X., Zhang, J., Aziz, A., Gahlmann, A. & Acton, S.T.
LCuts: Linear Clustering of Bacteria Using Recursive Graph Cuts. *2019 IEEE International*
Conference on Image Processing (ICIP), 1575-1579 (2019).

55. Source code available at: https://github.com/jwang-c/Postprocessing-using-LCuts.

56. Shi, J.B. & Malik, J. Normalized cuts and image segmentation. *Ieee Transactions on Pattern*
Analysis and Machine Intelligence **22**, 888-905 (2000).

57. Jaccard, P. The distribution of the flora in the alpine zone. *New Phytologist* **11**, 37-50 (1912).

58. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S. & Schmid, B. Fiji: an open-source platform for
biological-image analysis. *Nature methods* **9**, 676-682 (2012).

59. Ester, M., Kriegel, H.-P., Sander, J. & Xu, X. A density-based algorithm for discovering
clusters a density-based algorithm for discovering clusters in large spatial databases with
noise. in *Proceedings of the Second International Conference on Knowledge Discovery and*
Data Mining 226-231 (AAAI Press, Portland, Oregon, 1996).

60. Moen, E., Bannon, D., Kudo, T., Graf, W., Covert, M. & Van Valen, D. Deep learning for
cellular image analysis. *Nat Methods* (2019).

61. Caicedo, J.C., Roth, J., Goodman, A., Becker, T., Karhohs, K.W., Broisin, M., Molnar, C.,
McQuin, C., Singh, S., Theis, F.J. & Carpenter, A.E. Evaluation of Deep Learning Strategies
for Nucleus Segmentation in Fluorescence Images. *Cytometry Part A* **95**, 952-965 (2019).

62. Hartmann, R., Jeckel, H., Jelli, E., Singh, P.K., Vaidya, S., Bayer, M., Vidakovic, L., Díaz-
Pascual, F., Fong, J.C.N., Dragoš, A., Besharova, O., Nadell, C.D., Sourjik, V., Kovács,
Á.T., Yildiz, F.H. & Drescher, K. BiofilmQ, a software tool for quantitative image analysis
of microbial biofilm communities. *bioRxiv* (2019).

63. Reyer, M.A., McLean, E.L., Chennakesavalu, S. & Fei, J. An Automated Image Analysis
Method for Segmenting Fluorescent Bacteria in Three Dimensions. *Biochemistry* **57**, 209-
215 (2018).

64. Liu, J., Prindle, A., Humphries, J., Gabalda-Sagarra, M., Asally, M., Lee, D.Y., Ly, S.,
Garcia-Ojalvo, J. & Suel, G.M. Metabolic co-dependence gives rise to collective oscillations
within biofilms. *Nature* **523**, 550-554 (2015).

65. Prindle, A., Liu, J., Asally, M., Ly, S., Garcia-Ojalvo, J. & Suel, G.M. Ion channels enable
electrical communication in bacterial communities. *Nature* **527**, 59-63 (2015).

66. Humphries, J., Xiong, L., Liu, J., Prindle, A., Yuan, F., Arjes, H.A., Tsimring, L. & Suel,
G.M. Species-Independent Attraction to Biofilms through Electrical Signaling. *Cell* **168**,
200-209 e212 (2017).

67. Liu, J., Martinez-Corral, R., Prindle, A., Lee, D.-y.D., Larkin, J., Gabalda-Sagarra, M.,
Garcia-Ojalvo, J. & Süel, G.M. Coupling between distant biofilms and emergence of nutrient
time-sharing. *Science* **356**, 638-642 (2017).

68. Mitri, S. & Foster, K.R. The genotypic view of social interactions in microbial communities.
Annu Rev Genet **47**, 247-273 (2013).

69. Drescher, K., Nadell, C.D., Stone, H.A., Wingreen, N.S. & Bassler, B.L. Solutions to the
public goods dilemma in bacterial biofilms. *Current biology : CB* **24**, 50-55 (2014).

70. Persat, A., Nadell, C.D., Kim, M.K., Ingremeau, F., Siryaporn, A., Drescher, K., Wingreen,
N.S., Bassler, B.L., Gitai, Z. & Stone, H.A. The mechanical world of bacteria. *Cell* **161**, 988-
997 (2015).

71. Papenfort, K. & Bassler, B.L. Quorum sensing signal-response systems in Gram-negative
bacteria. *Nat Rev Microbiol* **14**, 576-588 (2016).

72. Kroos, L. Highly Signal-Responsive Gene Regulatory Network Governing Myxococcus
Development. *Trends Genet* **33**, 3-15 (2017).

73. Moon, S., Yan, R., Kenny, S.J., Shyu, Y., Xiang, L., Li, W. & Xu, K. Spectrally resolved,
functional super-resolution microscopy reveals nanoscale compositional heterogeneity in
live-cell membranes. *Journal of the American Chemical Society* **139**, 10944-10947 (2017).

74. Bramkamp, M. & Lopez, D. Exploring the existence of lipid rafts in bacteria. *Microbiol.
Mol. Biol. Rev.* **79**, 81-100 (2015).

75. Zou, S.B., Hersch, S.J., Roy, H., Wiggers, J.B., Leung, A.S., Buranyi, S., Xie, J.L., Dare, K.,
Ibba, M. & Navarre, W.W. Loss of elongation factor P disrupts bacterial outer membrane
integrity. *Journal of bacteriology* **194**, 413-425 (2012).

76. Gonelimali, F.D., Lin, J., Miao, W., Xuan, J., Charles, F., Chen, M. & Hatab, S.R.
Antimicrobial properties and mechanism of action of some plant extracts against food
pathogens and spoilage microorganisms. *Frontiers in microbiology* **9**, 1639 (2018).

77. Parasassi, T., De Stasio, G., d'Ubaldo, A. & Gratton, E. Phase fluctuation in phospholipid
membranes revealed by Laurdan fluorescence. *Biophysical journal* **57**, 1179-1186 (1990).

78. Strahl, H. & Hamoen, L.W. Membrane potential is important for bacterial cell division.
Proceedings of the National Academy of Sciences **107**, 12281-12286 (2010).

79. Prindle, A., Liu, J., Asally, M., Ly, S., Garcia-Ojalvo, J. & Süel, G.M. Ion channels enable
electrical communication in bacterial communities. *Nature* **527**, 59-63 (2015).

80. Weigert, M., Schmidt, U., Boothe, T., Müller, A., Dibrov, A., Jain, A., Wilhelm, B., Schmidt,
D., Broaddus, C. & Culley, S. Content-aware image restoration: pushing the limits of
fluorescence microscopy. *Nature methods* **15**, 1090-1097 (2018).

81. Krull, A., Vicar, T. & Jug, F. Probabilistic Noise2Void: Unsupervised content-aware
denoising. *arXiv preprint arXiv:1906.00651* (2019).

82. Krull, A., Buchholz, T.-O. & Jug, F. Noise2void-learning denoising from single noisy
images. in *Proceedings of the IEEE Conference on Computer Vision and Pattern
Recognition* 2129-2137 (2019).

83. Buchholz, T.-O., Prakash, M., Krull, A. & Jug, F. DenoiSeg: Joint Denoising and
Segmentation. *arXiv preprint arXiv:2005.02987* (2020).

Figure S1

Figure S1. Simulation of mixed labeling and mixed cell shape biofilms. (**a**) Cell arrangements (green indicates membrane labeled cells, purple indicates membrane labeled cells that simultaneously express interior fluorescence protein). (**b**) Fluorescence image based on the cell arrangements in (a) as displayed by the volume viewer plugin of Fiji[1]. (**c**) Cell arrangements (green indicates rod-shaped cells, magenta indicates spherical shaped cells). (**d**) Fluorescence image based on the cell arrangements in (c) as displayed by the volume viewer plugin of Fiji[1].

Figure S2

Figure S2. Binary segmentation result produced by thresholding the 'cell interior' confidence map at a high value (0.88-0.94). (**a**) Deconvolved fluorescence image. (**b**) 'Cell interior' confidence map. (**c**) Binary segmentation result (confidence threshold = 0.94). (**d** and **e**) Global voxel-level segmentation accuracy (y axis) versus the threshold of single-cell shape estimation (x axis, quantified as the single-cell Jaccard index) for cells labeled with cytosolic fluorophores (d) and cells labeled with membrane-localized fluorophores (e).

Figure S3

Figure S3. Determination of node direction in an outlier-removed neighborhood[2]. (**a**) A neighborhood of the target node (in red) is a sub-graph, where all adjacent nodes (in yellow) are connected via edges to the target node. Here, if the distance of two nodes is less than a chosen value (indicated by the dashed circle), these nodes are adjacent to each other. The blue dots are not part of the neighborhood. (**b**) A hop is defined as the number of edges that one has to traverse from one node to the other node in the graph. Here, the 5-hop neighborhood of the target node is shown. (**c**) The directional vectors are found from the target node to all the other nodes within in the 5-hop neighborhood (dashed lines). The nodes are classified as outliers if they have large relative angles compared to all the other directional vectors (red dashed lines). (**d**) Finally, the direction feature of the current node is evaluated as the major direction of the outlier removed neighborhood using principle component analysis.

Figure S4

Figure S4. Segmentation accuracies achieved for biofilm images with different SBR and cell densities. (**a**), (**b**), and (**c**) represent results for datasets with SBR 1.19, Density 54.5%, SBR 1.71, Density 54.5% and SBR 2.56, Density 54.5%, respectively. Each curve is plotted by averaging 10 different datasets.

Figure S5

Figure S5. Visual comparison of segmentation results achieved by previous segmentation approaches that rely solely on mathematical image processing. (**a**) Experimental dataset is a *E. coli* biofilm containing GFP expressing cells 10 hours after the inoculation. (**b**) Segmentation result obtained using *Seg3D*[3]. (**c**) Segmentation result obtained using the algorithm in Yan *et al.*[4]. (**d**) Segmentation result obtained using *BiofilmQ*[5].

Figure S6

Figure S6. Segmentation of mixed population biofilms containing spherical cells and rod-shaped cells. (**a**) Simulated fluorescence image of a mixture of spherical cells and rod-shaped cells. The mixing ratio is 50:50. (**b**) Segmentation result obtained by *BCM3D* and (**c**) ground truth. Three orthogonal planes are shown below each 3D image.

Figure S7

Figure S7. Segmentation of mixed population biofilms containing membrane-stained cells and membrane-stained cells that additionally express an intracellular fluorescent protein. (**a**) Simulated fluorescence image of a mixture of membrane-stained cells and membrane-stained cells that additionally express an intracellular fluorescent protein. The mixing ratio is 50:50. (**b**) Segmentation result obtained by *BCM3D* and (**c**) ground truth. Three orthogonal planes are shown below each 3D image.

Figure S8

Figure S8. **(a)** Experimental 2D slice of a mixed *E. coli* population containing membrane-stained cells and membrane-stained cells that additionally express an intracellular fluorescent protein. The mixing ratio at the time of inoculation was 50:50. All cells were labeled by the FM4-64 membrane-intercalating dye. **(b)** *BCM3D* segmentation result corresponding to the image shown in (a). Membrane-stained cells are displayed in green, and cells that were both membrane-stained and cytosolically-labeled are displayed in magenta.

References

1. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S. & Schmid, B. Fiji: an open-source platform for biological-image analysis. *Nature methods* **9**, 676-682 (2012).

2. Wang, J., Batabyal, T., Zhang, M.X., Zhang, J., Aziz, A., Gahlmann, A. & Acton, S.T. LCuts: Linear Clustering of Bacteria Using Recursive Graph Cuts. *2019 IEEE International Conference on Image Processing (ICIP)*, 1575-1579 (2019).

3. Reyer, M.A., McLean, E.L., Chennakesavalu, S. & Fei, J. An Automated Image Analysis Method for Segmenting Fluorescent Bacteria in Three Dimensions. *Biochemistry* **57**, 209-215 (2018).

4. Yan, J., Sharo, A.G., Stone, H.A., Wingreen, N.S. & Bassler, B.L. Vibrio cholerae biofilm growth program and architecture revealed by single-cell live imaging. *Proc Natl Acad Sci U S A* **113**, E5337-5343 (2016).

5. Hartmann, R., Jeckel, H., Jelli, E., Singh, P.K., Vaidya, S., Bayer, M., Vidakovic, L., Díaz-Pascual, F., Fong, J.C.N., Dragoš, A., Besharova, O., Nadell, C.D., Sourjik, V., Kovács, Á.T., Yildiz, F.H. & Drescher, K. BiofilmQ, a software tool for quantitative image analysis of microbial biofilm communities. BioRxiv 735423 (2019).