1

Non-Invasive Single-Cell Morphometry in Living Bacterial Biofilms

Mingxing Zhang[1,§], Ji Zhang[1,§], Yibo Wang[1,§], Jie Wang[3], Alecia M. Achimovich[2],

Scott T. Acton[3], and Andreas Gahlmann[1,2,*]

[1]Department of Chemistry, University of Virginia, Charlottesville, Virginia, USA

[2]Department of Molecular Physiology & Biological Physics, University of Virginia School of

Medicine, Charlottesville, Virginia, USA

[3]Department of Electrical & Computer Engineering, University of Virginia, Charlottesville,

Virginia, USA

§These authors contribute equally to this paper

*Corresponding Author : agahlmann@virginia.edu

Abstract

Fluorescence microscopy enables spatial and temporal measurements of live cells and cellular communities. However, this potential has not yet been fully realized for investigations of individual cell behaviors and phenotypic changes in dense, three-dimensional (3D) bacterial biofilms. Accurate cell detection and cellular shape measurement in densely packed biofilms are challenging because of the limited resolution and low signal to background ratios (SBRs) in fluorescence microscopy images. In this work, we present Bacterial Cell Morphometry 3D (*BCM3D*), an image analysis workflow that combines deep learning with mathematical image analysis to accurately segment and classify single bacterial cells in 3D fluorescence images. In *BCM3D*, deep convolutional neural networks (CNNs) are trained using simulated biofilm images with experimentally realistic SBRs, cell densities, labeling methods, and cell shapes. We systematically evaluate the segmentation accuracy of *BCM3D* using both simulated and experimental images. Compared to state-of-the-art bacterial cell segmentation approaches, *BCM3D* consistently achieves higher segmentation accuracy and further enables automated morphometric cell classifications in multi-population biofilms.

Introduction

Biofilms are multicellular communities of microorganisms that grow on biotic or abiotic surfaces[1-4]. In addition to cellular biomass, biofilms also contain an extracellular matrix (ECM) which is composed of polysaccharides, DNA, and proteins. Individual cells in biofilms interact with other cells, the ECM, or with the substrate surface, and the sum total of these interactions provide bacterial biofilms with emergent functional capabilities beyond those of individual cells. For example, biofilms are orders of magnitude more tolerant towards physical, chemical, and biological stressors, including antibiotic treatments and immune system clearance[1,2,5-8]. Understanding how such capabilities emerge from the coordination of individual cell behaviors requires imaging technologies capable of resolving and simultaneous tracking of individual bacterial cells in 3D biofilms.

Live cell-compatible imaging technologies, such as optical microscopy, can reveal the spatial and temporal context that affects cellular behaviors. However, conventional imaging modalities are not able to resolve individual cells within thick 3D biofilms over extended periods of time. For example, the diffraction-limited lateral x,y-resolution (~230 nm) of a confocal fluorescence microscope is barely sufficient to resolve bacterial cells positioned next to each other on flat glass coverslips. Even worse, the diffraction-limited axial z-resolution (570 nm) is comparable to the size of a single bacterial cell, so that densely-packed cells become unresolvable in the axial z-dimension[9,10]. Notable exceptions include loose biofilms (low cell density), spherical cell shapes[11,12], and mutant *Vibrio cholera* biofilms, in which cell-cell spacing is increased through the overproduction of ECM materials[13-15]. While single-cell resolved images have been obtained in such special situations, conventional optical microscopy modalities are not

generally capable to accurately resolve and quantitatively track individual cells in dense 3D

biofilms.

While super-resolution derivatives of confocal microscopy, known as Image Scanning

Microscopy[16], can improve spatial resolution, a perhaps more important limitation for long-term

live-cell imaging is photodamage to the specimen (phototoxicity) and to the fluorophores used

for labeling (photobleaching)[17-19]. In confocal microscopy-based approaches, undesired out-of-

focus fluorescence emission is filtered out by confocal pinholes to yield optically-sectioned

images with high contrast, i.e. high signal-to-background ratios (SBRs). However, repeated

illumination of out-of-focus regions during laser scanning and high light intensities at the focal

volume result in rapid photobleaching of fluorophores and unacceptable phototoxicity for light

sensitive specimens[17-19]. In fact, confocal fluorescence microscopy (as well as its super-

resolution derivatives) uses illumination light intensities that are two to three orders of

magnitude higher than the light intensities under which life has evolved[18]. The high rates of

phototoxicity and photobleaching make confocal-based microscopy unsuitable for high frame-

rate time-lapse imaging of living specimens over many hours and days[14,15,17,20,21].

In recent years, light sheet-based fluorescence excitation and imaging approaches have

been developed to overcome the drawbacks of confocal microscopy. Among these, lattice light

sheet microscopy (LLSM)[18,19] and field synthesis variants thereof[22], axially-swept light sheet

microscopy (ASLM)[23,24], dual-view light sheet microscopy[25,26], and single-objective oblique

plane light sheet microscopy[27-32] now combine excellent 3D spatial resolution with fast temporal

resolution and low phototoxicity at levels that cannot be matched by confocal microscopy.

Specifically, light sheet-based microscopy approaches can operate at illumination intensities that

are below the levels of cellular phototoxicity, even for notoriously light sensitive specimens, and

reduce fluorophore photobleaching by 20-50 times compared to confocal microscopy, while

maintaining comparable spatial resolution and contrast/SBR[18,28].

An additional challenge in high-resolution biofilm imaging is data quantification. Even if

sufficient resolution and high SBRs can be achieved to visually discern, i.e. qualitatively resolve

individual cells, robust computational algorithms are still needed for automated cell

segmentation and quantitative cell tracking. Towards this goal, image processing approaches

based on the watershed technique and intensity thresholding have been developed over the years

for single-cell segmentation in bacterial biofilms[13-15,21]. The broad applicability of watershed-

and threshold-based image processing algorithms is however limited, because these algorithms

require manual optimization of many user-selected parameters. Even with optimal parameters,

watershed- and threshold-based image processing methods often produce sub-optimal

segmentation results, especially when cell densities are high, when SBRs are low, and when

cellular fluorescence intensities are not uniform across the cytosol or the cell surface. To

overcome the drawbacks of traditional mathematical image processing approaches, automated

solutions based on supervised training of deep convolutional neural networks (CNNs) have been

used in recent years with great success for a wide range of problems in biomedical image

analysis[33].

Here, we present Bacterial Cell Morphometry 3D (*BCM3D*)[34], a generally applicable

workflow for single-cell segmentation and shape determination in high-resolution 3D images of

bacterial biofilms. *BCM3D* uses CNNs, *in silico*-trained with computationally simulated biofilm

images, in combination with mathematical image analysis to achieve accurate single cell

segmentation in 3D. The CNNs employed in *BCM3D* are based on the 3D U-Net architecture

and training strategy, which has achieved excellent performance in biomedical data analysis

benchmark tests[33]. The mathematical image analysis modules of *BCM3D* enable post-processing

of the CNN results to further improve the segmentation accuracy. We establish that experimental

bacterial biofilms images, acquired by lattice light sheet microscopy, can be successfully

segmented using CNNs trained with computationally simulated biofilm images, for which the

ground-truth voxel-level annotation maps are known accurately and precisely. By systematically

evaluating the performance of *BCM3D* for a range of SBRs, cell densities, and cell shapes, we

find that voxel-level segmentation accuracies of >80%, as well as cell counting accuracies of

>90%, can be robustly achieved. *BCM3D* consistently outperforms previously reported image

segmentation approaches that rely exclusively on conventional image processing approaches.

BCM3D also achieves higher segmentation accuracy on experimental 3D biofilm data than

Cellpose[35], a state-of-the-art, CNN-based, generalist algorithm for cell segmentation and the

algorithm used by Hartmann *et al.*[15], a specialized algorithm designed for bacterial cell

segmentation based on traditional mathematical image processing. We expect that *BCM3D*, and

CNN-based single-cell segmentation approaches in general, combined with non-invasive light

sheet-based fluorescence microscopy will enable accurate cell tracking over time in dense 3D

biofilms. This capability will launch a new era for bacterial biofilm research, in which the

emergent properties of microbial populations can be studied in terms of the fully-resolved

behavioral phenotypes of individual cells.

Results

Cell segmentation by thresholding CNN confidence maps

CNNs have been shown to perform well on pixel-level classification tasks for both 2D

and 3D data[36,37]. Bacterial biofilms, however, present a unique challenge in this context. The cell

shapes to be segmented are densely packed and barely resolvable even with the highest

resolution optical microscopes. Additionally, living biofilms in fluorescence microscopes can

only be imaged with low laser intensities to ameliorate phototoxicity and photobleaching

concerns. Unfortunately, low intensity fluorescence excitation also reduces the SBR in the

acquired images. So far, it remains unclear to what extent single-cell segmentation approaches

can accurately identify and delineate cell shapes in bacterial biofilm images obtained under low

intensity illumination conditions. To address this question, we implemented an *in silico* CNN

training strategy and systematically evaluated its voxel-level classification (cell morphometry)

and cell counting accuracies using simulated biofilm images with cell densities and SBRs similar

to those encountered in experimental data (see Methods).

We compared two commonly used cell labeling approaches, namely genetic labeling

through the expression of cell-internal fluorescent proteins and staining of the cell membranes

using fluorescent dyes (**Figure 1**). For both labeling approaches, voxel-level segmentation and

cell counting accuracies, obtained by thresholding CNN confidence maps (see Methods), depend

mostly on cell density, whereas the SBR plays a less important role (**Figure 2a-c** and **2d-f**). For

cell-internal labeling, SBRs of >1.7 and cell densities of <60% consistently produce voxel-level

classification accuracies of >80% and cell counting accuracies of >95%. On the other hand,

SBRs of <1.7 and cell densities of >60% lead to lower segmentation accuracies. While lower

segmentation accuracies are expected for higher cell densities and lower SBRs, the sharp drop-

offs observed here may indicate a fundamental performance limitation of the CNNs employed.

Still, the voxel-level classification and cell counting accuracies consistently surpass previous

approaches for bacterial cell segmentation for commonly encountered cell densities and SBRs.

Specifically, the cell counting accuracies obtained by Hartmann *et al.*[15], *Seg3D*[38], and Yan *et*

al.[13] quickly drop to zero as a function of increasing IoU matching threshold (a quantitative

measure of cell shape similarity relative to the ground truth, see Methods), indicating that cell

shapes are not accurately estimated by conventional image processing approaches (**Figure 2g-i**).

We also evaluated the segmentation accuracy of *Cellpose*, a recently developed, CNN-based

cellular segmentation algorithm[35]. The segmentation accuracy of *Cellpose* is comparable or

superior to the best-performing conventional image processing approaches – a considerable

achievement given that *Cellpose* was trained primarily on images of eukaryotic cells. However,

being a pre-trained generalist model, the segmentation accuracy of *Cellpose* is lower than the

accuracy achieved by the specialist *in silico*-trained CNNs of *BCM3D*, which were trained

specifically for 3D bacterial biofilm segmentation. Overall, the cell counting accuracies obtained

by *BCM3D* are higher than other methods and remain higher even for IoU matching thresholds

larger than 0.5, indicating that cell shapes are more accurately estimated by the *in silico*-trained

CNNs.

Figure 1. Simulation of fluorescent biofilms images and annotation maps used for CNN training. **(a)** Representative cell arrangements obtained by CellModeller. Due to the stochastic nature of biofilm growth, different cell arrangements are obtained in each new simulation. However, cell density is reproducible for each new simulated biofilm (typically $N = 10$ different biofilm simulations are used for CNN training, see Methods) **(b)** Simulated 3D fluorescence image based on the cell arrangements in **a**. **(c)** XY slice through the 3D simulated fluorescence image in **b** (upper panel shows cells expressing cytosolic fluorescent proteins, lower panel shows cells stained with membrane-intercalating dyes). **(d)** Ground truth cell arrangements giving rise to the image shown in **c**. Voxels are displayed as black (background), or in different colors (indicating different cells).

The accuracies of single-cell shape estimation and cell counting are predominantly

affected by cell density. The variation is more prominent for membrane-stained cells, because

inter-cellular fluorescence intensity minima are less pronounced when cell membranes are

labeled and cells physically contact each other (red arrow in **Figures 2c** and **2f**). By contrast,

intracellular fluorophores produce the highest intensities at the cell center, so that the gaps

between cells are more readily resolvable. Also noteworthy is the sharp drop-off in segmentation

accuracies for SBRs of <1.7 for all cases. In such low SBR regimes, fluorescence signals of the

cells become too difficult to be distinguished from the background. As a result, the CNNs falsely

identify random noisy patterns in the background as cells. Additionally, thresholding of the CNN

confidence maps often yields connected voxel clusters that contain multiple bacterial cells. False

identification and incomplete delineation of cells cause the pronounced decrease in segmentation

accuracy for SBRs of <1.7.

Figure 2. Performance of *BCM3D* using *in silico*-trained CNNs only on previously unseen simulated biofilm images. (a) The voxel-level segmentation accuracy quantifies whether each voxel has been assigned to the correct class ('cell interior', 'cell boundary', or 'background'). Solid circles represent the maximum local density and average SBRs encountered in experimental datasets (red, orange and blue: *E. coli* expressing GFP). **(b)** The cell counting accuracy (using an IoU matching threshold of 0.5 for each segmented object, see Methods) averaged over $N = 10$ replicate datasets for cells labeled with cytosolic fluorophores. **(c)** Example image of cells labeled with cytosolic fluorophores (Cell density = 60.0%, SBR = 1.34, indicated by white rectangle in panels **a** and **b**. Similar images were generated $N = 10$ times with different cell arrangements). **(d)** Voxel-level segmentation accuracy and **(e)** cell counting

accuracy averaged over $N = 10$ replicate datasets for cells labeled with membrane-localized

fluorophores. **(f)** Example image of cells labeled with membrane-localized fluorophores (Cell

density = 60.0%, SBR = 1.34, indicated by white rectangles in panels d and e. Similar images

were generated $N = 10$ times with different cell arrangements). The red arrows indicate a close

cell-to-cell contact point. **(g), (h) and (i)** Comparison of segmentation accuracies achieved by

conventional segmentation approaches (Hartmann *et al.*, *Seg3D*, Yan *et al.*), *Cellpose*, and

BCM3D (only using *in silico*-trained CNNs). Three simulated datasets (cytosolic fluorophores)

with different SBRs and cell densities are shown. Segmentation accuracy is parameterized in

terms of cell counting accuracy (*y* axis) and IoU matching threshold (*x* axis, a measure of cell

shape estimation accuracy). Each data point is the average of $N = 10$ independent biofilm

images. Data are presented as mean values ± one standard deviation indicated by error bars.

Curves approaching the upper right-hand corner indicate higher overall segmentation accuracy,

as indicated by the dashed arrows.

Post-processing of CNN confidence maps

To better identify individual cells in low SBR and high cell density datasets, we

developed a graph-based post-processing module (see Methods) that takes advantage of the fact

that bacterial cell shapes are highly conserved for a given species. Briefly, we transformed the

CNN 'cell interior' confidence maps into 3D point cloud data that trace out the central axes of

individual cells. This transformation was achieved by medial axis extraction using size-

constrained inscribed spheres[39] (**Figure S1a-c**). Single-cell axes are then identified as linearly

clustered data points by *LCuts* – a graph-based data clustering method designed to detect linearly

oriented groups of points[40]. The so-identified single-cell axes are then mapped back onto the

original segmentation volumes to obtain estimates of the 3D positions, shapes, and orientations

of the now separated cells (**Figure S1d**).

Post-processing with *LCuts* takes advantage of *a priori* knowledge about expected

bacterial cell sizes by removing erroneously segmented volumes that are significantly smaller

than the expected value and by splitting incompletely segmented volumes representing fused

cells. Improvements in cell counting accuracy of up to 15% and 36% are observed for cells

labeled with cytosolic fluorophores (**Figure 3a-c** and **Figure S2**) and membrane-localized

fluorophores (**Figure 3d-f**), respectively. The more substantial improvement for membrane-

stained cells is due to fact that CNNs trained on membrane-stained cells are more prone to

erroneously identifying speckled background noise as fluorescence signals in low SBR images.

In addition, membrane-intercalating fluorophores of two adjacent cells are in close proximity,

making it difficult to resolve fluorescence signals from two separate cells due to spatial signal

overlap (see the red arrow, **Figure 2c** and **2f**). *LCuts* thus provides an important benefit in

improving the cell counting accuracy to an extent not achieved by currently available

thresholding- or watershed-based post-processing algorithms (**Figure S3**).

Figure 3. Performance of *BCM3D* (*in silico*-trained CNNs and additional post-processing by *LCuts*) on previously unseen simulated data. (a) Voxel-level segmentation accuracy and **(b)** cell counting accuracy (using an IoU matching threshold of 0.5 for each segmented object) averaged over $N = 10$ replicate datasets for cells labeled with cytosolic fluorophores. **(c)** Improvement relative to *silico*-trained convolutional neural networks without post-processing. **(d)** Voxel-level segmentation accuracy and **(e)** cell counting accuracy averaged over $N = 10$ replicate datasets for cells labeled with membrane-localized fluorophores. **(f)** Improvements relative to *silico*-trained convolutional neural networks without post-processing.

Segmentation of experimental biofilm images

To test the performance of *BCM3D* on experimentally acquired biofilm images, we acquired time-lapse images of GFP expressing *E. coli* biofilms every thirty minutes for ten hours (see Methods). We then manually annotated one 2D slice in the 3D images at the t = 300, 360, and 600-minutes time points (see Methods). When referenced to these manual segmentation

results, the *LCuts*-processed CNN outputs consistently achieved better cell counting accuracies

than conventional segmentation methods (**Figure 4, Figure S4**). Initially, *Cellpose* and the

Hartmann *et al*. algorithm outperform the *in silico*-trained CNNs on two out of three of the test

images (*t* = 360 and 600 min), for which our *in silico*-trained CNNs struggle with

undersegmentation problems. However, mathematical post-processing of the CNN outputs by

LCuts corrects some of these errors, so that the integrated *BCM3D* workflow achieves improved

results compared to *Cellpose* and Hartmann *et al*. at each of the indicated time points. Visual

inspection of the segmentation results is also informative. *Cellpose* accurately segments

individual cells in low density regions, but suffers from oversegmentation errors in high density

biofilm regions (**Figure S4e**). The Hartmann *et al*. algorithm provides reasonable estimates of

cell positions in low and high density biofilm regions, but again struggles with cell shape

estimation (**Figure S4d**, see also **Figure 2g-i**). On the other hand, the integrated *BCM3D*

workflow (CNN + *LCuts*) produces biologically reasonable cell shapes regardless of cell density

(**Figure 4**).

Figure 4. **Comparison of segmentation accuracies achieved by conventional segmentation approaches (Hartmann *et al.*, *Seg3D*, Yan *et al.*), *Cellpose*, and *BCM3D*.** The estimated SBRs are 2.2, 1.8, and 1.3, respectively. The estimated cell densities are 54.8%, 59.0%, and 64.6%, respectively. **(a-c)** Three experimental *E. coli* datasets (cytosolic expression of GFP) acquired at different time points after inoculation of cells. Segmentation accuracy is parameterized in terms of cell counting accuracy (*y* axis) and IoU matching threshold (*x* axis). Each data point is the average of the cell counting accuracies calculated using annotation maps traced by $N = 3$ different researchers. Data are presented as mean values ± one standard deviation indicated by

error bars. Curves approaching the upper right-hand corner indicate higher overall segmentation

accuracy. **(d)** Comparison of segmentation results achieved at the $t = 600$ minutes time point by

manual annotation (shown is one of $N = 3$ researchers' annotation result, the other two

annotation results are shown in **Figure S4**), and by *BCM3D* using *in silico*-trained CNNs only

and after further refinement of CNN outputs using *LCuts*. Similar results were also obtained at

the $t = 300$ and $t = 360$ minute time points. Segmentation results of the other methods are shown

in **Figure S4**.

We attribute the more rapid drop-off of the cell counting accuracy as a function of

increasing IoU matching threshold in **Figure 4** to the following factors. First, human annotation

of experimentally acquired biofilm images differs from the ground truth segmentation masks that

are available for simulated data (**Figure S5**). The shape mismatches between algorithm

segmented and manually annotated cell shapes (**Figure S5** and **S6**) lead to a global lowering of

voxel-level segmentation accuracy and thus a more rapid drop-off of the cell counting accuracy

as a function of increasing IoU matching threshold. Because bacterial cell shapes are not

accurately captured by manual annotation (**Figures S5**), cell counting accuracies referenced to

manual annotations should be compared only at low IoU matching thresholds (0.1-0.3, shaded

grey in **Figure 4a-c**), as also pointed out previously[41]. We also note that bacterial cells in

experimental images appear motion-blurred if they are only partially immobilized and therefore

wiggle during image acquisition. Furthermore, optical aberrations and scattering effects were not

included in training data simulations, which may decrease the performance of the CNNs on

experimental data. Still, at IoU matching threshold < 0.3, the cell counting accuracy of *BCM3D*

remains above 75% while also producing biologically reasonable cell shapes. Thus, the bacterial

cell segmentation results of *BCM3D* represent a substantial improvement over other approaches

(**Figure 4** and **Figure S4**).

To demonstrate that *BCM3D* can achieve similarly high segmentation accuracies for

membrane-stained cells in different cellular arrangements, we analyzed a small patch of a

M. xanthus biofilm, which was stained with the membrane intercalating dye FM4-64 (**Figure**

5a). In contrast to *E. coli* biofilms, the submerged *M. xanthus* biofilm imaged here features cells

in a mesh-like arrangement with close cell-to-cell contacts, which presents a unique challenge for

3D single-cell segmentation. To obtain reference data for 3D segmentation accuracy

determination, we manually annotated each *xy*, *xz*, and *yz* slice of an entire 3D image stack

(**Figure 5b**). When referenced to this 3D manual segmentation result, *BCM3D* (**Figure 5c**)

produced cell counting accuracies above 70% at low (0.1-0.3) IoU matching thresholds, whereas

segmentation results obtained by conventional image processing (Hartmann *et al.*) and by

generalist CNN-processing (*Cellpose*) produced cell counting accuracies <50% in the same IoU

matching threshold region (**Figure 5d**). We note however that neither *Cellpose* nor the Hartmann

et al. algorithms were specifically optimized/designed for segmenting membrane-stained cells.

Indeed, the performance of *Cellpose* on this type of biofilm architecture is inferior to the results

achieved using the *in silico*-trained CNNs of *BCM3D* alone (without using *LCuts* post-

processing). One reason might be that the pre-trained, generalist *Cellpose* model has not been

trained sufficiently on long, thin, and highly interlaced rod-shaped cells, such as those contained

in a *M. xanthus* biofilm.

Figure 5. **3D Segmentation accuracy evaluation using *M. xanthus* biofilm images (cell density = 36.2%, and SBR = 1.58).** **(a)** Maximum intensity projection of a 3D *M. xanthus* fluorescence image. Cells were labeled with membrane-intercalating dye, FM4-64. Similar images were obtained at $N = 120$ different time points. **(b)** Maximum intensity projection of the manually obtained 3D segmentation result. **(c)** Maximum intensity projection of a CNN-based 3D segmentation result after *LCuts* post-processing. Cells that can be matched with the GT are displayed in the same colors as GT or otherwise colored in white. **(d)** Segmentation accuracy of compared algorithms parameterized in terms of cell counting accuracy (*y* axis) and IoU matching threshold (*x* axis).

Morphological separation of mixed cell populations

Given the improved segmentation results obtained using *BCM3D*, we reasoned that the same CNNs may have additional capacity to assign segmented objects to different cell types based on subtle morphological differences in the acquired images. Differences in the imaged cell morphologies arise due to physical differences in cell shapes (e.g. spherical vs. rod-shaped cells) or due to differences in the fluorescent labeling protocols (e.g. intracellular vs. cell membrane labeling), because fluorescence microscopes simply measure the spatial distributions of fluorophores in the sample. The ability to separate different cell morphologies is important for the study of multispecies biofilms, where interspecies cooperation and competition dictate population-level outcomes[3,42-49]. Separation of differentially labeled cells is also important for the study of gene activation in response to cell-to-cell signaling[50]. Expression of cytosolic fluorescent proteins by transcriptional reporter strains is a widely-used technique to visualize activation of a specific gene or genetic pathway in living cells. Such genetic labeling approaches can be complemented by chemical labeling approaches, e.g. using membrane intercalating chemical dyes that help visualize cells non-specifically or environmentally-sensitive membrane dyes that provide physiological information, including membrane composition[51,52], membrane organization and integrity[53-55], and membrane potential[56,57]. Chemical and genetic labeling approaches are traditionally implemented in two different color channels. However, there are important drawbacks to using multiple colors. First and foremost, the amount of excitation light delivered is increased by the necessity to excite differently colored fluorophores, raising phototoxicity and photobleaching concerns. Second, it takes *N* times as along to acquire *N*-color images (unless different color channels can be acquired simultaneously), making it challenging

354   to achieve high temporal sampling in time-lapse acquisition. For these reasons, methods that

355   extract complementary physiological information from a single-color image are preferable.

356   We evaluated the ability of *BCM3D* to automatically segment and identify rod-shaped

and spherical bacterial cells consistent with shapes of *E. coli* and *S. aureus* in simulated images

(**Figure S7**). To segment cells in two-population biofilms, we trained CNNs that classify pixels

into five different classes: 'background', 'cell interior of population 1', 'cell boundary of

population 1', 'cell interior of population 2' and 'cell boundary of population 2'. Thresholding

the CNNs confidence maps can achieve cell counting accuracies larger than 90% for both cell

types independent of their population fractions (**Figure 6a**). Post-processing of this result using

LCuts improved the cell counting accuracy by less than 0.5% on average, indicating that under-

segmented cell clusters are not prevalent in this dataset.

365   We next evaluated the ability of *BCM3D* to automatically segment and separate

membrane-stained cells that express cytosolic fluorescent proteins from those that do not (**Figure

S8**). Again, the cell counting accuracy is consistently above 80% for all tested mixing ratios

(**Figure 6b**). Finally, we applied *BCM3D* to experimentally acquired biofilm images of two

different *E. coli* strains. Both strains were stained by the membrane intercalating dye FM4-64,

but the second strain additionally expressed GFP (**Figure S9**). The cells were homogeneously

mixed prior to mounting to randomize the spatial distribution of different cell types in the

biofilm (see Materials and Methods). Multiple 2D slices from the 3D image stack were manually

annotated and compared with the results obtained by *BCM3D*. Consistent with the single-species

experimental data, a cell counting accuracy of 50% is achieved for each cell type at a 0.5 IoU

matching threshold and, at lower IoU matching thresholds, the counting accuracies increased to

60% to 70%, (**Figure 6cd**). Thus, using appropriately trained CNNs in *BCM3D* enables

automated and accurate cell type assignments based on subtle differences in cell morphologies in

mixed population biofilms – a capability not available using conventional image processing

methods.

Figure 6. Performance of *BCM3D* on mixed-population biofilm images. (a) Cell counting
accuracy of *BCM3D* on simulated images containing different ratios of rod-shaped and spherical
cells. Black diamonds represent the counting accuracy for $N = 10$ independently simulated
datasets. Green dots represent the cell density for each independent dataset. Error bars represent
± one standard deviation. **(b)** Cell counting accuracy of *BCM3D* on simulated images with
different ratios of membrane-labeled, and membrane-labeled and interior fluorescent protein
expressing cells. Black diamonds represent the counting accuracy for $N = 10$ independently
simulated datasets. Green dots represent the cell density for $N = 10$ independent datasets. Error
bars represent ± one standard deviation. **(c** and **d)** Cell counting accuracy of *BCM3D* on
experimental images of (c) membrane-labeled, and (d) membrane-labeled and interior
fluorescent protein expressing *E. coli* cells (mixing ratio ~ 1:1). Each data point is the average of
the cell counting accuracies calculated using annotation maps traced by three different
researchers ($N = 3$). Data are presented as mean values ± one standard deviation indicated by
error bars.

Discussion

CNNs have been successful applied to many different problems in biological image
analysis, but their ability to segment individual cells in 3D and time-lapse 3D bacterial biofilm
images has not yet been fully explored. Here, we demonstrated a CNN-based image analysis
workflow, termed *BCM3D*, for single-cell segmentation and shape classification (morphometry)
in 3D images of bacterial biofilms. In this work, we applied *BCM3D* to 3D images acquired by
lattice light sheet microscopy. However, *BCM3D* readily generalizes to 3D images acquired by
confocal microscopy or advanced super-resolution microscopy modalities, provided that realistic
image formation models are used to simulate the training datasets. The use of simulated training
data is a major advantage of *BCM3D*, because it overcomes inconsistencies inherent in manual
dataset annotation (**Figure S5** and **S6**) and thus solves the problem of obtaining sufficient

amounts of accurately annotated 3D image data. The ability to use simulated training data

provides needed flexibility not only in terms of the microscope platform used for imaging, but

also in terms of the bacterial cell shapes that are to be segmented.

We systematically investigated the advantages and limitations of *BCM3D* by evaluating

both voxel- and cell-level segmentation accuracies using simulated and experimental datasets of

different cell densities and SBRs. *BCM3D* enabled accurate segmentation of individual cells in

crowded environments and automatic assignments of individual cells to specific cell populations

for most of the tested parameter space. Such capabilities are not readily available when using

previously established segmentation methods that rely exclusively on conventional image and

signal processing algorithms.

While *BCM3D* surpasses the performance of previous approaches, we stress that further

improvements are possible and, for long-term, high frame-rate time-lapse imaging experiments,

absolutely needed. Our systematic analysis revealed that high cell density and low SBR datasets

are particularly challenging for the CNNs used in this work. Future work will therefore focus on

increasing the contrast and resolution in bacterial biofilm images. While, the use of optical super-

resolution modalities can provide higher spatial resolution, such resolution improvements often

come at a cost of reduced image contrast and faster photobleaching/phototoxicity. Software

solutions that can process images with limited resolution and low SBRs will therefore play a

tremendously important role in biological imaging. *BCM3D* is a general workflow that integrates

computational simulation of training data, *in silico*-training of CNNs for a *specific* task or a

specific cell type, and mathematical post-processing of the CNN outputs. Incorporating different

training strategies and different CNNs, such as the generalist CNN used in *Cellpose*[35], into the

BCM3D workflow will enable automated cross-validation of segmentation results when a ground

truth or manual annotation map is not available. Furthermore, CNN-based image processing

modules developed for contrast enhancement and denoising have also surpassed the performance

of conventional methods based on mathematical signal processing[58-61]. Incorporating these tools

into the *BCM3D* workflow promises to further improve the single-cell segmentation accuracies.

We anticipate that the ability to accurately identify and delineate individual cells in dense 3D

biofilms will enable accurate cell tracking over long periods of time. Detailed measurements of

behavioral single-cell phenotypes in larger bacterial communities will help determine how

macroscopic biofilm properties, such as its mechanical cohesion/adhesion and its biochemical

metabolism, emerge from the collective actions of *individual* bacteria.

Methods

Lattice Light Sheet Imaging of Bacterial Biofilms

Fluorescence images of bacterial biofilms were acquired on a home-built lattice light

sheet microscope (LLSM). LLSM enables specimen illumination with a thin light sheet derived

from 2D optical lattice[62,63]. Here, a continuous illumination light sheet was produced by a time-

averaged (dithered), square lattice pattern[18], and the illumination intensity at the sample was

<1 W/cm^2. The submicrometer thickness of the excitation light sheet is maintained over long

propagation distances (~30 µm), which enables optical sectioning, and thus high resolution, high

contrast imaging of 3D specimens comparable to confocal microscopy. However, fluorophore

excitation by a 2D light sheet reduces phototoxicity, because each excitation photon has multiple

opportunities to be absorbed by fluorophores in the excitation plane and produce in-focus

fluorescence. Widefield fluorescence images corresponding to each illuminated specimen plane

are recorded on a sCMOS detector (Hamamatsu ORCA Flash v2). In this work, 3D biofilm

images were acquired by translating the specimen through the light sheet in 200 nm steps using a

piezo nanopositioning stage (Physik Instrumente, P-621.1CD). The data acquisition program is

written in LabVIEW 2013 (National Instruments).

Ampicillin resistant *E.coli* K12, constitutively expressing GFP[64], were cultured at 37

degrees overnight in LB medium with 100 μg/ml ampicillin. Overnight cultures were diluted 100

times into the same culture medium, grown to an optical density at 600 nm (OD600) of 0.6 – 1.0,

and then diluted by an additional factor of 10. Round glass coverslips with the diameter of 5 mm

were put into a 24-well plate (Falcon) and 400 μL of cell culture was added to the well. Cells

were allowed to settle to the bottom of the well and adhere to the coverslip for 1 hour. The round

coverslips were then mounted onto a sample holder and placed into the LLSM sample-basin

filled with M9 medium. GFP fluorescence was excited using 488 nm light sheet excitation.

Biofilm growth was imaged at room temperature every 30 min for a total of 20 time points. At

each time point, a single 3D image stack contained 400 images, each acquired with a 15 ms

exposure time to avoid motion blur.

M.xanthus strain LS3908 expressing tdTomato under the control of the IPTG-inducible

promoter[65] and DK1622 (WT) were cultured in the nutrient rich CYE media at 30 degrees

Celsius until it reached an OD600 of 0.6 - 1.0. Media was supplemented with 1 mM IPTG for

tdTomato expressing cells. Chitosan (Thermo Fisher)-coated 5 mm round glass coverslips were

prepared by incubating coverslips with 1% (w/v) chitosan (1.5 % glacial acetic acid (v/v)) at

room temperature for 1 hour. Coverslips were then rinsed with water and placed into a 24-well

plate (Falcon) with 350-400 μL of undiluted cell culture. WT cells were stained directly in the 24

well plate with 5 ng/ml FM4-64 (Thermo Fisher) dye. Cells were allowed to settle and adhere to

the coverslip for 2 hours. After the settling period, the coverslip was gently rinsed with CYE

media to flush away unattached cells. The rinsed coverslip was then mounted onto a sample

holder and placed into the LLSM sample-basin filled with MC7 starvation buffer. tdTomato and

FM 4-64 fluorescence was excited using 561 nm light sheet excitation. The 3D image stack

contained 400 2D images. Each 2D slice was acquired with an exposure time of 30 ms.

For mixed population biofilm imaging, ampicillin resistant *E.coli* K12, constitutively

expressing GFP[64], and ampicillin resistant *E.coli* K12, expressing mScarlet (pBAD vector,

arabinose induce) were cultured separately at 37 degrees overnight in LB medium with 100

μg/ml ampicillin. Overnight cultures were diluted 100 times into the same culture medium,

grown to an optical density at 600 nm (OD600) of 0.6 – 1.0, and then diluted to an OD of 0.1.

After dilution, the two strains were mixed together. Round glass coverslips with the diameter of

5 mm were put into a 24-well plate (Falcon) and 500 μL of cell culture was added to the well.

Cells were allowed to settle to the bottom of the well and adhere to the coverslip for 1 hour. The

cell culture medium was then removed and replaced by 500 uL M9 medium containing 0.2%

(w/v) arabinose. The co-culture was incubated at 30 degrees overnight. 10 mins before imaging,

the co-culture was stained with 5 ng/ml FM4-64 (Thermo Fisher) dye. 3D image stacks of 20

planes with 5 ms exposure time per frame were acquired using 488 nm excitation.

Raw Data Processing

Raw 3D images were background subtracted and then deskewed and deconvolved[18,19].

The background was estimated by averaging intensity values of dark areas (devoid of cells) in

the field of view. Deconvolution was performed using the Richardson-Lucy algorithm with 10

iterations using experimentally measured point spread functions (PSFs) as the deconvolution

kernel. The experimentally measured PSFs were obtained separately for each color channel using

fluorescent beads (200 nm FluoSpheres®, Thermo Fisher) coated on a coverslip[66]. 3D images

were rendered using the 3D Viewer plugin in Fiji[67] or ChimeraX[68].

Generation of simulated biofilm images

To generate data for training of CNNs, we computationally simulated fluorescence

images of 3D biofilms, for which spatial arrangements among individual cells are known

precisely and accurately. Growth and division of individual rod-shaped cells in a population were

simulated using CellModeller, an individual-based computational model of biofilm growth

(**Figure 1a**)[69]. In individual-based biofilm growth models, cells are the basic modeling units.

Each cell is characterized by a set of parameters, including its 3D position, volume, and spatial

orientation. All the cells in the simulated biofilm are then allowed to evolve in time according to

predefined biological, chemical, and mechanical rules. For example, cells grow at a defined rate

and then divide after reaching a certain volume threshold. Cellular collisions that are due to cell

growth are alleviated by imposing a minimum distance criterion between cells at each time point.

For our simulations, we chose cell diameter and cell length (d, l) parameters consistent with a

given bacterial species, namely (1 μm, 3 μm) for *E. coli*[70], (0.7 μm, 6 μm) for *M. xanthus*[71], and

(1 μm, 1 μm) for spherically symmetric *S.aureus*[72].While the cell volume can be readily adjusted

in CellModeller, the cellular volume density, which is determined by the intercellular spacing, is

not directly adjustable. We therefore adjusted the cellular volume density after each simulation

by scaling the cellular positions (cell centroids) and thus the intercellular distances by a constant

factor, while leaving cell sizes, shapes, and orientations unchanged. This post-processing

procedure enabled simulation of the exact same 3D cell arrangements at adjustable cell volume

densities.

We fluorescently labeled simulated cell volumes and surfaces according to two

commonly used labeling strategies in fluorescence microscopy. To simulate expression of

intracellular fluorescent proteins, the fluorescence emitters were placed at random positions

within the cell volume. To simulate membrane staining, the fluorescence emitters were placed at

random positions on the cell surface. Each cell contained between 500 - 1000 fluorophores to

simulate expression level variations between cells, which is often observed in experimental

images. Once the fluorophore spatial distributions were determined, a 3D fluorescence image

(**Figure 1b**) was computationally generated. Each fluorophore was treated as an isotropic point

emitter, so that it would produce a diffraction-limited point-spread-function (PSF) on the

detector. Experimentally measured 3D PSF shapes (see Raw Data Processing) were used as the

convolution kernel. Next, the fluorescence signal intensity was scaled by multiplying the image

by a constant factor and then a constant background intensity was added to the image at ~200

photons per pixel, as measured in experimental data. This procedure enabled independent

adjustments of the fluorescence signal and background to obtain signal-to-background ratios

(SBRs) consistent with experimental data. In a final step, we introduced Poisson-distributed

counting noise, based on the summed background and signal intensities, as well as Gaussian-

distributed camera read-out noise (experimentally calibrated for our detector at 3.04 photons per

pixel on average)[73]. This resulting image data (**Figure 1c**) was then processed in the same

manner as experimental data (see Raw Data Processing). In contrast to experimental data,

generation of the corresponding voxel-level annotation maps is fast and error free, because the

underlying ground truth cell arrangements are known *a priori* (**Figure 1d**).

To mimic imaging of reporter gene expression in a subset of cells, we simulated biofilm

images, in which all cells were stained at the cell surface (e.g. with a membrane intercalating

fluorescent dye) and a subset of cells additionally contained intracellular fluorophores (e.g.

through the expression of an intracellular fluorescent protein) (**Figure S10ab**). The mixing ratios

between membrane-labelled, and membrane and interior labelled cells were 10:90, 30:70, 50:50,

70:30 and 90:10. Ten different cell arrangements containing ~300 cells were simulated for each

ratio. To train the CNNs (see next section), six datasets were used, all with a 50:50 mixing ratio.

To mimic imaging of cells with different morphologies, we simulated biofilms containing

spherical and rod-shaped cells (**Figure S10cd**). Cell arrangements were first simulated using rod

shaped cells and then a fraction of rod-shaped cells is replaced with spherical cells. The size of

the rod-shaped cells is that of *E. coli* (~3 × 1 μm, length by diameter). The size of the spherical

cells is that of *S. aureus* (~1 μm in diameter)[74]. Both cell types were labelled by intracellular

fluorophores, as described above. The mixing ratios between rod-shaped and spherical cells were

10:90, 30:70, 50:50, 70:30 and 90:10. Ten different cell arrangements containing ~300 cells were

simulated for each ratio. To train the CNNs (see next section), we picked one image from each

mixing ratio for a total of five images.

Training the convolutional neural networks We trained 3D U-Net CNNs for voxel-

level classification tasks[75] within the NiftyNet platform[76] (network architecture depth 4,

convolution kernel size 3, ReLU activation function, 32 initial feature maps, and random dropout

of 0.5 during training). To achieve robust performance, we trained these networks using five to

ten simulated biofilm images with randomly selected cell densities and signal-to-background

ratios (see Generation of simulated biofilm images). The same raw data processing steps used for

experimental data (see Raw Data Processing) were also applied to simulated data. 3D

deconvolved simulated data and their corresponding voxel-level annotations were used to train

the CNNs. Each image used for training contained ~9 million voxels. We trained CNNs by

classifying each voxel as 'background', 'cell interior' or as 'cell boundary' based on the

underlying cell arrangements. For mixed-species biofilms, two additional classes, 'cell interior'

and 'cell boundary' of the second species, were used. This type of annotation scheme has been

shown to increase separation of bacterial cells in 2D[77]. For data augmentation, we applied

NiftyNet's built-in scaling, rotation, and elastic deformation functions. Instead of the original

cross-entropy loss function combined with uniform sampling, we used the Dice loss function and

'balanced sampler', so that every label has the same probability of occurrence in training. All

networks were trained for 2000 to 3600 iterations with a learning rate of 0.0001. Using these

parameters, it took approximately 24 hours to train the CNNs on a NVIDIA Tesla V100 GPU

with 16 GB memory.

Thresholding of CNN-produced confidence maps

Voxel-level classification by CNNs generates different confidence maps (one confidence

map for each annotation class). The confidence values range between 0 and 1 and represent the

confidence of assigning individual voxels to a given class. After thresholding the 'cell interior'

confidence map to obtain a binary image (**Figure S11a-c**), connected voxel clusters can be

isolated and identified as single cell objects using 3D connected component labeling[78]. A

conservative size-exclusion filter was applied: small objects with a volume ~10 times less than

the expected cell size were considered background noise and filtered out using an area open

operator[78]. Since the cell-interior volumes do not contain the cell boundaries, we dilated each

object by 1-2 voxels to increase the cell volumes using standard morphological dilation[78]. The

threshold value to segment individual cell objects based on the 'cell interior' confidence map

was determined by plotting the overall voxel-level segmentation accuracy , quantified as the

Intersection-over-Union value (IoU value, aka Jaccard index[79]) versus the confidence value

thresholds (**Figure S11de**). Optimal voxel-level segmentation accuracies were consistently

obtained using confidence thresholds between 0.88 and 0.94. Throughout this work, we use 0.94

for cells labeled with intracellular fluorophores and 0.88 for cells labeled with membrane-

localized fluorophores.

Post-processing of U-Net result using a refined *LCuts* algorithm

Thresholding of the 'cell interior' confidence map produces a binary segmentation result

(background = 0, cell interior =1), where groups of connected, non-zero voxels identify

individual cells in most cases. However, when cells are touching, they are often not segmented as

individuals, but remain part of the same voxel cluster (undersegmentation). On the other hand, a

single cell may be erroneously split into smaller sub-cellular objects (oversegmentation). Finally,

in datasets with low SBR, connected voxel clusters may be detected that do not correspond to

cells and thus produce false positive objects (**Figure S1a**). To address these errors and improve

the segmentation accuracy further, we included additional mathematical image analysis steps to

post-process the CNN results and reduce undersegmentation and oversegmentation errors.

Step 1. False positive objects are identified by evaluating the coefficient of variation[80,81]

for each connected voxel cluster i:

$$CV_i = \frac{\sigma_i}{\mu_i} \quad (1)$$

where σ_i and μ_i denote the standard deviation and the mean of the intensity taken over all voxels

contained in connected voxel cluster i. If the coefficient of variation is larger than ρ, then the

current object will be classified as a false positive object and removed from the confidence map

by setting all its voxels to zero. The removed objects will then no longer be counted when

evaluating the cell counting accuracy. The value of ρ is selected based on the coefficient of

variation of the background. For the datasets analyzed here, this sample coefficient of variation

was determined to be $\rho = 1.1$. After CV-filtering, objects smaller than 25% of the expected

bacterial cell size are also removed by setting its voxels to zero. The remaining connected voxel

clusters are then considered for further processing (**Figure S1a**).

Step 2. To identify and delineate individual cells in the connected voxel clusters

identified in the previous step, we implemented medial axis extraction using the method of

inscribed spheres[39], with the constraint that the sphere radii do not exceed the expected diameter

of a single bacterial cell (e.g. $d = 0.8$ µm) (**Figure S1b** left). The set of N inscribed spheres are

tangent to the object's surface and parameterized by $(x_i, y_i, z_i; r_i<d/2)$ for $i = 1, …, N$.

Determination of the $(x_i, y_i, z_i; r_i)$ coordinates is achieved using the Euclidean distance transform

of the objects' boundary[82], so that the points with coordinates (x_i, y_i, z_i) reliably trace out the

central cell axes of individual bacterial cells (**Figure S1b** right).

Step 3. To separate different linear segments after cell axis extraction (**Figure S1c**), we

used a refined version of the linear cuts (*LCuts*) algorithm[40,83]. *LCuts* is a graph-based data

clustering method designed to detect linearly oriented groups of points with certain properties.

The fundamental elements of a weighted mathematical graph are nodes, edges, and edge weights.

Here, the points with coordinates (x_i, y_i, z_i) represent the graph nodes. Edges are the connections

among nodes. Edges are assigned weights, for example, to reflect the confidence that two nodes

belong to the same group. *LCuts* achieves grouping by assigning weights to edges in the fully

connected graph to reflect the similarity between two nodes. The features of each node include

its location and direction, where the location of each node is simply its Cartesian coordinates.

The direction of each node is found by first determining its 5-hop neighborhood, removing nodes

at large relative angles, and evaluating the major direction of the outlier removed neighborhood

(**Figure S12**).

The algorithm to separate the nodes into different groups is a recursive graph cutting

method[40]. Graph cuts (*e.g.* nCut[84]) disconnect the edges between two groups of nodes when the

combined weights of these edges are minimized. The weights, between node i and node j, are

calculated as follows:

$$w_{ij} = w_D \cdot w_T \quad (2)$$

where

$$w_D = \begin{cases} e^{-D_{ij}^2/\sigma_D^2} & \text{if } D_{ij}^2 \leq r \\ 0 & \text{if } D_{ij}^2 > r \end{cases} \quad (3)$$

$$w_T = e^{-(\cos(\theta_{ij})-1)^2/\sigma_T^2} \quad (4)$$

w_D weighs the distance between two nodes and w_T weighs difference between node

directions. D_{ij} is the Euclidean distance between node i and node j, and r is set to eliminate edges

between two far away nodes. θ_{ij} is the relative angle between the directions of nodes i and j. σ_D

and σ_T are adjustable parameters that control the rate of exponential decay. *LCuts* continues to

separate groups of nodes until each group satisfies a stopping criterion. The stopping criterion is

biologically inspired based on the expected length L of a single bacterial cell and a group's

linearity after each recursion. *LCuts* yields linearly oriented groups of points that trace out the

central axes of individual cells (**Figure S1c**). Importantly, cell separation is achieved without

having to specify the number of cells in the biofilm in advance. Furthermore, to limit the need

for optimization of postprocessing routines, the four adjustable parameters used in *LCuts*, namely

cell diameter d, the cell length L, and the decay parameters σ_D and σ_T are chosen based on *a*

priori knowledge about the bacterial cells under investigation. We found that the performance of

LCuts is not sensitive to the particular values of d, L, σ_D and σ_T as long as they are consistent

with the imaged bacterial cell sizes and shapes (**Figure S13**). Identification of single cells

provided by *LCuts* alleviates under-segmentation errors of the CNN-based segmentation.

Step 4. The final output of linear clustering can provide length, location and orientation

of each cell. Based on these linear clusters, the cellular architecture of the biofilms can be

reconstructed by placing geometrical models of cells in space as shown in **Figure S1d**. For fast

computation, spherocylinders are used as the geometrical model using a radius consistent with

the known sizes of bacterial cells. To further refine the cell surfaces to better align with the

CNN-segmented volumes, we enclosed the inscribed spheres found in Step 2 in a convex hull

(**Figure S1d**).

Performance Evaluation

We quantified segmentation accuracy both at the cell-level (object counting) and at the

voxel-level (cell shape estimation). To quantify the cell-level segmentation accuracy, we

designated segmented objects as true positive (*TP*) if their voxel overlap with the ground truth or

the manual annotation resulted in an IoU value larger than a particular IoU matching threshold.

This criterion ensures one-to-one matching. A threshold of 0.5 is typically chosen when

reporting single cell counting accuracy values [35,41]. We follow this convention here. If the

segmented cell object could not be matched to a ground truth/manually annotated cell volume,

then it was counted as a false positive (*FP*) and the IoU value of that segmented object was set to

zero. If a ground truth/manually annotated cell volume was not identified in the image, then it

was counted as false negative (*FN*). The cell (object) counting accuracy was then defined as

TP/(*TP*+*FP*+*FN*). The average IoU value over all segmented objects in the image quantifies the

voxel-level segmentation accuracy, i.e. the accuracy of cell shape estimation.

To evaluate the accuracy of cell segmentation on experimental data, three researchers

separately traced the cell contours on experimental 2D slices by using freehand selections in Fiji

ROI Manger[85]. Because human annotation is very time consuming (about 50 hours for a

complete 3D dataset containing ~300 cells in a 22 x 32 x 12 um^3 volume), one to three single 2D

slices were selected for each dataset. One exception is the 3D *M. xanthus*, for which the cell

outlines in all available x, y and z slices were traced manually (**Figure S14a**). For straight, rod-

shaped cells, the centroids of the resulting 2D cell contours all fall within the cell interior

volume. To group together the contours belonging to the same cells, the centroid of each contour

was projected along the x, y and z dimension. If the projected centroid was enclosed by any other

contour in a different slice, then the centroid of that contour was projected onto the plane of the

initial contour. Two contours were labeled as related if they contained each other's projected

centroids (**Figure S14b**). This process is repeated for all possible contour pairs and their

relationship is recorded in an adjacency matrix. Next, related contours were assigned to

individual cells (**Figure S14c**). To separate incorrectly grouped contours, we additionally

identified clusters of centroids using the DBSCAN point clustering algorithm[86] (**Figure S14d**).

In a final step, we manually removed incorrectly traced contours (**Figure S14e**). Cells are

reconstructed by creating convex hulls with the grouped contours (**Figure S14f** and **S14g**). This

procedure determined the approximate positions, shapes, and orientations of individual cells in

the 3D biofilm.

To estimate the SBRs of both simulated and experimental images, we manually selected

and determined the intensities of approximately ten 'signal' and ten 'background' regions in the

images. We computed the SBR as the mean signal intensity divided by the mean background

intensity. To estimate the local density of a biofilm, we partitioned the image into several 3D

tiles of size 64 by 64 by 8 voxels. We then estimated the local density as the total cell volume

contained in each tile divided by the tile volume. We calculated the mean density of the 10

densest tiles to define the 'local density' metric reported for each dataset in the paper. To

estimate the cell density in an experimentally acquired biofilm image, the same calculations were

performed on either 3D manual annotations (if available) or binary masks obtained by CNN-

processing.

Author Contributions

M.Z., J.Z., Y.W., J.W., S.T.A., and A. G. designed research;

M.Z., J.Z., Y.W., J.W., and A.M.A. performed research;

M.Z., J.Z., Y.W., J.W., S.T.A., and A. G. analyzed data;

and M.Z., J.Z., Y.W., J.W., S.T.A., and A. G. wrote the paper.

Competing Interests

The authors declare no competing interests.

Acknowledgements

This work was supported in part by the US National Institute of General Medical Sciences Grant No. 1R01GM139002 (A.G and S.T.A.) and by a Jeffress Memorial Trust Award in Interdisciplinary Research (to A.G and S.T.A.). M.Z. and J.W. were supported by a University of Virginia Presidential Fellowship in Data Science. Y.W. was supported in part by a Dean's M.S.-Ph.D. Fellowship in Data Science. We thank Lotte Søgaard-Andersen, Larry Shimkets, Huiwang Ai and Ingmar Riedel-Kruse for providing bacterial strains used in this work. We thank Karsten Siller, computational research consultant at the University of Virginia, for help with utilizing the Universities super-computing resources. We thank Knut Dresher and Eric Jelli for providing the segmentation results using the Hartmann *et al.* algorithm.

Biological materials availability

All strains used in this work are available from their original sources upon request.

Data availability

Source data are provided with this paper. All data used for generating the results presented in this paper are available from the corresponding author upon request.

Code availability

The code for running all the modules of BCM3D as well as training and test data is available at https://github.com/GahlmannLab/BCM3D.git[87], DOI: 10.5281/zenodo.4088658

References

1. Hall-Stoodley, L., Costerton, J.W. & Stoodley, P. Bacterial biofilms: from the natural
environment to infectious diseases. *Nat Rev Microbiol* **2**, 95-108 (2004).

2. O'Toole, G., Kaplan, H.B. & Kolter, R. Biofilm formation as microbial development. *Annual*
Review of Microbiology **54**, 49-79 (2000).

3. Nadell, C.D., Drescher, K. & Foster, K.R. Spatial structure, cooperation and competition in
biofilms. *Nat Rev Microbiol* **14**, 589-600 (2016).

4. Persat, A., Inclan, Y.F., Engel, J.N., Stone, H.A. & Gitai, Z. Type IV pili mechanochemically
regulate virulence factors in Pseudomonas aeruginosa. *Proc Natl Acad Sci U S A* **112**, 7563-
7568 (2015).

5. Zhang, Q., Lambert, G., Liao, D., Kim, H., Robin, K., Tung, C.-k., Pourmand, N. & Austin,
R.H. Acceleration of Emergence of Bacterial Antibiotic Resistance in Connected
Microenvironments. *Science* **333**, 1764-1767 (2011).

6. Donlan, R.M. Biofilms: microbial life on surfaces. *Emerg Infect Dis* **8**, 881-890 (2002).

7. Yan, J. & Bassler, B.L. Surviving as a Community: Antibiotic Tolerance and Persistence in
Bacterial Biofilms. *Cell host & microbe* **26**, 15-21 (2019).

8. Kovach, K., Davis-Fields, M., Irie, Y., Jain, K., Doorwar, S., Vuong, K., Dhamani, N.,
Mohanty, K., Touhami, A. & Gordon, V.D. Evolutionary adaptations of biofilms infecting
cystic fibrosis lungs promote mechanical toughness by adjusting polysaccharide production.
NPJ Biofilms Microbiomes **3**, 1 (2017).

9. Berk, V., Fong, J.C., Dempsey, G.T., Develioglu, O.N., Zhuang, X., Liphardt, J., Yildiz, F.H.
& Chu, S. Molecular architecture and assembly principles of *Vibrio cholerae* biofilms.
Science **337**, 236-239 (2012).

10. Vidakovic, L., Singh, P.K., Hartmann, R., Nadell, C.D. & Drescher, K. Dynamic biofilm
architecture confers individual and collective mechanisms of viral protection. *Nat Microbiol*
3, 26-31 (2018).

11. Stewart, E.J., Ganesan, M., Younger, J.G. & Solomon, M.J. Artificial biofilms establish the
role of matrix interactions in staphylococcal biofilm assembly and disassembly. *Sci Rep* **5**,
13081 (2015).

12. Stewart, E.J., Satorius, A.E., Younger, J.G. & Solomon, M.J. Role of environmental and
antibiotic stress on Staphylococcus epidermidis biofilm microstructure. *Langmuir* **29**, 7017-
7024 (2013).

13. Yan, J., Sharo, A.G., Stone, H.A., Wingreen, N.S. & Bassler, B.L. Vibrio cholerae biofilm
growth program and architecture revealed by single-cell live imaging. *Proc Natl Acad Sci U*
S A **113**, E5337-5343 (2016).

14. Drescher, K., Dunkel, J., Nadell, C.D., van Teeffelen, S., Grnja, I., Wingreen, N.S., Stone,
H.A. & Bassler, B.L. Architectural transitions in Vibrio cholerae biofilms at single-cell
resolution. *Proceedings of the National Academy of Sciences* **113**, E2066-E2072 (2016).

15. Hartmann, R., Singh, P.K., Pearce, P., Mok, R., Song, B., Díaz-Pascual, F., Dunkel, J. &
Drescher, K. Emergence of three-dimensional order and structure in growing biofilms.
Nature Physics **15**, 251-256 (2019).

16. Gregor, I. & Enderlein, J. Image scanning microscopy. *Curr Opin Chem Biol* **51**, 74-83
(2019).

17. Waldchen, S., Lehmann, J., Klein, T., van de Linde, S. & Sauer, M. Light-induced cell
damage in live-cell super-resolution microscopy. *Sci Rep* **5**, 15348 (2015).

18. Chen, B.C., Legant, W.R., Wang, K., Shao, L., Milkie, D.E., Davidson, M.W., Janetopoulos,
C., Wu, X.S., Hammer, J.A., 3rd, Liu, Z., English, B.P., Mimori-Kiyosue, Y., Romero, D.P.,
Ritter, A.T., Lippincott-Schwartz, J., Fritz-Laylin, L., Mullins, R.D., Mitchell, D.M.,
Bembenek, J.N., Reymann, A.C., Bohme, R., Grill, S.W., Wang, J.T., Seydoux, G., Tulu,
U.S., Kiehart, D.P. & Betzig, E. Lattice light-sheet microscopy: imaging molecules to
embryos at high spatiotemporal resolution. *Science* **346**, 1257998 (2014).

19. Li, D., Shao, L., Chen, B.C., Zhang, X., Zhang, M., Moses, B., Milkie, D.E., Beach, J.R.,
Hammer, J.A., 3rd, Pasham, M., Kirchhausen, T., Baird, M.A., Davidson, M.W., Xu, P. &
Betzig, E. Extended-resolution structured illumination imaging of endocytic and cytoskeletal
dynamics. *Science* **349**, aab3500 (2015).

20. Nadell, C.D., Ricaurte, D., Yan, J., Drescher, K. & Bassler, B.L. Flow environment and
matrix structure interact to determine spatial competition in Pseudomonas aeruginosa
biofilms. *Elife* **6**(2017).

21. Diaz-Pascual, F., Hartmann, R., Lempp, M., Vidakovic, L., Song, B., Jeckel, H., Thormann,
K.M., Yildiz, F.H., Dunkel, J., Link, H., Nadell, C.D. & Drescher, K. Breakdown of Vibrio
cholerae biofilm architecture induced by antibiotics disrupts community barrier function. *Nat*
Microbiol (2019).

22. Chang, B.J., Kittisopikul, M., Dean, K.M., Roudot, P., Welf, E.S. & Fiolka, R. Universal
light-sheet generation with field synthesis. *Nat Methods* **16**, 235-238 (2019).

23. Dean, K.M., Roudot, P., Welf, E.S., Danuser, G. & Fiolka, R. Deconvolution-free
Subcellular Imaging with Axially Swept Light Sheet Microscopy. *Biophysical journal* **108**,
2807-2815 (2015).

24. Chakraborty, T., Driscoll, M.K., Jeffery, E., Murphy, M.M., Roudot, P., Chang, B.J., Vora,
S., Wong, W.M., Nielson, C.D., Zhang, H., Zhemkov, V., Hiremath, C., De La Cruz, E.D.,
Yi, Y., Bezprozvanny, I., Zhao, H., Tomer, R., Heintzmann, R., Meeks, J.P., Marciano, D.K.,
Morrison, S.J., Danuser, G., Dean, K.M. & Fiolka, R. Light-sheet microscopy of cleared
tissues with isotropic, subcellular resolution. *Nat Methods* **16**, 1109-1113 (2019).

25. Qin, B., Fei, C., Bridges, A.A., Mashruwala, A.A., Stone, H.A., Wingreen, N.S. & Bassler,
B.L. Cell position fates and collective fountain flow in bacterial biofilms revealed by light-
sheet microscopy. *Science* (2020).

26. Wu, Y., Wawrzusin, P., Senseney, J., Fischer, R.S., Christensen, R., Santella, A., York, A.G.,
Winter, P.W., Waterman, C.M. & Bao, Z. Spatially isotropic four-dimensional imaging with
dual-view plane illumination microscopy. *Nature biotechnology* **31**, 1032-1038 (2013).

27. Dunsby, C. Optically sectioned imaging by oblique plane microscopy. *Optics Express* **16**,
20306-20316 (2008).

28. Voleti, V., Patel, K.B., Li, W., Perez Campos, C., Bharadwaj, S., Yu, H., Ford, C., Casper,
M.J., Yan, R.W., Liang, W., Wen, C., Kimura, K.D., Targoff, K.L. & Hillman, E.M.C. Real-
time volumetric microscopy of in vivo dynamics and large-scale samples with SCAPE 2.0.
Nat Methods **16**, 1054-1062 (2019).

29. Flemming, H.C. & Wuertz, S. Bacteria and archaea on Earth and their abundance in biofilms.
Nat Rev Microbiol **17**, 247-260 (2019).

30. York, A. High NA single-objective light-sheet. in
https://andrewgyork.github.io/high_na_single_objective_lightsheet/ (Github, 2020).

31. Voleti, V., Patel, K.B., Li, W., Campos, C.P., Bharadwaj, S., Yu, H., Ford, C., Casper, M.J.,
Yan, R.W. & Liang, W. Real-time volumetric microscopy of in vivo dynamics and large-
scale samples with SCAPE 2.0. *Nature methods* **16**, 1054-1062 (2019).

32. Bouchard, M.B., Voleti, V., Mendes, C.S., Lacefield, C., Grueber, W.B., Mann, R.S., Bruno,
R.M. & Hillman, E.M. Swept confocally-aligned planar excitation (SCAPE) microscopy for
high-speed volumetric imaging of behaving organisms. *Nature photonics* **9**, 113 (2015).

33. Ulman, V., Maska, M., Magnusson, K.E.G., Ronneberger, O., Haubold, C., Harder, N.,
Matula, P., Matula, P., Svoboda, D., Radojevic, M., Smal, I., Rohr, K., Jalden, J., Blau, H.M.,
Dzyubachyk, O., Lelieveldt, B., Xiao, P., Li, Y., Cho, S.Y., Dufour, A.C., Olivo-Marin, J.C.,
Reyes-Aldasoro, C.C., Solis-Lemus, J.A., Bensch, R., Brox, T., Stegmaier, J., Mikut, R.,
Wolf, S., Hamprecht, F.A., Esteves, T., Quelhas, P., Demirel, O., Malmstrom, L., Jug, F.,
Tomancak, P., Meijering, E., Munoz-Barrutia, A., Kozubek, M. & Ortiz-de-Solorzano, C. An
objective comparison of cell-tracking algorithms. *Nat Methods* **14**, 1141-1152 (2017).

34. Source code available at: https://github.com/GahlmannLab/BCM3D.git.

35. Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for
cellular segmentation. *bioRxiv*, 2020.2002.2002.931238 (2020).

36. Moen, E., Bannon, D., Kudo, T., Graf, W., Covert, M. & Van Valen, D. Deep learning for
cellular image analysis. *Nat Methods* (2019).

37. Caicedo, J.C., Roth, J., Goodman, A., Becker, T., Karhohs, K.W., Broisin, M., Molnar, C.,
McQuin, C., Singh, S., Theis, F.J. & Carpenter, A.E. Evaluation of Deep Learning Strategies
for Nucleus Segmentation in Fluorescence Images. *Cytometry Part A* **95**, 952-965 (2019).

38. Reyer, M.A., McLean, E.L., Chennakesavalu, S. & Fei, J. An Automated Image Analysis
Method for Segmenting Fluorescent Bacteria in Three Dimensions. *Biochemistry* **57**, 209-
215 (2018).

39. Choi, H.I., Choi, S.W. & Moon, H.P. Mathematical theory of medial axis transform. *Pacific*
Journal of Mathematics **181**, 57-88 (1997).

40. Wang, J., Batabyal, T., Zhang, M.X., Zhang, J., Aziz, A., Gahlmann, A. & Acton, S.T.
LCuts: Linear Clustering of Bacteria Using Recursive Graph Cuts. *2019 IEEE International*
Conference on Image Processing (ICIP), 1575-1579 (2019).

41. Weigert, M., Schmidt, U., Haase, R., Sugawara, K. & Myers, G. Star-convex Polyhedra for
3D Object Detection and Segmentation in Microscopy. in *2020 IEEE Winter Conference on*
Applications of Computer Vision (WACV) 3655-3662 (2020).

42. Liu, J., Prindle, A., Humphries, J., Gabalda-Sagarra, M., Asally, M., Lee, D.Y., Ly, S.,
Garcia-Ojalvo, J. & Suel, G.M. Metabolic co-dependence gives rise to collective oscillations
within biofilms. *Nature* **523**, 550-554 (2015).

43. Prindle, A., Liu, J., Asally, M., Ly, S., Garcia-Ojalvo, J. & Suel, G.M. Ion channels enable
electrical communication in bacterial communities. *Nature* **527**, 59-63 (2015).

44. Humphries, J., Xiong, L., Liu, J., Prindle, A., Yuan, F., Arjes, H.A., Tsimring, L. & Suel,
G.M. Species-Independent Attraction to Biofilms through Electrical Signaling. *Cell* **168**,
200-209 e212 (2017).

45. Liu, J., Martinez-Corral, R., Prindle, A., Lee, D.-y.D., Larkin, J., Gabalda-Sagarra, M.,
Garcia-Ojalvo, J. & Süel, G.M. Coupling between distant biofilms and emergence of nutrient
time-sharing. *Science* **356**, 638-642 (2017).

46. Mitri, S. & Foster, K.R. The genotypic view of social interactions in microbial communities.
Annu Rev Genet **47**, 247-273 (2013).

47. Drescher, K., Nadell, C.D., Stone, H.A., Wingreen, N.S. & Bassler, B.L. Solutions to the
public goods dilemma in bacterial biofilms. *Current biology : CB* **24**, 50-55 (2014).

48. Persat, A., Nadell, C.D., Kim, M.K., Ingremeau, F., Siryaporn, A., Drescher, K., Wingreen,
N.S., Bassler, B.L., Gitai, Z. & Stone, H.A. The mechanical world of bacteria. *Cell* **161**, 988-
997 (2015).

49. Papenfort, K. & Bassler, B.L. Quorum sensing signal-response systems in Gram-negative
bacteria. *Nat Rev Microbiol* **14**, 576-588 (2016).

50. Kroos, L. Highly Signal-Responsive Gene Regulatory Network Governing Myxococcus
Development. *Trends Genet* **33**, 3-15 (2017).

51. Moon, S., Yan, R., Kenny, S.J., Shyu, Y., Xiang, L., Li, W. & Xu, K. Spectrally resolved,
functional super-resolution microscopy reveals nanoscale compositional heterogeneity in
live-cell membranes. *Journal of the American Chemical Society* **139**, 10944-10947 (2017).

52. Bramkamp, M. & Lopez, D. Exploring the existence of lipid rafts in bacteria. *Microbiol.*
Mol. Biol. Rev. **79**, 81-100 (2015).

53. Zou, S.B., Hersch, S.J., Roy, H., Wiggers, J.B., Leung, A.S., Buranyi, S., Xie, J.L., Dare, K.,
Ibba, M. & Navarre, W.W. Loss of elongation factor P disrupts bacterial outer membrane
integrity. *Journal of bacteriology* **194**, 413-425 (2012).

54. Gonelimali, F.D., Lin, J., Miao, W., Xuan, J., Charles, F., Chen, M. & Hatab, S.R.
Antimicrobial properties and mechanism of action of some plant extracts against food
pathogens and spoilage microorganisms. *Frontiers in microbiology* **9**, 1639 (2018).

55. Parasassi, T., De Stasio, G., d'Ubaldo, A. & Gratton, E. Phase fluctuation in phospholipid
membranes revealed by Laurdan fluorescence. *Biophysical journal* **57**, 1179-1186 (1990).

56. Strahl, H. & Hamoen, L.W. Membrane potential is important for bacterial cell division.
Proceedings of the National Academy of Sciences **107**, 12281-12286 (2010).

57. Prindle, A., Liu, J., Asally, M., Ly, S., Garcia-Ojalvo, J. & Süel, G.M. Ion channels enable
electrical communication in bacterial communities. *Nature* **527**, 59-63 (2015).

58. Weigert, M., Schmidt, U., Boothe, T., Müller, A., Dibrov, A., Jain, A., Wilhelm, B., Schmidt,
D., Broaddus, C. & Culley, S. Content-aware image restoration: pushing the limits of
fluorescence microscopy. *Nature methods* **15**, 1090-1097 (2018).

59. Krull, A., Vicar, T. & Jug, F. Probabilistic Noise2Void: Unsupervised content-aware
denoising. *arXiv preprint arXiv:1906.00651* (2019).

60. Krull, A., Buchholz, T.-O. & Jug, F. Noise2void-learning denoising from single noisy
images. in *Proceedings of the IEEE Conference on Computer Vision and Pattern*
Recognition 2129-2137 (2019).

61. Buchholz, T.-O., Prakash, M., Krull, A. & Jug, F. DenoiSeg: Joint Denoising and
Segmentation. *arXiv preprint arXiv:2005.02987* (2020).

62. Chen, B.C., Legant, W.R., Wang, K., Shao, L., Milkie, D.E., Davidson, M.W., Janetopoulos,
C., Wu, X.S., Hammer, J.A. & Liu, Z. Lattice light-sheet microscopy: imaging molecules to
embryos at high spatiotemporal resolution. *Science* **346**, 1257998 (2014).

63. Valm, A.M., Cohen, S., Legant, W.R., Melunis, J., Hershberg, U., Wait, E., Cohen, A.R.,
Davidson, M.W., Betzig, E. & Lippincott-Schwartz, J. Applying systems-level spectral
imaging and analysis to reveal the organelle interactome. *Nature* **546**, 162 (2017).

64. Glass, D.S. & Riedel-Kruse, I.H. A synthetic bacterial cell-cell adhesion toolbox for
programming multicellular morphologies and patterns. *Cell* **174**, 649-658. e616 (2018).

65. Cotter, C.R., Schuttler, H.B., Igoshin, O.A. & Shimkets, L.J. Data-driven modeling reveals
cell behaviors controlling self-organization during Myxococcus xanthus development. *Proc*
Natl Acad Sci U S A **114**, E4592-E4601 (2017).

66. Gao, R., Asano, S.M., Upadhyayula, S., Pisarev, I., Milkie, D.E., Liu, T.-L., Singh, V.,
Graves, A., Huynh, G.H. & Zhao, Y. Cortical column and whole-brain imaging with
molecular contrast and nanoscale resolution. *Science* **363**, eaau8302 (2019).

67. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.Y., White, D.J., Hartenstein,
V., Eliceiri, K., Tomancak, P. & Cardona, A. Fiji: an open-source platform for biological-
image analysis. *Nature Methods* **9**, 676-682 (2012).

68. Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris, J.H. & Ferrin,
T.E. UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Sci*
27, 14-25 (2018).

69. Rudge, T.J., Steiner, P.J., Phillips, A. & Haseloff, J. Computational modeling of synthetic
microbial biofilms. *ACS Synth Biol* **1**, 345-352 (2012).

70. Reshes, G., Vanounou, S., Fishov, I. & Feingold, M. Cell shape dynamics in Escherichia
coli. *Biophysical journal* **94**, 251-264 (2008).

71. Starruß, J., Peruani, F., Jakovljevic, V., Søgaard-Andersen, L., Deutsch, A. & Bär, M.
Pattern-formation mechanisms in motility mutants of Myxococcus xanthus. *Interface focus* **2**,
774-785 (2012).

72. Harris, L.G., Foster, S. & Richards, R.G. An introduction to Staphylococcus aureus, and
techniques for identifying and quantifying S. aureus adhesins in relation to adhesion to
biomaterials: review. *Eur Cell Mater* **4**, 39-60 (2002).

73. Yan, T., Richardson, C.J., Zhang, M. & Gahlmann, A. Computational correction of spatially
variant optical aberrations in 3D single-molecule localization microscopy. *Optics Express* **27**,
12582-12599 (2019).

74. Muller, F.D., Schink, C.W., Hoiczyk, E., Cserti, E. & Higgs, P.I. Spore formation in
Myxococcus xanthus is tied to cytoskeleton functions and polysaccharide spore coat
deposition. *Mol Microbiol* **83**, 486-505 (2012).

75. Çiçek, Ö., Abdulkadir, A., Lienkamp, S.S., Brox, T. & Ronneberger, O. 3D U-Net: Learning
Dense Volumetric Segmentation from Sparse Annotation. 424-432 (Springer International
Publishing, Cham, 2016).

76. Gibson, E., Li, W., Sudre, C., Fidon, L., Shakir, D.I., Wang, G., Eaton-Rosen, Z., Gray, R.,
Doel, T., Hu, Y., Whyntie, T., Nachev, P., Modat, M., Barratt, D.C., Ourselin, S., Cardoso,

M.J. & Vercauteren, T. NiftyNet: a deep-learning platform for medical imaging. *Computer Methods and Programs in Biomedicine* **158**, 113-122 (2018).

77. Van Valen, D.A., Kudo, T., Lane, K.M., Macklin, D.N., Quach, N.T., DeFelice, M.M., Maayan, I., Tanouchi, Y., Ashley, E.A. & Covert, M.W. Deep Learning Automates the Quantitative Analysis of Individual Cells in Live-Cell Imaging Experiments. *PLoS Comput Biol* **12**, e1005177 (2016).

78. Acton, S.T. Fast Algorithms for Area Morphology. *Digit Signal Process* **11**, 187-203 (2001).

79. Jaccard, P. The distribution of the flora in the alpine zone. *New Phytologist* **11**, 37-50 (1912).

80. Yu, Y. & Acton, S.T. Speckle reducing anisotropic diffusion. *Trans. Img. Proc.* **11**, 1260-1270 (2002).

81. Tabassum, N., Vaccari, A. & Acton, S. Speckle removal and change preservation by distance-driven anisotropic diffusion of synthetic aperture radar temporal stacks. *Digit Signal Process* **74**, 43-55 (2018).

82. Yan, T.Q. & Zhou, C.X. A Continuous Skeletonization Method Based on Distance Transform. *Comm Com Inf Sc* **304**, 251-258 (2012).

83. Source code available at: https://github.com/jwang-c/Postprocessing-using-LCuts.

84. Shi, J.B. & Malik, J. Normalized cuts and image segmentation. *Ieee Transactions on Pattern Analysis and Machine Intelligence* **22**, 888-905 (2000).

85. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S. & Schmid, B. Fiji: an open-source platform for biological-image analysis. *Nature methods* **9**, 676-682 (2012).

86. Ester, M., Kriegel, H.-P., Sander, J. & Xu, X. A density-based algorithm for discovering clusters a density-based algorithm for discovering clusters in large spatial databases with noise. in *Proceedings of the Second International Conference on Knowledge Discovery and Data Mining* 226-231 (AAAI Press, Portland, Oregon, 1996).

87. Zhang, M., Zhang, J., Wang, Y., Wang, J., Achimovich, A.M., Acton, S.T. & Gahlmann, A. Non-Invasive Single-Cell Morphometry in Living Bacterial Biofilms, GitHub Repository: BCM3D, DOI:10.5281/zenodo.4088658. (2020).