Bioremediation potential of bacterial species from a contaminated aquifer undergoing intrinsic remediation

A contaminated aquifer and bioremediation potential of bacterial species

Daniel Abiriga[*], Andrew Jenkins, Live Semb Vestgarden and Harald Klempe

Department of Natural Sciences and Environmental Health, Faculty of Technology, Natural Sciences and Maritime Sciences, University of South-Eastern Norway, Campus Bø, Mid-Telemark, Norway.

* Corresponding author

Email: daniel.abiriga@usn.no (DA)

Abstract

As a mean to understand bacterial species involved in an ongoing bioremediation of a confined aquifer contaminated by a municipal solid waste landfill, culture-dependent and fluorescence microscopic techniques were used. Water samples from the contaminated aquifer and a background aquifer were extracted and subjected to chemical, bacteriological and microscopic analysis. Eighty-seven bacterial species were isolated, representing four phyla: *Actinobacteria* (25.3%), *Bacteroidetes* (16.1%), *Firmicutes* (3.4%) and *Proteobacteria* (55.2%). Among the *Proteobacteria* were *Alphaproteobacteria*, *Betaproteobacteria*, and *Gammaproteobacteria*, in order of increasing abundance. A clear distinction between uncontaminated and contaminated groundwater was observed. Water samples from the uncontaminated groundwater had both low cell density and low species richness, as is expected of oligotrophic aquifers. On the other hand, water samples from the contaminated groundwater had higher cell density and species richness. The highest species richness was recorded from the distal well. The difference observed between the uncontaminated and contaminated groundwater samples highlights the influence of the landfill leachate on the microbial ecology. Majority of the species detected in the contaminated groundwater represented taxa frequently recovered from contaminated environments, with 47% of these having documented bioremediation potential either at species or at genus level. It is likely that the landfill leachate promoted a community of mostly heterotrophic culturable bacteria, as comparison between direct microscopic and plate counts seems to suggest so.

Introduction

Groundwater is the main source of freshwater for drinking, agriculture and industry in many places globally [1, 2], but faces serious pollution challenges worldwide [3]. Among the

human activities that have caused severe damage to groundwater resources is landfill. All

over the world, landfills have served as the ultimate destination for municipal solid wastes [4],

and continue to do so [5]. In Norway, there was little recycling of wastes until the late 1990s

and most of the wastes from households and industries were deposited in municipal solid

waste landfills with no provision for treatment of the resultant leachate. Revdalen Landfill

represents one such historic site and was active from 1974 to 1997, which led to the

contamination of Revdalen Aquifer.

Several strategies exist to reclaim contaminated aquifers. They are broadly categorised as

natural and artificial. The latter, which includes the conventional pump and treat (P&T) are

faster, but require a major economic input for operation and maintenance [6]. Natural

attenuation such as *in situ* bioremediation on the other hand, offers inexpensive, eco-friendly

yet efficient remedies [7, 8]. In addition, unlike P&T, *in situ* bioremediation does not generate

secondary wastes. It is the most widely accessed in *in situ* remediation of groundwater [2].

However, *in situ* bioremediation, particularly intrinsic bioremediation is slow and the

groundwater remains polluted for long, although it may be speeded up by amended

bioremediation [7].

Traditionally, groundwater bioremediation has been demonstrated empirically by measuring

geochemical parameters. Over the years, however, it has become apparent that studying

microbial community composition in addition to geochemical measurements offers a more

complete picture of bioremediation. In order to make inferences about bioremediation and

effectively manage the processes, a survey to establish which microorganisms are responsible

is necessary [9-11]. Both past and recent studies have been conducted on microbiomes of

contaminated aquifers [12-20]. Nonetheless, this area still requires more elucidation [21, 22].

Bioremediation of hydrocarbon-polluted aquifers is well documented in the literature [14, 16,

19, 20, 23-25], while there is a dearth of studies on landfill-leachate contaminations. The bias

may reflect high-profile cases of hydrocarbon pollutions and the potential health hazard

presented by the concomitant xenobiotics, which are often toxic, mutagenic and carcinogenic

[7]. Moreover, hydrocarbons, at least are, easily degraded in the environment and their

compositions are less complex than effluents emanating from landfills. The complicated

attenuation processes in landfill-leachate-impacted groundwater makes assessments of

bioremediation processes a more difficult and less attractive venture, which therefore receives

less attention than it deserves.

In the present study, we studied microbial diversity of a landfill-leachate-contaminated

confined aquifer. The aquifer is undergoing intrinsic bioremediation and as a mean to

understand which microbes may be responsible for the bioremediation, we isolated and

characterised bacterial species from the aquifer. Parallel to this, we conducted direct

microscopic count as a check for bias from culture-dependent method. High throughput

sequencing is also being undertaken and is the subject of a future manuscript.

Materials and methods

Groundwater sampling and chemical analysis

Groundwater sampling is described elsewhere (Abiriga et al., unpublished). pH and electrical

conductivity were determined in the field using pH-110 meter (VWR International) and Elite

CTS Tester (Thermo Scientific, Singapore), respectively. Dissolved oxygen was determined

using the Winkler method. Alkalinity was measured upon arrival at the laboratory using

Mettler DL25 Titrator (Mettler Toledo, Switzerland). Major ions (sodium, potassium,

calcium, magnesium, chloride, nitrate and sulphate) were determined using Ion

Chromatography DIONX ICS-1100 (Thermo Scientific, USA). Total nitrogen and total

organic carbon were determined using FIAlyzer-1000 (FIAlab, USA) and TOC Fusion

(Teledyne Tekmar, USA), respectively. Iron and manganese were measured using AAnalyst-

400 (PerkinElmer, USA).

Bacteriology

Direct microscopic count

Water samples were collected in sterile 350 ml PETE bottles (VWR, UK). 4.5 ml water

samples were fixed with 2.5% phosphate-buffered glutaraldehyde and stained with 5 µg/ml

4′,6-diamidino-2-phenylindole (DAPI) [26]. The stained cells were filtered onto 0.2 µm black

polycarbonate Nuclepore membrane filters (Sigma-Aldrich, Germany), transferred onto

microscope slides and overlaid with antifade mountant oil (Citifluor AF87, EMS, PA, USA).

Cells were enumerated under × 100 oil objective using Olympus IX70 fluorescence

microscope (Tokyo, Japan). Ten fields were counted and the average count was used to

estimate bacterial density using the formula:

Bacteria (Cells/ml) = $(N \times A_t)/(V_f \times A_g \times d)$,

where N = average number of cells, A_t = effective area of the filter paper, V_f = volume of

water sample filtered, A_g = area of the counting grid, and d = dilution factor [26]. No

observable cells were found in our blanks and therefore correcting for background noise due

to contamination was not necessary.

Evaluation of growth media

Serial dilutions of up to 10^6 of representative water samples were prepared. 100 µl of the

dilutions were spread on nutrient agar, plate count agar and full-strength, half-strength, one-

third-strength and one-fourth-strength tryptic soy agar in triplicate. Inoculated plates were

incubated at 15 ℃ for >5 days.

Sulphate-reducing medium was prepared following method by Ňancucheo, Rowe (27), with

modification as follow; no addition of trace elements and pH adjusted to 6.5. Sulphide-

oxidising medium was prepared according to Gregorich and Carter (28). Lastly, iron-

oxidising medium was prepared according to Holanda, Hedrich (29), again adjusting to pH

6.5. The media were inoculated with 100 µl of water samples. Plates for sulphide-oxidising

were incubated at least one week, while sulphate-reducing and iron-oxidising plates were

incubated for at least four weeks. Sulphide-oxidising and iron-oxidising plates were incubated

under aerobic condition, while sulphate-reducing plates were incubated under anaerobic

condition.

Colony counting

Aerobes were counted following incubation under aerobic condition. Anaerobes were

enumerated in a parallel setup following incubation under anaerobic condition using GasPak

with EZ Anaerobe Container System (BD, USA). Two plates from two dilutions of each

sample were counted and the average was reported as colony forming units per ml.

Based on observable colony morphologies such as shape, elevation, margin, size, and colour,

etc., colonies covering the full diversity of colony morphologies were picked and purified by

repeated streaking and incubation until pure cultures were obtained. Pure cultures were then

subjected to wet field microscopy at × 1,000 (Olympus, CX22LEDRFS1, China) to observe

motility and cell shape, gram staining, oxidase activity, and catalase activity [30]. Strains

were stored at -70 ℃ in nutrient broth (Sigma, Switzerland) supplemented with 25% glycerol.

16S rRNA sequencing and strain identification

Approximately 5.0 µl of each pure culture was suspended in 750 µl PBS and centrifuged at

10,000 × g for 3 minutes. The supernatant discarded and the cell pellet stored at -20 ℃ prior

to DNA extraction. Frozen cell pellets were allowed to thaw and 2.5 µl of the suspension was

re-suspended in 100 µl of PCR-grade distilled water. The tubes were incubated at 99 ℃ for 10

minutes and then centrifuged at 10,000 × g for 3 minutes. The DNA in the supernatant was

quantified using Qubit® Flourometer 3.0 (Life Technologies, Malaysia).

A universal primer set (341f/926r) [31, 32] was used to amplify the 16S rRNA gene region

V3-V5. The PCR reaction was carried out in a 50 µl reaction mixture containing 10-50 ng

template DNA; 1X Perfecta SYBR Green PCR FastMix (Quanta Biosciences); 300 nM

forward and reverse primer (341f/926r, Invitrogen). Amplification was performed in a Real-

Time PCR format using the StepOne instrument (Applied Biosystems, Foster City, CA, USA)

with the following temperature profiles: initial holding at 95 ℃ for 10 min; 45 cycles of

denaturing (95 ℃, 15 sec), annealing (50 ℃, 30 sec) and extension (72 ℃, 2 min). A melting

curve was performed over the temperature range 60-95 ℃.

PCR product clean-up was conducted using one-step enzymatic EXO STAR solution (GE

Healthcare Illustra ExoProStar, UK) following manufacturer's instructions. Cycle Sequencing

was performed using BigDye Terminator v1.1 Sequencing Kit (ThermoFisher Scientific)

according to manufacturer's instructions. For clean-up of sequencing products, both ethanol

precipitation and BigDye XTerminator Purification Kit (ThermoFisher scientific) were

assessed. The former yielded better results and was subsequently preferred to the latter.

Purified Cycle Sequencing products were analysed using Genetic Analyser 3130xl from

Applied Biosystems. The 16S rRNA gene sequences of the 87 isolates have been deposited in

GenBank under Accession No. MT348616-MT348702.

Data analysis and statistics

Chromatogram files were processed using ChromasPro v2.1.8. Species identification was

conducted by BLAST search in NCBI database. Reference sequences of the species were

retrieved from GenBank and used to construct a phylogenetic tree, constructed using Mega X

[33].

Water chemistry and cell count data were analysed using R [34]. To compare water chemistry

between the background value and the contaminated water samples, one-tailed Wilcox signed

rank test was used, as the majority of the variables showed non-normal distribution. One-

tailed paired t-test was used for within-sample and overall comparison between plate and

microscopic counts. Comparison between aerobic and anaerobic counts was also done using

one-tailed paired t-test. Some of the data pairs for t-test were $\log_{10}$-transformed to conform to

normal distribution. Data distribution for normality was assessed graphically using histograms

and by Shapiro-Wilk normality test. Statistical significance was inferred at alpha = 0.05.

Results

Groundwater chemistry

Concentrations of most of the solutes in the groundwater have decreased greatly compared to

previous data (Abiriga et al., unpublished), but sulphate and nitrate continue to be leached at

higher amounts. Despite the decrease, most of the solutes still occur at levels significantly

above that of the background value (Table 1). Concentrations of the solutes decreased along

the groundwater flow path from R1 through R2 to R4, (henceforth referred to proximal,

intermediate and distal wells, respectively). Similarity of water chemistry was greatest

between the proximal and intermediate wells.

Table 1. Characteristics of the groundwater chemistry measured between 2018 and 2019.

Values from R0 were used as a benchmark against which values of R1, R2 and R4 were

compared. All units are in mg/l, except for pH (pH units), conductivity (μS/cm) and alkalinity

(mM).

	Mean	Range (min & max)		
	R0 (N=4)	R1 (N=20)	R2 (N=16)	R4 (N=12)
pH	4.9	6.0-7.7 [s]	5.9-7.1 [s]	5.6-6.2 [s]
Conductivity	35	145-251 [s]	102-220 [s]	78-190 [s]
Dissolved oxygen	4.14	0.20-3.42 [s]	0.59-4.79 [s]	1.14-9.12 [n]
Sodium	1.74	4.52-6.91 [s]	4.96-5.63 [s]	3.99-5.69 [s]
Potassium	0.48	5.00-7.52 [s]	4.81-7.49 [s]	6.38-9.92 [s]
Calcium	1.95	24.67-39.18 [s]	17.79-27.44 [s]	10.66-21.74 [s]
Magnesium	0.50	1.80-2.84 [s]	2.69-3.43 [s]	2.68-3.66 [s]
Ammonium	0.00 [a]	0.00-1.91 [s, a]	0.00-0.57 [s, a]	0.00-0.41 [s, a]
Iron	0.02 [d]	0.02-0.02 [d]	0.02-0.02 [d]	0.02-0.09 [n]
Manganese	0.04	0.04-0.65 [s]	0.03-0.31 [s]	0.01-1.42 [s]
Alkalinity	0.05	0.71-1.92 [s]	0.58-1.66 [s]	0.49-1.33 [s]
Sulphate	2.78	8.38-24.62 [s]	7.35-11.86 [s]	5.71-10.85 [s]
Nitrate (as Nitrogen)	1.51	1.26-4.83 [s]	0.38-3.71 [n]	1.28-3.95 [s]
Chloride	2.02	3.37-7.44 [s]	4.34-8.45 [s]	3.62-6.30 [s]
Total nitrogen	1.50	1.62-4.54 [s]	0.84-3.48 [s]	1.20-3.41 [s]

Total organic carbon	4.06	2.68-5.51 [n]	2.90-6.80 [n]	1.82-6.31 [n]

N = number of samples

[s] Values are significantly higher ($P < 0.05$) than the background value, except dissolved

oxygen which is lower than the background value.

[n] non-significant difference ($P > 0.05$)

[a] values below the limit of detection were treated as zero.

[d] Values are half the limit of detection (50 ppb).

Cell density and bacterial species

Selection of growth condition

Bacterial growth was examined on tryptic soy agar, plate count agar and nutrient agar. Among

these three nutrient media, tryptic soy agar gave the highest number of bacterial colonies. Best

growth was obtained with half-strength medium and was thus selected for the isolation of

bacteria. The cell density estimate from the plate count was in the range 1×10^2 - 3.2×10^5

(aerobic incubation) and 0 - 2.4×10^5 (anaerobic incubation) cfu/ml. Despite the comparable

maximum counts from the two growth conditions, anaerobic plate counts were significantly

lower than aerobic plate counts ($t = 3.628$, $df = 37$, $P < 0.05$). Isolates recovered under

anaerobic condition were all facultative anaerobes; the same strains dominated under both

aerobic and anaerobic conditions, however, some strains were isolated only under anaerobic

condition, even though they were facultative anaerobes. Such strains would have been missed

if only aerobic incubation was carryout. Wells in the contamination plume had higher plate

counts than the background well and the distal well had lower plate counts compared to the

intermediate and proximal wells (Fig 1).

Fig 1. Bacterial cell density in the groundwater samples estimated from aerobic plate count

(PC) and direct microscopic count (MC), error bars are mean + standard error. R0 is the

background well located in a nearby uncontaminated aquifer upstream of the landfill, while the

multilevel wells (R1 – R4) are located downstream of the landfill in the contaminated aquifer

along groundwater flow direction.

Selective culture for sulphate-reducing, sulphide-oxidising and iron-oxidising bacteria was

also attempted. Growth was observed in sulphate-reducing and sulphide-oxidising conditions,

although strains isolated under sulphate-reducing condition were non-sulphidogenic. No

growth was observed on iron-oxidising medium, except fungal mycelium and yeast.

Direct microscopic count

Microscopic counts were in the range 7×10^3 - 3.5×10^5 cells/ml. Cell morphologies observed

under fluorescence microscopy included small rods, long large rods, coccobacilli and vibrios.

Small rods were most frequently encountered. Water samples from the proximal and

intermediate wells apparently were dominated by short rods, vibrios and elongated narrow

rods, some occurring in chains. The distal well on the other hand, did not exhibit

predominance of any specific cell types, an observation that agrees with the more diverse

colony morphology observed in samples from this well.

There was a tendency of decreasing cell density with depth in the proximal well, while the

intermediate well showed an increase with depth. The distal well which showed very low

plate counts gave higher microscopic count. Within-sample comparison indicates that

microscopic counts were in all but one case higher than plate counts (Fig 1), but only

226 significantly so in 4 of the 13 samples ($P < 0.05$). Overall, microscopic count was higher than

227 plate count ($t = 6.935$, $df = 51$, $P < 0.05$) (S1 Fig).

Species composition

Small subunit rRNA gene sequencing of pure isolates revealed higher species richness in the

wells located in the contamination plume than in the background well (Fig 2). The distal well

had the highest species richness. Class- and family-level classifications are provided in S2 and

S3 Fig. The dominant phylum in the aquifer was *Proteobacteria*, representing 55.2% of the

total isolates. *Gamma (γ)-proteobacteria* and *Beta (β)-proteobacteria* were detected in all the

wells, while *Alpha (α)-proteobacteria* were detected only in the proximal and intermediate

wells (S2 Fig). Phylum *Bacteroidetes* appears to be enriched in the first two levels of the

proximal well. Forty-six genera and 87 species were identified (Fig 2, S1 Table and S4 Fig).

Genus *Pseudomonas* was the most abundant taxon (S5 Fig).

Fig 2. Bacterial genera recovered from the groundwater samples. Species in R0 represents

biodiversity of the uncontaminated aquifer located upstream of the landfill. Species

composition of the multilevel wells R1-R4 show biodiversity of the contaminated aquifer at

the proximal, intermediate and distal positions, respectively.

Of the eighty-seven species isolated, 7 were found only in the background aquifer, 77 found

only in the contaminated aquifer and three were found in both. These three species were

Rhodoferax ferrireducens and *Pseudomonas* sp. (2) detected in the background and distal

wells, and *Rugamonas* sp. detected in the background, intermediate and distal wells (S1

Table). As shown in Fig 3, 28 species were unique to the distal well. No single species was

found in all the four wells, although two genera (*Pseudomonas* and *Janthinobacterium*) were

common. Among the wells located in the contamination plume, only two species i.e.

Pseudomonas veronii and *Rhodococcus degradans* were found in all of them. The proximal-

intermediate was the most similar pair of wells, with 10 species in common.

Fig 3. Venn diagram showing relationships among the groundwater samples at genus-level

classification **(A)** and species-level classification **(B)**. R0 is the background well located

upstream of the landfill in a nearby aquifer, while R1, R2 and R4 are wells located

downstream of the landfill in the contamination plume along the groundwater flow direction.

Discussion

Cell density and species richness

Cell density estimates with plate count were generally lower than microscopic count, which is

consistent with the literature [28, 35]. However, this difference was not much greater than a

factor of two in most cases, and only in four of the thirteen samples was the difference

significant. This indicates that in most samples, a large proportion of the microbial population

is culturable. Presumably, the nutrients in the landfill leachate favour the growth of culturable

heterotrophic microorganisms. Microscopic examination of the water samples from the

proximal and intermediate wells showed a limited diversity of cell types, which is consistent

with the plate cultures, which were in most cases dominated by colonies of a few species:

Pseudomonas veronii, *Rhodococcus degradans*, *Pedobacter, Brevundimonas* and

Polaromonas.

The background aquifer has a low nutrient content and both the cell density and the microbial

diversity are low, although comparison of microscopic and plate counts suggest that a fairly

large proportion of the population is non-culturable. The contaminated aquifer, on the other

hand, is nutrient-rich and supports a denser and more diverse population. A low cell density

was also encountered at the deepest level of the proximal well (R105), although this location

is not nutrient-deficient. In this case, the low cell density is probably due to filtering effects of

the sediment, which, at this level is expected to contain much fine-grained material from the

till layer.

Groundwater quality was found to improve with distance from the landfill (Table 1), and this

was accompanied by an increase in biodiversity from proximal to distal wells. This may be

analogous to increase in diversity and community stability as leachate becomes less

contaminated over time [11]. We suggest that close to the landfill, only species resistant to the

toxic effects of the leachate are able to survive and grow, while at more distant locations,

toxic pollutants become attenuated by biodegradation, precipitation, sorption and dilution,

allowing the growth of more sensitive species. This might explain the observation in Fig 3,

where only 2 species were shared among the proximal, intermediate and distal wells, 10

species shared between the proximal and intermediate wells, and 28 species exclusive to the

distal well. This also suggests that the conditions prevailing in the proximal and intermediate

wells are relatively similar (which is reflected by the chemistry), thereby selecting for similar

taxa. Additional supporting information can be accessed from the online supplementary

material (S6 Fig).

Species composition and their bioremediation potential

Phylum *Actinobacteria*

The *Actinobacteria* in the genus *Arthrobacter* have been investigated for their potential in

biodegradation of xenobiotics and other harmful compounds [36]. Six known species and two

unidentified species of *Arthrobacter* were detected in Revdalen samples. The "*A. oxydans*

group" (*Pseudarthrobacter phenanthrenivorans, P. defluvii, P. sulfonivorans* and *P.*

siccitolerans), "*A. psychrolactophilus* group" (*A. livingstonensis*), and the "*A. globiformis*

group" (*A. globiformis*). *P. defluvii* is a 4-chlorophenol degrading microorganism [37], *P.*

phenanthrenivorans and *A. globiformis* are phenanthrene degrading bacteria [38, 39], and *P.*

sulfonivorans is dimethylsulfone degrading [40]. *P. siccitolerans,* and *A. livingstonensis* have

no known bioremediation potentials, except biogeochemical cycling, e.g. sulphate reduction

by *A. livingstonensis* and nitrate reduction by *P. siccitolerans* (S2 Table). Congeners of

polycyclic aromatic hydrocarbons (PAHs) have been detected in the contaminated

groundwater, albeit in minute concentrations (Abiriga et al., unpublished). This could have

contributed to the establishment of the *Arthrobacter* species with specialised metabolic

potentials.

Members of genus *Rhodococcus* (family *Nocardiaceae*) are widely distributed in various

types of environments including both pristine and hydrocarbon-contaminated soil, marine

sediments, sludge, freshwater and saltwater [41, 42]. The presence of aerobic alkane-

degrading *alkB* gene in almost every member of the genus have made *Rhodococcus*

biotechnologically valuable microorganisms [41]. Among the *Rhodococcus* isolated from

Revdalen Aquifer was *R. erythropolis*. This species possesses remarkable bioremediation

potential, being capable of degrading a wide range of halogenated hydrocarbons including

polychlorobiphenyls [36, 43], diesel oil, *normal-*, *cis-* and *cyclo*paraffins and aromatic

compounds e.g. naphthalene [44, 45]. Another *Rhodococcus* species recovered was *R.*

degradans, a species capable of degrading several complex organic compounds including

polychlorinated biphenyls (PCBs) and PAHs [42]. This was the dominant species in phylum

Actinobacteria and was recovered from all the wells in the contaminated aquifer. Other

species included *R. sovatensis*, so far with no reported bioremediation ability, but by the fact that the type species was characterised based on *alkB* gene [41] is an indication of its potential in biodegradation.

Other members of *Actinobacteria* recovered belong to family *Microbacteriaceae*, comprising of *Cryobacterium psychrotolerans, C. articum, Microbacterium hydrocarbonoxydans, M. maritypicum, Microtericola viridarii, Curtobacterium flaccumfaciens, Plantibacter auratus*, and *Leifsonia* spp. Bioremediation capabilities include degradation of alkane by *Cryobacterium*, toluene by *Plantibacter* [46], and crude-oil by *M. hydrocarbonoxydans* and *Leifsonia* sp. [47, 48].

Family *Nocardioidaceae* and *Streptomycetaceae* within clade *Actinobacteria* are each represented by a single species of respectively, *Aeromicrobium fastidiosum* and *Streptomyces* sp. No bioremediation potential have been described for either species yet. The latter might influence microbial interaction since the genus is a renowned biocin producer.

Phylum *Bacteroidetes*

Phylum *Bacteroidetes* is represented by *Sphingobacteriia, Cytophagia* and *Flavobacteria*. Members of *Bacteroidetes* may be strict or facultative anaerobes with hydrolytic and fermentative metabolism [11], but so far the known hydrocarbon degraders are aerobes [49].

Seven species of the genus *Pedobacter* (*Sphingobacteriia*) were recovered. These included *P. cryoconitis, P. africana, P. quisquiliarum, P. lusitanus, P. kyunghensis, P. panaciterae*, and *P. trunci*. The strains were originally isolated from respectively, alpine glacier cryoconite [50]; soil and activated sludge [51]; activated sludge [52]; deactivated uranium mine sludge [53]; soil of ginseng field [54]; soil [55]; and tree trunk [56]. *P. cryoconitis* was found to be capable of degrading diesel oil [50], a feature that is of importance in bioremediation. The fact

that some of the isolates were discovered from sludge suggests that they are capable of

biodegradation.

Members of class *Flavobacteria* detected included *F. aquidurense*, *Flavobacterium* sp.,

Cryseobacterium lactis and *C. viscerum*. Strains of genus *Flavobacterium* are capable of

degrading biomacromolecules [57], e.g. biphenyl and phenanthrene [58]. The type species of

Cryseobacterium lactis and *C. viscerum* were originally isolated from milk and diseased fish

respectively; to our knowledge this is the first report of their isolation from environmental

samples. Class *Cytophagia* comprised of two species of *Dyadobacter hamtensis* and

Dyadobacter sediminis, neither of which have documented bioremediation potentials.

Phylum *Proteobacteria*

Of the three classes of *Proteobacteria* found, *β-proteobacteria* was only represented by order

Burkholderiales, while *α-* and *γ-proteobacteria* were composed of three and four orders,

respectively: *Caulobacterales*, *Rhizobiales*, *Sphingomonadales*, *Aeromonadales*,

Enterobacterales, *Pseudomonadales* and *Xanthomonadales*.

Although members of *β-proteobacteria* may grow more slowly than pseudomonads, they may

be more abundant and efficient degraders in natural environments [59]. This is consistent with

our finding that order *Burkholderiales* was more abundant (40%) than *Pseudomonadales*

(23%). Its members constitute those with more versatile catabolic potentials than

Pseudomonadales. Members of family *Burkholderiaceae* with biotechnological relevance

were *Paraburkholderia xenovorans* and *Burkholderia pseudomultivorans*, the latter being

recovered from the background well. *P. xenovorans* is a PCB-degrading bacterium originally

isolated from PCB-contaminated landfill soil [60], and so far the best-known aerobic PCB

degrader [61]. However, PCBs have not been detected at Revdalen (Abiriga et al.,

unpublished). *B. pseudomultivorans* might be speculated to involve in biodegradation, given

that it is a member of the renowned *Burkholderia cepacia* complex [62]. Although such

degradation would only involve naturally occurring organic matter.

The number of taxa from family *Comamonadaceae* was greater than from *Burkholderiaceae.*

It included genus *Polaromonas,* which frequently inhabits cold oligotrophic environments and

was represented by four species: *P. naphthalenivorans, P. aquatica, P. jejuensis* and *P.*

vacuolata. Bioremediation potentials include degradation of biphenyl and benzoate by *P.*

naphthalenivorans [63, 64] and pyrene by *P. vacuolata* [65]. Another genus was

Janthinobacterium, represented by *J. lividum*, *J. svalbardensis*, and *J. agaricidamnosum*. The

last species has been found associated with a consortium involved in degradation of

phenanthrene [39]. Still among members of *Comamonadaceae* is *Simplicispira psychrophila*,

a species originally recovered from wastewater [66], which suggests its ability to biodegrade.

Lastly, *Variovorax boronicumulans* is a species known to bioaccumulate boron [67], but quite

recently, a *Variovorax* sp. was found to be responsible for biodegradation of benzene in a

coal-tar contaminated aquifer [68].

Isolates of *Pseudomonas* (*γ*-proteobacteria) included *P. silesiensis, P. veronii, P. brenneri, P.*

migulae, P. koreensis, P. frederiksbergensis, P. salomonii and *Pseudomonas* spp. *P.*

frederiksbergensis, and *P. salomonii* were recovered from the background well. Member of

the genus *Pseudomonas* are ubiquitous bacteria inhabiting a variety of environmental niches,

and are metabolically versatile microorganisms capable of utilising a wide range of simple, as

well as complex organic compounds [69, 70]. Genome analysis of *P. silesiensis* has revealed

bioremediation potential [70], while *P. veronii* is naphthalene and pentachlorophenol

degrading [71, 72]. *P. frederiksbergensis* is phenanthrene degrading [73], which illustrates the

intrinsic potential of native microorganisms to decontaminate environments, as the species

was recovered from the upstream background well. The other *Pseudomonas* species are normal freshwater microbes, except *P. salomonii,* which is a plant pathogen.

Tolumonas sp., a member of order *Aeromonadales* in *γ-proteobacteria* was also recovered from the water samples. Its closest relative *T. auensis* can transform phenylalanine, phenyl-acetate, phenyl-lactate and phenyl-pyruvate to toluene, and transform tyrosine to phenol [74].

Members of order *Xanthomonadales* recovered included *Luteibacter rhizovicinus*, *Rhodanobacter denitrificans* and *Rhodanobacter* sp. Bioremediation potential include phenanthrene degradation by an unidentified member of *Luteibacter* [75], while *R. denitrificans* as the name implies, could be involved in denitrification processes in the aquifer. Another study [76] reported *Rhodanobacter*-enriched microbial community in a metal-contaminated aquifer, implying high resistance to metal toxicity by members of the genus. Further, unidentified members of *Rhodanobacter* have been recovered from a Benzo[a]pyrene-degrading consortium [77]. Still within order *Xanthomonadales*, genus *Stenotrophomonas* was represented by four species. These included the clinically relevant and pyrene degrading *S. maltophilia* [78], *S. rhizophila* with antifungal activity [79], *S. tumulicola*, and unidentified *Stenotrophomonas* sp.

Order *Enterobacterales* included *Klebsiella aerogenes*, *Rouxiella badensis*, *Ewingella americana*, *Erwinia billingiae* and *Rahnella woolbedingensis*. While scientific interest in this order chiefly focuses on its medical significance, unidentified members of *Rahnella* and *Klebsiella* have been implicated in bioremediation of respectively, naphthalene [80] and crude oil [81].

Class *α-proteobacteria* were represented by species in four genera: *Brevundimonas mediterranea, Methylobacterium, Sphingomonas*, and *Phyllobacterium myrsinacearum. B.*

mediterranea is naphthalene degrading [71]. *Methylobacterium* sp. have been implied in

trichloroethene bioremediation [82]. *Sphingomonas* was represented by *S. echinoides* and an

unidentified *Sphingomonas* sp. which closely resembles *S. faeni*. Bioremediation potential of

genus *Sphingomonas* include degradation of trichloroethene [82], phenanthrene [39] and

dibenzofuran [83]. *P. myrsinacearum* has no known bioremediation potential, but may be

relevant for nutrient cycling particularly nitrogen and sulphur.

Phylum *Firmicutes*

418 The phylum *Firmicutes* was represented by three species; *Exiguobacterium acetylicum,*

419 *Staphylococcus warneri* and *Sporosarcina aquimarina*. *S. warneri* has been implicated in

420 degradation of diesel oil [84], while *E. acetylicum* as the name suggests degrades metabolic

421 intermediate products such as acetate. No literature is available on bioremediation potential of

422 *S. aquimarina*.

Biogeochemical transformation

Attempt to cultivate iron oxidising bacteria was unsuccessful, but a potential iron reducer,

Rhodoferax ferrireducens was recovered on nutrient medium. This species is also capable of

respiring on manganese and nitrate [85]. While iron levels were very low and often below the

limit of detection, manganese and nitrate were measurable. The maximum manganese

coincided with the presences of *R. ferrireducens* and principal component analysis showed a

positive correlation between this species and manganese (S6 Fig). This suggests that the

species is actively involved in biogeochemical cycling of manganese, and possibly nitrate.

For sulphur cycling, growth was observed under sulphide-oxidising and sulphate-reducing

conditions. Growth of sulphide-oxidising bacteria, such as *B. mediterranea*, which showed

high abundance in wells with higher sulphate concentration, is suggestive of *in situ* sulphide

oxidation. With two exceptions, all sulphate-reducing bacteria were isolated from the

proximal and intermediate wells. The low dissolved oxygen levels in these wells (<1 mg/L)

would be conducive for sulphate reduction, although the low levels of TOC may limit its

extent. Additional information on sulphur and nitrogen cycling potential of the strains is given

in S2 Table.

Strains with other nutrient cycling potentials included; *Herbaspirillum autotrophicum*, a

bacterium capable of using carbon dioxide as a sole carbon source [86], *Herminiimonas*

arsenicoxydans, a well-known biogeochemical cycler that transforms arsenite to arsenate

[87].

Conclusion

This study shows that nearly half of the culturable bacterial strains from a landfill-

contaminated confined aquifer had bioremediation potential. Preliminary results (Khan,

unpublished) suggest that some of these strains have the ability to grow on low density

polyethylene. Comparison of microscopic cell counts and plate counts indicated that the

strains in the most contaminated parts of the aquifer were mostly culturable heterotrophs,

while a higher proportion of non-culturable strains was present in the distal well, where

contamination was most attenuated. Biodiversity was lowest in the uncontaminated well,

moderate in the proximal and intermediate wells and highest in the distal well. The higher

biodiversity in the distal wells is possibly as a result of plentiful residual nutrients and low

levels of toxic contaminants. Each well had a distinctive bacterial flora, but the proximal and

intermediate wells seemed to be ecologically related as they had more taxa in common. There

was a clear difference between the flora of the contaminated wells and the uncontaminated

456 well. A more comprehensive characterisation of species by DNA metabarcoding is currently

457 underway.

Acknowledgement

459 We thank Frode Bergan and Tom Age Aarnes for their participation in fieldworks, and Karin

460 Brekke Li for technical assistance in water chemistry analysis.

References

1.   Mays D, Scheibe T. Groundwater Contamination, Subsurface Processes, and
Remediation Methods: Overview of the Special Issue of Water on Groundwater
Contamination and Remediation. Multidisciplinary Digital Publishing Institute; 2018.

2.   O'Connor D, Hou D, Ok YS, Song Y, Sarmah AK, Li X, et al. Sustainable in situ
remediation of recalcitrant organic pollutants in groundwater with controlled release
materials: A review. Journal of Controlled Release. 2018;283:200-13.

3.   Pous N, Balaguer MD, Colprim J, Puig S. Opportunities for groundwater microbial
electro-remediation. Microbial biotechnology. 2018;11(1):119-35.

4.   Reinhard M, Barker JF, Goodman NL. Occurrence and distribution of organic chemicals
in two landfill leachate plumes. Environmental Science & Technology. 1984;18(12):953-
61.

5.   Eggen T, Moeder M, Arukwe A. Municipal landfill leachates: a significant source for
new and emerging pollutants. Science of the Total Environment. 2010;408(21):5147-57.

6.   Hyldegaard BH, Jakobsen R, Weeth EB, Overheu ND, Gent DB, Ottosen LM. Challenges
in electrochemical remediation of chlorinated solvents in natural groundwater aquifer
settings. Journal of hazardous materials. 2019;368:680-8.

7.   Logeshwaran P, Megharaj M, Chadalavada S, Bowman M, Naidu R. Petroleum
hydrocarbons (PH) in groundwater aquifers: An overview of environmental fate, toxicity,
microbial degradation and risk-based remediation approaches. Environmental technology
& innovation. 2018;10:175-93.

8.   Nunes OC, Lopes AR, Manaia CM. Microbial degradation of the herbicide molinate by
defined cultures and in the environment. Applied microbiology and biotechnology.
2013;97(24):10275-91.

9.   Dlugonski J. Microbial biodegradation : from omics to function and application. Norfolk:
Caister Academic Press; 2016.

10.   Alfreider A, Vogt C, Babel W. Microbial diversity in an in situ reactor system treating
488   onochlorobenzene contaminated groundwater as revealed by 16S ribosomal DNA
489   analysis. Systematic and Applied Microbiology. 2002;25(2):232-40.

490   11.   Köchling T, Sanz JL, Gavazza S, Florencio L. Analysis of microbial community structure
491   and composition in leachates from a young landfill by 454 pyrosequencing. Applied
492   microbiology and biotechnology. 2015;99(13):5657-68.

493   12.   Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ, et al. Thousands
494   of microbial genomes shed light on interconnected biogeochemical processes in an
495   aquifer system. Nature communications. 2016;7.

496   13.   Cozzarelli IM, Suflita JM, Ulrich GA, Harris SH, Scholl MA, Schlottmann JL, et al.
497   Geochemical and microbiological methods for evaluating anaerobic processes in an
498   aquifer contaminated by landfill leachate. Environmental science & technology.
499   2000;34(18):4025-33.

500   14.   Harvey RW, Smith RL, George L. Effect of organic contamination upon microbial
501   distributions and heterotrophic uptake in a Cape Cod, Mass., aquifer. Applied and
502   Environmental Microbiology. 1984;48(6):1197-202.

503   15.   Hug LA, Thomas BC, Brown CT, Frischkorn KR, Williams KH, Tringe SG, et al. Aquifer
504   environment selects for microbial species cohorts in sediment and groundwater. The
505   ISME journal. 2015;9(8):1846-56.

506   16.   Kleikemper J, Pombo SA, Schroth MH, Sigler WV, Pesaro M, Zeyer J. Activity and
507   diversity of methanogens in a petroleum hydrocarbon-contaminated aquifer. Applied and
508   environmental microbiology. 2005;71(1):149-58.

509   17.   Ludvigsen L, Albrechtsen H-J, Ringelberg D, Ekelund F, Christensen T. Distribution and
510   composition of microbial populations in a landfill leachate contaminated aquifer
511   (Grindsted, Denmark). Microbial Ecology. 1999;37(3):197-207.

512   18.   Röling W, Van Breukelen B, Braster M, Goeltom M, Groen J, Van Verseveld H. Analysis
513   of microbial communities in a landfill leachate polluted aquifer using a new method for
514   anaerobic physiological profiling and 16S rDNA based fingerprinting. Microbial
515   Ecology. 2000;40(3):177-88.

516   19.   Tischer K, Zeder M, Klug R, Pernthaler J, Schattenhofer M, Harms H, et al. Fluorescence
517   in situ hybridization (CARD-FISH) of microorganisms in hydrocarbon contaminated
518   aquifer sediment samples. Systematic and applied microbiology. 2012;35(8):526-32.

519   20.   Watanabe K, Kodama Y, Hamamura N, Kaku N. Diversity, abundance, and activity of
520   archaeal populations in oil-contaminated groundwater accumulated at the bottom of an
521   underground crude oil storage cavity. Applied and environmental microbiology.
522   2002;68(8):3899-907.

523   21.   Meckenstock RU, Elsner M, Griebler C, Lueders T, Stumpp C, Aamand J, et al.
524   Biodegradation: updating the concepts of control for microbial cleanup in contaminated
525   aquifers. ACS Publications; 2015.

22. He Z, Zhang P, Wu L, Rocha AM, Tu Q, Shi Z, et al. Microbial functional gene diversity
predicts groundwater contamination and ecosystem functioning. MBio.
2018;9(1):e02435-17.

23. Dojka MA, Hugenholtz P, Haack SK, Pace NR. Microbial diversity in a hydrocarbon-and
chlorinated-solvent-contaminated aquifer undergoing intrinsic bioremediation. Applied
and Environmental Microbiology. 1998;64(10):3869-77.

24. Pickup R, Rhodes G, Alamillo M, Mallinson H, Thornton S, Lerner D. Microbiological
analysis of multi-level borehole samples from a contaminated groundwater system.
Journal of Contaminant Hydrology. 2001;53(3-4):269-84.

25. Rooney-Varga JN, Anderson RT, Fraga JL, Ringelberg D, Lovley DR. Microbial
communities associated with anaerobic benzene degradation in a petroleum-contaminated
aquifer. Applied and Environmental Microbiology. 1999;65(7):3056-63.

26. Kepner R, Pratt JR. Use of fluorochromes for direct enumeration of total bacteria in
environmental samples: past and present. Microbiol Mol Biol Rev. 1994;58(4):603-15.

27. Ňancucheo I, Rowe OF, Hedrich S, Johnson DB. Solid and liquid media for isolating and
cultivating acidophilic and acid-tolerant sulfate-reducing bacteria. FEMS microbiology
letters. 2016;363(10).

28. Gregorich EG, Carter MR. Soil sampling and methods of analysis: CRC press; 2007.

29. Holanda R, Hedrich S, Ňancucheo I, Oliveira G, Grail BM, Johnson DB. Isolation and
characterisation of mineral-oxidising "Acidibacillus" spp. from mine sites and
geothermal environments in different global locations. Research in microbiology.
2016;167(7):613-23.

30. Csuros M, Csuros C, Ver K. Microbiological examination of water and wastewater. Boca
Raton, Fla: Lewis Publishers; 1999.

31. Lane DJ, Pace B, Olsen GJ, Stahl DA, Sogin ML, Pace NR. Rapid determination of 16S
ribosomal RNA sequences for phylogenetic analyses. Proceedings of the National
Academy of Sciences. 1985;82(20):6955-9.

32. Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, et al. Evaluation of
general 16S ribosomal RNA gene PCR primers for classical and next-generation
sequencing-based diversity studies. Nucleic acids research. 2013;41(1):e1-e.

33. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: molecular evolutionary
genetics analysis across computing platforms. Molecular biology and evolution.
2018;35(6):1547-9.

34. R-Core-Team. R: A language and environment for statistical computing. R Foundation
for Statistical Computing. Vienna, Austria: https://www.R-project.org/; 2019.

35. Muyzer G, Smalla K. Application of denaturing gradient gel electrophoresis (DGGE) and
temperature gradient gel electrophoresis (TGGE) in microbial ecology. Antonie van
Leeuwenhoek. 1998;73(1):127-41.

36. Goodfellow M, Garrity GM, Bergey DH. Bergey's manual of systematic bacteriology :
Vol. 5 Pt. B : The Actinobacteria. 2nd ed. ed. New York: Springer; 2012.

37. Kim KK, Lee KC, Oh H-M, Kim MJ, Eom MK, Lee J-S. Arthrobacter defluvii sp. nov.,
4-chlorophenol-degrading bacteria isolated from sewage. International journal of
systematic and evolutionary microbiology. 2008;58(8):1916-21.

38. Kallimanis A, Kavakiotis K, Perisynakis A, Spröer C, Pukall R, Drainas C, et al.
Arthrobacter phenanthrenivorans sp. nov., to accommodate the phenanthrene-degrading
bacterium Arthrobacter sp. strain Sphe3. International journal of systematic and
evolutionary microbiology. 2009;59(2):275-9.

39. Bodour AA, Wang JM, Brusseau ML, Maier RM. Temporal change in culturable
phenanthrene degraders in response to long-term exposure to phenanthrene in a soil
column system. Environmental Microbiology. 2003;5(10):888-95.

40. Borodina E, Kelly DP, Schumann P, Rainey FA, Ward-Rainey NL, Wood AP. Enzymes
of dimethylsulfone metabolism and the phylogenetic characterization of the facultative
methylotrophs Arthrobacter sulfonivorans sp. nov., Arthrobacter methylotrophus sp.
nov., and Hyphomicrobium sulfonivorans sp. nov. Archives of microbiology.
2002;177(2):173-83.

41. Táncsics A, Máthé I, Benedek T, Tóth EM, Atasayar E, Spröer C, et al. Rhodococcus
sovatensis sp. nov., an actinomycete isolated from the hypersaline and heliothermal Lake
Ursu. International journal of systematic and evolutionary microbiology. 2017;67(2):190-
6.

42. Švec P, Černohlávková J, Busse H-J, Vojtková H, Pantu R, Cnockaert M, et al.
Classification of strain CCM 4446T as Rhodococcus degradans sp. nov. International
journal of systematic and evolutionary microbiology. 2015;65(12):4381-7.

43. Armfield SJ, Sallis PJ, Baker PB, Bull AT, Hardman DJ. Dehalogenation of haloalkanes
byRhodococcus erythropolis Y2. Biodegradation. 1995;6(3):237-46.

44. Laczi K, Kis Á, Horváth B, Maróti G, Hegedüs B, Perei K, et al. Metabolic responses of
Rhodococcus erythropolis PR4 grown on diesel oil and various hydrocarbons. Applied
microbiology and biotechnology. 2015;99(22):9745-59.

45. Meyer S, Moser R, Neef A, Stahl U, Kämpfer P. Differential detection of key enzymes
of polyaromatic-hydrocarbon-degrading bacteria using PCR and gene probes.
Microbiology. 1999;145(7):1731-41.

46. Giudice AL, Casella P, Caruso C, Mangano S, Bruni V, De Domenico M, et al.
Occurrence and characterization of psychrotolerant hydrocarbon-oxidizing bacteria from
surface seawater along the Victoria Land coast (Antarctica). Polar biology.
2010;33(7):929-43.

47. Schippers A, Bosecker K, Spröer C, Schumann P. Microbacterium oleivorans sp. nov.
and Microbacterium hydrocarbonoxydans sp. nov., novel crude-oil-degrading Gram-
positive bacteria. International journal of systematic and evolutionary microbiology.
2005;55(2):655-60.

48.   Dashti N, Khanafer M, El-Nemr I, Sorkhoh N, Ali N, Radwan S. The potential of oil-
605   utilizing bacterial consortia associated with legume root nodules for cleaning oily soils.
606   Chemosphere. 2009;74(10):1354-9.

607   49.   Prince RC, Amande TJ, McGenity TJ. Prokaryotic hydrocarbon degraders. Taxonomy,
608   Genomics and Ecophysiology of Hydrocarbon-Degrading Microbes. 2018:1-41.

609   50.   Margesin R, Spröer C, Schumann P, Schinner F. Pedobacter cryoconitis sp. nov., a
610   facultative psychrophile from alpine glacier cryoconite. International Journal of
611   Systematic and Evolutionary Microbiology. 2003;53(5):1291-6.

612   51.   Steyn P, Segers P, Vancanneyt M, Sandra P, Kersters K, Joubert J. Classification of
613   heparinolytic bacteria into a new genus, Pedobacter, comprising four species: Pedobacter
614   heparinus comb. nov., Pedobacter piscium comb. nov., Pedobacter africanus sp. nov. and
615   Pedobacter saltans sp. nov. proposal of the family Sphingobacteriaceae fam. nov.
616   International Journal of Systematic and Evolutionary Microbiology. 1998;48(1):165-77.

617   52.   Zhang B, Liu Z-Q, Zheng Y-G. Pedobacter quisquiliarum sp. nov., isolated from activated
618   sludge. International journal of systematic and evolutionary microbiology.
619   2017;68(1):438-42.

620   53.   Covas C, Caetano T, Cruz A, Santos T, Dias L, Klein G, et al. Pedobacter lusitanus sp.
621   nov., isolated from sludge of a deactivated uranium mine. International journal of
622   systematic and evolutionary microbiology. 2017;67(5):1339-48.

623   54.   Yang J-E, Shin J-Y, Park S-Y, Mavlonov GT, Yi E-J, Lee E-H, et al. Pedobacter
624   kyungheensis sp. nov., with ginsenoside converting activity. The Journal of general and
625   applied microbiology. 2012;58(4):309-16.

626   55.   Yoon M-H, Ten LN, Im W-T, Lee S-T. Pedobacter panaciterrae sp. nov., isolated from
627   soil in South Korea. International journal of systematic and evolutionary microbiology.
628   2007;57(2):381-6.

629   56.   Du J, Singh H, Ngo HT, Won K-H, Kim K-Y, Yi T-H. Pedobacter daejeonensis sp. nov.
630   and Pedobacter trunci sp. nov., isolated from an ancient tree trunk, and emended
631   description of the genus Pedobacter. International journal of systematic and evolutionary
632   microbiology. 2015;65(4):1241-6.

633   57.   Jean-François B, John PB. Bergey's Manual of Systematic Bacteriology (Genus
634   *Flavobacterium)*. 2nd ed ed. New York: Springer; 2011.

635   58.   Stucki G, Alexander M. Role of dissolution rate and solubility in biodegradation of
636   aromatic compounds. Appl Environ Microbiol. 1987;53(2):292-7.

637   59.   Tan WA, Parales RE. Hydrocarbon Degradation by Betaproteobacteria. Taxonomy,
638   Genomics and Ecophysiology of Hydrocarbon-Degrading Microbes. 2019:1-18.

639   60.   Goris J, De Vos P, Caballero-Mellado J, Park J, Falsen E, Quensen III JF, et al.
640   Classification of the biphenyl-and polychlorinated biphenyl-degrading strain LB400T
641   and relatives as Burkholderia xenovorans sp. nov. International journal of systematic and
642   evolutionary microbiology. 2004;54(5):1677-81.

61. Chain PS, Denef VJ, Konstantinidis KT, Vergez LM, Agulló L, Reyes VL, et al.
Burkholderia xenovorans LB400 harbors a multi-replicon, 9.73-Mbp genome shaped for
versatility. Proceedings of the National Academy of Sciences. 2006;103(42):15280-7.

62. Peeters C, Zlosnik JE, Spilker T, Hird TJ, LiPuma JJ, Vandamme P. Burkholderia
pseudomultivorans sp. nov., a novel Burkholderia cepacia complex species from human
respiratory samples and the rhizosphere. Systematic and applied microbiology.
2013;36(7):483-9.

63. Jeon CO, Park W, Ghiorse WC, Madsen EL. Polaromonas naphthalenivorans sp. nov., a
naphthalene-degrading bacterium from naphthalene-contaminated sediment.
International journal of systematic and evolutionary microbiology. 2004;54(1):93-7.

64. Yagi JM, Sims D, Brettin T, Bruce D, Madsen EL. The genome of Polaromonas
naphthalenivorans strain CJ2, isolated from coal tar-contaminated sediment, reveals
physiological and metabolic versatility and evolution through extensive horizontal gene
transfer. Environmental microbiology. 2009;11(9):2253-70.

65. Eriksson M, Dalhammar G, Mohn WW. Bacterial growth and biofilm production on
pyrene. FEMS microbiology ecology. 2002;40(1):21-7.

66. Grabovich M, Gavrish E, Kuever J, Lysenko AM, Podkopaeva D, Dubinina G. Proposal
of Giesbergeria voronezhensis gen. nov., sp. nov. and G. kuznetsovii sp. nov. and
reclassification of [Aquaspirillum] anulus,[A.] sinuosum and [A.] giesbergeri as
Giesbergeria anulus comb. nov., G. sinuosa comb. nov. and G. giesbergeri comb. nov.,
and [Aquaspirillum] metamorphum and [A.] psychrophilum as Simplicispira
metamorpha gen. nov., comb. nov. and S. psychrophila comb. nov. International journal
of systematic and evolutionary microbiology. 2006;56(3):569-76.

67. Miwa H, Ahmed I, Yoon J, Yokota A, Fujiwara T. Variovorax boronicumulans sp. nov.,
a boron-accumulating bacterium isolated from soil. International journal of systematic
and evolutionary microbiology. 2008;58(1):286-9.

68. Posman KM, DeRito CM, Madsen EL. Benzene degradation by a Variovorax species
within a coal tar-contaminated groundwater microbial community. Appl Environ
Microbiol. 2017;83(4):e02658-16.

69. Baïda N, Yazourh A, Singer E, Izard D. Pseudomonas brenneri sp. nov., a new species
isolated from natural mineral waters. Research in microbiology. 2001;152(5):493-502.

70. Kaminski MA, Furmanczyk EM, Sobczak A, Dziembowski A, Lipinski L. Pseudomonas
silesiensis sp. nov. strain A3T isolated from a biological pesticide sewage treatment plant
and analysis of the complete genome sequence. Systematic and applied microbiology.
2018;41(1):13-22.

71. Martirani-Von Abercron SM, Marín P, Solsona-Ferraz M, Castañeda-Cataña MA,
Marqués S. Naphthalene biodegradation under oxygen-limiting conditions: community
dynamics and the relevance of biofilm-forming capacity. Microbial biotechnology.
2017;10(6):1781-96.

72.   Nam I-H, Chang Y-S, Hong H-B, Lee Y-E. A novel catabolic activity of Pseudomonas
veronii in biotransformation of pentachlorophenol. Applied Microbiology and
Biotechnology. 2003;62(2-3):284-90.

73.   Andersen SM, Johnsen K, Sørensen J, Nielsen P, Jacobsen CS. Pseudomonas
frederiksbergensis sp. nov., isolated from soil at a coal gasification site. International
Journal of Systematic and Evolutionary Microbiology. 2000;50(6):1957-64.

74.   Fischer-Romero C, Tindall B, Jüttner F. Tolumonas auensis gen. nov., sp. nov., a toluene-
producing bacterium from anoxic sediments of a freshwater lake. International Journal of
Systematic and Evolutionary Microbiology. 1996;46(1):183-8.

75.   Muangchinda C, Pansri R, Wongwongsee W, Pinyakong O. Assessment of polycyclic
aromatic hydrocarbon biodegradation potential in mangrove sediment from D on H oi L
ot, S amut S ongkram P rovince, T hailand. Journal of applied microbiology.
2013;114(5):1311-24.

76.   Carlson HK, Price MN, Callaghan M, Aaring A, Chakraborty R, Liu H, et al. The
selective pressures on the microbial community in a metal-contaminated aquifer. The
ISME journal. 2019;13(4):937.

77.   Kanaly RA, Harayama S, Watanabe K. Rhodanobacter sp. strain BPC1 in a benzo [a]
pyrene-mineralizing bacterial consortium. Appl Environ Microbiol. 2002;68(12):5826-
33.

78.   Juhasz AL, Stanley G, Britz M. Microbial degradation and detoxification of high
molecular weight polycyclic aromatic hydrocarbons by Stenotrophomonas maltophilia
strain VUN 10,003. Letters in Applied Microbiology. 2000;30(5):396-401.

79.   Wolf A, Fritze A, Hagemann M, Berg G. Stenotrophomonas rhizophila sp. nov., a novel
plant-associated bacterium with antifungal properties. International journal of systematic
and evolutionary microbiology. 2002;52(6):1937-44.

80.   Ma Y, Wang L, Shao Z. Pseudomonas, the dominant polycyclic aromatic
hydrocarbon-degrading bacteria isolated from Antarctic soils and the role of large
plasmids in horizontal gene transfer. Environmental Microbiology. 2006;8(3):455-65.

81.   Odokuma L, Dickson A. Bioremediation of a crude oil polluted tropical rain forest soil.
Global Journal of Environmental Sciences. 2003;2(1):29-40.

82.   Connon SA, Tovanabootr A, Dolan M, Vergin K, Giovannoni SJ, Semprini L. Bacterial
community composition determined by culture-independent and-dependent methods
during propane-stimulated bioremediation in trichloroethene-contaminated groundwater.
Environmental Microbiology. 2005;7(2):165-78.

83.   Halden RU, Halden BG, Dwyer DF. Removal of Dibenzofuran, Dibenzo-p-Dioxin, and
2-Chlorodibenzo-p-Dioxin from Soils Inoculated withSphingomonas sp. Strain RW1.
Appl Environ Microbiol. 1999;65(5):2246-9.

84.   Saadoun I, Al-Akhras M-A, Abu-Ashour J. Bacterial degradation of hydrocarbons as
evidenced by respirometric analysis. Microbios. 1999;100(395):19-25.

85. Finneran KT, Johnsen CV, Lovley DR. Rhodoferax ferrireducens sp. nov., a psychrotolerant, facultatively anaerobic bacterium that oxidizes acetate with the reduction of Fe (III). International Journal of Systematic and Evolutionary Microbiology. 2003;53(3):669-73.

86. Aragno M, SCHLEGEL HG. Aquaspirillum autotrophicum, a new species of hydrogen-oxidizing, facultatively autotrophic bacteria. International Journal of Systematic and Evolutionary Microbiology. 1978;28(1):112-6.

87. Muller D, Simeonova DD, Riegel P, Mangenot S, Koechler S, Lievremont D, et al. Herminiimonas arsenicoxydans sp. nov., a metalloresistant bacterium. International journal of systematic and evolutionary microbiology. 2006;56(8):1765-9.

Supporting information

S1 Fig. Variation in mean bacterial cell count between microscopic and plate counts. Error bars are mean + standard error.

S2 Fig. Bacterial class-level taxonomic group detected from the water samples. R1, R2 and R4 are sampling wells located in the leachate plume, with the corresponding multilevel in each well. R0 is a reference well

S3 Fig. Bacterial family taxonomic group detected from the water samples. R1, R2 and R4 are sampling wells located in the leachate plume, while R0 is a background well.

S4 Fig. Maximum Likelihood tree as implemented in MEGA X based on 16S rRNA gene sequences of bacterial isolates, using Generalised Time Reversible Model. MSA was performed using muscle in MEGA X with default settings. Values at the nodes depicts bootstrap values (%) based on 1000 replications, those >50 are shown.

S5 Fig. Abundance of bacterial genera recovered from the water samples. A total of forty-six genera were recovered.

S6 Fig. PCA of site score as the gradient behind species abundance and water chemistry

variation. The two axes explained 41.7% of the observed variation in the species abundance.

Supplementary table captions

S1 Table. Bacterial species isolated from Revdalen groundwater samples

S2 Table. Biogeochemical cycling (nitrogen and sulphur) capabilities of the bacterial species

represented in Revdalen Aquifer.

Figure

Figure

Figure