1

Mfd regulates RNA polymerase association with hard-to-transcribe regions *in vivo*, especially those with structured RNAs

Mark N. Ragheb[1], Christopher Merrikh[2], Kaitlyn Browning[2], Houra Merrikh[2,*]

[1] Molecular and Cellular Biology Graduate Program and Medical Scientist Training Program, University of Washington, Seattle, WA, USA.
[2] Department of Biochemistry, Vanderbilt University, Nashville, TN 37205, USA

*Corresponding (Lead) author
Houra Merrikh
Email: houra.merrikh@vanderbilt.edu
Tel: 615-343-3846

Keywords: Mfd, RNA polymerase, regulatory RNAs, RNA secondary structure, toxin-antitoxins

Abstract

RNA polymerase (RNAP) encounters various roadblocks during transcription. These obstacles

can impede RNAP movement, influence transcription, ultimately necessitating the activity of

RNAP associated factors. One such factor is the bacterial protein Mfd; a highly conserved DNA

translocase and evolvability factor that interacts with RNAP. Although Mfd is thought to function

primarily in the repair of DNA lesions that stall RNAP, increasing evidence suggests that it may

also be important for transcription regulation. However, this is yet to be fully characterized.

To shed light on Mfd's *in vivo* functions, we identified the chromosomal regions where it

associates. We analyzed Mfd's impact on RNAP association and transcription regulation

genome-wide. We found that Mfd represses RNAP association at many chromosomal regions.

We found that these regions show increased RNAP pausing, suggesting that they are hard-to-

transcribe. Interestingly, we noticed that the majority of the regions where Mfd regulates

transcription contain highly structured regulatory RNAs. The RNAs identified regulate a myriad

of biological processes, ranging from metabolism, to tRNA regulation, to toxin-antitoxin (TA)

functions. We found that transcription regulation by Mfd, at least at some TA loci, is critical for

cell survival. Lastly, we found that Mfd promotes mutagenesis in at least one toxin gene,

suggesting that its function in regulating transcription may promote evolution of certain TA

systems, and other regions containing strong RNA secondary structures. We conclude that Mfd

is an RNAP co-factor that is important, and at times critical, for transcription regulation at hard-

to-transcribe regions, especially those that express structured regulatory RNAs.

Significance

The bacterial DNA translocase Mfd binds to stalled RNAPs and is generally thought to facilitate transcription-coupled DNA repair. Most of our knowledge about Mfd is based on data from biochemical studies. However, little is known about Mfd's function in living cells, especially in the absence of exogenous DNA damage. Here, we show that Mfd modulates RNAP association and alters transcription at a variety of chromosomal loci, especially those containing highly structured, regulatory RNAs. As such, this work improves our understanding of Mfd's function in living cells, and assigns it a new function as a transcription regulator.

Introduction

Timely and efficient transcription is a fundamental requirement for maintaining cellular

homeostasis. Previous work has shown that transcription elongation is discontinuous, with RNA

polymerase (RNAP) processivity being altered by a wide range of obstacles(1, 2). These

impediments vary in severity, from pause sites that slow the rate of RNAP movement(3–5) to

more severe obstacles, such as protein roadblocks and the replication fork. These impediments

can induce reverse translocation of RNAP with respect to both DNA and the nascent RNA

(RNAP backtracking)(6–9). The impact of these roadblocks on RNAP processivity are prevented

and resolved through various mechanisms including the coupling of transcription and

translation, as well as various cellular factors that help re-establish transcription elongation (i.e.

anti-backtracking factors such as GreA)(10).

In vitro work shows that the DNA translocase Mfd utilizes its RNAP binding properties and

forward translocase activity to rescue arrested RNAPs, restoring transcription elongation(11) as

well as promoting transcription termination(12). However, despite decades of research on the

biochemical characteristics of Mfd, the endogenous contexts in which its translocase and anti-

backtracking functions are critical for transcription remain elusive.

Mfd was initially described as a critical DNA repair factor *in vivo* that promotes transcription-

coupled repair (TCR)(13–15). In the TCR pathway, Mfd removes stalled RNAPs at bulky DNA

lesions and promotes nucleotide excision repair (NER) via its UvrA binding capacity (see review

(16)). However, cells lacking Mfd show little to no sensitivity to DNA damaging agents that

promote RNAP stalling(17, 18), especially relative to NER proteins(19, 20). Together with its

high degree of conservation, such data suggests that Mfd may have a broader cellular function

outside of DNA repair.

Mfd was shown to have other roles in the cell, such as regulating catabolite repression in

Bacillus subtilis(21, 22). Recent *in vitro* experiments have shown that Mfd is capable of

autonomously translocating on DNA in the absence of a lesion, but whether this occurs *in vivo* is

unclear(23). Interestingly, Mfd functions as an evolvability factor, promoting mutagenesis and

the rapid evolution of antibiotic resistance in diverse bacterial species(17). Mfd also promotes

mutagenesis during stationary-phase(24, 25) and at chromosomal regions where collisions

between the replication and transcription machineries occur(10). However, a comprehensive *in*

vivo study examining what genomic "hotspots" may be prone to Mfd's mutagenic activity has not

been performed.

Despite our limited understanding of Mfd's cellular functions, its high level of conservation in

bacteria implies a fundamental role for Mfd that may be separate from TCR. In addition, Mfd is

constitutively expressed, suggesting that it may have homeostatic role in transcription

regulation. Further work has recently suggested that Mfd plays a housekeeping function in cells

by associating with RNAP in the absence of exogenous stressors(26). However, the field still

lacks a clear understanding of the conditions and chromosomal features that prompt Mfd

binding and modulation of RNAP association and function.

In this work, we identify the genomic regions of Mfd association in both *B. subtilis* and

Escherichia coli. We also identify the regions where Mfd modulates RNAP association and

transcription. We find that Mfd alters RNAP density predominantly at regions containing highly

structured RNAs in both *B. subtilis* and *E. coli*. Our analysis of RNAP pausing experiments

strongly suggests that these sites are hard-to-transcribe. We find that Mfd regulates the

transcription of genes involved in various cellular functions including toxin-antitoxin (TA)

systems. Importantly, we observe that Mfd's regulatory activity, at least for TA systems, can be

essential: when we overexpress the toxin genes, cell viability is significantly compromised

without Mfd. Lastly, we find that Mfd promotes mutagenesis of at least the *txpA* toxin gene. We

conclude that RNA secondary structure is a major impediment to transcription *in vivo* and that

Mfd is an important RNAP co-factor that regulates transcription and RNAP association with such

loci.

Results

Mapping the genomic loci where Mfd associates

We began by identifying the chromosomal regions where Mfd associates using chromatin

immunoprecipitation followed by high-throughput sequencing (ChIP-seq). We constructed a *B.*

subtilis strain where Mfd is C-terminally Myc-tagged (Fig. S1). Prior work has shown that C-

terminally tagged Mfd retains its functionality *in vivo*(26) – (below we present data that confirms

the functionality of this Mfd-Myc in our system). To identify the chromosomal regions where Mfd

associates, we harvested exponentially growing cells and performed ChIP-seq analysis. We

controlled for potential ChIP artifacts by comparing this signal to ChIP-seq performed using Myc

antibody against *B. subtilis* cells lacking the Myc-tagged Mfd (Fig. 1*A*). Under these conditions,

489 out of 5755 genes analyzed in the *B. subtilis* genome exhibited preferential Mfd association

(defined as one standard deviation greater than the average Mfd ChIP signal across all genes)

(Dataset S1).

Mfd's genomic association pattern correlates with that of RNAP

Given the known physical interaction between Mfd and RNAP(27), we hypothesized that Mfd

binding sites may correspond to RNAP binding locations. To test this, we performed ChIP-seq

of RpoB, the β subunit of RNAP, using a native antibody. Indeed, we found that Mfd association

largely overlaps with RpoB occupancy (Pearson coefficient = 0.68) (Fig. 1*B* and Fig. S2),

consistent with Mfd's proposed function as an RNAP co-factor in *B. subtilis*.

Mfd requires interaction with RNAP for its association with all genomic loci

Mfd is a multi-modular protein, consisting of eight domains connected by flexible linkers(27). Of these domains, the RNAP interacting domain (RID) and the translocase module (composed of domains D5 and D6) are critical for Mfd's ability to rescue stalled transcription complexes(28–30). *In vitro*, Mfd is recruited to the identified genomic regions through its interaction with RpoB. We therefore tested whether the interaction of Mfd with RpoB is critical for its recruitment to the genomic loci we identified. Prior *in vitro* work suggested that RID mutations abrogate the interaction between Mfd and RNAP(27). We therefore constructed a strain containing a Myc tagged Mfd, with a mutation at the L522 residue to disrupt Mfd's binding to RpoB, without disrupting the stability or folding of Mfd, analogous to that described in *E. coli*(27). Upon confirming that the *B. subtilis* L522A mutation (analogous to L499R mutation described in *E. coli*) disrupted Mfd's interaction with RNAP via a bacterial 2-hybrid assay (Fig. S3), we performed ChIP-seq experiments with this mutant. The ChIP signal we detected in WT strains was abrogated in the strain expressing the L522A allele of *mfd* (Fig. 1*C*). These results strongly suggested that Mfd's interaction with RNAP is essential for its recruitment to all genomic loci identified in the previous experiments and that Mfd functions as a genome-wide RNAP co-factor *in vivo*.

Mfd's association with DNA requires transcription elongation

In vitro, Mfd helps promote the rescue of arrested transcription elongation complexes (TECs), yet how Mfd recognizes stalled RNAPs *in vivo* remains unclear. We therefore asked whether Mfd association with various genomic loci was facilitated via loading during the transcription initiation or elongation phase. To distinguish between transcription initiation and elongation, we utilized the antimicrobial rifampicin, which directly blocks transcription initiation(31, 32), subsequently eliminating the formation of transcription elongation complexes (TECs). We performed both Mfd and RpoB ChIP-seq studies in the presence of rifampicin and found that

this treatment largely eliminated both Mfd and RpoB ChIP-seq binding signal (Fig. 1*D-E*). This

finding is consistent with biochemical evidence showing that Mfd is unable to release RNAP at

initiation sites(11). These data suggest that Mfd associates with elongating rather than initiating

RNAPs *in vivo*.

Mfd decreases RNAP density at some genomic loci

Mfd can function to promote both transcription elongation as well as transcription termination, at

least *in vitro* (11, 12). However, the importance of these functions *in vivo* remains elusive.

Because of Mfd's effects on RNAP *in vitro*, we wondered if and how Mfd's close association with

RNAP occupancy *in vivo* altered RNAP association in living cells. We therefore performed ChIP-

seq of RpoB in WT and *Δmfd* strains and identified where RpoB occupancy is altered in the

absence of Mfd. ChIP-seq experiments did not detect alterations in RpoB occupancy at the

majority of genes where Mfd associates based on our Mfd ChIP-seq studies. This may be a

result of various factors, such as the existence of redundant transcription-associated factors, the

specific growth conditions of the experiment, or potential limitations of detection thresholds

which are possible in an ensemble assay such as ChIP-seq. However, we did find a number of

genes where RpoB occupancy either increased or decreased in the *Δmfd* strain compared to

WT (Fig. 2*A*). We specifically noticed a bias towards a greater number of genes that exhibited

increased rather than decreased RpoB occupancy in the *Δmfd* strain. Quantification of these

results revealed that a total of 116 genes exhibited at least a two-fold increase, while 53 genes

exhibited at least a two-fold decrease in RpoB occupancy without Mfd. Many of these genes are

within the same operon, and therefore are expressed as single transcripts. Thus, rather than

single genes, we grouped and analyzed the identified genes as transcription units (TUs)(33).

Our analysis revealed that *Δmfd* strains contain 71 TUs with at least one gene containing a

minimum of two-fold increase, and 31 TUs with at least one gene containing a minimum of two-

fold decrease in RpoB occupancy compared to WT (Table 1, Datasets S2 and S3).

We next wanted to determine whether the changes in RpoB occupancy observed in the Δ*mfd*

strain were directly due to Mfd's activity at those regions or whether we were detecting indirect

effects. To address this, we looked for a correlation between the regions where Mfd associates

(ChIP-seq experiments) and where there are also changes in RpoB association levels in the

absence of Mfd. We only observed Mfd association at the genes that showed increased RpoB

occupancy in the Δ*mfd* strain. The genes with decreased RpoB occupancy in the Δ*mfd* strain

did not show Mfd association (Fig. 2*B*). Specifically, we detected Mfd at 52 of the 116 genes

with increased RpoB association (35 of the 71 TUs) in the Δ*mfd* strain compared to WT. We did

not detect Mfd association at any of the 53 genes (31 TUs) with decreased RpoB occupancy.

Because these 31 TUs do not display Mfd association, we concluded that decreased RpoB

occupancy at these sites are not directly due to Mfd activity and likely reflect indirect effects on

transcription in the Δ*mfd* strains.

We next performed RpoB ChIP-qPCR analysis to confirm our ChIP-seq results. We chose two

sites from our candidate genes with Mfd binding and increased RpoB occupancy in the Δ*mfd*

strain. We found results consistent with our ChIP-seq studies (Fig. S4). We additionally tested

the functionality of the Myc tagged Mfd using RpoB occupancy as our readout. We did not find

changes in the RpoB levels in the tagged Mfd strain, confirming its functionality, at least in

regulating RNAP association (Fig. S4).

Based on these findings, we conclude that increased RNAP occupancy at the identified

chromosomal sites is a direct result of Mfd's function as either an RNAP termination (12) or

processivity factor (34). In other words, the data strongly suggest that under regular

circumstances, Mfd decreases RNAP association.

Regions where Mfd increased RNAP density are enriched for regulatory RNAs

In vitro, Mfd's translocase activity can help release RNAP or assist it with elongation when exposed to different obstacles. However, whether there are endogenous hotspots of RNAP stalling that require Mfd function remains unknown. Furthermore, if such hotspots exist, the nature of the potential obstacles remains to be determined. Intriguingly, 92% of the TUs that showed both an increase in RNAP density in the *Δmfd* strain and direct Mfd association express a minimum of one regulatory RNA (Table S1). These regulatory RNAs are a subset of the 1583 regulatory RNAs in *B. subtilis*, which encompass a wide variety of RNAs, including independent, non-coding transcripts, antisense RNAs, and multiple riboswitches(33). In comparison, only 39% of the TUs with decreased RNAP density in *Δmfd* strains contain regulatory RNAs (Table S2), which is consistent with the average percentage of TUs with predicted regulatory RNAs in *B. subtilis*(33, 35).

Sites of Mfd function contain highly structured regulatory RNAs

We hypothesized that Mfd function at the identified regions was related to RNA secondary structure impeding RNAP processivity. This hypothesis is consistent with changes in RNAP processivity due to secondary RNA structures, such as hairpins in the context of intrinsic transcription termination, and other transcription regulatory processes(36–38). Previous work characterized the predicted secondary structure for each regulatory RNA in *B. subtilis*(35). We sought to test whether the regions with increased RNAP in *Δmfd* strains are more prone to transcribing RNAs with more stable secondary structures. We determined the average minimum free energy (MFE) z-score for each of the RNAs at these regions as a proxy of RNA structure stability(39, 40). Specifically, we examined the regulatory RNAs in the TUs that had both increased RpoB density in the *Δmfd* strain and Mfd association. We then compared the results to TUs that showed no difference in RpoB density between WT and *Δmfd* strains. We found that the TUs where there is Mfd binding and increased RpoB density in the *Δmfd* strain have

significantly higher predicted RNA secondary structures relative to all other known regulatory

RNAs (Fig. 3*A*).

Many of these regulatory RNAs are not transcribed during standard growth conditions. To

control for the potential biases of including non-expressed RNAs, we stratified our analysis to

exclude RNAs that were not expressed under our growth conditions, as determined from our

RNA-seq data (see Fig. S8). Consistent with our global analysis, we find that expressed

regulatory RNAs associated with Mfd binding and increased RpoB density in *Δmfd* have

significantly higher MFE z-scores (Fig. 3*A*, Fig. S5)*.* We also find that within regulatory RNAs

associated with Mfd binding and increased RpoB density in the *Δmfd* strains, those that were

highly expressed (top 50%) have a higher MFE z-score compared to those that were not

expressed (Fig. 3*B*). These findings suggest that Mfd regulates RNAP at regions containing

highly structured RNAs.

Sites of Mfd activity correlate with difficult-to-transcribe regions

If sites of Mfd association and Mfd alteration in RpoB density were related to RNAP processivity,

we would expect these sites to be difficult-to-transcribe. To identify such regions, we utilized

experimental data from native elongating transcript sequencing (NET-seq) in *B. subtilis*(41).

Through the capture of nascent transcripts from tagged immunoprecipitation of RNAP

molecules, NET-seq allows for identification of RNAP pausing in living cells with high

resolution(41). We observed that many genes with increased RpoB levels without Mfd displayed

high NET-sig signal, and that the corresponding sequencing patterns closely aligned (Fig. 4*A*).

We quantified this pattern by analyzing the NET-seq read count across the previously identified

genes where Mfd associates and has increased RpoB signal either with or without Mfd. We

found that the median NET-seq signal is significantly higher, by ~3.7-fold, at genes with both

increased RpoB signal in *Δmfd* strains and Mfd binding relative to genes with decreased RpoB

signal (Fig. 4*B-C*).

Larson et al. identified 9,989 discrete pause sites in *B. subtilis*, defined as nucleotides with

greater than four standard deviations in the NET-seq read count(41). We therefore tested

whether sites with proposed Mfd activity due to RNA secondary structure contained a greater

number of pause sites relative to the average number across the genome. We indeed found that

there are roughly 3-fold greater number of discrete pause sites in genes with proposed Mfd

activity relative to the average number across the genome, with a notable increase in the

frequency of genes with more than one pause site per 100bp at these sites (Fig. 4*D-E*). We did

not find this pattern in genes with decreased RpoB association in *Δmfd* strains (Fig. 4*D-E*).

Critically, the discrete pause sites identified by Larson, et al. in *B. subtilis* are limited to coding

regions only, and do not include regulatory RNAs, likely underestimating the number of true

pause sites requiring Mfd activity. From these findings we conclude that Mfd functions mainly at

RNAP pause sites *in vivo*.

Mfd's effect on RNAP at sites of structured regulatory RNAs is conserved in *E. coli*

Mfd is highly conserved across all bacterial phyla, and functional homologs exist throughout all

domains of life(42–44). Prior work shows that Mfd function is also highly conserved, at least with

regards to its mutagenic activity(17). To test whether Mfd's genome-wide coupling with RNAP

was also conserved in gram-negative species, we performed Mfd and RpoB ChIP-seq

experiments in *E. coli*. We found that as in *B. subtilis*, Mfd and RpoB occupancy are highly

correlated (Pearson coefficient 0.98) in *E. coli* (Fig. S6*A-C*), showing that Mfd's function as an

RNAP co-factor is likely conserved.

We also determined whether altering RpoB occupancy at sites containing structured RNAs was

true in *E. coli*. We compared the signal of the ChIP-seq data for RpoB in WT and *Δmfd* strains in

E. coli and looked for genes where RpoB occupancy was altered in the absence of Mfd. As in *B.*

subtilis, we found a higher number of genes that exhibited at least a two-fold increase in RpoB

association in the absence of Mfd, compared to genes with decreased RpoB association (105

versus 24 respectively, Fig. S6*D*, Datasets S4 and S5). 49 of the 105 genes with increased

RpoB association also had Mfd association (Dataset S4). Of these 49 sites, we found that

roughly 40% either contain regulatory RNAs or are thought to contain a structural element.

Three of these sites are TA RNAs (*symR, sokA, sokC*)(45, 46). One is a tRNA (*metZ*) and two

are small regulatory RNAs (*spf, gadY*), all of which are thought to contain secondary

structures(47–49). Seven sites are repeated extragenic/intragenic palindromes (RIP/REP),

which contain extensive secondary structure and are thought to facilitate transcription

termination(50, 51), and six are ribosomal proteins which contain structured regulatory elements

within their 5' untranslated regions(52). In contrast, no regulatory RNAs, tRNAs, or ribosomal

proteins were found in the genes with decreased RpoB association in the *Δmfd* strains (Dataset

S5). These findings suggest Mfd's function at sites of structured regulatory RNAs is largely

conserved across species.

Genome-wide Mfd binding is correlated with RNA secondary structure

While various methods exist for computational predictions of RNA secondary structure,

gathering empirical data on RNA structure continues to be technically challenging, particularly in

a high-throughput fashion. The recent development of parallel analysis of the mRNA structure

(PARS) has allowed for experimental determination of RNA secondary structure in multiple

species, including in *E. coli*(53, 54). This method utilizes enzymatic digestion of single and

double-stranded RNA with RNA-specific nucleases and couples this with highly selective cDNA

generation of digested RNAs and subsequent deep sequencing(53). Del Campo et al. utilized

PARS to establish the RNA secondary structure for roughly 2,500 genes in *E. coli*(54). We

therefore wanted to determine whether sites experimentally confirmed to have high RNA

secondary structure in *E. coli* correlate with regions of Mfd binding. We observed that across

genomic sites containing a high PARS score, both the presence of Mfd association and the

magnitude of Mfd binding closely aligned with the magnitude of experimentally determined RNA

secondary structure (Fig. 5*A*). To further quantify this, we measured the area under the curve

(AUC) for both our Mfd ChIP-seq and the PARS data set to assess whether these two signals

were correlated. We indeed found a strong positive correlation between the AUC for our Mfd

ChIP-seq and the PARS score (r = 0.6) across the *E. coli* genome (Fig. 5*B*). These data

suggest that the presence of RNA secondary structure, at least in *E. coli*, at least partially

correlates with Mfd genome-wide association and function.

GreA does not preferentially effect transcription of regions containing structured RNAs

Various factors are known to help rescue arrested RNAP through different mechanisms. One of

the most well-known and highly conserved proteins is GreA, which functions as an RNAP

antibacktracking factor by cleaving the nascent 3' RNA that has extruded from the RNAP

catalytic channel during backtracking(10, 55). Furthermore, GreA promotes the escape of RNAP

from the promoter and suppression of promoter-proximal pausing during transcription

initiation(56). To test whether GreA activity also contributed to RNAP release at the loci

transcribing structured RNAs, we performed RpoB ChIP-seq of *B. subtilis* WT and a *ΔgreA*

strain. We found that the *ΔgreA* strain only had 12 genes (and six TUs) with increased RpoB

occupancy (Dataset S6, Fig. S7). Two of the six TUs transcribe regulatory RNAs and neither

contained significant predicted secondary structure. These results suggest that unlike Mfd, GreA

does not function in releasing RNAP from sites containing secondary structure. We also

observed that in the *ΔgreA* strain, a total 469 genes exhibited less than two-fold RpoB

occupancy (Dataset S6, Fig. S7). The high number of genes with decreased RpoB occupancy in

the absence of GreA may be due to decreased efficiency of RNAP promoter escape during

initiation and consequently decreased levels of elongating RNAP molecules, consistent with *in*

vitro findings(56, 57).

Mfd decreases expression at structured, regulatory RNAs

We tested the effect of Mfd on transcription at sites containing structured RNAs. We began by

performing RNA-seq of WT and *Δmfd* strains in *B. subtilis*. Differential expression analysis

between WT and *Δmfd* revealed a total of 378 genes with statistically significant lower RNA

levels (Dataset S7). Consistent with our WT and *Δmfd* RpoB ChIP-seq results, we found that

more genes were upregulated than downregulated in the *Δmfd* strain (240 genes upregulated

compared to 138 genes downregulated) (Fig. S8 and Dataset S7). When comparing our RpoB

ChIP-seq findings to RNA-seq, we found that of the 116 genes with greater than 2-fold RpoB

ChIP-seq signal in the *Δmfd* strain, 30 of them directly showed increased expression, while

none show decreased expression in the absence of Mfd. Analysis of genes with decreased

expression in the *Δmfd* strain showed an equal decrease throughout the gene body, suggesting

that Mfd suppresses full-length transcripts (Fig. S9). Because standard RNA sequencing

protocols are often not suitable for accurate measurement of small RNAs(58), we wondered if

there are additional genes with increased RpoB density in the *Δmfd* strain that have

corresponding increases in transcription, but were not accurately detected in our RNA-seq

analysis. We therefore directly measured RNA levels using qRT-PCR at three loci containing

non-coding RNAs (the *trnY* locus, *txpa-ratA*, and *bsrH-asBsrH*), all of which show Mfd binding

and increased RpoB signal in the *Δmfd* strain. We found that all three of these loci have

increased gene expression in the *Δmfd* strain compared to WT (Fig. S10). To confirm that the

Δmfd strain does not globally repress transcription, we performed qRT-PCR analysis on two

control loci, *rpoB* and *yolA*, and found no difference in RNA levels between WT and the *Δmfd*

strain (Fig. S10). These findings suggest that Mfd's *in vivo* effects on RNAP lead to decreased

transcription.

Cells lacking Mfd are highly sensitized to toxin overexpression

Aside from decreased rates of evolution, there are few phenotypic defects that have been

detected in the absence of Mfd, even upon exposure to DNA damage(17, 18, 24, 25, 59). We

wondered whether the transcriptional regulation activity of Mfd at regions we detected were

physiologically relevant. To address this, we focused on the highest structured regulatory RNAs

which had altered RpoB density in the *Δmfd* strain and were also sites of Mfd association.

These regulatory RNAs were present in two pairs of type I toxin-antitoxin (TA) loci in *B. subtilis*:

the *txpA/ratA* locus and the *bsrH/as-bsrH* locus. Type I TA loci are characterized by the

expression of a small toxic peptide and a noncoding RNA that neutralizes toxin expression by

direct binding and either inhibiting translation or promoting degradation of the toxin mRNA(60).

The cellular functions of type I TA loci remain unclear, but they have been proposed to be

important for diverse aspects of physiology, including persister formation(61), biofilm

formation(62), and prophage maintenance(63). Five type I TA loci have been identified in *B.*

subtilis(60) - we found that three of these loci have both Mfd binding and a minimum of two-fold

increase in RpoB density in the *Δmfd* strain, while a fourth locus, *yonT/as-YonT*, showed a

significant increase in RpoB density in the *Δmfd* strain (Table S3).

Transcription regulation at type I TA loci is essential for cell survival as overexpression of type I

toxins can be lethal(64). Based on our data, we hypothesized that cells lacking Mfd would be

sensitive to toxin over-expression, similar to ethanol stress and heat shock(60). Because of the

potential pleotropic effects, we chose to directly test our hypothesis by upregulating toxin

expression in a controlled fashion. We therefore constructed an integrative plasmid containing

the *txpA* toxin gene under an IPTG (isopropyl b-D-1-thiogalacto-pyranoside) inducible promoter

in both WT and *Δmfd* strains, and performed cellular viability assays. We found that cells lacking

Mfd are highly sensitized to both chronic (Fig. 6*A*) and acute (Fig. 6*C*) overexpression of TxpA.

Cells lacking Mfd showed up to five orders of magnitude sensitivity to overexpression of this

toxin. Similarly, we overexpressed the BsrH toxin in WT and *Δmfd* strains to test whether Mfd's

effect was conserved. Indeed, we see roughly four orders of magnitude sensitization of the

Δmfd strain to overexpression of BsrH in both chronic (Fig. 6*B*) and acute (Fig. 6*D*) conditions.

To test whether this effect was directly due to overexpression of BsrH or TxpA, we performed

qRT-PCR analysis in our WT and *Δmfd* strains containing the overexpression constructs. We

found that in both cases, toxin overexpression was increased by ~2-3-fold in the *Δmfd* strain

(Fig. 6*E-F*). To confirm that the effects we observed were not specific to our integration site or

promoter, we performed qRT-PCR analysis on strains containing an IPTG-inducible *lacZ* gene.

We find no effect on *lacZ* transcript levels between WT and the *Δmfd* strain (Fig. S11).

Interestingly, recent data has revealed that *Clostridium difficile* cells lacking Mfd have

morphologic differences and viability defects that are thought to be related to toxin

overexpression(65), underscoring the likely conserved and critical role that Mfd may play on the

expression of such regulatory RNAs. Overall, our findings suggest that Mfd plays an important

role in regulating transcription of TA loci in *B. subtilis*.

Mfd promotes mutagenesis of the *txpA* gene

Mfd is an evolvability factor(17) and promotes mutagenesis under various conditions, promoting

the development of antimicrobial resistance, and mutagenesis in general during nutrient

starvation, stationary-phase growth, and replication-transcription conflicts(24, 25, 59, 66). Most

of these studies utilize engineered reporter systems to study Mfd-mediated mutagenesis. It

remains unclear what endogenous sites are prone to Mfd-mediated mutagenesis. Given our

findings that Mfd is critical for transcriptional regulation at structured RNAs, we wondered if such

sites were particularly prone to Mfd-mediated mutagenesis. We tested this hypothesis by

performing Luria-Delbrück fluctuation assays(67) to select for toxin-resistant mutants and

assessed the mutation rate at our ectopic *txpA* locus. We plated cells on a high concentration of

IPTG (1mM) to ensure that only genetic revertants would produce viable colonies. We found

that the mutation rate of *txpA* is nearly 7-fold lower in the *Δmfd* compared to the WT strain (Fig.

S12). These differences are higher than previously reported mutation rate reductions in *Δmfd*

strains(17, 59). We conclude that Mfd promotes mutagenesis at TA loci in *B. subtilis* and given

our prior findings, it likely accelerates evolution at many of its endogenous targets.

Discussion

In this work, we identified the endogenous targets of Mfd and how Mfd impacts RNAP

association and transcription *in vivo*. We found that in both *E. coli* and *B. subtilis*, most of the

sites where Mfd associates and modulates transcription contain highly structured regulatory

RNAs. We found that RNAP modulation and transcriptional control by Mfd predominantly occur

at regions of frequent RNAP pausing. Our experiments demonstrated that Mfd is important for

maintaining cellular viability upon activation of TA systems that contain highly structured RNAs.

We also showed that Mfd promotes mutagenesis at these regions. Based on these findings, we

propose that Mfd is a RNAP co-factor which helps regulate transcription at many chromosomal

regions, especially those with highly structured RNAs.

Though Mfd is non-essential under many conditions, it becomes critical for cell viability when TA

systems are expressed. This suggests that there may be other situations when transcriptional

regulation by Mfd is important. It is not unreasonable to expect that Mfd would be critical under

conditions related to the functions of the genes where Mfd binds and regulates RNAP

association. For example, Mfd targets a mannose utilization operon, suggesting that efficient

growth on mannose may require transcriptional regulation by Mfd. By identifying the genomic

regions where Mfd binds, our study has provided key information regarding the conditions under

which Mfd function may become critical. Further work is needed to identify and characterize

these conditions.

Though Mfd clearly reduces RNAP density at regions containing structured RNAs, it is not clear

if this is due to RNAP rescue, early transcription termination, or a combination of both (Fig. 7).

Though Mfd can rescue or terminate transcription, the decision about which activity will occur is

thought to be related to the magnitude of the roadblock faced by RNAP, with more significant

roadblocks favoring termination(10, 21, 22, 34, 68, 69). Additionally, recent work from Le et al.

using *in vitro* translocase assays showed that Mfd rescues RNAP at pause sites, but that more

severe obstacles to RNAP movement lead to eventual termination(23).

Existing studies show that RNA secondary structure is capable of impeding RNAP processivity.

For example, RNAP pausing can be promoted by the formation of stable RNA hairpin structures

in the exit channel of RNAP and inhibit its movement(5, 70–72). Pausing via RNA secondary

structure is a mechanism that can regulate gene expression at riboswitches, promote coupling

of transcription and translation(73, 74), and is critical for the process of intrinsic transcription

termination in bacteria(12, 75).

The severity of stalling at structured RNAs is likely dependent on multiple factors including the

stability of the structure, length, and expression level. At certain sites, it is possible that such

pausing may induce RNAP backtracking, but mechanistic studies suggest that it is more

common for pausing to induce a "half translocated" state of RNAP(71) which inhibits its

immediate processivity(76). To our knowledge, it has not been shown that RNA secondary

structure is a significant enough roadblock to cause RNAP termination, suggesting that it is a

generally less severe of an impediment to RNAP movement. These findings would favor a

model whereby Mfd functions to promote RNAP rescue and elongation at structured RNAs,

leading to the decreased RNAP levels as we observe in our ChIP studies. However, we cannot

rule out the possibility that the regulatory RNAs identified in our studies promote a more severe

impediment to RNAP and that decreased RNAP levels reflect RNAP termination by Mfd.

Given the wide range of biological functions regulated by structured RNAs, Mfd's activity at the

sites we identified is likely to have a broad range of phenotypic effects. For example, we

identified multiple riboswitches containing TUs where Mfd binds and promotes RNAP release.

These TUs are involved in many critical metabolic processes, ranging from beta-glucoside

metabolism (*bglP-bglH-yxiE*)(77, 78), to the utilization of glycerol (*glpF-glpK* and *glpT-glpQ*)(79)

to purine metabolism (*purEKBCSQLFMNHD*)(80, 81). We also identified Mfd binding and RNAP

release at a locus containing a long cis-acting antisense RNA (*yabE*/S25) that is thought to play

a role in cell wall maintenance(82). Lastly, amongst many other sites, we identified Mfd binding

and RNAP termination at tRNA loci (the *trnY* locus in *B. subtilis* containing a highly structured

RNA of unknown function). The physiological relevance of Mfd's activity at these sites requires

further investigation that is outside of the scope of this study.

Various mechanisms of transcription-associated mutagenesis (TAM) exist(83, 84). Based on our

findings, we propose that the inherent structure of RNA may be an additional mechanism by

which transcription promotes mutagenesis, at least partially through Mfd. Interestingly, RNA

secondary structure has been reported to enhance mutation rates in replicating retroviruses(85),

suggesting that evolution via RNA secondary structure may be a universal mechanism. In

addition, a recent study by Thornlow et al.(86), using computational analyses, revealed that

tRNAs have higher mutation rates relative to other parts of the genome(86). They also

suggested that this phenomenon is linked to transcription. It is therefore quite possible that, at

least in bacteria, the evolution of tRNA structures is mediated by Mfd. By promoting

mutagenesis at sites of highly structured RNAs, Mfd may inherently alter the secondary

structure encoded at the site of its activity, leading to novel or altered functions of the RNA.

Additionally, non-coding RNAs are well-known to evolve very quickly(87), however, the

mechanisms by which this occurs is unknown. Our results suggest that Mfd may contribute to

the evolution of these regions via its mutagenic activity. Addressing this possibility would further

contribute to our mechanistic understanding of how non-coding RNAs evolve.

Materials and Methods

Strain constructions

All strains and plasmids used and constructed in this study are listed in Table S4 and primers

used are listed in Table S5. *B. subtilis* stains used in this study were derivates of the HM1

(JH642) parent strain(88). *E. coli* strains used were derivates of K-12 MG1655 (89).

Transformations into *B. subtilis* HM1 were performed under standard conditions as previously

described(90). Plasmids used in this study were grown in *E. coli* DH5α. Plasmids were cloned

using chemical transformations of competent *E. coli*. All plasmid purification was performed by

growth of appropriate *E. coli* strain overnight at 37° C in Luria-Bertani (LB) medium supplement

with the appropriate antibiotic and plasmids were subsequently purified using the GeneJet

Plasmid Miniprep Kit (Thermo). Further details on strain construction can be found in the SI

appendix.

Growth conditions

For experiments in *B. subtilis* and *E. coli*, cultures were grown as described unless otherwise

indicated. Cells were plated on LB supplemented with the appropriate antibiotic for isolation of

single colonies. Overnight cultures from single colonies were grown at 37° C in LB at 260 RPMs

and the following day cells were diluted back to OD600 0.05 and grown to exponential phase

(OD600 0.3-0.5) before harvesting. For acute rifampicin ChIP experiments, cultures were grown

in identical fashion until they reach OD600 0.3-0.5 and rifampicin was added at a concentration

of 100µg/mL for 5 minutes before harvesting.

ChIP-seq and ChIP-qPCR experiments

For *B. subtilis* Mfd ChIP experiments, c-Myc mouse monoclonal antibody (clone 9E10) was

used (Thermo). For *E. coli* ChIP experiments, custom polyclonal *E. coli* Mfd rabbit antisera was

produced and purified by Covance. ELISA titers were used to confirm Mfd titers from antisera.

For RpoB experiments, RNA polymerase beta mouse monoclonal antibody (clone 8RB13) was

used (Thermo).

ChIP experiments were performed as previously described(91, 92). Briefly, Cells were grown to

exponential phase as previously described and crosslinked with 1% formaldehyde v/v. After 20

minutes at room temperature, 0.5M final concentration of glycine was added and cells were

pelleted, washed in cold 1x PBS and pelleted again. Cells were resuspended in solution A (10

mM Tris pH 8.0, 10 mM EDTA, 50 mM NaCl, 20% sucrose), supplemented with 1 mg/ml

lysozyme and 1 mM AEBSF at 37° C for 30 minutes. 2x IP buffer (100 mM Tris pH 7.0, 10 mM

EDTA, 300 mM NaCl, 20% triton x-100), supplemented with 1mM AEBSF, was then added and

lysates were incubated on ice for 30 minutes. Cell lysates were sonicated four times at 30%

amplitude for ten seconds using a Fisher sonic dismembrator (Fisher FB120). Lysates were

centrifuged at 8000 RPMs for 15 minutes at 4° C. The supernatant was transferred into new

microfuge tubes.

ChIP lysates were split into a total DNA input control (40µl of lysate) and immunoprecipitation

(IP) (1mL of lysate). For Mfd ChIP experiments in *B. subtilis* 12µl anti-c-Myc antibody was

added to the IP samples. For Mfd ChIP-seq experiments in *E. coli*, 4 µl of native *E. coli* anti-Mfd

antibody was used. 2µl of anti-RpoB antibody was added for RpoB ChIPs in both *B. subtilis* and

E. coli. IP lysates were rotated overnight at 4° C. The following day, 30μl Protein A sepharose

beads (GE) were added to the IP samples and rotated for one hour at room temperature. Beads

were then pelleted with centrifugation at 2000 RPMs for one minute. Supernatant was decanted

and beads were subsequently washed six times with 1x IP buffer and one time with 1x TE pH

8.0. Beads were then pelleted and resuspended in 100μl of elution buffer (50mM Tris pH 8.0,

10mM EDTA), and 1% SDS and incubated at 65° C for 10 minutes. Beads were pelleted by

centrifugation and supernatant was transferred to a new microfuge tube. A second round of

elution was performed by resuspension of beads in 150μl of elution buffer II (10mM Tris pH 8.0,

1 mM EDTA, 0.67% SDS). Beads were pelleted and supernatant was transferred to microfuge

tube containing eluate from the first elution. IP samples were then incubated overnight at 65° C.

The following day, proteinase K was added at a final concentration 0.4 mg/mL and samples

were incubated for two hours at 37° C. Purification was performed by using the GeneJet PCR

Purification Kit (Thermo).

Library preparation for ChIP-seq was performed using the Nextera XT DNA Library Prep Kit

(Illumina) according to manufacturer's instructions. For ChIP-quantitative PCR (qPCR), Sso

Advanced Universal SYBR Green Supermix (BioRad) was used according to manufacturer's

instructions.

NET-seq analysis

Data from Larson et al. (41) was analyzed for determination of average NET-seq read counts in

addition to identification of pause sites. Data was collected from Gene Expression Omnibus

(accession number GSE56720) and plotted as a wiggle (.wig) file on the *B. subtilis* 168 genome.

Details of pause site determination are defined in the supplementary methods of Larson et al.

(41). In order to calculate the average read count for each gene, gene coordinates were

determined and NET-seq signal was averaged across these nucleotide positions. Average NET-

seq signal for each gene was calculated from the strand (plus or minus strand) that had the

highest NET-seq signal. The number of pause sites identified for each gene of interest was

normalized to gene length.

PARS-seq analysis

PARS data was kindly provided by Zoya Ignatova. Original PARS score data was refined to

remove duplicate data points. All positive PARS scores were then plotted as a wiggle (.wig) file

on the *E. coli* K-12 MG1655 genome sequence (Genbank: U00096.2). The average PARS

score for each annotated feature in the *E. coli* chromosome was calculated using custom

scripts.

Whole-genome sequencing analysis

ChIP-seq and RNA-seq samples were sequenced using the Illumina Nextseq 500/550

Sequencing system at the University of Washington Northwest Genomics Center and the

VANTAGE Sequencing Core at Vanderbilt University. After sequencing, sample reads from *B.*

subtilis were mapped to 168 genome (accession number: NC_000964.3) and from *E. coli* to the

MG1655 genome (accession number: NC_ U00096.2) using Bowtie2(93). For data visualization,

SAMtools was used to process SAM files(94) to produce wiggle plots(95). Wiggle files from all

ChIP samples were normalized to input samples (total input DNA subtracted from the ChIP

signal). For quantification of ChIP-seq and RNA-seq samples, BAM files were processed by the

featureCounts program to determine read counts per gene(96). To determine differential RNA-

seq expression (see supplementary materials and methods for RNA-seq experiments) and

differential ChIP-seq binding, read counts were analyzed by DEseq2 software(97). In order to

determine correlations between RpoB ChIP binding and Mfd ChIP, read counts generated by

featureCounts were divided by the total number of sequencing reads per sample. ChIP samples

were then divided by input samples and $\log_2$ normalized.

Quantitative RT-PCR assays

For quantification of RNA using qRT-PCR, *B. subtilis* cultures were grown to exponential phase

as described in the growth conditions sections. For experiments containing IPTG-inducible

constructs (P_{spank}-txpA, P_{spank}-bsrH, and P_{spank}-lacZ), expression was induced with 1mM IPTG for

5 minutes and RNA was subsequently extracted as described in the RNA-seq experiments

section (see supplementary materials and methods). Subsequently, 1µg of total RNA was

treated with DNaseI (Thermo) for one hour at 37° C. DNase denaturation was performed with

addition of 10mM EDTA and incubation at 65° C for 10 minutes. cDNA generation was

performed using the iScript Supermix (BioRad), according to the manufacturer's instructions.

Quantitative PCR (qPCR) was performed using the Sso Advanced Universal SYBR Green

Supermix (BioRad), according to manufacturer's instructions. For normalization of qRT-PCR,

primers to *B. subtilis* rRNA was used.

Cell survival assays

For chronic survival assays, strains were struck out on LB agar plates and *B. subtilis* cultures

were grown in 2mL LB until they reached an OD600 of 0.5-1.0. All cultures were normalized to

OD600 0.3 and serial dilutions were performed in 1x Spizizen's salts. 5µl of cells were plated on

control plates containing LB agar only and LB agar plates containing the designated

concentration of IPTG (see figure legends for concentrations). Plates were grown at 30° C

overnight and CFUs were enumerated the following day.

For acute survival assays, cultures were grown in 2ml until they reached an OD600 of 0.5-1.0

and then diluted back to OD600 0.05. Either 1mM or 0.1 mM IPTG was then added and cells

were grown for approximately 60 minutes (OD600~0.3). Cells were subsequently washed two

times with 1x Spizizen's salts to remove residual IPTG and were serially diluted. 5µl of cells

were plated on LB agar and plates were grown at 30° C overnight for CFU enumeration. For

both chronic and acute survival assays, images were taken using the BioRad Gel Doc XR+

Molecular Imager.

Acknowledgements

We thank Patrick Nugent and Ankunda Kariisa for assistance with strain construction and

bacterial 2-hybrid assays, Kevin Lang for helpful discussions, Sirena Tran for assistance with *E.*

coli ChIP-seq experiments, Zoya Ignatova for PARS data set, Charlie Lee at the University of

Washington and the VANTAGE genomics core at Vanderbilt for assistance with sequencing

experiments. This work was supported by the NIH National Institute of Allergy and Infectious

Diseases AI127422 and Training Program in Environmental Toxicology T32 ES007028 to K.B.

Footnotes

Author contributions: M.N.R., C.M., and H.M. conceived of and designed experiments. M.N.R.,

C.M., K.B. performed experiments. M.N.R., C.M., and H.M analyzed data and wrote the paper.

To whom the correspondence should be addressed. Email: houra.merrikh@vanderbilt.edu

The authors declare no conflict of interest

References

1.   N. Komissarova, M. Kashlev, Transcriptional arrest: Escherichia coli RNA polymerase

translocates backward , leaving the 3' end of the RNA intact. *Proc. Natl. Acad. Sci.* **94**,

1755–1760 (1997).

2.   P. Mcglynn, N. J. Savery, M. S. Dillingham, The conflict between DNA replication and

transcription. *Mol. Microbiol.* **85**, 12–20 (2012).

3.   I. Artsimovitch, R. Landick, Pausing by bacterial RNA polymerase is mediated by

mechanistically distinct classes of signals. *Proc. Natl. Acad. Sci.* **97**, 7090–7095 (2000).

4.   A. Mayer, H. M. Landry, L. S. Churchman, Pause & go: from the discovery of RNA

polymerase pausing to its functional implications. *Curr. Opin. Cell Biol.* **46**, 72–80 (2017).

5.   C. L. Chan, D. Wang, R. Landick, Multiple interactions stabilize a single paused

transcription intermediate in which hairpin to 3' end spacing distinguishes pause and

termination pathways. *J. Mol. Biol.* **268**, 54–68 (1997).

6.   D. A. Erie, The many conformational states of RNA polymerase elongation complexes

and their roles in the regulation of transcription. *Biochim. Biophys. Acta* **1577**, 224–239

(2002).

7.   A. C. M. Cheung, P. Cramer, Structural basis of RNA polymerase II backtracking, arrest

and reactivation. *Nature* **471**, 249–253 (2011).

8.   E. Nudler, A. Mustaev, E. Lukhtanov, A. Goldfarb, The RNA–DNA Hybrid Maintains the

Register of Transcription by Preventing Backtracking of RNA Polymerase. *Cell* **89**, 33–41

(1997).

9.   E. Nudler, RNA Polymerase Backtracking in Gene Regulation and Genome Instability.

Cell **149**, 1438–1445 (2012).

10. D. Dutta, K. Shatalin, V. Epshtein, M. E. Gottesman, E. Nudler, Linking RNA polymerase

backtracking to genome instability in E. coli. *Cell* **146**, 533–543 (2011).

11. J. Park, M. T. Marr, J. W. Roberts, E. coli Transcription Repair Coupling Factor (Mfd

Protein) Rescues Arrested Complexes by Promoting Forward Translocation. *Cell* **109**,

757–767 (2002).

12.   J.-S. Park, J. W. Roberts, Role of DNA bubble rewinding in enzymatic transcription

termination. *Proc. Natl. Acad. Sci.* **103**, 4870–4875 (2006).

13.   I. Mellon, G. Spivak, P. C. Hanawalt, Selective removal of transcription-blocking DNA

damage from the transcribed strand of the mammalian DHFR gene. *Cell* **51**, 241–249

(1987).

14.   C. Selby, A. Sancar, Molecular mechanism of transcription-repair coupling. *Science (80-.*

). **260**, 53 (1993).

15.   Christopher P. Selby, Mfd Protein and Transcription-Repair Coupling in E. coli.

Photochem Photobiol. **93**, 280–295 (2017).

16.   P. C. Hanawalt, G. Spivak, Transcription-coupled DNA repair: two decades of progress

and surprises. *Nat. Rev. Mol. Cell Biol.* **9**, 958–70 (2008).

17.   M. N. Ragheb, *et al.*, Inhibiting the Evolution of Antibiotic Resistance. *Mol. Cell* **73**, 157-

165.e5 (2019).

18.   V. Kamarthapu, *et al.*, ppGpp couples transcription to DNA repair in E.coli. *Science (80-.*

). (2016) https:/doi.org/10.1126/science.aad6945.

19.   V. Epshtein, *et al.*, UvrD facilitates DNA repair by pulling RNA polymerase backwards.

Nature **505**, 372–377 (2014).

20.   S. E. Cohen, *et al.*, Roles for the transcription elongation factor NusA in both DNA repair

and damage tolerance pathways in Escherichia coli. *Proc. Natl. Acad. Sci.* **107**, 15517–

15522 (2010).

21.   J. M. Zalieckas, L. V Wray, A. E. Ferson, S. H. Fisher, Transcription–repair coupling

factor is involved in carbon catabolite repression of the Bacillus subtilis hut and gnt

operons. *Mol. Microbiol.* **27**, 1031–1038 (1998).

22.   J. M. Zalieckas, L. V Wray, S. H. Fisher, Expression of the Bacillus subtilis acsA gene:

position and sequence context affect cre-mediated carbon catabolite repression. *J.*

Bacteriol. **180**, 6649–54 (1998).

23.   T. T. Le, *et al.*, Mfd Dynamically Regulates Transcription via a Release and Catch-Up

Mechanism. *Cell* **173**, 1823 (2018).

24.   C. Ross, *et al.*, Novel Role of mfd : Effects on Stationary-Phase Mutagenesis in Bacillus

subtilis. *J. Bacteriol.* **188**, 7512–7520 (2006).

25.   H. A. Martin, M. Pedraza-reyes, R. E. Yasbin, E. A. Robleto, Transcriptional De-

Repression and Mfd Are Mutagenic in Stressed Bacillus subtilis Cells. *J Mol Microbiol*

Biotechnol **21**, 45–58 (2011).

26.   H. N. Ho, A. M. van Oijen, H. Ghodke, The transcription-repair coupling factor Mfd

associates with RNA polymerase in the absence of exogenous damage. *Nat. Commun.*

9, 1–12 (2018).

27.   A. M. Deaconescu, *et al.*, Structural Basis for Bacterial Transcription-Coupled DNA

Repair. *Cell*, 507–520 (2006).

28.   A. L. Chambers, A. J. Smith, N. J. Savery, A DNA translocation motif in the bacterial

transcription-repair coupling factor, Mfd. *Nucleic Acids Res.* **31** (2003).

29.   A. J. Smith, N. J. Savery, RNA polymerase mutants defective in the initiation of

transcription-coupled DNA repair. *Nucleic Acids Res.* **33**, 755–764 (2005).

30.   C. Brugger, *et al.*, Molecular determinants for dsDNA translocation by the transcription-

repair coupling and evolvability factor Mfd. *Nat. Commun.* (2020)

https:/doi.org/10.1038/s41467-020-17457-1 (August 15, 2020).

31.   W. R. McClure, C. L. Cech, On the mechanism of rifampicin inhibition of RNA synthesis.

J. Biol. Chem. **253**, 8949–8956 (1978).

32.   E. A. Campbell, *et al.*, Structural mechanism for rifampicin inhibition of bacterial RNA

polymerase. *Cell* **104**, 901–912 (2001).

33.   P. Nicolas, *et al.*, Condition-dependent transcriptome reveals high-level regulatory

architecture in Bacillus subtilis. *Science* **335**, 1103–6 (2012).

34.   R. S. Washburn, Y. Wang, M. E. Gottesman, Role of E .coli Transcription-Repair

Coupling Factor Mfd in Nun-mediated Transcription Termination. *J. Mol. Biol.* **2836**, 655–

662 (2003).

35.   R. A. T. Mars, P. Nicolas, E. L. Denham, J. M. van Dijl, Regulatory RNAs in Bacillus

subtilis: a Gram-Positive Perspective on Bacterial RNA-Mediated Regulation of Gene

Expression. *Microbiol. Mol. Biol. Rev.* **80**, 1029–1057 (2016).

36.   A. Ray-Soni, M. J. Bellecourt, R. Landick, Mechanisms of Bacterial Transcription

Termination: All Good Things Must End. *Annu. Rev. Biochem.* **85**, 319–347 (2016).

37.   I. Gusarov, E. Nudler, The mechanism of intrinsic transcription termination. *Mol. Cell* **3**,

495–504 (1999).

38.   A. Weixlbaumer, K. Leon, R. Landick, S. A. Darst, Structural basis of transcriptional

pausing in bacteria. *Cell* **152**, 431–441 (2013).

39.   S. Y. Le, J. V Maizel, A method for assessing the statistical significance of RNA folding. *J.*

Theor. Biol. **138**, 495–510 (1989).

40.   E. Freyhult, P. P. Gardner, V. Moulton, A comparison of RNA folding measures. *BMC*

Bioinformatics **6**, 241 (2005).

41. M. H. Larson, *et al.*, A pause sequence enriched at translation start sites drives transcription dynamics in vivo. *Science (80-.).* **344**, 1042–1047 (2014).

42. C. P. Selby, A. Sancar, Cockayne syndrome group B protein enhances elongation by RNA polymerase II. *Proc. Natl. Acad. Sci.* **94**, 11205–11209 (1997).

43. P. C. Hanawalt, DNA repair. The bases for Cockayne syndrome. *Nature* **405**, 415–416 (2000).

44. H. Menoni, J. H. J. Hoeijmakers, W. Vermeulen, Nucleotide excision repair-initiating proteins bind to oxidative DNA lesions in vivo. *J. Cell Biol.* **199**, 1037–1046 (2012).

45. K. Pedersen, K. Gerdes, Multiple hok genes on the chromosome of Escherichia coli. *Mol. Microbiol.* **32**, 1090–1102 (1999).

46. M. Kawano, L. Aravind, G. Storz, An antisense RNA controls synthesis of an SOS-induced toxin evolved from an antitoxin. *Mol. Microbiol.* **64**, 738–754 (2007).

47. A. H. Potts, *et al.*, Global role of the bacterial post-transcriptional regulator CsrA revealed by integrated transcriptomics. *Nat. Commun.* **8**, 1–15 (2017).

48. T. Kenri, F. Imamoto, Y. Kano, Three tandemly repeated structural genes encoding tRNAf1Met in the metZ operon of Escherichia coli K-12. *Gene* **138**, 261–262 (1994).

49. C. Bækkedal, P. Haugen, The Spot 42 RNA: A regulatory small RNA with roles in the central metabolism. *RNA Biol.* **12**, 1071–1077 (2015).

50. V. Khemici, A. J. Carpousis, The RNA degradosome and poly(A) polymerase of Escherichia coli are required in vivo for the degradation of small mRNA decay intermediates containing REP-stabilizers. *Mol. Microbiol.* **51**, 777–790 (2004).

789   51.   W. Liang, K. E. Rudd, M. P. Deutscher, A Role for REP sequences in regulating

790   translation. *Mol. Cell* **58**, 431–439 (2015).

791   52.   Y. Fu, K. Deiorio-Haggar, J. Anthony, M. M. Meyer, Most RNAs regulating ribosomal

792   protein biosynthesis in Escherichia coli are narrowly distributed to Gammaproteobacteria.

793   *Nucleic Acids Res.* **41**, 3491–3503 (2013).

794   53.   Y. Wan, K. Qu, Z. Ouyang, H. Y. Chang, Genome-wide mapping of RNA structure using

795   nuclease digestion and high-throughput sequencing. *Nat. Protoc.* **8**, 849–869 (2013).

796   54.   C. Del Campo, A. Bartholomäus, I. Fedyunin, Z. Ignatova, Secondary Structure across

797   the Bacterial Transcriptome Reveals Versatile Roles in mRNA Regulation and Function.

798   *PLOS Genet.* **11**, e1005613 (2015).

799   55.   S. Borukhov, A. Polyakov, V. Nikiforov, A. Goldfarb, GreA protein: A transcription

800   elongation factor from Escherichia coli. *Proc. Natl. Acad. Sci.* **89**, 8899–8902 (1992).

801   56.   L. M. Hsu, N. V Vot, M. J. Chamberlint, Escherichia coli transcript cleavage factors GreA

802   and GreB stimulate promoter escape and gene expression in vivo and in vitro.

803   *Biochemistry* **92**, 11588–11592 (1995).

804   57.   E. Stepanova, *et al.*, Analysis of promoter targets for Escherichia coli transcription

805   elongation factor GreA in vivo and in vitro. *J. Bacteriol.* **189**, 8772–8785 (2007).

806   58.   F. Ozsolak, P. M. Milos, RNA sequencing: advances, challenges and opportunities. *Nat.

807   Rev. Genet.* **12**, 87–98 (2011).

808   59.   S. Million-Weaver, *et al.*, An underlying mechanism for the increased mutagenesis of

809   lagging-strand genes in Bacillus subtilis. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E1096-105

810   (2015).

811   60.   S. Brantl, N. Jahn, SRNAs in bacterial type I and type III toxin-antitoxin systems. *FEMS*

812   *Microbiol. Rev.* **39**, 413–427 (2015).

813   61.   T. Dörr, M. Vulić, K. Lewis, Ciprofloxacin Causes Persister Formation by Inducing the

814   TisB toxin in Escherichia coli. *PLoS Biol.* **8**, e1000317 (2010).

815   62.   J. Domka, J. Lee, T. Bansal, T. K. Wood, Temporal gene-expression in Escherichia coli

816   K-12 biofilms. *Environ. Microbiol.* **9**, 332–346 (2007).

817   63.   S. Durand, N. Jahn, C. Condon, S. Brantl, Type i toxin-antitoxin systems in Bacillus

818   subtilis. *RNA Biol.* **9**, 1491–1497 (2012).

819   64.   J. M. Silvaggi, J. B. Perkins, R. Losick, Small Untranslated RNA Antitoxin in Bacillus

820   subtilis. *J. Bacteriol.* **187**, 6641–6650 (2005).

821   65.   S. E. Willing, *et al.*, Increased toxin expression in a Clostridium difficile mfd mutant (2015)

822   https:/doi.org/10.1186/s12866-015-0611-5.

823   66.   H. Wimberly, *et al.*, R-loops and nicks initiate DNA breakage and genome instability in

824   non-growing Escherichia coli. *Nat. Commun.* **4**, 2115 (2013).

825   67.   S. Luria, M. Delbrück, Mutations of Bacteria from Virus Sensitivity to Virus Resistance.

826   *Genetics* **28**, 491–511 (1943).

827   68.   C. P. Selby, A. Sancar, Transcription-repair coupling and mutation frequency decline.

828   *Microbiol. Rev.* **175**, 7509–7514 (1993).

829   69.   S. C. Hung, M. E. Gottesman, Phage HK022 Nun Protein Arrests Transcription on Phage

830   lambda DNA in vitro and Competes with the Phage lambda N Antitermination Protein. *J.*

831   *Mol. Biol.*, 428–442 (1995).

832   70.   J. Y. Kang, *et al.*, RNA Polymerase Accommodates a Pause RNA Hairpin by Global

833   Conformational Rearrangements that Prolong Pausing. *Mol. Cell* **69**, 802-815.e1 (2018).

71.   M. L. Kireeva, M. Kashlev, Mechanism of sequence-specific pausing of bacterial RNA

polymerase. *Proc. Natl. Acad. Sci.* **106**, 8900–8905 (2009).

72.   V. R. Tadigotla, *et al.*, Thermodynamic and kinetic modeling of transcriptional pausing.

Proc. Natl. Acad. Sci. **103**, 4439–4444 (2006).

73.   M. E. Winkler, C. Yanofsky, Pausing of RNA polymerase during in vitro transcription of

the tryptophan operon leader region. *Biochemistry* **20**, 3738–44 (1981).

74.   A. V. Yakhnin, H. Yakhnin, P. Babitzke, RNA Polymerase Pausing Regulates Translation

Initiation by Providing Additional Time for TRAP-RNA Interaction. *Mol. Cell* **24**, 547–557

(2006).

75.   A. Ray-Soni, R. A. Mooney, R. Landick, Trigger loop dynamics can explain stimulation of

intrinsic termination by bacterial RNA polymerase without terminator hairpin contact.

Proc. Natl. Acad. Sci. **114**, E9233–E9242 (2017).

76.   J. Saba, *et al.*, The elemental mechanism of transcriptional pausing. *Elife* **8** (2019).

77.   S. Krüger, S. Gertz, M. Hecker, Transcriptional analysis of bglPH expression in Bacillus

subtilis: Evidence for two distinct pathways mediating carbon catabolite repression. *J.*

Bacteriol. **178**, 2637–2644 (1996).

78.   I. Irnov, C. M. Sharma, J. Vogel, W. C. Winkler, Identification of regulatory RNAs in

Bacillus subtilis. *Nucleic Acids Res.* **38**, 6637–6651 (2010).

79.   R.-P. Nilsson, L. Beijer, B. Rutberg, The glpT and glpQ genes of the glycerol regulon in

Bacillus subtilis. *Microbiology* **140**, 723–730 (1994).

80.   M. Mandal, B. Boese, J. E. Barrick, W. C. Winkler, R. R. Breaker, Riboswitches Control

Fundamental Biochemical Pathways in Bacillus subtilis and Other Bacteria. *Cell* **113**,

577–586 (2003).

81.   D. J. Ebbole, H. Zalkin, Cloning and characterization of a 12-gene cluster from Bacillus
858   subtilis encoding nine enzymes for de novo purine nucleotide synthesis. *J. Biol. Chem.*
859   **262**, 8274–8287 (1987).

860   82.   W. Eiamphungporn, J. D. Helmann, Extracytoplasmic Function σ Factors Regulate
861   Expression of the Bacillus subtilis yabE Gene via a cis-Acting Antisense RNA. *J.*
862   *Bacteriol.* **191**, 1101–1105 (2009).

863   83.   S. Jinks-Robertson, A. S. Bhagwat, Transcription-Associated Mutagenesis. *Annu. Rev.*
864   *Genet.* **48**, 341–359 (2014).

865   84.   N. Kim, S. Jinks-Robertson, Transcription as a source of genome instability. *Nat. Rev.*
866   *Genet.* **13**, 204–214 (2012).

867   85.   V. K. Pathakt, H. M. Temin, 5-Azacytidine and RNA Secondary Structure Increase the
868   Retrovirus Mutation Rate. *J. Virol.* **66**, 3093–3100 (1992).

869   86.   T. M. Lowe, *et al.*, Transfer RNA genes experience exceptionally elevated mutation rates.
870   *Proc. Natl. Acad. Sci.* **115**, 8996–9001 (2018).

871   87.   H. A. Dutcher, R. Raghavan, Origin, Evolution, and Loss of Bacterial Small RNAs.
872   *Microbiol. Spectr.* **6** (2018).

873   88.   S. P. Brehm, S. P. Staal, J. A. Hoch, Phenotypes of pleiotropic-negative sporulation
874   mutants of Bacillus subtilis. *J. Bacteriol.* **115**, 1063–70 (1973).

875   89.   F. R. Blattner, *et al.*, The complete genome sequence of Escherichia coli K-12. *Science*
876   *(80-.).* **277**, 1453–1462 (1997).

877   90.   C. . Harwood, S. M. Cutting, J. Wiley, *Molecular Biological Methods for Bacillus* (1990)
878   (April 11, 2019).

91.   K. S. Lang, *et al.*, Replication-Transcription Conflicts Generate R-Loops that Orchestrate
Bacterial Stress Survival. *Cell* **170**, 787-790.e18 (2017).

92.   H. Merrikh, C. Machón, W. H. Grainger, A. D. Grossman, P. Soultanas, Co-directional
replication-transcription conflicts lead to replication restart. *Nature* **470**, 554–7 (2011).

93.   B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,
357–359 (2012).

94.   H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078–2079 (2009).

95.   O. R. Homann, A. D. Johnson, MochiView: versatile software for genome browsing and
DNA motif analysis. *BMC Biol.* **8**, 49 (2010).

96.   Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

97.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

Figure Titles and Legends

Fig. 1. Mfd functions as an RNAP co-factor and requires transcription elongation for

association with DNA

(A) ChIP-seq plot of *B. subtilis* Mfd tagged with 1x myc (red) and of WT *B. subtilis* (light green)

using myc antibody. (B) ChIP-seq plot of WT *B. subtilis* RpoB. (C) ChIP-seq plot of *B. subtilis*

MfdL522A-myc point mutant. (D) *B. subtilis* Mfd-myc and (E) *B. subtilis* RpoB ChIP-seq after

treatment with 50µg/mL of rifampicin for five minutes. Plots are normalized to total DNA input

controls and are the average of at least two independent experiments.

Fig. 2. Mfd directly promotes release of RNAP *in vivo*

(A) RpoB ChIP-seq plots showing regions of RpoB enrichment in *Δmfd*. Top half of graph

(counts in green) reflects normalized RpoB ChIP-seq read counts where *B. subtilis Δmfd* had

increased signal relative to WT. Bottom half of graph (counts in red) reflects RpoB ChIP-seq

read counts where *Δmfd* had decreased signal relative to WT *B. subtilis*. High background

signal from ribosomal RNA was removed from plots for better visualization. (B) Scatter plot of

WT and *Δmfd* RpoB ChIP-seq measuring signal at each gene in *B. subtilis*. For quantification of

ChIP signal at each gene, read counts for each gene were normalized to total library counts and

IP samples were normalized to total DNA input to calculate an IP/Total DNA ratio. Ratios were

$\log_2$ normalized and averaged over at least two independent experiments. Data points above

and below colored shading indicate greater than two-fold increase and decrease in RpoB signal

in the *Δmfd* strain*,* respectively. Data points in red indicate genes bound by Mfd. Binding is

defined as one standard deviation greater than the average ChIP signal across all genes in *B.*

subtilis. Calculation of Mfd binding at each gene was determined as described for RpoB ChIP

samples.

Fig. 3. Transcription units with Mfd binding and increased RNAP density in *Δmfd* are

enriched for structured regulatory RNAs

(A) Scatter plot of the minimum free energy (MFE) Z-score for regulatory RNAs in *B. subtilis*.

Data points represent regulatory RNAs within TUs that have no observed change in RpoB

density between WT and *Δmfd* (grey points) and TUs that have increased RpoB density in *Δmfd*

and are also bound by Mfd (red points). The three scatter plots represent all regulatory RNAs

(left), only expressed regulatory RNAs (middle) and top 50% expressed regulatory RNAs

(right). Expression data determined from RNA-seq analysis. (B) Bar graph of regulatory RNAs

within TUs that have increased RpoB density in *Δmfd* and are bound by Mfd. Non-expressed

RNAs shown in light grey and top 50% of expressed RNAs shown in dark grey. Error bars

represent the standard error of the mean (SEM). Statistical significance was determined using

two-tailed Z-test for two population means (****p<0.0001).

Fig. 4. Genes with Mfd binding and increased RNAP density in *Δmfd* are enriched for

RNAP pause sites

(A) Read counts for RpoB ChIP-seq (red) and NET-seq (blue) across a 30kb window for four

representative genomic regions (*yabE/S25*, *manP* operon, *txpA*/ratA, and a tRNA locus).

Zoomed in plots showing the gene/operon only (y-axis scale is preserved). (B) Scatter plot of

the average NET-set read counts for genes with decreased RpoB association in *Δmfd* (black

dots, n=53) and increased RpoB association in *Δmfd* (red dots, n=52). Median and 95%

confidence interval shown. Statistical significance was determined using the nonparametric

Mann-Whitney test for two population medians (**** p<0.0001). (C) Line plot showing by genes

with increased (light grey) and decreased (dark grey) RpoB association in *Δmfd*, ranked by

increasing signal. Red data point in both plots represents the mean NET-seq read count across

the genome. (D) Average number of pause sites (per 100bp) for all genes in *B. subtilis* (black

bar), and genes with increased (red bar) and decreased RpoB association (grey bar) in *Δmfd,*

respectively. (E) histogram showing the relative frequency of pause sites (per 100bp) in genes

with increased (red bars) and decreased (grey bars) RpoB association in *Δmfd*.

Fig. 5. Genome wide Mfd association is correlated with sites of RNA secondary structure

based on PARS seq in *E. coli*

(A) Overlay of sequencing counts from Mfd ChIP-seq (dark blue) and PARS-seq (red) across

four 100kb regions across the *E. coli* genome. ChIP-seq plots averaged from at least two

independent experiments. (B) Linear regression analysis comparing Mfd ChIP-seq and PARS-

seq in *E. coli*. Read counts from ChIP-seq and PARS-seq were used to calculate the area under

the curve (AUC) for each gene in the *E. coli* genome. Pearson's correlation coefficient for *E. coli*

ChIP-seq and PARS-seq r = 0.6. Dotted lines represent 95% confidence intervals.

Fig. 6. Transcriptional regulation by Mfd at toxin-antitoxin loci is critical for cell survival

Survival assays under chronic (A) and acute (C) overexpression of TxpA toxin (MFE Z-score =

-6.77) in WT and *Δmfd* and survival assays under chronic (B) and acute (D) overexpression of

BsrH toxin (MFE Z-score = -7.51) in WT and *Δmfd*. For all figures, representative images shown

above, and quantification of data shown below. Error bars represent the SEM from at least three

independent experiments. (E and F) qRT-PCR analysis of txpA and bsrH overexpression in WT

and *Δmfd* strains. RNA values normalized to ribosomal RNA. Error bars represent the SEM from

at least two independent experiments. Statistical significance was determined using a two-tailed

Student's T-test (*p<0.05, **p<0.01, ***p<0.001).

Fig. 7. Model of Mfd activity at structured regulatory RNAs

During transcription, elongating RNAP (shown in blue) transcribes a highly structured RNA

sequence. The formation of a structured RNA within the exit channel of RNAP can pause RNAP

on DNA, leading to Mfd binding (shown in red). Two (non-mutually exclusive) models are shown

to explain Mfd's function once RNAP encounters a highly structured RNA. Bottom left: the

arrested complex is recognized by Mfd, which releases RNAP from the template and in doing so

represses transcription or bottom right: Mfd recognizes the arrested or stalled RNAP complex

and functions as an elongation factor, accelerating RNAP release off of the DNA template.

Tables

Table 1. Summary list of genes and previously defined TUs bound with changes in RpoB

density in *Δmfd* from ChIP-seq analysis. Changes in RpoB density and Mfd binding at TUs are

defined by changes in one or more genes corresponding to its associated TUs

	Increased RpoB density in *Δmfd*	Decreased RpoB density in *Δmfd*
Genes (and TUs)	116 (71)	53 (31)
Genes (and TUs) bound by Mfd	52 (35)	0 (0)

Fig. 1

Fig. 2

A

B

Fig. 3

Fig. 4

Fig. 5

Fig. 6

A Chronic TxpA overexpression

B Chronic BsrH overexpression

C Acute TxpA overexpression

D Acute BsrH overexpression

E

F

Fig. 7

Elongating RNAP

Mfd binds RNAP

RNAP stalls due to RNA
secondary structure

RNAP release   and/or   RNAP rescue