Title: Neurobiological underpinnings of rapid white matter plasticity during intensive

reading instruction

Abbreviated title: Neurobiology of white matter plasticity

Authors: Elizabeth Huber[1], Aviv Mezer[2], Jason D. Yeatman[3,4]

[1]Institute for Learning and Brain Sciences and Department of Speech and Hearing

Sciences, University of Washington, Seattle, WA, 98195

[2]Edmond & Lily Safra Center for Brain Sciences, The Hebrew University of Jerusalem,

Jerusalem, Israel

[3]Graduate School of Education, Stanford University, Stanford, CA, 94305

[4]Division of Developmental-Behavioral Pediatrics, Stanford University School of

Medicine, Stanford, CA, 95305.

Corresponding Author: Elizabeth Huber, ehuber@uw.edu

Conflict of Interest: The authors declare no financial or other conflict of interest.

Acknowledgements: This work was funded by NSF/BSF BCS #1551330/ #2015608,

Eunice Kennedy Shriver National Institute of Child Health and Human Development

Grant R01 HD09586101 and Jacobs Foundation Research Fellowship to JDY.

Abstract

Human white matter is remarkably plastic. Yet it is challenging to infer the biological underpinnings of this plasticity using non-invasive measurements like diffusion MRI. Here we capitalize on metrics derived from diffusion kurtosis imaging (DKI) to interpret previously reported changes in mean diffusivity throughout the white matter during an 8-week, intensive reading intervention. We then use an independent quantitative MRI measurement of R1 (1/T1 relaxation time) in the same white matter regions; since R1 closely tracks variation in myelin content, it provides complementary information about white matter microstructure. Behavioral measures, multi-shell diffusion MRI data, and quantitative T1 data were collected at regular intervals during the intervention in a group of 33 children with reading difficulties (7-12 years old), and over the same period in an age-matched non-intervention control group. Changes in DKI parameters modeled over the intervention were consistent with increased hindrance in the extra-axonal space, rather than a large-scale change in axon density and/or myelination. Supporting this interpretation, analysis of R1 values did not suggest a change in myelin, although R1 estimates were correlated with individual differences in reading skill. Together, these results suggest that large-scale changes in diffusivity observed over a short timescale during an intensive educational experience are most likely to reflect changes occurring in the extra-axonal space, in line with recent work highlighting the role of glial cells in experience-dependent plasticity and learning.

Keywords: Diffusion MRI; quantitative MRI; white matter modeling; plasticity; reading

Introduction

Experience can modify the microstructure of the white matter over remarkably short timescales (Taubert, Draganski et al. 2010, Blumenfeld-Katzir, Pasternak et al. 2011, Engvig, Fjell et al. 2012, Hofstetter, Tavor et al. 2013, Mamiya, Richards et al. 2016, Huber 2018). The majority of past work examining white matter plasticity in humans has relied on diffusion MRI (dMRI) and metrics derived from the diffusion tensor model (DTI; Basser et al., 1996; Beaulieu 2002), which are sensitive to myriad features of the white matter, including the number, size and branching of glial cells, the density and caliber of axons, the abundance of myelin, and the spatial arrangement of fibers within an imaging voxel (Basser and Pierpaoli 1996, Alexander, Lee et al. 2007, Walhovd, Johansen-Berg et al. 2014). The diffusion kurtosis model (DKI; Jensen et al., 2005) offers increased sensitivity to microstructural variation and, potentially, increased specificity, by modeling variation in the diffusion signal that is not considered by the tensor model (Cheung et al., 2009; Veraart et al., 2010; Steven et al., 2014), although DKI metrics (e.g., mean kurtosis, MK) are generally no more straightforward to interpret than those derived from DTI (e.g., mean diffusivity, MD). Thus, a challenge for diffusion MRI studies of white matter plasticity is to understand the neurobiological underpinnings of the observed changes in diffusion properties since a variety of distinct mechanisms could account for the data.

Over the last decade, a host of modeling approaches have been developed to exploit biologically informed priors in pursuit of a more accurate and interpretable parcellation of the diffusion signal (Jelescu and Budde 2017, Alexander, Dyrby et al. 2019). For example,

the recent white matter tract integrity (WMTI) model (Fieremans, Novikov et al. 2010,

Fieremans, Jensen et al. 2011) builds on DKI by defining separable axonal and extra-axonal

contributions to the diffusion signal, which are explicitly modeled to provide metrics such

as axonal water fraction (AWF) and extra-axonal diffusivity. These parameters have

previously been shown to be sensitive to individual differences in white matter as a

function of age (Chang, Owen et al. 2015, Jelescu, Veraart et al. 2015, Genc, Malpas et al.

2017) and cognitive performance (Chung, Fieremans et al. 2018), as well as white matter

pathology (Fieremans, Benitez et al. 2013, Benitez, Fieremans et al. 2014), including de-

myelination (Falangola, Guilfoyle et al. 2014, Jelescu, Veraart et al. 2015, Guglielmetti,

Veraart et al. 2016, Jelescu, Zurek et al. 2016, Kelm, West et al. 2016). With appropriate

pre-processing, AWF and extra-axonal diffusivity are highly reliable, even given the

constraints associated with data collection in young children (Huber 2018).

Microstructural modeling of the diffusion signal thus holds promise for illuminating the

biological underpinnings of white matter development, pathology, and plasticity. However,

even the most sophisticated models incorporate simplifying assumptions about the

underlying tissue, and many require that the user to set constraints, such as the absolute or

relative expected diffusivity of individual tissue compartments (Jelescu, Veraart et al.

2015). Although it is often possible to make principled choices, in some cases, the most

appropriate set of model assumptions is still up for debate: For example, while the WMTI

model has previously been validated under the assumption that extra-axonal diffusivity is

greater than intra-axonal diffusivity (Guglielmetti, Veraart et al. 2016, Jelescu, Zurek et al.

2016), recent work suggests that the opposite assumption may be more appropriate (intra-

axonal greater than extra-axonal diffusivity: (Jespersen, Olesen et al. 2018, Kunz, da Silva

et al. 2018). Thus, even state-of-the-art modeling techniques may be open to multiple

interpretations and, while model parameters can be used to inform new linking hypotheses,

it is not possible to directly link changes in any one parameter to a specific change in tissue

biology.

Multi-modal studies that combine microstructural modeling with complementary

measurements, such as quantitative T1 mapping, can provide critical insight beyond what

can be gleaned from diffusion alone (Cercignani and Bouyagoub 2018, Filo, Shtangel et

al. 2019, Takemura, Ogawa et al. 2019, Travis, Castro et al. 2019). For example, in a large

cross-sectional study, complementary DTI and quantitative T1 relaxation measurements

showed distinct trajectories over the lifespan (Yeatman et al., 2014). Quantitative T1

measurements capture the interaction between tissue density and the chemical composition

of that tissue (Mezer, Yeatman et al. 2013, Stuber, Morawski et al. 2014, Filo, Shtangel et

al. 2019), such that T1 values are more attenuated in highly myelination regions, while

diffusion measurements are influenced by any cell membrane capable of hindering the

diffusion process(Le Bihan 1995). Thus, the authors reasoned that the discrepancy between

measures could reflect changes in various non-neuronal cell types (microglia,

oligodendrocyte precursor cells (OPC), and/or astrocytes). Such changes are presumed to

occur alongside changes in myelination, but with distinct dynamics over the lifespan.

Experience-dependent plasticity in the white matter likely depends on processes no less

complex than those unfolding over maturation and aging. Learning related changes in white

matter diffusivity could theoretically reflect a number of distinct biological phenomena, such as activity-dependent changes in myelination or glial cell proliferation (Blumenfeld-Katzir, Pasternak et al. 2011, Lerch, Yiu et al. 2011, Sagi, Tavor et al. 2012, Sampaio-Baptista, Khrapitchev et al. 2013, Gibson, Purger et al. 2014). In animal models, short-term learning has been associated both with remodeling of myelin and proliferation of glial cells in the gray and white matter (Blumenfeld-Katzir, Pasternak et al. 2011, Lerch, Yiu et al. 2011, Sagi, Tavor et al. 2012, Sampaio-Baptista, Khrapitchev et al. 2013). In the white matter, the size of non-axonal effects can even exceed subsequent changes in myelination (Gibson, Purger et al. 2014), which has been interpreted as reflecting an initial over-production of glial cells associated with the early stages of the learning process. Although non-invasive MRI measurements cannot directly specify the exact cellular changes that are occurring, they can be used to reason about the underlying biology of plasticity measured in the human brain, and they provide a vital link between the literature on human learning and invasive studies examining plasticity in other species.

We previously reported that 8 weeks of intensive reading instruction prompts widespread changes in white matter diffusion properties, which track the learning process (Huber 2018). Given that these effects occurred rapidly, and were distributed throughout the white matter, we speculated that the changes in diffusivity might reflect an initial stage of the learning process (e.g., large-scale glial cell proliferation), rather than changes to signal conduction properties of functionally relevant axons via changes in myelination or axon caliber. Here, we first confirmed that the previously reported finding of spatially distributed intervention-driven changes in diffusivity holds in a larger sample of subjects (n=33). We

then fit the WMTI model to these data to test for intervention-driven changes in the

estimated axonal water fraction (AWF). Although we observed changes in diffusivity

within the extra-axonal space, we failed to detect changes in AWF. We next explored how

changing certain model assumptions affects the pattern of results in these data. Finally, we

used an independent quantitative MRI measurement of R1 (1/T1 relaxation time) to

examine the same white matter regions. Although R1 estimates were correlated with

individual differences in reading skill, analysis of R1 values did not suggest a change in

myelin content over the 8-week intervention period. Together, these results highlight the

value of multi-modal MRI data for constraining inferences about underlying white matter

biology, while pointing to the potential importance of non-neuronal cell types during the

early stages of learning.

Materials and Methods

Participants

A total of 149 behavioral and MRI sessions were conducted with a group of 33 children

ranging in age from 7 to 12 years, who participated in an intensive summer reading

intervention program. Of these subjects, 24 were included in a previous manuscript (Huber

2018). Members of the intervention group were recruited based on parent report of reading

difficulties and/or a clinical diagnosis of dyslexia. Multi-shell diffusion MRI and

behavioral data were collected before the intervention (baseline), after 3.62 (+/- 0.16)

weeks of intervention, after 6.71 (+/- 0.16) weeks of intervention, and at the end of the 8-

week intervention period. An additional 78 behavioral and MRI sessions were conducted with 29 participants, who were matched for age but not reading level. These subjects were recruited as a control group to assess the stability of our measurements over the repeated sessions. Control subjects participated in the same experimental sessions but did not receive the reading intervention. Some families of control group subjects were reluctant to commit to all four sessions, given that their children were not receiving an educational intervention. These families were given an opportunity to participate in 2 sessions. The interval for the two sessions was chosen in order to have balanced numbers of measurements at equivalent time points to the intervention group. In the intervention group, 5 subjects were unable to complete either the third of the fourth imaging session, and therefore participated in 3 sessions, total. The distribution of testing sessions for the intervention and control groups is summarized in **Table 1**.

	Baseline (0 days)	11-38	33-77	49-100
Intervention	33	33 (mean 25.36, std 6.70)	31 (mean 47.00, std 6.23)	30 (mean 72.70, std 10.91)
Control	29	20 (mean 22.25, std 6.51)	16 (mean 44.38, std 10.01)	12 (mean 58.33, std 5.31)

Table 1. Testing schedule for intervention and control groups. Experimental sessions were evenly spaced for each subject, with a baseline prior to the start of intervention, a second session within 11-38 days since the start of intervention (column 2), a third session within 33-77 days (column 3), and a fourth session within 49-100 days (column 4). Each cell gives the number of subjects sampled at each time bin. In the intervention group, some data sets are missing due to scheduling or data quality issues, while in the intervention group some subjects elected to complete only 2 sessions (see Methods for details). The mean number of days and standard deviation for each time bin are given in parenthesis.

All participants were native English speakers with normal or corrected-to-normal vision

and no history of neurological damage or psychiatric disorder. Subjects were screened

using a mock scanner to assess comfort and ability to hold still during the MRI sessions.

We obtained written consent from parents, and verbal assent from all child participants.

All procedures, including recruitment, consent, and testing, followed the guidelines of the

University of Washington Human Subjects Division and were reviewed and approved by

the UW Institutional Review Board. Subject demographics are given in **Table 2**.

	WJ-BRS	TOWRE	WJ-RF	WJ-CALC	WJ-MFF	Age (months)
Intervention	80.30/14.27	72.42/13.33	72.42/18.51	85.68/13.48	83.10/16.11	112.58/ 20.58
Control	96.55/19.59	86.74/22.31	91.74/22.33	97.45/13.72	92.67/18.00	117.03/ 14.38

Table 2. Demographic data for the intervention and non-intervention control groups. Each cell contains

the group mean and standard deviation for a given item. Subject groups were matched in age but not

reading skill. The first three columns give mean and standard deviation (mean / standard deviation)

within each group for three standard reading measures: The Basic Reading Skill composite from the

Woodcock Johnson Tests of Achievement (WJ-BRS), the Test of Word Reading Efficiency index

(TOWRE), and The Reading Fluency subtest of the Woodcock Johnson Tests of Achievement (WJ-RF).

Columns 4-5 give standard scores for the Woodcock Johnson Tests of Achievement Calculation (WJ-

CALC) and Math Facts Fluency (WJ-MFF), which measure efficiency and accuracy of math related

skills.

Reading intervention

Intervention subjects were enrolled in 8 weeks of the Seeing Stars: Symbol Imagery for

Fluency, Orthography, Sight Words, and Spelling (Bell 2007) program at three different

Lindamood-Bell Learning Centers in the Seattle area. The intervention program consists

of directed, one-on-one training in phonological and orthographic processing skills, lasting

four hours each day, five days a week. The curriculum uses an incremental approach,

building from letters and syllables to words and connected texts, emphasizing phonological

decoding skills as a foundation for spelling and comprehension. A hallmark of this

intervention program is the intensity of the training protocol (4 hours a day, 5 days a week)

and the personalized approach that comes with one-on-one instruction.

To test for longitudinal change in reading and non-reading (Calculation and Math Facts

Fluency subtests of the Woodcock Johnson Tests of Achievement) measures, we fit a linear

mixed effects model with a fixed effect of intervention time, in days (for control subjects,

this corresponds to days since the baseline session), and a random effect of subject. The

intervention group showed significant gains for all reading measures ($p < 0.05$ for WJ-

BRS; $p < 0.0001$ for TOWRE and WJ-RF). We found no significant growth in the non-

reading measures over the same time period. In a subset of reading-matched controls, who

theoretically have as much room to improve on the reading measures as the intervention

subjects (i.e., they are not approaching ceiling on the tests), we found no significant growth

in any reading measure. Within this group, the math fluency (WJ-MFF) did improve with

repeated testing ($p < 0.001$), presumably due to practice on the timed-test. Within the full

control sample (including typical and highly skilled readers), performance on timed tests

(TOWRE, WJ-RF, and WJ-MFF) improved with repeated testing (p < 0.01), although

reading accuracy (WJ-BRS) and calculation (WJ-CALC) showed no significant change.

Magnetic resonance imaging (MRI) acquisition protocol

All imaging data were acquired using a 3T Phillips Achieva scanner (Philips, Eindhoven,

Netherlands) at the University of Washington Diagnostic Imaging Sciences Center (DISC)

using a 32-channel head coil. An inflatable cap minimized head motion, and participants

were continuously monitored through a closed-circuit camera system.

Diffusion-weighted magnetic resonance imaging (dMRI) data were acquired at 2.0mm^3

spatial resolution with full brain coverage. Each session consisted of 3 DWI scans, one

with 32 non-collinear directions (b-value=800 s/mm^2), and a second with 64 non-collinear

directions (b-value=2,000 s/mm^2). Each of the DWI scans included 4 volumes without

diffusion weighting (b-value=0). We also collected one scan with 6 non-diffusion-

weighted volumes and a reversed phase encoding direction (posterior-anterior) to correct

for EPI distortions due to inhomogeneities in the magnetic field using FSL's topup tool

(Andersson, Skare et al. 2003). Additional pre-processing is carried out using tools in FSL

for motion and eddy current correction (Andersson and Sotiropoulos 2016). Data were

manually checked for imaging artifacts and excessive dropped volumes. Given that subject

motion can be especially problematic for the interpretation of group differences in DWI

data (Yendiki, Koldewyn et al. 2014), data sets with mean slice-by-slice displacement >

3mm are excluded from further analysis.

For quantitative T1 mapping, we followed protocol developed by (Mezer, Yeatman et al.

2013). We acquired 4 spoiled gradient echo recalled images using two different flip angles

(2 scans with 4° and 2 scans with 20°, all with TR = 14ms, TE = 2.3ms, and resolution of

1 mm^3). To correct the transmit coil inhomogeneity, we collected 4 spin echo inversion

recovery scans with EPI read-out (SEIR-EPI), with TR 6500, TE 6.46, inversion times of

50, 400, 1200, 2400 ms, and 2mm^2 inplane resolution with a slice thickness of 4 mm. We

then compared T1 fits estimated using the spoiled gradient echo images to fits estimated

using the unbiased (Barral, Gudmundson et al. 2010, Mezer, Yeatman et al. 2013, Mezer,

Rokem et al. 2016) SEIR-EPI images to characterize the inhomogeneity field and apply an

appropriate correction to the biased, high resolution spoiled gradient echo recalled images.

Modeling white matter tissue properties

All diffusion and quantitative T1 data were aligned to a common anatomical reference in

each subject's native, ACPC aligned space. Axonal water fraction and extra-axonal

diffusivities were modeled using the white matter tract integrity (WMTI) model

(Fieremans, Novikov et al. 2010, Fieremans, Jensen et al. 2011), after fitting the diffusion

kurtosis model (Jensen, Helpern et al. 2005). WMTI and DKI fitting was implemented in

DIPY (Garyfallidis, Brett et al. 2014). R1 maps were calculated by taking 1/T1 (seconds)

for each voxel.

All values were then mapped onto fiber tracts identified for each subject using the

Automated Fiber Quantification software package (Yeatman, Dougherty et al. 2012), after

initial generation of a whole-brain connectome using probabilistic tractography (MRtrix

3.0 (Tournier, Calamante et al. 2004)). Since the white matter tract integrity (WMTI)

assumes that fibers are relatively well aligned (Fieremans, Jensen et al. 2011), we followed

recommendations from previous work and restricted our analysis to voxels with fractional

anisotropy values greater than 0.3 (Jensen, Stickley et al. 2017, Jensen, McKinnon et al.

2017, Chung, Fieremans et al. 2018). Specifically, voxels with fractional anisotropy below

0.3 were removed and WMTI metrics were interpolated at each point on each fiber, and

then values were summarized along the fiber-tract core based on computing the median

value across fiber nodes. Our previous work has demonstrated that summarizing values

based on the median, rather than the mean, of WMTI metrics substantially increases the

reliability of an individual's data (Huber 2018). Data with outlying values (greater than 4

standard deviations from the sample mean) in the white matter for any of the fitted metrics

was excluded from further analysis. After excluding both outliers and individuals with

excessive motion (>3mm, see above), the final data set included 109 sessions from 32

intervention subjects and 66 sessions from 27 non-intervention control subjects.

Statistical Analysis

Statistical analysis was carried out using software written in Matlab (*draft code link:*

https://github.com/yeatmanlab/BioBasis.git). To assess change over the course of

intervention, we first averaged the middle 80% of each tract to create a single estimate of

each property for each subject and tract. We selected the middle portion to eliminate the

influence of crossing fibers near cortical terminations, and to avoid potential partial volume

effects at the white matter / gray matter border. Mean tract values were then entered into a

linear mixed effects model, with fixed effects of intervention time and a random effect of

subject. For quantifying intervention effects, we prefer to use 'hours of intervention', since

this variable directly reflects the intervention 'dose'. For analyses including the control

subjects, who did not participate in an intervention of any kind, we substitute 'session' for

'hours'. Since sessions were held at regular intervals, the two variables were highly

correlated (Pearson's $r = 0.98$, $p < 0.001$).

Results

Diffusion MRI

Significant changes in mean diffusivity (MD) were apparent throughout the white matter

in the intervention group (**Figure 1a-b**; tracts showing significant changes qFDR < 0.05).

In a group of age-matched control subjects who attended school as usual, we found no

significant changes in white matter diffusivity over the same time frame, and growth

estimates for the 16 pathways were distributed around zero (**Figure 1c**). A group

(intervention vs. non-intervention) by time (session number) interaction was significant (p

< 0.05, uncorrected) for the left arcuate and left inferior frontal-occipital fasciculus.

Figure 1. (A) White matter tracts are shown in two views   hemisphere, right hemisphere) for

an example subject. Color-coding is based on the coefficient   mean a linear mixed effects model

predicting mean diffusivity (MD) values as a function of in   ention hours at each location along

each tract. Numeric labels correspond to the tract names listed at right. (B) Coefficients from a

model (+/- 1 SE) substituting 'session' for 'hours of intervention', for the intervention group. Darker

shaded bars reflect significant change (qFDR < 0.05). (C) Results for the control subjects, using the

same statistical model as in (B). In the control group, growth estimates were distributed around zero

with none of the tracts showing a significant change in MD over an 8-week period.

Although mean diffusivity is a highly sensitive measure (De Santis, Drakesmith et al. 2014,

Huber 2018), it is not biologically specific. We next examined the effects of intervention

on parameters estimated from the WMTI model: axonal water fraction (AWF) and extra-

axonal mean diffusivity. As shown in **Figure 2**, intervention effects were limited to

parameters associated with the extra-axonal space: Extra-axonal MD effects mirror the MD

effects shown above. We saw no change in estimates of AWF over the intervention period.

Figure 2. Microstructural modeling of white matter plasticity. (A) Illustration of two scenarios in

which mean diffusivity would decline in a voxel: proliferation of glial cells within the extra axonal

space (top) or increasing axon caliber and myelination (bottom). (B, C) Plots show coefficients from

a linear mixed effects model predicting extra-axonal mean diffusivity and axon water fraction

(AWF) from intervention time (in hours; random effect of subjects). Tracts showing significant

change (*qFDR* < 0.05) are shaded.

We next examined how assumptions implemented in the WMTI model affect the results.

Typically, the WMTI model assumes higher diffusivity within the extra-axonal space

versus the intra-axonal space. This assumption is required for the model to converge on a

single solution (Fieremans, Jensen et al. 2011). However, recent work has called this

assumption into question (Jespersen, Olesen et al. 2018, Kunz, da Silva et al. 2018). Thus,

to examine how this choice might affect our interpretation of the data, we perform a

supplementary analysis in which we invert the assumed relationship between diffusivities

(intra- greater than extra-axonal). Inverting the assumed relationship between diffusivities

produced effects in intra-axonal diffusivity for the right IFOF and ILF, the left and right

SLF, and the right Arcuate (*p* < 0.05, uncorrected) and rendered the remaining effects non-

significant. Importantly, this change to the model does not affect our calculation of AWF

(Fieremans, Jensen et al. 2011), and so these results are the same under either set of

assumptions: We find no detectable difference in AWF over the course of the intervention.

Quantitative T1

The effects observed in the diffusion MRI data set suggest that learning related changes in

the white matter reflect increased hindrance of diffusion within the extra-axonal space,

rather than an increase in the total volume of myelin. We next test this idea using using a

separate qMRI data set. R1 measurements reflect both the total volume of tissue in a region,

and the molecular composition, such as lipid and iron, of that tissue (Mezer, Yeatman et

al. 2013, Stuber, Morawski et al. 2014, Filo, Shtangel et al. 2019). A decrease in the volume

of myelinated tissue (or, an increase in water) within a voxel would be associated with

lower measured R1 values, while an increase in myelinated tissue would be associated with

elevated R1 measurements. Consistent with the interpretation suggested by the DKI and

WMTI fits, we see no measureable change in R1 over the course of the intervention period

in either group (*qFDR* < 0.05, **Figure 3a-b**), even when using a more lenient statistical

threshold (*p* < 0.05, uncorrected). Meanwhile, tract-average R1 values measured at

baseline in the anterior callosal tract, left arcuate, and left and right ILF correlate with

individual differences in reading skill prior to the start of the intervention (*p* < 0.05,

uncorrected**, Figure 3c**), confirming that R1 is sensitive to behaviorally relevant properties

of the white matter.

Figure 3. Quantitative R1 values do not show significant change over the intervention period in the intervention group (A), or the non-intervention control group (B). Bar plots show coefficients from a linear mixed effects model predicting R1 from intervention time (session; random effect of subject). (C) Baseline R1 values correlate pre-intervention reading performance. Color coding reflects the coefficient from a linear model predicting tract-average R1 values from (age-standardized) pre-intervention reading scores. Regions without a significant ($p < 0.05$) relationship to behavior are colored gray.

Discussion

Here we use diffusion MRI and a model derived from diffusion kurtosis imaging (Fieremans, Jensen et al. 2011), alongside quantitative R1 (1/T1 relaxation time) measurements, to examine whether microstructural changes during a successful reading intervention are best interpreted as rapid, experience-dependent changes in the volume of axons and myelin, or as changes in diffusivity within the extra- and/or intra-axonal space. Our results support the latter interpretation. While we observed systematic changes in diffusivity during the intervention, we failed to detect a longitudinal change in restricted fraction of the diffusion signal (interpreted as axonal water fraction, AWF), which would

be expected to arise from a sufficiently large change in myelination (Jelescu, Zurek et al.

2016), or a change in axon caliber. Although we interpret this null result cautiously, we

have previously demonstrated that AWF can be estimated reliably in this age group, and

that AWF is highly sensitive to maturational changes that occur over the timescale of years

(Huber 2018). Consistent with the diffusion MRI results, we did not observe a change in

quantitative R1 values over the intervention period. However, in the same individuals (both

intervention and non-intervention control subjects), we found that R1 was correlated with

pre-intervention reading skill. This argues that R1 measurements capture behaviorally

relevant variation in white matter microstructure that are distinct from the biological

mechanisms associated with short-term plasticity and learning during an intensive

intervention.

We previously (Huber 2018) reported changes in mean diffusivity alongside growth in

reading skills during an intensive reading intervention. If these effects indeed reflect

properties of the extra-axonal space, what is the link to behavior? A systematic reduction

in extra-axonal diffusivity, without corresponding changes in axonal water fraction or

quantitative R1, could result from an increase in cell membranes hindering diffusion within

the extra-axonal space, without a corresponding change in the total volume of that space.

This might reflect proliferation of oligodendrocyte precursor cells (OPC), as seen in

previous animal work (Gibson, Purger et al. 2014), although extra-axonal diffusivity

estimates could also be influenced by changes in the size or distribution of astrocytes

(Sampaio-Baptista and Johansen-Berg 2017, Sepehrband, Cabeen et al. 2018).

The relationship between higher-level cognitive function, axonal, glial and vascular

properties is likely to be complex, and the process of maintaining and optimizing signaling

properties involves a number of distinct biological phenomena that operate over different

time scales. However, it is increasingly clear that activation and proliferation of glial cells

and their precursors is vital not only for maintenance of active connections within a circuit,

but also for the optimization of signaling properties. For example, oligodendrocytes

participate in myelin maintenance, repair, and use-related plasticity throughout the lifespan

(reviewed in (Nave 2010)), and have been shown to regulate axon caliber directly,

independent of myelination (Sanchez, Hassinger et al. 1996). This mechanism could

theoretically support fast, activity-dependent changes in signaling efficiency and

coordination. Neural activity, in turn, appears to promote oligodendrocyte precursor

proliferation (Barres and Raff 1993), perhaps increasing the potential malleability of highly

active circuits. Although the link between these biological phenomena and learning is not

yet well understood, animal models have demonstrated a critical role of glia for brain

function in health, and their dysfunction in disease states (Barres 2008). Studies employing

longitudinal measurements, coupled with increasingly sophisticated imaging and modeling

techniques, hold promise for revealing the interplay among distinct biological processes

that support learning and cognition. Our findings highlight the importance of considering

the often-ignored contribution of glial cells to diffusion measurements and associated

cognitive functions in humans across the lifespan.

Linking the diffusion process to tissue biology requires making certain assumptions about

the factors that contribute to measured diffusion signals (Novikov, Kiselev et al. 2018).

The WMTI model used here assumes well-aligned fibers as its inputs (Fieremans, Novikov

et al. 2010). We have tried to assure that this assumption is met in our analysis by sampling

voxels with fractional anisotropy values that fall within a range for which the model has

been validated (Jensen, McKinnon et al. 2017, Chung, Fieremans et al. 2018). In order to

fit the WMTI model with a single solution, one must further assume a higher rate of

diffusivity within either the intra-axonal or extra-axonal space for a given region of interest.

Previous work has validated the assumption of higher intrinsic diffusivity within the extra-

axonal-space (Guglielmetti, Veraart et al. 2016, Jelescu, Zurek et al. 2016), although recent

work has called this assumption into question (i.e., intra-axonal diffusivity may be greater

than extra-axonal diffusivity; (Jespersen, Olesen et al. 2018, Kunz, da Silva et al. 2018)).

Importantly, our calculation of AWF is robust, and does not depend on this assumption;

thus, our main finding of stable AWF does not depend how the relative diffusivities are

constrained. However, this choice does determine whether intervention-driven changes in

diffusivity are attributed to the intra- versus extra-axonal space, in some cases. Although it

is less clear what mechanism would alter the intrinsic diffusivity of the intra-axonal space

over the timescales considered here, it is important to note that we cannot differentiate

between these scenarios based solely on the WMTI model. Thus, our modeling results

should not be interpreted as conclusive evidence for any specific biological mechanism.

Maturational differences in diffusion within the white matter reflect pruning of connections

and changes in myelination that occur over the timescale of years (Chang, Owen et al.

2015, Jelescu, Veraart et al. 2015), which may in turn influence reading outcomes

(Yeatman, Dougherty et al. 2012). The current data support the notion that short-term

changes in diffusion properties reflect an initial stage of the learning process, but the

connection between rapid changes and long-term remodeling of axons and myelin is

currently unknown. Resolving the relationship between learning and plasticity at temporal

scales ranging from hours (Sagi, Tavor et al. 2012, Hofstetter, Tavor et al. , Hofstetter,

Friedmann et al.), to days (Huber 2018), to years (Yeatman, Dougherty et al. 2012, Wang,

Mauer et al. 2017), will require research aimed at forging a tighter link between education

and neuroscience.

References

Alexander, A. L., J. E. Lee, M. Lazar and A. S. Field (2007). "Diffusion tensor imaging of

the brain." Neurotherapeutics **4**(3): 316-329.

Alexander, D. C., T. B. Dyrby, M. Nilsson and H. Zhang (2019). "Imaging brain

microstructure with diffusion MRI: practicality and applications." NMR Biomed

32(4): e3841.

Andersson, J. L., S. Skare and J. Ashburner (2003). "How to correct susceptibility

distortions in spin-echo echo-planar images: application to diffusion tensor imaging."

Neuroimage **20**(2): 870-888.

Andersson, J. L. and S. N. Sotiropoulos (2016). "An integrated approach to correction

for off-resonance effects and subject movement in diffusion MR imaging."

Neuroimage **125**: 1063-1078.

Barral, J. K., E. Gudmundson, N. Stikov, M. Etezadi-Amoli, P. Stoica and D. G. Nishimura

(2010). "A robust methodology for in vivo T1 mapping." Magn Reson Med **64**(4):

1057-1067.

Barres, B. A. (2008). "The mystery and magic of glia: a perspective on their roles in

health and disease." Neuron **60**(3): 430-440.

Barres, B. A. and M. C. Raff (1993). "Proliferation of oligodendrocyte precursor cells

depends on electrical activity in axons." Nature **361**(6409): 258-260.

Basser, P. J. and C. Pierpaoli (1996). "Microstructural and physiological features of

tissues elucidated by quantitative-diffusion-tensor MRI." J Magn Reson B **111**(3):

209-219.

Bell, N. (2007). <u>Seeing stars</u>. San Luis Obispo, CA, Gander.

Benitez, A., E. Fieremans, J. H. Jensen, M. F. Falangola, A. Tabesh, S. H. Ferris and J. A.

Helpern (2014). "White matter tract integrity metrics reflect the vulnerability of late-

myelinating tracts in Alzheimer's disease." <u>Neuroimage Clin</u> **4**: 64-71.

Blumenfeld-Katzir, T., O. Pasternak, M. Dagan and Y. Assaf (2011). "Diffusion MRI of

structural brain plasticity induced by a learning and memory task." <u>PLoS One</u> **6**(6):

e20678.

Cercignani, M. and S. Bouyagoub (2018). "Brain microstructure by multi-modal MRI:

Is the whole greater than the sum of its parts?" <u>Neuroimage</u> **182**: 117-127.

Chang, Y. S., J. P. Owen, N. J. Pojman, T. Thieu, P. Bukshpun, M. L. Wakahiro, J. I. Berman,

T. P. Roberts, S. S. Nagarajan, E. H. Sherr and P. Mukherjee (2015). "White Matter

Changes of Neurite Density and Fiber Orientation Dispersion during Human Brain

Maturation." <u>PLoS One</u> **10**(6): e0123656.

Chung, S., E. Fieremans, N. E. Kucukboyaci, X. Wang, C. J. Morton, D. S. Novikov, J. F.

Rath and Y. W. Lui (2018). "Working Memory And Brain Tissue Microstructure: White

Matter Tract Integrity Based On Multi-Shell Diffusion MRI." <u>Sci Rep</u> **8**(1): 3175.

De Santis, S., M. Drakesmith, S. Bells, Y. Assaf and D. K. Jones (2014). "Why diffusion

tensor MRI does well only some of the time: variance and covariance of white matter

tissue microstructure attributes in the living human brain." <u>Neuroimage</u> **89**: 35-44.

Engvig, A., A. M. Fjell, L. T. Westlye, T. Moberget, O. Sundseth, V. A. Larsen and K. B.

Walhovd (2012). "Memory training impacts short-term changes in aging white

matter: a longitudinal diffusion tensor imaging study." <u>Hum Brain Mapp</u> **33**(10):

2390-2406.

Falangola, M. F., D. N. Guilfoyle, A. Tabesh, E. S. Hui, X. Nie, J. H. Jensen, S. V. Gerum, C.

Hu, J. LaFrancois, H. R. Collins and J. A. Helpern (2014). "Histological correlation of

diffusional kurtosis and white matter modeling metrics in cuprizone-induced corpus

callosum demyelination." NMR Biomed **27**(8): 948-957.

Fieremans, E., A. Benitez, J. H. Jensen, M. F. Falangola, A. Tabesh, R. L. Deardorff, M. V.

Spampinato, J. S. Babb, D. S. Novikov, S. H. Ferris and J. A. Helpern (2013). "Novel white

matter tract integrity metrics sensitive to Alzheimer disease progression." AJNR Am J

Neuroradiol **34**(11): 2105-2112.

Fieremans, E., J. H. Jensen and J. A. Helpern (2011). "White matter characterization

with diffusional kurtosis imaging." Neuroimage **58**(1): 177-188.

Fieremans, E., D. S. Novikov, J. H. Jensen and J. A. Helpern (2010). "Monte Carlo study

of a two-compartment exchange model of diffusion." NMR Biomed **23**(7): 711-724.

Filo, S., O. Shtangel, N. Salamon, A. Kol, B. Weisinger, S. Shifman and A. A. Mezer

(2019). "Disentangling molecular alterations from water-content changes in the

aging human brain using quantitative MRI." Nat Commun **10**(1): 3403.

Garyfallidis, E., M. Brett, B. Amirbekian, A. Rokem, S. van der Walt, M. Descoteaux, I.

Nimmo-Smith and C. Dipy (2014). "Dipy, a library for the analysis of diffusion MRI

data." Front Neuroinform **8**: 8.

Genc, S., C. B. Malpas, S. K. Holland, R. Beare and T. J. Silk (2017). "Neurite density

index is sensitive to age related differences in the developing brain." Neuroimage

148: 373-380.

Gibson, E. M., D. Purger, C. W. Mount, A. K. Goldstein, G. L. Lin, L. S. Wood, I. Inema, S.

E. Miller, G. Bieri, J. B. Zuchero, B. A. Barres, P. J. Woo, H. Vogel and M. Monje (2014).

"Neuronal activity promotes oligodendrogenesis and adaptive myelination in the mammalian brain." Science **344**(6183): 1252304.

Guglielmetti, C., J. Veraart, E. Roelant, Z. Mai, J. Daans, J. Van Audekerke, M. Naeyaert, G. Vanhoutte, Y. P. R. Delgado, J. Praet, E. Fieremans, P. Ponsaerts, J. Sijbers, A. Van der Linden and M. Verhoye (2016). "Diffusion kurtosis imaging probes cortical alterations and white matter pathology following cuprizone induced demyelination and spontaneous remyelination." Neuroimage **125**: 363-377.

Hofstetter, S., N. Friedmann and Y. Assaf (2017). "Rapid language-related plasticity: microstructural changes in the cortex after a short session of new word learning." Brain Struct Funct **222**(3): 1231-1241.

Hofstetter, S., I. Tavor, S. Tzur Moryosef and Y. Assaf (2013). "Short-term learning induces white matter plasticity in the fornix." J Neurosci **33**(31): 12844-12850.

Huber, E., Donnelly, P.M., Rokem, A., Yeatman, J.D. (2018). "Rapid and widespread white matter plasticity during an intensive reading intervention." Nature Communications.

Huber, E., Henriques, R.N., Owen, J.P., Rokem, A., Yeatman, J.D. (2018). "Applying biophysical models to understand the role of white matter in cognitive development." bioRxiv.

Jelescu, I. O. and M. D. Budde (2017). "Design and validation of diffusion MRI models of white matter." Front Phys **28**.

Jelescu, I. O., J. Veraart, V. Adisetiyo, S. S. Milla, D. S. Novikov and E. Fieremans (2015). "One diffusion acquisition and different white matter models: how does

microstructure change in human early development based on WMTI and NODDI?"

Neuroimage **107**: 242-256.

Jelescu, I. O., M. Zurek, K. V. Winters, J. Veraart, A. Rajaratnam, N. S. Kim, J. S. Babb, T.

M. Shepherd, D. S. Novikov, S. G. Kim and E. Fieremans (2016). "In vivo quantification

of demyelination and recovery using compartment-specific diffusion MRI metrics

validated by electron microscopy." Neuroimage **132**: 104-114.

Jensen, A., T. Stickley, W. Torrissen and K. Stigmar (2017). "Arts on prescription in

Scandinavia: a review of current practice and future possibilities." Perspect Public

Health **137**(5): 268-274.

Jensen, J. H., J. A. Helpern, A. Ramani, H. Lu and K. Kaczynski (2005). "Diffusional

kurtosis imaging: the quantification of non-gaussian water diffusion by means of

magnetic resonance imaging." Magn Reson Med **53**(6): 1432-1440.

Jensen, J. H., E. T. McKinnon, G. R. Glenn and J. A. Helpern (2017). "Evaluating kurtosis-

based diffusion MRI tissue models for white matter with fiber ball imaging." NMR

Biomed **30**(5).

Jespersen, S. N., J. L. Olesen, B. Hansen and N. Shemesh (2018). "Diffusion time

dependence of microstructural parameters in fixed spinal cord." Neuroimage **182**:

329-342.

Kelm, N. D., K. L. West, R. P. Carson, D. F. Gochberg, K. C. Ess and M. D. Does (2016).

"Evaluation of diffusion kurtosis imaging in ex vivo hypomyelinated mouse brains."

Neuroimage **124**(Pt A): 612-626.

Kunz, N., A. R. da Silva and I. O. Jelescu (2018). "Intra- and extra-axonal axial

diffusivities in the white matter: Which one is faster?" Neuroimage **181**: 314-322.

Le Bihan, D. (1995). "Molecular diffusion, tissue microdynamics and microstructure."

<u>NMR Biomed</u> **8**(7-8): 375-386.

Lerch, J. P., A. P. Yiu, A. Martinez-Canabal, T. Pekar, V. D. Bohbot, P. W. Frankland, R.

M. Henkelman, S. A. Josselyn and J. G. Sled (2011). "Maze training in mice induces MRI-

detectable brain shape changes specific to the type of learning." <u>Neuroimage</u> **54**(3):

2086-2095.

Mamiya, P. C., T. L. Richards, B. P. Coe, E. E. Eichler and P. K. Kuhl (2016). "Brain white

matter structure and COMT gene are linked to second-language learning in adults."

<u>Proc Natl Acad Sci U S A</u> **113**(26): 7249-7254.

Mezer, A., A. Rokem, S. Berman, T. Hastie and B. A. Wandell (2016). "Evaluating

quantitative proton-density-mapping methods." <u>Hum Brain Mapp</u> **37**(10): 3623-

3635.

Mezer, A., J. D. Yeatman, N. Stikov, K. N. Kay, N. J. Cho, R. F. Dougherty, M. L. Perry, J.

Parvizi, H. Hua le, K. Butts-Pauly and B. A. Wandell (2013). "Quantifying the local

tissue volume and composition in individual brains with magnetic resonance

imaging." <u>Nat Med</u> **19**(12): 1667-1672.

Nave, K. A. (2010). "Myelination and support of axonal integrity by glia." <u>Nature</u>

468(7321): 244-252.

Novikov, D. S., V. G. Kiselev and S. N. Jespersen (2018). "On modeling." <u>Magn Reson</u>

<u>Med</u> **79**(6): 3172-3193.

Sagi, Y., I. Tavor, S. Hofstetter, S. Tzur-Moryosef, T. Blumenfeld-Katzir and Y. Assaf

(2012). "Learning in the fast lane: new insights into neuroplasticity." <u>Neuron</u> **73**(6):

1195-1203.

Sampaio-Baptista, C. and H. Johansen-Berg (2017). "White Matter Plasticity in the

Adult Brain." Neuron **96**(6): 1239-1251.

Sampaio-Baptista, C., A. A. Khrapitchev, S. Foxley, T. Schlagheck, J. Scholz, S. Jbabdi, G.

C. DeLuca, K. L. Miller, A. Taylor, N. Thomas, J. Kleim, N. R. Sibson, D. Bannerman and

H. Johansen-Berg (2013). "Motor skill learning induces changes in white matter

microstructure and myelination." J Neurosci **33**(50): 19499-19503.

Sanchez, I., L. Hassinger, P. A. Paskevich, H. D. Shine and R. A. Nixon (1996).

"Oligodendroglia regulate the regional expansion of axon caliber and local

accumulation of neurofilaments during development independently of myelin

formation." J Neurosci **16**(16): 5095-5105.

Sepehrband, F., R. P. Cabeen, J. Choupan, G. Barisano, M. Law and A. W. Toga (2018).

"A systematic bias in DTI findings." bioRxiv.

Stuber, C., M. Morawski, A. Schafer, C. Labadie, M. Wahnert, C. Leuze, M. Streicher, N.

Barapatre, K. Reimann, S. Geyer, D. Spemann and R. Turner (2014). "Myelin and iron

concentration in the human brain: a quantitative study of MRI contrast." Neuroimage

93 Pt 1: 95-106.

Takemura, H., S. Ogawa, A. A. Mezer, H. Horiguchi, A. Miyazaki, K. Matsumoto, K.

Shikishima, T. Nakano and Y. Masuda (2019). "Diffusivity and quantitative T1 profile

of human visual white matter tracts after retinal ganglion cell damage." Neuroimage

Clin **23**: 101826.

Taubert, M., B. Draganski, A. Anwander, K. Muller, A. Horstmann, A. Villringer and P.

Ragert (2010). "Dynamic properties of human brain structure: learning-related

changes in cortical areas and associated fiber connections." J Neurosci **30**(35): 11670-

11677.

Tournier, J. D., F. Calamante, D. G. Gadian and A. Connelly (2004). "Direct estimation

of the fiber orientation density function from diffusion-weighted MRI data using

spherical deconvolution." Neuroimage **23**(3): 1176-1185.

Travis, K. E., M. R. H. Castro, S. Berman, C. K. Dodson, A. A. Mezer, M. Ben-Shachar and

H. M. Feldman (2019). "More than myelin: Probing white matter differences in

prematurity with quantitative T1 and diffusion MRI." Neuroimage Clin **22**: 101756.

Walhovd, K. B., H. Johansen-Berg and R. T. Karadottir (2014). "Unraveling the secrets

of white matter--bridging the gap between cellular, animal and human imaging

studies." Neuroscience **276**: 2-13.

Wang, Y., M. V. Mauer, T. Raney, B. Peysakhovich, B. L. C. Becker, D. D. Sliva and N.

Gaab (2017). "Development of Tract-Specific White Matter Pathways During Early

Reading Development in At-Risk Children and Typical Controls." Cereb Cortex **27**(4):

2469-2485.

Yeatman, J. D., R. F. Dougherty, M. Ben-Shachar and B. A. Wandell (2012).

"Development of white matter and reading skills." Proc Natl Acad Sci U S A **109**(44):

E3045-3053.

Yeatman, J. D., R. F. Dougherty, N. J. Myall, B. A. Wandell and H. M. Feldman (2012).

"Tract profiles of white matter properties: automating fiber-tract quantification."

PLoS One **7**(11): e49790.

Yendiki, A., K. Koldewyn, S. Kakunoori, N. Kanwisher and B. Fischl (2014). "Spurious

group differences due to head motion in a diffusion MRI study." <u>Neuroimage</u> **88**: 79-

90.