Frequency-independent biological signal identification (FIBSI): A free program that simplifies intensive analysis of non-stationary time series data

Ryan M. Cassidy[1], Alexis G. Bavencoffe[2], Elia R. Lopez[2], Sai S. Cheruvu[2], Edgar T. Walters[2], Rosa A. Uribe[3], Anne Marie Krachler[4], Max A. Odem[4]*

[1]Medical Scientist Training Program, M.D. Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences, Houston, TX 77030, USA.

[2]Department of Integrative Biology and Pharmacology, McGovern Medical School at UTHealth, Houston, TX 77030, USA.

[3]Department of Biosciences, Rice University, Houston, TX, USA

[4]Department of Microbiology and Molecular Genetics, McGovern Medical School at UTHealth, Houston, TX 77030, USA.

*Correspondence should be addressed to: Max Odem: Phone: +1(713)500-5466; Email: Max.Odem@uth.tmc.edu

Ryan M. Cassidy contributed to conceptualization, data curation, funding acquisition, formal analysis, methodology, resources, software development, validation, visualization, and writing – original draft preparation.

Alexis G. Bavencoffe contributed to data curation, funding acquisition, investigation, resources, validation, and writing – review and editing.

Elia R. Lopez contributed to data curation, investigation, resources, validation, and writing – review and editing.

Sai S. Cheruvu contributed to software development, visualization, and writing – review and editing.

Edgar T. Walters contributed to conceptualization, funding acquisition, methodology, project administration, resources, supervision, and writing – review and editing.

Rosa Uribe contributed to funding acquisition, methodology, project administration, resources, supervision, and writing – review and editing.

Anne Marie Krachler contributed to conceptualization, funding acquisition, methodology, project administration, resources, supervision, and writing – review and editing.

Max A. Odem contributed to conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, software development, supervision, validation, visualization, and writing – original draft preparation.

Abstract (300/300 words)

Extracting biological signals from non-linear, dynamic and stochastic experimental data can be challenging, especially when the signal is non-stationary. Many currently available methods make assumptions about the data structure (e.g., signal is periodic, sufficient recording time) and modify the raw data in pre-processing using filters and/or transformations. With an agnostic approach to biological data analysis as a goal, we implemented a signal detection algorithm in Python that quantifies the dimensional properties of waveform deviations from baseline via a running fit function. We call the resulting free program frequency-independent biological signal identification (FIBSI). We demonstrate the utility of FIBSI on two disparate types of experimental data: *in vitro* whole-cell current-clamp electrophysiological recordings of rodent sensory neurons (i.e., nociceptors) and *in vivo* fluorescence image time-lapse movies capturing gastrointestinal motility in larval zebrafish. In rodent nociceptors, depolarizing fluctuations in membrane potential are irregular in shape and difficult to distinguish from noise. Using FIBSI, we determined that nociceptors from naïve mice generate larger, more frequent fluctuations compared to naïve rats, suggesting species-specific specializations in rodent nociceptors. In zebrafish, measuring gut motility is a useful tool for addressing developmental and disease-related mechanisms associated with gut function. However, available methods are laborious, technically complex, and/or not cost-effective. We developed and tested a novel assay that can characterize intestinal peristalsis using imaging time series datasets. We used FIBSI to identify muscle contractions in the fluorescence signals and compared their frequencies in unfed and fed larvae. Additionally, FIBSI allowed us to discriminate between peristalsis and oscillatory sphincter-like movements in functionally distinct gut segments (foregut, midgut, and cloaca). We conclude that FIBSI, which is freely available via GitHub, is widely useful for the unbiased analysis of non-stationary signals and extraction of biologically meaningful information from experimental time series data and can be employed for both descriptive and hypothesis-driven investigations.

Author Summary (172/200 words)

Biologists increasingly work with large, complex experimental datasets. Those datasets often encode biologically meaningful signals along with background noise that is recorded along with the biological data during experiments. Background noise masks the real signal but originates from other sources, for example from the equipment used to perform the measurements or environmental disturbances. When it comes to analyzing the data, distinguishing between the real biological signals and the background noise can be very challenging. Many existing programs designed to help scientists with this problem are either difficult to use, not freely available, or only appropriate to use on very specific types of datasets. The research presented here embodies our goal of helping others to analyze their data by employing a powerful but novice-friendly program that describes multiple features of biological activity in its raw form without abstract transformations. We show the program's applicability using two different kinds of biological activity measured in our labs. It is our hope that this will help others to analyze complex datasets more easily, thoroughly, and rigorously.

1. Introduction

It can be difficult to identify and characterize biological signals encoded within non-linear, dynamic

and stochastic experimental data using analytical methods without introducing bias due to *a priori*

assumptions about the data structure and nature of the signals. This is a challenge faced widely

across biological disciplines as many biological datasets are non-ideal (e.g., the frequency

resolution is low due to undersampling of the time domain, or the signal-to-noise ratio is low due

to limited measurement sensitivity) and the encoded signals are often of irregular shape. The

Fourier transform and derivative analytical tools [1], which are widely used to analyze biological

time series, have limited capacity to accurately identify and characterize biological signals

encoded in such data. A wealth of non-linear methods exist that aim at analyzing these kinds of

data series (e.g., using filters and transformations [2,3], multifractals for invariant structures [4]),

but most are field-specific and only applicable to a specific type of experimental data (e.g.,

electroencephalogram and electrocardiogram recordings [5,6]), and would require major

optimization prior to application to other types of experimental data with different data structures

and signal shapes. Information about a signal of interest is obtained from data that has been

subject to differing degrees of pre- and post-hoc processing. We contend that biologically

meaningful, easily interpretable information can be readily derived from signals in their raw form,

even if they are non-stationary.

In this paper, we demonstrate the utility of a novel frequency-independent biological signal

identification (FIBSI) program as a first-line tool for isolating biological signals from unprocessed

(raw) time series data. We demonstrate the utility of FIBSI using two different types of

experimental data series that were acquired using unrelated techniques: *in vitro* whole-cell

current-clamp electrophysiological recordings of depolarizing spontaneous fluctuations (DSFs) of

membrane potential from dissociated rodent putative C-type sensory neurons (primarily

nociceptors) and *in vivo* fluorescence time-lapse imaging data of motility in the larval zebrafish

intestine. Each type of data presents commonly encountered challenges that deter many

researchers. The DSFs in nociceptors are irregular in waveform and occurrence, small in

magnitude, and thus can be difficult to distinguish as a biologically relevant signal from internal

and external noise in neuronal recordings (i.e., low signal-to-noise ratios). For the analysis of gut

motility, the signal-to-noise ratios can vary substantially between regions of the intestinal tract and

between experimental animals. Motility frequencies can also differ within and between regions

within specimens. Challenges in quantifying fluorescence signals are similar to those encountered

with identifying electrophysiological signals, with the additional hurdle of reliably identifying a

region of interest (ROI) that shows the signal of interest without high dilution due to background

noise. Previously published methods circumvent these issues by different means. For example,

some focus on low frequency, high signal-to-noise ratio data collected from ROIs with substantial

physical separation from each other [7,8], use post-hoc analyses with a second identification

method (e.g., antibody staining or tight localization to neuronal nuclei) to tie ROIs to relevant

physical locations [9,10], use a short stimulus and test window to isolate activity [11], and/or use

phasic stimulations to identify ROIs and the Fourier transform to reduce noise [12–14]. In many

systems, however, the choice of ROIs is not flexible, and ROIs containing the relevant biological

signal of interest are intrinsically noisy, so the above-mentioned approaches are not applicable.

Here we demonstrate how FIBSI overcomes these common challenges without modifications to

the raw data and minimal ad-hoc assumptions. The signal (event) detection algorithm core to

FIBSI adapts the Ramer-Douglas-Peucker algorithm [15,16] to break down waveforms above and

below a time-dependent running fit function. The dimensional properties of the isolated signals

(events) can then be used for descriptive and hypothesis-driven statistical analyses.

2. Results

2.1. FIBSI allows an unbiased comparative analysis of membrane potential fluctuations in rodent

nociceptors

We have characterized two predominant types of small-diameter C-type sensory neurons in

rodents using whole-cell current-clamp electrophysiology: rapidly-accommodating (RA) and non-

accommodating (NA) neurons [17,18]. Here, we used a subset of RA and NA neurons dissociated

from 16 male Sprague-Dawley rats and 1 male and 4 female C57BL/6 wild-type mice for post-hoc

analysis. These rodents were used as naïve controls in prior studies performed by our research

group [18,19]. No major differences in electrophysiological properties were reported between

genders [18], so the neurons isolated from male and female mice were pooled. Like in previous

studies, these neurons were distinguished by injecting 2 s pulses of current at increasing

increments of 5 pA until each neuron reached rheobase by eliciting an action potential (AP). All

RA neurons elicited a single AP at the beginning of the pulse (**Fig 1A**), while a majority of NA

neurons exhibited delayed AP firing at rheobase (**Fig 1B-C**). At twice rheobase, RA neurons only

fired a single AP at the beginning of the pulse (**Fig 1D**), while NA neurons elicited repetitive firing

(**Fig 1E-F**). Additionally, both RA and NA neurons generated irregular, low-frequency DSFs in

membrane potential. In NA neurons, the DSFs were notably larger in amplitude and those that

reached threshold potential generated an AP. Prolonged depolarization to -45 mV increases the

likelihood of rodent nociceptors to be active and generate DSFs, and is used as an experimental

setting to model inflammatory and pain-like conditions *in vitro* [17,18]. However, no direct

comparisons of the DSFs across rodent species have been performed. To make the experimental

conditions comparable between neuron types and rodent species, we chose a subset of neurons

that were depolarized to -45 mV for 30 s [17,18] for FIBSI analysis. Both RA and NA neurons are

abundant in rats (~30% and ~70% respectively) [17], but RA neurons are sparse in mice (~6%)

[18], and too few had been sampled to warrant further analysis. Rat RA neurons did not exhibit

ongoing activity when depolarized to -45 mV, and DSFs were small in magnitude (**Fig 1G**). A

fraction (~20%) of the rat and mouse NA neurons exhibited ongoing activity and larger DSFs (**Fig**

1H-I), as reported previously [17,18]. We also tested for other signs of increased excitability in NA

neurons using standard membrane property measures. The mouse NA neurons had significantly

lower membrane capacitances, lower rheobase values, and depolarized resting membrane

potentials in comparison to rat NA neurons (**Fig 1J-L**), but their AP thresholds were similar (**Fig**

1M). In comparison to rat RA neurons, rat NA neurons had lower rheobase values and

hyperpolarized AP thresholds, like reported previously [17].

Fig 1. Membrane potential fluctuations, action potentials, and excitability properties in

nociceptors dissociated from rodents.

Example response patterns of RA and NA neurons at rheobase (A-C) and 2x rheobase (D-F). (A

inset) Neurons were incrementally injected with brief pulses of depolarizing current from a -60 mV

holding potential. (G-I) Prolonged depolarization to a -45 mV holding potential was used to

measure DSFs under comparable conditions across all neurons. Large DSFs marked in NA

neurons (red arrows). (J-M) Membrane properties measured in neurons (rat RA n = 20, rat NA n

= 27, mouse NA n = 28). The mouse NA neurons exhibited increased excitability compared to the

rat NA neurons as indicated by a significantly smaller membrane capacitance (J), lower rheobase

(K), and depolarized membrane potential (L). The rat RA neurons were less excitable with higher

rheobase values (K) and depolarized AP thresholds (M). Data in (J-M) shown as the median ±

95% CI with all points included. Data in (J) were analyzed using a 1-way ANOVA ($F_{(2, 72)}$ =

15.05, $P < 0.0001$) followed by Dunnett's multiple comparisons test. Data in (K) were analyzed

using a Brown-Forsythe 1-way ANOVA ($F_{(2.00, 30.78)} = 31.95$, $P < 0.0001$) followed by Dunnett's

T3 multiple comparisons test. Data in (L-M) were analyzed using Kruskal-Wallis tests (KW =

26.12, $P < 0.0001$ and KW = 16.37, $P = 0.0003$, respectively) followed by Dunn's multiple

comparisons test. **$P < 0.01$, ***$P < 0.001$. ANOVA, analysis of variance; CI, confidence interval;

MP, membrane potential.

Finally, it is assumed that signal analysis techniques like the fast Fourier transform (FFT)

capture the dominant frequencies associated with membrane potential oscillations (or fluctuations

as we refer to them, to include highly irregular changes in potential) in rodent sensory neurons

[20–27]. To test this assumption, a subset of the -45 mV recordings (n = 10 neurons per group)

were analyzed using the FFT. The dominant frequencies were highly variable (ranges: rat RA =

0.2-50 Hz, rat NA = 0.15-15 Hz, mouse NA = 0.15-10 Hz; FFT results are available online, see

Program and Data Availability section in Methods) and did not completely agree with the

published FFT results of 15-107 Hz for NA-like small-diameter neurons isolated from rats [20].

This suggested the FFT may not be suitable for making inferences about the DSFs.

Next, we used FIBSI to isolate the basic dimensional properties of all DSFs in the RA and

NA neuron datasets to determine whether DSFs differed between neuron types and rodent

species (**Fig 2A-C**). Current-clamp recordings had been performed using the same equipment as

reported previously [17,18], so we used published cutoffs (≥1.5 mV and ≥20 ms) to eliminate from

analysis low-amplitude signals that are likely noise from equipment. For FIBSI analysis, a running

median window of 1 s corresponding to 20,000 samples along the x-axis was used, and no filters

or transformations were applied to the raw data prior to signal detection.

Fig 2. Processing whole-cell current-clamp recordings to measure membrane potential

fluctuations in rodent nociceptors.

(A-C) Event detection workflow in a whole-cell current-clamp recording measured from a rat NA

neuron (APs are truncated). Order of operations: (A) calculate the running median (red line,

window = 1 s) based on the raw voltage and time coordinate data; (B) convert the running median

199 to y = 0 and calculate residuals, identify peaks of DSFs (blue X) that meet user-defined cutoffs

200 (≥3.0 mV amplitude, ≥20 ms duration in this example), and manually approximate the amplitude

201 of suprathreshold DSFs (red X) that produce an AP (see Methods for amplitude substitutions);

202 (C) visual confirmation of the DSF dimensional characteristics. Descriptive information for the

203 subthreshold DSFs include start, peak, and end times, amplitude, duration, and AUC (blue

204 highlight) to the converted running median at y = 0. Descriptive information for the AP waveform

205 (pink highlight) includes the aforementioned properties. (D) Scatter plot of DSF dimensions and

206 exponential plateau curves (cutoffs: ≥1.5 mV and ≥20 ms; rat RA n = 20 neurons, rat NA n = 27

207 neurons, mouse NA n = 28 neurons). Data were fit using exponential plateau models for each

208 neuron type; a single model did not fit all data ($F_{(6, 5689)}$ = 6.696, P < 0.0001). Refer to **S1 Table**

209 for model parameters. (E) The AUC of the DSFs ≥3 mV was significantly larger in rat NA neurons

210 compared to rat RA neurons. The DSFs in Mouse NA neurons were also significantly larger than

211 in rat NA neurons. (F) The NA neurons exhibited rightward shifts in DSF frequency (%) per

212 amplitude bin compared to the RA neurons. In NA neurons, DSFs ≥5 mV triggered APs and the

213 AP probability rapidly increased to 1 at 7-9 mV. Raster plots depicting frequencies of DSFs ≥3

214 mV in (G) rat RA neurons, (H) rat NA neurons, and (I) mouse NA neurons. The rat and mouse

215 NA neurons with suprathreshold DSFs and ongoing activity at -45 mV correspond to numbers 22-

216 27 and 23-28 on the y-axis of (H-I), respectively. (J) The mean frequencies for DSFs ≥3 mV were

217 significantly higher in the NA neurons. Data in (E) shown as the median ± 95% CI with all points

218 included. The raw AUC data collected for (E) were log transformed for the use of parametric

219 statistics. Transformed results were analyzed using a 1-way ANOVA ($F_{(2, 971)}$ = 38.90, *P*

220 <0.0001) followed by Sidak's multiple comparisons test. Data in (J) shown as medians (gray line)

221 and analyzed using a Kruskal-Wallis test (KW = 23.85, *P* < 0.0001) followed by Dunn's multiple

222 comparisons test. **P < 0.05. AUC, area under the curve.

To our knowledge, no statistical model has been reported that describes the relationship

between DSF amplitude and duration. Such a model would provide insight into the channel

opening and closing dynamics driving DSF generation in nociceptors. This model would also

provide a reference against which the effects of various pharmaceuticals can be compared and

future treatments for pain may be developed. In order to approach this is an unbiased manner,

we employed the Akaike information criterion to compare the quality of fit of several different

nonlinear regression models to the neuron datasets. Overall, the exponential plateau model was

a better fit for the RA and NA neurons compared to the Gompertz, Logistic, and Malthusian

models (see **S1 Table** for model parameters). We then asked whether the exponential plateau

model could fit all data points using the same set of fitting parameters, but a separate set of

parameters was necessary for each neuron type (F(6, 5689) = 6.696, *P* < 0.0001; **Fig 2D**). All

models tested, including the exponential plateau model for each neuron type, failed to pass for

homoscedasticity (*P* < 0.0001 for each model), and the 95% CI for the RA neuron model could

not be calculated by the GraphPad Prism software. These results suggest interpretations based

solely on statistical models should be made with caution, but they might provide some insight into

the relationship between DSF amplitude and duration.

Making comparisons based solely on DSF amplitude (like done previously [17,18]) limits

interpretation of the results because it omits any influence duration may have on DSF size, as the

DSFs do not exhibit regular waveforms. Therefore, we asked whether the total AUC of DSFs

differed between neuron types. A cutoff of ≥3 mV was chosen because it is at the lower bound of

the range of amplitudes that may be functionally relevant under pain-associated conditions when

neurons are depolarized [17,18]. The AUC of DSFs in rat NA neurons was larger than in RA

neurons, and the AUC of DSFs in mouse NA neurons was the largest (**Fig 2E**). We then binned

DSFs by amplitude to determine their frequency of occurrence (reported as a percentage of total

DSFs generated in each amplitude bin), and the amplitude necessary to generate APs (**Fig 2F**).

The mean frequency of DSFs per amplitude bin was higher in NA neurons than RA neurons for

rats. No DSFs triggered APs in the RA neurons, but a fraction of the large amplitude DSFs (≥5

mV) in NA neurons did trigger APs (**Fig 2F**). Mouse NA neurons tended to require larger DSFs to

trigger APs before the AP probability plateaued. Finally, we used raster plots to depict the

differences in frequencies of DSFs ≥3 mV between RA and NA neurons (**Fig 2G-I**). Few RA

neurons generated DSFs ≥3 mV (**Fig 2G**). The frequency of DSFs ≥3 mV was significantly higher

in NA neurons from mice (**Fig 2I**) than rats (**Fig 2H-J**). The frequencies of DSFs ≥3 mV detected

in NA neuron recordings by FIBSI were orders of magnitude lower than the oscillation frequencies

reported based on similar measurements using the FFT [20].

2.2. Using fluorescence contrasted visualization and FIBSI to analyze gut motility in the larval

zebrafish

Direct intestinal injections and oral gavage can be used to deliver dyes into zebrafish larvae

[28,29]. However, these methods are invasive, can be deterrents without adequate technical

expertise, and may directly influence motility. We first determined whether immersing awake

larvae in embryo medium (E3) media supplemented with Nile Red dye – without using an egg

emulsion to promote feeding [30] – was a viable non-invasive alternative for staining the intestinal

luminal space (**Fig 3A-B**). No acute or long-term (≥24 hours) effects of exposure on general

activity levels and behavior, and no mortality were observed at the Nile Red dose used (0.01

μg/mL). Video playback of the fluorescence time-lapses (**S2 Movies**) confirmed contractions

propagated at different rates in the different regions of the gut, and in opposing directions (i.e.,

retrograde and anterograde [31]) from a site of pacemaker-like activity near the end of the foregut

[32]. A total of 19 ROIs were evenly positioned along the foregut, midgut, and hindgut regions of

the intestinal tract (**Fig 3B-C**). We predicted that the Nile Red fluorescence intensity would change

periodically, reflecting contraction waves traveling through the gut (**Fig 3D-E**); retrograde

contractions coincide with a decrease in intensity in the foregut, anterograde contractions coincide

with an increase in intensity in the midgut and hindgut, and rapid movement of the cloaca coincide

with high-frequency oscillations in intensity in the hindgut. Indeed, contractions were visible in the

fluorescence intensity recordings as they propagated rhythmically through the intestinal tract at

<0.1 Hz, and the directional changes in intensity matched our predictions (**Fig 3F**).

Fig 3. Fluorescent-contrasted visualization of gut motility in larval zebrafish.

(A) Wild-type *AB* zebrafish larvae at 8 dpf. The intestinal tract is outlined and highlighted in blue.

(B) Merged bright field and red fluorescence (Nile Red dye) image with different functional

segments of the intestinal tract labeled. (C) Fluorescence image of (B) depicting positions of ROIs

used to measure changes in fluorescence intensity resulting from muscle contractions. (D-E)

Models and predicted time series depicting fluorescent fluctuations from baseline in ROIs during

propagation of retrograde contractions in the foregut and anterograde contractions in the

midgut/hindgut. The high-frequency movements of the cloaca were predicted to generate

oscillatory changes in fluorescence. (F) Time course of raw fluorescence intensity changes

measured from 20 ROIs in (C) in a single larva (traces color coded for visual differentiation). dpf,

days post-fertilization.

During our development process, we noticed that some recording periods were primarily

high fluorescence, with contractions signaled by a sudden reduction in fluorescence (a trough

event), whereas other recordings were primarily low fluorescence, punctuated by sudden

increases in fluorescence (a peak event). This is likely related to differing contraction physiology

in the foregut, midgut, and hindgut. In order to increase the comparability of these events and

standardize the identification of amplitude, we implemented a second normalization procedure

that traces a fitted line peak-to-peak (or trough-to-trough) of events identified in the first round of

normalization (i.e., sliding median). This new line is used as the reference against which the data

is compared and events identified again. The peak-to-peak function was applied to the foregut

(ROIs 1-7) and hindgut (ROIs 17-19), and the trough-to-trough function was applied to the midgut

(ROIs 8-16). The following running median window sizes were used: 50 x-samples for ROIs 1-7

and 10 x-samples for ROIs 8-19.

Processed fluorescence intensity data for the foregut (**Fig 4A-C**), midgut (**Fig 4D-F**), and

hindgut ROIs (**Fig 4G-I**) revealed several features: foregut contractions were overall larger in

amplitude, foregut and midgut contractions had similar durations, and the high-frequency

oscillations coinciding with cloaca movement in the hindgut were overall shorter and smaller.

These features mirrored observations made during video playback and predictions about signal

shape (**Fig 3D-E**). A cutoff of ≥0.05 amplitude (F/F0 units) and duration of ≥10 s (duration cutoff

inferred from published contraction frequencies and intervals, see [31,33–35]) was used to isolate

low-frequency muscle contractions (quadrant 4, Q4) in ROIs 1-19 from noise (Q1-3) Here, noise

refers to signal artifacts and movements of the gut not related to the muscle contractions of

interest observed in the videos. A separate cutoff of ≥0.05 amplitude and duration window of 5 s

≤ x ≤ 10 s was used to isolate the rapid movements of the cloaca present in ROIs 17-19. Raster

plots for Q1-3 (**S3 Figure**) did not depict rhythmic activity indicative of peristalsis in either unfed

or fed larvae. In contrast, raster plots for Q4 in unfed (**Fig 4J**) and fed larvae (**Fig 4K**) exhibited

key characteristics of peristalsis also observed in spatiotemporal maps [31]: anterograde (purple

arrows) and retrograde (green arrows) waves traveled rhythmically through the intestinal tract.

These waves originated from a pacemaker-like region near the end of the foregut (blue highlight)

[31,32].

We then asked whether the spatiotemporal information extracted from our time-lapse

datasets using FIBSI could be used to detect differences in gut motility between unfed and fed

larvae. A recent study using image velocimetry and spectral analysis reports differences in motility

between unfed and fed zebrafish larvae at 7 dpf [35], but the study is limited to the midgut, and

no other comparable datasets have been published. First, we confirmed that muscle contraction

frequencies did not differ between unfed *AB* (0.026 ± 0.001 Hz, n = 8) and unfed transgenic *Tg(-*

8.3phox2bb:Kaede) larvae (0.028 ± 0.001 Hz, n = 8; unpaired *t* test *P* = 0.158). Contraction

frequencies in the cloaca also did not differ between the two lines (0.046 ± 0.004 Hz and 0.051 ±

0.003 Hz, respectively; unpaired *t* test *P* = 0.407). Therefore, the two lines were pooled to increase

statistical power for comparison to fed *AB* larvae. Muscle contraction frequencies were higher in

fed compared to unfed larvae (**Fig 4L**).

Fig 4. Processing fluorescence time series datasets using FIBSI to analyze gut motility in

larval zebrafish.

Event detection workflow for fluorescence time series corresponding to ROIs #1-7 in the foregut

(A-C), #8-6 in the midgut (D-F), and #17-19 in the hindgut (G-I). The residuals were calculated

from the peak-to-peak or trough-to-trough trend line (red) and the trend line was normalized to y

= 0. Event peaks (blue X) were identified only for the waveforms with start/end times that crossed

y = 0 (red X). After processing, event durations and amplitudes were plotted for inspection and

filtering (1 dot = 1 event, events are color coded by ROI). Cutoffs (dotted red lines, amplitude of

≥0.05 and duration ≥10 s) were applied to isolate low-frequency contractions in Q4 of al 19 ROIs.

Oscillatory movement of the cloaca was isolated in ROIs 17-19 using a cutoff window (dotted

black line, amplitude cutoff of ≥0.05 amplitude and 5 s ≤ x ≤ 10 s duration). Raster plots of events

in Q4 and cloaca in unfed (J) and fed (K) larvae depict low-frequency retrograde (green lines) and

anterograde (purple lines) contractions traveling through the gut. Contractions originated from a

region of pacemaker-like activity near the end of the foregut (blue shading at ROIs #6-8). (L) Mean

contraction frequencies across the length of the whole gut (ROIs #1-19) were significantly

increased in fed (black circles) compared to unfed (white circles) larvae. (M) Fed larvae had

significantly increased mean contraction frequencies in the foregut (ROIs #1-7), but exhibited no

differences from unfed larvae along the midgut (ROIs #8-16) and hindgut (ROIs #17-19). (N)

Mean contraction frequencies for the cloaca were significantly decreased in fed larvae. Points in

(L-N) represent the mean per unfed (n = 16) and fed (n = 13) larvae and shown as the group

mean ± SEM. The initial comparison between groups in (L) was made using an unpaired t test

and gut region-specific comparisons in (M) were made using a 2-way ANOVA (Group factor F(1,

78) = 6.116, P = 0.0156; Region factor F(2, 78) = 188.6, P < 0.0001; Interaction F(2, 78) = 7.460,

P = 0.0011) followed by Sidak's post-hoc test. Data in (N) were compared using an unpaired t

test. *P < 0.05, **P < 0.01, ***P < 0.001. SEM, standard error of the mean.

We then asked whether motility in fed larvae was higher along the entire gut, or if the

increased motility was limited to a specific functional gut segment. The frequency of retrograde

contractions in the foregut was higher in fed compared to unfed larvae (**Fig 4M**). In contrast to the

prior study [35], we observed no differences between unfed and fed groups for anterograde

contractions in the midgut or hindgut (**Fig 4M**). Finally, we observed a significant decrease in the

frequency of cloaca movements in fed compared to unfed larvae (**Fig 4N**). The data collected

using our fluorescence contrast-based assay of gut motility, in conjunction with signal detection

and data analysis using FIBSI provided novel insights into individual functional gut segments of

larval zebrafish. We were able to characterize peristalsis along the whole intestinal tract as well

as high-frequency, low-amplitude oscillatory movements of the cloaca.

3. Discussion

3.1. Overview

In this study we introduce a new signal detection algorithm implemented in the free program FIBSI

that can be used to analyze a diverse array of non-linear, dynamic and stochastic experimental

datasets by extracting biologically meaningful signals from background noise. As experimental

biologists, we wanted to develop FIBSI to be particularly suited to the needs of other

experimentalists. As such, strengths of FIBSI include its capability to process non-ideal datasets

such as those we included here to demonstrate the utility of the program. These may include

datasets with low signal-to-noise ratios, irregular signal shapes, and low-resolution (i.e., under-

sampled) time series. In contrast to many other signal detection programs, FIBSI requires no

modification of the raw data, and minimal ad-hoc assumptions regarding data structure and signal

shape are needed prior to processing and signal detection. That is, FIBSI captures quantitative

information for all signals contained with the dataset. We validated the performance of FIBSI using

two different types of measurements: electrophysiological whole-cell current-clamp recordings

from rodent sensory neurons and fluorescence intensity imaging data capturing muscle

contractions in the larval zebrafish intestinal tract. Hypothesis testing based on prior biological

knowledge – two prior pain-related rodent studies [17,18] and established methods for measuring

intestinal motility in vertebrates [31–40] – was used to draw novel conclusions about the analyzed

datasets. The results of these tests are summarized and discussed below.

3.2 Membrane potential fluctuations differ between nociceptors in two rodent species

We were the first to automate the quantification of DSFs and demonstrate their functional

importance for AP generation in putative C-type nociceptors using the prototype algorithm upon

which FIBSI is based [17]. We showed that DSFs bridge the gap between the resting membrane

potential and AP threshold of a neuron and presumably play a role in driving ongoing pain-related

information from nociceptors [41,42]. We have also begun to identify associated cellular

mechanisms (e.g., dependence upon cell signals such as cyclic AMP-dependent protein kinase

and exchange protein activated by cyclic AMP-associated pathways) [17,18,43] to better

understand how DSFs are generated and under what conditions they can be inhibited. Others

have also noted their potential role in driving other pain-related conditions in rodents [44] and in

humans [45]. Given the limited research on DSFs, our current study is the first to make direct

comparisons between similar nociceptors in two rodent species commonly used in pain-related

studies. This was done to 1) confirm whether nociceptors in rats and mice function similarly, and

2) to validate FIBSI in its new, generalized form using the same data type as in our original studies

[17,18].

The NA neurons isolated from rats and mice recapitulated several of the specializations

we have described that promote ongoing activity and ongoing pain [17,18,41–43]. Under naïve

conditions (i.e., no injury), rat NA neurons are clearly more excitable than RA neurons. A novel

finding presented here is that mouse NA neurons are more excitable than rat NA neurons with

enhanced rheobase values and depolarized resting membrane potentials. A plausible explanation

for the species-specific differences in NA neuron excitability is that the neurons in mice may be

smaller in size, as suggested by the smaller membrane capacitance measures. This raises the

interesting question of whether the membrane densities and compositions of the channels

underlying the DSFs differ between species; this is of ongoing interest in the Walters lab and

under investigation. These specializations are expected to prime the NA neurons to activate more

readily in response to intra- and extracellular signals associated with nociceptive and potentially

painful stimuli. This hypothesis is further supported by our analysis of DSFs using FIBSI. The

mouse NA neurons generate larger, more frequent DSFs than rat NA neurons. However, the

greater DSF size and frequency, and necessity for larger-amplitude DSFs to generate APs in

mice compared to rats is unexpected because naïve rats and mice have similar AP voltage

thresholds while the mice have a more depolarized RMP compared to rats. We also compared

exponential growth models to describe the dimensional properties of the DSFs and found different

models were necessary for each neuron type. These models need to be compared further under

other pharmacological and pain-related conditions in order to better understand the differences

between species. Finally, the frequency results we obtained in our FFT analysis did not

recapitulate the results for the same type of neuron [20], and other studies do not fully report their

results for comparison [21–27]. One possible explanation for the discrepancy is that the longer

sampling duration used (30 s compared to 1-4 s in the original study [20]) gave a more

representative image of the fluctuations and activity in each neuron, which may influence the

measured frequencies. Using FIBSI, frequencies for DSFs ≥3 mV in NA neurons depolarized to -

45 mV peaked at 1-2 Hz. This finding is similar to our spinal cord injury and serotonin-induced

inflammatory conditions, under which frequencies for DSFs ≥5 mV peak at ~1.5 Hz [17]. Given

these similarities, it is unlikely we would observe a shift in the FFT signal to 20-107 Hz for NA

neurons depolarized to -45 mV, even if we used the same injury condition as the prior study

(sciatic nerve injury [20]). Taken together, these observations suggest the FFT is not an effective

tool for analyzing highly irregular DSFs in membrane potential, as opposed to the sinusoidal

oscillations examined primarily in large DRG neurons previously (most of which are not

nociceptors) [20–27]. Unlike the FFT, the descriptive data generated by FIBSI can be concretely

linked to precise DSFs of interest and dissociated from other biological and non-biological signals

in the current-clamp recordings.

Our findings suggest two assumptions that should be avoided when investigating DSF

mechanisms: 1) DSFs of one species are indistinguishable from DSFs of another, even in what

appear to be functionally equivalent nociceptors, and 2) DSFs in their measured form represent

unitary events. Future studies can test whether DSFs usually represent summation of discrete

events and if underlying unitary components can be identified based on waveform characteristics

(see [46,47]). The use of FIBSI and similar analytical approaches used to study DSFs will improve

our understanding of basic nociceptor physiology in multiple species and help to identify

fundamental mechanisms for driving pain that may be shared by rodents and humans.

3.3 Validation of a simple, fast, and cost-effective assay of gut motility in larval zebrafish

The current approach for assessing gut motility in vertebrates *in vivo* and *ex vivo* is to generate

spatiotemporal maps of the intestinal tract and measure changes in pixel opaqueness as indices

of movement [31,32,34,36–40]. Inspection of these maps can be done manually but the process

is laborious and possibly subject to human unconscious bias. More recent studies have improved

the analysis of gut motility in that the process has been semi-automated, which also enables

assessment of a wider range of parameters (e.g., contraction amplitude in μm). However, these

methods require custom microscopy equipment [48], complex image velocimetry and spectral

analysis [35], and/or use of customized "in-house" [49] or proprietary software [33,40]. Here, we

developed a simpler, faster, and more cost-effective assay using fluorescence contrast-based

imaging to visualize gut motility *in vivo*. The resulting time series datasets can be processed using

FIBSI to analyze muscle contractions. While 8- and 16-bit grayscale bright field images can also

be analyzed using FIBSI, fluorescence intensity measurements have a larger dynamic range, and

the resulting time series have a higher signal-to-noise ratio (i.e., contractions are more visible)

The advantage of FIBSI over existing programs used to analyze gut motility is its user-friendliness

and free availability.

Our attempts to validate the fluorescence contrast-based gut motility assay and the

performance of FIBSI against prior studies was challenging due to the fact that many reports do

not include sufficient experimental details (e.g., feeding regimen, gut region imaged, and

methodology used to calculate contraction frequencies). It is important that future studies are

explicit in their reporting to facilitate more direct comparisons between studies and method

development. Some studies measure motility in only select regions of the gut [29,35,50] while

others measure motility throughout the whole gut and pool the results [33]. We contend calculating

the frequency of retrograde and anterograde contractions separately [31,34] gives a more

accurate representation of gut function and how its luminal contents may be processed

mechanically. Our gut motility assay and FIBSI captured important characteristic features of

peristalsis also observed in the spatiotemporal maps: anterograde and retrograde contractions

originated from a site of pacemaker-like activity posterior to the intestinal bulb [31,32] near our

delineation of the foregut and midgut. Observation of these features increased our confidence in

the results. The contraction frequencies obtained using FIBSI recapitulated the published range

of frequencies for unfed and fed larvae [33,34,50]. The only published study to directly compare

gut motility in unfed and fed zebrafish larvae used measurements from the midgut, and it reports

feeding at 5-6 dpf increases motility at 7 dpf from 0.035 Hz to 0.041 Hz [35]. We observed similar

rates (~0.03 Hz) in the midgut, but feeding zebrafish live paramecia 1 day before testing did not

increase the rate of anterograde contractions in the midgut. Instead, we found that feeding

increased retrograde contractions in the foregut. Another novel observation was that feeding

decreased the rate of oscillatory movements of the cloaca. Adult shorthorn sculpin – another

teleost species – respond to feeding in a similar manner; orally-directed (retrograde) standing

contractions increase in frequency while slow, anally-propagating (anterograde) contractions

occur less frequently [49]. We did not observe a decrease in anterograde contractions, but it is

likely that the increase in retrograde contractions in sculpin and zebrafish serve a similar function

to displace and propel large quantities of luminal contents [49]. In zebrafish larvae, the oscillatory

motions of the cloaca may constitute sphincter-like contractions that control the timing of

defecation by keeping the luminal contents inside the fish rectum. Evidence suggests the caudal

segments of the adult zebrafish intestine may share functional similarities with the human rectum

[51], but to our knowledge there are no published reports of sphincter-like function in the hindgut

of larval zebrafish. It is possible that a reduction in the oscillations (i.e., relaxation) may facilitate

more frequent removal of waste products, but this has yet to be confirmed using transit

experiments. Our findings suggest that larval zebrafish are capable of generating functionally

complex gut motility patterns in response to feeding, possibly similar to other adult teleost species.

This adds to their tractability as a powerful model organism for *in vivo* imaging. The significance

of these gut region-specific phenomena is becoming increasingly apparent (for review [52]), but

these complex motility patterns require further attention using methods that yield functional

information of the whole gut. Future improvements to our motility assay and FIBSI could link

changes in contraction amplitudes to changes in the luminal dimensions of the intestinal tract,

similar to using image velocimetry to track movement [35].

There is a pressing need for improved imaging and analysis techniques that can be used

to address gut motility in altered physiological states and disease [53]. Analysis with FIBSI yields

experimentally valid and reproducible results, and it should be useful to researchers measuring

gut motility in zebrafish as a functional output of the enteric nervous system (ENS). The ENS

regulates multiple gastrointestinal functions (e.g., gut motility, hormone secretion). Deficiencies in

the development and function of the ENS can lead to debilitating neurological disorders (e.g.,

Hirschsprung's disease) [54–58]. Other factors of interest associated with ENS function (e.g.,

stress, microbiome, inflammation, visceral pain) [59–61] have traditionally been tested using

rodent models. Recent reviews have highlighted the validity of using zebrafish as an analogous

model to address some of the above questions [53,62–65], and the application of our motility

assay and FIBSI to this system will facilitate such research.

3.4. Current and future applications of FIBSI

We originally designed the prototype version of FIBSI (first report [17]) for the singular purpose of

quantifying DSFs and APs in nociceptors and incorporating the methodologies of others (e.g.,

estimating AP threshold [66]) within the field. However, collaborative efforts led us to realize the

broader applicability of the program to other researchers and types of datasets. Additional

features (e.g., filtering methods, down-sampling, trimming) are included for those interested, and

we encourage others to make modifications to the existing code (see availability section in

Methods) and/or insert their own functions. The signal detection algorithm was tested solely on

datasets with fixed sampling rates. Although the minimum requirement for identifying an event is

two consecutive x values with the same sign, it is possible some assumptions we have made in

other aspects of the algorithm may lead to inaccurate identification of events in datasets with

variable sampling rates. Testing and implementing different methods of data interpolation could

alleviate this limitation but has not been done by our research group.

We speculate that FIBSI could theoretically work with almost any type of x,y dataset where

the measured parameter exhibits deviations from baseline, as FIBSI is adapted to account for

changes in the baseline level of activity via the running fit function. This is an improvement over

arbitrary decisions for baseline or rudimentary measures (e.g., simple average) that assume the

system being studied is fixed. Although we have only fully tested FIBSI for processing of voltage

fluctuation time series and fluorescence intensity time series, ongoing projects are currently

adapting FIBSI to process other types of datasets (e.g., *in vivo* nerve recordings and calcium

imaging in rodent and zebrafish neurons), and to investigate physiological states and disease

models pertinent to our research groups. Other examples of biological datasets that may be

processed and analyzed using FIBSI include animal vibrations (e.g., honeybee waggle dance

[67]) and sounds (e.g., ultrasonic vocalizations [68,69]). In summary, there exists broad potential

for FIBSI to facilitate signal analysis of diverse types of raw data by biologists having quite different

backgrounds in quantitative analysis.

4. Methods

4.1. Description of frequency-independent biological signal identification (FIBSI) program

The signal detection algorithm used by FIBSI was generalized from our initial study where we

identified DSFs in nociceptors [17]. It is an extension of the Ramer-Douglas-Peucker algorithm

for identifying significant shapes that ought to be retained in an image while reducing the total

number of points needed to represent said image [15,16]. The FIBSI program was developed and

tested on Ubuntu Linux and Windows 10 operating systems. The data processing is as follows:

•   A vector Y_{trace} in which t = time points and y = independent measured variable in an x, y

series is received as input

554
$$Y_{trace} = (y_{t1}, y_{t2}, \dots y_{tn})_{trace}$$

553   (1)

555   •   The vector can be normalized to an external reference if selected

557
$$Y_{trace(normalized)} = (y_{t1}, y_{t2}, \dots y_{tn})_{trace} / (y_{t1}, y_{t2}, \dots y_{tn})_{reference}$$

556   (2)

558   •   A fitted line is generated where f = function used (e.g., running median, least-squares linear

559   regression)

561
$$Y_{fit} = f[Y_{trace(normalized)}]$$

560   (3)

562   •   The residuals to the fitted line at each given point are calculated

564
$$Y_{\Delta fit} = Y_{trace(normalized)} - Y_{fit}$$

563   (4)

565   •   Signal/event detection is performed by dividing $Y_{\Delta fit}$ into discrete local maxima 'above' and

566   local minima 'below' waveforms in relation to Y_{fit}:

567   **a.** We first define that y values indexed at time points where $y \geq 0$ are elements of

568   'above' time points, and y values at time points where $y < 0$ are elements of 'below'

569   time points:

570
$$y_{ta} \in y_{tn} \; if \; y \geq 0$$

571
$$y_{tb} \in y_{tn} \; if \; y < 0$$

572   **b.** We then define the 'above' and 'below' vectors in $Y_{\Delta fit}$ where m is the length of the

573   'above' vector:

574
$$Y_{\Delta fit(above)} = (y_{ta1}, y_{ta2}, \dots y_{tam})$$

575
$$Y_{\Delta fit(below)} = (y_{tb1}, y_{tb2}, \dots y_{ta(n-m)})$$

576   **c.** We then define the set of criteria for discrete 'above' events, where $y = f(T)$, and $t_i \in$

577   T where t_i is a local maxima if the following conditions are met:

578
$$f(t_i) > 0$$

Start (rise) time: $t_R \in T$ such that $t_R < t_i$ and $f(t_R) = 0$

End (fall) time: $t_F \in T$ such that $t_F > t_i$ and $f(t_F) = 0$

Components of the rising phase: $R = \{t \in T \mid 0 < t_R < t_i\}$

Components of the falling phase: $F = \{t \in T \mid t_i < t_F < t_n\}$

R must not contain numbers that cross zero: $(\forall_t \in R)f(t) > 0$

F must not contain numbers that cross zero: $(\forall_t \in F)f(t) > 0$

An 'above' event, Z, is the union of components in R and F: $Z = R \cup F$

$(\forall_t \in Z)\nexists_t$ where $f(t) > f(t_i) \Rightarrow t_i$ is the time of the local maximum, t_R is the start time, and t_F

is the end time

$\therefore Z$ is the set of numbers contained within a single 'above' event

d. The local minima for a 'below' event is therefore defined as $f(t_i) < 0$ and a similar

set of rules:

591
$$(\forall_t \in Z)\nexists_t \text{ where } f(t) < f(t_i)$$

• Events can be excluded via user-defined duration and amplitude cutoffs, and replaced with

the line of best fit connecting the start and end points of the excluded event; this is a

generalization of the AP exclusion method described previously [17]

•   Refitting can be applied; this allows additional use of a trend line calculated by tracing the

peak-to-peak pathway of identified 'above' or 'below' events; steps (3-4) are repeated to

generate a new $Y_{fit(peak)}$ and $Y_{\Delta fit(peak)}$; troughs can also be used to generate $Y_{fit(trough)}$ and

$Y_{\Delta fit(trough)}$

•   If exclusions and/or refitting methods are applied, then event detection can be performed

again

•   Event directionality (above or below), dimensions (duration, amplitude, AUC), and indices

(start, peak, end) are tabulated; graphs and comma-separated x, y series can be generated

as optional outputs

4.2. Data analysis

Analysis of raw electrophysiological and fluorescence time series data was performed with FIBSI

written using the Anaconda v2019.7.0.0 (Anaconda, Inc, Austin, TX) distribution of Python v3.5.2,

and the NumPy and matplotlib.pyplot libraries. Custom scripts were also written to analyze the

electrophysiological data using the FFT. Statistical analyses of the DSF and gut motility data

generated by FIBSI were performed using Prism v8.2.1 (GraphPad Software, Inc, La Jolla, CA).

Decisions to compare the DSFs in NA neurons between rats and mice, and to compare the DSFs

between the NA and RA neurons in rats were made *a priori*. Amplitude substitutions for

suprathreshold DSFs were calculated manually as described previously [17,18] (see also the

FIBSI tutorial included online). One zebrafish (fed) was omitted from analysis due to a lack of

muscle movements within the gut, but blood flow was still observed. This was confirmed during

video playback and inspection of the raw fluorescence intensity time series. Statistical significance

was set at $P < 0.05$ and all reported P values are two-tailed.

4.3. Program and data availability

The FIBSI source code, experimental data used in this study, and a tutorial for using FIBSI are

available on a GitHub repository titled "FIBSI Program" by user rmcassidy

(https://github.com/rmcassidy/FIBSI_program). This has been done to promote the free use,

modification, and distribution of FIBSI as a useful tool for researchers. Forking the repository is

encouraged and changes will, to the extent possible, be pulled into the original repository as an

update.

4.4. Description of sensory neuron electrophysiological data used for post-hoc analysis

A subset of electrophysiological data obtained from dorsal root ganglia (DRG) sensory neurons

dissociated from 16 male Sprague-Dawley rats and 1 male and 4 female C57BL/6 wild-type mice

were used for post-hoc analysis. These rodents were used as naïve controls in prior studies

performed by our research group, and detailed methods have been published [18,19]. Rodent

experiments were performed in accordance with the International Association for the Study of

Pain, the Guide for the Care and Use of Laboratory Animals, and were approved by the

Institutional Animal Welfare Committee of the University of Texas Health Science Center at

Houston, protocol numbers AWC-18-0035, AWC-15-0122, AWC-18-0134, and AWC-15-0160.

Briefly, the DRG were harvested below thoracic vertebrate levels T9 (mice) and T10 (rats) and

sensory neurons were enzymatically dissociated for overnight incubation at 37 °C with 5% CO_2.

After 18-24 hours, whole-cell current-clamp recordings were performed. To consistently analyze

the DSFs in different neurons and rodent species, only recordings during which neurons were

injected with the current necessary to hold the membrane potential at -45 mV for 30 seconds to

model pain-associated conditions [17,18] were used. Neurons were sampled at 20 kHz with a 10

kHz Bessel filter, and the time and voltage coordinate data were extracted to comma separated

value (.csv) files using PatchMaster v2x90.1 software (HEKA Elektronik, Holliston, MA).

4.5. Zebrafish handling and care

All zebrafish care, breeding, and experiments were performed in accordance with the Guide for

the Care and Use of Laboratory Animals and were approved by the Institutional Animal Welfare

Committee of the University of Texas Health Science Center at Houston, protocol number AWC-

19-0078. Adult *AB* wild-type zebrafish (*Danio rerio*) were housed inside the Center for Laboratory

Animal Medicine and Care specific pathogen-free aquatic facility in recirculating tanks maintained

on a 14 h/10 h light/dark cycle at pH 7.2-7.4 and 26 °C. Zebrafish eggs were obtained following

natural spawning; breeding groups (4:3 female to male ratio) were placed into separate breeding

tanks the night before spawning. Eggs were first rinsed with sterile E3 (prepared in double distilled

water with 10 mM HEPES, 5 mM NaCl, 0.17 mM KCl, 0.40 mM $CaCl_2$, and 0.67 mM $MgSO_4$, pH

7.2) and then raised in petri dishes filled with E3 supplemented with 0.033 mg/mL methylene blue

and 0.013 mg/mL 1-phenyl-2-thiourea to prevent melanin synthesis. The eggs were incubated at

28.5 °C in a diurnal incubator (14 h/10 h light/dark cycle) and E3 was exchanged daily until all

embryos hatched, then E3 was exchanged every other day. For feeding, a subset of larvae were

incubated with the live prey *Paramecium caudatum* for 2 h at 6 days dpf [70]. The remaining

larvae were not fed. In some experiments involving unfed larvae, transgenic *Tg(-*

8.3phox2bb:Kaede) zebrafish larvae [71,72] were used in place of wild-type *AB* larvae. Adult

transgenic zebrafish were housed and spawned in accordance with IACUC-approved protocol

1144724 by the Uribe lab at Rice University. Fertilized embryos were delivered to the Krachler

lab at 0 dpf for rearing. Transgenic embryos were handled the same as wild-type *AB* larvae.

4.6. Using fluorescence microscopy to image gut motility in larval zebrafish

At 7 dpf, zebrafish larvae were transferred to 25 cm^2 (T25) culture flasks (10-20 larvae per flask)

filled with 15 mL of E3 supplemented with 0.01 µg/mL of Nile Red (Sigma Aldrich, St. Louis, MO).

Flasks were incubated at 28.5 °C. Following incubation (2-4 h), larvae were anesthetized with

0.16 mg/mL tricaine (prepared in double distilled water and buffered to pH 7.2 using 0.5 M PIPES

and 1 M NaOH; Sigma Aldrich) and sorted for uptake of Nile Red in the intestinal tract. Larvae

with intestinal tracts clearly outlined with Nile Red dye were embedded in 1% low melting point

agarose (Promega, Madison, WI) in E3 supplemented with 0.16 mg/mL tricaine in 6-well glass

bottom plates (#1.5 high performance cover glass, 0.17±0.005 mm; Cellvis, Mountainview, CA).

The intestinal tracts of the larvae were imaged for 6-10 min at 22 °C using an inverted confocal

laser scanning microscope (Olympus FluoView FV3000; Olympus Corporation of the Americas

Headquarters, Center Valley, PA) using the following settings: 10X objective lens with x1.0 zoom,

1024 x 768 x 1 pixel resolution (1.406 µm/pixel), 1.25-2.5 frames/s sampling rate, 561 nm

excitation laser (Galvano) set to 0.5% transmission and 650 V, and a 570-670 nm emission

detection window. Post-acquisition, a total of 19 analysis ROIs (fixed position) were drawn over

the length of the intestinal tract to measure changes in fluorescence during gut motility, and 1

reference ROI was drawn outside the intestinal tract to correct for background fluorescence (see

Results for ROI positioning). Care was taken to use the same ROI dimensions and to position the

ROIs consistently between larvae (ROI center-to-center distance = ~55 µm). The Nile Red dye

was used as a contrast agent to outline the intestinal tract [30]. Changes in fluorescence during

gut motility were due to muscle contractions widening, closing, and moving the intestinal tract.

Total fluorescence intensity and time coordinate data per ROI were extracted to comma separated

value (.csv) files using Olympus FV31S-SW v2.3.2.169 software. All larvae were euthanized at

the end of experiments at 7-8 dpf using 1.6 mg/mL buffered tricaine.

Acknowledgements

The authors would like to thank members of the Uribe, Krachler, and Walters labs for technical

assistance.

Funding and Conflicts of Interest

Development and testing of FIBSI: R.M. Cassidy was supported by a UTHealth Center for Clinical

and Translational Sciences (4TL1TR000369) and NIH-NIDA NRSA F30 (F30DA047030); M.A.

Odem was supported by a Zilkha Family Fellowship. Zebrafish experiments were supported by a

John S. Dunn Collaborative Research Award to R.A. Uribe and A.M. Krachler, and NIH grant R01

AI132354 to A.M. Krachler. R.A. Uribe is a CPRIT Scholar in Cancer Research (RR170062),

which provided startup funds. Rodent experiments were supported by NIH-NINDS R01

(NS091759) to C.W. Dessauer and E.T. Walters, a subcontract to E.T. Walters from Manzanita

Pharmaceuticals Inc. and US Army Medical Research Grant W81XWH-12-1-0504 to E.T. Walters,

and two grants from Mission Connect – a TIRR Foundation Program – #017-107 to A.G.

Bavencoffe and #016-113 C.W. Dessauer and J. Herrera. The authors report no conflicts of

interest.

References

1.   Kumar GG, Sahoo SK, Meher PK. 50 Years of FFT Algorithms and Applications. Circuits, Syst Signal
      Process. 2019;38: 5665–5698. doi:10.1007/s00034-019-01136-8

2.   Joy BR, Amara A, Nakhmani A. Transform with no Parameters Based on Extrema Points for Non-
      stationary Signal Analysis. Circuits, Syst Signal Process. 2018;37: 2535–2547. doi:10.1007/s00034-
      017-0676-5

3.   Tsakanikas P, Sigalas C, Rigas P, Skaliora I. High-Throughput Analysis of in-vitro LFP
      Electrophysiological Signals: A validated workflow/software package. Sci Rep. 2017;7: 1–11.
      doi:10.1038/s41598-017-03269-9

4.   Fayyaz Z, Bahadorian M, Doostmohammadi J, Davoodnia V, Khodadadian S, Lashgari R.
      Multifractal detrended fluctuation analysis of continuous neural time series in primate visual
      cortex. J Neurosci Methods. 2019;312: 84–92. doi:10.1016/j.jneumeth.2018.10.039

5.   Mei Z, Zhao X, Chen H, Chen W. Bio-signal complexity analysis in epileptic seizure monitoring: A
      topic review. Sensors (Switzerland). 2018;18: 1–27. doi:10.3390/s18061720

6.   Nayak SK, Bit A, Dey A, Mohapatra B, Pal K. Review Article A Review on the Nonlinear Dynamical
      System Analysis of Electrocardiogram Signal. 2018;2018. doi:10.1155/2018/6920420

7.   Baraban M, Koudelka S, Lyons DA. Ca 2+ activity signatures of myelin sheath formation and
      growth in vivo. Nat Neurosci. 2018;21: 19–25. doi:10.1038/s41593-017-0040-x

8.   Patel JM, Swanson J, Ung K, Herman A, Hanson E, Ortiz-Guzman J, et al. Sensory perception
      drives food avoidance through excitatory basal forebrain circuits. Elife. 2019;8: 1–28.
      doi:10.7554/eLife.44548

9.   Lovett-Barron M, Andalman AS, Allen WE, Vesuna S, Kauvar I, Burns VM, et al. Ancestral Circuits
      for the Coordinated Modulation of Brain State. Cell. 2017;171: 1411-1423.e17.
      doi:10.1016/j.cell.2017.10.021

730   10.   Andalman AS, Burns VM, Lovett-Barron M, Broxton M, Poole B, Yang SJ, et al. Neuronal Dynamics

731   Regulating Brain and Behavioral State Transitions. Cell. 2019;177: 970-985.e20.

732   doi:10.1016/j.cell.2019.02.037

733   11.   Mu Y, Bennett D V., Rubinov M, Narayan S, Yang CT, Tanimoto M, et al. Glia Accumulate Evidence

734   that Actions Are Futile and Suppress Unsuccessful Behavior. Cell. 2019;178: 27-43.e19.

735   doi:10.1016/j.cell.2019.05.050

736   12.   Migault G, van der Plas TL, Trentesaux H, Panier T, Candelier R, Proville R, et al. Whole-Brain

737   Calcium Imaging during Physiological Vestibular Stimulation in Larval Zebrafish. Curr Biol.

738   2018;28: 3723-3735.e6. doi:10.1016/j.cub.2018.10.017

739   13.   Chen X, Mu Y, Hu Y, Kuan AT, Nikitchenko M, Randlett O, et al. Brain-wide Organization of

740   Neuronal Activity and Convergent Sensorimotor Transformations in Larval Zebrafish. Neuron.

741   2018;100: 876-890.e5. doi:10.1016/j.neuron.2018.09.042

742   14.   Vladimirov N, Wang C, Höckendorf B, Pujala A, Tanimoto M, Mu Y, et al. Brain-wide circuit

743   interrogation at the cellular level guided by online analysis of neuronal function. Nat Methods.

744   2018;15: 1117–1125. doi:10.1038/s41592-018-0221-x

745   15.   Douglas DH, Peucker TK. Algorithms for the reduction of the number of points required to

746   represent a digitized line or its caricature. Cartogr Int J Geogr Inf Geovisualization. 1973;10: 112–

747   122. doi:https://doi.org/10.3138/FM57-6770-U75U-7727

748   16.   Ramer U. An iterative procedure for the polygonal approximation of plane curves. Comput Graph

749   Image Process. 1972;1: 244–256. doi:10.1016/S0146-664X(72)80017-0

750   17.   Odem MA, Bavencoffe AG, Cassidy RM, Lopez ER, Tian J, Dessauer CW, et al. Isolated nociceptors

751   reveal multiple specializations for generating irregular ongoing activity associated with ongoing

752   pain. Pain. 2018;159: 2347–2362. doi:10.1097/j.pain.0000000000001341

18.   Berkey SC, Herrera JJ, Odem MA, Rahman S, Cheruvu SS, Cheng X, et al. EPAC1 and EPAC2

promote nociceptor hyperactivity associated with chronic pain after spinal cord injury. Neurobiol

Pain. 2020;7. doi:10.1016/j.ynpai.2019.100040

19.   Odem MA, Lacagnina MJ, Katzen SL, Li J, Spence EA, Grace PM, et al. Sham surgeries for central

and peripheral neural injuries persistently enhance pain-avoidance behavior as revealed by an

operant conflict test. Pain. 2019;160: 2440–2455. doi:10.1097/j.pain.0000000000001642

20.   Amir R, Michaelis M, Devor M. Membrane potential oscillations in dorsal root ganglion neurons:

role in normal electrogenesis and neuropathic pain. J Neurosci. 1999;19: 8589–8596.

doi:https://doi.org/10.1523/jneurosci.19-19-08589.1999

21.   Amir R, Michaelis M, Devor M. Burst discharge in primary sensory neurons: triggered by

subthreshold oscillations, maintained by depolarizing afterpotentials. J Neurosci. 2002;22: 1187–

1198. doi:https://doi.org/10.1523/jneurosci.22-03-01187.2002

22.   Liu C-N, Michaelis M, Amir R, Devor M. Spinal nerve injury enhances subthreshold membrane

potential oscillations in DRG neurons: Relation to neuropathic pain. J Neurophysiol. 2000;84:

205–215. doi:10.1152/jn.2000.84.1.205

23.   Mathers DA, Barker JL. Spontaneous voltage and current fluctuations in tissue cultured mouse

dorsal root ganglion cells. Brain Res. 1984;293: 35–47. doi:https://doi.org/10.1016/0006-

8993(84)91450-1

24.   Wang YY, Wen ZH, Duan JH, Zhu JL, Wang WT, Dong H, et al. Noise enhances subthreshold

oscillations in injured primary sensory neurons. NeuroSignals. 2011;19: 54–62.

doi:10.1159/000324519

25.   Xing JL, Hu SJ, Long KP. Subthreshold membrane potential oscillations of type A neurons in

injured DRG. Brain Res. 2001;901: 128–136. doi:10.1016/S0006-8993(01)02329-0

26.   Xing JL, Hu SJ, Jian Z, Duan JH. Subthreshold membrane potential oscillation mediates the

excitatory effect of norepinephrine in chronically compressed dorsal root ganglion neurons in the

rat. Pain. 2003;105: 177–183. doi:10.1016/S0304-3959(03)00200-8

27.   Yang RH, Wang WT, Chen JY, Xie RG, Hu SJ. Gabapentin selectively reduces persistent sodium

current in injured type-A dorsal root ganglion neurons. Pain. 2009;143: 48–55.

doi:10.1016/j.pain.2009.01.020

28.   Cocchiaro JL, Rawls JF. Microgavage of zebrafish larvae. J Vis Exp. 2013; 1–12. doi:10.3791/4434

29.   Shi Y, Zhang Y, Zhao F, Ruan H, Huang H, Luo L, et al. Acetylcholine serves as a derepressor in

Loperamide-induced Opioid-Induced Bowel Dysfunction (OIBD) in zebrafish. Sci Rep. 2014;4: 1–

12. doi:10.1038/srep05602

30.   Cloney K, Steele SL, Stoyek MR, Croll RP, Smith FM, Prykhozhij S V., et al. Etiology and functional

validation of gastrointestinal motility dysfunction in a zebrafish model of CHARGE syndrome.

FEBS J. 2018;285: 2125–2140. doi:10.1111/febs.14473

31.   Holmberg A, Olsson C, Hennig GW. TTX-sensitive and TTX-insensitive control of spontaneous gut

motility in the developing zebrafish (Danio rerio) larvae. J Exp Biol. 2007;210: 1084–1091.

doi:10.1242/jeb.000935

32.   Rich A, Gordon S, Brown C, Gibbons SJ, Schaefer K, Hennig G, et al. Kit signaling is required for

development of coordinated motility patterns in zebrafish gastrointestinal tract. Zebrafish.

2013;10: 154–160. doi:10.1089/zeb.2012.0766

33.   James DM, Kozol RA, Kajiwara Y, Wahl AL, Storrs EC, Buxbaum JD, et al. Intestinal dysmotility in a

zebrafish (Danio rerio) shank3a;shank3b mutant model of autism. Mol Autism. 2019;10: 1–15.

doi:10.1186/s13229-018-0250-4

34.   Heanue TA, Boesmans W, Bell DM, Kawakami K, Vanden Berghe P, Pachnis V. A Novel Zebrafish

ret Heterozygous Model of Hirschsprung Disease Identifies a Functional Role for mapk10 as a

Modifier of Enteric Nervous System Phenotype Severity. PLoS Genet. 2016;12: 1–23.

doi:10.1371/journal.pgen.1006439

35.   Ganz J, Baker RP, Hamilton MK, Melancon E, Diba P, Eisen JS, et al. Image velocimetry and

spectral analysis enable quantitative characterization of larval zebrafish gut motility.

Neurogastroenterol Motil. 2018;30: 1–12. doi:10.1111/nmo.13351

36.   Hennig GW, Costa M, Chen BN, Brookes SJH. Quantitative analysis of peristalsis in the guinea-pig

small intestine using spatio-temporal maps. J Physiol. 1999;517: 575–590. doi:10.1111/j.1469-

7793.1999.0575t.x

37.   Roberts RR, Ellis M, Gwynne RM, Bergner AJ, Lewis MD, Beckett EA, et al. The first intestinal

motility patterns in fetal mice are not mediated by neurons or interstitial cells of Cajal. J Physiol.

2010;588: 1153–1169. doi:10.1113/jphysiol.2009.185421

38.   Roach G, Wallace RH, Cameron A, Ozel RE, Hongay CF, Baral R, et al. Loss of ascl1a prevents

secretory cell differentiation within the zebrafish intestinal epithelium resulting in a loss of distal

intestinal motility. Dev Biol. 2013;376: 171–186. doi:10.1038/jid.2014.371

39.   Kendig DM, Hurst NR, Grider JR. Spatiotemporal mapping of motility in Ex Vivo preparations of

the intestines. J Vis Exp. 2016;2016: 1–11. doi:10.3791/53263

40.   Hibberd TJ, Feng J, Luo J, Yang P, Samineni VK, Gereau RW, et al. Optogenetic Induction of

Colonic Motility in Mice. Gastroenterology. 2018;155: 514-528.e6.

doi:10.1053/j.gastro.2018.05.029

41.   Bedi SS, Yang Q, Crook RJ, Du J, Wu Z, Fishman HM, et al. Chronic spontaneous activity generated

in the somata of primary nociceptors is associated with pain-related behavior after spinal cord

injury. J Neurosci. 2010;30: 14870–14882. doi:10.1523/JNEUROSCI.2428-10.2010

42.   Yang Q, Wu Z, Hadden JK, Odem MA, Zuo Y, Crook RJ, et al. Persistent pain after spinal cord injury

is maintained by primary afferent activity. 2014;34: 10765–10769. doi:10.1523/JNEUROSCI.5316-

13.2014

43.   Bavencoffe A, Li Y, Wu Z, Yang Q, Herrera J, Kennedy EJ, et al. Persistent electrical activity in

primary nociceptors after spinal cord injury is maintained by scaffolded adenylyl cyclase and

protein kinase A and is associated with altered adenylyl cyclase regulation. J Neurosci. 2016;36:

1660–1668. doi:10.1523/JNEUROSCI.0895-15.2016

44. Guo Z, Qiu CS, Jiang X, Zhang J, Li F, Liu Q, et al. TRESK K+ channel activity regulates trigeminal

nociception and headache. eNeuro. 2019;6. doi:10.1523/ENEURO.0236-19.2019

45. North RY, Li Y, Ray P, Rhines LD, Tatsui CE, Rao G, et al. Electrophysiological and transcriptomic

correlates of neuropathic pain in human dorsal root ganglion neurons. Brain. 2019;142: 1215–

1226. doi:10.1093/brain/awz063

46. Gotman J, Gloor P. Automatic recognition and quantification of interictal epileptic activity in the

human scalp EEG. Electroencephalogr Clin Neurophysiol. 1976;41: 513–529. doi:10.1016/0013-

4694(76)90063-8

47. Greaney JL, Kenney WL. Measuring and quantifying skin sympathetic nervous system activity in

humans. J Neurophysiol. 2017;118: 2181–2193. doi:10.1152/jn.00283.2017

48. Baker RP, Taormina MJ, Jemielita M, Parthasarathy R. A combined light sheet fluorescence and

differential interference contrast microscope for live imaging of multicellular specimens. J

Microsc. 2015;258: 105–112. doi:10.1111/jmi.12220

49. Brijs J, Hennig GW, Axelsson M, Olsson C. Effects of feeding on in vivo motility patterns in the

proximal intestine of shorthorn sculpin (Myoxocephalus scorpius). J Exp Biol. 2014;217: 3015–

3027. doi:10.1242/jeb.101741

50. Bates JM, Mittge E, Kuhlman J, Baden KN, Cheesman SE, Guillemin K. Distinct signals from the

microbiota promote different aspects of zebrafish gut differentiation. Dev Biol. 2006;297: 374–

386. doi:10.1016/j.ydbio.2006.05.006

51. Wang Z, Du J, Lam SH, Mathavan S, Matsudaira P, Gong Z. Morphological and molecular evidence

for functional organization along the rostrocaudal axis of the adult zebrafish intestine. BMC

Genomics. 2010;11. doi:10.1186/1471-2164-11-392

52. Rao M. An increasingly complex view of intestinal motility. Nat Rev Gastroenterol Hepatol. 2019;

epub ahead of print. doi:10.1038/s41575-019-0249-0

53. Ganz J. Gut feelings: Studying enteric nervous system development, function, and disease in the

zebrafish model system. Dev Dyn. 2018;247: 268–278. doi:10.1002/dvdy.24597

54. Brosens E, Burns AJ, Brooks AS, Matera I, Borrego S, Ceccherini I, et al. Genetics of enteric

neuropathies. Dev Biol. 2016;417: 198–208. doi:10.1016/j.ydbio.2016.07.008

55. Rao M, Gershon MD. The bowel and beyond: the enteric nervous system in neurological

disorders. Nat Rev Gastroenterol Hepatol. 2016;13: 517–528. doi:10.1038/nrgastro.2016.107

56. Rao M, Gershon MD. Enteric nervous system development: what could possibly go wrong? Nat

Rev Neurosci. 2018;19: 552–565. doi:10.1038/s41583-018-0041-0

57. Burzynski G, Shepherd IT, Enomoto H. Genetic model system studies of the development of the

enteric nervous system, gut motility and Hirschsprung's disease. Neurogastroenterol Motil.

2009;21: 113–127. doi:10.1111/j.1365-2982.2008.01256.x

58. Burns AJ, Roberts RR, Bornstein JC, Young HM. Development of the enteric nervous system and

its role in intestinal motility during fetal and early postnatal stages. Semin Pediatr Surg. 2009;18:

196–205. doi:10.1053/j.sempedsurg.2009.07.001

59. Moloney RD, Johnson AC, O'Mahony SM, Dinan TG, Greenwood-Van Meerveld B, Cryan JF. Stress

and the Microbiota-Gut-Brain Axis in Visceral PaRelevance to Irritable Bowel Syndrome. CNS

Neurosci Ther. 2016;22: 102–117. doi:10.1111/cns.12490

60. Mahony SMO, Dinan TG, Cryan JF. The microbiota gut brain axis as a key regulator of visceral

pain. Pain. 2017;158: 19–28. doi:10.1177/2398212817705279

61. Escalante J, McQuade RM, Stojanovska V, Nurgali K. Impact of chemotherapy on gastrointestinal

functions and the enteric nervous system. Maturitas. 2017;105: 23–29.

874   doi:10.1016/j.maturitas.2017.04.021

875 62. Flores EM, Nguyen AT, Odem MA, Eisenhoffer GT, Krachler AM. The zebrafish as a model for

876   gastrointestinal tract–microbe interactions. Cell Microbiol. 2020;22. doi:10.1111/cmi.13152

877 63. Bao W, Volgin AD, Alpyshov ET, Friend AJ, Strekalova T V., de Abreu MS, et al. Opioid

878   Neurobiology, Neurogenetics and Neuropharmacology in Zebrafish. Neuroscience. 2019;404:

879   218–232. doi:10.1016/j.neuroscience.2019.01.045

880 64. Douglas AE. Simple animal models for microbiome research. Nat Rev Microbiol. 2019;17: 764–

881   775. doi:10.1038/s41579-019-0242-1

882 65. de Abreu MS, Giacomini ACVV, Sysoev M, Demin KA, Alekseeva PA, Spagnoli ST, et al. Modeling

883   gut-brain interactions in zebrafish. Brain Res Bull. 2019;148: 55–62.

884   doi:10.1016/j.brainresbull.2019.03.003

885 66. Sekerli M, Del Negro C a., Lee RH, Butera RJ. Estimating action potential thresholds from

886   neuronal time-series: New metrics and evaluation of methodologies. IEEE Trans Biomed Eng.

887   2004;51: 1665–1672. doi:10.1109/TBME.2004.827531

888 67. Nieh JC, Tautz J. Behavior-locked signal analysis reveals weak 200-300 Hz comb vibrations during

889   the honeybee waggle dance. J Exp Biol. 2000;203: 1573–1579.

890 68. Han JS, Bird GC, Li W, Jones J, Neugebauer V. Computerized analysis of audible and ultrasonic

891   vocalizations of rats as a standardized measure of pain-related behavior. 2005;141: 261–269.

892   doi:10.1016/j.jneumeth.2004.07.005

893 69. Reno JM, Marker B, Cormack LK, Schallert T, Duvauchelle CL. Automating ultrasonic vocalization

894   analyses: The WAAVES program. J Neurosci Methods. 2013;219: 155–161.

895   doi:10.1016/j.jneumeth.2013.06.006

896 70. Flores E, Thompson L, Sirisaengtaksin N, Nguyen AT, Ballard A, Krachler AM. Using the protozoan

897   Paramecium caudatum as a vehicle for food-borne infections in zebrafish larvae. J Vis Exp.

2019;2019: 1–7. doi:10.3791/58949

71.   Harrison C, Wabbersen T, Shepherd IT. In vivo visualization of the development of the enteric

nervous system using a Tg(-8.3bphox2b: Kaede) transgenic zebrafish. Genesis. 2014;52: 985–990.

doi:10.1002/dvg.22826

72.   Uribe RA, Hong SS, Bronner ME. Retinoic acid temporally orchestrates colonization of the gut by

vagal neural crest cells. Dev Biol. 2018;433: 17–32. doi:10.1016/j.ydbio.2017.10.021

Supporting Information Captions

S1 table. Parameters for DSF exponential growth models.

* Denotes probability the model is correct in comparison to the exponential plateau. AICc, Akaike's

criterion; df, degrees of freedom; k, rate constant; YM, max y value; Y0, starting y value.

S2 movies. Fluorescence contrasted contractions during gut motility in unfed and fed

zebrafish larvae.

(A) Unfed larva corresponding to Fig 3B-C; 1 frame = 0.653 s. (B-C) Unfed larvae; 1 frame =

0.659 s. (D-E) Fed larvae; 1 frame = 0.592 s. Playback set to 30 frames/s for all movies.

S3 figure. Events detected in quadrants 1-3 for unfed and fed zebrafish larvae do not

exhibit peristalsis rhythmicity.

(A-C) Raster plots of Q1-3 corresponding to the unfed larva depicted in Fig 4J. (D) Time course

of fluorescence intensity changes measured from 20 ROIs corresponding to the fed larva depicted

in Fig 4K (traces color coded for visual differentiation). (E) Data corresponding to (D) post-

processing with FIBSI. The same cutoffs (dotted red lines, duration ≥10 s and amplitude ≥0.05)

were applied to distinguish between background noise (Q1-3) and low-frequency contractions.

Events are color coded according to their ROI of origin in (D). (F-H) Raster plots for Q1-3

corresponding to the data in (E).

Figure 1

Figure 2

A Intestinal tract Cloaca · *AB*, 8 dpf

B Foregut · Midgut · Hindgut · 100 μm

C 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 · Analysis ROIs · Reference ROI 20 · Red = Nile Red staining in intestinal tract

D Foregut · Midgut/hindgut · Hindgut/cloaca · Lumen · ROI · Time

E Foregut · Baseline · Contraction · Time · Midgut/hindgut · Contraction · Baseline · Time · Hindgut/cloaca · Time · Predicted fluorescence intensity

F Raw fluorescence time series per ROI · Total intensity (AU) · ROI #1-7 · ROI #8-19 · ROI #20 · Time (s)

Figure 3

Figure 4