Title: The effect of resource limitation on the temperature-dependence of mosquito
population fitness

Paul J. Huxley[1], Kris A. Murray[1, 2], Samraat Pawar[3], Lauren J. Cator[3]
[1]MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease
Epidemiology, School of Public Health, Imperial College London, London, UK.
[2]MRC Unit The Gambia at London School of Hygiene & Tropical Medicine, Banjul, The
Gambia.
[3]Department of Life Sciences, Imperial College London, Ascot, UK.

ABSTRACT

Laboratory-derived temperature dependencies of life history traits are increasingly being used
to make mechanistic predictions for how climatic warming will affect vector-borne disease
dynamics, partially by affecting abundance dynamics of the vector population. These
temperature-trait relationships are typically estimated from populations reared on optimal
resource supply, even though natural populations of vectors are expected to experience
variation in resource supply, including intermittent resource limitation. Using laboratory
experiments on the mosquito *Aedes aegypti*, a principal arbovirus vector, combined with
stage-structured population modelling, we show that low-resource supply significantly
depresses the vector's maximal population growth rate across the entire temperature range
(22–32°C) and causes it to peak at a lower temperature than at high-resource supply. This
effect is primarily driven by an increase in juvenile mortality and development time,
combined with an exaggerated decrease in adult size with temperature at low-resource
supply. Our study suggests that projections of vector abundance and disease transmission
based on laboratory studies are likely to substantially underestimate how resource supply can
modulate the temperature-dependency of population-level fitness through its influence on
juvenile survival and development time. Our results provide compelling evidence for future
studies to consider resource supply when predicting the effects of climate and habitat change
on disease vectors and transmission.

Key words: climatic warming, abundance, temperature, population fitness, resource
limitation, vector-borne disease

INTRODUCTION

The global burden of human, animal, and plant vector-borne diseases has increased substantially in recent decades [1,2]. The transmission patterns of these diseases are strongly linked to spatio-temporal distribution and abundance of their vectors [3,4]. Therefore, there is growing concern that climate and land-use change coupled with rapid globalization may shift the distributions and abundances of vector species and thus, the diseases they transmit [5,6]. However, we currently lack a mechanistic understanding of how changes in multiple abiotic environmental drivers affect the abundance of disease vectors [7–9].

Because most disease vectors are small ectotherms, interest is growing rapidly in the effects of environmental temperature in particular on their population fitness [5,6,10–15]. Biological rates ("functional traits", such as metabolic, development, and mortality rate [7]) of ectotherms increase approximately exponentially with temperature up to some optimum before declining to zero [16]. Therefore, a number of recent studies have used laboratory data on the thermal responses of functional traits of vectors to predict how temperature will affect vector abundance and disease transmission in the field, leading to interesting new insights [8]. However, laboratory-derived temperature-trait relationships are generally measured in populations reared on optimal resource supply. Yet, natural populations of vectors are expected to experience variation in resource supply, including intermittent resource limitation.

Indeed, along with temperature, resource supply is another ubiquitous environmental driver that is expected to limit the fitness of vector populations in nature [17–19]. Moreover, temperature and resource supply are expected to act interactively [9,20]. The primary reason for this, is that while the energy cost of somatic maintenance, growth and ontogenetic development of individuals generally increases with temperature [21,22], the ability to meet this increasing demand depends on resource supply. If the resources available to an individual do not keep pace with increasing energy requirements, its growth, development, and survival would be compromised. Ultimately, this should negatively affect fitness, with the severity of these effects increasing with temperature. While the importance of resource supply in mediating the effect of temperature on population abundance may seem obvious, this problem remains largely unresolved theoretically and empirically, not just in vector-borne disease research, but in thermal ecology in general. For example, Ecological Metabolic Theories (including the Metabolic Theory of Ecology (MTE) and Dynamic Energy Budget (DEB) theory), which seek to link organismal metabolic rates to ontogenetic and population growth, generally assume that resource supply is not a limiting factor [21–24].

Additionally, resource supply can also interact with temperature to affect population fitness of ectotherms by determining size at maturity. Generally, size at maturity decreases with rising temperature (the size-temperature rule [25], which also applies to disease vectors, such as mosquitoes [26]). However, the size-temperature rule too remains largely untested under resource limitation in disease vectors and other ectotherms [25,27]. For vectors specifically, female size is demographically and epidemiologically important because it is associated with longevity, fecundity, and biting behaviour [28,29].

In general, the question of whether and how temperature and resource supply may together modulate disease transmission through underlying traits remains open. Here we seek to fill this gap in knowledge by investigating the effect of realistic variation in resource supply on the temperature-dependence of population-level fitness in *Aedes aegypti*, a principal

mosquito vector of human arboviruses (e.g. dengue, yellow fever and zika; [38]). Because
resource competition between larvae is expected to be a major regulator of adult mosquito
abundance, many studies have examined how resource supply and larval density interact to
affect fitness [30–33], while others have investigated the effect of resource supply and larval
density separately [34,35]. However, none of these studies have considered environmental
temperature. On the other hand, studies that have considered temperature have not examined
how the effects of temperature and resource supply together affect fitness through traits
[36,37]. By taking a trait-based approach, we seek to gain general, mechanistic insights into
how resource availability and temperature may together affect the abundance of disease
vectors in the field.

METHODS

To investigate the effects of temperature and resource supply on mosquito life history, we
employed a 3×2 factorial design comprised of three temperatures (22, 26, and 32°C) and two
resource supply levels: 0.1 (low-resource supply) and 1 mg larva^{-1} day^{-1} (high-resource
supply). We conducted a preliminary assay to determine the adequacy of this replication level
to detect statistically significant effect sizes (electronic supplementary material, Tables S4,
S5). These experimental temperatures span the range of average annual temperatures [39]
that this strain of *Ae. aegypti* is likely to experience in the wild (F16-19 originating from Fort
Meyer, FL; [40]). This low-resource supply level was chosen because previous work has
found that it is the highest resource limitation that can be applied to this species without
resulting in complete juvenile mortality [17,18]; a level of limitation that might be expected
in wild populations. We also determined that the low-resource level was appropriate with a
preliminary assay (electronic supplementary material, Tables S4, S5). The high-resource
supply level corresponds to the upper mid-range of the high-resource supply levels used in
Arrivillaga and Barrera [17] and Barrera et al. [18], and is consistent with the levels of
resource supply commonly used in laboratory studies on this species [36,41].

Batches of approximately 300 *Ae. aegypti* eggs were randomly assigned to one of the three
experimental temperatures and immersed in plastic tubs containing 150 ml of tap water. Each
tub was provided with a pinch of powdered fish food (Cichlid Gold®, Hikari, Kyrin Food
Industries Ltd., Japan) to stimulate overnight hatching. The tubs were then submerged in
water baths (Grant Instruments: JAB Academy) set at either 22, 26, or 32°C. Water baths
were situated in a 20°C climate-controlled insectary with a 12L:12D photoperiod and 30
minutes of gradual transition of light levels to simulate sunrise and sunset. On the following
day, first instar larvae were separated into cohorts of 30 and held in tubs containing 150 ml of
water. We created three replicate tubs per treatment (90 individuals/treatment). Low-resource
supply treatments were provided 3 mg of food and high-resource supply treatments received
30 mg. Thereafter, resource levels were adjusted daily according to the number of living
individuals in each tub prior to feeding each day such that resource levels were maintained at
an approximately constant level during the juvenile lifespan. Rearing tubs were cleaned and
refilled with fresh tap water daily. Water volumes were also adjusted daily in accordance
with mortality to maintain larval density (0.2 larvae $\times$ ml^{-1}). Figure S1 is a schematic of the
experimental design and the traits measured.

Fitness calculation

To calculate population-level fitness, we used our data to parameterise stage-structured
matrix projection models [42], which describe change in a population over time:

$$\mathbf{N}_{t+1} = \mathbf{M}\mathbf{N}_t, \qquad\qquad (1)$$

120
where $\mathbf{N}_t$ is a vector of abundances in the stage classes at time t and $\mathbf{M}$ is the population
projection matrix. The first row of $\mathbf{M}$ is populated by daily fecundity (the number of female
offspring produced per female at age i). The sub-diagonal of $\mathbf{M}$ is populated with the
probabilities of survival from age i to age $i+1$. Multiplying the transition matrix ($\mathbf{M}$; eqn. 1)
and stage-structured population size vector ($\mathbf{N}_t$; eqn. 1) sequentially across time intervals
yields the stage-structured population dynamics. Once the stable stage distribution of the
abundance vector is reached, the dominant eigenvalue of the system is the finite population
rate of increase (λ) [42]. Then, the intrinsic rate of population growth is

$$r_{\max} = \log(\lambda).$$

This is a population's inherent capacity to reproduce, and therefore a measure of population-
level fitness [23,43,44]. Negative $r_{\max}$ values indicate decline and positive ones, growth. The
projection matrices were built and analysed using the `popbio` R package [45,46].

Parameterisation

Immature development time and immature and adult survival proportions
Matrix survival elements (the sub-diagonal of the matrix $\mathbf{M}$; eqn. 1) were populated with
continuous survival proportions estimated using the Kaplan-Meier survival function in the
`survival` R package [45,47]. We assumed life stage duration (i.e. larva-to-pupa-to-adult)
was the mean duration of transitioning into and out of that stage, and a fixed age of adult
emergence at the mean age of emergence. Adult survival elements were populated with the
Kaplan-Meier proportions. Hatching-to-adult development times were calculated by
recording the day and time that egg eclosion, pupation and adult emergence occurred for each
individual. Upon pupation, mosquitoes were held in individual falcon tubes containing 5 ml
of tap water. This enabled pupa-to-adult development durations and the lifespans of
individual starved adults to be recorded. Starvation forces adults to metabolise the nutritional
reserves accumulated during larval development, so starved lifespan should increase with
body size. Therefore, starved adult lifespan is a useful indicator of the carry over effects of
temperature and resource availability in the larval habitat [48,49].

Daily fecundity rate
The use of scaling relationships between fecundity and size is common in predictions of
population growth in *Aedes* mosquitoes [50,51]. A detailed description of our method for
estimating fecundity is provided in the electronic supplementary material (Fig. S2). Briefly,
we measured wing length as a proxy for body size, and estimated lifetime fecundity using
previously published datasets on the temperature- and resource supply-dependent scaling
between wing length and lifetime fecundity [41,49]. Daily fecundity rate is required for the
first row of $\mathbf{M}$ (eqn. 1), so lifetime fecundity was divided by lifespan and multiplied by 0.5
(assuming a 1:1 male-to-female offspring ratio) to give temperature-specific individual daily
fecundity.

Parameter sensitivity
We used the delta method to approximate 95% confidence intervals (CIs) for our fitness
calculations [42,52] to account for how uncertainty in survival and fecundity estimates is
propagated through to the $r_{\max}$ estimate. This method requires the standard errors of the
survival and fecundity element estimates. For survival, we used the standard errors estimated
by the Kaplan-Meier survival function in the `survival` R package. For fecundity, we
calculated the standard errors of the mean daily fecundity rates (electronic supplementary

material, Table S2) for each treatment using the `Rmisc` R package [53]. As an additional
sensitivity analysis, we recalculated fitness using the upper and lower 95% CIs of the
exponents for the scaling of wing length and lifetime fecundity (Fig. 3, electronic
supplementary material, Fig. S2).

Elasticity analysis

Elasticities were used to quantify the proportional contributions of individual life history
traits to r_{max}. Elasticity, e_{ij}, measures the proportional effect on λ of an infinitesimal change in
an element of **M** (eqn. 1) with all other elements held constant (the partial derivative) [54,55].
This partial derivative of λ, with respect to each element of **M**, is $s_{ij} = \partial\lambda/\partial a_{ij} = v_i w_j$ with the
dot product $\langle \mathbf{w}, \mathbf{v} \rangle = 1$. Here, **w** is the dominant right eigenvector (the stage distribution
vector of **M**), **v** is the dominant left eigenvector (the reproductive value vector of **M**), and a_{ij}
is the $i{\times}j^{th}$ element of **M**. Elasticities can then be calculated using the relationship: $e_{ij} = a_{ij}/\lambda \times$
s_{ij}. Multiplying an elasticity by λ gives the absolute contribution of its corresponding a_{ij} to λ
[54,55]. Absolute contributions for juvenile and adult elements were summed and changed
proportionally to quantify the sensitivity of r_{max} to these traits.

Statistical analyses

All statistical analyses were conducted using R [45]. The trait data (development time,
lifespan and wing length) were nonlinear, positive and right skewed, so we used full factorial
generalized linear models (GLM) with gamma distributions and identity link functions
(predictor effects were considered additive) to determine the significance of each predictor on
the thermal response of each of these traits. Replicate was included in these GLMs as a fixed
effect.

We investigated the effect of temperature and resource supply on juvenile mortality rate by
fitting a set of candidate distributions (exponential, log-logistic, Gompertz and Weibull) to
the survival data with R package `flexsurv` [56]. The Gompertz survival function was the
best fit to these data according to the Akaike Information Criterion (AIC) (electronic
supplementary material, Table S3). The final mortality model was obtained by dropping
terms from the full model (consisting of temperature $\times$ resource supply + replicate as fixed
effect predictors). If removing a term did not improve model fit (ΔAIC > -2), it was removed
from the full model (electronic supplementary material, Table S3). For each treatment,
maximum likelihood methods executed in `flexsurv` estimated mortality parameters (and
their 95% CIs) of the Gompertz model, $\mu_x = ae^{bx}$, where a is the baseline morality rate, and b
is the change in mortality rate with time. These parameter estimates were then used to
determine the significance of the effects of temperature and resource supply on juvenile
mortality.

RESULTS

All trait responses varied significantly with temperature and resource supply, with a
significant interaction between the two environmental variables (Figs. 1, 2; electronic
supplementary material, Tables S1, S2). Thus, the realised effect of temperature on trait
responses was consistently and significantly mediated by resource supply.

At low-resource supply, daily juvenile mortality rates, μ_x, increased with time at all
temperatures, whereas, at high-resource-supply, they decreased with time (Fig. 1e). Baseline
juvenile mortality rates, a, were significantly lower at low-resource supply than at high-
resource supply as temperatures increased from 22 to 32°C (non-overlapping 95% CIs at 26

and 32°C; Fig 1c). Mortality rate trajectories, b, were significantly lower at high-resource
supply than at low-resource supply as temperatures increased from 22 to 32°C (non-
overlapping 95% CIs at 26 and 32°C; Fig 1d).

Development time varied significantly with the interaction between temperature and resource
supply (ANOVA; $F_{2,0.75} = 24.11$, $p<0.001$; electronic supplementary material, Table S1).
Whereas development time decreased both at warmer temperatures and at high-resource
supply, the decrease with temperature was greater at low-resource supply than at high-
resource supply. At low-resource supply, development time decreased by 15.45 days as
temperatures increased from 22 to 32°C, whereas at high-resource supply, it decreased by
6.38 days across this range (Fig. 2a, electronic supplementary material, Table S2).

Adult lifespan varied significantly with the interaction between temperature and resource
supply (ANOVA; $F_{2, 2.41} = 14.95$, $p<0.001$; electronic supplementary material, Table S1).
Although lifespan decreased both at warmer temperatures and at low-resource supply, the
decrease with temperature was greater at high-resource supply than at low-resource supply.
High-resource supply lifespan decreased by 8.89 days, whereas low-resource supply lifespan
decreased by 4.71 days as temperatures increased from 22 to 32°C (Fig 2b, electronic
supplementary material, Table S2).

The interaction between temperature and resource supply resulted in significant variation in
size at maturity (wing length) between resource levels (ANOVA; $F_{2,0.03} = 4.36$, $p=0.01$;
electronic supplementary material, Table S1). Adult size decreased both at warmer
temperatures and at low-resource supply, though the decrease with temperature was greater at
high-resource supply than at low-resource supply. At low-resource supply, size decreased by
0.37 mm as temperatures increased from 22 to 32°C, while at high-resource supply, size
decreased by 0.54 mm (Fig 2c, electronic supplementary material, Table S2).

Population fitness (r_{max})
Resource limitation depressed r_{max} to negative values at all temperatures, with a unimodal
relationship of r_{max} with temperature (Fig. 3, electronic supplementary material, Table S2).
Low-resource supply r_{max} increased from –0.09 at 22°C to –0.03 at 26°C and then decreased
acutely to –0.38 at 32°C. In contrast, at high-resource supply, r_{max} was always positive and
increased monotonically with temperature from 0.17 at 22°C to maximal growth (0.28) at
32°C.

Elasticity analysis
Juvenile survival was the most important contributor to r_{max} (electronic supplementary
material, Fig. S3). For example, at low-resource supply at 32°C, a 0.5 proportional increase
in juvenile survival would almost halve the rate of decline from –0.384 to –0.204 (electronic
supplementary material, Fig. S3a). In contrast, for the same treatment, a proportional increase
of the same magnitude for adult survival would increase r_{max} from –0.384 to –0.379 (Fig.
S3b), and fecundity would increase r_{max} from –0.384 to –0.376 (Fig. S3c). This underlines
how the temperature-dependence of r_{max} derives mainly from how resource supply level
impacts juvenile mortality and development, which determine the number of reproducing
individuals and the timing of reproduction, respectively. Fecundity and adult survival, on the
other hand, have relatively negligible effects on r_{max}, which suggests that the carry over effect
of reduced size at maturity on r_{max} is relatively weak.

DISCUSSION

Our results show that juvenile resource regimes can have far-reaching effects on the temperature-dependence of population-level fitness, r_{max}. Differences between the thermal response of traits at low- versus high-resource supply resulted in a marked divergence of the temperature-dependence of r_{max} between the two resource levels (Fig. 3). At low-resource supply, fitness was negative throughout and unimodal (declining steeply from 26 to 32°C). This indicates that population fitness becomes increasingly, and non-linearly constrained by resource limitation as temperatures increase. Fitness at high-resource supply was positive and monotonically increasing. This lack of unimodality indicates that the temperature at which fitness peaks is higher than 32°C when resources are abundant.

The elasticity analysis shows that the primary mechanism underlying the divergent temperature-dependence of r_{max} across resource levels is decreased juvenile survival at low resources, which decreased population-level reproductive output (Figs. 1, 3). At low-resource supply, the daily mortality rate started low and then increased over time, while at high-resource supply, it started high and then decreased to very low levels (Fig. 1e). The effect of resource limitation on fitness was further compounded by the increase in juvenile development time (Fig. 1a), which delayed the onset of reproduction.

Fecundity and adult lifespan had comparatively negligible effects on r_{max}, which suggests that the carry over effect of reduced size at maturity on r_{max} is relatively weak. For example, at high-resource supply, adult lifespan and body size were greater at 26°C than at 32°C, yet fitness at 32°C was predicted to be 25% higher (Figs. 2c, 3). This is because high-resource supply and increased temperature minimised juvenile mortality and optimised development rate. This allowed faster recruitment at 32°C, leading to increased fitness as greater numbers of individuals could contribute to population growth through reproductive output sooner than for other treatments. This is consistent with general studies of ectotherm fitness [57], including mosquitoes [30]. This is a key finding, as it implies that predictions about the effect of warming on vector abundance and disease transmission based on laboratory-derived trait data (which are generally from populations under high- or optima resource supply) likely underestimate the effect of temperature on development time and juvenile survival, and overestimate effects of temperature on lifespan and fecundity.

Indeed, the trait-level responses of our high-resource supply treatments correspond with studies that have synthesised laboratory-derived trait responses to temperature to estimate vector fitness and R_0. In these studies, the juvenile development rate of most mosquito vectors is expected to increase from ~0.07 day^{-1} at 22°C to ~0.14 day^{-1} at 32°C [8]. In the present study, development rate (1/development time, Fig. 1a) increased by a similar margin (~0.08 to ~0.17 day^{-1}) across the same temperature range. In contrast, at low-resource supply, we found juvenile development rate was ~0.05 day^{-1} at 32°C (Fig. 1a). Such differences in juvenile trait responses are likely to substantially alter predictions about the temperature-dependence of R_0. This underlines the importance of considering resource supply when predicting the temperature-dependence of R_0 for vector-borne diseases.

Juvenile survival decreased significantly with temperature, and was overall significantly lower at low-resource supply (Fig. 1). This is probably because somatic maintenance costs increase with metabolic rate [21], which cannot be met below a threshold resource supply level. This explains why the highest level of mortality occurred at 32°C at low-resource supply, where the energy supply-demand deficit was expected to be the largest.

The Gompertz-shaped juvenile survival curves observed at 22 and 26°C at low-resource supply (Fig. 1a) may well arise from the amount of resource being sufficient for somatic maintenance, but not for development. This could be a key line of future investigation because it points to the importance of understanding how resource availability combines with temperature and other environmental factors to affect natural mosquito populations. For example, the negative effects of resource limitation on population growth through increased juvenile development time and mortality may be exacerbated, as individuals remain in the vulnerable juvenile stages for longer, which may increase predation threat [58]. If resource availability increases with climatic warming, the negative effects of predation on population growth could be offset by increased development and recruitment rates [59]. Alternatively, population growth could be dampened, if climate change reduces the quantity of food available to ectotherms [60,61].

We did not measure the effect of temperature and resource supply on fecundity directly, but used the size-scaling of this trait to estimate this effect. This is because most of the effect of resource limitation on juveniles is expected to affect adult mosquitoes indirectly by reducing size at emergence and lifespan [29,49]. Predicted fecundity increased nonlinearly with temperature, mediated by resource supply levels (Fig 1d, electronic supplementary material, Table S1). Across both resource levels, these fecundity estimates are similar to datasets that are used to parameterise mosquito-borne disease transmission models (e.g. [62]). However, even substantial under- or overestimation of fecundity by our size-scaling predictions and the use of starved adult lifespans, would not affect our main conclusions. This is because predicted fitness was relatively insensitive to these traits (Figs. 3, S3).

While the increased negative carry over effects of temperature at resource limitation on adult traits may have had a relatively weak impact on fitness compared to juvenile traits, temperature × resource supply interactions may have important effects on other components of vector-borne disease transmission [63]. For example, larger individuals may have greater transmission potential because they are more likely to outlive a pathogen's external incubation period [64]. On the other hand smaller individuals may bite more frequently, which can increase transmission probability [65]. Also, larval nutrition [35] and temperature [66] can independently influence within-vector parasite development, but future studies could consider how the combined effects of temperature and resource supply affect this, and other important transmission traits.

In this study, we have not considered the temperature-dependence of resource supply itself (supply was held constant across temperatures in our experiments). In nature, the availability of resources may in fact be temperature-dependent. This is because microbial growth rates increase with temperature to some optimum, which may increase the concentration of food in the environment [9,67,68]. For example, *Anopheles* [69] and *Aedes* [70] mosquitoes can be reared exclusively on cultures of *Asaia* bacteria. We have also not directly addressed larval competition for resources by manipulating the number of larvae for a given resource supply level. Variation in larval density may introduce additional fitness constraints through interference and exploitative competition. It could also interact with temperature-dependent resource supply because a higher larval density will increase accumulation of waste products. These are interesting and potentially important avenues for future investigation.

We note that experimenting with more resource levels would not change our qualitative results, which we have shown to be robust using thorough sensitivity analyses. Indeed, the two resource levels we have chosen represent extremes, and it is reasonable to conclude that

mosquito population fitness in the field fluctuates, with resource fluctuations, between the radically different temperature responses as we have found here (Fig. 3). One avenue for future work is to find more accurate methods to estimate effective temperature-dependent fitness values in the field, accounting for resource fluctuations.

Organisms experience significant resource limitation over space and time in nature. This is particularly true for insects such as mosquitoes, which have juvenile stages restricted to small, ephemeral aquatic habitats that are susceptible to resource fluctuations [17–19,71]. Our study underlines the importance of the effects of resource supply on the temperature-dependence of population-level fitness of an important disease vector. In doing so, our findings suggest that current projections of how climatic warming affects vector-borne disease transmission may prove inaccurate because they generally fail to consider resource limitations. Our findings also underline the need for future research effort to be directed at better understanding how temperature and resource supply interact in the field, and how this, and interactions between other environmental factors, may influence other components of vector-borne disease systems.

References

1. Stanaway JD *et al.* 2016 The global burden of dengue: An analysis from the Global Burden of Disease Study 2013. *Lancet Infect. Dis.* **16**, 712–723. (doi:10.1016/S1473-3099(16)00026-8)

2. World Health Organization. 2014 A global brief on vector-borne diseases. *World Heal. Organ.* **WHO/DCO/WH**, 1–56.

3. Li R, Xu L, Bjørnstad ON, Liu K, Song T, Chen A, Xu B, Liu Q, Stenseth NC. 2019 Climate-driven variation in mosquito density predicts the spatiotemporal dynamics of dengue. *Proc. Natl. Acad. Sci.* **116**, 3624–3629. (doi:10.1073/pnas.1806094116)

4. Ng K-C, Chaves L, Tsai K-H, Chuang T-W. 2018 Increased Adult Aedes aegypti and Culex quinquefasciatus (Diptera: Culicidae) Abundance in a Dengue Transmission Hotspot, Compared to a Coldspot, within Kaohsiung City, Taiwan. *Insects* **9**, 98. (doi:10.3390/insects9030098)

5. Mordecai EA, Ryan SJ, Caldwell JM, Shah MM, LaBeaud AD. 2020 Climate change could shift disease burden from malaria to arboviruses in Africa. *Lancet Planet. Heal.* **4**, e416-423. (doi:10.1016/S2542-5196(20)30178-9)

6. Ryan SJ, Carlson CJ, Tesla B, Bonds MH, Ngonghala CN, Mordecai EA, Johnson LR, Murdock CC. 2021 Warming temperatures could expose more than 1.3 billion new people to Zika virus risk by 2050. *Glob. Chang. Biol.* **27**, 84–93. (doi:10.1111/gcb.15384)

7. Cator LJ, Johnson LR, Mordecai EA, Moustaid F El, Smallwood TRC. 2020 The role of vector trait variation in vector-borne disease dynamics. **8**, 1–25. (doi:10.3389/fevo.2020.00189)

8. Mordecai EA *et al.* 2019 Thermal biology of mosquito-borne disease. *Ecol. Lett.* **22**, 1690–1708. (doi:10.1111/ele.13335)

9. Cross WF, Hood JM, Benstead JP, Huryn AD, Nelson D. 2015 Interactions between temperature and nutrients across levels of ecological organization. *Glob. Chang. Biol.* **21**, 1025–1040. (doi:10.1111/gcb.12809)

10. Iwamura T, Guzman-Holst A, Murray KA. 2020 Accelerating invasion potential of disease vector Aedes aegypti under climate change. *Nat. Commun.* **11**, 2130. (doi:10.1038/s41467-020-16010-4)

11. Parham PE, Michael E. 2010 Modeling the Effects of Weather and Climate Change on Malaria Transmission. *Environ. Health Perspect.* **118**, 620–626. (doi:10.1289/ehp.0901256)

12. Liu-Helmersson J, Stenlund H, Wilder-Smith A, Rocklö J. 2014 Vectorial Capacity of Aedes aegypti: Effects of Temperature and Implications for Global Dengue Epidemic Potential. *PLoS One* **9**. (doi:10.1371/journal.pone.0089783)

13. Miazgowicz KL *et al.* 2020 Age influences the thermal suitability of Plasmodium falciparum transmission in the Asian malaria vector Anopheles stephensi. *Proc. R. Soc. B Biol. Sci.* **287**,

392		20201093. (doi:10.1098/rspb.2020.1093)
393	14.	Shocket MS, Verwillow AB, Numazu MG, Slamani H, Cohen JM, El Moustaid F, Rohr J,
394		Johnson LR, Mordecai EA. 2020 Transmission of West Nile and five other temperate
395		mosquito-borne viruses peaks at temperatures between 23°C and 26°C. *Elife* **9**, 1–67.
396		(doi:10.7554/eLife.58511)
397	15.	Hamlet A, Jean K, Perea W, Yactayo S, Biey J, Van Kerkhove M, Ferguson N, Garske T.
398		2018 The seasonal influence of climate and environment on yellow fever transmission across
399		Africa. *PLoS Negl. Trop. Dis.* **12**. (doi:10.1371/journal.pntd.0006284)
400	16.	Dell AI, Pawar S, Savage VM. 2011 Systematic variation in the temperature dependence of
401		physiological and ecological traits. *Proc. Natl. Acad. Sci.* **108**, 10591–10596.
402		(doi:10.1073/pnas.1015178108)
403	17.	Arrivillaga J, Barrera R. 2004 Food as a limiting factor for Aedes aegypti in water-storage
404		containers. *J. Vector Ecol.* **29**, 11–20.
405	18.	Barrera R, Amador M, Clark GG. 2006 Ecological factors influencing Aedes aegypti (Diptera:
406		Culicidae) productivity in artificial containers in Salinas, Puerto Rico. *J. Med. Entomol.* **43**,
407		484–492. (doi:10.1603/0022-2585(2006)43[484:EFIAAD]2.0.CO;2)
408	19.	Walsh RK, Facchinelli L, Ramsey JM, Bond JG, Gould F. 2011 Assessing the impact of
409		density dependence in field populations of Aedes aegypti. *J. Vector Ecol.* **36**, 300–307.
410		(doi:10.1111/j.1948-7134.2011.00170.x)
411	20.	Thomas MK, Aranguren-Gassis M, Kremer CT, Gould MR, Anderson K, Klausmeier CA,
412		Litchman E. 2017 Temperature-nutrient interactions exacerbate sensitivity to warming in
413		phytoplankton. *Glob. Chang. Biol.*, March. (doi:10.1111/gcb.13641)
414	21.	Kooijman SALM. 2000 *Dynamic energy and mass budgets in biological systems*. Cambridge:
415		Cambridge University Press.
416	22.	West GB, Brown JH, Enquist BJ. 2001 A general model for ontogenetic growth. *Nature* **413**,
417		628–631. (doi:10.1038/35098076)
418	23.	Savage VM, Gillooly JF, Brown JH, West GB, Charnov EL. 2004 Effects of Body Size and
419		Temperature on Population Growth. *Am. Nat.* **163**, 429–441. (doi:10.1086/381872)
420	24.	Amarasekare P, Savage V. 2012 A Framework for Elucidating the Temperature Dependence
421		of Fitness. *Am. Nat.* **179**, 178–191. (doi:10.1086/663677)
422	25.	Atkinson D. 1994 Temperature and Organism Size—A Biological Law for Ectotherms? In
423		*Adv. Ecol. Res.*, pp. 1–58. Academic Press.
424	26.	Rueda LM, Patel KJ, Axtell RC, Stinner RE. 1990 Temperature-Dependent Development and
425		Survival Rates of Culex quinquefasciatus and Aedes aegypti (Diptera: Culicidae). *J. Med.*
426		*Entomol.* **27**, 892–898. (doi:10.1093/jmedent/27.5.892)
427	27.	Forster J, Hirst AG, Atkinson D. 2012 Warming-induced reductions in body size are greater in
428		aquatic than terrestrial species. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 19310–19314.
429		(doi:10.1073/pnas.1210460109)
430	28.	Nasci RS. 1991 Influence of Larval and Adult Nutrition on Biting Persistence in Aedes aegypti
431		(Diptera: Culicidae). *J. Med. Entomol.* **28**, 522–526. (doi:10.1093/jmedent/28.4.522)
432	29.	Steinwascher K. 1982 Relationship Between Pupal Mass and Adult Survivorship and
433		Fecundity for Aedes aegypti. *Environ. Entomol.* **11**, 150–153. (doi:10.1093/ee/11.1.150)
434	30.	Dye C. 1984 Models for the Population Dynamics of the Yellow Fever Mosquito, Aedes
435		aegypti. *J. Anim. Ecol.* **53**, 247. (doi:10.2307/4355)
436	31.	Gilpin ME, McClelland GA. 1979 Systems analysis of the yellow fever mosquito Aedes
437		aegypti. *Fortschr. Zool.* **25**, 355–388.
438	32.	Walsh RK, Aguilar CL, Facchinelli L, Valerio L, Ramsey JM, Scott TW, Lloyd AL, Gould F.
439		2013 Regulation of aedes aegypti population dynamics in field systems: Quantifying direct and
440		delayed density dependence. *Am. J. Trop. Med. Hyg.* **89**, 68–77. (doi:10.4269/ajtmh.12-0378)
441	33.	Gimnig JE, Ombok M, Otieno S, Kaufman MG, Vulule JM, Walker ED. 2002 Density-
442		Dependent Development of Anopheles gambiae (Diptera: Culicidae) Larvae in Artificial
443		Habitats. *J. Med. Entomol.* **39**, 162–172. (doi:10.1603/0022-2585-39.1.162)
444	34.	Romeo Aznar V, Alem I, De Majo MS, Byttebier B, Solari HG, Fischer S. 2018 Effects of
445		scarcity and excess of larval food on life history traits of Aedes aegypti (Diptera: Culicidae). *J.*
446		*Vector Ecol.* **43**, 117–124. (doi:10.1111/jvec.12291)

447   35.   Shapiro LLM, Murdock CC, Jacobs GR, Thomas RJ, Thomas MB. 2016 Larval food quantity
448   affects the capacity of adult mosquitoes to transmit human malaria. *Proc. R. Soc. B Biol. Sci.*
449   **283**, 20160298. (doi:10.1098/rspb.2016.0298)
450   36.   Couret J, Dotson E, Benedict MQ. 2014 Temperature, larval diet, and density effects on
451   development rate and survival of Aedes aegypti (Diptera: Culicidae). *PLoS One* **9**, 1–9.
452   (doi:10.1371/journal.pone.0087468)
453   37.   Padmanabha H, Correa F, Legros M, Nijhout HF, Lord C, Lounibos LP. 2012 An eco-
454   physiological model of the impact of temperature on Aedes aegypti life history traits. *J. Insect*
455   *Physiol.* **58**, 1597–1608. (doi:10.1016/j.jinsphys.2012.09.015)
456   38.   Brady OJ, Hay SI. 2020 The global expansion of dengue: How Aedes aegypti mosquitoes
457   enabled the first pandemic arbovirus. *Annu. Rev. Entomol* **65**, 191–208.
458   39.   Arguez A, Durre I, Applequist S, Vose RS, Squires MF, Yin X, Heim RR, Owen TW. 2012
459   Noaa's 1981-2010 U.S. climate normals. *Bull. Am. Meteorol. Soc.* **93**, 1687–1697.
460   40.   Bargielowski IE, Lounibos LP, Carrasquilla MC. 2013 Evolution of resistance to satyrization
461   through reproductive character displacement in populations of invasive dengue vectors. *Proc.*
462   *Natl. Acad. Sci. U. S. A.* **110**, 2888–2892. (doi:10.1073/pnas.1219599110)
463   41.   Farjana T, Tuno N. 2012 Effect of body size on multiple blood feeding and egg retention of
464   Aedes aegypti (L.) and Aedes albopictus (Skuse) (Diptera: Culicidae). *Med. Entomol. Zool.* **63**,
465   123–131. (doi:10.7601/mez.63.123)
466   42.   Caswell H. 1989 *Matrix population models construction, analysis, and interpretation.*
467   Sunderland, Massachusettes: Sinauer Associates.
468   43.   Birch LC. 1948 The Intrinsic Rate of Natural Increase of an Insect Population. *J. Anim. Ecol.*
469   **17**, 15. (doi:10.2307/1605)
470   44.   Cole LC. 1954 The population consequences of life history phenomena. *Q. Rev. Biol.* **29**, 103–
471   137. (doi:10.1086/400074)
472   45.   R Core Team. 2018 R: A language and environment for statistical computing.
473   46.   Stubben C, Milligan B. 2007 Estimating and Analyzing Demographic Models Using the
474   popbio Package in R. *J. Stat. Softw.* **22**, 1–23. (doi:10.18637/jss.v022.i11)
475   47.   Therneau TM. 2020 A package for survival analyis in R.
476   48.   Agnew P, Hide M, Sidobre C, Michalakis Y. 2002 A minimalist approach to the effects of
477   density-dependent competition on insect life-history traits. *Ecol. Entomol.* **27**, 396–402.
478   (doi:10.1046/j.1365-2311.2002.00430.x)
479   49.   Briegel H. 1990 Metabolic relationship between female body size, reserves, and fecundity of
480   Aedes aegypti. *J. Insect Physiol.* **36**, 165–172. (doi:10.1016/0022-1910(90)90118-Y)
481   50.   Focks DA, Haile D., Daniels E, Mount GA. 1993 Dynamic Life Table Model for Aedes
482   aegypti (Diptera: Culicidae): Simulation Results and Validation. *J. Med. Entomol.* **30**, 1018–
483   1028. (doi:10.1093/jmedent/30.6.1018)
484   51.   Juliano SA. 1998 Species Introduction and Replacement among Mosquitoes: Interspecific
485   Resource Competition or Apparent Competition? *Ecology* **79**, 255. (doi:10.2307/176880)
486   52.   Skalski JR, Millspaugh JJ, Dillingham P, Buchanan RA. 2007 Calculating the variance of the
487   finite rate of population change from a matrix model in Mathematica. *Environ. Model. Softw.*
488   **22**, 359–364. (doi:10.1016/j.envsoft.2005.12.003)
489   53.   Hope RM. 2013 Rmisc: Rmisc: Ryan Miscellaneous. *R Packag. version 1.5*
490   54.   Caswell H, Naiman RJ, Morin R. 1984 Evaluating the consequences of reproduction in
491   complex salmonid life cycles. *Aquaculture* **43**, 123–134. (doi:10.1016/0044-8486(84)90016-4)
492   55.   de Kroon H, Plaisier A, van Groenendael J, Caswell H. 1986 Elasticity: The Relative
493   Contribution of Demographic Parameters to Population Growth Rate. *Ecology* **67**, 1427–1431.
494   (doi:10.2307/1938700)
495   56.   Jackson C. 2016 flexsurv: A platform for parametric survival modeling in R. *J. Stat. Softw.* **70**,
496   1–33.
497   57.   Huey RB, Berrigan D. 2001 Temperature, Demography, and Ectotherm Fitness. *Am. Nat.* **158**,
498   204–210. (doi:10.1086/321314)
499   58.   Benrey B, Denno RF. 1997 The Slow-Growth--High-Mortality Hypothesis: A Test Using the
500   Cabbage Butterfly. *Ecology* **78**, 987. (doi:10.2307/2265852)
501   59.   Culler LE, Ayres MP, Virginia RA. 2015 In a warmer Arctic, mosquitoes avoid increased

mortality from predators by growing faster. *Proc. R. Soc. B Biol. Sci.* **282**, 20151549. (doi:10.1098/rspb.2015.1549)

60. Lister BC, Garcia A. 2018 Climate-driven declines in arthropod abundance restructure a rainforest food web. *Proc. Natl. Acad. Sci.* **115**, E10397–E10406. (doi:10.1073/pnas.1722477115)

61. Huey RB, Kingsolver JG. 2019 Climate warming, resource availability, and the metabolic meltdown of ectotherms. *Am. Nat.* **194**. (doi:10.1086/705679)

62. Yang HM, Macoris MLG, Galvani KC, Andrighetti MTM, Wanderley DMV. 2009 Assessing the effects of temperature on the population of Aedes aegypti, the vector of dengue. *Epidemiol. Infect.* **137**, 1188–1202. (doi:10.1017/S0950268809002040)

63. Parham PE *et al.* 2015 Climate, environmental and socio-economic change: weighing up the balance in vector-borne disease transmission. *Philos. Trans. R. Soc. B Biol. Sci.* **370**, 20130551. (doi:10.1098/rstb.2013.0551)

64. Bellan SE. 2010 The Importance of Age Dependent Mortality and the Extrinsic Incubation Period in Models of Mosquito-Borne Disease Transmission and Control. *PLoS One* **5**, e10165. (doi:10.1371/journal.pone.0010165)

65. Scott TW, Morrison AC, Lorenz LH, Clark GG, Strickman D, Kittayapong P, Zhou H, Edman JD. 2000 Longitudinal Studies of Aedes aegypti (Diptera: Culicidae) in Thailand and Puerto Rico: Population Dynamics. *J. Med. Entomol.* **37**, 77–88. (doi:10.1603/0022-2585-37.1.77)

66. Shapiro LLM, Whitehead SA, Thomas MB. 2017 Quantifying the effects of temperature on mosquito and parasite traits that determine the transmission potential of human malaria. *PLOS Biol.* **15**, e2003489. (doi:10.1371/journal.pbio.2003489)

67. Craine JM, Fierer N, McLauchlan KK. 2010 Widespread coupling between the rate and temperature sensitivity of organic matter decay. *Nat. Geosci.* **3**, 854–857. (doi:10.1038/ngeo1009)

68. Smith TP, Thomas TJH, García-Carreras B, Sal S, Yvon-Durocher G, Bell T, Pawar S. 2019 Community-level respiration of prokaryotic microbes may rise with global warming. *Nat. Commun.* **10**, 5124. (doi:10.1038/s41467-019-13109-1)

69. Chouaia B *et al.* 2012 Delayed larval development in Anopheles mosquitoes deprived of Asaia bacterial symbionts. *BMC Microbiol.* **12**, S2. (doi:10.1186/1471-2180-12-S1-S2)

70. Souza RS, Virginio F, Riback TIS, Suesdek L, Barufi JB, Genta FA. 2019 Microorganism-Based Larval Diets Affect Mosquito Development, Size and Nutritional Reserves in the Yellow Fever Mosquito Aedes aegypti (Diptera: Culicidae). *Front. Physiol.* **10**, 1–24. (doi:10.3389/fphys.2019.00152)

71. Subra R, Mouchet J. 1984 The regulation of preimaginal populations of Aedes aegypti (L.) (Diptera: Culicidae) on the Kenya coast. *Ann. Trop. Med. Parasitol.* **78**, 63–70. (doi:10.1080/00034983.1984.11811774)

72. van den Heuvel MJ. 1963 The Effect of Rearing Temperature on the Wing Length, Thorax Length, Leg Length and Ovariole Number of the Adult Mosquito, Aedes Aegypti (L.). *Trans. R. Entomol. Soc. London* **115**, 197–216. (doi:10.1111/j.1365-2311.1963.tb00819.x)

73. Schneider CA, Rasband WS, Eliceiri KW. 2012 NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* **9**, 671–675. (doi:10.1038/nmeth.2089)

544
Fig. 1 | The effect of resource supply on the temperature-dependence of juvenile survival.
Survival curves at (**a**) low- and (**b**) high- resource supply by temperature with 95% confidence
bounds. **c,** Baseline mortality rates, a, by resource supply level across temperatures with 95% CIs.
Mortality rates were significantly lower at low-resource supply than at high-resource supply as
temperatures increased from 22 to 32°C (95% CIs at 26 and 32°C do not overlap). **d,** Change in
mortality rate trajectories, b, by resource supply level across temperatures with 95% CIs. Rate
trajectories were significantly lower at high-resource supply than at low-resource supply as
temperatures increased from 22°C (95% CIs at 26 and 32°C do not overlap). **e,** Logged daily mortality
rates ($\ln(\mu_x) = \ln(a) + bx$) show how mortality rates at low-resource supply started low and increased
with time; at high-resource supply they started high and decreased with time.

555
Fig. 2 | The combined effect of temperature and resource supply level on *Ae. aegypti* life history
traits. a-c, Resource supply significantly modulated the effect of temperature on all directly measured
traits, except adult lifespan at 26°C (**b**). Symbols denote the GLM-estimated means with 95% CIs
calculated from the standard errors (electronic supplementary material, Table S2) for the resource
supply levels at each temperature (circles, low-resource supply; triangles, high-resource supply). The
resulting ANOVAs of the GLMs for each trait are presented in Table S1 (electronic supplementary
material). **d,** Predicted fecundity increased significantly with temperature at both resource levels (non-
overlapping 95% CIs). The extent to which fecundity increased with temperature and resource level

was only significant at 26°C. The numbers of females that survived to adulthood (*n*) in each treatment
were: 22°C at low-resource supply *n* = 23, at high-resource supply *n* = 37; 26°C at low-resource
supply *n* = 29, at high-resource supply *n* = 30; 32°C at low-resource supply *n* = 10, at high-resource
supply *n* = 27. Boxplot horizontal lines represent medians. Lower and upper hinges are the 25th and
75th percentiles. Upper whiskers extend from the hinge to the largest value no further than $1.5 \times$ inter-
quartile range (IQR) from the hinge. The lower whisker extends from the hinge to the smallest value
at most $1.5 \times$ IQR of the hinge.

571
Fig 3. | Population-level *Ae. aegypti* fitness (r_{max}) by resource supply (0.1 (low) or 1 mg larva^{-1}
day^{-1} (high)) across temperatures. Fitness estimates for each treatment, with 95% confidence
intervals (CIs). The three data points for each treatment represent r_{max} estimated using the 95% CI
bounds of the exponents for the scaling of lifetime fecundity with wing length (electronic
supplementary material, eqn. S1, Fig. S2). The lightest greyscale hue estimates derive from the lower
95% CIs, the midrange hue estimates with trend lines derive from the slopes and the darkest hue
derive from the upper 95% CIs.