Title: The effect of resource limitation on the temperature-dependence of mosquito population fitness

Paul J. Huxley[1], Kris A. Murray[1, 2], Samraat Pawar[3], Lauren J. Cator[3]

[1]MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease Epidemiology, School of Public Health, Imperial College London, London, UK. [2]MRC Unit The Gambia at London School of Hygiene & Tropical Medicine, Banjul, The Gambia. [3]Department of Life Sciences, Imperial College London, Ascot, UK.

Email addresses of authors: Paul Huxley (p.huxley@imperial.ac.uk), Kris Murray (kris.murray@imperial.ac.uk), Samraat Pawar (s.pawar@imperial.ac.uk), Lauren Cator (l.cator@imperial.ac.uk)

Key words: climatic warming, abundance, temperature, population-level fitness, resource limitation, vector-borne disease

Correspondence should be sent to:

Paul J. Huxley
MRC Centre for Global Infectious Disease Analysis
Department of Infectious Disease Epidemiology
Imperial College London
Norfolk Place, London
W2 1PG
p.huxley@imperial.ac.uk

Statement of authorship: All authors designed the experiment, LJC provided the mosquitoes, PJH and SP performed the modelling, PJH collected data and analysed output data. PJH wrote the first draft of the manuscript, and all authors contributed substantially to revisions.

Data accessibility statement: Data will be made available on request, if required during the review process. Data will be made publicly available at publication from the Dryad Digital Repository.

Acknowledgements:

PJH, KAM, SP and LJC were funded by a Natural Environment Research Council grant [NE/L002515/1].

ABSTRACT

Laboratory-derived temperature dependencies of life history traits are increasingly being used
to make mechanistic predictions for how climatic warming will affect vector-borne disease
dynamics by affecting abundance dynamics of the vector population. These temperature-trait
relationships are typically estimated from populations reared on optimal resource supply,
even though natural populations of vectors are expected to experience variation in resource
supply, including intermittent resource limitation.

Using laboratory experiments on the mosquito *Aedes aegypti*, a principal arbovirus vector,
combined with stage-structured population projection modelling, we show that low-resource
supply significantly depresses the vector's maximal population growth rate across the entire
temperature range (22–32°C) and causes it to peak at a lower temperature than at high-
resource supply. This effect is primarily driven by an increase in juvenile mortality and
development time, combined with an exaggerated decrease in size of adults with temperature
at low-resource supply. Our study suggests that by ignoring resource limitation, projections
of vector abundance and disease transmission based on laboratory studies are likely to
substantially underestimate the effect of temperature on development time and juvenile
survival, and overestimate the effect of temperature on lifespan, size and fecundity. Our
results provide compelling evidence for future studies to consider resource supply when
predicting the effects of climate and habitat change on disease vectors.

INTRODUCTION

The global public health burden of some vector-borne diseases has increased substantially in recent decades [1,2]. The geographical distribution and abundance of vectors is strongly linked to transmission patterns of these diseases [3,4]. There is growing concern that climate and land-use change, coupled with rapid globalization, may result in shifts in both the distributions of vector species and the diseases they transmit [5]. However, mechanistic models that account for how fundamental environmental drivers (temperature, rainfall and resource supply) may independently and interactively affect population-level vector fitness are largely absent [6–8].

Ecological Metabolic Theory (EMT, including the Metabolic Theory of Ecology (MTE) and Dynamic Energy Budget (DEB) theory) links body size and environmental temperature through metabolic rate to individual-level trait performance and population-level fitness [9–11]. For example, the biological rates (e.g. metabolic, development, mortality) of ectotherms increase approximately exponentially with temperature up to some optimum before declining back to zero at a thermal maximum to affect population growth in a similar way [9,12,13]. While this core EMT prediction is well-supported, models linking metabolic rates to population-level fitness generally assume that resource supply is not a limiting factor [10,14]. Recent efforts to incorporate the effects of temperature and resource availability have focused on individual traits [15], or, when estimating population growth rates, have assumed that mortality is independent of resource availability [16].

Temperature and resource supply may interact to significantly affect fitness and its component traits, because the energy cost of somatic maintenance, development and growth generally increases with temperature [11,17]. The ability to meet this increasing demand is influenced by resource supply from an individual's environment. If the resources available to an individual do not keep pace with these increasing maintenance and growth requirements, its growth, development, and survival should be compromised. Ultimately, these effects should negatively affect fitness, and the severity of these effects should increase with temperature [15].

The thermal responses of individual traits have been used extensively in the rapidly expanding body of research on the responses of disease transmission rates to climate change [18–21]. The classical EMT framework has recently provided important contributions in this area [22–24], including vector-borne diseases [19,20]. However, despite the potential importance of resource supply, very little is known about how it interacts with temperature to influence vector demographics and transmission capacity (Mordecai et al. 2019). For example, Mordecai et al. [19,20] recently used a MTE trait-based framework to predict the temperature-dependence of the basic reproduction number (R_0), but all the parameterisations for the temperature-dependence of component traits come from mosquitoes provided with optimal food quantities. Thus, the question of whether and how temperature and resource supply may together modulate disease transmission through underlying traits remains open.

Additionally, temperature indirectly affects ectotherm fitness by determining size at maturity. Generally, size at maturity decreases with rising temperature (the size-temperature rule, [25], which also applies to disease vectors, such as mosquitoes [26]. For vectors specifically, female size is demographically and epidemiologically important because it is associated with longevity, fecundity, and biting behaviour. For example, individuals reared at warmer temperatures emerge earlier and are smaller, shorter-living and less fecund than conspecifics reared at cooler temperatures [26,27]. Reduced development time, due to increased temperature, can increase population-level fitness by decreasing generation time [28], but

cooler temperatures yield larger, more fecund and longer-living females, which are all traits
that generally increase fitness [27,29,30]. Furthermore, since larger females live longer, they
are more likely to outlive a pathogen's external incubation period, and hence have greater
transmission potential than smaller individuals [31]. However, smaller females may bite more
frequently, which can increase transmission probability [32]. All these lines of evidence
underline the importance of understanding the governing physiological mechanism(s) of the
size-temperature rule and determining how it relates to mosquito fitness and disease
transmission. However, it remains largely untested under resource limitation [25,33].

Studies on the effects of resource availability on mosquito traits generally report a positive
association between food level and juvenile development and survival [34–36], as well as
adult body size, nutritional reserves and fecundity. Many studies have examined how
resource quantity effects fitness in interaction with larval density, as competition for limited
resources between larvae is generally believed to be a major regulator of adult mosquito
abundance [37–39]. However, studies on the effects of resource supply on fitness have not
generally included temperature.

Given that wild mosquitoes are likely to experience variations of temperature and resource
availability [40–42], measuring the effects of both variables directly is important for
predicting their population dynamics. In this study, we investigated whether and how fitness
and its component traits are affected by the interaction between realistic temperature and
resource supply variation and interactions between these two factors in *Aedes aegypti*, a
principal mosquito vector of human arboviruses (e.g. dengue, yellow fever and zika; [43]).
We show that these factors can each act alone and interact with each other to significantly
modify the responses of body size, life history traits and fitness to temperature.

METHODS

To investigate the effects of temperature and resource supply on mosquito life history, we
employed a 3×2 factorial design comprised of three temperatures (22, 26, and 32°C) and
two resource supply levels: 0.1 (low-resource supply) and 1 mg larva^{-1} day^{-1} (high-resource
supply). These experimental temperatures span the range of average annual temperatures [44]
that this strain of *Ae. aegypti* is likely to experience in the wild (F16-19 originating from Fort
Meyer, FL; [45]). Our low-resource supply level was chosen because it is a level of limitation
often experienced by wild populations, and because previous work has demonstrated that it is
the lowest resource limitation that can be applied to this species without resulting in almost
complete juvenile mortality [40,41]. The high-resource supply level corresponds to the upper
mid-range of the high-resource supply levels used in Arrivillaga & Barrera [40] and Barrera
et al. [41], and is consistent with the levels of resource supply commonly used in laboratory
studies on this species [35,46].

Batches of approximately 300 *Ae. aegypti* eggs were randomly assigned to one of the three
experimental temperatures and immersed in plastic tubs containing 150 ml of tap water. Each
tub was provided with a pinch of powdered fish food (Cichlid Gold®, Hikari, Kyrin Food
Industries Ltd., Japan) to stimulate overnight hatching. The tubs were then submerged in
water baths (Grant Instruments: JAB Academy) set at either 22, 26, or 32°C. Water baths
were situated in a 20°C climate-controlled insectary with a 12L:12D photoperiod and 30
minutes of gradual transition of light levels to simulate sunrise and sunset. On the following
day, first instar larvae were separated into cohorts of 30 and held in tubs containing 150 ml of
water. We created three replicate tubs per treatment (90 individuals/treatment). Low-resource
supply treatments were provided 3 mg of powdered fish food and high-resource supply
treatments received 30 mg. Thereafter, resource levels were adjusted daily according to the

number of living individuals in each tub prior to feeding each day such that resource levels
were maintained at an approximately constant level during the juvenile lifespan. Rearing tubs
were cleaned and refilled with fresh tap water daily. Water volumes were also adjusted daily
in accordance with mortality to maintain larval density (0.2 larvae $\times$ ml^{-1}).

Fitness calculation

148 To calculate population-level fitness, we used our data to parameterise age-structured (Leslie)
149 population projection matrices [47], which describe change in a population over time:

$$\mathbf{N}_{t+1} = \mathbf{AN}_t, \qquad (1)$$

where $\mathbf{N}_t$ is a vector of the age structure at time t and $\mathbf{A}$ is the population projection matrix.
The first row of elements in $\mathbf{A}$ is populated with fecundity data. Fecundity is the number of
female offspring produced per female at age i. The sub-diagonal of $\mathbf{A}$ is populated with the
probability of survival from age i to age $i + 1$. Multiplying the transition matrix ($\mathbf{A}$; eqn. 1)
and age-structured population size vector ($\mathbf{N}_t$; eqn. 1) sequentially across time intervals yields
population dynamics. Once the stable age (stage) distribution of the abundance vector is
reached, the dominant eigenvalue of the system is the finite population rate of increase (λ)
[47]. Then, the intrinsic rate of population growth,

$$r_{\max} = \log(\lambda).$$

158 This is a population's inherent capacity to reproduce, and therefore a measure of population-
159 level fitness [10,29,48]. Negative $r_{\max}$ values indicate decline and positive ones, growth. The
160 projection matrices were built and analysed using the `popbio` R package [49,50].

Parameterisation

Immature development time and immature and adult survival proportions

Matrix survival elements (the sub-diagonal of the matrix $\mathbf{A}$; eqn. 1) were populated with
continuous survival proportions estimated using the Kaplan-Meier survival function in the
`survival` R package [49,51]. We assumed life stage duration (i.e. larva-to-pupa-to-adult)
was the mean duration of transitioning into and out of that stage, and a fixed age of adult
emergence at the mean age of emergence. Adult survival elements were populated with the
Kaplan-Meier proportions. Hatching-to-adult development times were calculated by
recording the day and time that egg eclosion, pupation and adult emergence occurred for each
individual. Upon pupation, mosquitoes were held in individual falcon tubes containing 5ml of
tap water. This enabled pupa-to-adult development durations and the lifespans of individual
starved adults to be recorded. In the absence of food, this condition forces adults to
metabolise the nutritional reserves accumulated during larval development. Therefore,
starved adult lifespan is a useful indicator of the carry over effects of temperature and
resource availability in the larval habitat [34,52].

Fecundity (the first row of $\mathbf{A}$; eqn. 1)

Allometric relationships between fecundity and size are commonly used in predictions of
population growth in *Aedes* mosquitoes [53,54]. Here, we estimated fecundity using
temperature and resource supply-dependent scaling between wing length and fecundity [46].
Wing length (L in eqn. 3) was measured to the nearest 0.01 mm from the alula to the wing tip,
excluding the apical fringe [55]. Wings (one per female) were removed, mounted onto glass

slides, photographed using a dissecting microscope and then measured with ImageJ software
[56].

Our analysis of the dataset indicated that the scaling between size and fecundity changes
significantly with temperature but not resource supply (Fig. S1). So, to estimate lifetime
fecundity (F in eqn. 3) from wing length for mosquitoes that we reared at 22 and 32°C at
high- and low-resource supply, we used the scaling coefficients from the Farjana & Tuno (50;
$n = 264$, $R^2 = 0.87$, $p<0.001$; eqn. 2) dataset at their at 20 and 30°C, respectively. For
mosquitoes that we reared at 26°C, there is no corresponding temperature treatment in the
Farjana & Tuno (50) dataset, so we used the scaling coefficients from the Briegel dataset at
27°C (56; $n = 206$, $R^2 = 0.77$, $p<0.001$; eqn. 2). Lifetime fecundity (F) was then divided by
lifespan and multiplied by 0.5 (assuming a 1:1 male-to-female offspring ratio) to obtain
temperature-specific individual daily fecundity. Later, we show that this level of variation in
the scaling of fecundity does not change our results qualitatively.

$$22°C, F = 0.93 + 3.16 \log(L)$$
$$26°C, F = 0.40 + 3.80 \log(L) \quad (2)$$
$$32°C, F = 0.26 + 4.08 \log(L)$$

Parameter sensitivity

We used the delta method to approximate 95% confidence intervals (CIs) for our fitness
calculations [47,57] to account for how uncertainty in survival and fecundity estimates is
propagated through to the r_{max} estimate. This method requires the standard errors of the
survival and fecundity element estimates. For survival, we used the standard errors estimated
by the Kaplan-Meier survival function. For fecundity, we used the standard errors of the
mean daily egg production rates for each treatment. As additional sensitivity analyses, we
recalculated fitness using the upper and lower 95% CIs of the exponents (Figs. 3, S1) for the
scaling of lifetime fecundity with wing length. The standard error for the scaling exponent at
27°C is not provided in Briegel [52], though the model's R^2 is satisfactory (0.77). Thus, to
determine the sensitivity of r_{max} to uncertainty in fecundity estimates at 26°C, we assumed a
similar 95% CI (± 0.25) to those determined through our analysis of the Farjana & Tuno [46]
dataset at 20 and 30°C.

Elasticity analysis

Elasticities were used to quantify the proportional contributions of individual life history
traits to r_{max}. Elasticities can provide valuable insights into the sensitivity of population
growth to hypothetical environmental perturbations during different life stages. Elasticity, e_{ij},
measures the proportional effect on λ of an infinitesimal change in an element of **A** (eqn. 1)
with all other elements held constant (the partial derivative) [58,59]. This partial derivative of
λ, with respect to each element of **A**, is $s_{ij} = \partial\lambda/\partial a_{ij} = v_i w_j$ with the dot product $\langle \mathbf{w}, \mathbf{v} \rangle = 1$.
Here, **w** is the dominant right eigenvector (the stage distribution vector of **A**), **v** is the
dominant left eigenvector (the reproductive value vector of **A**), and a_{ij} is the $i \times j^{th}$ element of
A. Elasticities can then be calculated using the relationship: $e_{ij} = a_{ij}/\lambda \times s_{ij}$. Multiplying an
elasticity by λ gives the absolute contribution of its corresponding a_{ij} to λ [58,59]. Absolute
contributions for juvenile survival and fecundity elements were summed and changed
proportionally to quantify the sensitivity of r_{max} to these traits.

Statistical analyses

All analyses were conducted using R [49]. In the first instance, to assess how temperature, resource supply, their interaction (whether low-resource supply treatments exhibited different thermal responses) and replicate influenced development time, lifespan and wing length, we fitted linear regression models. Model diagnostics and Shapiro-Wilk [60] tests were used to assess data normality. These data were nonlinear, positive and right skewed. Therefore, we used full factorial generalized linear models (GLM) with gamma distributions and identity link functions (predictor effects were considered additive) to determine the significance of each predictor on the thermal response of each of these traits. Replicate was included in these GLMs as a fixed effect. Goodness-of-fit was assessed using model deviance/d.f. scores and model residuals.

We investigated the effect of temperature and resource supply on juvenile mortality rate by fitting a set of candidate distributions (exponential, log-logistic, Gompertz and Weibull) to the survival data with R package `flexsurv`. The Gompertz survival function was the best fit to these data according to the Akaike Information Criterion (AIC) (Table S3). The final mortality model was obtained by dropping terms from the full model (consisting of temperature × resource supply + replicate as fixed effect predictors). If removing a term did not improve model fit (ΔAIC > −2), it was removed from the full model (Table S3). For each treatment, maximum likelihood methods executed in `flexsurv` estimated mortality parameters (and their 95% CIs) of the Gompertz model, $\mu_x = ae^{bx}$, where (a) is the baseline morality rate, and (b) is the change in mortality rate with time (i.e. the slope of the mortality trajectory). These parameter estimates were then used to determine the significance of the effects of temperature and resource supply on juvenile mortality.

RESULTS

All trait responses varied significantly with temperature and resource supply, with a significant interaction between the two environmental variables (Tables S1, S2; Figs. 1, 2). Thus, the realised effect of temperature on trait responses was consistently and significantly mediated by resource supply.

Baseline juvenile mortality rates, a, were significantly lower at low-resource supply than at high-resource supply as temperatures increased from 22 to 32°C (non-overlapping 95% CIs at 26 and 32°C; Fig 1c). Mortality rate trajectories, b, were significantly lower at high-resource supply than at low-resource supply as temperatures increased from 22 to 32°C (non-overlapping 95% CIs at 26 and 32°C; Fig 1d).

Development time varied significantly with the interaction between temperature and resource supply (ANOVA; $F_{2,0.75} = 24.11$, $p<0.001$; Table S1). Whereas development time decreased both at warmer temperatures and at high-resource supply, the rate of decrease with temperature was greater at low-resource supply than at high-resource supply. At low-resource supply, development time decreased by 17.01 days as temperatures increased from 22 to 32°C, whereas at high-resource supply, it decreased by 6.38 days across this range (Table S2, Fig. 2a).

Lifespan varied significantly with the interaction between temperature and resource supply (ANOVA; $F_{2,\ 2.41} = 15.34$, $p<0.001$; Table S1). Although lifespan decreased both at warmer temperatures and at low-resource supply, the rate of decrease with temperature was greater at high-resource supply than at low-resource supply. High-resource supply lifespan decreased

by 8.89 days, whereas low-resource supply lifespan decreased by 4.71 days as temperatures
increased from 22 to 32°C (Table S2, Fig 2b).

The interaction between temperature and resource supply resulted in significant variation in
size at maturity (wing length) between resource levels (ANOVA; $F_{2,0.03} = 4.36$, p=0.01; Table
S1). Size decreased both at warmer temperatures and at low-resource supply, though the rate
of decrease with temperature was greater at high-resource supply than at low-resource
supply. At low-resource supply, length decreased by 0.37 mm as temperatures increased from
22 to 32°C, while at high-resource supply, length decreased by 0.54 mm (Table S2, Fig 2c).

Population fitness (r_{max})

Resource limitation depressed r_{max} to negative values at all temperatures, with a unimodal
relationship of r_{max} with temperature (Fig. 3, Table S2). Low-resource supply r_{max} increased
from –0.09 at 22°C to –0.03 at 26°C and then decreased acutely to –0.38 at 32°C. In contrast,
at high-resource supply, r_{max} was always positive and increased monotonically with
temperature from 0.17 at 22°C to maximal growth (0.28) at 32°C.

Elasticity analysis

Juvenile survival was the most important contributor to r_{max} (Fig. 4). For example, at low-
resource supply at 32°C, a 0.5 proportional increase in juvenile survival would almost halve
the rate of decline from –0.384 to –0.204 (Fig. 4a). In contrast, for the same treatment, a
proportional increase of the same magnitude for adult survival would increase r_{max} from
–0.384 to –0.379 (Fig. 3b), and fecundity would increase r_{max} from –0.384 to –0.376 (Fig.
3c). This underlines how the temperature-dependence of r_{max} derives mainly from how
resource supply level impacts juvenile mortality and development, which determine the
number of reproducing individuals and the timing of reproduction, respectively. Fecundity
and adult survival, on the other hand, have relatively negligible effects on r_{max}, which
suggests that the carry over effect of reduced size at maturity on r_{max} is relatively weak.

DISCUSSION

Our results show that the temperature-dependence of population-level fitness and its
component life history traits can be significantly modulated by resource supply. When adult
lifespan and juvenile development time were expressed as rates (adult mortality and juvenile
development rate, respectively), all traits at high-resource supply met EMT expectations by
being positive and increasing nonlinearly with temperature (Fig. 2a, c) [61]. This result is
consistent with previous studies that have used laboratory-derived trait responses to
temperature to estimate *Ae. aegypti* fitness and R_0; these studies report similar responses of
traits to our high-resource supply responses. For example, in Mordecai et al.'s [20]
temperature-dependent R_0 predictions, juvenile *Ae. aegypti* development rate increased from
approximately 0.07 day^{-1} at 22°C to 0.14 day^{-1} at 32°C. In the present study, development
rate (Fig. 1a, Table S2) increased by a similar margin (~0.08 to ~0.17 day^{-1}) across the same
temperature range.

In contrast, at low-resource supply, although trait thermal responses qualitatively met EMT
predictions, they deviated markedly from the results of previous laboratory experiments. For
example, in the present study, at low-resource supply, juvenile development rate was ~0.03
day^{-1} at 22°C, which is 57% less than the rate (~0.07 day^{-1}) used at this temperature in
Mordecai et al. [20] to predict R_0. Similarly, at low-resource supply at 32°C, we found
juvenile development rate was ~0.05 day^{-1}; whereas in Mordecai et al. [20] it was 64%

greater (~0.14 day^{-1}). Juvenile survival proportions at low-resource supply also differed
markedly from the high-resource supply proportions used in Mordecai et al. [20]. In the
present study, survival was 0.51 at 22°C, 0.64 at 26°C and 0.22 at 32°C (Fig. 1a); whereas in
Mordecai et al. [20] it was approximately 0.83 between 20-22°C, 0.76 at 25°C, and 0.72 at
32°C.

These differences in juvenile trait responses are likely to substantially alter temperature-
dependent R_0 predictions. For example, when the abundance components of Mordecai et al.'s
R_0 framework [7] are parameterised with juvenile trait responses at our low-resource supply,
the estimated number of mosquitoes at 32°C decreases by almost one order of magnitude
(from ~257.68 to ~28.12). This underlines the importance of considering resource supply
when predicting temperature-dependent R_0.

The differences between the temperature-dependence of traits at low- versus high-resource
supply resulted in a marked divergence of the temperature-dependence of r_{max} (Fig. 3)
between the two treatments. At low-resource supply, fitness was negative throughout and
unimodal (declining steeply from 26 to 32°C). This indicates that population fitness becomes
increasingly constrained by resource limitation as temperatures increase. Fitness at high-
resource supply was positive, increasing and linear. This absence of unimodality suggests that
the temperature at which fitness peaks (T_{pk}) may be higher than 32°C when resources are
abundant. However, the effects of temperature and resource availability across a broader
upper range of temperatures would be required to verify this.

The elasticity analysis shows that the primary mechanism underlying the divergent trends of
of r_{max} across resource supply levels is decreased juvenile survival, which decreased
population-level fertility and, ultimately, fitness (Figs. 1, 3). As temperatures increased from
22 to 32°C, baseline juvenile mortality rates were significantly lower at low-resource supply
than at high-resource supply (Fig. 1c), but, at high-resource supply, rate trajectories (Fig. 1d)
were negative and significantly lower than at low-resource supply as temperatures increased
across this range. The effect of resource limitation on fitness was further compounded by the
increase in juvenile development time (Fig. 1a), which delayed the onset of reproduction.

Development time at both resource supply levels decreased nonlinearly with temperature,
though it decreased more steeply with temperature at low-resource supply than at high-
resource supply (Table S2, Fig. 1a). Development time was highest at 22°C for both resource
supply levels. However, high-resource supply possibly minimised development cost at 22°C,
which reduced high-resource supply development time (by 67%) relative to low-resource
supply at this temperature. At both resource supply levels, the steepest declines in
development time occurred as temperatures increased from 22 to 26°C, likely because this
change in temperature increased metabolic rate at the cost of development (thus reducing
development cost).

Juvenile survival decreased significantly with temperature, and it was significantly lower at
low-resource supply (Fig. 1). This is probably because somatic maintenance cost increases as
metabolic rate increases [11,17]; if the amount of resource available cannot satisfy basal
requirements, mortality will increase. This explains why the highest level of mortality
occurred at 32°C at low-resource supply, where the energy supply-demand deficit was likely
largest. The Gompertz-shaped juvenile survival curves observed at 22 and 26°C at low-
resource supply (Fig. 1a) may well arise from the amount of resource being enough for
somatic maintenance but insufficient for development. This could be a key line of future
investigation because it points to the importance of understanding how resource availability
combines with temperature and other environmental factors to affect natural mosquito

populations. For example, the negative effects of resource limitation on population growth through increased juvenile development time and mortality may be exacerbated, as individuals remain in the vulnerable juvenile stages for longer, which may increase predation threat [62]. If resource availability increases with climatic warming, the negative effects of predation on population growth could be offset by increased development and recruitment rates [63]. Alternatively, population growth could be dampened, if climate change reduces the quantity of food available to ectotherms [15,64].

The effects of resource supply on adult traits (size at emergence, adult lifespan and fecundity) were weak compared to its effects on juvenile traits (Fig. 4). For example, at high-resource supply, adult lifespan and body size were higher at 26°C than at 32°C, yet fitness at 32°C was predicted to be 39% higher (Fig. 3). This is because high-resource supply and increased temperature minimised juvenile mortality and optimised development rate. This allowed faster recruitment at 32°C, leading to increased fitness as greater numbers of individuals could contribute to growth through reproductive output sooner than for other treatments. This finding is consistent with general studies of ectotherm fitness [65], including mosquitoes [37]. Therefore, projections of vector abundance and disease transmission that assume high-resource supply are likely to underestimate the effect of temperature on development time and juvenile survival, and overestimate effects of temperature on lifespan and fecundity.

Size at emergence decreased significantly with temperature for both resource supply levels (Table S1, Fig. 1c), consistent with the size-temperature rule [25]. Resource supply drove significant intercept variation in the size-temperature relationship, which reflects how high-resource supply allowed larvae to accumulate greater energy reserves and grow to become larger adults than their less well-fed conspecifics that were exposed to the same temperature. Size decreased more steeply with temperature at high-resource supply than at low-resource supply. Wing length decreased by ~0.20 mm at both low- and high-resource supply as temperatures increased from 22 to 26°C (Table S2). However, as temperatures increased from 26 to 32°C, wing length decreased by ~0.17 and ~0.34 mm at low- and high-resource supply, respectively. For size at emergence, the decline with temperature at high-resource supply may reflect the optimal (or approaching optimal) attainable size at each temperature.

Like size at emergence, adult lifespan decreased more acutely with temperature at high-resource supply than at low-resource supply (Table S1, Fig. 1b). This derived from the difference in lifespan between resource supply levels at 22°C. As temperatures increased from 26 to 32°C, lifespans declined to a similar extent at both resource supply levels. At low-resource supply, lifespan decreased by ~3.69 days, whereas at high-resource supply it decreased by ~4.24 days (Table S2). However, as temperatures increased from 22 to 26°C, lifespan decreased by ~1.02 days at low-resource supply and by ~4.65 days at high-resource supply. This relatively flatter decline at low-resource supply may reflect how larvae respond to cooler temperatures when resources are limited. Cooler temperatures prolong development by slowing underlying rate processes [66]. These processes may have been slowed further by metabolic down-regulation in response to resource limitation, which is predicted to delay development but increase survival, as individuals invest available energy in maintenance [67,68]. At high-resource supply, lifespan declined linearly, which may reflect how higher levels of resource supply allow temperature to optimise trait responses. For lifespan, the decline with temperature at high-resource supply may reflect the optimal (or approaching optimal) attainable lifespan at each temperature.

We did not measure the effect of temperature directly on fecundity but used the size-scaling of this trait to estimate this effect. This is because most of the effect of resource limitation on juveniles is expected to affect adult mosquitoes indirectly, by reducing size at emergence,

with few direct carry over effects [52,69]. Predicted fecundity increased nonlinearly with
temperature, mediated by resource supply levels (Fig 1d, Table S1). Across both resource
levels, these fecundity estimates are similar to those reported in past studies. For example, in
Mordecai et al. [20], *Ae. aegypti* fecundity increased from approximately 3.58 to 11.20 eggs
female^{-1} day^{-1} as temperatures increased from 22 to 32°C compared to an increase from 4.29
to 14.17 eggs female^{-1} day^{-1} at high-resource supply and 5.05 to 12.25 eggs female^{-1} day^{-1} at
low supply, across the same temperature range. Any differences likely arise from the fact that
our size-scaling-based predictions ignore additional effects on fecundity, such as adult food
supply. However, even substantial under- or overestimation of fecundity by our size-scaling
predictions and the use of starved adult lifespans, would not affect our main conclusions
because predicted fitness was relatively insensitive to these traits (Figs. 3, 4).

Although we have not considered the temperature-dependence of resource supply itself
(supply was held constant across temperatures in our experiments), in nature, the availability
of resources may in fact be temperature-dependent. This is because microbial growth rates
increase with temperature to some optimum, which may increase the concentration of food in
the environment [8,70,71]. For example, *Anopheles* [72] and *Aedes* [73] mosquitoes can be
reared exclusively on cultures of *Asaia* bacteria. The temperature-dependence of resource
supply is an interesting and potentially important avenue of investigation that would further
elucidate how the interaction between temperature and resource availability affects the fitness
of mosquitoes.

We note that experimenting with more resource levels would not change our qualitative
results, which we have shown to be robust using thorough sensitivity analyses. Indeed, the
two resource levels we have chosen represent extremes, and it is reasonable to conclude that
mosquito population fitness in the field fluctuates, with resource fluctuations, between
radically different temperature responses as we have found here (Fig. 3). One avenue for
future work is to find more accurate methods to estimate effective temperature-dependent
fitness values in the field, accounting for resource fluctuations.

Organisms experience significant resource limitation over space and time in nature. This is
particularly true for insects such as mosquitoes, which have juvenile stages restricted to
small, ephemeral aquatic habitats that are susceptible to resource fluctuations [40–42,74]
Overall, our results provide compelling evidence for the importance of resource availability
in larval habitats in *Ae. aegypti* and other mosquito vectors in mediating the effect of climatic
warming on their population growth and, therefore, disease transmission. More generally,
considering the effect of resource availability is likely to be important for further advancing
Ecological Metabolic Theory, and improving its utility for understanding the responses of
holometabolous insects to environmental change.

References

1.   Stanaway JD *et al.* 2016 The global burden of dengue: An analysis from the Global Burden of Disease
Study 2013. *Lancet Infect. Dis.* **16**, 712–723. (doi:10.1016/S1473-3099(16)00026-8)

2.   World Health Organization. 2014 A global brief on vector-borne diseases. *World Heal. Organ.*
WHO/DCO/WH, 1–56.

3.   Li R, Xu L, Bjørnstad ON, Liu K, Song T, Chen A, Xu B, Liu Q, Stenseth NC. 2019 Climate-driven
variation in mosquito density predicts the spatiotemporal dynamics of dengue. *Proc. Natl. Acad. Sci.*
116, 3624–3629. (doi:10.1073/pnas.1806094116)

4.   Ng K-C, Chaves L, Tsai K-H, Chuang T-W. 2018 Increased Adult Aedes aegypti and Culex
quinquefasciatus (Diptera: Culicidae) Abundance in a Dengue Transmission Hotspot, Compared to a
Coldspot, within Kaohsiung City, Taiwan. *Insects* **9**, 98. (doi:10.3390/insects9030098)

5.   Mordecai EA, Ryan SJ, Caldwell JM, Shah MM, LaBeaud AD. 2020 Climate change could shift disease
burden from malaria to arboviruses in Africa. *Lancet Planet. Heal.* **4**, e416-423. (doi:10.1016/S2542-
5196(20)30178-9)

6.   Cator LJ, Johnson LR, Mordecai EA, Moustaid F El, Smallwood TRC. 2020 The role of vector trait
variation in vector-borne disease dynamics. **8**, 1–25. (doi:10.3389/fevo.2020.00189)

7.   Mordecai EA *et al.* 2019 Thermal biology of mosquito-borne disease. *Ecol. Lett.* **22**, 1690–1708.
(doi:10.1111/ele.13335)

8.   Cross WF, Hood JM, Benstead JP, Huryn AD, Nelson D. 2015 Interactions between temperature and
nutrients across levels of ecological organization. *Glob. Chang. Biol.* **21**, 1025–1040.
(doi:10.1111/gcb.12809)

9.   Brown JH, Gillooly JF, Allen AP, Savage VM, West GB. 2004 Toward a metabolic theory of ecology.
Ecology **85**, 1771–1789. (doi:10.1890/03-9000)

10. Savage VM, Gillooly JF, Brown JH, West GB, Charnov EL. 2004 Effects of Body Size and
Temperature on Population Growth. *Am. Nat.* **163**, 429–441. (doi:10.1086/381872)

11. Kooijman SALM. 2000 *Dynamic energy and mass budgets in biological systems*. Cambridge:
Cambridge University Press.

12. Gillooly JF. 2001 Effects of Size and Temperature on Metabolic Rate. *Science (80-.).* **293**, 2248–2251.
(doi:10.1126/science.1061967)

13. Gillooly JF, Charnov EL, West GB, Savage VM, Brown JH. 2002 Effects of size and temperature on
developmental time. *Nature* **417**, 70–73. (doi:10.1038/417070a)

14. Amarasekare P, Savage V. 2012 A Framework for Elucidating the Temperature Dependence of Fitness.
Am. Nat. **179**, 178–191. (doi:10.1086/663677)

15. Huey RB, Kingsolver JG. 2019 Climate warming, resource availability, and the metabolic meltdown of
ectotherms. *Am. Nat.* **194**. (doi:10.1086/705679)

16. Thomas MK, Aranguren-Gassis M, Kremer CT, Gould MR, Anderson K, Klausmeier CA, Litchman E.
2017 Temperature-nutrient interactions exacerbate sensitivity to warming in phytoplankton. *Glob.*
Chang. Biol., March. (doi:10.1111/gcb.13641)

17. Kooijman SALM. 1988 The von Bertalanffy growth rate as a function of physiological parameters: a
comparative analysis. In *Mathematical Ecology* (eds TG Hallam, LJ Gross, SA Levin), pp. 3–45.
Singapore.

18. Iwamura T, Guzman-Holst A, Murray KA. 2020 Accelerating invasion potential of disease vector
Aedes aegypti under climate change. *Nat. Commun.* **11**, 2130. (doi:10.1038/s41467-020-16010-4)

19. Mordecai EA *et al.* 2013 Optimal temperature for malaria transmission is dramatically lower than
previously predicted. *Ecol. Lett.* **16**, 22–30. (doi:10.1111/ele.12015)

20. Mordecai EA *et al.* 2017 Detecting the impact of temperature on transmission of Zika, dengue, and

chikungunya using mechanistic models. *PLoS Negl. Trop. Dis.* **11**, e0005568.
(doi:10.1371/journal.pntd.0005568)

21.   Parham PE, Michael E. 2010 Modeling the Effects of Weather and Climate Change on Malaria
Transmission. *Environ. Health Perspect.* **118**, 620–626. (doi:10.1289/ehp.0901256)

22.   Kirk D, Jones N, Peacock S, Phillips J, Molnár PK, Krkošek M, Luijckx P. 2018 Empirical evidence
that metabolic theory describes the temperature dependency of within-host parasite dynamics. *PLOS*
Biol. **16**, e2004608. (doi:10.1371/journal.pbio.2004608)

23.   Kirk D, Luijckx P, Stanic A, Krkošek M. 2019 Predicting the Thermal and Allometric Dependencies of
Disease Transmission via the Metabolic Theory of Ecology. *Am. Nat.* **193**, 661–676.
(doi:10.1086/702846)

24.   Molnár PK, Kutz SJ, Hoar BM, Dobson AP. 2013 Metabolic approaches to understanding climate
change impacts on seasonal host-macroparasite dynamics. *Ecol. Lett.* **16**, 9–21. (doi:10.1111/ele.12022)

25.   Atkinson D. 1994 Temperature and Organism Size—A Biological Law for Ectotherms? In *Adv. Ecol.*
Res., pp. 1–58. Academic Press.

26.   Rueda LM, Patel KJ, Axtell RC, Stinner RE. 1990 Temperature-Dependent Development and Survival
Rates of Culex quinquefasciatus and Aedes aegypti (Diptera: Culicidae). *J. Med. Entomol.* **27**, 892–898.
(doi:10.1093/jmedent/27.5.892)

27.   Costanzo KS, Westby KM, Medley KA. 2018 Genetic and environmental influences on the size-
fecundity relationship in Aedes albopictus (Diptera: Culicidae): Impacts on population growth
estimates? *PLoS One* **13**, e0201465. (doi:10.1371/journal.pone.0201465)

28.   Kingsolver JG, Huey RB. 2008 Size, temperature, and fitness: Three rules. *Evol. Ecol. Res.* **10**, 251–
268.

29.   Birch LC. 1948 The Intrinsic Rate of Natural Increase of an Insect Population. *J. Anim. Ecol.* **17**, 15.
(doi:10.2307/1605)

30.   Livdahl TP, Sugihara G. 1984 Non-Linear Interactions of Populations and the Importance of Estimating
Per Capita Rates of Change. *J. Anim. Ecol.* **53**, 573. (doi:10.2307/4535)

31.   Bellan SE. 2010 The Importance of Age Dependent Mortality and the Extrinsic Incubation Period in
Models of Mosquito-Borne Disease Transmission and Control. *PLoS One* **5**, e10165.
(doi:10.1371/journal.pone.0010165)

32.   Scott TW, Morrison AC, Lorenz LH, Clark GG, Strickman D, Kittayapong P, Zhou H, Edman JD. 2000
Longitudinal Studies of Aedes aegypti (Diptera: Culicidae) in Thailand and Puerto Rico: Population
Dynamics. *J. Med. Entomol.* **37**, 77–88. (doi:10.1603/0022-2585-37.1.77)

33.   Forster J, Hirst AG, Atkinson D. 2012 Warming-induced reductions in body size are greater in aquatic
than terrestrial species. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 19310–19314.
(doi:10.1073/pnas.1210460109)

34.   Agnew P, Hide M, Sidobre C, Michalakis Y. 2002 A minimalist approach to the effects of density-
dependent competition on insect life-history traits. *Ecol. Entomol.* **27**, 396–402. (doi:10.1046/j.1365-
2311.2002.00430.x)

35.   Couret J, Dotson E, Benedict MQ. 2014 Temperature, larval diet, and density effects on development
rate and survival of Aedes aegypti (Diptera: Culicidae). *PLoS One* **9**, 1–9.
(doi:10.1371/journal.pone.0087468)

36.   Romeo Aznar V, Alem I, De Majo MS, Byttebier B, Solari HG, Fischer S. 2018 Effects of scarcity and
excess of larval food on life history traits of Aedes aegypti (Diptera: Culicidae). *J. Vector Ecol.* **43**, 117–
124. (doi:10.1111/jvec.12291)

37.   Dye C. 1984 Models for the Population Dynamics of the Yellow Fever Mosquito, Aedes aegypti. *J.*
Anim. Ecol. **53**, 247. (doi:10.2307/4355)

38.   Gilpin ME, McClelland GA. 1979 Systems analysis of the yellow fever mosquito Aedes aegypti.
Fortschr. Zool. **25**, 355–388.

39. Southwood TR, Murdie G, Yasuno M, Tonn RJ, Reader PM. 1972 Studies on the life budget of Aedes
aegypti in Wat Samphaya, Bangkok, Thailand. *Bull. World Health Organ.* **46**, 211–226.

40. Arrivillaga J, Barrera R. 2004 Food as a limiting factor for Aedes aegypti in water-storage containers. *J.*
Vector Ecol. **29**, 11–20.

41. Barrera R, Amador M, Clark GG. 2006 Ecological factors influencing Aedes aegypti (Diptera:
Culicidae) productivity in artificial containers in Salinas, Puerto Rico. *J. Med. Entomol.* **43**, 484–492.
(doi:10.1603/0022-2585(2006)43[484:EFIAAD]2.0.CO;2)

42. Walsh RK, Facchinelli L, Ramsey JM, Bond JG, Gould F. 2011 Assessing the impact of density
dependence in field populations of Aedes aegypti. *J. Vector Ecol.* **36**, 300–307. (doi:10.1111/j.1948-
7134.2011.00170.x)

43. Brady OJ, Hay SI. 2020 The global expansion of dengue: How Aedes aegypti mosquitoes enabled the
first pandemic arbovirus. *Annu. Rev. Entomol* **65**, 191–208.

44. Arguez A, Durre I, Applequist S, Vose RS, Squires MF, Yin X, Heim RR, Owen TW. 2012 Noaa's
1981-2010 U.S. climate normals. *Bull. Am. Meteorol. Soc.* **93**, 1687–1697.

45. Bargielowski IE, Lounibos LP, Carrasquilla MC. 2013 Evolution of resistance to satyrization through
reproductive character displacement in populations of invasive dengue vectors. *Proc. Natl. Acad. Sci. U.*
S. A. **110**, 2888–2892. (doi:10.1073/pnas.1219599110)

46. Farjana T, Tuno N. 2012 Effect of body size on multiple blood feeding and egg retention of Aedes
aegypti (L.) and Aedes albopictus (Skuse) (Diptera: Culicidae). *Med. Entomol. Zool.* **63**, 123–131.
(doi:10.7601/mez.63.123)

47. Caswell H. 1989 *Matrix population models construction, analysis, and interpretation*. Sunderland,
Massachusettes: Sinauer Associates.

48. Cole LC. 1954 The population consequences of life history phenomena. *Q. Rev. Biol.* **29**, 103–137.
(doi:10.1086/400074)

49. R Core Team. 2018 R: A language and environment for statistical computing.

50. Stubben C, Milligan B. 2007 Estimating and Analyzing Demographic Models Using the popbio Package
in R. *J. Stat. Softw.* **22**, 1–23. (doi:10.18637/jss.v022.i11)

51. Therneau TM. 2020 A package for survival analyis in R.

52. Briegel H. 1990 Metabolic relationship between female body size, reserves, and fecundity of Aedes
aegypti. *J. Insect Physiol.* **36**, 165–172. (doi:10.1016/0022-1910(90)90118-Y)

53. Focks DA, Haile DG, Daniels E, Mount GA. 1993 Dynamic Life Table Model for Aedes aegypti
(Diptera: Culicidae): Analysis of the Literature and Model Development. *J. Med. Entomol.* **30**, 1003–
1017. (doi:10.1093/jmedent/30.6.1003)

54. Juliano SA. 1998 Species Introduction and Replacement among Mosquitoes: Interspecific Resource
Competition or Apparent Competition? *Ecology* **79**, 255. (doi:10.2307/176880)

55. van den Heuvel MJ. 1963 The Effect of Rearing Temperature on the Wing Length, Thorax Length, Leg
Length and Ovariole Number of the Adult Mosquito, Aedes Aegypti (L.). *Trans. R. Entomol. Soc.*
London **115**, 197–216. (doi:10.1111/j.1365-2311.1963.tb00819.x)

56. Schneider CA, Rasband WS, Eliceiri KW. 2012 NIH Image to ImageJ: 25 years of image analysis. *Nat.*
Methods **9**, 671–675. (doi:10.1038/nmeth.2089)

57. SKALSKI J, MILLSPAUGH J, DILLINGHAM P, BUCHANAN R. 2007 Calculating the variance of
the finite rate of population change from a matrix model in Mathematica. *Environ. Model. Softw.* **22**,
359–364. (doi:10.1016/j.envsoft.2005.12.003)

58. Caswell H, Naiman RJ, Morin R. 1984 Evaluating the consequences of reproduction in complex
salmonid life cycles. *Aquaculture* **43**, 123–134. (doi:10.1016/0044-8486(84)90016-4)

59. de Kroon H, Plaisier A, van Groenendael J, Caswell H. 1986 Elasticity: The Relative Contribution of

Demographic Parameters to Population Growth Rate. *Ecology* **67**, 1427–1431. (doi:10.2307/1938700)

60.   Shapiro SS, Wilk MB. 1965 An Analysis of Variance Test for Normality (Complete Samples).
Biometrika **52**, 591. (doi:10.2307/2333709)

61.   Dell AI, Pawar S, Savage VM. 2011 Systematic variation in the temperature dependence of
physiological and ecological traits. *Proc. Natl. Acad. Sci.* **108**, 10591–10596.
(doi:10.1073/pnas.1015178108)

62.   Benrey B, Denno RF. 1997 The Slow-Growth-High-Mortality Hypothesis: A Test Using the Cabbage
Butterfly. *Ecology* **78**, 987. (doi:10.2307/2265852)

63.   Culler LE, Ayres MP, Virginia RA. 2015 In a warmer Arctic, mosquitoes avoid increased mortality
from predators by growing faster. *Proc. R. Soc. B Biol. Sci.* **282**, 20151549.
(doi:10.1098/rspb.2015.1549)

64.   Lister BC, Garcia A. 2018 Climate-driven declines in arthropod abundance restructure a rainforest food
web. *Proc. Natl. Acad. Sci.* **115**, E10397–E10406. (doi:10.1073/pnas.1722477115)

65.   Huey RB, Berrigan D. 2001 Temperature, Demography, and Ectotherm Fitness. *Am. Nat.* **158**, 204–210.
(doi:10.1086/321314)

66.   Angilletta MJ. 2004 Temperature, Growth Rate, and Body Size in Ectotherms: Fitting Pieces of a Life-
History Puzzle. *Integr. Comp. Biol.* **44**, 498–509. (doi:10.1093/icb/44.6.498)

67.   Moore CG, Whitacre DM. 1972 Competition in Mosqviitoes. 2. Production of Aedes aegypti1 Larval
Growth Retardant at Various Densities and Nutrition Levels. *Ann. Entomol. Soc. Am.* **65**, 915–918.
(doi:10.1093/aesa/65.4.915)

68.   Storey KB, Storey JM. 1990 Metabolic Rate Depression and Biochemical Adaptation in Anaerobiosis,
Hibernation and Estivation. *Q. Rev. Biol.* **65**, 145–174. (doi:10.1086/416717)

69.   Steinwascher K. 1982 Relationship Between Pupal Mass and Adult Survivorship and Fecundity for
Aedes aegypti. *Environ. Entomol.* **11**, 150–153. (doi:10.1093/ee/11.1.150)

70.   Craine JM, Fierer N, McLauchlan KK. 2010 Widespread coupling between the rate and temperature
sensitivity of organic matter decay. *Nat. Geosci.* **3**, 854–857. (doi:10.1038/ngeo1009)

71.   Smith TP, Thomas TJH, García-Carreras B, Sal S, Yvon-Durocher G, Bell T, Pawar S. 2019
Community-level respiration of prokaryotic microbes may rise with global warming. *Nat. Commun.* **10**,
5124. (doi:10.1038/s41467-019-13109-1)

72.   Chouaia B *et al.* 2012 Delayed larval development in Anopheles mosquitoes deprived of Asaia bacterial
symbionts. *BMC Microbiol.* **12**, S2. (doi:10.1186/1471-2180-12-S1-S2)

73.   Souza RS, Virginio F, Riback TIS, Suesdek L, Barufi JB, Genta FA. 2019 Microorganism-Based Larval
Diets Affect Mosquito Development, Size and Nutritional Reserves in the Yellow Fever Mosquito
Aedes aegypti (Diptera: Culicidae). *Front. Physiol.* **10**, 1–24. (doi:10.3389/fphys.2019.00152)

74.   Subra R, Mouchet J. 1984 The regulation of preimaginal populations of Aedes aegypti (L.) (Diptera:
Culicidae) on the Kenya coast. *Ann. Trop. Med. Parasitol.* **78**, 63–70.
(doi:10.1080/00034983.1984.11811774)

Fig. 1 | The effect of resource supply on the temperature-dependence of juvenile survival across time (days) with 95% confidence bounds. Survival curves at (a) low- and (b) high- resource supply by temperature. c, Baseline mortality rates, a, by resource supply level across temperatures with 95% CIs. Mortality rates were significantly lower at low-resource supply than at high-resource supply as temperatures increased from 22 to 32°C (95% CIs at 26 and 32°C do not overlap). d, Change in mortality rate trajectories, b, by resource supply level across temperatures with 95% CIs. Rate trajectories were significantly lower at high-resource supply than at low-resource supply as temperatures increased from 22°C (95% CIs at 26 and 32°C do not overlap).

Fig. 2 | The combined effect of temperature and resource supply level on *Ae. aegypti* life history traits. a-c, Resource supply significantly modulated the effect of temperature on all directly measured traits, except adult lifespan at 26°C (**b**). Symbols denote the GLM-estimated means with 95% CIs (Table S2) for the resource supply levels at each temperature (circles are low-resource supply treatments; triangles are high-resource supply treatments). The resulting ANOVAs of the GLMs for each trait are presented in Table S1. **d,** Predicted fecundity increased significantly with temperature at both resource levels (non-overlapping 95% CIs). The extent to which fecundity increased with temperature and resource level was only significant at 26°C. The numbers of females that survived to adulthood (*n*) in each treatment were: 22°C at low-resource supply *n* = 23, at high-resource supply *n* = 37; 26°C at low-resource supply *n* = 29, at high-resource supply *n* = 30; 32°C at low-resource supply *n* = 10, at high-resource supply *n* = 27.

Fig 3. | Population-level *Ae. aegypti* fitness (r_{max}) by resource supply (0.1 (low) or 1 mg larva^{-1} day^{-1} (high)) across temperatures. Fitness estimates for each treatment, with 95% confidence intervals (CI). The three data points for each treatment represent r_{max} estimated using the 95% CI bounds of the exponents for the scaling of lifetime fecundity with wing length (eqn. 2, Fig. S1). The lightest greyscale hue estimates derive from the lower 95% CIs, the midrange hue estimates with trend lines derive from the slopes and the darkest hue derive from the upper 95% CIs.

Fig. 4 | Sensitivity of r_{max} to proportional changes in juvenile survival (a) adult survival (b) and
fecundity (c) for each *Ae. aegypti* population. Juvenile survival was the most important contributor
to r_{max}, as relatively small changes in the summed matrix elements for this trait would result in
relatively large changes in r_{max}. Sensitivity of r_{max} to adult traits was much weaker compared to
sensitivity to juvenile traits. Solid black vertical lines denote r_{max} at zero as represented by the
midrange grey hue estimates in Fig. 3. Dark blue lines denote high-resource supply (1 mg larva^{-1} day^{-1}); light blue lines denote low-resource supply (0.1 mg larva^{-1} day^{-1}).

Supplementary Information

Table S1 | Type II Analysis of Variance results from GLMs fitted to the responses of life history traits to temperature and resource levels. Significant effects are shown in boldface type. * $\Rightarrow$ $p<0.05$; ** $\Rightarrow$ $p<0.01$; *** $\Rightarrow$ $p<0.001$. Each pseudo R^2 (= residual deviance/null deviance) approximates how much variance the model was able to capture.

Trait	Predictor	χ^2	df	*F* value	p-value
Juvenile development time	**Temperature**	**9.74**	**2**	**314.64**	**<0.001 *****
$R^2 = 0.96$	**RS**	**50.30**	**1**	**3251.01**	**<0.001 *****
	Temperature: RS	**0.75**	**2**	**24.11**	**<0.001 *****
	Replicate	0.01	2	0.18	0.83
	Residuals	2.30	148		
Lifespan	**Temperature**	**34.54**	**2**	**209.76**	**<0.001 *****
$R^2 = 0.74$	**RS**	**3.18**	**1**	**34.57**	**<0.001 *****
	Temperature: RS	**2.41**	**2**	**15.34**	**<0.001 *****
	Replicate	0.40	2	2.47	0.08
	Residuals	11.94	148		
Body size	**Temperature**	**0.65**	**2**	**111.58**	**<0.001 *****
$R^2 = 0.79$	**RS**	**0.92**	**1**	**314.28**	**<0.001 *****
	Temperature: RS	**0.03**	**2**	**4.36**	**0.01 ***
	Replicate	0.003	2	0.48	0.62
	Residuals	0.41	141		

654 **Table S2 | Comparison of the effect of resource supply on the temperature-dependence of fitness**
655 **and its component traits.** The means with 95% CIs for development time, lifespan and wing were
656 estimated by the GLMs (Table 1). For fecundity, these were estimated using the `Rmisc` package in R.
657 For r_{max}, 95% CIs were approximated using the method described in [57].

Trait	Temperature (°C)	Mean ± 95% Confidence intervals	
		Low-resource supply	High-resource supply
	22	36.45 ± 1.78	12.19 ± 0.47
Development time (days)	26	27.86 ± 1.22	8.53 ± 0.37
	32	19.47 ± 1.53	5.81 ± 0.26
	22	6.61 ± 0.72	11.35 ± 0.99
Lifespan (days)	26	5.59 ± 0.54	6.70 ± 0.64
	32	1.90 ± 0.38	2.46 ± 0.28
	22	2.65 ± 0.06	3.13 ± 0.06
Wing length (mm)	26	2.45 ± 0.05	2.93 ± 0.06
	32	2.28 ± 0.07	2.59 ± 0.06
	22	5.05 ± 1.71	4.29 ± 0.28
Fecundity (eggs female^{-1} day^{-1})	26	4.42 ± 0.50	6.79 ± 0.46
	32	12.25 ± 5.00	14.17 ± 3.03
	22	−0.09 ± 0.02	0.17 ± 0.01
Population-level fitness (r_{max})	26	−0.03 ± 0.02	0.21 ± 0.03
	32	−0.38 ± 0.06	0.28 ± 0.05

658 **Table S3 | Simplification of the Gompertz juvenile survival model**. The full model includes the
659 effects of temperature × resource supply and replicate on mortality. The interaction model without
660 replicate won because it had the lowest AIC. ΔAICs were calculated as differences from the
661 interaction model.

Model terms	Model name	AIC	ΔAIC	df
Temperature × RS	**Interaction**	**1050.85**	**0**	**7**
Temperature × RS + replicate	Full	1053.05	+2.2	9
Temperature + RS	No interaction	1053.50	+2.7	5
Temperature	Temperature only	1055.99	+5.1	4
Resource	Resource only	1110.30	+59.5	3
None	Null	1108.70	+57.9	2

 Sensitivity of r_{max} to uncertainty in predicted fecundity

As an additional sensitivity analysis, we recalculated fitness using the upper and lower 95%
CIs of the exponents for the scaling of lifetime fecundity with wing length (eqn. 2). These are
shown in Fig. S1. The standard error for the scaling exponent at 27°C was not provided in
Briegel [52] but the model's R^2 is quite high (0.77). Thus, to determine the sensitivity of r_{max}
to uncertainty in fecundity estimates at 26°C, we assumed a similar 95% CI ($\pm$ 0.25) to that of
the Farjana et al. [46] dataset at 20 and 30°C.

Fig. S1 | a, Analysis of the Farjana et al. [46] dataset shows that the scaling of wing length and
fecundity in *Ae. Aegypti* is temperature-dependent. The scaling exponents (slopes) for both
resource supply levels are significantly higher at 30°C than at 20°C. However, the effect of resource
supply on fecundity is non-significant at the temperature level (not shown). The standard error for the
scaling exponent at 27°C is not shown because it is not provided in [52], so for 26°C, we assumed a
similar 95% CI to those in Farjana et al. ([46]; 3.80 $\pm$ 0.25). Despite these assumptions relating to
fecundity, our r_{max} calculations are robust to uncertainty/variation in the underlying scaling and
temperature dependences (Figs. 3 and 4).