A high-quality genome assembly and annotation of the gray mangrove, *Avicennia marina*

Running title: Genome assembly of the gray mangrove

Guillermo Friis[1*], Joel Vizueta[2], David R. Nelson[1], Basel Khraiwesh[1,3], Enas Qudeimat[1,3],

Kourosh Salehi-Ashtiani[1], Alejandra Ortega[4], Alyssa Marshell[5], Carlos M. Duarte[4], John A.

Burt[1]

[1]Center for Genomics and Systems Biology, New York University - Abu Dhabi, PO Box 129188,

Abu Dhabi, United Arab Emirates

[2]Departament de Genètica, Microbiologia i Estadística and Institut de Recerca de la

Biodiversitat (IRBio), Universitat de Barcelona, Barcelona, Spain

[3]Center for Desert Agriculture, Division of Biological and Environmental Sciences and

Engineering, King Abdullah University of Science and Technology (KAUST), Thuwal, 23955-

6900, Saudi Arabia

[4] Red Sea Research Center (RSRC) and Computational Bioscience Research Center, King

Abdullah University of Science and Technology (KAUST), Thuwal, 23955-6900, Saudi Arabia

[5]Department of Marine Science and Fisheries, College of Agricultural and Marine Sciences,

Sultan Qaboos University, Muscat, Oman

[*]Corresponding author: Center for Genomics and Systems Biology, New York University —

Abu Dhabi, PO Box 129188, Abu Dhabi, United Arab Emirates; Email: gfm3@nyu.edu; Tel:

+97126286739.

Abstract

The gray mangrove [*Avicennia marina* (Forsk.) Vierh.] is the most widely distributed mangrove

species, ranging throughout the Indo-West Pacific. It presents remarkable levels of geographic

variation both in phenotypic traits and habitat, often occupying extreme environments at the

edges of its distribution. However, subspecific evolutionary relationships and adaptive

mechanisms remain understudied, especially across populations of the West Indian Ocean. High-

quality genomic resources accounting for such variability are also sparse. Here we report the first

chromosome-level assembly of the genome of *A. marina*. We used a previously release draft

assembly and proximity ligation libraries Chicago and Dovetail HiC for scaffolding, producing a

456,526,188 bp long genome. The largest 32 scaffolds (22.4 Mb to 10.5 Mb) accounted for 98 %

of the genome assembly, with the remaining 2% distributed among much shorter 3,759 scaffolds

(62.4 Kb to 1 Kb). We annotated 23,331 protein-coding genes using tissue-specific RNA-seq

data, from which 13,312 were associated to GO terms. Genome assembly and annotated set of

genes yield a 96.7% and 92.3% completeness score, respectively, when compared with the

eudicots BUSCO dataset. Furthermore, an F_{ST} survey based on resequencing data successfully

identified a set of candidate genes potentially involved in local adaptation, and revealed patterns

of adaptive variability correlating with a temperature gradient in Arabian mangrove populations.

Our *A. marina* genomic assembly provides a highly valuable resource for genome evolution

analysis, as well as for identifying functional genes involved in adaptive processes and

speciation.

Key words: gray mangrove, *Avicennia marina*, genome assembly, HiC, Arabia

Introduction

Mangroves are a polyphyletic group of trees and shrubs that inhabit the intertidal zone of the tropic and sub-tropic coasts (Primavera et al. 2018; Hogarth 2015; Nagelkerken et al. 2008; Polidoro et al. 2010). Mangroves share several morphological and physiological adaptations to their harsh intertidal habitat, including traits for tolerance of high salinity, alternating dessication and submergence of soils across tidal cycles, and exposure of roots to hypoxic, sulfide-rich soils (Tomlinson 2016). Mangroves are of great ecological and economic importance, providing key functions such as high productivity, much of which is exported to surrounding ecosystems, and acting as breeding, nesting, nursery, foraging and shelter areas for a range of biota (Lee et al. 2014; Carugati et al. 2018; Primavera et al. 2018; Nagelkerken et al. 2008). Mangroves also serve as an important carbon sink, supporting climate change mitigation and adaptation potential (Duarte et al. 2013). The diversity of evolutionary origins and adaptive mechanisms found in mangroves provide compelling systems for studying patterns of trait evolution, lineage divergence and speciation (e.g., Zhou et al. 2007; Urashi et al. 2013; Xu et al. 2017b).

Of the approximately 70 mangrove species described, the gray mangrove *Avicennia marina* has the broadest latitudinal and longitudinal distribution (Tomlinson 2016; Spalding et al. 2010; Hogarth 2015). At least three sub-specific, partially allopatric taxa or 'varieties' have been described: *A. marina* var. *australasica*, restricted to southeastern Australia and New Zealand; *A. marina* var. *eucalyptifolia*, in northern regions of Australasia; and *A. marina* var. *marina*, that ranges from New Caledonia in the Pacific and across the entire Indian Ocean (Tomlinson 2016; Duke 1991; Li et al. 2016). The broad geographic distribution of *A. marina* is reflected in its presence across diverse environmental gradients (e.g. temperature, freshwater, sediment and

nutrient supply, salinity, tidal range) and spatial settings (e.g. open coastlines, coastal lagoons,

estuaries, deltas, coral fringes) (Duke 1990; Quisthoudt et al. 2012). Along with its widespread

distribution, several aspects of *A. marina* biology make it a promising model organism among

mangroves for the study of evolution based on genomic and molecular approaches. First, the

broad environmental gradients present across the *A. marina* range are mirrored by remarkable

geographic variation in phenotypic and life-history traits (Duke 1990; Tomlinson 2016), which

makes them an suitable system for studying evolutionary processes related to dispersal,

directional selection, and neutral evolution. Previous studies have reported the phylogenetic

relationships for the sub-varieties of *A. marina* and congeneric species (Duke et al. 1998; Li et al.

2016; Nettel et al. 2008), yet the specific drivers of lineage diversification remain understudied.

In particular, the extensive gray mangrove populations from the Eastern African and Arabian

coasts have rarely been included in reported DNA sequence-based analyses, either for diversity

screening or other purposes (e.g., Duke et al. 1998; Li et al. 2016). Second, *A. marina* can

tolerate highly variable and extreme environmental conditions, and often occupies marginal,

biologically limiting environments at the edges of its distribution (Morrisey et al. 2010). While

several biological structures and mechanisms of the gray mangrove physiology have been

described (Nguyen et al. 2015; Naidoo 2016), the genetic basis and pathways underlying such

tolerance are still largely unknown (Mehta et al. 2005; Jithesh et al. 2006; Xu et al. 2017a).

Understudied Arabian populations represent a particular gap due to their occurrence at the

extremes of air and water temperature, salinity, and aridity (Ben-Hasan and Christensen 2019;

Camp et al. 2018). Third, the *A. marina* genome is moderately small and structurally simple

compared with other eukaryotes, and presents a limited amount of transposable elements (Lyu et

al. 2018; Xu et al. 2017a; Das et al. 1995), which facilitates the identification of short

polymorphisms and structural variants.

Previous studies on *A. marina* using high-throughput sequencing techniques and genomic

approaches have been released. Two multi-species studies, including *A. marina,* explored

patterns of convergent evolution at functional genes (Xu et al. 2017a) and transposable elements

loads (Lyu et al. 2018), based on draft genomes that were recently made available online. Whole-

genome assemblies of *A. marina* and several other mangrove taxa have recently been used for

demographic inference (Guo et al. 2018) and convergent evolution analysis (He et al. 2020), but

the underlying genomic data are not publicly accessible.

Here, we report a high-quality, chromosome-level assembly obtained using proximity ligation

libraries as a resource for genome-based studies on *A. marina* and related mangrove species. An

structural and functional annotation based on RNA-seq data from multiple tissues is also

provided. In addition, we generated whole-genome shotgun data for a set of resequenced

individuals from six different populations along the coasts of the Arabian Peninsula. We used

these data to evaluate the performance of the assembly as a reference to study patterns of

adaptive variability at the genomic level. Resequencing data is also used for organelle

assembling.

Materials and Methods

Genome sequencing and assembly

A high-quality genome was produced combining the previously released draft assembly of

Avicennia marina (Xu et al. 2017a; GenBank assembly accession: GCA_900003535.1,

GCA_900174615.1; Lyu et al. 2018) and newly generated sequence data from proximity ligation

libraries. Preparation of proximity ligation libraries Chicago and HiC as well as scaffolding with

the software pipeline HiRise (Putnam et al. 2016) was conducted at Dovetail Genomics, LLC.

The sequenced sample sequenced was leaf tissue obtained from an individual located at Ras

Ghurab Island in the Arabian Gulf (Fig 1; Abu Dhabi, United Arab Emirates; 24.601°N,

4.566 °E). Briefly, for Chicago and the Dovetail HiC library preparation, chromatin was fixed

with formaldehyde. Fixed chromatin was then digested with DpnII and free blunt ends were

ligated. Crosslinks were reversed, and the DNA purified from protein, which was then sheared to

~350 bp mean fragment size. Libraries were generated using NEBNext Ultra enzymes and

Illumina-compatible adapters, and sequencing was carried out on an Illumina HiSeq X platform.

Chicago and Dovetail HiC library reads were then used as input data for genome assembly for

HiRise, a software pipeline designed specifically for using proximity ligation data to scaffold

genome assemblies (Putnam et al. 2016). A previously reported draft genome of *Avicennia*

marina (Lyu et al. 2018; Xu et al. 2017a; GenBank assembly accession: GCA_900003535.1)

was used in the assembly pipeline, excluding scaffolds shorter than 1Kb since HiRise does not

assemble them. Further details are provided in the Supplementary Information.

The mitochondrial genome was assembled using NOVOplasty2.7.2 (Dierckxsens et al. 2017) and

resequencing data based on Illumina paired-end 150 bp libraries from a conspecific individual

(See below; Supplementary Information). The maturase (matR) mitochondrial gene available in

NCBI (GenBank accession no. AY289666.1) was used for the input seed sequence. The

assembly of the chloroplast yielded a high number of alternative, unsolved assemblies and is

thereby no reported.

Fig. 1. Geography and adaptive variability in Arabian gray mangroves. (A) Locations of the six

stands sampled for whole genome resequencing (colored circles) and for RNA-seq (green star).

(B) F_{ST} genome scan based on 22,181 windows of 20 Kb. Boxplot outliers (coefficient=1.5) are

marked in red (C) t-SNE based on 200 SNP outliers linked to functional genes. The background

shows the correlation between t-SNE1 and t-SNE2 with the annual temperature range registered

in each one of the sampling locations. Temperature depicted in the legend is in °C.

Genome annotation

We performed the annotation of the *A. marina* genome using mRNA data from a set of tissues of

conspecific individuals. Samples were collected on the coast of the Eastern Central Red Sea

north of Jeddah in the Kingdom of Saudi Arabia (22.324 °N, 39.100 °E; Figure 1A). Total RNA

was isolated from root, stem, leaf, flower, and seed using TRIzol reagent (Invitrogen, USA).

RNA-seq libraries were prepared using TruSeq RNA sample prep kit (Illumina, Inc.), with

inserts that range in size from approximately 100-400 bp. Library quality control and

quantification were performed with a Bioanalyzer Chip DNA 1000 series II (Agilent), and

sequenced in a HiSeq2000 platform (Illumina, Inc.). Messenger RNA reads were mapped with

HISAT2 (Kim et al. 2015), and genome-referenced transcripts for each tissue were produced and

merged with StringTie (Pertea et al. 2015). Prediction of coding regions was performed with

TransDecoder (Haas and Papanicolaou 2015). The obtained gene annotation gff3 file was

validated and used to generate the reported gene annotation statistics with GenomeTools

(Gremme et al. 2013) and in-house Perl scripts. The Trinotate (Haas 2015) pipeline was then

implemented to conduct a homology-based functional annotation by using Swiss-Prot (Bairoch

and Apweiler 2000) and pfam (Bateman et al. 2002) databases, generating a final set of

annotated functional genes. Further details on mRNA sequencing and annotation scripts are

provided in the Supplementary Information.

Repetitive regions were first modelled *ab initio* using RepeatModeler v2.0.1 (Flynn et al. 2019)

in all scaffolds longer than 100 Kbp with default options. The resulting repeat library was used to

annotate and soft-mask repeats in the genome assembly with RepeatMasker 4.0.9 (Smit et al.

2015).

Gene completeness assessment

We assessed gene completeness in the genome assembly, and gene annotation, using BUSCO

(Benchmarking Universal Single-Copy Orthologs) v4.0.5 (--auto-lineage-euk option;

Waterhouse et al. 2018). BUSCO evaluations were conducted using the 255 and 2,326 single-

copy orthologous genes in Eukaryota_odb10 and Eudicots_odb10 datasets, respectively.

Adaptive variability analysis and functional assessment of A. marina *genome*

To test the potential of the assembly and annotation reported here as a resource for genomic-

based studies, we checked for regions of high divergence across the genome of *A. marina* using

newly generated whole-genome data. We resequenced 60 individuals from six different

populations from each of the major seas bordering Arabia (Figure 1A; Table S1, Supplementary

Information), including populations in the Red Sea, the Arabian Gulf and the Arabian Sea/Sea of

Oman. Arabia's regional seas present extreme, but divergent, environmental conditions for

mangrove growth. The northern Red Sea is characterized by cold winter temperatures and high

salinity (Carvalho et al. 2019), while the southern Persian/Arabian Gulf is the world's hottest sea

each summer and is also hypersaline, with both areas considered arid to hyperarid with limited

(<200 mm) rainfall (Vaughan et al. 2019). In contrast, the Arabian Sea and Sea of Oman have

normal oceanic salinity, and summer temperatures that are buffered by cold-water upwelling as a

result of the Indian Ocean monsoon, resulting in more benign environmental conditions

(Claereboudt 2019). Using vcftools (Danecek et al. 2011), we conducted an F_{ST} survey across the

six sampled populations based on 20 Kb sliding windows, and identifiied outlier loci associated

with functional genes. We then used these loci to explore geographic patterns of adaptive

variability by means of t-SNE analysis, testing for correlations between variability in sea surface

temperature and t-SNE scores. Details on sequencing, variant calling, and analytical procedures

are available in the Supplementary Information.

Data availability

Genome assembly, genome annotation and related supporting resources for facilitating the use of

these data have been deposited at DRYAD (doi:10.5061/dryad.3j9kd51f5). SNP matrix from

resequenced individuals used in adaptive variation analyses is also provided in this dataset in vcf

format. Raw resequencing data have been deposited in GenBank under the accessions SRA:

SRP265707; BioProject: PRJNA629068.

The genome assembly has also been deposited at DDBJ/ENA/GenBank along with raw sequence

data from Chicago and HiC libraries under the accession JABGBM000000000, SRA:

SRP265707; BioProject: PRJNA629068.

Datasets relating to the RNA-seq analysis have been deposited in Mendeley

(doi:10.17632/9tsp7fr28r). The RNA-seq reads have been deposited at Genbank under the

Bioproject (SRA) accession: PRJNA591919; Biosample accession: SAMN13391520.

Results and Discussion

Sequencing and genome assembly

We sequenced and assembled a reference genome of a gray mangrove individual from the

Arabian Gulf, an extreme environment at the northern limit of the species' distribution. Chicago

and Dovetail HiC libraries produced 235 million and 212 million 2x150 bp paired-end reads,

respectively, and 134.1 Gb data overall. Genome scaffolding with HiRise yielded an assembly of

456.5 Mb for a final sequence coverage of 293X; an L50/N50 equal to 15 scaffolds/13.98 Mb

and a relatively large number of ambiguous bases (i.e., N) inserted in the genome (10.6%; Table

1). The scaffold length distribution was heavily skewed towards extreme values (Table 1, Figure

2). The 32 longest scaffolds, ranging from 22.40 Mb to 10.58 Mb (median = 13.44 Mb)

accounted for 98.03% of the whole assembled genome, congruent with a chromosome number

2N = 64 reported earlier (He et al. 2020). The remaining 1.97% of the genome was distributed

among another 3,759 scaffolds ranging from 62.5 Kb to over 1 Kb (median = 1.8 Kb). The large

number of small scaffolds may be due to the high fragmentation of the draft genome used in the

assembly pipeline (Dovetails, personal communication). The integrity assessment of the *A.*

marina genome retrieved a 98.8% and a 96.7% of the tested BUSCO groups for the eukaryote

and the eudicots databases, respectively (Table 1). The remarkable discontinuity in length sizes,

as well as the integrity and high quality of the scaffolding lends considerable support to the

hypothesis of 32 chromosomes; further sequencing efforts involving long-read sequencing are

warranted for confirmation. The mitochondrial genome assembly was 22,019 pb long with a

46.4% GC content.

Table 1. Summary statistics for the genome assembly and annotation of *A. marina*. CDS

indicates protein-coding sequences.

Genome assembly	
Total length	456,526,188 bp
Number of scaffolds	3,791
N50/L50	13,979,447 bp/15 scaffolds
N90/L90	11,144,373 bp/29 scaffolds
Chromosome scale	10,583,658 bp/32 scaffolds
Longest scaffold	22,400,447 bp
Missingness	10.6%
GC content	35.2 %
BUSCO eukaryota database	C:98.8% [S:81.2%, D:17.6%], F:0.8%, M:0.4%, N:255
BUSCO eudicots database	C:96.7% [S:89.2%, D:7.5%], F:0.8%, M:2.5%, N:2,326
Genome annotation	
Number of genes	23,331
Number of annotated genes	21,147
Number of genes with GOs	13,312
Average gene length	5,383.7 bp
Number of CDS	59,445
Average CDS length (bp)	11,46.33 bp
Number of exons	493,357
Average exon length (bp)	307.06 bp
Number of introns	433,912
Average intron length (bp)	625.19 bp
BUSCO eukaryota database	C:97.2% [S:78.8%, D:18.4%], F:1.6%, M:1.2%, N:255
BUSCO eudicots database	C:92.3% [S:85.2%, D:7.1%], F:1.6%, M:6.1%, N:2,326

BUSCO parameters are C: Complete BUSCO; S: Complete and single-copy BUSCOs; D:

Complete and duplicated BUSCOs; F: Fragmented BUSCOs; M: Missing BUSCOs; and N:

Total BUSCO groups searched. CDS indicates protein-coding sequences.

Fig. 2. Length bar-plot of the longest 40 scaffolds arranged by decreasing size. The genome was

sequenced using proximity ligation libraries Chicago and Dovetails HiC, and the assembly was

carried out with HiRise pipeline.

Genome annotation

We identified 23,331 protein-coding genes for which 21,147 orthologs from other species were

identified, and 13,312 were associated to GO terms. The average gene length was 5.38 Kb, with

a mean of 8.3 exons and 18.6 introns per gene. BUSCO integrity analysis reported a 97.2% of

recovered BUSCOs for the eukaryota database, and a 92.3% in the case of the eudicots (Table

1). We also found that a total of 40.20% (188.5 Mb) of the *A. marina* assembly consisted of

repetitive elements, a value moderately larger than the 30.4% previously reported for the species

(Xu et al. 2017a). The greatest proportions corresponded to long terminal repeats and

unclassified elements (20% and 16.7%, respectively; Table S2, Supplementary Information).

Adaptive variability analysis and functional assessment of A. marina *genome*

We resequenced 60 individuals of *A. marina* from six different populations across the

environmentally diverse coasts of the Arabian Peninsula (Figure 1A) with a coverage of 85X.

After SNP calling and a strict filtering for quality and missing data, we obtained a dataset of

538,185 SNPs for 56 individuals. An F_{ST} scan based on sliding 20 Kb windows revealed a

heterogeneous landscape of differentiation and detected a peak of high divergence at the Scaffold

20 (Figure 1B). A total of 200 highly divergent loci were identified, from which 43 overlapped

with annotated genes associated to GO terms (Table S2, Figure S1). Several of these genes are

involved in the development of shoots, leaves and flowers (BAM2 and BAM7; DeYoung et al.

2006; Reinhold et al. 2011), root and seeds (FEI1; Basu et al. 2016), as well as in protein storage

(VTI12; Sanmartín et al. 2007). Importantly, we also found signals of differentiation in genes

involved in plant sensitivity to salt and osmotic stress (WRKY40; Chen et al. 2010), drought and

palatability to detritivorous crustaceans (LOX6; Grebner et al. 2013), supporting the role of

abiotic and biotic factors in the differentiation of Arabian mangroves (Table S3, Supplementary

Information). A t-SNE based 324 SNPs extracted from the functionally annotated, highly

divergent loci showed clear clustering patterns among sampled populations (Figure 1C). Loading

scores of retained t-SNE axes showed high correlation with the gradient of sea surface

temperature (SST; p-values below $8.3x10^{-14}$ and $2.0x10^{-16}$ for t-SNE1 and t-SNE2, respectively),

also congruent with a differentiation process driven by environmental factors. In light of these

results, further research into gene evolution and biological pathways involved in local adaptation

to the extreme environment of the Arabian mangroves is in process. These questions are,

however, beyond the scope of this report and thus will be presented elsewhere.

In conclusion, we report the first chromosome-scale assembly for the *Avicennia marina* genome

along with a comprehensive annotation based on tissue-specific RNA-seq data. The genome is

highly contiguous and complete, and we demonstrated that it is a valuable resource for variant

calling and the identification of functional, candidate genes underlying phenotypic and

environmental divergence among mangrove taxa. Improved scaffolding also enables the

identification of regions putatively under selection, including structural variants such as

chromosome rearrangements or copy number variations, all relevant for investigating questions

related to evolutionary biology and molecular ecology in this ecological and socioeconomically

important species.

Acknowledgements

The authors are grateful to NYUAD's Core Bioinformatics for analysis and computational

assistance. We thank Mark Priest, Dain McParland, Noura Al-Mansoori, Anique Ahmad, and

Ada Kovaliukaite for their participation in the fieldwork. We gratefully acknowledge the

Environment Agency Abu Dhabi (permit reference number 20181823a), and the Oman Ministry

of Environment and Climate Affairs, Director General of Nature Conservation (permit number:

6210/10/75), for providing permits for research related to this manuscript. This work was

supported by New York University Abu Dhabi Institute Grant (73 71210 CGSB9), NYUAD

Faculty Research Funds (AD060) and King Abdullah University of Science and Technology

baseline funding to CMD. We dedicate this manuscript to the memory of Basel Khraiwesh, who

had produced the RNA-seq data used for annotation in this study.

References

Bairoch, A., and R. Apweiler, 2000 The SWISS-PROT protein sequence database and its

supplement TrEMBL in 2000. *Nucleic Acids Research* 28 (1):45-48.

Basu, D., L. Tian, T. Debrosse, E. Poirier, K. Emch *et al.*, 2016 Glycosylation of a fasciclin-like

arabinogalactan-protein (SOS5) mediates root growth and seed mucilage adherence via a

cell wall receptor-like kinase (FEI1/FEI2) pathway in Arabidopsis. *PLOS One* 11 (1).

Bateman, A., E. Birney, L. Cerruti, R. Durbin, L. Etwiller *et al.*, 2002 The Pfam protein families

database. *Nucleic Acids Research* 30 (1):276-280.

Ben-Hasan, A., and V. Christensen, 2019 Vulnerability of the marine ecosystem to climate

change impacts in the Arabian Gulf—an urgent need for more research. *Global ecology*

and conservation 17:e00556.

Camp, E.F., V. Schoepf, P.J. Mumby, L.A. Hardtke, R. Rodolfo-Metalpa *et al.*, 2018 The future

of coral reefs subject to rapid climate change: lessons from natural extreme

environments. *Frontiers in Marine Science* 5:4.

Carugati, L., B. Gatto, E. Rastelli, M.L. Martire, C. Coral *et al.*, 2018 Impact of mangrove

forests degradation on biodiversity and ecosystem functioning. *Scientific Reports* 8

(1):13298.

Carvalho, S., B. Kürten, G. Krokos, I. Hoteit, and J. Ellis, 2019 The Red Sea, pp. 49-74 in *World*

Seas: An Environmental Evaluation, edited by C. Sheppard. Elsevier.

Chen, H., Z. Lai, J. Shi, Y. Xiao, Z. Chen *et al.*, 2010 Roles of Arabidopsis WRKY18, WRKY40

and WRKY60 transcription factors in plant responses to abscisic acid and abiotic stress.

BMC plant biology 10 (1):281.

Claereboudt, M.R., 2019 Oman, pp. 25-48 in *World Seas: An Environmental Evaluation*, edited

by C. Sheppard. Elsevier.

Danecek, P., A. Auton, G. Abecasis, C.A. Albers, E. Banks *et al.*, 2011 The variant call format

and VCFtools. *Bioinformatics* 27 (15):2156-2158.

Das, A.B., U.C. Basak, and P. Das, 1995 Variation in nuclear DNA content and karyotype

analysis in three species of Avicennia, a tree mangrove of coastal Orissa. *Cytobios* 84:93-

102.

DeYoung, B.J., K.L. Bickle, K.J. Schrage, P. Muskett, K. Patel *et al.*, 2006 The CLAVATA1-

related BAM1, BAM2 and BAM3 receptor kinase-like proteins are required for meristem

function in Arabidopsis. *The Plant Journal* 45 (1):1-16.

Dierckxsens, N., P. Mardulyn, and G. Smits, 2017 NOVOPlasty: de novo assembly of organelle

genomes from whole genome data. *Nucleic Acids Research* 45 (4):e18-e18.

Duarte, C.M., I.J. Losada, I.E. Hendriks, I. Mazarrasa, and N. Marbà, 2013 The role of coastal

plant communities for climate change mitigation and adaptation. *Nature Climate Change*

3 (11):961-968.

Duke, N., 1990 Phenological trends with latitude in the mangrove tree Avicennia marina. *The*

Journal of Ecology:113-133.

Duke, N., 1991 A systematic revision of the mangrove genus Avicennia (Avicenniaceae) in

Australasia. *Australian Systematic Botany* 4 (2):299-324.

Duke, N.C., J.A. Benzie, J.A. Goodall, and E.R. Ballment, 1998 Genetic structure and evolution

of species in the mangrove genus Avicennia (Avicenniaceae) in the Indo-West Pacific.

Evolution 52 (6):1612-1626.

Flynn, J.M., R. Hubley, C. Goubert, J. Rosen, A.G. Clark *et al.*, 2019 RepeatModeler2:

automated genomic discovery of transposable element families. *bioRxiv*:856591.

Grebner, W., N.E. Stingl, A. Oenel, M.J. Mueller, and S. Berger, 2013 Lipoxygenase6-

dependent oxylipin synthesis in roots is required for abiotic and biotic stress resistance of

Arabidopsis. *Plant Physiology* 161 (4):2159-2170.

Gremme, G., S. Steinbiss, and S. Kurtz, 2013 GenomeTools: a comprehensive software library

for efficient processing of structured genome annotations. *IEEE/ACM Transactions on*

Computational Biology and Bioinformatics 10 (3):645-656.

Guo, Z., X. Li, Z. He, Y. Yang, W. Wang *et al.*, 2018 Extremely low genetic diversity across

mangrove taxa reflects past sea level changes and hints at poor future responses. *Global*

Change Biology 24 (4):1741-1748.

Haas, B., 2015 Trinotate: Transcriptome functional annotation and analysis.

Haas, B., and A. Papanicolaou, 2015 TransDecoder (find coding regions within transcripts).

Github, nd https://github. com/TransDecoder/TransDecoder (accessed May 17, 2018).

He, Z., S. Xu, Z. Zhang, W. Guo, H. Lyu *et al.*, 2020 Convergent adaptation of the genomes of

woody plants at the land-sea interface. *National Science Review*.

Hogarth, P.J., 2015 *The biology of mangroves and seagrasses*: Oxford University Press.

Jithesh, M., S. Prashanth, K. Sivaprakash, and A. Parida, 2006 Monitoring expression profiles of

antioxidant genes to salinity, iron, oxidative, light and hyperosmotic stresses in the highly

salt tolerant grey mangrove, Avicennia marina (Forsk.) Vierh. by mRNA analysis. *Plant*

cell reports 25 (8):865-876.

Kim, D., B. Langmead, and S. Salzberg, 2015 hisat2. *Nature methods*.

Lee, S.Y., J.H. Primavera, F. Dahdouh-Guebas, K. McKee, J.O. Bosire *et al.*, 2014 Ecological

role and services of tropical mangrove ecosystems: a reassessment. *Global Ecology and*

Biogeography 23 (7):726-743.

Li, X., N.C. Duke, Y. Yang, L. Huang, Y. Zhu *et al.*, 2016 Re-evaluation of phylogenetic

relationships among species of the mangrove genus Avicennia from Indo-West Pacific

based on multilocus analyses. *PLOS One* 11 (10):e0164453.

Lyu, H., Z. He, C.I. Wu, and S. Shi, 2018 Convergent adaptive evolution in marginal

environments: unloading transposable elements as a common strategy among mangrove

genomes. *New phytologist* 217 (1):428-438.

Mehta, P.A., K. Sivaprakash, M. Parani, G. Venkataraman, and A.K. Parida, 2005 Generation

and analysis of expressed sequence tags from the salt-tolerant mangrove species

Avicennia marina (Forsk) Vierh. *Theoretical and Applied Genetics* 110 (3):416-424.

Morrisey, D.J., A. Swales, S. Dittmann, M.A. Morrison, C.E. Lovelock *et al.*, 2010 The ecology

and management of temperate mangroves, pp. 43-160. CRC Press, Boca Raton(USA).

Nagelkerken, I., S. Blaber, S. Bouillon, P. Green, M. Haywood *et al.*, 2008 The habitat function

of mangroves for terrestrial and marine fauna: a review. *Aquatic Botany* 89 (2):155-185.

Naidoo, G., 2016 The mangroves of South Africa: An ecophysiological review. *South African*

Journal of Botany 107:101-113.

Nettel, A., R.S. Dodd, Z. Afzal-Rafii, and C. Tovilla-Hernández, 2008 Genetic diversity

enhanced by ancient introgression and secondary contact in East Pacific black

mangroves. *Molecular Ecology* 17 (11):2680-2690.

Nguyen, H.T., D.E. Stanton, N. Schmitz, G.D. Farquhar, and M.C. Ball, 2015 Growth responses

of the mangrove Avicennia marina to salinity: development and function of shoot

hydraulic systems require saline conditions. *Annals of Botany* 115 (3):397-407.

Pertea, M., G.M. Pertea, C.M. Antonescu, T.-C. Chang, J.T. Mendell *et al.*, 2015 StringTie

enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature*

Biotechnology 33 (3):290.

Polidoro, B.A., K.E. Carpenter, L. Collins, N.C. Duke, A.M. Ellison *et al.*, 2010 The loss of

species: mangrove extinction risk and geographic areas of global concern. *PLOS One* 5

(4):e10095.

Primavera, J.H., D.A. Friess, H. Van Lavieren, and S.Y. Lee, 2018 The Mangrove Ecosystem,

pp. 18-51 in *World Seas: An Environmental Evaluation: Volume II: The Indian Ocean to*

the Pacific, edited by C. Sheppard. Academic Press.

Putnam, N.H., B.L. O'Connell, J.C. Stites, B.J. Rice, M. Blanchette *et al.*, 2016 Chromosome-

scale shotgun assembly using an in vitro method for long-range linkage. *Genome*

Research 26 (3):342-350.

Quisthoudt, K., N. Schmitz, C.F. Randin, F. Dahdouh-Guebas, E.M. Robert *et al.*, 2012

Temperature variation among mangrove latitudinal range limits worldwide. *Trees* 26

(6):1919-1931.

Reinhold, H., S. Soyk, K. Šimková, C. Hostettler, J. Marafino *et al.*, 2011 β-Amylase–Like

proteins function as transcription factors in Arabidopsis, controlling shoot growth and

development. *The Plant Cell* 23 (4):1391-1403.

Sanmartín, M., A. Ordóñez, E.J. Sohn, S. Robert, J.J. Sánchez-Serrano *et al.*, 2007 Divergent

functions of VTI12 and VTI11 in trafficking to storage and lytic vacuoles in Arabidopsis.

Proceedings of the National Academy of Sciences 104 (9):3645-3650.

Smit, A., R. Hubley, and P. Green, 2015 RepeatMasker Open-4.0. 2013–2015.

Spalding, M., M. Kainuma, and L. Collins, 2010 *World atlas of mangroves*. London, UK:

Earthscan.

Tomlinson, P.B., 2016 *The botany of mangroves*: Cambridge University Press.

Urashi, C., K.M. Teshima, S. Minobe, O. Koizumi, and N. Inomata, 2013 Inferences of

evolutionary history of a widely distributed mangrove species, Bruguiera gymnorrhiza, in

the Indo-West Pacific region. *Ecology and Evolution* 3 (7):2251-2261.

Vaughan, G.O., N. Al-Mansoori, and J.A. Burt, 2019 The Arabian Gulf, pp. 1-23 in *World Seas:*

An Environmental Evaluation, edited by C. Sheppard. Elsevier.

Waterhouse, R.M., M. Seppey, F.A. Simão, M. Manni, P. Ioannidis *et al.*, 2018 BUSCO

applications from quality assessments to gene prediction and phylogenomics. *Molecular*

Biology and Evolution 35 (3):543-548.

Xu, S., Z. He, Z. Guo, Z. Zhang, G.J. Wyckoff *et al.*, 2017a Genome-wide convergence during

evolution of mangroves from woody plants. *Molecular Biology and Evolution* 34

(4):1008-1015.

Xu, S., Z. He, Z. Zhang, Z. Guo, W. Guo *et al.*, 2017b The origin, diversification and adaptation

of a major mangrove clade (Rhizophoreae) revealed by whole-genome sequencing.

National Science Review 4 (5):721-734.

Zhou, R.C., K. Zeng, W. Wu, X.S. Chen, Z.H. Yang *et al.*, 2007 Population genetics of

speciation in nonmodel organisms: I. Ancestral polymorphism in mangroves. *Molecular*

Biology and Evolution 24 (12):2746-2754.

445