Title: Threshold assessment, categorical perception, and the evolution of reliable signaling

Running Title: Threshold assessment and honest signals

Authors: James H. Peniston*[1], Patrick A. Green[2,3], Matthew N. Zipple[3], Stephen Nowicki[3]

1. Department of Biology, University of Florida, Gainesville, FL, USA

2. Centre for Ecology and Conservation, College of Life and Environmental Sciences, University
of Exeter, Penryn, Cornwall, UK

3. Department of Biology, Duke University, Durham, NC, USA

* Corresponding author: jimmpen@ufl.edu

Author Contributions: All authors conceptualized the study together and contributed to model
design and interpretation. J.H.P. constructed the models and wrote the first draft of the
manuscript. P.A.G., M.N.Z., and S.N. contributed to the writing of the manuscript.

Acknowledgments: We are grateful to E.M. Caves, S. Peters, S. Johnsen, and S. Zlotnik for
providing value advice throughout this project. We also thank N. Kortessis and the members of
the UF SRNE journal club for providing comments on earlier versions of this manuscript.
Funding was provided by the Duke University Office of the Provost. PAG was supported by a
Human Frontier Science Program Fellowship # LT000460/2019-L. M.N.Z. was supported by an
NSF graduate research fellowship.

Abstract:

Animals often use assessment signals to communicate information about their quality to a

variety of receivers, including potential mates, competitors, and predators. But what maintains

reliable signaling and prevents signalers from signaling a better quality than they actually have?

Previous work has shown that reliable signaling can be maintained if signalers pay fitness costs

for signaling at different intensities and these costs are greater for lower quality individuals than

higher quality ones. Models supporting this idea typically assume that continuous variation in

signal intensity is perceived as such by receivers. In many organisms, however, receivers have

threshold responses to signals, in which they respond to a signal if it is above a threshold value

and do not respond if the signal is below the threshold value. Here, we use both analytical and

individual-based models to investigate how such threshold responses affect the reliability of

assessment signals. We show that reliable signaling systems can break down when receivers

have an invariant threshold response, but reliable signaling can be rescued if there is variation

among receivers in the location of their threshold boundary. Our models provide an important

step towards understanding signal evolution when receivers have threshold responses to

continuous signal variation.

Keywords: Animal communication, honest signals, mate choice, sensory ecology, sexual

selection, signaling theory

Introduction:

In contexts ranging from mate choice to aggression, and in taxa spanning vertebrates and invertebrates, animals use signals to assess each other (Maynard Smith and Harper 2003; Searcy and Nowicki 2005; Seyfarth et al. 2010). These signals can include any behavioral or morphological trait of one individual (the "signaler") that has evolved to convey information to another individual (the "receiver"), such as the song of a bird, the sex pheromone of a moth, or the aposematic coloration of a poison frog (Bradbury and Vehrencamp 2011). The information conveyed in signals often regards the quality of the signaler, such as the signaler's size (e.g., call frequency in frogs; Ryan 2001) or physiological condition (e.g., plumage brightness in some birds; Lindström and Lundström 2000). A central question for researchers studying animal signaling systems is, what maintains the reliability of assessment signals (Searcy and Nowicki 2005)? That is, if it benefits signalers to produce a signal indicating a better quality than they actually have, why do signalers produce a signal that reliably (i.e., "honestly") communicates quality?

In a seminal verbal model, Zahavi (1975) suggested that assessment signals could be reliable indicators of quality if signalers pay costs for expressing signals. This idea was termed the "handicap principle" in that signalers of the highest quality could afford to pay greater handicaps to produce a signal than could signalers of lower quality. Although reliable signaling did not evolve when Zahavi's original assumptions were made explicit in genetic models (Maynard Smith 1976; Kirkpatrick 1986), Grafen (1990*a*, 1990*b*) later showed, using both population genetic and game theory models, that the handicap principle can lead to the evolution of reliable signaling if two conditions are met. First, the costs of signaling at a given intensity (e.g., producing a large or colorful signal) must be greater for signalers of lower quality

compared to signalers of higher quality. Second, receivers should be able to assess continuous

variation in signal intensities and thereby gauge signaler quality; that is, as signal intensity

increases continuously, so does the benefit the signaler gains from receivers (We note that Penn

and Számadó (2020) suggested that Grafen's models are substantively different from Zahavi's

original conceptualization. Here, we refer to them both as the "handicap principle" for

consistency with the literature). Johnstone (1997) later developed graphical versions of Grafen's

(1990*a*, 1990*b*) mathematical models, illustrating how the optimal level of signaling—the

equilibrium point at which the net benefits are greatest—will be different for lower *versus* higher

quality individuals. This variation in equilibrium points leads to reliable signaling (Johnstone

1997).

The cost-based reliable signaling theory developed by Zahavi (1975), Grafen (1990*a*,

1990*b*), and Johnstone (1997) has played a crucial role in our understanding of the evolution of

animal signals (reviewed in: Maynard-Smith and Harper 2003; Searcy and Nowicki 2005). Other

studies have recognized, however, that some assumptions of these models do not necessarily

reflect the reality of animal signaling systems. For example, the initial models of Grafen (1990*a*,

1990*b*) did not incorporate the concept of perceptual error—when a signaler's true signal value is

not perceived as such by a receiver. In a subsequent series of models, Grafen and Johnstone

showed that adding perceptual error into models of signal evolution makes otherwise continuous

signaling systems more discrete, such that fewer equilibrium signaling values exist (Johnstone

and Grafen 1992; Grafen and Johnstone 1993; Johnstone 1994). Reliable signaling can be

maintained in these models as long as signalers with greater signal intensity are more likely to be

perceived as having greater intensity signals (Johnstone and Grafen 1992). Furthermore, a

continuum of equilibrium signaling values reemerges if perceptual errors are more common at

higher signal intensities or if signaling costs increase more rapidly at higher signal intensities

(Johnstone 1994). Other researchers have found results similar to those of Grafen and Johnstone

when incorporating other aspects of biological realism into models of signal evolution: when the

assumptions of the classic handicap models are relaxed, signaling systems are altered, but

reliability can be maintained (e.g., Lachmann and Bergstrom 1998; Proulx 2001).

The aforementioned studies have advanced our understanding of signal evolution, but

they have all maintained the assumption that continuous variation in signal intensity is perceived

as such by receivers. Some organisms have threshold responses to assessment signals, however,

in which signal receivers respond in a binary manner to continuous variation in signal intensity

(e.g., Masataka 1983; Zuk et al. 1990; Reid and Stamps 1997; Stoltz and Andrade 2010; Beckers

and Wagner 2011; Robinson et al. 2011; Roff 2015). In these systems, receivers respond to a

signal if it is above a threshold value and do not respond if the signal is below the threshold

value. For example, female variable field crickets (*Gryllus lineaticeps*) prefer males producing

chirps at rates above 3.0 chirps per second but do not discriminate between chirp rate variants

that both lie either above or below this threshold (Beckers and Wagner 2011).

Threshold responses may reflect a behavioral decision by receivers (e.g., Beckers and

Wagner 2011), but they may also be linked to an animal's perceptual system. Categorical

perception, for example, occurs when continuous variation in a stimulus is perceived as

belonging to distinct categories, with individuals showing an increased capacity to discriminate

stimuli that fall into different categories as compared to stimuli that fall in the same category

(Harnad 1987; Green et al. 2020). For example, a recent study found that female zebra finches

(*Taeniopygia guttata*) show categorical perception of the orange to red color continuum

representative of male beak color: females labeled eight color variants as lying in two categories

and showed better discrimination of cross-category variants as compared to equally-spaced

within-category variants (Caves, Green et al. 2018). Categorical perception of signal variation,

such as that found in zebra finches (Caves, Green et al. 2018), túngara frogs (Baugh et al. 2008),

and sparrows (Nelson and Marler 1989), would likely lead to a threshold response if categories

are treated in an binary fashion, thereby violating the continuous assumptions of cost-based

models of reliable signaling.

While some models have investigated when different types of threshold responses will

evolve (Janetos 1980; Real 1990; Svennungsen et al. 2011; Bleu et al. 2012), few studies have

explored how threshold responses influence the evolution of reliable signaling. Broom and

Ruxton (2011) showed that when receivers respond to signals in a threshold manner it leads to

the evolution of "all-or-nothing" signaling systems. That is, signalers with a quality below a

critical value all produce the same cheap signal while signalers with a quality above the critical

value all produce the same expensive signal. The model of Broom and Ruxton (2011) assumed

that receivers all had the same threshold value. However, it is conceivable that receivers could

vary in their threshold values, for example due to environmental conditions or developmental

history. Here, we explore the evolutionary implications of such inter-individual variation in

threshold values.

We begin by developing a model of signal evolution that assumes receivers respond to

continuous signal variation in a continuous fashion (akin to the models of Grafen 1990a, 1990b;

Johnstone 1997). This model requires some simplifying assumptions, but it provides a clear

demonstration of how reliable signals evolve when there is not a threshold response (the typical

assumption). We then show how threshold responses to continuous signal variation can remove

the variation in equilibrium signal intensities central to current models of reliable signaling, but

how introducing inter-individual variation in threshold responses can rescue the evolution of

reliable signals. In addition to these analytical models, we also develop individual-based

simulations to assess the robustness of our conclusions under more realistic ecological and

genetic assumptions, as well as to ask what role additional complexities in receiver choice can

have on the evolution of signals responded to in a threshold fashion. While most of our models

assume that the mean threshold value of receivers is fixed, and thus ignore evolution in receivers,

we also ran additional individual-based simulations in which we allowed both signalers' signals

and receivers' threshold values to evolve. The results of these co-evolutionary models are

consistent with our simpler models, and confirm that our general conclusions hold even when

receivers and signalers coevolve.

Continuous Assessment Model

We start by constructing a model of signal evolution that assumes receivers respond to

continuous signal variation in a continuous fashion (i.e., there is not a threshold response). This

model can be thought of as an intermediate between those of Grafen (1990a, 1990b) and

Johnstone (1997): it can be mathematically difficult to incorporate ecological complexity into

Grafen's models, while Johnstone's model is purely graphical and thus cannot be analytically

evaluated. This model, and our other analytical models below, are rather simple models in which

we do not model the evolution of receivers. Instead, we assume that receivers have a fixed

reaction norm which is a continuous function that scales positively with signal intensity (i.e.,

receivers are more likely to respond to signals of greater intensity). We also assume that the

optimal signal intensity for a signaler is that which maximizes its net benefit. Therefore, the

equilibrium signal intensity is defined as the maximization of the difference between $b(s)$ and

$c(s)$, where $b(s)$ is the fitness benefit received by a signaler for producing a signal intensity s and

$c(s)$ is the fitness cost associated with producing signal intensity s.

In line with previous models (Grafen 1990*a*, 1990*b*; Johnstone 1997), we assume that

fitness costs increase linearly with signal intensity and that these costs increase more rapidly for

low-quality signalers than high-quality ones (the same qualitative results should apply for any

monotonically increasing cost function). Let $c(s) = \alpha(q)s$, where $\alpha(q)$ is a function relating a

signaler's quality, q, to its cost of signaling. For analytical tractability, we assume that signaling

cost decreases linearly with quality and that $0 \leq q \leq 1$ ($q = 0$ are the lowest quality individuals

and $q = 1$ are the highest quality individuals). Therefore,

172
$$c(s) = [ac_0 + a(1-q)]s , \tag{1}$$

173   where a is the rate at which the slope of the cost function decreases with quality, and c_0

174   determines the minimum possible cost of signaling, ac_0. Following the graphical representation

175   of Johnstone (1997), we assume that signaler benefit is a saturating function of signal intensity

176   such that

177
$$b(s) = \frac{rs}{1+ds} , \tag{2}$$

where r is the rate that benefit increases with signal intensity and d is the degree of saturation

with signal intensity (plots of equations [1] and [2] can be seen in Figure 1A). More

mechanistically, the benefit function can be thought of as a function of the receivers' reaction

norm, which is not explicitly modeled here but the same qualitative results should apply for any

continuous, monotonically increasing benefit function.

Maximizing the difference between equation (1) and (2) gives the equilibrium signal

intensity

$$s^* = \frac{\sqrt{\frac{r}{a(1 + c_0 - q)}} - 1}{d}$$

(3)

(we provide a derivation in the appendix, sec. A1). This equation shows that higher quality

signalers will have greater signal intensities than lower quality ones (Figure 1B). In other words,

signal intensity is a reliable indicator of quality. Figure 1A demonstrates this pattern for two

different qualities, showing that, for high-quality individuals, the maximum net benefit is at a

higher signal intensity than for low-quality individuals. With greater values of c_0, the

relationship between signaler quality and signal intensity becomes more linear and less steep

(Figure 1B). For simplicity, in all other figures, we assume that $c_0 = 0$.

The results of this model are in line with those of Grafen (1990*a*, 1990*b*) and Johnstone

(1997). This modeling approach also provides a useful framework that can be adapted to

analytically explore animal signaling in different biological scenarios, such as threshold

assessment.

Figure 1. The evolution of reliable signals in a model with continuous assessment. A)

Relationship between signal intensity and fitness costs or benefits for signalers of low ($q = 0.2$)

and high ($q = 0.6$) quality, given $c_0 = 0$. Arrows denote the equilibrium signal intensity, s^*, for

low- and high-quality individuals, which occurs where the difference between benefit and cost is

the greatest. B) Relationship between signaler quality and equilibrium signal intensity given by

equation (3). Three different values for c_0 are shown. Note that higher quality individuals signal

at greater signal intensities and that panel A closely matches Johnstone's graphical model (1997,

Figure 7.2). In both panels, $a = 0.5$, $r = 4$, and $d = 4$.

Threshold Assessment Model

Fixed Threshold

We next adapt the above model so that receivers respond to signal variation in a binary

fashion (i.e., a threshold response). That is, if a signal is below a threshold value T, the receiver

assesses the signal as low-quality and if the signal is above T, the receiver assesses the signal as

high-quality. Because a signaler's benefit depends on the receiver's assessment, the benefit of

signaling at a given intensity can be thought of as the proportion of the receiver population that

assesses the signal as high-quality. For example, in a mating context where males are signaling

to females, a male's benefit of signaling would be the proportion of females in the population

that assess it as a high-quality mate and therefore mate with it. Alternatively, in a predator-prey

context, the signaler's (prey's) benefit is the proportion of predators that identify the prey as not

worth pursuing, e.g., because the prey is too toxic or too able to escape. We assume that signal

intensities at or above the receiver threshold gain maximum benefits, while signal intensities

below the threshold gain no benefit. Therefore,

221
$$b(s) \begin{cases} 0 & \text{if } s < T, \\ 1 & \text{if } s \geq T, \end{cases} \tag{4}$$

where 1 is the maximum benefit at which every receiver assesses the signaler as high-quality.

In this model, there are only two equilibrium signal intensities: the baseline signal

intensity 0 and the threshold value T. This is an example of an all-or-nothing signaling system

and thus our model agrees with that of Broom and Ruxton (2011). If the costs to an individual of

signaling above the threshold are so high that there is never a net benefit of signaling, that

individual will signal at the baseline signal intensity of 0. In all other cases, individuals signal

exactly at the threshold value (Figure 2A,B). This makes intuitive sense: there is no benefit to

signaling below the threshold value because these signals are all assessed as low-quality, but

there is also no benefit of signaling any higher than exactly at the threshold value and accruing

additional signaling costs. The result of this effect is that signalers on either side of the threshold

will have identical signal intensities of either zero below the threshold or at the threshold value if

above the threshold (Figure 2B). In fact, if we assume that for all qualities q there is some signal

intensity s where $b(s) > c(s)$, all individuals will signal exactly at the threshold value as is shown

in Figure 2C,D. Furthermore, because individuals signaling at an intensity of 0 are never

assessed as high-quality, a signal intensity of 0 would only be evolutionarily stable if being

assessed as high-quality was not strictly necessary for reproduction. Otherwise, all signalers

would signal at the threshold value, irrespective of costs. When all signalers signal at the

threshold value (e.g., Figure 2D), there is no information provided by the signal and thus, if

receivers' threshold responses are not ecologically, physiologically, or genetically constrained,

we would expect receivers to evolve to ignore the signals all together (an outcome conceptually

similar to the lek paradox; Borgia 1979; Tomkins et al. 2004; Kotiaho et al. 2008).

We focus on the case of a receiver with a binary response, but it is worth noting that the

results of this model can be generalized for categorical responses in which receivers have

multiple possible assessment categories (e.g., lowest-quality, low-quality, high-quality, highest-

quality). As with the binary case, the number of equilibria signal intensities will be equal to the

number of assessment categories.

Figure 2. In a model with strict threshold assessment, signalers either evolve 'all-or-nothing'

signals (A and B) or all signalers signal at the same value (C and D). A and C show graphical

depictions of the relationship between signal intensity and fitness costs or benefits for signalers

of low ($q = 0.2$) and high ($q = 0.8$) quality. The dashed gray lines indicate the receiver threshold

value. Arrows denote the equilibrium signal intensity, s^*, for low- and high-quality individuals,

which occurs where the difference between benefit and cost is the greatest. Note that in C the

equilibrium signal intensity is the same for low- and high-quality individuals. B and D show the

relationship between signaler quality and signal intensity. In A and B, $a = 2.0$ and in C and D, a

$= 0.5$. In all panels, $T = 1$ and $c_0 = 0$.

Variable threshold

The model above assumes a fixed threshold value for all receivers, but it is likely that

threshold values will vary among receivers. For example, the sample of female crickets tested by

Beckers & Wagner (2011) showed a threshold of chirp rate for mate choice decisions, but there

was also within-sample variance around this threshold value. Similarly, predators may vary in

the threshold value of an aposematic signal required to induce avoidance (for examples see

Endler and Mappes 2004). To model a variable threshold, we now assume that the threshold

values of the receiver population are normally distributed with mean $\bar{T}$ and variance σ^2. Note

that this convention implies that receivers can have negative threshold values: if a receiver's

threshold value is less than or equal to 0, it evaluates all signalers as high-quality because the

minimum signal intensity is 0. Once again assuming that the benefit of the signal is the

proportion of the receiver population that assesses the signal as high quality, it follows that

272
$$b(s) = \int_{-\infty}^{s} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left[\frac{-(T-\bar{T})^2}{2\sigma^2}\right] dT \, , \tag{5}$$

273   which simply gives that the benefit of signaling at intensity *s* is the area less than *s* under a

274   normal distribution with mean $\bar{T}$ and variance σ^2. For example, if a signal intensity of *s* is

275   assessed as high-quality by 20% of the receiver population, $b(s) = 0.2$.

276   Given equation (5) and assuming that costs still follow equation (1), the equilibrium

277   signal intensity is

278
$$s^* = \bar{T} + \sqrt{-2\sigma^2 \ln\left(a(1+c_0-q)\sqrt{2\pi\sigma^2}\right)} \, . \tag{6}$$

As with equation (3), this equation provides the equilibrium signaling intensity for different

qualities of the signalers. However, if $b(0) > s^*$ or if equation (6) is undefined, which occurs if

$a(1 + c_0 - q)\sqrt{2\pi\sigma^2} \geq 1$, the equilibrium signaling value is instead 0 (see the appendix, sec. A1,

for more details and for the derivation of equation [6]).

Equation (6) shows that incorporating variance into the receivers' threshold values

restores the reliability of the signaling system such that signalers of different qualities receive

maximum net benefits at different signal intensities (Figure 3A). Furthermore, with greater

variance, signal intensity increases faster with increasing quality. That is, for signalers of the

same difference in quality, increasing variance in receiver thresholds leads to a larger difference

in signal intensity (Figure 3B–D).

After introducing inter-individual variation in the threshold value, the only case in which

signalers of the different qualities signal at exactly the same value is when they are signaling at

the baseline value 0 (e.g., Figure 3A, $\sigma^2 = 0.4$ at $q < 0.028$). Because we assume that threshold

values follow a normal distribution, there is always at least a small proportion of the receiver

population that have a threshold value less than 0 and thus assess all signalers as high-quality.

Therefore, there is always some benefit to signaling at the baseline value, and this benefit

increases with increased variance in threshold values because more of the receiver population

will have a threshold value less than 0.

Interestingly, in this model, the equilibrium signaling value is never between 0 and $\overline{T}$

(i.e., equation [6] is always greater than $\overline{T}$ or undefined). In other words, individuals either

evolve to signal at the baseline value of 0 or to signal at some intensity above the mean threshold

value $\overline{T}$. This is surprising because, in this model, there is a benefit to signaling between 0 and

the mean threshold value. However, further exploration reveals that the net benefit is always

greater to signal either above $\overline{T}$ or at 0. This is due to the sigmoidal shape of the benefit

function: below $\overline{T}$, the benefit function curves towards lower values, but above $\overline{T}$ it curves

towards higher values (Figure 3B–D). The sigmoidal shape of the benefit function emerges as a

result of the normal distribution of the receiver variance. Because costs are linear, this sigmoidal

shape results in the net benefit being greater at values above $\overline{T}$ than below it. If the variation in

receiver thresholds has a different distribution, however, it is possible for an equilibrium

signaling value to fall between 0 and $\overline{T}$ (see the appendix [sec. A3, fig. A1], for an example

assuming a gamma distribution).

Figure 3. Variation in threshold values restores reliability of signaling systems. A) Relationship

between quality and equilibrium signal intensity given by equation (6). Dashed black line

denotes the mean threshold value. Colored curves indicate different relationships for different

values of variation in threshold value (σ^2). B–C) Relationship between signal intensity and

fitness costs or benefits for signalers of low ($q = 0.2$) and high ($q = 0.6$) quality for three different

degrees of variation in the threshold value (σ^2). Gray distributions represent the distributions of

threshold values in the receiver population. Arrows denote the equilibrium signal intensity, s^*,

for low- and high-quality individuals, which occurs where the difference between benefit and

cost is the greatest. Parameters: $\overline{T} = 1$, $a = 0.5$, and $c_0 = 0$.

Individual-based Simulations

The analytical models above make many simplifying assumptions (e.g., ignoring genetics

and assuming infinite population sizes). To assess the robustness of our results and to explore

further ecological complexity, we developed stochastic individual-based simulations. These

simulations track individuals' genotypes and phenotypes and model the evolution of signaling

behavior. Unlike in the analytical models, these simulations actually model adaptation via

changes in allelic values rather than simply maximizing the net benefit. For these simulations we

chose to model the evolution of a mating signal in a sexually reproducing species with two sexes

(signalers and receivers). In principle, this same modeling framework could be adapted to fit

other types of signaling systems (e.g., intraspecific competition or predator-prey interactions).

Methods

Each run of the simulation modeled a population in which each individual was either a

signaler or a receiver. Each signaler had a quality q (ranging from 0 to 1), which determined its

cost of signaling. Each receiver had a threshold T, which was used to evaluate signalers (details

below). Generations were overlapping and population size was regulated by the number of

mating sites available K. The simulation was broken up into time steps, which we will refer to as

years. There was an annual sequence of events which happened in the following order: mortality,

filling of mating sites, and then reproduction (including signal assessment and mating, Figure 4).

Mortality occurred at the start of each year independent of age or reproductive status. For

signalers, quality and signal intensity affected mortality such that $M = [ac_0 + a(1-q)]s + m_0$,

where m_0 is the baseline mortality, a is the rate at which the slope of the cost function decreases

with quality, c_0 determines the minimum cost of signaling, and s is the signal intensity.

Receivers died with a fixed mortality of $2m_0$, which was higher than the baseline signaler

mortality in order to keep sex ratios relatively balanced.

Receivers occupied mating sites, from which they evaluated the signals of signalers.

There were only K mating sites available each year. If there were more than K receivers in the

population, K receivers were randomly selected (without replacement) to occupy mating sites

that year. Unselected receivers did not mate that year. We assumed that there was no spatial

structure to the population such that all individuals had the same probability of arriving at any

site and all sites became unoccupied at the end of the year.

Receivers selected signalers as mates by evaluating their signal intensity, s. Receivers had

threshold assessment of signals such that if a signaler's signal intensity was above the receiver's

threshold value T, the receiver evaluated the signaler as a high-quality mate and if the signal

intensity was below T, the receiver evaluated the signaler as low-quality. Receivers mated with

the first signaler they evaluated as high-quality and did not mate again that year. Each year,

receivers evaluated up to N signalers. If by that point no signalers were evaluated as high-quality,

the receiver mated with the final signaler that it evaluated regardless of quality (i.e., threshold

with last-chance option; Janetos 1980). The threshold value of each receiver was randomly

assigned at birth (regardless of genotype) from a normal distribution with mean $\overline{T}$ and variance

σ^2. Each mated pair produced B offspring per year and all offspring survived with a probability

0.5.

Each signaler had a trait, $z(q)$, which determined its signal intensity as a function of its

quality. This trait was genetically determined by 11 additive diploid loci, each with infinitely

many allelic values. Each locus determined the signaling value when the signaler had a specific

quality (broken up by increments of 0.1). That is, locus 1 determined the signal intensity when

the signaler's quality was 0.0, locus 2 determined the signal intensity when the signaler's quality

was 0.1, and so on. This implementation allows signal intensity to be almost any function of

quality. The phenotypic values of signal intensity for a given quality were determined by adding

together the genotypic values of each haplotype at the respective locus plus a random

environmental component drawn from a zero-mean normal distribution with variance e^2. If an

individual's genotypic value was less than 0, its phenotype was assigned to be 0. An individual's

phenotype (signal intensity) was determined at birth and did not change throughout its lifetime.

Note that, because q is a continuous quantity, but the trait $z(q)$ is in increments of 0.1, signalers

evaluated their quality by rounding to the nearest increment of 0.1. Therefore, signalers could not

perfectly evaluate their own quality, which is a realistic assumption for many organisms (e.g.,

Percival and Moore 2010).

At birth, each offspring was randomly assigned to either the signaling sex (signalers) or

the receiving sex (receivers) and each signaler was randomly (uniform distribution, 0–1)

assigned a quality q. Each haplotype of the offspring was assigned by randomly selecting one of

the alleles at each locus of the respective parent (i.e., independent recombination). During birth,

a mutation occurred on each locus with probability μ. If a mutation occurred, a random value

drawn from a zero-mean normal distribution with variance ρ^2 was added to the value at that

locus. Receivers were only carriers for this trait, and it did not affect their preference.

Each run of the simulation was initiated with $2K$ individuals whose genotypes were

randomly assigned from a normal distribution with mean $\bar{T}$ and variance 0.2 (sex, quality, and

signal were randomly assigned in the same manner as they were assigned at birth). To allow

populations to reach selection-mutation-drift balance, there was a 10,000-year "burn-in" period

at the beginning of the simulation. We used two different methods for this burn-in period. In the

first, receivers randomly selected mates independent of signal and there was no cost of signaling.

This simply allowed the population to reach mutation-drift balance so that the initially assigned

genotypes did not affect results. In the second method, receivers assessed signals in a continuous

manner, where the probability of mating with a signaler was $[(rs)/(1+ds)]+0.05$, where 0.05 is

the baseline probability of mating and r and d are the same as in equation (2). This latter method

models how signals evolve if receivers evolve threshold assessment after previously having had

continuous assessment. In both cases, following the 10,000-year burn-in period, simulations then

ran for 20,000 years with threshold assessment. Results were qualitatively similar using both

methods for the 10,000-year burn-in period. For simplicity, all presented results are for the case

in which receivers responded in a continuous fashion during the burn-in period because there

was less variation among runs of those simulations (see figure A2 for results of simulations with

random mating during the burn-in period).

Figure 4. Life-history diagram for the individual-based simulations. The star indicates the start

of the annual cycle. Silhouettes of fiddler crabs (genus *Uca*) are included as visual examples for

signalers and receivers in a mating context.

Results

The results of our individual-based simulations matched the qualitative predictions of our

analytical model. As predicted by the analytical model, with little variance in threshold values,

signalers evolved to signal at relatively similar values whether they were low- or high-quality;

however, with greater variance, the difference between signal values became greater (Figure 5).

411   There was greater variance in the signal intensity of low-quality individuals compared

412 with high-quality ones both within and among runs of the simulation (among run variation

413 shown in Figure 5). This is likely because there was more genetic drift at the loci associated with

414 signaling from low-quality individuals. Because low-quality individuals had higher mortality

415 rates by definition, fewer survived until mate selection and thus the stochasticity of mortality had

416 a greater influence on which low-quality individuals reproduced than it did for individuals of

417 higher quality. Although low-quality individuals signaled at lower intensity than high-quality

418 ones on average, in some runs of the simulation, genetic drift led to low-quality individuals

419 signaling at unexpectedly high signal intensities.

Figure 5. The relationship between signaler quality and signal intensity in the individual-based

422 simulations. Compare to the analytical model results shown in Figure 3A. The dashed black line

423 denotes the mean threshold value. Shaded areas represent the mean of 10 runs of the simulation

424 plus or minus one standard deviation. Different colored shading represents different levels of

425 threshold variation (σ^2). Negative genotypic values were interpreted as zeros, as that is how

they affected phenotype. Parameters: $\bar{T}=1$, $a=0.5$, $c_0=0$, $K=300$, $u=0.25$, $\rho^2=0.01$,

$e^2=0.0001$, $B=5$, $N=10$.

The individual-based simulations provide additional information about the phenotypic

distribution of signalers in the population that cannot be gained by our analytical methods

(Figure 6). These distributions show that even if the mean genotype of the population was to

always signal above the threshold value (as was the case for $\sigma^2=0.004$ and $\sigma^2=0.04$ in Figure

5), there were still individuals in the populations with signal intensities below the threshold

value. In other words, as long as there is variation in threshold values, signal intensities can vary

along a continuum (Figure 6). Furthermore, the greater the variation in threshold values, the

more uniform the distribution of signal intensities (Figure 6). For all simulation parameters, there

was a peak in the number of individuals signaling at an intensity of 0. This is because 0 is the

minimum signal intensity and thus the tail of the phenotypic distribution that would be below 0

was all clustered into that phenotypic value.

One major difference between our analytical model and individual-based simulations is

that the simulations did not show the sudden change from signaling at the baseline value to

signaling above the threshold value that was seen the analytical models (compare $\sigma^2=0.4$ in

Figure 3A and Figure 5). This disagreement can be explained by considering the ecological

assumptions of our models. Recall in the analytical model that there was always some benefit to

signaling at the baseline value because threshold values were normally distributed and thus some

proportion of the receiver population assessed all signal intensities as high-quality. In essence,

this logic assumes that the population is infinitely large, which was obviously not the case in the

individual-based simulations.

However, in our individual-based simulations, a benefit of signaling at the baseline value

emerged in another way, which was by receivers mating with the final signaler that they

evaluated regardless of its quality (i.e., the last-chance option). This provided signalers some

chance of mating, even if their signal intensity was below the threshold value of all receivers.

The lower the maximum number of signalers a receiver evaluated N, the greater this baseline

benefit, because there was an increased chance that any given signaler was the final one a

receiver evaluated. Reducing the value of N in the simulations resulted in sudden changes in

signal intensities similar to those seen in the analytical model (Figure 7). Furthermore, the lower

the value of N, the higher the quality at which this sudden change occurred. This pattern is

intuitive because with a greater baseline benefit, populations should evolve to only signal above

the baseline value when there is a larger net-benefit, which occurs at high qualities. Indeed, in

the extreme case of $N = 1$, which would amount to random mating, we would expect all

individuals to signal at the baseline value and thus sustain the minimum signaling costs.

We also restructured simulations so that receivers did not mate with the last mate that

they evaluated and instead did not mate at all if none of the N signalers evaluated had a signal

intensity above their threshold value (i.e., fixed-threshold without a last-chance option; Roff

2015). In these simulations, mean signaling values below the threshold value were less likely to

evolve (Figure A3). This is logical, because as with increasing the value of N in the simulations

with a last-chance option, this scenario decreases the benefit of signaling at the baseline value.

The only way a signaler can mate is if its signal intensity is above the threshold value of a

receiver.

We also ran simulations in which variation was incorporated into the signalers instead of

the receivers. Biologically, this occurs when a signal (a phenotype) is influenced by local

environment conditions (i.e., phenotypic plasticity). In these simulations, all receivers had the

same threshold value and we incorporated variation into the signalers by increasing the random

environmental component of the phenotype e^2, which is equivalent to decreasing the heritability

of the signaling trait. As with variation in threshold values, in these simulations, signals once

again evolved so that higher quality individuals signaled at greater signal intensities than lower

quality ones, and the strength of this pattern increased with more variation in the random

environmental component (Figure A4). This result occurs because there is selection for high-

quality individuals to signal farther above the threshold value to ensure that their phenotype (and

their offspring's phenotypes) are above the threshold value regardless of the environmental

effect. As above, in these simulations, receivers mated with the last signaler evaluated, so there

was some benefit for low-quality individuals to signal below the threshold value because they

could reduce the costs of signaling and still have a chance of being the last signaler a receiver

evaluated.

Finally, we ran simulations in which we allowed the threshold values of receivers to

evolve, relaxing our assumption that the mean threshold value of the receiver population was

fixed. A full description of the methods for these simulations is provided in the appendix, sec.

A4. Briefly, these simulations assumed that (1) a receiver's threshold value was determined by a

single-locus quantitative trait and (2) receivers had greater fecundity if they mated with a higher

quality signaler. Genetic variation in receiver thresholds was introduced via mutation, and mean

population threshold values evolved to an optimum value (Figure A5). Additional variation in

threshold values was introduced as variance in the environmental component of the threshold

phenotype. As in our other models, these simulations showed that increased environmental

variance in threshold values leads to a greater difference between the signaling values of low-

and high-quality signalers (Figure A6), thus demonstrating that the qualitative results of our

previous models can still hold if receivers' threshold and signalers' signals are allowed to

coevolve.

Figure 6. The phenotypic distribution of signalers in the individual-based simulations. Each

panel shows the distributions for the cumulative number of individuals in 5 runs of the

simulation. The solid vertical line indicates the mean threshold value of the receiver population.

Parameters: $\bar{T} = 1$, $K = 300$, $u = 0.25$, $\rho^2 = 0.01$, $e^2 = 0.0001$, $B = 5$, $N = 10$.

Figure 7. Results of the individual-based simulations for different values for the maximum

number of mates a receiver could evaluate N. If a receiver did not encounter a signaler with a

signal intensity above the receiver's threshold, it mated with the final signaler it evaluated

regardless of signal intensity. Each line shows the mean genotypic values for 10 runs of the

simulation. Parameters: $\bar{T} = 1$, $\sigma^2 = 0.004$, $K = 300$, $u = 0.25$, $\rho^2 = 0.01$, $e^2 = 0.0001$, $B = 5$.

Discussion:

Most previous models of signal evolution have assumed that receivers assess continuous

variation in a signals in a continuous manner (e.g., Grafen 1990*a*, 1990*b*; Johnstone 1994). In

many species, however, receivers exhibit threshold responses in which receivers respond to

continuously varying signals in an binary fashion (e.g., Masataka 1983; Zuk et al. 1990; Reid

and Stamps 1997; Stoltz and Andrade 2010; Beckers and Wagner 2011; Robinson et al. 2011).

We have shown that (1) invariant threshold assessment of signals leads to all-or-nothing signals

(Broom and Ruxton 2011) or a breakdown of reliable signaling systems, but (2) reliable

signaling can be restored if variation is introduced either in the value of the threshold boundary

among receivers or in the translation of genotype to phenotype among signalers. In addition, (3)

when reliable signaling evolves, the mean signal intensity of signalers will typically be above the

mean threshold value of receivers, but (4) the population of signalers will still show a continuum

of signal intensities from well below to well above the population mean threshold.

Our models emphasize the importance of variation among receivers in maintaining

reliable signaling systems. In natural populations, it seems highly likely that threshold values

will vary among receivers to some degree because local environmental conditions affect

phenotypes. Indeed, in a mating context, there is often considerable variation in mate choice

among receivers due to genetic, developmental, or environmental differences (Jennions and

Petrie 1997). In animals exhibiting categorical perception in particular, there is evidence of

variation in category boundaries (Caves et al. 2018; Zipple et al. 2019; Caves et al. *in review*) as

well as context-dependence of categorical boundaries (Lachlan and Nowicki 2015). Future

empirical work should quantify the degree of variation among receivers' threshold values and

more directly investigate its effect on signal evolution. Empirical work could also test

conclusions (3) and (4) above by comparing the distribution of signal intensities and threshold

values in threshold signaling systems.

536   Although our models explicitly considered signals whose reliability is maintained as a

537 result of high- and low-quality signalers facing differential costs of signaling, a similar

538 framework also applies to signals whose reliability is maintained as a result of high- and low-

539 need signalers receiving differential benefits of signaling (Johnstone 1997). An example of this

540 latter case would be nestling birds begging for food from their parents. In this scenario, the

541 benefit of signaling (begging) increases more rapidly for high-need nestlings than low-need ones.

542 As is the case for models with continuous assessment, the same general conclusions emerge from

543 our threshold assessment models whether we assume signalers have differential costs or benefits

544 (see the appendix [sec. A2, fig. A7 and fig. A8] for differential benefits results).

545   Similarly, while we have only discussed among-receiver variation in threshold values

546 (i.e., inter-receiver variation), the same general patterns should hold if individual receivers vary

547 in their threshold values over time (intra-receiver variation). This intra-receiver variation might

548 emerge from changes during development or changes in environmental conditions, but it could

549 also be a result of variation in an individual receiver's ability to detect a threshold boundary.

550 That is, a receiver might mistakenly assess a signal as above its threshold when in actuality it

551 was not or *vice versa*. Our analytical model treats all these types of variation identically;

552 therefore, it predicts that either inter- or intra-receiver variation in threshold values will maintain

553 signal reliability. This suggests that while perceptual errors by the receiver decrease the number

554 of equilibrium signaling values in models with continuous assessment (Johnstone and Grafen

555 1992), in models with threshold assessment, perceptual errors might in fact *increase* the number

556 of equilibrium signaling values. Our individual-based simulations only explicitly modeled inter-

557 receiver variation, however. It would be useful for future studies to explore intra-receiver

558 variation more explicitly.

Our analytical models focus on pure fixed threshold assessment, in which individual

receivers evaluate any signal above their threshold value as high-quality and below their

threshold as low-quality. However, there are different variants of threshold assessment in which

receivers responses follow a general threshold rule but are context-dependent (reviewed by Roff

2015). For instance, there is evidence that female variable field crickets, *Gryllus lineaticeps*,

evaluate mates using a threshold strategy with a last-chance option in which females use a

threshold rule, but if a female has evaluated N males and none of them were above the threshold

value, it mates with the final male encountered (Janetos 1980; Beckers and Wagner 2011). The

results of our individual-based simulations show that while our general conclusions about signal

evolution hold whether receivers follow as a pure fixed threshold strategy or have a last-chance

option, the specific relationship between signaler quality and signal intensity will differ

depending on the strategy. Moreover, with the last-chance option, this relationship depends on

the number of signalers evaluated before mating with the final signaler. We have not evaluated

other context-dependent threshold strategies (Roff 2015), but it is likely that these will also alter

the specific relationship between signaler quality and signal intensity.

In addition, we have only considered the evolution of signals under relatively simple

ecological scenarios in which reliable signaling is maintained by differential signaling costs (or

benefits), but in more complicated scenarios, reliable signals can evolve even without immediate

signaling costs. For example, reliable signaling can evolve without signaling costs when there

are repeated interactions (Rich and Zollman 2016). Similarly, reliable signaling might evolve via

kin selection when individuals are signaling among close relatives (e.g., begging in birds; Caro et

al. 2016). Furthermore, our models do not explain the evolution of dishonest signals (in our

models, signals either evolve to be reliable or to contain no information). Dishonest signals (or

"bluffs") can be maintained in otherwise reliable signaling systems by the existence of frequency

dependent selection (Számadó 2017), a factor that is not present in our models.

It is important to recognize that most of our models assume there is a fixed mean

threshold value for the receiver population. If variations in threshold values among receivers are

due to heritable differences, threshold values themselves will be subject to natural selection and

potentially change over time (Janetos 1980; Real 1990; Bleu et al. 2012). We would expect the

mean threshold value to remain stable if threshold values are either physiologically constrained,

as might be the case in some species with categorical perception, or ecological constrained, as

might be the case when the same perceptual machinery is used to assess signals in addition to

being used in some other ecological context (e.g., female guppies, *Poecilia reticulata*, use the

color orange for mate assessment and food detection; Rodd et al. 2002).

The coevolution of threshold and signal does not change the qualitative results of our

models. Our individual-based simulations with coevolution confirmed that the qualitative

conclusions of our models can still hold when selection causes receiver's thresholds to evolve

(Figure A6). However, we only explored a limited set of ecological and genetic assumptions in

those simulations, and additional work is necessary to more fully understand how the

coevolution between signalers' signals and receivers' thresholds affects the reliability of

signaling systems. For instance, in intra-specific signaling systems (e.g., mate choice), there is

often a genetic correlation between receiver preference (here, the threshold value) and signaling

traits, which can lead to emergent evolutionary phenomena such as Fisherian runaway selection

(Fisher 1930; Lande 1981; Kirkpatrick 1982). Future studies should explore how threshold

responses by receivers affects these processes.

References

Baugh, A. T., K. L. Akre, and M. J. Ryan. 2008. Categorical perception of a natural, multivariate signal: Mating call recognition in túngara frogs. PNAS 105:8985–8988.

Beckers, O. M., and W. E. Wagner. 2011. Mate sampling strategy in a field cricket: evidence for a fixed threshold strategy with last chance option. Animal Behaviour 81:519–527.

Bleu, J., C. Bessa-Gomes, and D. Laloi. 2012. Evolution of female choosiness and mating frequency: effects of mating cost, density and sex ratio. Animal Behaviour 83:131–136.

Borgia, G. 1979. Sexual selection and the evolution of mating systems. Pp. 19–80 *in* M. S. Blum and N. A. Blum, eds. Sexual selection and the evolution of mating systems. Academic Press, New York, New York.

Bradbury, J. W., and S. L. Vehrencamp. 2011. Principles of Animal Communication. 2nd ed. Oxford University Press, Sunderland, MA.

Broom, M., and G. D. Ruxton. 2011. Some Mistakes Go Unpunished: The Evolution of "All or Nothing" Signalling. Evolution 65:2743–2749.

Caro, S. M., S. A. West, and A. S. Griffin. 2016. Sibling conflict and dishonest signaling in birds. Proceedings of the National Academy of Sciences 113:13803–13808.

Caves, E. M., P. A. Green, M. N. Zipple, S. Peters, S. Johnsen, and S. Nowicki. 2018. Categorical perception of colour signals in a songbird. Nature 560:365–367.

Endler, J. A., and J. Mappes. 2004. Predator Mixes and the Conspicuousness of Aposematic Signals. The American Naturalist 163:532–547. The University of Chicago Press.

Fisher, Ronald. A. 1930. The genetical theory of natural selection. Oxford: Claredon Press, Oxford, England.

Grafen, A. 1990a. Biological signals as handicaps. Journal of Theoretical Biology 144:517–546.

Grafen, A. 1990b. Sexual selection unhandicapped by the fisher process. Journal of Theoretical Biology 144:473–516.

Grafen, A., and R. A. Johnstone. 1993. Why we need ESS signalling theory. Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences 340:245–250.

Green, P. A., N. C. Brandley, and S. Nowicki. 2020. Categorical perception in animal communication and decision-making. Behav Ecol, doi: 10.1093/beheco/araa004.

Harnad, S. R. (ed). 1987. Categorical perception: the groundwork of cognition. Cambridge University Press, New York.

Janetos, A. C. 1980. Strategies of Female Mate Choice: A Theoretical Analysis. Behavioral Ecology and Sociobiology 7:107–112.

Jennions, M. D., and M. Petrie. 1997. Variation in mate choice and mating preferences: a review of causes and consequences. Biological Reviews 72:283–327.

Johnstone, R. A. 1994. Honest signalling, perceptual error and the evolution of 'all-or-nothing' displays. Proceedings of the Royal Society of London. Series B: Biological Sciences 256:169–175.

Johnstone, R. A. 1997. The evolution of animal signals. Pp. 155–178 *in* J. R. Krebs and N. B. Davies, eds. Behavioural ecology: an Evolutionary Approach. Blackwell Publishing, Malden, MA, USA.

Johnstone, R. A., and A. Grafen. 1992. Error-prone signalling. Proceedings of the Royal Society of London. Series B: Biological Sciences 248:229–233.

Kirkpatrick, M. 1982. Sexual Selection and the Evolution of Female Choice. Evolution 36:1–12.

Kirkpatrick, M. 1986. The Handicap Mechanism of Sexual Selection Does Not Work. The American Naturalist 127:222–240.

Kotiaho, J. S., N. R. LeBas, M. Puurtinen, and J. L. Tomkins. 2008. On the resolution of the lek paradox. Trends in Ecology & Evolution 23:1–3.

Lachmann, M., and C. T. Bergstrom. 1998. Signalling among Relatives: II. Beyond the Tower of Babel. Theoretical Population Biology 54:146–160.

Lande, R. 1981. Models of speciation by sexual selection on polygenic traits. Proceedings of the National Academy of Sciences 78:3721–3725.

Lindström, K., and J. Lundström. 2000. Male greenfinches (*Carduelis chloris*) with brighter ornaments have higher virus infection clearance rate. Behav Ecol Sociobiol 48:44–51.

Masataka, N. 1983. Categorical responses to natural and synthesized alarm calls in Goeldi's monkeys (*Callimico goeldii*). Primates 24:40–51.

Maynard Smith, J. 1976. Sexual selection and the handicap principle. Journal of Theoretical Biology 57:239–242.

Maynard Smith, J., and D. Harper. 2003. Animal Signals. Oxford University Press, New York.

Nelson, D. A., and P. Marler. 1989. Categorical perception of a natural stimulus continuum: birdsong. Science 244:976–978.

Penn, D. J., and S. Számadó. 2020. The Handicap Principle: how an erroneous hypothesis became a scientific principle. Biological Reviews 95:267–290.

Percival, D., and P. Moore. 2010. Shelter size influences self-assessment of size in crayfish, *Orconectes rusticus*: Consequences for agonistic fights. Behaviour 147:103–119.

Proulx, S. R. 2001. Can behavioural constraints alter the stability of signalling equilibria? Proceedings of the Royal Society of London. Series B: Biological Sciences 268:2307–2313.

Real, L. 1990. Search theory and mate choice. i. models of single-sex discrimination. The American Naturalist 136:376–405.

Reid, M. L., and J. A. Stamps. 1997. Female mate choice tactics in a resource-based mating system: field tests of alternative models. The American Naturalist 150:98–121.

Rich, P., and K. J. S. Zollman. 2016. Honesty through repeated interactions. Journal of Theoretical Biology 395:238–244.

Robinson, E. J. H., N. R. Franks, S. Ellis, S. Okuda, and J. A. R. Marshall. 2011. A simple threshold rule is sufficient to explain sophisticated collective decision-making. PLOS ONE 6:e19981.

Rodd, F. H., K. A. Hughes, G. F. Grether, and C. T. Baril. 2002. A possible non-sexual origin of mate preference: are male guppies mimicking fruit? Proceedings of the Royal Society of London. Series B: Biological Sciences 269:475–481.

Roff, D. A. 2015. The evolution of mate choice: a dialogue between theory and experiment. Annals of the New York Academy of Sciences 1360:1–15.

Ryan, M. J. (ed). 2001. Anuran communication. Smithsonian Institution Press, Washington, DC.

Searcy, W. A., and S. Nowicki. 2005. The Evolution of Animal Communication: Reliability and Deception in Signaling Systems. Princeton University Press, Princeton, New Jersey.

Seyfarth, R. M., D. L. Cheney, T. Bergman, J. Fischer, K. Zuberbühler, and K. Hammerschmidt. 2010. The central importance of information in studies of animal communication. Animal Behaviour 80:3–8.

Stoltz, J. A., and M. C. B. Andrade. 2010. Female's courtship threshold allows intruding males to mate with reduced effort. Proc. R. Soc. B 277:585–592.

Svennungsen, T. O., Ø. H. Holen, and O. Leimar. 2011. Inducible Defenses: Continuous
Reaction Norms or Threshold Traits? The American Naturalist 178:397–410. The
University of Chicago Press.
Számadó, S. 2017. When honesty and cheating pay off: the evolution of honest and dishonest
equilibria in a conventional signalling game. BMC Evol Biol 17:270.
Tomkins, J. L., J. Radwan, J. S. Kotiaho, and T. Tregenza. 2004. Genic capture and resolving the
lek paradox. Trends in Ecology & Evolution 19:323–328.
Zahavi, A. 1975. Mate selection—A selection for a handicap. Journal of Theoretical Biology
53:205–214.
Zipple, M. N., E. M. Caves, P. A. Green, S. Peters, S. Johnsen, and S. Nowicki. 2019.
Categorical colour perception occurs in both signalling and non-signalling colour ranges
in a songbird. Proc. R. Soc. B 286:20190524.
Zuk, M., K. Johnson, R. Thornhill, and J. D. Ligon. 1990. Mechanisms of female choice in red
jungle fowl. Evolution 44:477–485.
710