Patterns of speciation are similar across mountainous and lowland regions for a Neotropical plant radiation (Costaceae: *Costus*)

Running Title: Ecogeographic speciation in *Costus*

Oscar M. Vargas[1,2], Brittany Goldston[3], Dena L. Grossenbacher[3], Kathleen M. Kay[1]

[1]Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, Santa Cruz, California 95060

[2]Email: ovargash@ucsc.edu

[3]Department of Biology, California Polytechnic State University, San Luis Obispo, California 93401

Author Contributions: Study was designed by OMV, DLG and KMK. KMK curated the plant collection used for genomic sampling. OMV, DLG and KMK collected the data. Phylogenetic analyses were conducted by OMV. Species richness and topographic complexity analyses were conducted by BG; all other geographic and climatic analyses were conducted by DLG. Manuscript was written jointly by OMV, DLG, and KMK.

Acknowledgments: This research was funded by NSF Dimensions of Biodiversity grant DEB-1737878 to KMK, DLG, Jennifer Funk (Chapman University), Santiago Ramirez (UC Davis), and Carlos Garcia-Robledo (U Conn). We thank the Organization for Tropical Studies,

Tortuguero National Park, Estación Biológica Monteverde, and Estación Tropical La Gamba for facilitating field work in Costa Rica. We also thank the Smithsonian Tropical Research Institute for facilitating field work in Panama. We thank Julia Harenčár, Rossana Maguiña, and Pedro Juarez (UCSC) for help with field collecting, and Edgardo Ortiz for assistance with the analysis of the transcriptomic data. All field research was conducted with appropriate collection permits in Costa Rica (M-P-SINAC-PNI-ACAT-026-2018, ACC-PI-027-2018, M-PC-SINAC-PNI-ACLAP-020-2018, INV-ACOSA-076-18, M-PC-SINAC-PNI-ACTo-020-18, and CONAGEBIO R-056-2019-OT and R-058-2019-OT), Panama (SE/AP-13-19), and Peru (Resolución Jefatural N° 007-2017-SERNANP-RNAM-JEF and Resolución de Dirección General N° 231-2017-SERFOR/DGGSPFFS both granted by Servicio Nacional Forestal y de Fauna Silvestre del Estado Peruano). Additional plant samples were graciously contributed by John Kress, Douglas Schemske, Dave Skinner, the National Tropical Botanical Garden, and the MICH, MO, UC, and US herbaria. We thank Jim Velzy, Sylvie Childress, and Sarah Ashlock for greenhouse care of the living *Costus* collection, Hannah Thacker and Jaycee Favela (UCSC) for help with lab work, and Russell White and Andrew Fricker for help with the topographic complexity analysis.

Data Accessibility Statement: All data and code will be archived at

https://bitbucket.org/oscarvargash/costus_speciation and DRYAD upon acceptance.

Patterns of speciation are similar across mountainous and lowland regions for a Neotropical plant radiation (Costaceae: Costus)

Abstract

High species richness and endemism in tropical mountains are recognized as major contributors to the latitudinal diversity gradient. The processes underlying mountain speciation, however, are largely untested. The formation of steep ecogeographic gradients and the geographic isolation of populations by geological features are predicted to promote speciation in mountains relative to lowland regions. We evaluate these processes in a species-rich Neotropical genus of understory herbs, *Costus*, that range from the lowlands to montane forests. We test for an association between topography and species richness, infer a biogeographical history, and ask whether the contributions of niche divergence, geographical isolation, and pollination shifts differ between mountain-influenced and Amazonian sister pairs inferred from a 756-gene phylogeny. We find that species richness is highest in topographically complex regions and infer that early *Costus* ancestors diverged in Central America during a period of mountain formation in the last 3 My. We find evidence for climatic divergence, geographical isolation, and pollination shifts in general, but do not find significant differences in these factors between mountain-influenced and Amazonian sister pairs. We conclude that despite higher indices of climatic niche and species diversity in the mountains, speciation patterns in *Costus* appear similar across tropical lowland and mountainous regions.

Keywords: Ecological specialization, diversification, geographic isolation, Neotropics, pollination shifts, spiral gingers.

Introduction

Tropical mountains exhibit extreme species richness and endemism, contribute substantially to latitudinal diversity gradients, and are thought to be cradles of recent speciation (Rahbek et al. 2019a, Rahbek et al. 2019b). The Neotropics, for example, contain three of the five most species-rich hotspots for plant diversity worldwide, including Costa Rica-Chocó, Atlantic Brazil, and Tropical Eastern Andes (Barthlott 2005), all of which are associated with mountain ranges. While phylogenetic studies have linked the timing of montane diversifications with mountain building (Luebert and Wigend 2014), the mechanisms by which tropical mountains promote speciation remain unclear.

Mountains are hypothesized to play two major roles in the process of speciation: the generation of ecogeographic gradients and the geographic isolation of populations by geological features. Steep montane and premontane gradients, in factors such as precipitation and temperature, can promote speciation by ecogeographic divergence without sustained allopatry (Gentry 1982; Angert and Schemske 2005; Hughes and Atchison 2015; Pyron et al. 2015). Biotic interactions can also promote ecogeographic specialization (Dobzhansky 1950), and faunal communities change with elevational gradients. For example, hummingbird assemblages turnover rapidly in tropical mountains (Stiles 1981) and often pollinate species-rich montane plant taxa (Gentry 1982; Kay et al. 2005; Lagomarsino et al. 2016). In contrast, geological features can drive allopatric speciation by serving as dispersal barriers even without ecogeographic divergence among populations. If tropical organisms have narrower climatic tolerances than temperate ones, as hypothesized, the effect of geological features on allopatric isolation may be greatly magnified in tropical mountains (Janzen 1967; Ghalambor et al. 2006; Cadena et al. 2012; Guarnizo and Canatella 2013). Both ecogeographic gradients and

topographic dispersal barriers may lead to frequent progenitor-derivative, or budding speciation, in mountain-influenced areas, in which new species arise from a marginal or founder population of an established widespread species. Moreover, it is unknown whether there is a dominant mechanism driving speciation in tropical montane and premontane environments or if, alternatively, these mechanisms work synergistically.

Geographic isolation and ecogeographic adaptation predict that mountain-influenced sister species will be allopatric (Fig. 1D), will differ in climatic niche (Fig. 1E), or both (Fig. 1F). Under a model of allopatric speciation, reproductive isolation in mountains can arise when a species' range is dissected by a newly formed barrier or when individuals disperse to suitable habitat that is geographically isolated by the surrounding topographic complexity (Fig. 1D). Ecogeographic adaptation and specialization in mountains can promote reproductive isolation when a marginal population adapts to novel climate conditions at the upper or lower elevation boundary of a species' range, at the cost of adaptation to climatic conditions in the remainder of the species' range (Fig. 1E) (Angert et al. 2008). If biotic communities also vary with climate gradients, ecogeographic specialization in mountains may prominently feature adaptation to other organisms, such as pollinators, that contribute to reproductive isolation. It is also possible that geographic isolation and ecological specialization work together to promote speciation in mountains, with geographically isolated populations adapting to new climate niches (Fig. 1F). We propose that these scenarios for speciation (Fig. 1D-F) should be much more common in mountainous regions than speciation without climate niche divergence and/or substantial geographic isolation (Fig. 1C).

In contrast, low climatic complexity in Amazonia predicts that sister pairs there will occupy similar climatic niches and may or may not be allopatric (Figs. 1A-B). Under a model of

allopatric speciation in Amazonia, reproductive isolation can arise when populations are geographically isolated by non-climatic features like rivers and edaphic gradients (Fig 1B) (Fine et al. 2005, Misiewicz and Fine 2014, Weir et al. 2015). Alternatively, reproductive isolation could arise without geographic isolation, again due to pollinator shifts or other biotic drivers; however, we expect pollinator shifts to be less frequent without steep climatic gradients underlying turnover in biotic communities (Fig. 1A). The absence of steep montane and premontane gradients in Amazonia predicts that Amazonian sister pairs will have less overall climatic and geographic divergence than mountain-influenced pairs and be less likely to undergo budding speciation.

Here we test for unique signatures of speciation in mountains relative to lowlands in spiral gingers (*Costus*), a diverse Neotropical clade. *Costus* comprises approximately 59 species of understory herbs found from sea level to cloud forests throughout tropical Central and South America. Neotropical *Costus* are widely interfertile (Kay and Schemske 2008) with stable ploidy (Maas 1972; 1977), and prior studies have suggested a prominent role for prezygotic reproductive barriers, including ecogeographic isolation (Chen and Schemske 2015), differences in pollination syndrome (orchid bee v. hummingbird; Kay and Schemske 2003), and floral divergence within a pollination syndrome (Kay 2006; Chen 2013). Here, we begin by testing whether there is an association between *Costus* species richness and topographical complexity. Then, we infer a multi-locus phylogeny for *Costus* that we use to reconstruct the biogeographic history, the timing of divergence, and the evolution of pollination syndromes in the genus. Finally, we use sister species comparisons to test for differences in speciation processes in montane and Amazonian regions using climatic, geographic, and pollination data. We discuss how our results shed light on speciation in the Neotropics.

FIGURE 1. Testing the role of mountains in speciation. Hypothetical ranges of sister species, as black and red filled circles, overlayed on a landscape where lighter grays indicate higher elevations. Sister species in Amazonia, a region with low topographic complexity and relatively homogeneous climate, are expected to be sympatric (A) or allopatric (B), and isolated by lowland geological features, fine-scale habitat divergence, or biotic interactions. Mountain-influenced sister species may also show these patterns (C), however, we predict that geographic isolation (D), climate niche divergence (E), or both (F) will be more common in mountains than in the lowlands (indicated by bolded boxes).

Material and Methods

Data and control files including commands with all parameters used in our analyses can be accessed at https://bitbucket.org/XXXXXX/XXXXXXX

OCCURRENCE DATA

We downloaded all known occurrence records for the species in our study from the Global Biodiversity Information Facility (GBIF, http://www.gbif.org). We supplemented these with occurrences from our own field studies in 2018 and 2019, as well as additional herbarium (Cornell, MSU, and UC JEPS) and iNaturalist data (https://www.inaturalist.org). All occurrences were then filtered for quality by excluding records without decimal accuracy in latitude and longitude, and with coordinates failing to match the locality description. To avoid potential

taxonomic misidentifications, we retained only occurrences where the identification was made by one of three taxonomic experts: Paul Maas, Dave Skinner, or KMK. We checked species' epithets against the most recently published taxonomies and corrected synonyms and spelling errors. The final filtered dataset included 4,834 unique occurrences for 61 taxa (mean per taxa=79, range=1-593, SD=121).

TOPOGRAPHIC COMPLEXITY AND SPECIES RICHNESS

To quantify topographic complexity across our study region, we used the terrain ruggedness index (TRI) function in ArcGIS (ESRI 2018). TRI is a measure of topographic heterogeneity that takes the sum in elevation change between a focal grid cell and all neighboring grid cells (Riley et al. 1999). To calculate TRI, we used raw elevation data at 1km resolution from earthenv.org/topography and projected it to 1600 and 6400 km² grids. The two grid sizes allowed us to assess whether our results were sensitive to spatial scale. Species richness per grid cell was recorded as the number of unique species occurrences in the filtered occurrence data described above (Point Statistics function in ArcGIS). Richness, TRI values, and X,Y coordinates for each grid cell were extracted across the two spatial scales and saved for downstream analysis (Extract Multi Values to Point function in ArcGIS).

To determine whether species richness is predicted by the terrain ruggedness index (TRI), we used both an ordinary least squares regression model and a spatial autoregressive lag model. Ecological data is typically affected by spatial autocorrelation (SAC), with nearby localities being more similar than expected when random (Kissling and Carl 2008). As a result of species distributions being spatially constructed by nature, our data will likely have residual SAC, breaking an assumption of linear regression. To check for residual SAC between TRI and species richness we used Moran's I coefficient. A Moran's I coefficient near zero indicates no SAC,

while positive and negative values indicate positive or negative autocorrelation (Bhattarai et al. 2004). Spatial autoregressive models are commonly used to mitigate known SAC in the residuals. Accordingly, we used a spatial autoregressive lag model to relate species richness and TRI. Due to the ongoing debate of incorporating spatial autocorrelation into the analysis of species distribution data, we chose to compare our model results to an ordinary least squares regression model which ignores spatial autocorrelation (Dormann 2007). Model selection was accomplished using the Akaike information criterion (AIC).

PHYLOGENETIC ANALYSIS

To infer a well resolved phylogeny we employed targeted sequencing, capturing 853 genes in 113 samples representing 57 species, including outgroups (Table S1) and, when possible, samples from different geographic locations for widespread species and putative progenitor-derivative species pairs. We selected the genes based on six transcriptomes belonging to neotropical *Costus*, five newly sequenced (Table S2) and one published (GenBank BioSample: SAMN00991785). We extracted total RNA using the RNeasy Plant Mini Kit (Qiagen, San Diego, California) from fresh tissue; Poly-A enriched libraries (with an insert size of 300 bp) were prepared by the DNA Technologies Core at the University of California, Davis, with subsequent paired-end 150 bp HiSeq4000 sequencing (Illumina Inc.). We employed SeqyClean v.1.10.07 (Zhbannikov et al. 2017) to remove low quality reads and read tails using default parameters and a cutoff Phred score of 20. Poly A/T tails were also trimmed. Assembly of transcripts was performed with Trinity v2.8.4 (Grabherr et al. 2011). With the transcriptomic dataset, we employed BaitBox (https://github.com/edgardomortiz/baitbox) to select the genes for sequencing. Briefly, BaitBox first used VSearch V.2.10.3 (Edgar 2010) to deduplicate individual transcriptomes (query_cov [overlap] = 0.99, id [similarity] = 0.995) and performed clustering

among the transcripts of all samples (query_cov = 0.75, id = 0.75) outputting a fasta file for every cluster. Clusters were subsequently aligned with MAFFT v7.407 (Katoh and Standley 2013) using the "--auto" mode. Only genes for which a single copy was found in the alignments were used for subsequent subselection, filtering out possible paralogs. Final gene selection for sequencing in the phylogenetic analysis was based on transcript length (len = 720-2400), transcript presence in a minimum of four species (spp= 4,6), transcript presence in a focal species (*Costus pulverulentus*, foc = 1), percentage of gaps in the alignment not exceeding 50% (gap = 50), an average pairwise percentage identity range of 75–99.6% (pid = 75-99.6), and allowing a maximum of 15% of short introns per gene (< 120 bp) (psr = 0.15). Finally, allowing for a G-C content of 30–70% and a tailing percentage overlap of 66.55, BaitBox designed 16,767 baits of 120bp in length for the 853 genes selected.

We extracted DNA from recently collected field and greenhouse samples using NucleoSpin Plant Mini Kit II (Macherey-Nagel, Düren, Germany) according to the manufacturer's protocol, adding 5 uL proteinase K (20 mg/mL) to the digestion step and increasing the digestion incubation time to an hour. For herbarium specimens we used the MagPure Plan DNA LQ kit (Angen Biotech, Guangdong, China). Library preparation and sequencing for the 853 targeted genes was performed by Rapid Genomics (Gainesville, Florida). We employed HybPiper v1.3.1 (Johnson et al. 2016) to assemble the targeted genes, and MAFFT using the "linsi" exhaustive algorithm to align the matrices containing concatenated exons and introns. Problematic sections in the alignments were trimmed with the "-automated1" option of trimAl v1.4.rev22 (Capella-Gutiérrez et al. 2009). Rogue taxa were removed with the "-resoverlap 0.75 -seqoverlap 75" arguments of trimAl.

To filter out tentative paralog genes unidentified by BaitBox, we excluded genes with extreme variation in branch lengths based on the assumption that ingroup branches should not be extremely long considering the recent diversification of Neotropical Costus (Kay et al. 2005). To identify genes with extreme in-group branch length variation, we first inferred trees for each alignment with IQ-Tree v1.6.12 (Nguyen et al. 2015) using a GTR+G model and 1000 ultrafast bootstraps. Then, after removing outgroups with pxrmt (Brown et al. 2017) and outlier sequences with TreeShrink "-q 0.10" (Mai and Mirabab 2018), we calculated the variation in branch lengths using SortaDate (Smith et al. 2018), and then sorted genes accordingly. Visual examination of genes with extreme variation in branch lengths revealed possibly paralogy. Therefore, to be conservative we filtered out genes in the top 10% distribution of branch-length variation, resulting in 756 genes for subsequent phylogenetic analysis. Visual examination of remaining genes after filtering revealed no potential paralog issues.

We used concatenated- and coalescent-based approaches for the inference of species trees. Before concatenation, gene alignments were filtered from outlier sequences flagged previously by TreeShrink in our gene trees. A matrix containing sequences for 756 genes was used to infer a concatenated-based species tree, using IQ-Tree with an independent GTR+G model of sequence evolution for each gene partition and 1000 ultrafast bootstraps. We calculated the number of gene trees supporting a given node in the concatenated topology by employing phyparts (Smith et al. 2015), and results were plotted with phypartspiecharts.py (https://github. com/mossmatters/MJPythonNotebooks). For the coalescent inference, we inferred a species tree based on the 756 IQ-Tree-inferred topologies with ASTRAL v.5.6.3 (Zhang et al. 2017), collapsing branches with less than 90 ultrafast-bootstrap support and removing from each input tree the taxa flagged as having outlier branch lengths.

To determine whether Costus diversification coincided with substantial mountain uplift, and to estimate divergence time for sister species, we calibrated our concatenated topology using fossils and external non-*Costus* Zingiberales sequences (Table S3). The inclusion of non-*Costus* sequences was necessary because of the absence of *Costus* fossils. First, we identified the top 50 most clock-like genes in our dataset using the metrics outputted by SortaDate, considering in order of priority branch-length variance (low variance prefered), root-to-tip length (high length preferred), and topological similarity with the concatenated topology (high similarity preferred). Then, we mapped filtered transcriptomic reads from nine Zingiberales species and two outgroups to the top 50 most clock-like genes using reads2sam2consensus_baits.py (Vargas et al. 2019), which wraps sam2consensus.py (https://github.com/edgardomortiz/sam2consensus). The resulting matrices with *Costus* and non-*Costus* sequences were filtered for those containing all non-*Costus* taxa and reduced by leaving only one sample per monophyletic species. We aligned the 27 resultant gene matrices with MAFFT using the "linsi" algorithm and filtered the alignments for 95% column occupancy with the command "pxclsq" of Phyx (Brown et al. 2017). The filtered 27 alignments were concatenated and analyzed with BEAST v.2.6.1 (Bouckaert et al. 2014) with independent GTR+G models for each gene partition. As a prior, we used a chronogram with relationships fixed based on a Zingiberales phylo-transcriptomic analysis (Carlsen et al. 2018) and our concatenated tree. Branch lengths for the prior tree were calculated with IQ-Tree and later parameterized with TreePL (Smith and O'Meara 2012). Calibrations points were set as follows: 69 Mya (CI = 63–76) to the stem node of the Zingiberaceae based on the fossil *Zingiberopsis magnifolia* (Hickey and Peterson, 1978), and 77 Mya (CI = 69–86) to the crown clade of the Zingiberales based on *Spirematospermum chandlerae* (Friis 1988). BEAST was set to run for 100 m generations sampling every 1 k, and using a birth-death model. We

calculated the chronogram by combining 6 sets of 2500 trees from independent runs with

LogCombiner v.2.6.1 and imputing those in TreeAnnotator v.2.6.1 (Bouckaert et al. 2014) after

checking for chain convergence and a minimum effective sample size of 200 for all parameters

with Tracer 1.7.1 (Rambaut et al 2014). We used FigTree v1.4.4

(https://github.com/rambaut/figtree/releases) and ggtree to produce the tree figures (Yu et al.

2017).

BIOGEOGRAPHIC ANALYSIS

To test whether early ancestors of *Costus* originated in the mountains or lowlands, we

inferred the biogeographic history of the group. We first scored the absence and presence of

extant species in four bioregions, Central America + Choco (C), West Indies (W), Andean (A),

and Amazonian (M), based on our curated occurrence dataset. We then performed an ancestral

range reconstruction using the dispersal-extinction-cladogenesis model (DEC; Ree and Smith

2008), a likelihood version (DIVALIKE) of the dispersal-vicariance model (Ronquist 1997), and

a likelihood implementation (BAYAREALIKE) of the BAYAREA model (Landis 2013) as

implemented in BIOGEOBEARS (Matzke 2013). We abstained from using the founder J

parameter given its caveats (Ree and Sanmartín 2018). BIOGEOBEARS infers ancestral areas

using the aforementioned models and compares them based on likelihoods. Bioregions were

modified from a previous biogeographic study of the Neotropical region (Morrone 2014),

considering the distribution of *Costus*. Our input tree was the chronogram inferred after time

calibration analysis. The most likely reconstruction was selected based on the corrected Akaike

Information Criterion (AICc).

ESTIMATING CLIMATE NICHE

To estimate the climate niche of each species, we obtained four variables representing aspects of temperature and precipitation (http://www.worldclim.org/): mean annual temperature, mean annual precipitation, temperature seasonality, and precipitation seasonality. All data were projected into a South America Albers Equal Area Conic projection and resampled to a 1 km x 1 km grid cell size. Realized niche position of each species was estimated by circumscribing each species' occurrence-based niche relative to all occupied niche space across Neotropical *Costus* using the "PCA-env" ordination technique implemented in the ecospat package (Broennimann et al. 2012; Broennimann et al. 2018). Here, the dimensions of the environmental space for *Costus* were reduced to the first and second axes from a principal components analysis (PCA). The PCA of the four climate variables was constructed using all curated Neotropical *Costus* occurrences, subsampled to one occurrence per grid cell (N=2743 grid cells total). We then created a grid with 100 x 100 PCA unit grid cells and used the species' presence data to project the density of each species into environmental space using a kernal density function (Broennimann et al. 2012). Niche position for each species was estimated as the mean of PC1 and PC2.

To determine whether mountain-influenced taxa occupy a larger volume of climate niche space overall, we tested whether the variation in species' mean niche values differ by region (Amazonian, mountain-influenced). We visualized the evolution of climate niches by projecting the phylogeny on species' mean values for PC1 and PC2 using Phytools (Revell 2012), and we used Levene's tests on PC1 and PC2 separately to compare the variance in mean niche values between regions (leveneTest function, car package in R, Fox and Weisberg 2019). Species were categorized as either Amazonian (all occurrences contained in the Amazon and/or West Indies

bioregions) or mountain-influenced (occurrences fully or partially contained in the Central America + Choco or Andean bioregions)

EVOLUTION OF POLLINATION SYNDROMES

We evaluated whether hummingbird pollination and/or shifts from orchid bee to hummingbird pollination are more prevalent in mountain-influenced than Amazonian taxa. Orchid bee- or hummingbird-pollination syndromes were assigned to taxa based on previous studies in the genus (Maas 1972; Maas 1977; Kay and Schemske 2003; Kay et al. 2005) and KMK expertise. Although only a subset of species have pollination data, pollination syndromes accurately predict whether orchid bees v. hummingbirds are the primary pollinator (Kay and Schemske 2003). Thus, pollination syndromes serve as a tractable proxy for an important biotic interaction that could contribute to ecogeographic divergence. We first used a chi-squared test to determine whether hummingbird pollination is more frequent among mountain-influenced species than Amazonian species, regardless of their phylogenetic history. We then performed a character reconstruction of pollination syndromes on the phylogeny using phytools (Revell 2012) and used another chi-squared test to determine whether reconstructed shifts from bee to hummingbird pollination are more likely in mountain-influenced vs. Amazonian ancestors. Ancestors and species were categorized as stated in the previous paragraph.

IDENTIFYING SISTER TAXA

We identified sister taxa for comparisons of range overlap, climate niche divergence, and pollination shifts in recent and phylogenetically independent speciation events. Sister taxa were identified from the 1000 rapid bootstraps of the concatenated alignment by first pruning the trees to the reduced taxon set used for the time-calibrated phylogeny and then counting the frequency of all sister pairs across bootstrap replicates with a custom R script (R Core team 2020). This

frequency was used as a weighting factor in downstream analyses to account for uncertainty in tree topology. Sister pairs were categorized as either mountain-influenced (one or both species occurred in Central America, Choco, or the Andes) or as Amazonian (both species occurred in the lowland Amazon, or one species in the Amazon and the other in the West Indies). We flagged potential cases of budding speciation in the phylogeny when we observed a widespread species having a taxon nested in it with a smaller range area; we arbitrarily chose a minimum asymmetry ratio of 5 (large / small range) as a cut off.

ESTIMATING SISTER PAIR RANGE OVERLAP, RANGE ASYMMETRY, AND CLIMATIC NICHE DIVERGENCE

For each sister pair, we used the filtered occurrence data to estimate the degree of range overlap using a grid approach. We divided the Neotropics into a series of cells by grid lines that follow degree longitude and latitude using the "raster" R package version 2.9-5 (Hijmans 2016). We calculated range overlap as the summed area of grid cells occupied by both species, divided by the summed area of occupied grid cells for the smaller ranged species. Thus, range overlap could range between 0 (no range overlap) and 1 (the smaller-ranged species is found only within the range of the larger-ranged species) (Barraclough and Vogler 2000; Fitzpatrick and Turelli, 2006). We calculated range size asymmetry as the summed area of grid cells occupied by the larger ranged species divided by the summed area of grid cells for the smaller ranged species (Fitzpatrick and Turrelli 2006). In order to assess whether the ensuing analyses were sensitive to spatial scale, range overlap and size asymmetry was calculated at two cell sizes, 0.05 and 0.1 decimal degrees, representing grid cells of approximately 33 and 131 km^2 respectively (exact value varies by latitude). Sister pairs lacking adequate geographic data (fewer than 4 known

occurrences for one or both species) and those taxonomically poorly understood were excluded from all downstream analyses (Table S4).

COMPARING MOUNTAIN-INFLUENCED AND AMAZONIAN SISTER PAIRS

We performed a series of analyses to determine whether climate divergence, geographic isolation, or speciation mode differs between mountain-influenced and Amazonian sister pairs. We predicted that mountain-influenced pairs would have greater and more variable niche divergence, greater and more variable geographic isolation, and more frequent parapatric and budding speciation than Amazonian pairs. We also predicted that mountain-influenced pairs would have more frequent shifts to hummingbird pollination than Amazonian pairs, but were unable to make this comparison with sister pairs because of the small number of shifts occurring at the tips of the phylogeny.

To compare climate niche divergence between regions, we compared the frequency of climate niche equivalency, the degree of climate niche divergence, and the variance in climate niche divergence for mountain-influenced v. Amazonian sister pairs. To estimate climate niche equivalency, we determined whether each sister pair occupies statistically equivalent niches, i.e., that the niche overlap between sister species is equal to that of two species occupying random niches in the same range of environmental conditions that are available to the species in question (Warren et al. 2008; Broennimann et al. 2012). This was performed using the function ecospat.niche.equivalency.test (ecospat package, Broennimann et al. 2018), whereby the observed overlap is compared to a null distribution of simulated overlaps when randomly reallocating the occurrences of both species among the joint distribution of occurrences. Only pairs where each member species occupied at least 5 grid cells were used in this analysis (N=22 sister pairs). The frequency of climate niche equivalency was compared between regions

(mountain-influenced, Amazonian) with a weighted chi-squared test, weighted by the number of bootstrapped trees containing a given sister pair (function wtd.chisq, weights package, Pasek 2020). To compare the mean and variance in climate niche divergence of sister pairs between regions, we calculated climate niche divergence for each pair as the euclidean distance between mean PC1 and PC2 for each species and then used a two sample T-test and a Levene's test, respectively, each weighted by the number of bootstrapped trees containing a given sister pair (wtd.t.test function, weights package, Pasek 2020; leveneTest function, car package, Fox and Weisberg 2019).

To compare geographic isolation between regions, we first quantified the current range overlap for sister pairs between regions and then examined how range overlap varies with divergence time. To quantify the mean and variance of current range overlap, we used a two sample T-test and a Levene's test, each weighted by the number of bootstrapped trees containing a given sister pair (wtd.t.test function, weights package, Pasek 2020; leveneTest function, car package, Fox and Weisberg 2019). We note that current overlap may differ from overlap at the time of speciation due to post-speciation range expansions, contractions and shifts; however, by using only sister species and comparing regions, we can infer differences in geographic isolation between regions for the most recently diverged species pairs in *Costus*.

To determine whether allopatric or parapatric speciation predominates in the mountains or Amazonian regions, we tested whether sister pair range overlap was predicted by divergence time, region, and their interaction using a linear model (lm function, weighted by the number of bootstrapped trees containing a given sister pair, R). If allopatric speciation is dominant, then more recently-diverged species pairs should be allopatric, whereas older pairs might show range overlap due to range shifts since speciation (Fitzpatrick and Turelli 2006). Contrastingly, if

sympatric/parapatric speciation is dominant, younger sister species pairs should show range overlap whereas older pairs should show a variety of configurations (Fitzpatrick and Turelli 2006; Anacker and Strauss 2014). A significant interaction between divergence time and region would indicate that the predominant geographic mode of speciation differs by region.

Finally, to determine whether budding speciation occurs and whether this phenomenon varies by region, we examined the relationship between range asymmetry and divergence time, and we also looked for evidence of nested phylogenetic relationships that would indicate a small-ranged taxon was derived from within a widespread progenitor taxon. We first tested whether sister pair range asymmetry was predicted by divergence time, region, and their interaction using a generalized linear model with a natural log link function, gamma distribution suitable for left-skewed response variables such as range size asymmetry (glm function, weighted by the number of bootstrapped trees containing a given sister pair). If budding speciation is common, then range size asymmetry is predicted to be greatest for the youngest sister pairs and to decrease on average with time, as ranges undergo expansion or contraction following the initial budding speciation event (Fitzpatrick and Turelli 2006; Grossenbacher et al. 2014). A significant interaction between divergence time and region would indicate that the signature of budding speciation differs by region. Significance of predictors was assessed by likelihood ratio chi-squared tests (LR) using single term deletions. Our phylogenetic sampling also allowed us to assess nested phylogenetic relationships indicative of budding speciation in four putative cases, three mountain-influenced and one Amazonian (*C. scaber* in Central America – *C. ricus, C. pulverulentus* – *C.* sp. nov. 18020/18049, *C. laevis* – *C. wilsonii*, and *C. scaber* in South America – *C. spicatus,* respectively). We did not know in advance which, if any, species were produced

by budding speciation, but we attempted to sample across the known geographic distributions of multiple widespread taxa.

Because we only have a modest set of sister pairs that are used repeatedly in the analyses above, we present exact p-values and describe effect sizes, without using a strict $\alpha = 0.05$ to determine significance.

Results

Topographic complexity and species richness

We find a positive association between topographic complexity and species richness in *Costus*, consistent with mountains playing a key role in diversification of this clade (Fig. 2). We recover a significant spatial autocorrelation between Terrain Ruggedness Index (TRI) and species richness (Fig. 2; Moran's I > 0.12, P < 0.001, across both spatial scales). Ordinary least squares regression shows a significant positive relationship between TRI and species richness (1600 km^2, F = $7.56_{1,973df}$, P = 0.006; 6400 km^2, F = $6.82_{1,597df}$, P = 0.009). Simultaneous Autoregressive (SAR) lag models also shows a positive relationship between TRI and species richness, however, significance varies by scale (1600 km^2, z=1.89, P = 0.138, pseudo-r squared = 0.227; 6400 km^2, z = 1.89, P = 0.049, pseudo-r squared=0.246). SAR lag models are favored over ordinary least squares regression using AIC (1600 km2, Likelihood Ratio = 1282.51, P < 0.001; 6400 km2, Likelihood Ratio = 170.55, P < 0.001). Overall, Costus shows a center of species richness in the Costa Rica-Choco and northern Andean floristic regions, moderate richness in the Guiana Shield and eastern slope of the southern Andes, and relatively low richness in the Amazon basin. It is important to note that our sampling relied on collection efforts carried out by previous collectors and our team, which largely focused on sampling

Central America, and therefore it is possible that Andean and Amazonian *Costus* diversity is underestimated.

FIGURE 2. Species richness mapped onto a grid measuring topographic complexity using the Terrain Ruggedness Index. Warmer colors indicate higher topographic complexity, and circles are scaled to show the number of reported species per grid.

Phylogenetic analysis

We infer a robust species-level phylogeny for *Costus*. Our concatenated matrix has 95.3% cell occupancy comprising 133 samples by 756 genes in 1,474,816 aligned columns (Tables S5–6). Average length per gene, including partial introns, is 1,951 bp. Samples have an average of 728 genes. We calculate two species trees, the first based on a concatenated alignment

and a second based on individual gene trees in a coalescent framework. Both species trees have robust support and similar topologies (Figs. 3,S1). Because of the general low molecular divergence found in the ingroup (average pairwise identity = 90.0%) and the low signal found in individual genes (Fig. S2), we selected the concatenated topology as the best phylogenetic hypothesis for the remainder of this study (Fig. 3). Our *Costus* phylogeny is robust with most nodes presenting full or high support (>=95 ultrafast bootstrap, >=90 Shimodaira-Hasegawa approximate likelihood ratio test) and generally agrees with previous phylogenetic studies in *Costus* (Kay et al 2005, André et al 2016).

TIME-CALIBRATION OF THE PHYLOGENY

Our time-calibrated phylogeny dates the crown clade of Neotropical *Costus* to 3.0 Mya with a 95% CI = 1.50–4.87 (Fig 4). Our matrix for the calculation of the chronogram, which included Zingiberales sequences for each one of its families and a reduced *Costus* sampling, is composed of 69 taxa by 27 clock-like genes, comprising 22,237 aligned columns (Table S7). Our time-estimation for the origin of *Costus* in the Neotropics is consistent with a previous dating of 1.1–5.4 Mya (Kay et al. 2005) but younger than another of ~7 Mya (André et al. 2016).

BIOGEOGRAPHIC ANALYSIS

Ancestral range reconstruction suggests that the Central American region has dominated the biogeographical history of the genus in the Neotropics (Fig 4). Based on AICc, the model that best fits the reconstruction is DIVALIKE (AIC =185.5), followed by DEC (AIC=188.2), and BAYAREALIKE (AIC=230.5). The DIVALIKE reconstruction shows that nearly half of the ancestors in the phylogeny were distributed in Central America (25 out of 54 ancestors with a > 0.90 probability) with the vast majority of the early ancestors estimated as Central American. Colonization out of Central America is inferred to have happened around 1.5 Mya to the Andean

and the Amazon regions. Similar biogeographical patterns are also found in the reconstruction

inferred with the DEC but not BAYAREALIKE models (Figs. S3,S4).

FIGURE 3. Phylogram inferred from the concatenated matrix of 756 genes for *Costus*. Node numbers indicate ultrafast bootstrap (left) and a Shimodaira-Hasegawa approximate likelihood ratio test (right). nodes without values shown are fully supported at 100. Outgroup branch lengths are reduced to save space. Terminal clades are color coded according to the geographic region on the map. Stars indicate species with evidence for budding speciation.

FIGURE 4. Time calibrated phylogeny, DIVALIKE biogeographic historical inference, and ancestral pollination character reconstruction for Neotropical *Costus*. Pie charts represent percent probabilities of areas for the ranges of ancestors. C: Central America and/or Choco, W: West Indies, A: Andes, M: Amazon. Mixed colors indicate combined bioregions. Pollination syndromes are indicated for every tip taxon along with inferred transitions from bee to hummingbird pollination on internal branches. The most recent common ancestor is reconstructed as bee pollinated. Gray horizontal bars represent 95% confidence intervals for node ages.

CLIMATE NICHE OF MOUNTAIN-INFLUENCED AND AMAZONIA SPECIES

Principal component analysis reveals the first two climate niche axes explain 42 and 23% of the variation among all *Costus* occurrences, respectively. PC1 primarily describes variation in mean annual precipitation and seasonality: low values indicate greater precipitation and high values indicate higher seasonality in both temperature and precipitation. PC2 primarily describes variation in mean annual temperature: low values indicate cooler environments. Overall, we found that the variance in species' mean niche values was greater among mountain-influenced than among Amazonian species for PC2 (Fig 5; PC2 F = $10.05^{1,46df}$, *P* = 0.003), but not PC1

(Levene's test: PC1 F $= 0.05^{1,46df}$, $P = 0.824$), indicating that, together, mountain influenced taxa

are occupying a larger temperature niche space.

FIGURE. 5. *Costus* phylogeny projected onto the first two principal components for climatic niche. Species are categorized into mountain-influenced (red) and Amazonian (blue). The variance in PC2, but not PC1, is significantly different between categories (see main text).

EVOLUTION OF POLLINATION SYNDROMES IN MOUNTAINS VERSUS THE AMAZON

The most likely scenario for the evolution of pollination syndromes in our phylogeny involves 11 shifts from orchid bee- to hummingbird-pollination, with seven shifts happening in recent divergence events and four along internal branches of the phylogeny. We find no significant difference in the proportion of hummingbird-pollinated taxa between mountain-influenced and Amazonian taxa (21 out of 39 taxa v. 5 out of 16 taxa; X^2 (1 , N=55) = 1.51, $P = 0.220$; Fig S5A). Similarly, we find no significant difference in the frequency of shifts to

hummingbird pollination in ancestors characterized as mountain-influenced or Amazonian (8 shifts along 81 branches v. 3 shifts along 28 branches; X^2 (NA , N=109) = 0.02, P = 1; Fig S5B).

MOUNTAIN-INFLUENCED AND AMAZONIAN SISTER PAIR COMPARISON

Climate niche divergence does not differ between pair types. Thirty-one sister pairs were inferred from the bootstrap replicates of the phylogenetic analysis, and 24 have enough geographic and taxonomic information for comparative analyses (Table S4, Fig. S6). Of these, 15 pairs are identified as mountain-influenced and 9 pairs as Amazonian. We test for niche equivalency for 22 out of our 24 pairs (2 taxa have less than 5 occurrences) rejecting niche equivalency for 60% of sister pairs (Table S4, Fig. S7). We find no difference in the proportion of mountain-influenced and Amazonian pairs with equivalent niches (X^2 (1 , N=22) = 14.29, P = 0.217). Climatic niche divergence is on average 46% greater for mountain influenced pairs relative to Amazonian pairs; however, this difference is not significant (weighted t-test, t = -1.47$^{20.2df}$, P = 0.157; Figs. 6, 7). Similarly, the variance in climatic niche divergence is 33% greater among mountain influenced relative to Amazonian pairs; however, this difference also is not significant (Levene's test, F = 1.21^{22df}, P = 0.283).

We examine two aspects of speciation mode: the prevalence of parapatric speciation and evidence for budding speciation. We find no support for the hypothesis that geographic mode of speciation (allopatric or paratric) differs in the mountains versus lowlands. Overall, there is generally little range overlap for sister pairs and no evidence that range overlap varies by region. Average range overlap is 0.09 for mountain influenced and Amazonian pairs at the fine spatial scale (weighted t-test, t = -0.56^{22df}, P = 0.584; Levene's test, F = 3.51^{22df}, P = 0.074; Figs. 6, 7). Range overlap is not predicted by divergence time, region, or their interaction (Fig 6; linear model: divergence time F = 0.50^{1,20df}, P = 0.487; region F = 0.78^{1,20df}, P = 0.386; divergence time

by region F = 0.006^{1,20df}, P = 0.937). We note that there are two outlier data points with range

overlap >0.3 (*C. montanus* – *C. barbatus*, *C. ricus* – *C. scaber*), breaking model assumptions.

When those points are removed from the analysis, range overlap is predicted by divergence time,

but not by region or their interaction (divergence time F =6.63^{1,18df}, P = 0.019; region F =

0.44^{1,18df}, P = 0.515; divergence time by region F = 1.35^{1,18df}, P = 0.261). We urge caution

interpreting this result, since there is no biological justification for excluding the two outliers.

We find limited support for a budding model of speciation being more common in the

mountains than lowlands. Range size asymmetry is greater for younger relative to older

mountain-influenced pairs, while the opposite is observed for amazonian pairs (Fig 6; GLM:

divergence time X^2 = 0.039, P = 0.843; region X^2 = 1.747, P = 0.186; divergence time by region

X^2 = 4.34, P = 0.037). In light of multiple comparisons between sister pair types, we treat this

result with caution. Note that we present only range overlap and asymmetry results for the fine

spatial scale above (~33km2). Results at the course spatial scale were qualitatively similar for all

tests (Fig S8). Finally, we assess evidence for nested phylogenetic relationships between four

taxon pairs, three mountain-influenced and one Amazonian (*C. scaber* in Central America – *C.*

ricus, C. pulverulentus – *C.* sp. nov.18020/18049, *C. laevis* – *C. wilsonii*, and *C. scaber* in South

America – *C. spicatus*, respectively). The first two of those pairs show paraphyly consistent with

budding speciation, whereas the latter two are reciprocally monophyletic, although they all show

substantial range asymmetry (Fig. 2, Table S4).

FIGURE. 6. Mountain-influenced and Amazonian sister pair comparisons. A) Climate niche euclidean distance by region. B) Range overlap v. divergence time (Mya) by region with marginal boxplots indicating differences in range overlap by region. C) Range size asymmetry v. divergence time (Mya) by region, with marginal boxplots indicating differences in range size asymmetry by region. Range overlap and asymmetry were calculated on a grid of 0.05 decimal degrees. In scatterplots, every pair is represented by one dot and its size is proportional to its bootstrap support. Regression lines based on predicted values are included for model effects with uncorrected P<0.05. See main text for statistical results.

Discussion

Mountains are associated with exceptionally high plant diversity in the Neotropics, with tree species richness peaking in the forests nestled at the eastern base of the Northern Andes (ter Steege et al. 2010) and some of the fastest known plant radiations on earth occurring in high elevation Neotropical habitats (Drummond et al. 2012; Madriñán et al. 2013, Uribe-Convers and Tank 2015, Lagomarsino et al. 2016; Vargas et al. 2017, Contreras-Ortiz et al. 2018; Morales-Briones et al. 2018). We see a similar pattern in Neotropical *Costus*, with species richness positively correlated with topographic complexity across its geographic range in Central and South America. Despite the important contribution of tropical mountains to the latitudinal diversity gradient, the mechanisms underlying this remain unclear. Gentry (1982) hypothesized that recent mountain uplift in the Andes and southern Central America promoted rapid diversification of plants with short generation times (e.g., herbs, shrubs, vines and epiphytes) by

providing strong ecogeographic gradients in both climate and pollinators, especially

hummingbirds. In contrast, Janzen (1967; reviewed in Sheldon et al. 2018) proposed that the lack

of strong temperature seasonality in the tropics leads to narrow physiological tolerances and

greater potential for allopatric speciation due to topographic dispersal barriers in mountains. The

relative importance of these two mechanisms in driving species richness is unclear—

investigation requires understanding species-level relationships in clades that span both montane

and lowland environments, and until now we have generally been left comparing Amazonian

trees (e.g., Fine et al. 2005; Vargas and Dick 2020) to high elevation shrubs and herbs (e.g.,

Contreras-Ortiz et al. 2017; Vargas and Simpson 2019). *Costus* provides a unique opportunity to

use a species-level phylogeny to examine possible speciation mechanisms in a clade that spans

lowlands to cloud forests, albeit for a single herbaceous life form.

We first examine ecogeographic divergence caused by macroclimatic conditions and

adaptation to different functional groups of pollinators, orchid bees and hummingbirds. We find

that both mountain-influenced and Amazonian pairs experience climate niche divergence (Fig. 6.

Table S4), with only a marginal trend of greater climatic niche divergence in mountain-

influenced pairs, a remarkable result given the steep gradients in climate in tropical mountains.

Nevertheless, montane species occupy a significantly greater amount of climatic niche space,

primarily due to expansion into cooler environments (Fig. 5). Taken together, our results suggest

that climatic divergence occurs in both mountain-influenced and Amazonian pairs, and that

mountain-influenced taxa occupy the greater temperature variation the mountains offer (Rahbek

et al. 2019a). Despite hummingbird pollination being common in mountain-influenced species, it

is not proportionally more common than in Amazonian species. Moreover, we observe similar

proportions of pollination shifts throughout the tree when ancestors are categorized as mountain-

influenced or Amazonian based on our biogeographic reconstruction (Fig. 4,S5). Thus, while hummingbird pollination may be an important driver of diversification in the mountain-influenced pairs, our results show that it is similarly important in lowland Amazonian pairs.

We further predicted that mountain-influenced pairs would have less range overlap than Amazonian pairs if mountains serve as dispersal barriers and cause long-lasting allopatric separation. We find instead that there is generally little range overlap for sister species, regardless of whether they are mountain-influenced or Amazonian (Fig. 6). Although more mountain-influenced than Amazonian pairs have complete allopatric separation (e.g., *C. amazonicus – C. malortieanus*; Fig. 7, right panel), there is no significant difference between regions. This is remarkable given the height and the ruggedness of the Northern Andes and Central American Cordillera, yet may simply reflect the importance of geographic isolation for most, if not all, speciation. Our results contrast with a previous study in *Costus*, which used species distribution models to predict co-occurrence across all nodes in the phylogeny and found extensive sympatry (André et al. 2016). Species distribution models may lead to dramatic overestimates of actual co-occurrence (Guisan and Rahbek 2011), particularly in topographically complex landscapes where dispersal is likely limited. Furthermore, because geographic signatures of speciation erode over time as ranges expand, contract, and shift (Fitzpatrick and Turelli 2006), the use of sister species comparisons, rather than all nodes in the phylogeny is more likely to yield information regarding speciation itself.

Additionally, we examined evidence for budding speciation, which may be common when speciation is driven by ecological specialization across steep environmental gradients or founder effect speciation across topographic dispersal barriers (Anacker and Strauss 2014; Grossenbacher et al. 2014). Indirect evidence of budding speciation could come from range size

asymmetry decreasing over time-since-divergence, since derivative species should start from small marginal populations. We find a weak pattern of this being the case in mountain-influenced, but not Amazonian pairs (Fig. 6C). Further evidence for budding speciation could come from geographically intensive phylogenetic sampling of sister pairs showing that the smaller-ranged species is nested within the widespread, paraphyletic progenitor (Baldwin 2005). While we find support for this pattern in two mountain-influenced pairs, our phylogenetic sampling was not extensive enough to make statistical comparisons with Amazonian taxa. Thus, while budding speciation likely occurs in *Costus*, our results are not sufficient to draw robust conclusions about differences in the frequency of budding speciation between regions.

We see examples of how climatic niche divergence, range overlap and asymmetry, and pollination shifts can be similar between mountain-influenced and Amazonian pairs (Fig. 7). *Costus* laevis is a widespread bee-pollinated lowland species whose range abuts its restricted (but not phylogenetically nested) montane hummingbird-pollinated sister, *C. wilsonii*, in southern Costa Rica. In this case, speciation may be explained by upslope adaptation to a colder drier environment accompanied by a shift to hummingbird pollination in *C. wilsonii*, perhaps the quintessential ecological specialization model of divergence that Gentry (1982) envisioned for Neotropical mountains (Fig. 1E; Fig. 7 center). In contrast, the lowland Amazonian pair *C. erythrophyllus – C. spiralis* shows comparable levels of range overlap, range asymmetry and climate niche divergence, in this case along PC1 (precipitation and seasonality) rather than PC2 (temperature), as well as a shift in pollination syndrome, but without the direct influence of mountains (Fig. 7 left panel). These examples illustrate that multiple ecogeographic factors can promote lineage splitting in both mountains and lowlands.

How can the similarity in patterns of speciation between mountain-influenced and Amazonian pairs be reconciled with the pattern of increased species richness in mountainous regions we see in *Costus*? Our biogeographic reconstructions suggest it may be a result of happenstance and timing—*Costus* likely first established ca. 3 Mya in Central America when the Talamanca Cordillera started to uplift (Driese et al. 2007), and the genus diversified in this region for close to 1.5 My before colonizing the already elevated Andes cordillera (Gregory-Wodzicki 2000) and the Amazon lowlands (Fig. 4). In fact, we find that Amazonian sister pairs are significantly younger than mountain-influenced pairs (Fig. S5A), consistent with the Amazon basin as a site of recent and rapid diversification. In general, *Costus* has had more time to diversify in Central America and northwestern South America than in the Amazon basin, without the need for invoking different modes of speciation.

To what extent are our results likely to apply to other Neotropical plant lineages? The role of mountains in the diversification of *Costus*, which is restricted to < ca. 2000 m, may be different from other montane lineages. Studies of high elevation Andean paramo plant groups have found substantial allopatry (*Espeletia*: Diazgranados and Barber 2017; *Linochilus*: Vargas and Simpson 2019) and ecological divergence (Campanulaceae: Lagomarsino et al. 2016; *Lupinus*: Nevado et al. 2016, *Espeletia*: Cortés et al. 2018). The young geological age of these alpine habitats may spur rapid diversification simply through the opening of new niche space, regardless of being positioned in the tropics (Weir and Schluter 2008). In sharp contrast with the mountains, plant lineages in the Amazonian lowlands can date back to the Paleocene (Dick and Pennington 2019), although some are much younger (e.g., *Ruellia*: Tripp and Tsai 2017). Diversification studies of plants in the Amazon have found a large role for edaphic ecological specialization (*Protium*: Fine et al. 2014), other fine-scale habitat divergence (Gesneriaceae:

Roalson and Roberts 2016), and biotic interactions, (*Pitcairnia:* Palma-Silva et al. 2011, *Ruellia*: Tripp and Tsai 2017). Both abiotic and biotic conditions vary across lowland forests, and speciation may be rapid even without the influence of mountains.

Taken together, our study suggests that ecogeographic differentiation happens similarly in mountainous regions and tropical lowlands. It may be that wet tropical forests have ongoing and often rapid diversification regardless of topography. Although mountains provide a larger overall climate niche landscape (Rahbek et al. 2019a), we find no evidence that the diversification processes are fundamentally different. However, we caution that these results may not apply to tropical alpine clades that are able to colonize geologically young and spatially disjunct ecosystems above treeline, and these clades may contribute disproportionately to the species richness and endemism of tropical mountains (Hughes and Atchinson 2015). Further studies with a similar framework to ours are needed to determine whether our conclusions can be generalized across tropical organisms in montainous and lowland regions. Our study demonstrates the potential of combining species-level phylogenomics with spatial and ecological data to test longstanding hypotheses about diversification in tropical mountains.

FIGURE. 7. Range maps, climate niches, and inflorescence photos of representative sister species pairs. Left: an Amazonian pair. Center and right: mountain influenced pairs. Top row: occurrences in the map. Central row: comparisons between the climate niches of the sister species (PC axes correspond to Fig. 5). Photo credits from left to right: *Costus spiralis* and *C. erythrophyllus* by KMK, *C. laevis* by R. Maguiña*, C. wilsonii* by P. Juarez, *C. malortieanus* by DLG, and *C. amazonicus* by KMK.

Literature

Anacker, B. L., and S. Y. Strauss. 2014. The geography and ecology of plant speciation: Range overlap and niche divergence in sister species. Proc. R. Soc. B Biol. Sci. 281.

André, T., S. Salzman, T. Wendt, and C. Specht. 2016. Speciation dynamics and biogeography of Neotropical spiral gingers (Costaceae). Mol. Phylogenet. Evol. 103:55–63.

Angert, A. L., and D. W. Schemske. 2005. The evolution of species' distributions: reciprocal transplants across the elevation ranges of *Mimulus* Cardinalis and *M*. lewisii. Evolution. 59:1671.

Angert, A. L., H. D. Bradshaw, and D. W. Schemske. 2008. Using experimental evolution to investigate geographic range limits in monkeyflowers. Evolution. 62:2660–2675.

Barthlott, W., J. Mutke, D. Rafiqpoor, G. Kier, and H. Kreft. 2005. Global Centers of Vascular Plant Diversity. Nov. Acta Leopoldina NF 92:61–83.

Baldwin, B. G. 2005. Origin of the serpertine-endemic herb *Layia discoidea* from the widespread *L. glandulosa* (Compositae). Evolution. 59:2437–2479.

Barraclough, T. G., and A. P. Vogler. 2000. Detecting the geographical pattern of speciation from species-level phylogenies. Am. Nat. 155:419–434.

Bhattarai, K. R., O. R. Vetaas, and J. A. Grytnes. 2004. Fern species richness along a central Himalayan elevational gradient, Nepal. J. Biogeogr. 31:389–400.

Bouckaert, R., J. Heled, D. Kühnert, T. Vaughan, C. H. Wu, D. Xie, M. A. Suchard, A. Rambaut, and A. J. Drummond. 2014. BEAST 2: A Software Platform for Bayesian Evolutionary Analysis. PLoS Comput. Biol. 10:1–6.

Broennimann, O., M. C. Fitzpatrick, P. B. Pearman, B. Petitpierre, L. Pellissier, N. G. Yoccoz, W. Thuiller, M. J. Fortin, C. Randin, N. E. Zimmermann, et al. 2012. Measuring ecological niche overlap from occurrence and spatial environmental data. Glob. Ecol. Biogeogr. 21:481–497.

Broennimann, V. D. C., and A. Guisan. 2018. ecospat: Spatial Ecology Miscellaneous Methods. R package version 3.0. https://CRAN.R-project.org/package=ecospat

Brown, J. W., J. F. Walker, and S. A. Smith. 2017. Phyx: Phylogenetic tools for unix. Bioinformatics 33:1886–1888.

Cadena, C. D., K. H. Kozak, J. L. Parra, C. M. Mccain, J. P. Go, C. K. Bowie, A. C. Carnaval, C. Moritz, C. Rahbek, T. E. Roberts, et al. 2012. Latitude, elevational climatic zonation and speciation in New World vertebrates. Proc. R. Soc. B Biol. Sci. 279:194–201.

Capella-Gutiérrez, S., J. M. Silla-Martínez, and T. Gabaldón. 2009. trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972–1973.

Carlsen, M. M., T. Fér, R. Schmickl, J. Leong-Škorničková, M. Newman, and W. J. Kress. 2018. Resolving the rapid plant radiation of early diverging lineages in the tropical Zingiberales: Pushing the limits of genomic data. Mol. Phylogenet. Evol. 128:55–68.

Chen, G. F. 2013. Sexual isolation in two bee-pollinated *Costus* (Costaceae). Plant Reprod. 26:3–16.

Chen, G. F., and D. W. Schemske. 2015. Ecological differentiation and local adaptation in two sister species of Neotropical *Costus* (Costaceae). Ecology 96:440–449.

Contreras-Ortiz, N., G. Atchison, C. Hughes, and S. Madriñán. 2018. Convergent evolution of high elevation plant growth forms and geographically structured variation in Andean *Lupinus* (Leguminosae). Bot. J. Linn. Soc. 187:118–136.

Cortés, A. J., L. N. Garzón, J. B. Valencia, and S. Madriñán. 2018. On the causes of rapid diversification in the páramos: isolation by ecology and genomic divergence in *Espeletia*. Front. Plant Sci. 9.

Diazgranados, M., and J. Barber. 2017. Geography shapes the phylogeny of frailejones (Espeletiinae Cuatrec., Asteraceae): a remarkable example of recent rapid radiation in sky islands. PeerJ 5:e2968.

Dick, C. W., and R. T. Pennington. 2019. History and geography of Neotropical tree diversity. Annu. Rev. Ecol. Evol. Syst. 50:579–301.

Dobzhansky, T. 1950. Evolution in the Tropics. Am. Sci. 38:208–221.

Dormann, C. F. 2007. Effects of incorporating spatial autocorrelation into the analysis of species distribution data. Glob. Ecol. Biogeogr. 16:129–138.

Driese, S. G., K. H. Orvis, S. P. Horn, Z. H. Li, and D. S. Jennings. 2007. Paleosol evidence for Quaternary uplift and for climate and ecosystem changes in the Cordillera de Talamanca, Costa Rica. Palaeogeogr. Palaeoclimatol. Palaeoecol. 248:1–23.

Drummond, C. S., R. J. Eastwood, S. T. S. Miotto, and C. E. Hughes. 2012. Multiple continental radiations and correlates of diversification in *Lupinus* (Leguminosae): testing for key innovation with incomplete taxon sampling. Syst. Biol. 61:443–460.

Edgar, R. C. 2010. Search and clustering orders of magnitude faster than BLAST. Bioinformatics 26:2460–2461.

ESRI. 2018. Environmental System Research Institute (ESRI) ArcGIS release 10.6.1. ESRI, Redlands, CA.

Fine, P. V. A., D. C. Daly, G. Villa, I. Mesones, K. M. Cameron, G. V. Muñoz, I. Mesones, and K. M. Cameron. 2005. The contribution of edaphic heterogeneity to the evolution and diversity of Burseraceae trees in the western Amazon. Evolution 59:1464.

Fine, P. V. A., F. Zapata, and D. C. Daly. 2014. Investigating processes of neotropical rain forest tree diversification by examining the evolution and historical biogeography of the Protieae (Burseraceae). Evolution 68:1988–2004.

Fitzpatrick, B. M., and M. Turelli. 2006. The geography of mammalian speciation: mixed signals from phylogenies and range maps. Evolution 60:601–615.

Friis, E. 1988. *Spirematospermum chandlerae* sp. nov., an extinct species of Zingiberaceae from the North American Cretaceous. Tertiary Res. 9:7–12.

Fox, J., and S. Weisberg. 2019. An R Companion to Applied Regression, Third Edition. Sage, Thousand Oaks, CA.

Gentry, A. H. 1982. Neotropical Floristic Diversity□: Phytogeographical Connections Between Central and South America , Pleistocene Climatic Fluctuations , or an Accident of the Andean Orogeny? Ann. Missouri Bot. Gard. 69:557–593.

Ghalambor, C. K., R. B. Huey, P. R. Martin, J. Tweksbury, and G. Wang. 2006. Are mountain passes higher in the tropics? Janzen's hypothesis revisited. Integr. Comp. Biol. 46:5–17.

Grabherr, M. G., B. J. Haas, M. Yassour, J. Z. Levin, D. A. Thompson, I. Amit, X. Adiconis, L. Fan, R. Raychowdhury, Q. Zeng, et al. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat. Biotechnol. 29:644–652.

Gregory-Wodzicki, K. M. 2000. Uplift history of the Central and Northern Andes: a review. Geol. Soc. Am. Bull. 112:1091–1105.

Grossenbacher, D. L., S. D. Veloz, and J. P. Sexton. 2014. Niche and range size patterns suggest that speciation begins in small, ecologically diverged populations in north american monkeyflowers (*Mimulus* spp.). Evolution 68:1270–1280.

Guarnizo, C. E., and D. C. Cannatella. 2013. Genetic divergence within frog species is greater in topographically more complex regions. J. Zool. Syst. Evol. Res. 51:333–340.

Guisan, A., and C. Rahbek. 2011. SESAM - a new framework integrating macroecological and species distribution models for predicting spatio-temporal patterns of species assemblages. J. Biogeogr. 38:1433–1444.

Hickey, L.J., Peterson, R.K., 1978. *Zingiberopsis*, a fossil genus of the ginger family from Late Cretaceous to early Eocene sediments of western interior North America. Can. J. Bot. 56, 1136–1152.

Hijmans, R. J. 2019. Raster: geographic data and modelling. R package version 2.9-5. https://CRAN.R-project.org/package=raster

Hughes, C. E., and G. W. Atchison. 2015. The ubiquity of alpine plant radiations : from the Andes to the Hengduan Mountains. New Phytol. 207:275–282.

Janzen, D. H. 1967. Why mountain passes are higher in the tropics. Am. Nat. 101:233–249.

Johnson, M. G., E. M. Gardner, Y. Liu, R. Medina, B. Goffinet, A. J. Shaw, N. J. C. Zerega, and N. J. Wickett. 2016. HybPiper: Extracting Coding Sequence and Introns for Phylogenetics from High-Throughput Sequencing Reads Using Target Enrichment. Appl. Plant Sci. 4:1600016.

Katoh, K., and D. M. Standley. 2013. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Mol. Biol. Evol. 30:772–780.

Kay, K. M. 2006. Reproductive isolation between two closely related hummingbird-pollinated Neotropical gingers. Evolution 60:538.

Kay, K. M., and D. W. Schemske. 2003. Pollinator assemblages and visitation rates for 11 species of Neotropical *Costus* (Costaceae). Biotropica 35:198.

Kay, K. M., and D. W. Schemske. 2008. Natural selection reinforces speciation in a radiation of neotropical rainforest plants. Evolution 62:2628–2642.

Kay, K. M., P. A. Reeves, R. G. Olmstead, and D. W. Schemske. 2005. Rapid speciation and the evolution of hummingbird pollination in neotropical *Costus* subgenus *Costus* (Costaceae): Evidence from nrDNA its and ETS sequences. Am. J. Bot. 92:1899–1910.

Kissling, W.D., and G. Carl. 2008. Spatial autocorrelation and the selection of simultaneous autoregressive models. Glob. Ecol. Biogeogr. 17:59–71.

Lagomarsino, L. P., F. L. Condamine, A. Antonelli, A. Mulch, and C. C. Davis. 2016. The abiotic and biotic drivers of rapid diversification in Andean bellflowers (Campanulaceae). New Phytol. 210:1430–1442.

Landis, M. J., N. J. Matzke, B. R. Moore, and J. P. Huelsenbeck. 2013. Bayesian analysis of biogeography when the number of areas is large. Syst. Biol. 62:789–804.

Luebert, F., and M. Weigend. 2014. Phylogenetic insights into Andean plant diversification. Front. Ecol. Evol. 2:1–17.

Maas, P. J. M. 1972. Costoideae (Zingiberaceae). Flora Neotrop. 8:1–139.

Mass, P. J. M. 1977. *Renealmia* (Zingiberaceae - Zingiberoideae) Costoideae (additions) (Zingiberaceae). Flora Neotrop. 18:1–218.

Madriñán, S., A. J. Cortés, and J. E. Richardson. 2013. Páramo is the world's fastest evolving and coolest biodiversity hotspot. Front. Genet. 4:192.

Mai, U., and S. Mirarab. 2018. TreeShrink: Fast and accurate detection of outlier long branches in collections of phylogenetic trees. BMC Genomics 19:272.

Matzke, N. J. 2013. BioGeoBEARS: Biogeography with Bayesian (and likelihood) evolutionary analysis in R Scripts, CRAN: The Comprehensive R Archive Network, Vienna, Austria.

Misiewicz, T. M., and P. V. A. Fine. 2014. Evidence for ecological divergence across a mosaic of soil types in an Amazonian tropical tree: *Protium subserratum* (Burseraceae). Mol. Ecol. 23:2543–2558.

Morales-Briones, D. F., K. Romoleroux, F. Kolář, and D. C. Tank. 2018. Phylogeny and evolution of the neotropical radiation of *Lachemilla* (Rosaceae): uncovering a history of reticulate evolution and implications for infrageneric classification. Syst. Bot. 43:17–34.

Morrone, J. J. 2014. Biogeographical regionalisation of the neotropical region. Zootaxa 3782:1–110.

Nevado, B., G. W. Atchison, C. E. Hughes, and D. A. Filatov. 2016. Widespread adaptive evolution during repeated evolutionary radiations in New World lupins. Nat. Commun. 7:12384.

Nguyen, L. T., H. A. Schmidt, A. Von Haeseler, and B. Q. Minh. 2015. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol. Biol. Evol. 32:268–274.

Palma-Silva, C., T. Wendt, F. Pinheiro, T. Barbará, M. F. Fay, S. Cozzolino, and C. Lexer. 2011. Sympatric bromeliad species (*Pitcairnia* spp.) facilitate tests of mechanisms involved in species cohesion and reproductive isolation in Neotropical inselbergs. Mol. Ecol. 20:3185–3201.

Pasek, J. 2020. Weights: Weighting and Weighted Statistics. R package version 1.0.1. https://CRAN.R-project.org/package=weights

Pyron, R. A., G. C. Costa, M. A. Patten, and F. T. Burbrink. 2015. Phylogenetic niche conservatism and the evolutionary basis of ecological speciation. Biol. Rev. 90:1248–1262.

R Core Team. 2020. R: A Language and Environment for Statistical Computing. Vienna, Austria.

Rahbek, C., M. K. Borregaard, A. Antonelli, R. K. Colwell, B. G. Holt, D. Nogues-Bravo, C. M. Ø. Rasmussen, K. Richardson, M. T. Rosing, R. J. Whittaker, and J. Fjeldså. 2019a. Building mountain biodiversity: Geological and evolutionary processes. Science 365:1114–1119.

Rahbek, C., M. K. Borregaard, R. K. Colwell, B. Dalsgaard, B. G. Holt, N. Morueta-Holme, D. Nogues-Bravo, R. J. Whittaker, and J. Fjeldså. 2019. Humboldt's enigma: What causes global patterns of mountain biodiversity? Science 365:1108–1113.

Rambaut, A., M. A. Suchard, D. Xie, and A. J. Drummond. 2014. Tracer v1.6.

    http://beast.bio.ed.ac.uk/Tracer

Ree, R. H., and I. Sanmartín. 2018. Conceptual and statistical problems with the DEC+J

    model of founder-event speciation and its comparison with DEC via model selection. J.

    Biogeogr. 45:741–749.

Ree, R. H., and S. A. Smith. 2008. Maximum Likelihood Inference of Geographic Range

    Evolution by Dispersal, Local Extinction, and Cladogenesis. Syst. Biol. 57:4–14.

Revell, L. J. 2012. phytools: An R package for phylogenetic comparative biology (and other

    things). Methods Ecol. Evol. 3:217–223.

Riley, S. J., S. D. DeGloria, and R. Elliot. 1999. A terrain ruggedness index that quantifies

    topographic heterogeneity. Intermountain Journal of Sciences 5:1–4.

Roalson, E. H., and W. R. Roberts. 2016. Distinct processes drive diversification in

    different clades of Gesneriaceae. Syst. Biol. 65:662–684.

Ronquist, F. 1997. Dispersal-Vicariance Analysis: A new approach to the quantification of

    historical biogeography. Syst. Biol. 46:195–203.

Sheldon, K. S., R. B. Huey, M. Kaspari, and N. J. Sanders. 2018. Fifty years of mountain

    passes: a perspective on Dan Janzen's classic article. Am. Nat. 191:553–565.

Smith, S. A., and B. C. O'Meara. 2012. TreePL: Divergence time estimation using

    penalized likelihood for large phylogenies. Bioinformatics 28:2689–2690.

Smith, S. A., M. J. Moore, J. W. Brown, and Y. Yang. 2015. Analysis of phylogenomic

    datasets reveals conflict, concordance, and gene duplications with examples from

    animals and plants. BMC Evol. Biol. 15:150.

Smith, S. A., J. W. Brown, and J. F. Walker. 2018. So many genes, so little time: A

    practical approach to divergence-time estimation in the genomic era. PLoS One 13.

Stiles, G. F. 1981. Geographical aspects of bird-flower coevolution, with particular

    reference to Central America. Ann. Missouri Bot. Gard. 68:323–351.

ter Steege, H and Amazon Tree Diversity Network. 2010. RAINFOR (Amazon Forest

    Inventory Network). Pp. 349–359 *in* C. Hoorn and F. Wesselingh eds. Amazonia:

    Landscape and Species Evolution. Wiley, Oxford, UK.

Tripp, E. A., and Y. H. E. Tsai. 2017. Disentangling geographical, biotic, and abiotic

    drivers of plant diversity in neotropical *Ruellia* (Acanthaceae). PLoS One 12:1–17.

Uribe-Convers, S., and D. C. Tank. 2015. Shifts in diversification rates linked to

    biogeographic movement into new areas: An example of a recent radiation in the

    andes. Am. J. Bot. 102:1854–1869.

Vargas, O. M., E. M. Ortiz, and B. B. Simpson. 2017. Conflicting phylogenomic signals

    reveal a pattern of reticulate evolution in a recent high-Andean diversification

    (Asteraceae: Astereae: *Diplostephium*). New Phytol. 214:1736–1750.

Vargas, O. M., M. Heuertz, S. A. Smith, and C. W. Dick. 2019. Target sequence capture in the Brazil nut family (Lecythidaceae): Marker selection and in silico capture from genome skimming data. Mol. Phylogenet. Evol. 135:98–104.

Vargas, O. M., and C. W. Dick. 2020. Diversification history of neotropical Lecythidaceae, an ecologically dominant tree family of Amazon Rain Forest. Pp. 791–809 *in* V. Rull and A. C. Carnaval, eds. Neotropical Diversification. Springer, Cham, Switzerland.

Vargas, O. M., and B. B. Simpson. 2019. Allopatric speciation is more prevalent than parapatric ecological divergence in tropical montane systems (Asteraceae: *Piofontia*). BioRxiv 1–32. https://doi.org/10.1101/868216

Warren, D. L., R. E. Glor, and M. Turelli. 2008. Environmental niche equivalency versus conservatism: Quantitative approaches to niche evolution. Evolution 62:2868–2883.

Weir, J. T., and D. Schulter. 2008. The latitudinal gradient in recent speciation and extinction rates of birds and mammals. Science 315:1574–1577.

Weir, J. T., M. S. Faccio, P. Pulido-Santacruz, A. O. Barrera-Guzmán, and A. Aleixo. 2015. Hybridization in headwater regions, and the role of rivers as drivers of speciation in Amazonian birds. Evolution 69:1823–1834.

Yu, G., D. K. Smith, H. Zhu, Y. Guan, and T. T. Y. Lam. 2017. Ggtree: an R package for visualization and annotation of phylogenetic trees with their covariates and other associated data. Methods Ecol. Evol. 8:28–36.

Zhang, C., E. Sayyari, and S. Mirarab. 2017. ASTRAL-III: increased scalability and impacts of contracting low support branches. Pp. 53–75 *in* J. Meidanis and L. Nakhleh, eds. Comparative Genomics. Springer, Cham, Switzerland.

Zhbannikov, I. Y., S. S. Hunter, J. A. Foster, and M. L. Settles. 2017. SeqyClean: A Pipeline for High-throughput Sequence Data Preprocessing. Pp. 407–416 *in* Proceedings of the 8th ACM International Conference on Bioinformatics, Computational Biology, and Health Informatics. ACM, New York, NY, USA.