Serratia marcescens **secretes proteases and chitinases with larvicidal activity**

against *Anopheles dirus*

Natapong Jupatanakul[1,*], Jutharat Pengon[1], Shiela Marie Gines Selisana[2#], Waeowalee

Choksawangkarn[3], Nongluck Jaito[4], Atiporn Saeung[5], Ratchanu Bunyong[1], Navaporn

Posayapisit[1], Khrongkhwan Thammatinna[6], Nuttiya Kalpongnukul[7,8], Kittipat Aupalee[5], Trairak

Pisitkun[8] and Sumalee Kamchonwongpaisan[1]

[1] Protein-Ligand Engineering and Molecular Biology Research Team, National Center for Genetic

Engineering and Biotechnology (BIOTEC), National Science and Technology Development Agency

(NSTDA), Pathumthani, Thailand

[2] Institute of Biological Sciences, College of Arts and Sciences, University of the Philippines Los Baños,

Laguna, Philippines

[3] Department of Biochemistry, Faculty of Science, Burapha University, Chonburi, Thailand

[4] Enzyme Technology Research Team, National Center for Genetic Engineering and Biotechnology

(BIOTEC), National Science and Technology Development Agency (NSTDA), Pathumthani, Thailand

[5] Department of Parasitology, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand

[6] Department of Biology, Faculty of Science, Prince of Songkhla University, 15 Karnjanavanich Rd., Hat

Yai, Songkhla, Thailand

[7] Inter-Department Program of Biomedical Sciences, Faculty of Graduate, Chulalongkorn University,

Bangkok, Thailand

[8] Center of Excellence in Systems Biology, Faculty of Medicine, Chulalongkorn University, Bangkok,

Thailand

*Corresponding author: Email: natapong.jup@biotec.or.th

#Current address: School of Life Science and Technology, Tokyo Institute of Technology,

Nagatsuta-cho, Midori-ku, Yokohama, Kanagawa 226-8501, Japan.

ABSTRACT

Vector control, the most efficient tool to reduce mosquito-borne disease transmission, has been compromised by the rise of insecticide resistance. Recent studies suggest the potential of mosquito-associated microbiota as a source for new biocontrol agents or new insecticidal chemotypes. In this study, we identified a strain of *Serratia marcescens* that has larvicidal activity against *Anopheles dirus*, an important malaria vector in Southeast Asia. This bacterium secretes heat-labile larvicidal macromolecules when cultured under static condition at 25°C but not 37°C. Two major protein bands of approximately 55 kDa and 110 kDa were present in spent medium cultured at 25°C but not at 37°C. The Liquid Chromatography-Mass Spectrometry (LC-MS) analyses of these two protein bands identified several proteases and chitinases that were previously reported for insecticidal properties against agricultural insect pests. The treatment with protease and chitinase inhibitors led to a reduction in larvicidal activity, confirming that these two groups of enzymes are responsible for the macromolecule's toxicity. Taken together, our results suggest a potential use of these enzymes in the development of larvicidal agents against *Anopheles* mosquitoes.

Keywords: *Anopheles*; microbiota; larvicidal; mass spectrometry; entomopathogen

1. Introduction

Aside from nuisances, mosquitoes are vectors of a wide range of human pathogens including *Plasmodium* parasites that cause malaria, filarial nematodes and viruses such as dengue, Zika, yellow fever, and chikungunya. These pathogens cause diseases that are major public health and economical burdens causing approximately 500,000 deaths each year (Shaw and Catteruccia, 2019), with malaria alone accounting for 90% of the deaths. *Anopheles dirus* has been incriminated as the primary vector of the malarial protozoa *P. falciparum* and *P. vivax* in Southeast Asia (Manguin et al., 2008). Although malaria cases and disease burden have significantly decreased over the past decades owing to an extensive effort in mosquito control and antimalarial drug treatment, malaria elimination faces obstacles such as the increasing insecticide resistance of mosquitoes as well as the emergence of antimalarial drug resistance of *Plasmodium* parasites (Whitty and Ansah, 2019). While there are several new classes of antimalarial drugs in the clinical pipelines ("MMV-supported projects | Medicines for Malaria Venture," 2020; Wells et al., 2015), insecticides with new mode of action are much less developed (Shaw and Catteruccia, 2019; Whitty and Ansah, 2019).

Mosquitoes live in constant interactions with microorganisms in their environment and throughout their life cycle (Gendrin and Christophides, 2013; Wang et al., 2011), and in particular, the aquatic stages of the mosquito larvae because they are submerged in water (Dickson et al., 2017). Although microbiota are needed for the healthy development of the mosquito larva (Chouaia et al., 2012; Coon et al., 2014; Dickson et al., 2017; Valzania et al., 2018), certain microbes can be pathogenic to them (Baumann et al., 1991; Laurence et al., 2011). These entomopathogenic microorganisms, such as *Bacillus thuringiensis* and *Bacillus sphaericus* or their by-products, have long been used for mosquito population control (Baumann et al., 1991; Laurence et al., 2011). In addition to these *Bacillus* species, recent studies have identified entomopathogenic bacteria from the mosquito gut microbiota such as *Chromobacterium* spp. and *Serratia marcescens* (Bahia et al., 2014; Ramirez et al., 2014; Short et al., 2018). *Chromobacterium* spp. isolated from the mosquito gut from Panama *(Csp_P)* has insecticidal properties and has potential to be developed as a mosquito larvicidal

agent (Caragata et al., 2020). This suggests that the natural microbial communities associated

with mosquitoes are invaluable resources for the discovery of biocontrol agents with new mode

of actions and the development of novel insecticides.

Our laboratory has recently set up a new insectary facility to expand our drug development

pipeline to test antimalarial compounds against sexual stage of *Plasmodium* parasites. During

the optimization of *Anopheles dirus* rearing condition, we found that if the larval food particle

size was not fine enough, the food sedimented to the bottom of the pan. Due to the surface

feeding nature of *An. dirus*, the sedimented food led to a bacterial growth and biofilm at the

bottom of the pan. Mortality of the mosquito larva was observed a few days after the biofilm

formed at the bottom of the pan. From this observation, we hypothesized that certain microbes

in the larval rearing pan secrete molecule(s) that have larvicidal activity against *An. dirus* after

the bacteria grow to a certain density when culture under static condition. In this study, we

isolated and identified culturable bacteria with distinct morphology from the *An. dirus* larval

rearing pan where larval death was being observed. Of seven isolates, we found a strain of

Serratia marcescens that secreted heat-labile larvicidal macromolecules into spent medium

when it was cultured under static condition at 25°C. We then used mass spectrometry to

identify chitinases and proteases as potential larvicidal molecules. Chitinase and protease

inhibitors were then used to confirm that these molecules contributed to larvicidal activity. This

study demonstrates that the chitinases and proteases uncovered in this study could potentially

be used as larvicides against *Anopheles* mosquito.

2. Materials and Methods

2.1 Ethics statement

Animal work in this study was carried out in accordance to the recommendations

provided in the Guide for the Care and Use of Laboratory Animals of the National Institutes of

Health. Mice were used only as a blood source for mosquito rearing, according to the approved

protocol (BT-Animal 26/2560). Larvicidal activity assays were performed following the

approved protocol (BT-Animal 20/2560). Both animal protocols were approved by the BIOTEC

Committee for use and care of laboratory animals.

2.2 Mosquito rearing condition

Specimens of laboratory strain of *An. dirus* were originally collected in Mae Sod

District, Tak Province, Thailand. A colony of *An. dirus* was maintained at the Department of

Parasitology, Chiang Mai University and BIOTEC's insectary at 27 °C with 75% humidity and

a 12-h day/night, 30-min dusk/ dawn lighting cycle. The larvae were fed a diet of powdered

fish food (Tetrabit, Germany). Adults were fed on a 10% sucrose solution supplemented with

5% multivitamin syrup (Sevenseas, UK) *ad libitum* (Choochote and Saeung, 2013). Female

ICR mice were anesthetized using 2% Tribromoethanol and used as a blood source to

maintain mosquito colony (approved protocol BT-Animal 26/2560)

2.3 Bacterial isolation and identification

Bacteria were isolated from two biological niches, larval rearing water and bacterial

biofilm at the bottom of the rearing pan. The rearing water was 10-fold serially diluted with

sterile 1X Phosphate buffered saline (PBS). The biofilm was gently rinsed five times with 1X

PBS then the biofilm was scraped with sterile plastic scraper, resuspended with 20 mL of

sterile PBS then 10-fold serially diluted. The serial dilutions were then plated on Luria-Bertani

(LB) and incubated at room temperature for 2 days. Bacterial colonies with distinct

morphologies from each ecological niche were then streaked on LB agar plate. A single colony

from each bacterial isolate was then propagated and used for molecular identification using

16s rRNA gene. Briefly, a colony of each isolate was resuspended in 50 µL of sterile water

then 1 µL of resuspended bacteria was used in a PCR reaction using Phusion® High-Fidelity

DNA Polymerase (New England Biolabs) with forward (16s-8F: 5'

AGAGTTTGATCCTGGCTCAG 3') and reverse (16s-1492R: 5'

CGGTTACCTTGTTACGACTT3') primers (Eden et al., 1991). PCR amplification was

performed with an initial denaturation of 2 minutes 30 seconds at 98°C, followed with 30 cycles

of denaturation at 98°C for 10 seconds, annealing at 52°C for 15 seconds and an extension

at 72°C for 1 minute. The amplicons were then purified using QIAquick PCR purification kit

then submitted for Sanger DNA sequencing. The obtained bacterial 16s rRNA gene

sequences from the forward and reverse primers were manually trimmed and assembled. The

assembled sequences were then searched on the NCBI's Basic Local Alignment Search Tool

(BLAST, https://blast.ncbi.nlm.nih.gov/Blast.cgi) (Altschul et al., 1990).

2.4 Bacteria culture and bacterial spent media preparation

All isolated bacteria were cultured in LB medium under specific conditions as

mentioned for each experiment. To start a culture, a colony of each bacterium was inoculated

in 10 mL of LB medium then cultured at 25°C, 200 rpm overnight. The absorbance at 600 nm

(OD_{600}) was measured for the overnight culture, then 1 mL of culture at OD_{600} = 1 was

inoculated into 5 mL of fresh LB medium. For static culture, 18 mL of inoculated culture was

added to a 90 mm diameter petri dish then incubated without shaking at specified temperature.

For planktonic culture, 18 mL of inoculated culture was added to a 50 mL conical tube then

agitated at 200 rpm at specified temperature. After incubation, bacterial cells were removed

from the spent media by centrifugation at 4,415 x g 4 °C for 45 minutes. Supernatants were

then filtered through 0.22 μm syringe filter (Sartorius). The sterile spent media were then kept

at 4 °C until further use.

2.5 Larvicidal activity assay

Second instar larvae of *An. dirus* were filtered and rinsed twice with 250 mL sterile

distilled water. Larvicidal assays were performed with 20 mosquitoes in 4 mL of total water

volume per treatment per replicate in each well of 6-well plate. Filtered bacteria spent media

were added to each well at 10% v/v and mortality was recorded for 72 hours. For the group

with antibiotics treatment, Penicillin/Streptomycin solution (Gibco) were added to the well at

final concentration of 100 U/mL and 100 μg/mL, respectively.

For *Pseudomonas* sp. challenge, the bacterium was cultured overnight at 25 ºC then

the cells were collected by centrifugation at 1500 x *g*, 4 ºC for 15 minutes. Bacterial cells were

then washed twice with 1X PBS, diluted then added to larvicidal plate at concentrations as

specified in the results section.

2.6 Characterization of physical property of secreted larvicidal molecules and preparation of

concentrated macromolecule fraction

We first evaluated whether the secreted larvicidal molecules were proteins

(macromolecules) or small molecules because they would require different methods of

identification. Thus, the larvicidal molecules were characterized based on the heat stability

and molecular weight.

The heat stability of the secreted larvicidal molecules was determined by heating the

spent media at 45, 70, and 95 ºC for one hour. Heated spent media were then used to test for

remaining larvicidal activity.

Proteins (macromolecules) and small molecules were separated using Amicon® Ultra-

4 10K Centrifugal Filter Device following the manufacturer's instructions. Briefly, 4 ml of spent

media were added to the filter unit then centrifuged at 1500 x *g*, 4 ºC for 30 minutes. The flow

through was collected as small molecule fraction while the molecules retained on the filter

device were used as macromolecule fraction. The concentrated macromolecule fraction was

reconstituted with either LB or 1X PBS to concentrations as indicated in each experiment.

2.7 Cytotoxicity in C6/36 insect cell line

Cytotoxicity of *S. marcescens* spent medium in insect cell line was determined by

sulforhodamine B (SRB) assay according to previously published protocol (Vichai and

Kirtikara, 2006) with slight modification for C6/36 cell line. The SRB assay measured cellular

protein content to determine cell survival after spent medium treatment. Briefly, C6/36 cells

were seeded into each well of 96-well plate at 30,000 cells/well in 100 µl of 10% (v/v) *S.*

marcescens spent medium in complete L15 medium (L15 supplemented with 10% Fetal

bovine serum, 1% MEM non-essential amino acids, 2% Tryptose phosphate broth, 100 U/mL

Penicillin and 100 µg/mL Streptomycin) and incubated at 28 ºC for 72 hours. After incubation,

cell monolayers were fixed with 10% (w/v) trichloroacetic acid for 45 minutes at 4 ºC and rinsed

4 times with tap water. Fixed cell monolayer was then stained with 0.057% (w/v) SRB in 1%

acetic acid for 30 minutes at room temperature in the dark. Stained cell monolayers were then

washed with 100 µl 1% acetic acid for four times to remove unbound dye. To solubilize protein-

bound SRB, 200 µl of 10 mM Tris (pH 10.5) was added to each well then the plate was shaken

for 30 minutes at room temperature. Absorbance at 510 nm was then measured in a

microplate reader to determine cell viability. LB was used as an untreated control and 5 µM

cycloheximide was used as a positive control of the cytotoxicity assay.

2.8 In-gel digestion of secreted S. marcescens macromolecules for LC-MS/MS analysis

The cell-free macromolecules of *S. marcescens* static culture at 25 and 37 ºC were

mixed with Laemmli buffer, denatured at 95 ºC for 10 minutes then subjected to SDS-PAGE.

After the SDS-PAGE fractionation, protein bands present in the 25 but not in 37 ºC culture

condition were excised from the gel for in-gel tryptic digestion followed by LC-MS/MS analyses

(Shevchenko et al., 2007). Briefly, the gel bands were excised and dehydrated in 100%

acetonitrile. Proteins were reduced by 10 mM dithiothreitol at 56°C for 30 min and alkylated

by 20 mM chloroacetamide at room temperature for 20 min. The gel bands were further

destained using 50 mM NH_4HCO_3 in 50% acetonitrile for 30 min, followed by dehydration in

neat acetonitrile. Digestion was carried out by adding 13 ng/µL trypsin in 50 mM NH_4HCO_3 to

cover the gel pieces, and the reaction was incubated at 37°C overnight. Extraction of cleaved

peptides was performed by adding extraction buffer (70% acetonitrile/1% formic acid) to the

digests at the ratio of 2:1 (v/v). Then the extracts were lyophilized and redissolved in 0.1%

formic acid ready for LC-MS/MS analyses.

2.9 LC-MS/MS analyses of secreted S. marcescens macromolecules

The digested peptides (200 ng) were analyzed by an EASY-nLC 1000 HPLC system

(Thermo Fisher Scientific) interfaced with a Q Exactive™ Hybrid Quadrupole-Orbitrap mass

spectrometer (Thermo Fisher Scientific) via an EASY-Spray ion source (Thermo Fisher

Scientific). The separation was performed for 60 min at the flow rate of 300 nL/min using C18-

reverse phase column (EASY-Spray™, Thermo Fisher Scientific). The stepwise gradient was

set from 5% solvent B (0.1% formic acid in acetonitrile)/ 95% solvent A (0.1% formic acid in

H_2O) to 20% solvent B/ 80% solvent A for 43 min and then to 40% solvent B within the next

10 min, followed by holding at 95% for 55 - 60 min. Data-dependent acquisition was performed

wherein the ten most abundant precursor ions were selected for fragmentation. The precursor

masses were scanned by the Orbitrap mass analyzer at the mass range of 400 – 1600 m/z

and the resolution of 70,000. Higher-energy collisional dissociation was employed for

fragmentation with an exclusion of +1 charged ions. All data were acquired using Thermo

Xcalibur 2.2 software.

Mass spectra were searched against NCBI database of *Serratia* sp. (August, 2018)

using Proteome Discoverer 2.2 connected to SEQUEST search engine (Eng et al., 1994).

Precursor mass tolerance was set to 10 ppm and the product ion mass tolerance was 0.02

Dalton. Trypsin was specified with maximum 2 missed cleavages. Carbamidomethylation was

chosen as a fixed modification; while oxidation of methionine was allowed as variable

modification. Decoy search was performed using the false discovery rate of 0.01 as a

threshold.

2.10 Chitinase activity assay

Chitinase secretion of *S. marcescens* when cultured in different culture temperature

was demonstrated by the ability of the bacterium to generate clear zone on LB agar

supplemented with 0.5% chitin. *S. marcescens* inoculated plates were cultured at 25°C or

37°C for 10 days before visualization of clear zone.

Chitinase activity of spent medium was determined by measuring the ability of the

spent medium to digest chitin into a reducing sugar, *N*-acetyl-*D*-glucosamine. To set up the

chitinase activity assay, 100 µL of cell-free spent culture media served as a crude enzyme

source was mixed with 100 µL of 1% (w/v) chitin powder (Sigma, USA) in 50 mM sodium

phosphate buffer (pH 6.5). The reaction mixture was incubated at 30ºC for 90 min. The amount

of reducing sugar was then determined by using *p*-hydroxybenzoic acid hydrazide (pHBAH)

assay (Lever, 1972). A stock solution of 5% (w/v) pHBAH (Sigma, USA) was prepared in acid

medium (0.5 M HCl) and diluted at the moment of analysis to 1% (w/v) in alkaline medium (0.5

M NaOH). Samples were mixed with 1% (w/v) pHBAH in alkaline solution (0.5 M NaOH) at a

ratio of 1:3 and the mixture was heated in the boiling bath for 5 min to develop the color. After

cooling, the solution was centrifuged at 13,800 x *g* for 2 min. The absorbance at 410 nm of

the supernatant was then measured using spectrophotometer. *N*-acetyl-*D*-glucosamine

standard solution was prepared in the same manner at concentrations of 0, 0.2, 0.4, 0.6, 0.8,

and 1 mg/ml in distilled water. One unit of enzyme activity was defined as the amount of

enzyme which releases 1 µg of reducing sugar per hour under the assay conditions.

2.11 Statistical analyses

Statistical significance of the survival analyses was calculated using Log-rank test

followed by a Bonferroni post-hoc analysis using 'survminer ' package (Kassambara et al.,

2019) in R (R Core Team, 2020).

Concentration of *Pseudomonas* sp. treatment resulting in 50% larval mortality (LC_{50})

and 90% larval mortality (LC_{90}) was calculated by Probit analysis with Abbot's modification

using 'ecotox' package (Scholefield et al., 2008) in R (R Core Team, 2020).

Statistical analysis comparing chitinase activity level and inhibition of chitinase activity

was performed using One-way ANOVA with Tukey multiple comparison test using

multcompView (Graves and Dorai-Raj, 2019) package in R (R Core Team, 2020).

3. Results

3.1 Isolation of bacteria from An. dirus larval rearing pan

A total of seven bacteria species were isolated and identified from the rearing pan by

16s rRNA gene sequencing. These include one Actinobacteria: *Leucobacter* spp. (Leu_spp),

two Bactereoidetes: *Sphingobacterium* spp. (Sph_spp) and *Sphingobacterium multivorum*

(Sch_m); and four proteobacteria: *Acinetobacter soli* (Aci_s), *Ochrobactrum* spp. (Och_spp),

Serratia marcescens (Ser_m), and *Stenotrophomonas maltophilia* (Ste_m) as summarized in

Table 1.

Table 1. Bacterial isolates from *An. dirus* larval rearing pan identified by 16s rRNA gene

sequencing

Abbreviation	Species	Phylum	Top known hit	Identities	Gaps
Leu_spp	*Leucobacter* spp.	Actinobacteria	*Leucobacter* sp. Re-6	1332/1332 (100%)	0/1332 (0%)
Sph_spp	*Sphingobacterium* spp.	Bacteroidetes	*Sphingobacterium* sp. strain JPB23	1305/1340 (97%)	0/1340 (0%)
Sph_m	*Sphingobacterium multivorum*	Bacteroidetes	*Sphingobacterium multivorum* strain GZT-2	1341/1342 (99%)	1/1342 (0%)
Ser_m	*Serratia marcescens*	Proteobacteria	Serratia sp. strain p2	1346/1346 (100%)	0/1346 (0%)
Aci_s	*Acinetobacter soli*	Proteobacteria	*Acinetobacter soli* strain GFJ2	1347/1349 (99%)	0/1349 (0%)
Ste_m	*Stenotrophomonas maltophilia*	Proteobacteria	*Stenotrophomonas maltophilia* strain D299	1259/1350 (93%)	1/1350 (0%)
Och_spp	*Ochrobactrum* spp.	Proteobacteria	*Ochrobactrum* sp. Azn6.2	1296/1307 (99%)	10/1307 (0%)

3.2 S. marcescens secreted larvicidal molecule into spent medium when cultured under static

condition

Because larval mortality was observed a few days after bacterial bloom, we

hypothesized that the isolated bacteria could secrete larvicidal molecule(s) after the bacteria

grow to a certain density; at a time when biofilm was observed at the bottom of the pan for a

few days. Therefore, the isolated bacteria were cultured under either static or planktonic

conditions for three days before spent media collection for the screening of larvicidal activity.

None of the cell-free spent media obtained from planktonic condition had any toxicity

against *An. dirus* larvae (Figure 1A). However, for the stationary culture, only the cell-free

spent media from *S. marcescens* had a detrimental effect on the survival of the larvae (Figure

1A). Next, spent medium of *S. marcescens* cultured under planktonic and static conditions

was collected over-time after 1, 2, or 3 days of incubation. Again, the larvicidal molecule was

secreted into the *S. marcescens* only under static culture condition and the larvicidal activity

of the spent medium increased over time starting from the second day after inoculation (Figure

1B).

To investigate if the *S. marcescens* cells are entomopathogenic to *An. dirus* larvae,

the bacterium was cultured under a static condition then washed with 1X PBS. Washed

bacterial cells were then added to larval culture at various concentrations (Figure 1C). Even

at OD_{600} of 5, *S. marcescens* cells did not cause significant mortality to *An. dirus* larvae. The

Escherichia coli cells used as a control also did not cause larval mortality. These results

suggested that *S. marcescens* only cause larval mortality through the secretion of

entomopathogenic molecules.

Figure 1. Survival analyses of *An. dirus* mosquito larvae after treatment with bacterial spent

media at 10% (v/v) concentration. (A) Survival curve of *An. dirus* larvae after treatment with

bacterial spent media cultured under planktonic condition or static condition. The spent media

were collected at three days after incubation under planktonic or static state at 25 ºC Ech_c:

Escherichia coli strain XL10, Aci_s: *Acinetobacter soli*, Leu_spp: Leucobacter spp., Och_spp:

Ochrobatrum spp., Ser_m: *Serratia marcescens*, Sph_m: *Sphingobacterium multivorum,*

Sph_spp: *Sphingobacterium* spp., Ste_m: *Stenotrophomonas maltophilia*. LB medium or

water was used as a control at final concentration of 10% (v/v). The graphs represent data

from three independent biological replicates (B) Survival curve of *An. dirus* larvae after

treatment with *S. marcescens* spent medium cultured under planktonic or static condition and

collected after 1, 2, or 3 days after inoculation. The graphs represent data from two

independent biological replicates. (C) Survival curve of *An. dirus* larvae after treatment with *E.*

coli or *S. marcescens* cells at various concentration. Bacterial cells were cultured under static

condition at 25 ºC then washed twice with PBS. The graphs represent data from three

independent biological replicates. Statistical significance of the survival analyses was

calculated using Log-rank test followed by a Bonferroni post-hoc analysis using 'survminer

'package in R. Significance level: **: p<0.01, ****: p<0.0001.

3.3 Larvicidal molecules from S. marcescens are heat sensitive macromolecules

Because the nature of larvicidal molecule would dictate subsequent methods for

identification and characterization, the larvicidal molecule was then next broadly characterized

based on heat sensitivity and molecular weight. The *S. marcescens* filtered spent medium was

heated at 45, 70 and 95 ºC for one hour then used to determine larvicidal activity at 10% (v/v)

(Figure 2A). The larvicidal molecules secreted by *S. marcescens* are heat-sensitive as the

heat treatment at 45 ºC for 1 hour significantly reduced the 3-day mortality rate of treated

larvae from 100% to 35%. Heat treatment at 70 and 95 ºC completely destroyed the larvicidal

activity. To further characterize if the heat-labile active larvicidal elements were small or large

macromolecules, we fractionated the active spent media by size exclusion using Amicon®

Ultra-4 10K filter device and tested their larvicidal activity. The result shows that only the

macromolecule fraction with molecular weight larger than 10 kDa possesses larvicidal activity

(Figure 2B).

Figure 2. Larvicidal molecule(s) secreted by *S. marcescens* are heat sensitive macromolecule(s). (A) Heat sensitivity of larvicidal activity from *S. marcescens* spent medium. The *S. marcescens* spent medium was heated at 45, 70, and 95 ºC for one hour before testing for larvicidal activity at 10% (v/v) final concentration. LB medium was used as a control at a final concentration of 10% (v/v). The graphs represent data from two independent biological replicates (B) Larvicidal activity of *S. marcescens* spent medium after separation of small (<10 kDa) and macromolecules (>10 kDa) using Amicon® Ultra-4 10K filter device at 10% (v/v). LB medium was used as a control at final concentration of 10% (v/v). The graphs represent data from two independent biological replicates. Statistical significance of the survival analyses was calculated using Log-rank test followed by a Bonferroni post-hoc analysis using 'survminer' package in R.

3.4 Larvicidal activity of S. marcescens spent medium is not likely to be caused by cytotoxicity.

To test the cytotoxicity of *S. marcescens* spent medium against insect cells, the spent medium was used to treat C6/36 cells at the final concentration of 10% (v/v) spent medium in complete L15 medium at 28 ºC for 72 hours. When compared to LB medium treatment, the cell viability of C6/36 treated with *E. coli* spent medium, *S. marcescens* spent medium, and cycloheximide were $111.25\pm8.23\%$, $76.79\pm15.02\%$, and $7.82\pm2.06\%$, respectively (Figure 3). Although statistically significant, the low cytotoxicity level of *S. marcescens* spent medium against C6/36 cells could not support the strong larvicidal activity as shown in our experiments. This could be because the cell does not carry the essential molecule/pathway targeted by *S.*

marcescens' entomopathogenic molecules. Another possible explanation is that the death of

An. dirus larvae is due to damage at the physiological level rather than direct toxicity to insect

cells.

Figure 3. C6/36 cell viability assay of *E. coli* spent medium (Ech_c), *S. marcescens* spent

medium (Ser_m), and 5 µM cycloheximide. The cell viability was shown relative to LB medium

treatment. Spent medium and LB were used at 10% (v/v) final concentration. P-values were

calculated using raw data from sulforhodamine B assay by One-way ANOVA with TukeyHSD

post hoc test with multcompView package in R software.

3.5 S. marcescens spent medium can directly kill An. dirus larvae and also enhance larvae's

susceptibility to bacterial infection.

In previous experiments, LB medium was used as a control in the larvicidal assays.

We observed that the presence of LB or filtered spent medium allowed bacteria in the

environment to grow in these treatments (Figure 1 and 2). Although these bacteria did not

cause larval death in the LB-treated group, they might interfere with the larvicidal activity

caused by *S. marcescens* spent medium treatment.

To test whether these naturally growing bacteria have any impact on larvicidal activity

by *S. marcescens*, penicillin (100 U/mL) and streptomycin (100 µg/mL) were added in the

larvicidal assay experiment to prevent bacterial growth. The addition of antibiotics significantly

inhibited bacterial contamination because water in each well of the larvicidal assay remained

clear throughout the experiment. In the absence of natural bacterial growth, the

macromolecules secreted by *S. marcescens* can still kill *An. dirus* larvae confirming its intrinsic

toxicity (Figure 4A). Interestingly, the observed mortality in the antibiotics-treated group was

significantly lower than the group without antibiotics treatment. In addition to antibiotics

treatment, resuspension of *S. marcescens* macromolecules in PBS instead of LB also reduced

natural bacterial growth in larvicidal assays. Even without obvious natural bacterial growth, *S.*

marcescens macromolecule resuspended in PBS could kill *An. dirus* larvae in a concentration-

dependent manner (Figure 4B). These results suggest that the larvicidal molecule has intrinsic

larvicidal activity, and also increases larvae's susceptibility to bacterial infection.

To confirm the hypothesis that *S. marcescens* secreted macromolecules increases

larvae's susceptibility to bacterial infection, we pretreated *An. dirus* larvae with a sublethal

concentration of *S. marcescens* secreted macromolecule (0.5X diluted in PBS; see Figure 4B)

for 24 hours then added an entomopathogenic strain of *Pseudomonas* sp. at the final bacterial

density of OD_{600} = 5, 1, 0.4, 0.2, and 0.08 (Figure 4C). *An. dirus* larvae pretreated with *S.*

marcescens secreted macromolecules significantly increased the mortality rate caused by

Pseudomonas sp. in all treatment except at OD 5, which the entomopathogenicity was too

high to observe the difference (Figure 4D). The *Pseudomonas* sp. LC_{50} of the *S. marcescens*

macromolecule-treated group was 0.115 OD_{600}, a 55% decrease from 0.258 OD_{600} in the PBS-

treated group. Similarly, the *Pseudomonas* sp. LC_{90} of the *S. marcescens* macromolecule-

treated group was 0.287 OD_{600}, a 70% decrease from 0.962 OD_{600} in the PBS-treated group

(Figure 4E and 4F). These results confirm significant effect of *S. marcescens* macromolecules

on susceptibility of *An. dirus* larvae to bacterial infection, even at sublethal concentration.

Figure 4. Intrinsic larvicidal activity of *S. marcescens* macromolecule and its enhancing effect

on larvae's susceptibility to bacterial infection. (A) Survival of *An. dirus* larvae treated with 2X

concentration of macromolecules from *S. marcescens* spent medium diluted in LB with or

without antibiotics. LB with or without antibiotics were used as a control. The graphs represent

data from two independent biological replicates (B) *An. dirus* larvae treated with *S.*

marcescens macromolecules reconstituted to the final concentration of 0.5X, 1X, and 2X in

PBS. The graphs represent data from two independent biological replicates. (C) Schematic

diagram of the treatment of *An. dirus* larvae with sublethal concentration of *S. marcescens*

secreted macromolecule followed by *Pseudomonas* sp. treatment. The larvae were treated for

24 hours with 0.5x concentration of macromolecules from *S. marcescens* diluted in PBS buffer

before addition of *Pseudomonas* sp. at final concentration of OD_{600} = 0.08, 0.2, 0.4, 1, and 5.

Mortality of the larvae was monitored every 3 hours for 24 hours. The graphs represent data

from three independent biological replicates. (D) Kaplan-Meier curve of *An. dirus* larvae from

Pseudomonas sp. after treatment with sublethal concentration of *S. marcescens*

macromolecules comparing to the PBS control. (E) Probit analysis of *An. dirus* survival

probability at 24 hours after treatment of *S. marcescens* macromolecules. (F) LC_{50} and LC_{90}

of *Pseudomonas* sp. treatment in *An. dirus* calculated from Probit analysis.

3.6 S. marcescens secreted different sets of macromolecules when cultured under different

conditions

Because we observed different levels of larvicidal activity from *S. marcescens* spent

medium when cultured under different conditions, the comparison of macromolecules from

these culture conditions would help us to pinpoint macromolecule(s) that has larvicidal activity.

Although the planktonic condition did not induce secretion of a larvicidal molecule, *S.*

marcescens barely secreted proteins under planktonic condition. Further comparison between

these two conditions would yield several protein candidates for further studies.

To identify other culture conditions that might result in differential larvicidal activity, *S.*

marcescens was cultured under static condition at different temperatures, 25 ºC and 37 ºC.

Here, *S. marcescens* spent medium from 25 ºC incubation could kill *An. dirus* larvae while

those cultured at 37 ºC could not (Figure 5A).

Pattern of secreted protein from *S. marcescens* cultured at 25 ºC and 37 ºC was then

compared by SDS-PAGE (Figure 5B). Two major protein bands (approximately 110 and 55

kDa) were observed in the spent medium from 25ºC culture but not 37ºC (Figure 5B). These

two protein bands were then excised from the gel then subjected to LC-MS/MS for

identification.

The 110 kDa protein band consists of several protease enzymes including

aminopeptidase N, serralysin, and peptidase M60 viral enhancin protein; chitinase enzymes

including chain A chitobiase; cell surface protein; DUF4214 domain-containing protein;

hemolysin; DUF4214 domain-containing protein; peptide ABC transporter substrate-binding

protein; aconitate hydratase; DNA-directed RNA polymerase; formate dehydrogenase-N

subunit alpha; glycine dehydrogenase; translation elongation factor Tu; phosphoenolpyruvate

carboxykinase; LysR family transcriptional regulator; short-chain dehydrogenase/reductase

SDR; phosphorylcholine phosphatase; acyl-CoA dehydrogenase (Supplementary table 1).

The 55 kDa protein band also consists of proteases including serralysin, proline

dipeptidase, extracellular serine protease precursor, secreted alkaline metalloproteinase, and

Xaa-Pro aminopeptidase; chitinase enzyme; serine 3-dehydrogenase; DUF4214 domain-

containing protein; arginine-tRNA ligase; succinylarginine dihydrolase; cell surface protein;

histidine ammonia-lyase; capsule assembly Wzi family protein; glutathione reductase;

phosphoenolpyruvate carboxykinase; flagellin; F0F1 ATP synthase subunit beta; and

hypothetical protein (Supplementary table 2).

Figure 5. *S. marcescens* secreted larvicidal molecules when cultured under static condition at

25 ºC but not 37 ºC. (A) Survival curve of *An. dirus* larvae after treatment with *S. marcescens*

spent media cultured under static condition at 25 ºC or 37 ºC. The graphs represent data from

three independent biological replicates. Statistical significance of the survival analyses was

calculated using Log-rank test followed by a Bonferroni post-hoc analysis using 'survminer

'package in R. (B) SDS-PAGE of *S. marcescens* spent media cultured under static condition

at 25 ºC or 37 ºC. Two red arrows indicate major protein bands that were expressed when the

bacterium was cultured at 25 ºC but not 37 ºC.

Supplementary Table 1. LC-MS analysis of 110 kDa protein band showing the NCBI

accession number, protein description, percentage of the protein sequence covered by

identified peptides (% sequence coverage), the number of identified peptides (# Peptides),

total number of identified peptide spectra matched for each identified protein (#PSM),

number of amino acid residues in each identified protein (#AA), molecular weight of proteins

(MW), calculated pI of proteins (Calc. pI) and the identification score of proteins calculated

by Sequest HT algorithms (Protein score).

Accession number	Protein description	% Sequence coverage	# Peptides	# PSMs	# AAs	MW [kDa]	Calc. pI	Protein score

1081143318	cell surface protein [*Serratia* sp. HMSC15F11]	27	18	84	998	100.9	4.79	210.67
1333427106	DUF4214 domain-containing protein [*Serratia marcescens*]	10	7	65	1000	101.1	4.83	191.57
1047610669	aminopeptidase N [*Serratia marcescens* subsp. marcescens]	15	10	11	872	98.1	5.49	16.76
1184111806	peptidase M60 viral enhancin protein [*Serratia marcescens*]	11	8	9	855	94.3	6.39	16.36
983153020	peptidase M60 viral enhancin protein [*Serratia marcescens*]	11	8	9	855	94.2	6.32	16.29
1091769957	hemolysin [*Serratia marcescens*]	11	13	15	1608	165	8.73	10.8
1027975265	peptide ABC transporter substrate-binding protein [*Serratia marcescens*]	4	2	2	535	59.9	7.2	4.13
542019733	protease [*Serratia marcescens* EGD-HP20]	4	3	3	964	107.6	6.2	3.55
944700465	aconitate hydratase [*Serratia* sp. Leaf50]	2	1	1	891	97.3	5.85	2.77
291423462	DNA-directed RNA polymerase, beta subunit [*Serratia* odorifera DSM 4582]	5	5	6	1365	153	5.2	2.73

1194684911	MULTISPECIES: formate dehydrogenase-N subunit alpha [*Serratia*]	1	1	1	1015	112.7	7.14	2.7
1340368250	glycine dehydrogenase (aminomethyl-transferring), partial [*Serratia* sp. SSNIH2]	4	1	1	246	26.4	8.73	2.45
1047638873	hypothetical protein AN660_0220605 [*Serratia marcescens* subsp. *marcescens*]	2	1	1	718	81.3	5	2.45
1000823267	Translation elongation factor Tu [*Serratia rubidaea*]	2	1	1	394	43.2	5.31	2.3
1047650606	phosphoenolpyruvate carboxykinase (ATP) [*Serratia marcescens* subsp. *marcescens*]	7	4	4	539	59.3	5.5	1.97
1001786477	LysR family transcriptional regulator [*Serratia rubidaea*]	4	1	1	306	33.2	9	1.92
1006552443	DNA-directed RNA polymerase subunit beta' [*Serratia ficaria*]	3	3	3	1408	155	6.74	1.78
1195419500	short-chain dehydrogenase/reductase SDR [*Serratia marcescens*]	2	1	1	297	31.7	7.15	1.71

363988229	succinyl-CoA synthetase alpha subunit [*Serratia symbiotica* str. *'Cinara cedri'*]	3	1	1	291	30.5	7.72	1.7
640696690	MULTISPECIES: phosphorylcholine phosphatase [Serratia]	2	1	1	353	39.8	6.58	1.69
1048498251	Serralysin precursor [*Serratia marcescens*]	9	3	3	487	52.2	4.89	1.67
408489453	Chain A, 1 Chitobiase	6	4	5	858	95.7	6.57	1.64
977285585	Chitobiase precursor [*Serratia marcescens*]	8	5	6	885	98.3	6.52	1.64
948122997	acyl-CoA dehydrogenase [*Serratia* sp. Leaf51]	1	1	1	822	90.2	7.06	1.64

Supplementary Table 2. LC-MS analysis of 55 kDa protein band showing the NCBI

accession number, protein description, percentage of the protein sequence covered by

identified peptides (% sequence coverage), the number of identified peptides (# Peptides),

total number of identified peptide spectra matched for each identified protein (#PSM),

number of amino acid residues in each identified protein (#AA), molecular weight of proteins

(MW), calculated pI of proteins (Calc. pI) and the identification score of proteins calculated

by Sequest HT algorithms (Protein score).

Accession number	Protein description	% Sequence coverage	# Peptides	# PSMs	# AAs	MW [kDa]	Calc. pI	Protein score
1048498251	Serralysin precursor [*Serratia marcescens*]	36	15	76	487	52.2	4.89	166.7

1205048185	serine 3-dehydrogenase [*Serratia nematodiphila*]	31	13	43	472	50.4	4.75	96.43
452850059	insecticidal serralysin-like protein, partial [*Serratia marcescens*]	52	14	69	320	35	5.33	144.33
1333426919	serine 3-dehydrogenase [*Serratia marcescens*]	25	12	31	472	50.4	4.74	71.38
1268850421	serine 3-dehydrogenase [*Serratia* sp. BW106]	18	8	29	487	51.9	4.82	67.01
1122396880	DUF4214 domain-containing protein [*Serratia marcescens*]	4	3	12	1000	101	4.87	39.48
1047617935	arginine--tRNA ligase [*Serratia marcescens* subsp. marcescens]	22	12	12	576	63.9	5.55	18.71
1039033801	succinylarginine dihydrolase [*Serratia marcescens*]	27	12	12	446	49.4	6.77	11.96
1081143318	cell surface protein [*Serratia* sp. HMSC15F11]	10	6	8	998	100.9	4.79	7.17
1334464241	histidine ammonia-lyase [*Serratia marcescens*]	19	7	8	514	54.9	6	12.68
1127886605	Proline dipeptidase [*Serratia marcescens* SMB2099]	18	7	9	444	50.2	6.38	11.32
977423541	Extracellular serine protease precursor [*Serratia marcescens*]	12	8	9	1034	107.8	5.39	4.24
1280179082	capsule assembly Wzi family protein [*Serratia* sp. TKO39]	17	6	6	478	52.7	5.41	6.61
1027970958	Xaa-Pro aminopeptidase [*Serratia marcescens*]	11	4	5	438	49	5.38	7.85

1027968317	glutathione reductase [*Serratia marcescens*]	16	5	5	450	48.5	6.38	6.32
1001783318	phosphoenolpyruvate carboxykinase (ATP) [*Serratia rubidaea*]	5	2	2	539	59	5.25	1.95
1216656901	flagellin [Serratia marcescens]	11	4	5	423	44.4	5.24	1.99
493760228	F0F1 ATP synthase subunit beta [*Serratia symbiotica*]	7	2	2	460	50.2	5.12	2.55
1141208419	serine 3-dehydrogenase [*Serratia oryzae*]	6	3	7	522	55.7	4.54	11.77
1028171175	hypothetical protein [*Serratia* sp. YD25]	4	2	2	568	60.8	4.65	2.17
676309841	N-succinylglutamate 5-semialdehyde dehydrogenase [*Serratia* sp. SCBI]	2	1	1	639	68.5	9.8	2.13
383294472	glucose-6-phosphate isomerase [*Serratia* sp. M24T3]	4	2	2	549	61.2	6.09	1.64
1000823334	Secreted alkaline metalloproteinase [*Serratia rubidaea*]	4	2	6	497	53.2	4.7	9.45
1057964136	chitinase [*Serratia* sp. YD25]	3	2	2	480	51.8	5.62	2
1195419500	short-chain dehydrogenase/reductase SDR [*Serratia marcescens*]	2	1	3	297	31.7	7.15	5.5
1142792582	serine 3-dehydrogenase [*Serratia* sp. S119]	3	1	1	479	52	4.81	1.8

1006546552	NADP-dependent isocitrate dehydrogenase [*Serratia ficaria*]	3	1	1	417	45.6	5.68	2.03

3.7 Protease inhibitors reduced larvicidal activity of S. marcescens spent medium.

The LC-MS results indicated that *S. marcescens* secreted several proteases when

cultured at 25 ºC under a static condition. Some of these proteases have been reported for

insecticidal activity in other insects (Baumann, 2013; Wang and Granados, 1997). Therefore,

we added the Complete protease inhibitor (Roche) supplemented with 5 mM EDTA in the

larvicidal assays to determine whether these proteases contributed to *S. marcescens* larvicidal

activity. We found that protease inhibitors significantly reduced the three-day mortality of the

larvae to 50% as compared to 87.5% of the control group without protease inhibitors. This

suggests that the larvicidal activity of proteases was partially involved in the observed larval

mortality (Figure 6).

Figure 6. Protease inhibitor reduced larvicidal activity of *S. marcescens* secreted

macromolecule. The graphs represent data from four independent biological replicates.

Statistical significance of the survival analyses was calculated using Log-rank test followed by

a Bonferroni post-hoc analysis using 'survminer 'package in R.

3.8 Chitinase inhibitor reduced larvicidal activity of S. marcescens spent medium.

Because the LC-MS analysis suggested that our strain of *S. marcescens* expressed

chitinase and chitobiose enzymes when cultured under static condition at 25 ºC but not 37 ºC,

we then inoculated the bacterium on LB agar supplemented with 0.5% chitin then incubated

at 25 ºC or 37 ºC to determined chitinase secretion. After ten days of incubation, *S.*

marcescens cultured at 25 ºC produced a clear zone while the incubation at 37 ºC did not

(Figure 7A). Chitinase activity was also determined in the liquid static culture by pHBAH assay.

Chitinase activity of *S. marcescens* spent medium increased overtime during the incubation

under static condition at 25 ºC and saturated after 3 days of incubation (Figure 7B). This result

coincides well with the increase in larvicidal activity during the first three days of *S.*

marcescens incubation (Figure 1B).

Cyclic dipeptides such as Cyclo-Gly-Pro have previously been reported to inhibit

chitinase enzyme activity (Houston et al., 2004). This compound was then used to confirm that

the larvicidal activity was due to chitinase secretion. The results in Figure 6C show that Cyclo-

Gly-Pro can inhibit chitinase activity of *S. marcescens* spent medium in a concentration-

dependent manner. At 8 mM, Cyclo-Gly-Pro inhibited chitinase activity by 53.36±12.22%,

which was not significantly different from 56.81±8.34% inhibition by 10 mM Cyclo-Gly-Pro

(Figure 7C). Thus, 8 mM Cyclo-Gly-Pro was used in the following larvicidal assays. Larvicidal

activity was significantly reduced when 8 mM Cyclo-Gly-Pro was added to the spent medium

compared to the group without the chitinase inhibitor, albeit not to the same level as the LB

control (Figure 7D). This might be due to the incomplete inhibition of the chitinase activity or

other enzymes might be contributing to the larvicidal activity in addition to chitinases.

Figure 7. *S. marcescens* secreted macromolecules have chitinase activity and chitinase

inhibitor reduced larvicidal activity. (A) *S. marcescens* secreted chitinase that resulted in a

clear zone around bacterium colony when cultured at 25 ºC but not 37 ºC. (B) *S. marcescens*

spent medium accumulated chitinase activity over time and saturated after three days of

incubation. One unit of chitinase activity was defined as the amount of enzyme which releases

1 µg of reducing sugar per hour as measured by pHBAH assay. (C) Cyclo-Gly-Pro inhibit

chitinase activity in a concentration-dependent manner. Statistical analyses comparing

chitinase activity and inhibition of chitinase activity were performed using One-way ANOVA

with Tukey multiple comparison test using multcompView package in R. (D) Chitinase inhibitor

reduced larvicidal activity of the *S. marcescens* secreted macromolecule. The graphs

represent data from two independent biological replicates. Statistical significance of the

survival analyses was calculated using Log-rank test followed by a Bonferroni post-hoc

analysis using 'survminer 'package in R.

3.9 Combination of protease inhibitors and chitinase inhibitor could not completely inhibit

larvicidal activity.

Since both the protease inhibitors and chitinase inhibitors could partially reduced

larvicidal activity, the combination of both inhibitors rewas used to determine if simultaneous

inhibition of proteases and chitinases could completely inhibit the larvicidal activity (Figure 8).

The combination of protease and chitinase inhibitors significantly reduced larvae mortality to

33.33% from 93.33% in the no inhibitor control. Although not significantly different, the

combination of protease and chitinase inhibitors has a trend to further reduce larvicidal activity

compared to the treatment with single class of inhibitors (56.66% in the protease inhibitor

treatment and 48.33% in the chitinase inhibitor treatment).

Figure 8: Combination of protease inhibitor and chitinase inhibitor could not completely inhibit

larvicidal activity. The graph represents data from three independent biological replicates.

Statistical significance of the survival analyses was calculated using Log-rank test followed by

a Bonferroni post-hoc analysis using 'survminer 'package in R.

4. Discussion

In this study, a strain of non-prodigiosin producing *S. marcescens* was identified and

found to secrete molecules with larvicidal activity against *Anopheles dirus*, a malaria vector in

Southeast Asia. *S. marcescens* is a Gram-negative bacterium commonly found in soil and

water but not as a human fecal microbiome (Donnenberg, 2015). With such natural habitats,

it is often associated with insects including mosquitoes (Gendrin and Christophides, 2013;

Wang et al., 2017, 2011). Interestingly, different strains of *S. marcescens* can have different

relationship with *Anopheles* mosquitoes raging from symbiotic (Wang et al., 2017) to

entomopathogenic (Bahia et al., 2014). The cells of *S. marcescens* isolated in the present

study did not kill *An. dirus* larvae directly, but instead can secrete larvicidal molecules into the

culture medium when cultured under static condition for at least 2-3 days. The results were in

accordance with our original observations that the larval death occurred a few days after larval

water became turbid. The fact that the larvicidal molecules were secreted is an advantage for

application purposes, given that the secreted factors can be applied instead of the bacterium.

Thus preventing or limiting the risk of nosocomial infections (Golemi-Kotra, 2008). The

secreted factors can also be expressed heterologously in other expression systems that are

more suited for industrial scale production.

S. marcescens has been extensively studied for its potential as insect pest control

agent in agriculture due to its insecticidal property (Bidari et al., 2017; Li et al., 2011; Miller et

al., 2008; Pineda-Castellanos et al., 2015; Tu et al., 2010). The bacterium was known to

secrete many insect virulence factors including several enzymes, as well as secondary

metabolite including a red pigment prodigiosin. Prodigiosin is a metabolite possessing several

bioactivities including antimalarial and insecticidal/mosquito larvicidal properties (Bidari et al.,

2017; Darshan and Manonmani, 2015; Patil et al., 2011; Suryawanshi et al., 2015). The fact

that our *S. marcescens* isolate doesn't produce red pigment suggested that the bacterium

secretes other mosquito larvicidal molecules. This finding is in line with a previous study

reporting that level of prodigiosin production did not correlate with *S. marcescens* insecticidal

activity against silk worm larva (Zhou et al., 2016).

The heat-labile larvicidal macromolecules were secreted only when *S. marcescens*

was cultured under static condition at lower temperature of 25 ºC but not 37 ºC nor at 25 ºC

under planktonic condition. These results suggested that the metabolic activity of this

bacterium was influenced by culture conditions, consequently affecting the subsets of

molecules secreted by this strain of *S. marcescens*. Comparison of protein expression profiles

between the 25 ºC and 37 ºC static cultures helped us to identify potential proteins with

larvicidal activity which include many proteases and chitinases. Protease and chitinase

activities were then associated with larvicidal activity against *An. dirus* as supplementation of

chitinase or protease inhibitors diminished the larvicidal activity. These proteases and

chitinases are likely to damage the larvae at physiological levels rather than at the cellular

level because the secreted macromolecules lack cytotoxicity against mosquito cells.

Serralysin, one of the proteases identified from the mass spectrometry analysis of *S.*

marcescens secreted macromolecules, has previously been extensively characterized.

Serralysin is a metalloprotease first discovered in the culture medium of *Serratia* sp. (Miyata

et al., 1970) then later found in other Gram negative bacteria (Baumann, 2013). The protease

is secreted as an inactive enzyme and autocatalytically activated in the presence of divalent

cations (Baumann, 2013). The enzyme was reported to have entomopathogenic property

against larvae of scarab beetle (Pineda-Castellanos et al., 2015). Interestingly, serralysin was

found to have immunosuppression property in silk worm and considered as a virulence factor

(Ishii et al., 2014). The mechanism of immunosuppression in the study was through the

degradation of hemocyte adhesion molecule. Another study demonstrated that a serralysin-

type metalloprotease could cleave other proteins involved in insect immunity found in

Manduca sexta hemolymph (Felföldi et al., 2009)**.** The immune suppression by serralysin

could be one of the underlying mechanisms by which *An. dirus* larvae were rendered more

susceptible to bacterial infection.

One of the proteases that was detected in our LC-MS analysis is Peptidase M60

enhancin protein, which is a metalloprotease that requires zinc for its function. Enhancin was

first described in baculovirus (Hashimoto et al., 1991) then later found in several bacteria and

fungi (Nakjang et al., 2012; Slavicek, 2012). Mucin is an important protein component of insect

peritrophic matrix including *Anopheles* mosquito (Dinglasan et al., 2009). Enhancin from

baculoviruses could degrade intestinal mucin of the insects thus making them more

susceptible to viral infection (Toprak et al., 2012; Wang and Granados, 1997) and Bt toxin

(Granados et al., 2001). Bacterial enhancin could also digest mucin (Nakjang et al., 2012), but

the impact on baculovirus and Bt toxin toxicity was inconclusive (Fang et al., 2009; Galloway

et al., 2005; Zhao et al., 2011).

Our results indicate that chitinase and protease inhibitors could only partially reduce

the larvicidal activity of the spent medium against *An. dirus* larvae. This suggests that the

effect was likely to be a result from a combination of both chitinase and protease enzymes.

Although larvicidal activity caused by macromolecules was further reduced when protease

inhibitors and chitinase inhibitors were used in combination, the combinatorial effect was

marginal and could not completely inhibited the larvicidal activity. Therefore, it is possible that

macromolecules other than proteases and chitinases also contributed to the larvicidal activity.

This could also be explained by the incomplete inhibition of proteases and chitinases by the

inhibitors. As discussed above, several previous studies suggested that larvicidal activity of

the identified enzymes were marginal on their own. Contribution of each individual enzyme on

larvicidal activity should therefore be further investigated. This can be done by heterologous

expression of each enzyme to determine lethal concentrations individually and in combination.

Chitin, a polymer of *N*-acetylglucosamine, is the major component of mosquito

extracellular structures including the cuticle of the exoskeleton and the peritrophic matrix (PM)

of the midgut in arthropods (Liu et al., 2019). These exoskeletal structures provide strength to

the body structure, and serve as muscle attachment sites which allow locomotion and flight

(Liu et al., 2019). They are also important barriers for the protection of arthropods from the

surrounding environment, mechanical damage, heat, desiccation, and invasion of

microorganisms (Liu et al., 2019). With all these important functions, damage to the

exoskeleton can be fatal to mosquitoes. Several techniques have been developed to

compromise the insects' exoskeleton including inhibition of chitin synthesis (Belinato et al.,

2013; Doucet and Retnakaran, 2012; Zhang et al., 2010), and degradation of the chitin

structure (Berini et al., 2019; Singh and Arya, 2019). The later strategy has been exploited

using chitinase enzymes or chitinase-producing microorganisms including those from *S.*

marcescens (Aggarwal et al., 2015; Binod et al., 2007). However, many studies suggested

that chitinases might not be ideal to be used as a stand-alone insecticide for some insects

because the larvicidal activity from chitinase can be marginal (Bahar et al., 2012; Zhong et al.,

2015). Rather, the enzymes could be used to enhance the insecticidal activity of other control

agents including Bt toxin, entomopathogenic virus and chemical insecticides (Kelkenberg et

al., 2015; Ramírez-Suero et al., 2011; Zhong et al., 2015). Previous research has

demonstrated that chitinases can also help overcome any resistance phenotype developed

against a biocontrol strategy. Cai et al. showed that *Bacillus sphaericus* (Bs) toxin-resistant

Culex quinquefasciatus became susceptible to the Bs treatment when Chitinase AC was

overexpressed in *B. sphaericus* (Cai et al., 2007). In addition to the insecticidal property,

chitinases can also inhibit fungus by destroying chitinous cell wall structure (Berini et al., 2017;

Singh and Arya, 2019; Zarei et al., 2011). *S. marcescens* has been tested to be used against

plant fungal pathogen due to its fungicidal properties (Purkayastha et al., 2018) making it an

interesting candidate to be used against several classes of agricultural diseases.

Peritrophic matrix (PM) is an acellular structure inside the mosquito gut that

compartmentalize gut content from the midgut epithelium (Richards and Richards, 1977;

Terra, 2001). It serves as a barrier to protect the insect from the digestive enzymes, abrasion

from food particles, and invading parasites (Terra, 2001). PM structure consist of chitin and

proteins network, with peritrophins as major protein components (Terra, 2001). Several

previous studies have shown the insecticidal activity of microbial chitinases and proteases,

and their ability to degrade the exoskeleton and perithrophic matrix (Berini et al., 2019, 2018;

Chang et al., 2013; Li et al., 2008; Singh and Arya, 2019). Thus, the larvicidal acticity of our

Serratia culture-derived fractions in combination to the reduction of larvicidal activity when

exposed to chitinase and protease inhibitors hints that this is one if the underlying mechanisms

of larval killing. However, more histological-based studies would be needed to concretely

confirm this assertion. In addition to direct killing, the compromised peritrophic matrix can be

responsible for enhanced susceptibility to bacterial infection observed in our study. For

instance, in *Drosophila*, disruption of the peritrophic matrix formation renders the fruit fly more

susceptible to bacterial infection (Buchon et al., 2013). The disruption of PM may be the

underlying mechanism of the increased larvae's susceptibility to *Pseudomonas* sp. after *S.*

marcescens spent medium treatment.

Since chitin and protein are fundamental building blocks of several organisms, the

secreted larvicidal chitinase and proteases can have unintended effects to non-target

organisms in the same ecological niches. When chitinases are used for agricultural insect pest

management, the enzymes can have additional benefit in fungal disease control as chitin also

serve as exoskeleton of fungi (Singh and Arya, 2019). However, *An. dirus* used in this study

reside in forestry areas (Obsomer et al., 2007). The application of non-specific chitinase and

protease larvicides in such area can affect biodiversity in the forestry environment. On the

contrary to the *An. dirus* in Southeast Asia, several reports found adaptation of *Anopheline*

mosquitoes in urban environment in Africa (Mattah et al., 2017; Takken and Lindsay, 2019).

These urban *Anopheline* mosquitoes might be better targets for the protease- or chitinase-

based insecticide due to less potential impact on biodiversity but the efficiency of larvicidal

activity has to be confirmed in these species. The shorter duration of the larvicidal effect from

enzymatic-based larvicidal molecules suggests environmentally friendly alternative to long-

lasting chemical pesticides (Winding et al., 2004).

In summary, this study has identified a strain of *S. marcescens* from a larval rearing

habitat in the insectary, that secretes chitinases and proteases that are pathogenic to *An. dirus*

larvae. In addition to their intrinsic toxicity, these molecules have potential to increase the

larvae's susceptibility to natural entomopathogenic microorganisms, and other mosquito

control strategies currently in use. Although the secreted larvicidal proteases and chitinases

have potential for *Anopheles* mosquito control, further studies on concentration optimization,

formulation of enzyme mixtures, and enzyme modifications to enhance heat stability are

needed for its development into field applications.

Funding

This work was supported by the National Center for Genetic Engineering and Biotechnology

(P1751424) given to N. Jupatanakul. The funders had no role in study design, data collection

and interpretation.

Author contributions

Natapong Jupatanakul: Conceptualization, Methodology, Validation, Formal analysis,

Investigation, Resources, Writing - Original Draft, Writing - Review & Editing, Visualization,

Supervision, Project administration, Funding acquisition

Jutharat Pengon: Methodology, Validation, Formal analysis, Investigation, Writing - Review

& Editing

Shiela Marie Gines Selisana: Methodology, Validation, Formal analysis, Investigation,

Writing - Review & Editing

Waeowalee Choksawangkarn: Methodology, Validation, Formal analysis, Investigation,

Writing - Original Draft, Writing - Review & Editing

Nongluck Jaito: Methodology, Validation, Formal analysis, Investigation, Writing - Original

Draft

Atiporn Saeung: Methodology, Resources, Writing - Review & Editing

Ratchanu Bunyong: Investigation

Navaporn Posayapisit: Investigation

Khrongkhwan Thammatinna: Investigation

Nuttiya Kalpongnukul: Investigation

Kittipat Aupalee: Investigation

Trairak Pisitkun: Methodology, Resources,

Sumalee Kamchonwongpaisan: Resources, Writing - Review & Editing

Declaration of Competing Interest

All authors declare that we have no conflict of interest, financial or otherwise.

Acknowledgements

We thank Jose Luis Ramirez (USDA) for the suggestions and editorial help with the

manuscript. We are grateful to Asst. Prof. Dr. Patchara Sriwichai from Department of Medical

Entomology, Faculty of Tropical Medicine, Mahidol University, who kindly provided *An. dirus*

adult used to establish the colonies used in this study to A. Saeung.

References:

Aggarwal, C., Paul, S., Tripathi, V., Paul, B., Khan, Md.A., 2015. Chitinolytic activity in *Serratia*

marcescens (strain SEN) and potency against different larval instars of Spodoptera litura with

effect of sublethal doses on insect development. BioControl 60, 631–640.

https://doi.org/10.1007/s10526-015-9674-3

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search

tool. Journal of Molecular Biology 215, 403–410. https://doi.org/10.1016/S0022-

2836(05)80360-2

Bahar, A.A., Sezen, K., Demirbağ, Z., Nalçacioğlu, R., 2012. The relationship between insecticidal

effects and chitinase activities of Coleopteran-originated entomopathogens and their

chitinolytic profile. Ann Microbiol 62, 647–653. https://doi.org/10.1007/s13213-011-0301-y

Bahia, A.C., Dong, Y., Blumberg, B.J., Mlambo, G., Tripathi, A., BenMarzouk-Hidalgo, O.J., Chandra,

R., Dimopoulos, G., 2014. Exploring *Anopheles* gut bacteria for *Plasmodium* blocking activity.

Environmental Microbiology 16, 2980–2994. https://doi.org/10.1111/1462-2920.12381

Baumann, P., Clark, M.A., Baumann, L., Broadwell, A.H., 1991. *Bacillus sphaericus* as a mosquito

pathogen: properties of the organism and its toxins. Microbiology and Molecular Biology

Reviews 55, 425–436.

Baumann, U., 2013. Serralysin and Related Enzymes, in: Handbook of Proteolytic Enzymes. Elsevier,

pp. 864–867.

Belinato, T.A., Martins, A.J., Lima, J.B.P., Valle, D., 2013. Effect of triflumuron, a chitin synthesis

inhibitor, on *Aedes aegypti, Aedes albopictus* and *Culex quinquefasciatus* under laboratory

conditions. Parasites & Vectors 6, 83. https://doi.org/10.1186/1756-3305-6-83

Berini, F., Casartelli, M., Montali, A., Reguzzoni, M., Tettamanti, G., Marinelli, F., 2019. Metagenome-

Sourced Microbial Chitinases as Potential Insecticide Proteins. Front. Microbiol. 10.

https://doi.org/10.3389/fmicb.2019.01358

Berini, F., Katz, C., Gruzdev, N., Casartelli, M., Tettamanti, G., Marinelli, F., 2018. Microbial and viral

chitinases: Attractive biopesticides for integrated pest management. Biotechnology Advances,

Prospects in Biotechnology 36, 818–838. https://doi.org/10.1016/j.biotechadv.2018.01.002

Berini, F., Presti, I., Beltrametti, F., Pedroli, M., Vårum, K.M., Pollegioni, L., Sjöling, S., Marinelli, F.,

2017. Production and characterization of a novel antifungal chitinase identified by functional

screening of a suppressive-soil metagenome. Microbial Cell Factories 16, 16.

https://doi.org/10.1186/s12934-017-0634-8

Bidari, F., Shams-Bakhsh, M., Mehrabadi, M., 2017. Isolation and characterization of a *Serratia*

marcescens with insecticidal activity from *Polyphylla olivieri* (Col.: Scarabaeidae) 142, 162–

172. https://doi.org/10.1111/jen.12421

Binod, P., Sukumaran, R.K., Shirke, S.V., Rajput, J.C., Pandey, A., 2007. Evaluation of fungal culture

filtrate containing chitinase as a biocontrol agent against *Helicoverpa armigera*. Journal of

Applied Microbiology 103, 1845–1852. https://doi.org/10.1111/j.1365-2672.2007.03428.x

Buchon, N., Osman, D., David, F.P.A., Fang, H.Y., Boquete, J.-P., Deplancke, B., Lemaitre, B., 2013.

Morphological and Molecular Characterization of Adult Midgut Compartmentalization in

Drosophila. CellReports 3, 1725–1738. https://doi.org/10.1016/j.celrep.2013.04.001

Cai, Y., Yan, J., Hu, X., Han, B., Yuan, Z., 2007. Improving the Insecticidal Activity against Resistant

Culex quinquefasciatus Mosquitoes by Expression of Chitinase Gene *chiAC* in *Bacillus*

sphaericus. Applied and Environmental Microbiology 73, 7744–7746.

https://doi.org/10.1128/aem.01510-07

Caragata, E.P., Otero, L.M., Carlson, J.S., Dizaji, N.B., Dimopoulos, G., 2020. A non-live preparation

of *Chromobacterium* sp. Panama (Csp_P) is a highly effective larval mosquito biopesticide.

Appl. Environ. Microbiol. https://doi.org/10.1128/AEM.00240-20

Chang, Y.-T., Hsieh, C., Wu, L.-C., Chang, H.C., Kao, S.-S., Meng, M., Hsieh, F.-C., 2013.

Purification and Properties of an Insecticidal Metalloprotease Produced by *Photorhabdus*

luminescens Strain 0805-P5G, the Entomopathogenic Nematode Symbiont. International

journal of molecular sciences 14, 308–321. https://doi.org/10.3390/ijms14010308

Choochote, W., Saeung, A., 2013. Systematic Techniques for the Recognition of *Anopheles* Species

Complexes, in: Manguin, S. (Ed.), *Anopheles* Mosquitoes - New Insights into Malaria Vectors.

InTech. https://doi.org/10.5772/54853

Chouaia, B., Rossi, P., Epis, S., Mosca, M., Ricci, I., Damiani, C., Ulissi, U., Crotti, E., Daffonchio, D.,
757   Bandi, C., Favia, G., 2012. Delayed larval development in *Anopheles* mosquitoes deprived of
758   *Asaia* bacterial symbionts. BMC Microbiology 12, S2. https://doi.org/10.1186/1471-2180-12-
759   S1-S2

760 Coon, K.L., Vogel, K.J., Brown, M.R., Strand, M.R., 2014. Mosquitoes rely on their gut microbiota for
761   development. Molecular Ecology 23, 2727–2739. https://doi.org/10.1111/mec.12771

762 Darshan, N., Manonmani, H.K., 2015. Prodigiosin and its potential applications. Journal of Food
763   Science and Technology 52, 5393–5407. https://doi.org/10.1007/s13197-015-1740-4

764 Dickson, L.B., Jiolle, D., Minard, G., Moltini-Conclois, I., Volant, S., Ghozlane, A., Bouchier, C., Ayala,
765   D., Paupy, C., Moro, C.V., Lambrechts, L., 2017. Carryover effects of larval exposure to
766   different environmental bacteria drive adult trait variation in a mosquito vector. Science
767   Advances 3, e1700585. https://doi.org/10.1126/sciadv.1700585

768 Dinglasan, R., Devenport, M., Florens, L., Johnson, J., McHugh, C., Donnelly-Doman, M., Carucci, D.,
769   Yates, J., Jacobs-Lorena, M., 2009. The *Anopheles gambiae* Adult Midgut Peritrophic Matrix
770   Proteome. Insect Biochem Mol Biol 39, 125–134. https://doi.org/10.1016/j.ibmb.2008.10.010

771 Donnenberg, M.S., 2015. Enterobacteriaceae, in: Principles and Practice of Infectious Diseases. p.
772   2503.

773 Doucet, D., Retnakaran, A., 2012. Insect Chitin: Metabolism, Genomics and Pest Management, in:
774   Dhadialla, T.S. (Ed.), Advances in Insect Physiology, Insect Growth Disruptors. Academic
775   Press, pp. 437–511. https://doi.org/10.1016/B978-0-12-391500-9.00006-1

776 Eden, P.A., Schmidt, T.M., Blakemore, R.P., Pace, N.R., 1991. Phylogenetic Analysis of
777   *Aquaspirillum magnetotacticum* Using Polymerase Chain Reaction-Amplified 16S rRNA-
778   Specific DNA. International Journal of Systematic Bacteriology 41, 324–325.
779   https://doi.org/10.1099/00207713-41-2-324

780 Eng, J.K., McCormack, A.L., Yates, J.R., 1994. An approach to correlate tandem mass spectral data
781   of peptides with amino acid sequences in a protein database. J Am Soc Mass Spectr 5, 976–
782   89. https://doi.org/10.1016/1044-0305(94)80016-2

783 Fang, S., Wang, L., Guo, W., Zhang, X., Peng, D., Luo, C., Yu, Z., Sun, M., 2009. *Bacillus*
784   *thuringiensis* Bel Protein Enhances the Toxicity of Cry1Ac Protein to *Helicoverpa armigera*

Larvae by Degrading Insect Intestinal Mucin. Appl. Environ. Microbiol. 75, 5237–5243.

https://doi.org/10.1128/AEM.00532-09

Felföldi, G., Marokházi, J., Képiró, M., Venekei, I., 2009. Identification of Natural Target Proteins

Indicates Functions of a Serralysin-Type Metalloprotease, PrtA, in Anti-Immune Mechanisms.

Appl. Environ. Microbiol. 75, 3120–3126. https://doi.org/10.1128/AEM.02271-08

Galloway, C.S., Wang, P., Winstanley, D., Jones, I.M., 2005. Comparison of the bacterial Enhancin-

like proteins from *Yersinia* and *Bacillus* spp. with a baculovirus Enhancin. Journal of

Invertebrate Pathology 90, 134–137. https://doi.org/10.1016/j.jip.2005.06.008

Gendrin, M., Christophides, G.K., 2013. The *Anopheles* Mosquito Microbiota and Their Impact on

Pathogen Transmission, in: Manguin, S. (Ed.), Anopheles Mosquitoes - New Insights into

Malaria Vectors. InTech. https://doi.org/10.5772/55107

Golemi-Kotra, D., 2008. *Serratia, Edwardsiella* and *Morganella* Infections, in: Enna, S.J., Bylund, D.B.

(Eds.), XPharm: The Comprehensive Pharmacology Reference. Elsevier, New York, pp. 1–6.

https://doi.org/10.1016/B978-008055232-3.63827-9

Granados, R.R., Fu, Y., Corsaro, B., Hughes, P.R., 2001. Enhancement of *Bacillus thuringiensis*

Toxicity to *Lepidopterous* Species with the Enhancin from *Trichoplusia* ni Granulovirus.

Biological Control 20, 153–159. https://doi.org/10.1006/bcon.2000.0891

Graves, S., Dorai-Raj, H.-P.P. and L.S. with help from S., 2019. multcompView: Visualizations of

Paired Comparisons.

Hashimoto, Y., Corsaro, B.G., Granados, R.R., 1991. Location and nucleotide sequence of the gene

encoding the viral enhancing factor of the *Trichoplusia* ni granulosis virus. Journal of General

Virology 72, 2645–2651. https://doi.org/10.1099/0022-1317-72-11-2645

Houston, D.R., Synstad, B., Eijsink, V.G.H., Stark, M.J.R., Eggleston, I.M., Aalten, D.M.F. van, 2004.

Structure-based exploration of cyclic dipeptide chitinase inhibitors. 47, 5713–5720.

https://doi.org/10.1021/jm049940a

Ishii, K., Adachi, T., Hamamoto, H., Sekimizu, K., 2014. *Serratia marcescens* Suppresses Host

Cellular Immunity via the Production of an Adhesion-inhibitory Factor against

Immunosurveillance Cells. J. Biol. Chem. 289, 5876–5888.

https://doi.org/10.1074/jbc.M113.544536

Kassambara, A., Kosinski, M., Biecek, P., 2019. survminer: Drawing Survival Curves using "ggplot2"

815   [WWW Document]. URL https://CRAN.R-project.org/package=survminer

816   Kelkenberg, M., Odman-Naresh, J., Muthukrishnan, S., Merzendorfer, H., 2015. Chitin is a necessary

817   component to maintain the barrier function of the peritrophic matrix in the insect midgut.

818   Insect Biochemistry and Molecular Biology 56, 21–28.

819   https://doi.org/10.1016/j.ibmb.2014.11.005

820   Laurence, D., Christophe, L., Roger, F., 2011. Using the Bio-Insecticide *Bacillus Thuringiensis*

821   Israelensis in Mosquito Control, in: Stoytcheva, M. (Ed.), Pesticides in the Modern World -

822   Pests Control and Pesticides Exposure and Toxicity Assessment. InTech.

823   https://doi.org/10.5772/17005

824   Lever, M., 1972. A new reaction for colorimetric determination of carbohydrates. Analytical

825   Biochemistry 47, 273–279. https://doi.org/10.1016/0003-2697(72)90301-6

826   Li, B., Yu, R., Liu, B., Tang, Q., Zhang, G., Wang, Y., Xie, G., Sun, G., 2011. Characterization and

827   Comparison of *Serratia marcescens* Isolated from Edible Cactus and from Silkworm for

828   Virulence Potential and Chitosan Susceptibility 42, 96–104. https://doi.org/10.1590/s1517-

829   83822011000100013

830   Li, H., Tang, H., Sivakumar, S., Philip, J., Harrison, R.L., Gatehouse, J.A., Bonning, B.C., 2008.

831   Insecticidal activity of a basement membrane-degrading protease against *Heliothis virescens*

832   (Fabricius) and *Acyrthosiphon pisum (*Harris). Journal of Insect Physiology 54, 777–789.

833   https://doi.org/10.1016/j.jinsphys.2008.02.008

834   Liu, X., Zhang, J., Zhu, K.Y., 2019. Chitin in Arthropods: Biosynthesis, Modification, and Metabolism,

835   in: Yang, Q., Fukamizo, T. (Eds.), Targeting Chitin-Containing Organisms, Advances in

836   Experimental Medicine and Biology. Springer, Singapore, pp. 169–207.

837   https://doi.org/10.1007/978-981-13-7318-3_9

838   Manguin, S., Garros, C., Dusfour, I., Harbach, R.E., Coosemans, M., 2008. Bionomics, taxonomy,

839   and distribution of the major malaria vector taxa of *Anopheles* subgenus Cellia in Southeast

840   Asia: An updated review. Infection, Genetics and Evolution 8, 489–503.

841   https://doi.org/10.1016/j.meegid.2007.11.004

842   Mattah, P.A.D., Futagbi, G., Amekudzi, L.K., Mattah, M.M., de Souza, D.K., Kartey-Attipoe, W.D.,

843   Bimi, L., Wilson, M.D., 2017. Diversity in breeding sites and distribution of *Anopheles*

mosquitoes in selected urban areas of southern Ghana. Parasites & Vectors 10, 25.

https://doi.org/10.1186/s13071-016-1941-3

Miller, T.A., Lauzon, C.R., Lampe, D.J., 2008. Technological advances to enhance agricultural pest

management. 627, 141–150. https://doi.org/10.1007/978-0-387-78225-6_12

Miyata, K., Maejima, K., Tomoda, K., Isono, M., 1970. *Serratia* Protease. Agricultural and Biological

Chemistry 34, 310–318. https://doi.org/10.1080/00021369.1970.10859598

MMV-supported projects | Medicines for Malaria Venture [WWW Document], 2020. URL

https://www.mmv.org/research-development/mmv-supported-projects (accessed 4.5.20).

Nakjang, S., Ndeh, D.A., Wipat, A., Bolam, D.N., Hirt, R.P., 2012. A Novel Extracellular

Metallopeptidase Domain Shared by Animal Host-Associated Mutualistic and Pathogenic

Microbes. PLOS ONE 7, e30287. https://doi.org/10.1371/journal.pone.0030287

Obsomer, V., Defourny, P., Coosemans, M., 2007. The *Anopheles dirus* complex: spatial distribution

and environmental drivers. Malaria Journal 6, 26–16. https://doi.org/10.1186/1475-2875-6-26

Patil, C.D., Patil, S.V., Salunke, B.K., Salunkhe, R.B., 2011. Prodigiosin produced by *Serratia*

marcescens NMCC46 as a mosquito larvicidal agent against *Aedes aegypti* and *Anopheles*

stephensi. Parasitology Research 109, 1179–1187. https://doi.org/10.1007/s00436-011-2365-

9

Pineda-Castellanos, M.L., Rodríguez-Segura, Z., Villalobos, F.J., Hernández, L., Lina, L., Nuñez-

Valdez, M.E., 2015. Pathogenicity of Isolates of *Serratia marcescens* towards Larvae of the

Scarab *Phyllophaga blanchardi* (Coleoptera). Pathogens 4, 210–228.

https://doi.org/10.3390/pathogens4020210

Purkayastha, G.D., Mangar, P., Saha, A., Saha, D., 2018. Evaluation of the biocontrol efficacy of a

Serratia marcescens strain indigenous to tea rhizosphere for the management of root rot

disease in tea. PLOS ONE 13, e0191761. https://doi.org/10.1371/journal.pone.0191761

R Core Team, 2020. R: The R Project for Statistical Computing. R Foundation for Statistical

Computing, Vienna, Austria.

Ramirez, J.L., Short, S.M., Bahia, A.C., Saraiva, R.G., Dong, Y., Kang, S., Tripathi, A., Mlambo, G.,

Dimopoulos, G., 2014. *Chromobacterium Csp*_P Reduces Malaria and Dengue Infection in

Vector Mosquitoes and Has Entomopathogenic and In Vitro Anti-pathogen Activities. PLoS

Pathogens 10, e1004398-13. https://doi.org/10.1371/journal.ppat.1004398

Ramírez-Suero, M., Valerio-Alfaro, G., Bernal, J.S., Ramírez-Lepe, M., 2011. Synergisitic effect of

chitinases and *Bacillus thuringiensis* israelensis spore-toxin complex against *Aedes aegypti*

larvae. The Canadian Entomologist 143, 157–164. https://doi.org/10.4039/n10-051

Richards, A.G., Richards, P.A., 1977. The Peritrophic Membranes of Insects. Annu. Rev. Entomol. 22,

219–240. https://doi.org/10.1146/annurev.en.22.010177.001251

Scholefield, R.J., Slaght, K.S., Stephens, B.E., 2008. Seasonal Variation in Sensitivity of Larval Sea

Lampreys to the Lampricide 3-Trifluoromethyl-4-Nitrophenol. North American Journal of

Fisheries Management 28, 1609–1617. https://doi.org/10.1577/M06-178.1

Shaw, W.R., Catteruccia, F., 2019. Vector biology meets disease control: using basic research to fight

vector-borne diseases. Nature Microbiology 4, 20–34. https://doi.org/10.1038/s41564-018-

0214-7

Shevchenko, A., Tomas, H., Havli, J., Olsen, J.V., Mann, M., 2007. In-gel digestion for mass

spectrometric characterization of proteins and proteomes. Nat Protoc 1, 2856–2860.

https://doi.org/10.1038/nprot.2006.468

Short, S.M., Tol, S. van, MacLeod, H.J., Dimopoulos, G., 2018. Hydrogen cyanide produced by the

soil bacterium *Chromobacterium* sp. Panama contributes to mortality in *Anopheles gambiae*

mosquito larvae. Nature Publishing Group 8, 1–13. https://doi.org/10.1038/s41598-018-

26680-2

Singh, G., Arya, S.K., 2019. Antifungal and insecticidal potential of chitinases: A credible choice for

the eco-friendly farming. Biocatalysis and Agricultural Biotechnology 20, 101289.

https://doi.org/10.1016/j.bcab.2019.101289

Slavicek, J.M., 2012. Baculovirus Enhancins and Their Role in Viral Pathogenicity, in: Molecular

Virology. IntechOpen.

Suryawanshi, R.K., Patil, C.D., Borase, H.P., Narkhede, C.P., Salunke, B.K., Patil, S.V., 2015.

Mosquito larvicidal and pupaecidal potential of prodigiosin from *Serratia marcescens* and

understanding its mechanism of action. Pesticide Biochemistry and Physiology 123, 49–55.

https://doi.org/10.1016/j.pestbp.2015.01.018

Takken, W., Lindsay, S., 2019. Increased Threat of Urban Malaria from *Anopheles stephensi*

Mosquitoes, Africa. Emerg. Infect. Dis. 25, 1431–1433.

https://doi.org/10.3201/eid2507.190301

Terra, W.R., 2001. The origin and functions of the insect peritrophic membrane and peritrophic gel.

Arch. Insect Biochem. Physiol. 47, 47–61. https://doi.org/10.1002/arch.1036

Toprak, U., Harris, S., Baldwin, D., Theilmann, D., Gillott, C., Hegedus, D.D., Erlandson, M.A., 2012.

Role of enhancin in *Mamestra configurata* nucleopolyhedrovirus virulence: selective

degradation of host peritrophic matrix proteins. Journal of General Virology, 93, 744–753.

https://doi.org/10.1099/vir.0.038117-0

Tu, S., Qiu, X., Cao, L., Han, R., Zhang, Y., Liu, X., 2010. Expression and characterization of the

chitinases from Serratia marcescens GEI strain for the control of Varroa destructor, a honey

bee parasite 104, 75–82. https://doi.org/10.1016/j.jip.2010.02.002

Valzania, L., Martinson, V.G., Harrison, R.E., Boyd, B.M., Coon, K.L., Brown, M.R., Strand, M.R.,

2018. Both living bacteria and eukaryotes in the mosquito gut promote growth of larvae. PLoS

Neglected Tropical Diseases 12, e0006638. https://doi.org/10.1371/journal.pntd.0006638

Vichai, V., Kirtikara, K., 2006. Sulforhodamine B colorimetric assay for cytotoxicity screening. Nature

Protocols 1, 1112–1116. https://doi.org/10.1038/nprot.2006.179

Wang, P., Granados, R.R., 1997. An intestinal mucin is the target substrate for a baculovirus

enhancin. PNAS 94, 6977–6982. https://doi.org/10.1073/pnas.94.13.6977

Wang, S., Dos-Santos, A.L.A., Huang, W., Liu, K.C., Oshaghi, M.A., Wei, G., Agre, P., Jacobs-

Lorena, M., 2017. Driving mosquito refractoriness to *Plasmodium falciparum* with engineered

symbiotic bacteria. Science 357, 1399–1402. https://doi.org/10.1126/science.aan5478

Wang, Y., Iii, T.M.G., Kukutla, P., Yan, G., Xu, J., 2011. Dynamic Gut Microbiome across Life History

of the Malaria Mosquito *Anopheles gambiae* in Kenya. PLOS ONE 6, e24767.

https://doi.org/10.1371/journal.pone.0024767

Wells, T.N.C., van Huijsduijnen, R.H., Van Voorhis, W.C., 2015. Malaria medicines: a glass half full?

Nature Reviews Drug Discovery 14, 424–442. https://doi.org/10.1038/nrd4573

Whitty, C.J.M., Ansah, E., 2019. Malaria control stalls in high incidence areas. BMJ 365.

https://doi.org/10.1136/bmj.l2216

Winding, A., Binnerup, S.J., Pritchard, H., 2004. Non-target effects of bacterial biological control

agents suppressing root pathogenic fungi. FEMS Microbiol Ecol 47, 129–141.

https://doi.org/10.1016/S0168-6496(03)00261-7

Zarei, M., Aminzadeh, S., Zolgharnein, H., Safahieh, A., Daliri, M., Noghabi, K.A., Ghoroghi, A.,

Motallebi, A., 2011. Characterization of a chitinase with antifungal activity from a native

Serratia marcescens B4A. Braz J Microbiol 42, 1017–1029. https://doi.org/10.1590/S1517-

838220110003000022

Zhang, X., Zhang, J., Zhu, K.Y., 2010. Chitosan/double-stranded RNA nanoparticle-mediated RNA

interference to silence chitin synthase genes through larval feeding in the African malaria

mosquito (*Anopheles gambiae*). Insect Molecular Biology 19, 683–693.

https://doi.org/10.1111/j.1365-2583.2010.01029.x

Zhao, D., Guo, W., Sun, W., Xu, D., Liu, D., 2011. Identification of a novel enhancin-like gene from

Bacillus thuringiensis. Front. Agric. China 5, 423–429. https://doi.org/10.1007/s11703-011-

1117-z

Zhong, W., Ding, S., Guo, H., Zhong, W., Ding, S., Guo, H., 2015. The chitinase C gene PsChiC from

Pseudomonas sp. and its synergistic effects on larvicidal activity. Genetics and Molecular

Biology 38, 366–372. https://doi.org/10.1590/S1415-475738320140320

Zhou, W., Li, J., Chen, J., Liu, X., Xiang, T., Zhang, L., Wan, Y., 2016. The red pigment prodigiosin is

not an essential virulence factor in entomopathogenic *Serratia marcescens*. Journal of

Invertebrate Pathology 136, 92–94. https://doi.org/10.1016/j.jip.2016.03.011

950