coronapp: A Web Application to Annotate and Monitor

SARS-CoV-2 Mutations

Daniele Mercatelli[1,#], Luca Triboli[1], Eleonora Fornasari[1], Forest Ray[2], Federico M.

Giorgi[1,*]

[1] *Department of Pharmacy and Biotechnology, University of Bologna, Bologna,*

40126, Italy

[2] *Department of Systems Biology, Columbia University Medical Center, New York*

City, 10032, United States

[*] Corresponding author.

E-mail: federico.giorgi@unibo.it (Giorgi FM)

Running title: *Mercatelli D et al / coronapp – monitoring SARS-CoV-2 mutations*

Word number: 3529

Figure number: 3

Abstract

The avalanche of genomic data generated from the SARS-CoV-2 virus requires the

development of tools to detect and monitor its mutations across the world. Here, we

present a webtool, *coronapp*, dedicated to easily processing user-provided

SARS-CoV-2 genomic sequences and visualizing current worldwide status of

SARS-CoV-2 mutations.

The webtool allows users to highlight mutations and categorize them by frequency,

country, genomic location and effect on protein sequences, and to monitor their

presence in the population over time.

The tool is available at http://giorgilab.unibo.it/coronapp/ for the worldwide dataset

and at http://giorgilab.unibo.it/coronannotator/ for the annotation of user-provided

sequences.   The   full   code   is   freely   shared   at

https://github.com/federicogiorgi/giorgilab/tree/master/coronapp

KEYWORDS: COVID-19; SARS-CoV-2; mutations; genetics; web application

Data Availability Statement

The data that support the findings of this study derive from the GISAID consortium

and are openly available in Github , in Rdata format for the R environment, in files

results.rda   and   metadata.rda,   at   the   following   link:

https://github.com/federicogiorgi/giorgilab/tree/master/coronapp/data

1 Introduction

SARS-CoV-2 is a novel pathogenic enveloped RNA beta-coronavirus causing a
severe illness in human hosts known as coronavirus disease-2019 (COVID-19). The
predominant COVID-19 illness is a viral pneumonia, often requiring hospitalization
and in some cases intensive care [1]. With almost 27.5 million laboratory-confirmed
positive cases worldwide as of 9 September 2020 and an estimated case fatality rate
across 204 countries of 5.2%, COVID-19 has become a global health challenge in
only a few months [2]. SARS-CoV-2 infection depends on the recognition of host
angiotensin converting enzyme 2 (ACE2), exposed on the cell surface in human lung
tissues [3,4]. SARS-CoV-2 spike glycoprotein binds ACE2, mediating membrane
fusion and cell entry [5]. Upon cell entry, the virus subverts host cell molecular
processes, inducing interferon responses and eventually apoptosis [6].

To date, much effort has been made to develop therapeutic strategies to limit
SARS-CoV-2 transmission and replication, but no treatment or vaccine has proven
effective against the virus, and repurposing of approved therapeutic agents has been
the main practical approach to manage the emergency so far [7]. As viruses mutate
during replication, the emergence of SARS-CoV-2 sub-strains and the challenge of a
probable antigenic drift require attention, especially for vaccine development [8].

Although sequence analyses of SARS-CoV-2 have shown that genomic variability
is very low [9], new SARS-CoV-2 mutation hotspots are emerging due to the high
number of infected individuals across countries and to viral replication rates [10].
Three major SARS-CoV-2 clades known as clade G, V, and S have emerged, showing
a different geographical prevalence [10]. The most frequent mutation detected so far
defines the G clade and causes an aminoacidic change, aspartate (D) or glycine (G), at
position 614 (D614G) of the viral Spike protein [11].

Continual genomic surveillance should be considered to monitor the possible
appearance of viral subtypes characterized by altered tropism, or causing more
aggressive symptoms. Constant and widespread monitoring of mutations is also a
powerful means of informing drug development and global or local pandemic

management. The Global Initiative on Sharing All Influenza Data (GISAID) has

collected to date (9 September 2020) over 90,000 publicly accessible SARS-CoV-2

sequences. The GISAID effort has made it possible to compare genomes on a

geographical and temporal scale and an increasing number of laboratories have started

to sequence COVID-19 patient samples worldwide [12,13]. Several online tools have

been developed to monitor the evolution of the virus from a phylogenetic perspective,

such as Nextstrain [14], or to visualize epidemiological data such as number of cases

and deaths [15]. However, no online tool currently exists to annotate user-provided

SARS-CoV-2 genomic sequences, which may derive from specific GISAID subsets

or from sequencing efforts of individual laboratories. Neither does any tool

specifically monitor the prevalence of specific SARS-CoV-2 mutations associated to

particular geographic regions or protein locations, nor their frequency in the

population over time.

To overcome these limitations, we have developed *coronapp*, a web application

with two purposes: real-time tracking of SARS-CoV-2 mutational status and

annotation of user-provided viral genomic sequences. Our tool enables users to easily

perform genomic comparisons and provides an instrument to monitor SARS-CoV-2

genomic variance, both worldwide and by uploading custom and locally produced

genomic sequences. The webtool is available at http://giorgilab.dyndns.org/coronapp/

and   the   full   source   code   is   shared   on   Github

https://github.com/federicogiorgi/giorgilab/tree/master/coronapp

Results

24 The   webtool   *coronapp*   is   available   at   the   website

25 http://giorgilab.dyndns.org/coronapp/ and it automatically provides the user with the

26 current status of SARS-CoV-2 mutations worldwide. The app also allows users to

27 annotate user-provided sequences (Figure 1 A). There are multiple functionalities of

28 *coronapp*, described in the following paragraphs.

Current Status of SARS-CoV-2 mutational data

A worldwide analysis is shown, generated using data from GISAID. Specifically, we processed all SARS-CoV-2 complete (>29,000 sequenced nucleotides) genomic sequences, excluding low-quality sequences (>5% undefined nucleotide "N") and viruses extracted from non-human hosts.

The underlying database is updated weekly, and we provide the date of the last version as a reference for studies based on the data provided. We indicate the number of samples processed and the total number of mutational events detected (Figure 1 A). We also show the number of distinct mutated loci. Currently, this number is slightly below 20,000, meaning that two thirds of the original Wuhan SARS-CoV-2 genome has been affected by mutations and/or sequencing errors (the full length of the reference genome is 29,903 nucleotides, based on sequence id NC_045512.2).

Mutation frequency in SARS-CoV-2 proteins

We show the frequency of mutations along the length of every SARS-CoV-2 protein, reporting in the X-axis the amino acid position and on the Y-axis its frequency, either as number of observed samples carrying the mutation, the vase 10 logarithm of that number, or the percentage over all sequenced samples. In the example in Figure 1 B, we show the most frequent mutations affecting the viral Spike protein S, distinguishing silent mutations and amino acid-changing mutations (including the introduction of STOP codons). For Spike, the mutations appear to be evenly distributed in frequency along the protein length, with the most frequent mutation being the aforementioned D614G. Mouse-over functionality is provided to allow the user to identify the selected mutation (e.g. N439K in Figure 1 B).

The SARS-CoV-2 mutation table

The user can visualize or download the full table of mutations on which the webtool operates (Figure 2 A). This table is frequently updated and allows the user to specify a worldwide or a country-specific dataset. The table also provides a Search function to

look for specific variants or sample ids, and it can be viewed online or downloaded in

full as a Comma-Separated Values (CSV) file.

The table shows every mutation in a specific geographical area, reporting:

•   the GISAID sample ID (useful for cross-reference with the GISAID database

and other analyses based on it, e.g. Nexstrain).

•   The country where the sample was collected.

•   The position of the mutation, on the reference genome (refpos) and on the

sample (qpos).

•   The sequence at the mutation site, on the reference genome (refvar) and on the

sample (qvar).

•   The length of the sample genome (qlength); the reference genome is 29,903

nucleotides long.

•   The protein affected by the mutation or, if the mutation is extragenic, the

denomination of the untranslated region (UTR), e.g. 5'UTR or 3'UTR.

•   The effect of the mutation on the amino acid sequence of the protein (variant).

This uses the canonical mutational standard, indicating the original amino

acid(s), the position on the protein, and the mutated amino acid(s). An asterisk

(*) indicates a STOP codon, while the letters indicate amino acids in IUPAC

code. E.g. a mutation P315L indicates a leucine mutation (L) on the amino

acid location 315, normally occupied by a proline (P). Nucleotide mutations

can be silent, i.e. not yielding any aminoacidic change, e.g. the mutation

F106F, where the codon of phenylalanine 106 is affected but without changing

the corresponding amino acid. As in the previous column, mutations affecting

UTR regions are simply reported as the location of the nucleotide affected.

•   The class of the mutation, of which there are currently 10 types:

26   o SNP: a change of one or more nucleotides, determining a change in

27   amino acid sequence.

28   o SNP_stop: a change of one or more nucleotides, yielding the generation

29   of one or more STOP codons.

- o SNP_silent: a change of one or more nucleotides with no effect in protein sequence.

- o Insertion: the insertion of 3 (or multiples of 3) nucleotides, causing the addition of 1 or more amino acids to the protein sequence.

- o Insertion_stop: the insertion of 3 (or multiples of 3) nucleotides, causing the generation of a novel STOP codon.

- o Insertion_frameshift: the insertion of nucleotides not as multiples of 3, causing a frameshift mutation.

- o Deletion: the deletion of 3 (or multiples of 3) nucleotides, causing the removal of 1 or more amino acids to the protein sequence.

- o Deletion_stop: the removal of 3 (or multiples of 3) nucleotides, causing the generation of a novel STOP codon.

- o Deletion_frameshift: the deletion of nucleotides not as multiples of 3, causing a frameshift mutation.

- o Extragenic: a mutation affecting intergenic or UTR regions.

- The extended annotation of the protein region affected by the mutation (e.g. "Spike" for "S" or "Predicted phosphoesterase, papain-like proteinase" for NSP3, the Non-Structural Protein 3).

- The full name of the variant (varname), in the format proteinName:AApositionAA, to allow for unique denomination of viral proteome variants.

Mutational overview

The user is also provided with a general overview of the mutational status of the selected country or the entire world (Figure 2 B). Six bar plots provide a summary and highlights of the dataset, specifically:

- The most mutated samples, indicating which samples (in GISAID IDs) carry the highest number of mutations

- The overall mutations per sample, indicating the distributions of mutations per sample. It has been previously reported [10] that the current mode for mutation number compared to the reference NC_045512.2 genome is 7.5.

- The most frequent events per class. Classes are the same as reported in the mutation table and are described in the previous paragraph.

- The most frequent events per type. Individual mutation types are shown as specific nucleotides events, e.g. cytosine to thymidine transitions (C>T), guanosine to thymidine transversion (G>T) or even multinucleotide mutations (e.g. GGG>AAC, observed in the Nucleocapsid protein). As reported before, nucleotide transitions seem to be the most abundant SARS-CoV-2 type of mutational event worldwide [11].

- The most frequent events, either in nucleotide coordinates or in aminoacidic coordinates. Currently, the most frequent events are four mutations affecting SARS-CoV-2 genomes belonging to clade G, which is the most sequenced worldwide and predominant in Europe. These mutations are A23403G (associated to the already mentioned D614G mutation in the Spike protein), C3037T, C14408T and C241T.

Analysis of mutations over time

The *coronapp* webtool allows users to monitor the abundance and frequency of any SARS-CoV-2 mutation in any country specified (Figure 3 A). Both plots in this section report continuous dates on the X-axis, starting on the day of the first collected SARS-CoV-2 genome available on GISAID: December 24, 2019.

The "abundance" plot reports on the Y-axis the number of samples carrying a selected mutation in a particular day, in the specified country or worldwide. Since the date reported is the collection date (not the submission date to the GISAID database), there is usually a drop towards the right part of the plot, as there are fewer sequences collected approaching the day of the analysis. The "frequency" plot on the other hand normalizes the abundance of mutations by the total number of sequences generated on

each day. The plot currently shows a sharp increase in clade G-associated mutations

(e.g. S:D614G), as these mutations are most frequent in countries where sequencing is

more pervasive (e.g. United Kingdom).

Annotation of user-provided SARS-CoV-2 genomic sequence.

coronapp provides the user with an optional tool, *coronannotor,* providing the

optional possibility of uploading one or more SARS-CoV-2 genomic sequences,

which can be complete or partial. The format of the sequences is standard FASTA,

and an example input FASTA containing 12 sequences is provided (Figure 3 B). The

analysis is almost instantaneous and shows an overall breakdown of the most mutated

samples and most frequent mutations in the dataset. Moreover, a full table of all

detected mutations is provided: this can be visualized and searched on the web

browser or downloaded as a standard CSV file. Finally, a mutation frequency plot is

provided, allowing the user to visualize mutation frequency in selected proteins.

The user can easily return to the worldwide status of the app by refreshing or

reopening the page.

Discussion

Our webtool *coronapp* provides a fast, simple tool to annotate user-provided

SARS-CoV-2 genomes and visualize all mutations currently present in viral

sequences collected worldwide. The results provided by this instrument can have

several applications. The main purpose of *coronapp* is to help medical laboratories at

the front lines of COVID-19 fight with the opportunity to quickly define the

mutational status of their sequences, even without dedicated bioinformaticians.

Additionally, it enables scientists to perform mutational co-variance analyses and

to identify present and future significant functional interactions between viral

mutations, as previously attempted for the influenza virus and the human

immunodeficiency virus (HIV) [16]. Another application is the identification of the

most frequent mutations in specific protein regions: for example, our tool can quickly

identify that the most frequent mutation in the Spike protein, D614G, lies outside the

known interaction domain with the human protein ACE2, which spans roughly

between Spike amino acids 330 and 530 [17].

A recently published structural model simulating the effect of the D614G mutation

on the 3D structure of the spike protein has suggested that this mutation may result in

a viral particle which binds ACE2 receptors less efficiently, due to the masking of the

host receptor binding site on viral spikes [18]. The same researchers have reported a

possible correlation of the D614G form with increased case fatality rates,

hypothesizing that this mutation may lead to a viral form which is better suited to

escape immunologic surveillance by eliciting a lower immunologic response [18].

The *coronapp* analysis highlighted in Figure 1 B shows that a mutation located within

the Spike/ACE2 interaction domain is the change of Asparagine (N) to a Lysine (K)

in position 439 of the Spike sequence; this mutation could affect the protein folding or

its affinity with ACE2, as Asparagine is less charged than the basic amino acid

Lysine.

One of *coronapp*'s key strengths is to help prioritize scientific efforts on specific

aminoacidic variations that could affect the efficacy of anti-viral strategies or the

development of a vaccine by tracking the most frequent mutations in the population.

A further novelty of *coronapp* is that it provides a mean to assess the growth or

decline of specific mutations over time, in order to identify possible viral adaptation

mechanisms.

We provide not only the webtool, but also all the underlying code for the

annotation and visualization steps on a public Github repository, in order to help other

computational scientists in the ongoing battle against COVID-19. Furthermore, the

coronapp structure and concept could be expanded to other current and future

pathogens as well (e.g. the seasonal influenza or HIV), in order to monitor the

mutational status across proteins, countries and time.

Materials and methods

The webtool *coronapp* has been developed using the programming language R and is

based on a Shiny server (current version 1.4.0.2) running on R version 3.6.1. The app

is based on two distinct files, server.R and ui.R, managing the server functionalities

and the browser visualization processes, respectively. The results visualization utilizes

both basic R functions and Shiny functionalities; for tooltip functionality, *coronapp*

uses the R package *googleVis* v0.6.4, which provides an interface between R and the

Google visualization API [19].

The core of the annotation of the user-provided sequences rests in the NUCMER

(Nucleotide Mummer) alignment tool, version 3.1 [20]. Nucmer output is processed

by UNIX and R scripts provided in Github within the server.R file.

Authors' contributions

DM drafted the manuscript and performed the mutational analysis and literature search. LT developed the user interface code and drafted the methodological parts of the manuscript. EF worked on graphical interface of the webtool. FR wrote the manuscript and performed literature search. FMG designed the study, developed the server code, finalized the manuscript and provided financial support. All authors tested the webtool and provided original contributions to its development. All authors read and approve the final manuscript.

Competing interests

The authors have declared no competing interests.

Acknowledgements

We thank the Italian Ministry of University and Research for their support, under the Montalcini Grant 2016.

Figures and Legends

Figure 1 Overview of *coronapp*

A. Screenshot of the entry page of *coronapp* showing the basic tool description, the

interface to upload user-provided sequences and the overall summary of the mutations

detected worldwide. **B**. Common interface showing mutation frequency in

SARS-CoV-2 proteins, with occurrence of the mutation on the Y-axis and protein

coordinate on the Y-axis. Red dots indicate amino acid (aa)-changing mutations, and

blue dots indicate silent mutations. Tooltip functionality is also provided to identify

and quantify each mutation on mouse-over.

Figure 2 Mutation table and overview in *coronapp*

A. Result table of *coronapp*, available both for worldwide-precomputed and

user-input analyses. A "download full table" button is provided to allow the user to

perform larger-scale analyses autonomously. **B**. Barplots showing the most mutated

samples, overall sample mutations and most frequent mutation events, classes and

types. This analysis is also available both for worldwide-precomputed and user-input

analyses.

Figure 3 Analysis of mutations over time

A. Chronological analysis made by *coronapp*, showing the frequency of each

user-specified mutation in any user-specified country (or worldwide). The graph

shows the same data normalized by total number of samples, as the percentage of

samples sequenced in a specific day and carrying the mutation. **B**. Screenshot of the

coronannotor companion tool, allowing users to annotate their own SARS-CoV-2

sequences (provided as FASTA files).

References

[1] Guan W-J, Ni Z-Y, Hu Y, Liang W-H, Ou C-Q, He J-X, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. N Engl J Med 2020;382:1708–20. https://doi.org/10.1056/NEJMoa2002032.

[2] Phua J, Weng L, Ling L, Egi M, Lim C-M, Divatia JV, et al. Intensive care management of coronavirus disease 2019 (COVID-19): challenges and recommendations. Lancet Respir Med 2020. https://doi.org/10.1016/S2213-2600(20)30161-2.

[3] Zhang H, Penninger JM, Li Y, Zhong N, Slutsky AS. Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. Intensive Care Med 2020;46:586–90. https://doi.org/10.1007/s00134-020-05985-9.

[4] Guzzi PH, Mercatelli D, Ceraolo C, Giorgi FM. Master Regulator Analysis of the SARS-CoV-2/Human Interactome. J Clin Med 2020;9:982. https://doi.org/10.3390/jcm9040982.

[5] Ou X, Liu Y, Lei X, Li P, Mi D, Ren L, et al. Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-CoV. Nat Commun 2020;11:1620. https://doi.org/10.1038/s41467-020-15562-9.

[6] Blanco-Melo D, Nilsson-Payant BE, Liu W-C, Uhl S, Hoagland D, Møller R, et al. Imbalanced Host Response to SARS-CoV-2 Drives Development of COVID-19. Cell 2020;181:1036-1045.e9. https://doi.org/10.1016/j.cell.2020.04.026.

[7] Tu Y-F, Chien C-S, Yarmishyn AA, Lin Y-Y, Luo Y-H, Lin Y-T, et al. A Review of SARS-CoV-2 and the Ongoing Clinical Trials. Int J Mol Sci 2020;21. https://doi.org/10.3390/ijms21072657.

[8] Koyama T, Weeraratne D, Snowdon JL, Parida L. Emergence of Drift Variants That May Affect COVID-19 Vaccine Development and Antibody Treatment. Pathog Basel Switz 2020;9. https://doi.org/10.3390/pathogens9050324.

[9] Ceraolo C, Giorgi FM. Genomic variance of the 2019□nCoV coronavirus. J Med Virol 2020;92:522–8. https://doi.org/10.1002/jmv.25700.

[10] Mercatelli D, Giorgi FM. Geographic and Genomic Distribution of SARS-CoV-2 Mutations. Front Microbiol 2020;11. https://doi.org/10.3389/fmicb.2020.01800.

[11] Chiara M, Horner DS, Gissi C, Pesole G. Comparative genomics suggests limited variability and similar evolutionary patterns between major clades of SARS-CoV-2. BioRxiv; 2020. https://doi.org/10.1101/2020.03.30.016790.

[12] Gudbjartsson DF, Helgason A, Jonsson H, Magnusson OT, Melsted P, Norddahl GL, et al. Spread of SARS-CoV-2 in the Icelandic Population. N Engl J Med 2020. https://doi.org/10.1056/NEJMoa2006100.

[13] Fauver JR, Petrone ME, Hodcroft EB, Shioda K, Ehrlich HY, Watts AG, et al. Coast-to-Coast Spread of SARS-CoV-2 during the Early Epidemic in the United States. Cell 2020;181:990-996.e5. https://doi.org/10.1016/j.cell.2020.04.021.

[14] Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C, et al.
Nextstrain: real-time tracking of pathogen evolution. Bioinformatics
2018;34:4121–3. https://doi.org/10.1093/bioinformatics/bty407.
[15] Max Roser EO-O Hannah Ritchie, Hasell J. Coronavirus Pandemic
(COVID-19). Our World Data 2020.
[16] Sruthi CK, Prakash MK. Statistical characteristics of amino acid covariance as
possible descriptors of viral genomic complexity. Sci Rep 2019;9:18410.
https://doi.org/10.1038/s41598-019-54720-y.
[17] Lan J, Ge J, Yu J, Shan S, Zhou H, Fan S, et al. Structure of the SARS-CoV-2
spike receptor-binding domain bound to the ACE2 receptor. Nature
2020;581:215–20. https://doi.org/10.1038/s41586-020-2180-5.
[18] Becerra-Flores M, Cardozo T. SARS-CoV-2 viral spike G614 mutation exhibits
higher case fatality rate. Int J Clin Pract 2020.
https://doi.org/10.1111/ijcp.13525.
[19] Gesmann M, de Castillo D. Using the Google visualisation API with R. R J
2011;3:40–44.
[20] Delcher AL, Salzberg SL, Phillippy AM. Using MUMmer to Identify Similar
Regions in Large Sequence Sets. Curr Protoc Bioinforma
2003;00:10.3.1-10.3.18. https://doi.org/10.1002/0471250953.bi1003s00.

22