**Divergence and introgression in small apes, the genus**

***Hylobates*, revealed by reduced representation sequencing**

Authors: Kazunari Matsudaira[1]*, Takafumi Ishida[1]

Affiliation:

[1] Department of Biological Sciences, School of Science, The University of

Tokyo, Tokyo 113-0033, Japan

Corresponding Author:

Kazunari Matsudaira

Department of Biological Sciences, School of Science, The University of

Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan

Tel.: +81-3-5841-4496

kaz.matsudaira@gmail.com

Running title: Introgression among *Hylobates* gibbons

Word count for main text: 5,498

**Abstract**

Gibbons in the genus *Hylobates,* which live in Southeast Asia, show

great diversity, comprising seven to nine species. Natural hybridisation has

been observed in the species contact zones, although the roles played by

hybridisation and introgression in the evolution of these species remain

unclear. To uncover the divergence history and the contributions of

hybridisation and introgression to the evolution of *Hylobates*, random

amplicon sequencing-direct (GRAS-Di) analysis was employed to genotype

47 gibbons, representing eight species from three genera. After quality

filtering, over 300,000 autosomal single-nucleotide variant (SNV) sites were

identified. The SNV-based autosomal phylogeny, together with the

mitochondrial phylogeny, supported a divergence pattern beginning

approximately 4.3 million years ago. First, the mainland species, *H. pileatus*

and *H. lar*, consecutively diverged from the Sundaic island species. Second,

*H. moloch,* in Java (and likely *H. klossii,* in the Mentawai Islands) diverged

from the other species. Third, *H. muelleri,* in Borneo, and *H. agilis/H.*

*albibarbis,* in Sumatra and southwestern Borneo, diverged. Lastly, *H. agilis*

and *H. albibarbis* diverged from each other. The Patterson's D-statistics

indicated significant introgression between *H. lar* and *H. pileatus*, between

*H. lar* and *H. agilis*, and between *H. albibarbis* and *H. muelleri*, and weak

introgression was identified between *H. moloch* and *H. albibarbis*, and

between *H. moloch* and *H. muelleri abbotti*, suggesting incomplete

reproductive barriers among *Hylobates* species and that hybridisation and

introgression occur whenever the distribution ranges contact. Some

candidates for introgressed genomic regions were detected, and the

functions of these would be revealed by further genome-wide studies.

Keywords:

gibbons, GRAS-Di, hybridisation, introgression

**Introduction**

Primates is a highly diversified mammalian order, with an

estimated 500 (and a minimum of 350) extant species (Groves 2001;

Rylands and Mittermeier 2014). Recently, genome-wide studies of

non-human primates have become possible, revealing evolutionary

processes, such as speciation, and the demographic histories of various

taxa (Prado-Martinez et al. 2013; Xue et al. 2016; Nater et al. 2017).

Historical hybridisation and introgression have been detected in many cases

(e.g., chimpanzees and bonobos [de Manuel et al. 2016], macaques [Fan et

al. 2018], baboons [Rogers et al. 2019], and Dryas and green monkeys [van

der Valk et al. 2020]); thus, these processes are recognised as common

phenomena.

Moreover, the potential roles played by hybridisation and

introgression in the evolution of non-human primates, such as adaptive

introgression, have been widely discussed (Nye et al. 2018; Rogers et al.

2019; van der Valk et al. 2020), although further studies remain necessary

to reveal how hybridisation occurs and the degree to which introgression

has affected primate evolution. Among primates, Hylobatidae, which is a

family of small apes or gibbons, is considered to be a good model for

studying the contributions of divergence and hybridisation/introgression to

evolution, especially as an analogy for evolution in hominins (Zichello

2018).

Gibbons are small apes that live in the Indomalaya realm and show

great diversity among four extant genera, which each have different

numbers of chromosomes (*Hoolock* [2n = 38], *Hylobates* [2n = 44],

*Symphalangus* [2n = 50], and *Nomascus* [2n = 52]), comprising 15 to 20

species (Roos 2016; Fan et al. 2017). Among these four genera, *Hylobates*

shows the highest taxonomic diversity, followed by *Nomascus*. *Hylobates* is

distributed throughout both the mainland and islands of Southeast Asia,

and seven allopatric species are recognised, and characterised by various

pelage colours and vocalisations: *Hylobates pileatus* (pileated gibbons), *H.*

*lar* (white-handed gibbons), *H. agilis* (agile gibbons), *H. albibarbis* (Bornean

white-bearded gibbons), *H. muelleri* (Müller's gibbons), *H. moloch* (Javan

gibbons), and *H. klossii* (Kloss's gibbons) (Fig. 1; Groves 2001).

Some studies have proposed to elevate three subspecies of *H.*

*muelleri*, *H. muelleri muelleri* (Müller's gibbons), *H. muelleri abbotti*

(Abbot's gibbons) and *H. muelleri funereus* (North Bornean grey gibbons),

to independent species (Thinh et al. 2010a; Roos 2016); however, no

consensus has been reached, thus far. We used the recognised

seven-species classification, in this study. In addition, in this study, we refer

to *H. pileatus* and *H. lar* as mainland species for simplicity, although some *H.*

*lar* is known to be distributed in northern Sumatra, and refer to the other

species as Sundaic species.

The phylogeny of the *Hylobates* species has been studied

according to morphology (Groves 1972; Creel and Preuschoft 1984;

Geissmann 2002), vocalisation (Geissmann 2002), mitochondrial DNA

(mtDNA) (Garza and Woodruff 1992; Hayashi et al. 1995; Takacs et al. 2005;

Chatterjee 2006; Whittaker et al. 2007; Chan et al. 2010; Matsudaira and

Ishida 2010; Thinh et al. 2010a), and nuclear DNA (Israfil et al. 2011; Kim et

al. 2011; Chan et al. 2012, 2013). A whole-mitochondrial genome

(mitogenome) study successfully resolved the mtDNA phylogeny of

*Hylobates*, which demonstrated that *H. pileatus* diverged first and *H. lar*

diverged second, followed by the divergence between proto-*H. agilis*/*H.

muelleri* and proto-*H. moloch*/*H. klossii* (Chan et al. 2010 and its correction).

This study suggested that divergence began among mainland species and

was followed by the Sundaic species, which was supported a scenario in

which the common ancestor of *Hylobates* originated on the mainland and

that the distribution later expanded to the islands of Southeast Asia

(Groves 1972; Chivers 1977; Whittaker et al. 2007). The same and other

studies estimated that mtDNA divergence began approximately 3.5 to 3.9

million years ago (MYA) and was completed in a relatively short time (Chan

et al. 2010; Matsudaira and Ishida 2010; Thinh et al. 2010a).

In contrast with the mitogenome studies, nuclear DNA markers

have yielded ambiguous results on species phylogeny (Kim et al. 2011;

Chan et al. 2013), likely due to the rapid divergence that occurred among

*Hylobates* species, as suggested by mtDNA, resulting in the incomplete

lineage sorting of many nuclear genetic markers. In addition, because

hybridisation has been observed in the contact zones between *Hylobates*

species (Brockelman and Gittins 1984; Marshall and Sugardjito 1986), many

historical hybridisations and introgression may have occurred after the

initial divergence of each species; thus, their nuclear genomes may have

become admixed. In a previous study, using 14 autosomal loci, covering

approximately 11.5 kb, the presence of historical introgression between

some *Hylobates* species was detected (Chan et al. 2013). However, the

numbers of nuclear DNA markers used in previous studies were likely too

few to assess the complicated evolutionary history of the *Hylobates* species.

Genome-wide studies that utilise larger numbers of nuclear markers are

expected to overcome these potential issues, revealing the detailed

evolution of *Hylobates* gibbons.

To date, several approaches have been used to conduct

genome-wide studies of non-model organisms (Beichman et al. 2018). The

whole-genome sequencing (WGS) of several tens of individuals in each

species can cover all of the information in the genomes and can account for

genetic diversity in each species (e.g., de Manuel et al. 2016; Nater et al.

2017). Although the cost of WGS has been dropping rapidly, studying

multiple individuals of many species, such as of *Hylobates,* using WGS

remains expensive and is also costly in terms of computational resources.

Reduced-representation sequencing methods, such as

restriction-site-associated DNA sequencing (RAD-seq) and genotype by

sequencing (GBS), are alternative methods that limit the number of

sequencing loci across genomes, facilitating the effective acquisition of

sufficient numbers of genetic markers from large numbers of samples,

supporting a variety of genome-wide studies (Andrews et al. 2016).

Genotyping by random amplicon sequencing-direct (GRAS-Di) is an

alternative method that uses polymerase chain reaction (PCR) and a set of

primers to amplify several tens of thousands of loci, by randomly annealing

the primers to genomes (Enoki and Takeuchi 2018). This method has been

successfully applied to a variety of fish species and was able to detect

several thousands of single nucleotide variants (SNVs) in each species

(Hosoya et al. 2019).

In this study, we conducted a GRAS-Di analysis of gibbon species,

to uncover the species phylogeny and the evolutionary history of

divergence and hybridisation/introgression events among *Hylobates*

gibbons.

**Materials and Methods**

Samples

DNA was extracted from 46 Epstein Barr virus-transformed B

lymphoblastoid cell lines and one muscle tissue sample, using the QIAamp

DNA Mini Kit (Qiagen, Germany). The DNA samples consisted of two

samples from *H. agilis*, three from *H. albibarbis*, 20 from *H. lar*, one from *H.*

*moloch*, four from *H. muelleri* (consisting of two from *H. muelleri muelleri*

and two from *H. muelleri abbotti*), 10 from *H. pileatus*, one from *Nomascus*

*leucogenys* (northern white-cheeked gibbons), and six from *Symphalangus*

*syndactylus* (siamangs) (Supplementary Table S1). The samples were

derived from captive zoo gibbons, as described in previous studies (Ishida

et al. 1985; Nakayama and Ishida 2006).

At the time of sample collection, the species status and key

external characteristics were not well-recorded for some individuals.

Therefore, we confirmed the maternal species status of all 47 gibbons by

comparing the *cytochrome b* (*cytb*) sequence from mtDNA with publicly

available data (see details below). To minimise the inclusion of

unidentifiable hybrid individuals, we analysed multiple individuals from

each species/subspecies, when possible.

Mitochondrial DNA sequencing and analysis

We determined the *cytb*, tRNA-Thr, tRNA-Pro, and partial D-loop

sequences (approximately 1.9 kb) of all 47 gibbons and determined the

whole-mitogenome sequences (approximately 16.5 kb) of a subset of

individuals (n = 6). Approximately 8-kb (covering the *cytb* to D-loop region)

and 10-kb fragments of mtDNA were PCR amplified, using the following

primers:             5'-GGCTTTCTCAACTTTTAAAGGATA-3'          and

5'-TGTCCTGATCCAACATCGAG-3'         (8-kb         fragment),        and

5'-CCGTGCAAAGGTAGCATAATC-3' and 5'-TTACTTTTATTTGGAGTTGCACCA-3'

(10-kb fragment) (Finstermeier et al. 2013).

PCR was performed in a 25 µl mixture, consisting of 0.5 units KOD

FX DNA polymerase (Toyobo, Japan), 1× PCR Buffer for KOD FX, 0.4 mM of

each dNTP, 7.5 pmol of each primer, and 1 µl DNA solution. The PCR

thermal cycle conditions were as follows: the initial incubation was

performed at 94°C, for 2 min, followed by 35 cycles consisted of a

denaturation step at 98°C, for 10 s, and an annealing/extension step at

68°C for 8 min (8-kb fragment) or an annealing step at 63°C for 30 s and

extension at 68°C for 8 min (10-kb fragment), followed by a final extension

step at 68°C for 10 min. PCR amplification was confirmed by agarose gel

electrophoresis.

The PCR products were purified using a FavorPrep PCR Clean-Up

Mini Kit (Favorgen, Taiwan) and used as templates for sequencing reactions.

Sanger sequencing was outsourced to Fasmac (Japan), with internal

primers, including three forward primers (5'-CCACGACCAATGATACGAAA-3';

5'-AGACAACGCCACACTCACAC-3'; and 5'-CTTCACCCTCAGCACCCAAAGC-3'

[Andayani et al. 2001]) and four reverse primers

(5'-ATGAGGAGGAGGAGAAATAGTCCTAG-3';

5'-GGGGTGGAAGGTAATTTTGTC-3'; 5'-CGGCTTACAAGACCGGTGT-3'; and

5'-AAGACAGATACTGCGACATAGG-3' [Matsudaira et al. 2013]), to determine

the *cytb* to D-loop region. Other internal primer sequences are available, on

reasonable request, from the authors. Sequences were called from

spectrograms, using MEGA 6 (Tamura et al. 2013).

To confirm the maternal species status, we assessed the mtDNA

phylogenetic positions of all 47 *cytb* sequences (1,140 bases) by comparing

them with the published *cytb* sequences of 175 gibbons (Arnerson et al.

1996; Chan et al. 2010; Matsudaira and Ishida 2010; Thinh et al. 2010a;

Thinh et al. 2010b; Finstermeier et al. 2013; Fan et al. 2017) and eight other

primate species: *Homo sapiens*, *Pan troglodytes*, *Pan paniscus*, *Gorilla*

*gorilla*, *Pongo pygmaeus*, *Pongo abelli*, *Macaca mulatta,* and *Papio anubis*

(Horai et al. 1995; Xu and Arnason 1996; Zinner et al. 2013; Liedigk et al.

2014) (Supplementary Table S2). We constructed a maximum likelihood

(ML) tree, using IQ-TREE (Nguyen et al. 2015). The TN+F+I+G4 substitution

model was selected by ModelFinder (Kalyaanamoorthy et al. 2017). An

ultrafast bootstrap approximation test (Minh et al. 2013) was performed,

using 1,000 iterations.

Previous whole-mitogenome studies did not analyse *H. albibarbis*

(Chan et al. 2010; Matsudaira and Ishida 2010). Thus, we newly determined

the whole-mitogenome sequences of one *H. albibarbis* and five other

*Hylobates* gibbons. We analysed these six sequences, together with the

published whole-mitogenome sequences of 44 gibbons (which included

one *S. syndactylus* in the GRAS-Di data set [Sample ID: G75]) (Arnerson et al.

1996; Chan et al. 2010; Matsudaira and Ishida 2010; Finstermeier et al.

2013; Fan et al. 2017) and eight outgroup species, similar to the *cytb*

analysis (Supplementary Table S3). After aligning the sequences using

MUSCLE (Edgar 2004), which was implemented in MEGA, we extracted the

sequences of 13 protein-coding genes and two rRNA-coding regions, which

resulted in 13,798-bp sequences. The ML tree was constructed using

IQ-TREE. Among the 15 identified genes, seven partitions with different

substitution models were selected by ModelFinder (Supplementary Table

S4). An ultrafast bootstrap test was conducted, for 1,000 iterations.

Divergence time estimation was conducted using BEAST 2.6.0.

(Bouckaert et al. 2014). The relaxed, lognormal clock model and birth-death

model tree priors were selected. The partition scheme and the substitution

models used in the ML tree inference were also applied. Three divergence

time priors were used, assuming normal distribution: (1) Hominoids and

Cercopithecoids, as 30.5 MYA (95% lower and upper limit: 25.0–36.0)

(Stevens et al. 2013; Bond et al. 2015), (2) *Pongo* and *Homo/Pan/Gorilla,* as

15.5 MYA (13.0–18.0) (Kelley 2002; Besenbacher et al. 2019), (3) *Homo* and

*Pan,* as 6.5 MYA (6.0–7.0) (Senut et al. 2001; Brunet et al. 2002). Four

independent Markov chain Monte Carlo (MCMC) runs were performed for

25,000,000 generations, and parameters were recorded once in every

1,000 generations. The conversions of the results were assessed by Tracer

1.7.1. (Rambaut et al. 2018). The first 25% samples from each run were

discarded as burn-in, and the remaining results were merged, using

LogCombiner in BEAST. A consensus tree was constructed from 75,004

trees, using TreeAnotator in BEAST, and was visualised using FigTree v1.4.4

(https://github.com/rambaut/figtree/releases).

GRAS-Di

The RNA in the extracted DNA samples was digested by RNase A

(Nippon Gene, Japan), for 1 h. Then, ethanol precipitation was performed,

and the DNA was eluted using Low EDTA TE (Thermo Scientific, USA). We

measured the DNA concentrations of the samples using a Qubit 4

fluorometer, with the Qubit dsDNA BR Assay Kit (Thermo Scientific), and

prepared 100 ng of DNA (> 15 ng/µl) per sample. Library preparation and

sequencing using the GRAS-Di method were outsourced to GeneBay, Inc.

(Japan). The protocols for the library preparation were those described by

Hosoya et al. (2019), except that the PCR reaction volume was 25 µl during

the first PCR run and 50 during the second PCR run, instead of 10 µl. The

sequences of the 64 primers used to obtain amplicons during the first PCR

run were the same as those described in Hosoya et al. (2019). The library

was sequenced on a HiSeq 4000 (Illumina, USA), with 151-bp paired-end

reads.

Read mapping and genotyping

Sequence reads were de-multiplexed to each individual and

provided as fastq files. We removed 3' adapter sequences and low-quality

reads using cutadapt 1.12 (Martin 2011), with the following command:

(cutadapt        -a        CTGTCTCTTATACACATCTCCGAGCCCACGAGAC        -A

CTGTCTCTTATACACATCTGACGCTGCCGACGA --overlap 10 --minimum-length

51 --quality-cutoff 20). Cleaned reads were mapped to the reference

genome sequence of *N. leucogenys*, nomLeu3 (Carbone et al. 2014), using

Burrows-Wheeler Aligner (BWA-MEM) 0.7.17 (Li 2013). Generated sam files

were converted into bam files and sorted using SortSam command of

Genome Analysis Toolkit (GATK) 4.1.0.0. (McKenna et al. 2010). After

adding read group information, using GATK AddOrReplaceReadGroups,

each bam file was indexed by samtools 1.9 (Li et al. 2009). GATK

HaplotypeCaller with --emit-ref-confidence and GVCF options, was used to

call variants in each individual. The generated gvcf files for the 47 gibbons

were merged using GATK CombineGVCFs. GATK GenotyepGVCFs, with the

-all-sites option, was used for joint genotyping variants among the 47

gibbons.

SNV sites were extracted using GATK SelectVariants. Then, GATK

hard filters were applied to remove ambiguous results (QD < 2.0; FS > 60.0;

MQ < 40.0; MQRankSum < -12.5; ReadPosRankSum < -8.0). Subsequently,

we extracted SNV sites that satisfied the following criteria: (1) genotyped in

all 47 gibbons, with at least 5× depth of coverage for each gibbon; (2)

biallelic (i.e., no more than two alleles); and (3) variable among the 47

gibbons. Last, we extracted only those SNV sites located on the 25

autosomal long scaffolds of nomLeu3. We used GATK VariantFiltration,

vcftools 0.1.17 (Danecek et al. 2011), and vcflib

(https://github.com/vcflib/vcflib) to perform the filtering. The generated

dataset consisted of 358,142 SNV sites.

Species tree and network construction

We constructed a phylogenetic tree of the autosomal SNVs, using

ML and Bayesian frameworks. The ML tree was constructed using IQ-TREE.

Because the dataset consisted of only SNV sites, we used an ascertainment

bias correction (ASC) model (Lewis 2001) for the ML tree construction. In

the ASC model, heterozygous and invariant sites were ignored; thus,

196,186 of 358,142 SNV sites were used for the analysis. The

TVM+F+ASC+R2 model was selected by ModelFinder, and an ultrafast

bootstrap approximation test was performed, with 1,000 iterations.

A Bayesian tree was constructed, using MrBayes 3.2.7a (Ronquist

et al. 2012). The GTR+G model was selected by MrModeltest 2.4 (Nylander

2004). Four independent MCMC runs were performed, consisting of one

heat chain and three cold chains. Each run comprised 10,000,000

generations, and the parameters were sampled every 5,000 generations.

Convergence among the four runs was assessed by Tracer. The first 25% of

samples from each run were discarded as burn-in, and the results of the

four runs (i.e., 6,004 samples) were merged. The tree was visualised using

FigTree.

To assess the reticulate evolution among *Hylobates*, a phylogenetic

network of autosomal SNVs was generated, using the NeighborNet

algorithm in SplitsTree 4.15.1 (Huson and Bryant 2006). To clarify the

results, we only included *Hylobates* gibbons in this analysis.

Allele-sharing distance

To uncover the genetic relationships among individuals, we

calculated the allele-sharing distances (ASD) (Bowcock et al. 1994) between

each pair of individuals, using asd v1.0.0 (https://github.com/szpiech/asd).

Based on the ASD matrix, multi-dimensional scaling (MDS) plots

(two-dimensional plots) were drawn, using R 3.6.2 (R Core Team 2019).

Detection of introgression

The presence of introgression among *Hylobates* was assessed by

Patterson's D-statistics tests (also known as ABBA-BABA tests) (Green et al.

2010; Patterson et al. 2012). Because the phylogenetic relationships among

*Hylobates* species/subspecies have been inferred from both mitogenomes

and autosomal SNVs and are well-supported and consistent, (except for one

mitogenome from *H. agilis*; see Results), we used the autosomal tree

topology and calculated the D-statistics by setting the outgroup as either *N.*

*leucogenys* or *S. syndactylus*. The D-statistics were calculated for both

combinations of one individual per species/subspecies and the

combinations of population data for species/subspecies. The D-statistics

were calculated using AdmixTools (Patterson et al. 2012), with the R

package admixr (Petr et al. 2019).

Potentially introgressed loci

When an introgression event occurs, most genomic regions from

other species are expected to be removed, due to incompatibility and

deleteriousness. However, some regions may remain as introgressed loci

because of adaptive advantages. Potentially introgressed regions can be

detected by comparing D-statistics and genetic divergence in each segment

of the genome (Smith and Kronfrost 2013). Because introgressed regions

should have shorter coalescent times than average, the regions with high

absolute D-statistics and low genetic divergence can be considered

candidates for introgressed regions.

To detect the potentially introgressed loci among *Hylobates*

species/subspecies, we calculated the corrected D-statistics, $f_d$, (Martin et

al. 2015), and genetic divergence $d_{xy}$ (Smith and Kronfrost 2013), using

ABBABABAwindows.py            and            popgenWindows.py

([https://github.com/simonhmartin/genomics_general](https://github.com/simonhmartin/genomics_general)),            respectively.

Because of limitations in the number of SNV sites in our data set, we

calculated the values for regions with a large window size of 1,000,000

bases, with at least 100 SNVs per window.

**Results**

Genotyping

We obtained a mean $\pm$ standard deviation (SD) of 8.6 million (M) $\pm$

0.87 (min = 6.6 M; max = 10.2 M) raw read pairs, per sample. Adapter

trimming and quality filtering by cutadapt yielded 8.5 M ± 0.85 (min = 6.5

M; max = 10.0 M) read pairs, per sample. In the nomLeu3 genome, 3.6% to

5.2% (95,030,378 to 139,035,756 sites) of the 2,654,007,897 autosomal

sites were mapped by at least one read in each sample, and 1.2% to 1.8%

(31,448,837 to 47,600,791 sites) were mapped by at least five reads in each

sample. The mean depth of coverage at the mapping sites in each sample

was 16.3× ± 1.4. After hard filtering by GATK, a data set containing 358,142

biallelic, autosomal, SNV sites that were successfully genotyped in all 47

gibbons was generated.

We calculated the nucleotide diversity (Nei and Li 1979) of each

species/subspecies across the 358,142 autosomal SNV sites (Table 1). *H.*

*pileatus* showed the lowest nucleotide diversity (0.0198 ± 0.0811). The

three Bornean species/subspecies (*H. albibarbis*, *H. muelleri muelleri*, and *H.*

*muelleri abbotti*) showed the highest nucleotide diversity (0.0598–0.0694).

The other three *Hylobates* species (*H. moloch*, *H. lar*, and *H. agilis*), *S.*

*syndactylus,* and *N. leucogenys* were in the middle (0.0371–0.0511).

Phylogenetic tree, divergence time, ASD, and phylogenetic network

Phylogenetic relationships among *Hylobates* were well resolved,

both in the mitogenome and the autosomal SNV trees (Fig. 2), and the

topologies of species/subspecies relationships were consistent, except for

one mitogenome sequence from *H. agilis* (Sample ID: G80). This individual

clustered with the other *H. agilis* individuals in the autosomal SNV tree (Fig.

2B) but clustered with *H. lar* individuals and separated from other *H. agilis*

individuals in the mitogenome tree (Fig. 2A; labelled as *H. agilis* [group2]).

In the *cytb* tree, which consisted of the reference sequences of four *H. lar*

subspecies, this *H. agilis* individual tightly clustered with the sequences

from *H. lar vestitus* (Sumatran white-handed gibbons) (Supplementary Fig.

S1). Because the mitogenome represents the genealogy of only a single

locus, we considered the autosomal SNV tree to represent the true species

tree and used the autosomal SNV tree for all further analyses.

Both the autosomal and mitogenome phylogenetic trees

demonstrated that the divergence of *Hylobates* species started in the

mainland species, followed by the Sundaic species/subspecies. Based on

the estimated divergence time indicated by the mitogenomes (Fig. 2A,

Supplementary Table S5), among *Hylobates*, divergence first occurred

between *H. pileatus* and the common ancestor of the other *Hylobates*

species, approximately 4.3 MYA (95% highest posterior density credibility

interval [HPD CI]: 3.7 to 5.0). Next, *H. lar* diverged from the Sundaic species,

approximately 3.8 MYA (3.3 to 4.4). Then, *H. moloch* (and *H. klossii*)

diverged from the Sumatran/Bornean species, approximately 3.5 MYA (3.0

to 4.0). *H. muelleri* and *H. agilis*/*H. albibarbis* diverged approximately 3.2

MYA (2.8 to 3.7). *H. agilis* and *H. albibarbis* diverged approximately 1.4 MYA

(1.2 to 1.7). *H. muelleri muelleri* and *H. muelleri abbotti* diverged

approximately 1.4 MYA (1.1 to 1.7).

MDS plots of ASD also revealed similar relationships among the

*Hylobates* species/subspecies. In the MDS plot containing all *Hylobates*

gibbons (n = 40), three clusters appeared. *H. pileatus* formed one cluster, *H.*

*lar* formed another cluster, and the Sundaic *Hylobates* species/subspecies

formed the final cluster (Fig. 3A). In addition, among the Sundaic *Hylobates*

species/subspecies, *H. moloch* separated from the others, which is

consistent with the autosomal and mitogenome phylogeny results. When

we plotted only the Sundaic *Hylobates* species/subspecies (n = 10), the

species/subspecies were all separated from each other (Fig. 3B). *H. agilis*, *H.*

*albibarbis*, and *H. muelleri* (*muelleri* and *abbotti* subspecies) were plotted

on a line, with *H. agilis* at one end and *H. muelleri* at the other end. *H.*

*albibarbis* was located between the two species, closer to *H. agilis* than to

*H. muelleri*. Furthermore, *H. muelleri muelleri* (n = 2) and *H. muelleri*

*abbotti* (n = 2) were slightly separated from each other in the plot, which

indicated the presence of genetic structure-divergence between the two

subspecies.

The phylogenetic network (Supplementary Fig. S2) showed

consistent relationships among the *Hylobates* species/subspecies, as shown

in the phylogenetic trees and the MDS plots of ASD. A clear reticulation,

which indicates the introgression, was observed in *H. albibarbis* and *H.*

*muelleri* (both *muelleri* and *abbotti* subspecies). In addition, the two *H.*

*agilis* individuals (Sample ID: G80 and G83) formed a reticulation, which

may be related to differences in their mitogenomes (Fig. 2A). Furthermore,

the network highlighted the presence of a population structure within *H.*

*lar*, in which 14 of 20 individuals were tightly clustered together. This result

was similar to the *cytb* tree, in which the same 14 individuals and one

additional individual were clustered more tightly compared with the other

five individuals (Supplementary Fig. S1).

Introgression

The D-statistics calculated for species quartets showed signals of

introgression (i.e., absolute Z-score > 3) between *H. albibarbis* and *H.*

*muelleri* (both *muelleri* and *abbotti* subspecies), between *H. pileatus* and *H.*

*lar*, between *H. lar* and *H. agilis*, between *H. albibarbis* and *H. moloch*, and

between *H. muelleri abbotti* and *H. moloch* (Table 2, Supplementary Table

S6). These results were consistent, regardless of the outgroup species (*N.*

*leucogenys* and *S. syndactylus*) used for the analysis. No other signals of

introgression were detected among the other testable quartets of

*Hylobates* species/subspecies. The D-statistics calculated for individual

quartets also showed signals of introgression in the same pairs of

species/subspecies (and several cases between the *H. muelleri abbotti* and

*H. lar* individuals) (Fig. 4, Supplementary Fig. S3, Supplementary Table S7).

However, the differences in the genetic makeups of the two *H. agilis*

individuals (Sample ID: G80 and G83) affected the D-statistics and Z-scores,

and the signals could be both higher and lower than the threshold

(absolute Z-score > 3), in some cases, such as between *H. albibarbis* and *H.*

*moloch*.

Potentially introgressed loci

459           Calculations of $f_d$ and $d_{xy}$ were performed for the pairs of species

whose introgression was detected by D-statistics. We considered autosomal

regions with the top 5% $f_d$ values and the bottom 5% $d_{xy}$ values to be

potentially introgressed loci (Supplementary Table S8). Among those, three

loci (chr7b: 50,000,001-51,000,000; chr7b: 51,000,0001–52,000,000; and

chr10: 68,000,0001-69,000,000) were detected as candidates between *H.*

*lar* and *H. pileatus,* in all of the tested quartets (Fig. 5, Supplementary Table

S8). Furthermore, two loci (chr4: 151,000,001–152,000,000; chr16:

55,000,001–56,000,000) were detected between *H. agilis* and *H. lar,* in

several tested quartets (Supplementary Table S8). Note that the genomic

positions shown here are those for *N. leucogenys* (2n = 52) and, thus, differ

from the exact genomic positions found in *Hylobates* (2n = 44).

**Discussion**

The autosomal phylogeny identified in this study was consistent

with the previously reported mitogenome phylogeny (Chan et al. 2010 and

its correction) and the mitogenome phylogeny identified in this study,

which supports the following scenario: the most recent common ancestor

of *Hylobates* originated on the mainland in Southeast Asia, expanded their

distribution southward and emerged in Sundaland during the Pliocene era

(approximately 3.0 to 4.3 MYA), and speciated due to subsequent isolations

(Whittaker et al. 2007; Thinh et al. 2010a; Chan et al. 2013).

The early divergence of *H. moloch* on Java, relative to the other

Sundaic species living in Sumatra and Borneo, which was previously

uncovered by the earlier mitogenome study (Chan et al. 2010) was also

supported by the present autosomal phylogeny study. We could not include

*H. klossii* on the Mentawai islands in the GRAS-Di analysis; thus, the close

relationship between *H. moloch* and *H. klossii* that has been suggested by

mtDNA phylogeny (Takacs et al. 2005; Chan et al. 2010) and vocalisation

analysis (Geissmann 2002) should be tested in the future.

Both the mitogenome and the autosomal trees suggested that

soon after the divergence of *H. moloch*, *H. agilis*/*H. albibarbis* and *H.*

*muelleri* diverged from each other. ASD also suggests that *H. albibarbis* has

a closer relationship with *H. agilis* than with *H. muelleri* (*muelleri* and

*abbotti* subspecies). This result is consistent with a previous

population-based genetic study (Hirai et al. 2009), a *cytb* phylogeny analysis

(Thinh et al. 2010a), and a previous taxonomy classification, which included

*H. albibarbis* as a subspecies of *H. agilis,* based on vocalisations (Marshall

and Sugardjito 1986).

At present, *H. albibarbis* and *H. muelleri* are distributed in Borneo,

whereas *H. agilis* is distributed in Sumatra. The current distribution may

have occurred as follows; the common ancestor of *H. agilis* and *H.*

*albibarbis* was first distributed only in Sumatra, and approximately 1.4 MYA,

the proto-*H. albibarbis* extended their distribution to southwestern Borneo

and either replaced or admixed with *Hylobates* gibbons (likely *H. muelleri*)

that once lived in that region (Thinh et al. 2010a). Lower sea levels and the

appearance of land bridges in the Sundaic region during glacial periods

(Woodruff 2010) likely made the migration between Sumatra and Borneo

possible.

Alternatively, the common ancestor of *H. agilis* and *H. albibarbis*

may have widely distributed, from Sumatra to southwestern Borneo, after

the divergence from proto-*H. muelleri,* approximately 3.2 MYA, and 1.4

MYA may represent the time point at which the forests of Sumatra and

southwestern Borneo became disconnected. Interestingly, the estimated

mtDNA divergence time between *H. muelleri muelleri* and *H. muelleri*

*abbotti* was also approximately 1.4 MYA. Thus, the fragmentation of forests

and the isolation of gibbon populations in the Sundaic region may have

become more significant during this period than before.

Both the mtDNA divergence time and the ASD of autosomal SNVs

between *H. muelleri muelleri* and *H. muelleri abbotti* (0.0697 to 0.0701; n =

4 pairs) were similar to (or slightly larger than) those between *H. agilis* and

*H. albibarbis* (0.0671–0.0693; n = 6 pairs). Therefore, these results suggest

that these two taxa may be better classified as different species (i.e., *H.*

*muelleri* and *H. abbotti*), which was proposed by Thinh et al. (2010a).

However, these results are not conclusive, because the sample size of this

study was limited and only two individuals in each subspecies were

analysed. In addition, *H. muelleri funereus* was not analysed in this study.

The analysis of a larger number of samples, including *H. muelleri funereus*,

remains necessary to determine whether our results successfully captured

the genetic diversity among *H. muelleri* and to draw conclusions regarding

their taxonomic statuses.

The GRAS-Di analysis detected introgression signals between

several pairs of *Hylobates* species/subspecies in Patterson's D-statistics. The

signals of introgression between *H. lar* and *H. pileatus*, between *H.*

*albibarbis* and *H. muelleri* (both *muelleri* and *abbotti* subspecies), and

between *H. lar* and *H. agilis* were detected in every individual of each

species. Interestingly, these three species pairs have been known to form

contact/hybrid zones at the species boundaries (Brockelman and Gittins

1984); however, at present, the contact/hybrid zones are limited to small

areas. Because every individual of the species demonstrated signals of

introgression, the limited hybridisation and introgression that has recently

been observed in the contact/hybrid zones appears insufficient to explain

the results. Following the glacial cycles that formed the Sundaland and

fragmented the forests (Woodruff 2010), the shapes of species boundaries

and hybrid zones are likely to have changed over time.

The degree of admixed ancestry that can be identified in

individuals from the same species/subspecies may not be uniform across

their distribution. Two individuals of *H. agilis* (Sample ID: G80 and G83)

showed different levels of D-statistic and Z-score values in several

combinations, which likely reflects differences in the amounts of *H.*

*lar*-derived genetic components in their genetic makeups. Individual G80,

which has a mitogenome that is closely related to *H. lar* (*vestitus*

subspecies), showed higher absolute D-statistics in tests with *H. lar*. This

type of mtDNA has been observed in several *H. agilis* individuals from

Sumatra, in a previous study (Hirai et al. 2009). Thus, the mtDNA of G80

likely introgressed from *H. lar* to *H. agilis* in the past, similar to the

observed mtDNA introgression from *H. pileatus* to *H. lar* (Matsudaira et al.

2013). The frequency and distribution of this mtDNA group among *H. agilis*

are not known, but the differences in the mtDNA and the D-statistics may

reflect the distance from the species boundary between *H. lar* and *H. agilis,*

which is located around Lake Toba. Because differences in the genetic

makeup between G80 and G83 affected the D-statistic values, future

studies of *Hylobates* gibbons should be aware of potential hidden genetic

structures within a species.

Although the introgression signals were not robust among every

individual, introgression between *H. albibarbis* and *H. moloch* and between

*H. muelleri abbotti* and *H. moloch* were also suggested. For these two

species pairs, the current distributions are separated by seas; however,

these distributions may have contacted each other on land bridges, when

the sea level was low. These results suggested that whenever the

distribution range of two *Hylobates* species formed contact regions, hybrid

zones were formed and that the reproductive barriers between *Hylobates*

species, other than geographical barriers, are incomplete. However, the

degrees of introgression may vary among hybrid zones, as suggested by

previous observations. Although the degree of introgression may be limited

between *H. lar* and *H. pileatus* within the hybrid zone at Khao Yai, Thailand

(Brockelman and Gittins 1984), a hybrid swarm (a stable hybrid/admixed

population) has formed between *H. albibarbis* and *H. muelleri* within a

hybrid zone at the Barito headwaters, central Borneo (Mather 1992).

Previous studies have suggested possible factors that may limit

hybridisation and introgression in these contact/hybrid zones, such as

positive assortative mating and the reduced fitness of hybrid gibbons

(Brockelman and Gittins 1984, Mather 1992, Suwanvecho and Brockelman

2012; Asensio et al. 2017). Further genetic studies that focus on such

hybrid zones are expected to reveal factors that affect the frequency of

hybridisation and the degree of introgression among species.

Some potentially introgressed regions were detected by assessing

their $f_d$ and $d_{xy}$ values; however, this analysis was unable to detect

particular genes or sets of introgressed SNVs because the density of SNVs

available in this study was low; thus, we were only able to conduct the test

using large window sizes. Intensive sequencing efforts, such as the WGS of

*Hylobates* species, will reveal additional details regarding the introgressed

regions detected in this study, and will likely detect additional regions

across the genome.

Overall, the GRAS-Di analysis, which successfully identified

hundreds of thousands of autosomal SNVs, effectively resolved the species

phylogeny of *Hylobates* gibbons (except *H. klossii*) and revealed the

presence of hybridisation and introgression throughout the evolution of

*Hylobates*. Importantly, this study suggested that the phenotypic diversity

of *Hylobates* gibbons that is currently observed, such as in pelage colour

and vocalisation, represent the products of both divergence and

hybridisation/introgression. This study supports the current view of

hybridisation/introgression as driving factors in the evolution of primates

and other taxa. Future studies of *Hylobates* gibbons that include more

samples with known origins and WGS studies will likely uncover additional

details regarding the contributions of genetic changes that have occurred

during divergence and introgression to species/subspecies differences and

diversity among *Hylobates*.

**Acknowledgements**

This study was supported by JSPS KAKENHI [Grant Numbers JP18H02516

and JP19K16240].

**Conflict of interest**

The authors declare that they have no conflicts of interest.

**Data archiving**

The raw read files produced by the GRAS-Di analysis were deposited in

DDBJ Sequence Read Archive (DRA), with BioProject Accession No.

PRJDB9913. The mtDNA sequences were deposited in the

DDBJ/EMBL/NCBI nucleotide database, with Accession No.

LC548011-LC548056. Please see Supplementary Table S1 for the details.

**References**

Andayani N, Morales JC, Forstner MRJ, Supriatna J, Melnick DJ (2001)

Genetic variability in mtDNA of the silvery gibbon: implications for the

conservation of a critically endangered species. *Conserv Biol* **15**:

770-775.

Andrews KR, Good JM, Miller MR, Luikart G, Hohenlohe PA (2016)

Harnessing the power of RADseq for ecological and evolutionary

genomics. *Nat Rev Genet* **17**: 81-92.

Arnason U, Gullberg A, Xu X (1996) A complete mitochondrial DNA

molecule of the white-handed gibbon, *Hylobates lar*, and comparison

among individual mitochondrial genes of all hominoid genera.

*Hereditas* **124**: 185-189.

Asensio N, José-Domínguez JM, Kongrit C, Brockelman WY (2017) The

ecology of white-handed and pileated gibbons in a zone of overlap and

hybridization in Thailand. *Am J Phys Anthropol* **163**: 716-728.

Beichman AC, Huerta-Sanchez E, Lohmueller KE (2018) Using genomic data

to infer historic population dynamics of nonmodel organisms. *Annu Rev*

*Ecol Evol Syst* **49**: 433-456.

Besenbacher S, Hvilsom C, Marques-Bonet T, Mailund T, Schierup MH

(2019) Direct estimation of mutations in great apes reconciles

phylogenetic dating. *Nat Ecol Evol* **3**: 286-292.

Bond M, Tejedor MF, Campbell KEJ, Chornogubsky L, Novo N, Goin F (2015)

Eocene primates of South America and the African origins of New

World monkeys. *Nature* **520**: 538-541.

Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu CH, Xie D et al. (2014)

BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS*

*Comput Biol* **10**: e1003537.

Bowcock AM, Ruiz-Linares A, Tomfohrde J, Minch E, Kidd JR, Cavalli-Sforza

LL (1994) High resolution of human evolutionary trees with

polymorphic microsatellites. *Nature* **368**: 455-457.

Brockelman WY, Gittins SP (1984) Natural hybridization in the *Hylobates lar*

species group: implications for speciation in gibbons. In: Brockelman

WY, Preuschoft H, Chivers DJ, Creel N (eds) *The Lesser Apes:*

*Evolutionary and Behavioural Biology*, Edinburgh University Press:

Edinburgh. pp 498–532.

Brunet M, Guy F, Pilbeam D, Mackaye HT, Likius A, Ahounta D et al. (2002) A

new hominid from the Upper Miocene of Chad, Central Africa. *Nature*

**418**: 145-51.

Carbone L, Harris RA, Gnerre S, Veeramah KR, Lorente-Galdos B,

Huddleston J et al. (2014) Gibbon genome and the fast karyotype

evolution of small apes. *Nature* **513**: 195-201.

Chan Y-C, Roos C, Inoue-Murayama M, Inoue E, Shih C-C, Pei KJ-C et al.

(2010) Mitochondrial genome sequences effectively reveal the

phylogeny of *Hylobates* gibbons. *PLoS ONE* **5**: e14419.

Chan Y-C, Roos C, Inoue-Murayama M, Inoue E, Shih C-C, Pei KJ-C et al.

(2013) Inferring the evolutionary histories of divergences in *Hylobates*

and *Nomascus* gibbons through multilocus sequence data. *BMC Evol*

*Biol* **13**: 82.

Chan Y-C, Roos C, Inoue-Murayama M, Inoue E, Shih C-C, Vigilant L (2012) A

comparative analysis of Y chromosome and mtDNA phylogenies of the

*Hylobates* gibbons. *BMC Evol Biol* **12**: 150.

Chatterjee HJ (2006) Phylogeny and biogeography of gibbons: a

dispersal-vicariance analysis. *Int J Primatol* **27**: 699-712.

Chivers DJ (1977) The lesser apes. In: Prince Rainier III of Monaco HSH,

Bourne GH (eds) *Primate Conservation*, Academic Press: New York. pp

539-598.

Creel N, Preuschoft H (1984) Systematics of the lesser apes: a quantitative

taxonomic analysis of craniometric and other variables. In: Brockelman

WY, Preuschoft H, Chivers DJ, Creel N (eds) *The Lesser Apes:*

*Evolutionary and Behavioural Biology*, Edinburgh University Press:

Edinburgh. pp 562–613.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA et a.

(2011) The variant call format and VCFtools. *Bioinformatics* **27**:

2156-2158.

de Manuel M, Kuhlwilm M, Frandsen P, Sousa VC, Desai T, Prado-Martinez J

et al. (2016) Chimpanzee genomic diversity reveals ancient admixture

with bonobos. *Science* **354**: 477-481.

Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy

and high throughput. *Nucleic Acids Res* **32**: 1792-1797.

Enoki H, Takeuchi Y (2018) New genotyping technology, GRAS-Di, using next

generation sequencer. Proceedings of the Plant and Animal genome

conference XXVI. San Diego, CA.

https://pag.confex.com/pag/xxvi/meetingapp.cgi/Paper/29067

Fan P-F, He K, Chen X, Ortiz A, Zhang B, Zhao C et al. (2017) Description of a

new species of *Hoolock* gibbon (Primates: Hylobatidae) based on

integrative taxonomy. *Am J Primatol* **79**: e22631.

Fan Z, Zhou A, Osada N, Yu J, Jiang J, Li P et al. (2018) Ancient hybridization

and admixture in macaques (genus *Macaca*) inferred from whole

genome sequences. *Mol Phylogenet Evol* **127**: 376-386.

Finstermeier K, Zinner D, Brameier M, Meyer M, Kreuz E, Hofreiter M et al.

(2013) A mitogenomic phylogeny of living primates. *PLoS ONE* **8**:

e69504.

Garza JC, Woodruff D (1992) A phylogenetic study of the gibbons

(*Hylobates*) using DNA obtained noninvasively from hair. *Mol*

*Phylogenet Evol* **1**: 202-210.

Geissmann T (2002) Taxonomy and evolution of gibbons. *Evol Anthropol*

*Supple* **1**: 28-31.

Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M et al. (2010) A

draft sequence of the Neandertal genome. *Science* **328**: 710-722.

Groves CP (1972) Systematics and phylogeny of gibbons. In: Rumbaugh DM

(ed) *Gibbon and Siamang: A Series of Volumes on the Lesser Apes Vol 1*,

Karger: Basel. pp. 1-89.

Groves CP (2001) *Primate Taxonomy*, Smithsonian Institution Press:

Washington, DC.

Hayashi S, Hayasaka K, Takenaka O, Horai S (1995) Molecular phylogeny of

gibbons inferred from mitochondrial DNA sequences: preliminary

report. *J Mol Evol* **41**: 359-365.

Hirai H, Hayano A, Tanaka H, Mootnick AR, Wijayanto H,

Perwitasari-Farajallah D (2009) Genetic differentiation of agile gibbons

between Sumatra and Kalimantan in Indonesia. In: Lappan S, Whittaker

DJ (eds) *The Gibbons: New Perspectives on Small Ape Socioecology and*

*Population Biology*, Springer: New York. pp 37-50.

Horai S, Hayasaka K, Kondo R, Tsugane K, Takahata N (1995) Recent African

origin of modern humans revealed by complete sequences of

hominoid mitochondrial DNAs. *Proc Natl Acad Sci USA* **92**: 532-536.

Hosoya S, Hirase S, Kikuchi K, Nanjo K, Nakamura Y, Kohno H et al. (2019)

Random PCR-based genotyping by sequencing technology GRAS-Di

(genotyping by random amplicon sequencing, direct) reveals genetic

structure of mangrove fishes. *Mol Ecol Resour* **19**: 1153-1163.

Huson DH, Bryant D (2006) Application of phylogenetic networks in

evolutionary studies. *Mol Biol Evol* **23**: 254-267.

Israfil H, Zehr SM, Mootnick AR, Ruvolo M, Steiper ME (2011) Unresolved

molecular phylogenies of gibbons and siamangs (Family: Hylobatidae)

based on mitochondrial, Y-linked, and X-linked loci indicate a rapid

Miocene radiation or sudden vicariance event. *Mol Phylogenet Evol* **58**:

447-455.

Ishida T, Yamamoto K, Ishimoto G, Shotake T, Takenaka O, Nozawa K et al.

(1985) A field study of infection with human T-cell leukemia virus

among Asian primates. *Microbiol Immunol* **29**: 839-46.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017)

ModelFinder: fast model selection for accurate phylogenetic estimates.

*Nat Methods* **14**: 587-589.

Kelley J (2002) The hominoid radiation in Asia. In: Hartwig W (ed) *The*

*Primate Fossil Record*, Cambridge University Press: Cambridge. pp

369–384.

Kim SK, Carbone L, Becquet C, Mootnick AR, Li DJ, de Jong PJ et al. (2011)

Patterns of genetic variation within and between gibbon species. *Mol*

*Biol Evol* **28**: 2211-2218.

Lewis PO (2001) A likelihood approach to estimating phylogeny from

discrete morphological character data. *Syst Biol* **50**: 913-925.

Li H. (2013) Aligning sequence reads, clone sequences and assembly contigs

with BWA-MEM. arXiv:1303.3997v2 [q-bio.GN].

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N et al. (2009) The

Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**:

2078-9.

Liedigk R, Roos C, Brameier M, Zinner D (2014) Mitogenomics of the Old

World monkey tribe Papionini. *BMC Evol Biol* **14**: 176.

Marshall J, Sugardjito J (1986) Gibbon systematics. In: Swindler DR, Erwin J

(eds) *Comparative Primate Biology, Volume I: Systematics, Evolution,*

*and Anatomy*, Alan R. Liss: New York. pp 137-185.

Martin M (2011) Cutadapt removes adapter sequences from

high-throughput sequencing reads. *EMBnet.journal* **17**: 10-12.

Martin SH, Davey JW, Jiggins CD (2015) Evaluating the use of ABBA–BABA

statistics to locate introgressed loci. *Mol Biol Evol* **32**: 244-257.

Mather R (1992) A Field Study of Hybrid Gibbons in Central Kalimantan,

Indonesia. PhD thesis, Cambridge University.

Matsudaira K, Ishida T (2010) Phylogenetic relationships and divergence

dates of the whole mitochondrial genome sequences among three

gibbon genera. *Mol Phylogenet Evol* **55**: 454-459.

Matsudaira K, Reichard UH, Malaivijitnond S, Ishida T (2013) Molecular

evidence for the introgression between *Hylobates lar* and *H. pileatus* in

the wild. *Primates* 54: 33-37.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A et al.

(2010) The Genome Analysis Toolkit: a MapReduce framework for

analyzing next-generation DNA sequencing data. *Genome Res* **20**:

1297-303.

Minh BQ, Nguyen MAT, von Haeseler A (2013) Ultrafast approximation for

phylogenetic bootstrap. *Mol Biol Evol* **30**: 1188-1195.

Nakayama K, Ishida T (2006) Alu-mediated 100-kb deletion in the primate

genome: the loss of the agouti signaling protein gene in the lesser apes.

*Genome Res* **16**: 485-490.

Nater A, Mattle-Greminger MP, Nurcahyo A, Nowak MG, de Manuel M,

Desai T et al. (2017) Morphometric, behavioral, and genomic evidence

for a new orangutan species. *Curr Biol* **27**: 1-12.

Nei M, Li W-H (1979) Mathematical model for studying genetic variation in

terms of restriction endonucleases. *Proc Natl Acad Sci USA* **76**:

5269-5273.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ (2015) IQ-TREE: a fast

and effective stochastic algorithm for estimating maximum-likelihood

phylogenies. *Mol Biol Evol* **32**: 268-274.

Nye J, Laayouni H, Kuhlwilm M, Mondal M, Marques-Bonet T, Bertranpetit J

(2018) Selection in the introgressed regions of the chimpanzee genome.

*Genome Biol Evol* **10**: 1132-1138.

Nylander, JAA (2004) MrModeltest v2. Program distributed by the author.

Evolutionary Biology Centre, Uppsala University.

https://github.com/nylander/MrModeltest2

Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y et al. (2012)

Ancient admixture in human history. *Genetics* **192**: 1065-1093.

Petr M, Vernot B, Kelso J (2019) admixr-R package for reproducible analyses

using ADMIXTOOLS. *Bioinfomatics* **35**: 3194-3195.

Prado-Martinez J, Sudmant PH, Kidd JM, Li H, Kelley JL, Lorente-Galdos B et

al. (2013) Great ape genetic diversity and population history. *Nature*

**499**: 471-475.

R Core Team (2019) R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria.

http://www.R-project.org/

Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA (2018) Posterior

summarization in Bayesian phylogenetics using Tracer 1.7. *Syst Biol* **67**:

901-904.

Rogers J, Raveendran M, Harris RA, Mailund T, Leppälä K, Athanasiadis G et

al. (2019) The comparative genomics and complex population history

of *Papio* baboons. *Sci Adv* **5**: eaau6947.

Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S et al.

(2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and

model choice across a large model space. *Syst Biol* **61**: 539–542.

Roos C (2016) Phylogeny and classification of gibbons (Hylobatidae). In:

Reichard UH, Hirai H, Barelli C (eds) *Evolution of Gibbons and Siamang:*

*Phylogeny, Morphology, and Cognition*, Springer: New York. pp

151–164.

Rylands AB, Mittermeier RA (2014) Primate taxonomy: species and

conservation. *Evol Anthropol* **23**: 8-10.

Senut B, Pickford M, Gommery D, Mein P, Cheboi K, Coppens Y (2001). First

hominid from the Miocene (Lukeino formation, Kenya). *C R Acad Sci*

**332**: 137–144.

Smith J, Kronfrost MR (2013) Do *Heliconius* butterfly species exchange

mimicry alleles? *Biol Lett* **9**: 20130503.

Stevens NJ, Seiffert ER, O'Connor PM, Roberts EM, Schmitz MD, Krause C

(2013) Palaeontological evidence for an Oligocene divergence between

Old World monkeys and apes. *Nature* **497**: 611-614.

Suwanvecho U, Brockelman WY (2012) Interspecific territoriality in gibbons

(*Hylobates lar* and *H. pileatus*) and its effects on the dynamics of

interspecies contact zones. *Primates* **53**: 97-108.

Takacs Z, Morales JC, Geissmann T, Melnick DJ (2005) A complete

species-level phylogeny of the Hylobatidae based on mitochondrial

ND3–ND4 gene sequences. *Mol Phylogenet Evol* **36**: 456-467.

Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6:

molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* **30**:

2725-2729.

Thinh VN, Mootnick AR, Geissmann T, Li M, Ziegler T, Agil M et al. (2010a)

Mitochondrial evidence for multiple radiations in the evolutionary

history of small apes. *BMC Evol Biol* **10**: 74.

Thinh VN, Rawson B, Hallam C, Kenyon M, Nadler T, Walter L et al. (2010b)

Phylogeny and distribution of crested gibbons (genus *Nomascus*) based

on mitochondrial cytochrome b gene sequence data. *Am J Primatol* **72**:

1047-1054.

van der Valk T, Gonda CM, Silegowa H, Almanza S, Sifuentes-Romero I, Hart

TB et al. (2020) The genome of the endangered Dryas monkey provides

new insights into the evolutionary history of the vervets. *Mol Biol Evol*

**37**: 183-194.

Whittaker DJ, Morales JC, Melnick DJ (2007) Resolution of the *Hylobates*

phylogeny: congruence of mitochondrial D-loop sequences with

molecular, behavioral, and morphological data sets. *Mol Phylogent Evol*

**45**: 620-628.

Woodruff DS (2010) Biogeography and conservation in Southeast Asia: how

2.7 million years of repeated environmental fluctuations affect today's

patterns and the future of the remaining refugial-phase biodiversity.

*Biodivers Conserv* **19**: 919-941.

Xu X, Arnason U (1996) The mitochondrial DNA molecule of Sumatran

orangutan and a molecular proposal for two (Bornean and Sumatran)

species of orangutan. *J Mol Evol* **43**: 431-437.

Xue C, Raveendran M, Harris RA, Fawcett GL, Liu X, White S et al. (2016)

The population genomics of rhesus macaques (*Macaca mulatta*) based

on whole-genome sequences. *Genome Res* **26**: 1651-1662.

Zichello JM (2018) Look in the trees: Hylobatids as evolutionary models for

extinct hominins. *Evol Anthropol* **27**: 142-146.

Zinner D, Wertheimer J, Liedigk R, Groeneveld LF, Roos C (2013) Baboon

phylogeny as inferred from complete mitochondrial genomes. *Am J*

*Phys Anthropol* **150**: 133-140.

Fig. 1 Distribution of extant gibbons. Adapted from Thinh et al. 2010a,

2010b; Fan et al. 2017.

Fig. 2 Phylogenetic trees of gibbons. Values on the nodes (shown in blue) are bootstrap values for the ML analysis and the posterior probabilities for the Bayesian analysis. (A) Mitogenome phylogeny and divergence times, as estimated by Bayesian analysis, using BEAST 2. The outgroup species were omitted from this figure. (B) Species phylogeny, based on the autosomal SNVs, produced by ML analysis using IQ-TREE. * indicates the individuals (n = 7) that were analysed both for mitogenomes and autosomal SNVs.

Fig. 3 MDS plot of ASD. (A) A two-dimensional plot of ASD for seven

*Hylobates* species/subspecies (n = 40). (B) A two-dimensional plot of ASD

for five Sundaic *Hylobates* species/subspecies (n = 10).

Fig. 4 The ranges of Z-scores for Patterson's D-statistics among *Hylobates*

species/subspecies. *N. leucogenys* was using as an outgroup. Species

quartets are shown as (P1, P2; P3, Outgroup). Positive Z-scores (> 3)

indicate introgression between P1 and P3. Negative Z-scores (< -3) indicate

introgression between P2 and P3. Hagi: *H. agilis*, Halb: *H. albibarbis*, Hlar: *H.*

*lar*, Hmab: *H. muelleri abbotti*, Hmmu: *H. muelleri muelleri*, Hmol: *H.*

*moloch*, Hpil: *H. pileatus*, Nleu: *N. leucogenys*.

Fig. 5 Potentially introgressed genomic regions between *H. lar* and *H.*

*pileatus*. (A) $f_d$ (Martin et al. 2015) and (B) $d_{xy}$ were calculated for 1 M

base-windows, containing more than 100 SNV sites. Genomic regions with

the top 5% of $f_d$ values are shown as black dots, whereas all others are

shown as grey dots. The genomic regions are presented using the reference

genome of *N. leucogenys* (nomLeu3; 2n = 52) and, thus, do not match the

chromosomal conditions of *Hylobates* (2n = 44).

Table 1. Nucleotide diversity (π) across the 358 142 SNV sites.

| Species | Sample size | π (± SD) |
|---|---|---|
| *H. pileatus* | 10 | 0.0198 ± 0.0811 |
| *H. lar* | 20 | 0.0489 ± 0.1240 |
| *H. moloch* | 1 | 0.0371 ± 0.1891 |
| *H. agilis* | 2 | 0.0511 ± 0.1605 |
| *H. albibarbis* | 3 | 0.0646 ± 0.1543 |
| *H. muelleri muelleri* | 2 | 0.0598 ± 0.1718 |
| *H. muelleri abboti* | 2 | 0.0694 ± 0.1814 |
| *S. syndactylus* | 6 | 0.0415 ± 0.1151 |
| *N. leucogenys* | 1 | 0.0462 ± 0.2100 |

910 Table 2. Patterson's D-statistics by species. *N. leucogenys* was used as an

911 outgroup. Positive Z-scores (> 3) indicate the introgression between P1 and

912 P3. Negative Z-scores (< -3) indicate the introgression between P2 and P3.

| P1 | P2 | P3 | Outgroup | D | SE | Z |
|---|---|---|---|---|---|---|
| *H. moloch* | *H. lar* | *H. pileatus* | *N. leucogenys* | -0.11 | 0.01 | **-9.28** |
| *H. agilis* | *H. lar* | *H. pileatus* | *N. leucogenys* | -0.11 | 0.01 | **-9.22** |
| *H. albibarbis* | *H. lar* | *H. pileatus* | *N. leucogenys* | -0.11 | 0.01 | **-9.89** |
| *H. m. muelleri* | *H. lar* | *H. pileatus* | *N. leucogenys* | -0.11 | 0.01 | **-9.88** |
| *H. m. abbotti* | *H. lar* | *H. pileatus* | *N. leucogenys* | -0.11 | 0.01 | **-9.68** |
| *H. agilis* | *H. moloch* | *H. pileatus* | *N. leucogenys* | 0.01 | 0.01 | 1.00 |
| *H. albibarbis* | *H. moloch* | *H. pileatus* | *N. leucogenys* | 0.00 | 0.01 | -0.01 |
| *H. m. muelleri* | *H. moloch* | *H. pileatus* | *N. leucogenys* | 0.00 | 0.01 | 0.02 |
| *H. m. abbotti* | *H. moloch* | *H. pileatus* | *N. leucogenys* | 0.00 | 0.01 | 0.26 |
| *H. albibarbis* | *H. agilis* | *H. pileatus* | *N. leucogenys* | -0.02 | 0.01 | -1.66 |
| *H. m. muelleri* | *H. agilis* | *H. pileatus* | *N. leucogenys* | -0.02 | 0.01 | -1.29 |
| *H. m. abbotti* | *H. agilis* | *H. pileatus* | *N. leucogenys* | -0.01 | 0.01 | -0.99 |
| *H. m. muelleri* | *H. albibarbis* | *H. pileatus* | *N. leucogenys* | 0.00 | 0.01 | 0.05 |
| *H. m. abbotti* | *H. albibarbis* | *H. pileatus* | *N. leucogenys* | 0.00 | 0.01 | 0.41 |
| *H. m. muelleri* | *H. m. abbotti* | *H. pileatus* | *N. leucogenys* | 0.00 | 0.01 | -0.37 |
| *H. agilis* | *H. moloch* | *H. lar* | *N. leucogenys* | 0.07 | 0.01 | **5.86** |
| *H. albibarbis* | *H. moloch* | *H. lar* | *N. leucogenys* | 0.01 | 0.01 | 1.07 |

| | | | | | | |
|---|---|---|---|---|---|---|
| *H. m. muelleri* | *H. moloch* | *H. lar* | *N. leucogenys* | 0.01 | 0.01 | 0.87 |
| *H. m. abbotti* | *H. moloch* | *H. lar* | *N. leucogenys* | 0.02 | 0.01 | 1.66 |
| *H. albibarbis* | *H. agilis* | *H. lar* | *N. leucogenys* | -0.08 | 0.01 | **-7.96** |
| *H. m. muelleri* | *H. agilis* | *H. lar* | *N. leucogenys* | -0.07 | 0.01 | **-6.34** |
| *H. m. abbotti* | *H. agilis* | *H. lar* | *N. leucogenys* | -0.06 | 0.01 | **-5.59** |
| *H. m. muelleri* | *H. albibarbis* | *H. lar* | *N. leucogenys* | 0.00 | 0.01 | -0.17 |
| *H. m. abbotti* | *H. albibarbis* | *H. lar* | *N. leucogenys* | 0.01 | 0.01 | 0.97 |
| *H. m. muelleri* | *H. m. abbotti* | *H. lar* | *N. leucogenys* | -0.01 | 0.01 | -1.14 |
| *H. albibarbis* | *H. agilis* | *H. moloch* | *N. leucogenys* | 0.04 | 0.01 | **4.07** |
| *H. m. muelleri* | *H. agilis* | *H. moloch* | *N. leucogenys* | 0.03 | 0.01 | 2.29 |
| *H. m. abbotti* | *H. agilis* | *H. moloch* | *N. leucogenys* | 0.05 | 0.01 | **3.88** |
| *H. m. muelleri* | *H. albibarbis* | *H. moloch* | *N. leucogenys* | -0.01 | 0.01 | -0.94 |
| *H. m. abbotti* | *H. albibarbis* | *H. moloch* | *N. leucogenys* | 0.01 | 0.01 | 1.04 |
| *H. m. muelleri* | *H. m. abbotti* | *H. moloch* | *N. leucogenys* | -0.02 | 0.01 | -2.06 |
| *H. m. muelleri* | *H. m. abbotti* | *H. agilis* | *N. leucogenys* | 0.02 | 0.01 | 1.97 |
| *H. m. muelleri* | *H. m. abbotti* | *H. albibarbis* | *N. leucogenys* | 0.02 | 0.01 | 2.06 |
| *H. agilis* | *H. albibarbis* | *H. m. muelleri* | *N. leucogenys* | -0.13 | 0.01 | **-15.43** |
| *H. agilis* | *H. albibarbis* | *H. m. abbotti* | *N. leucogenys* | -0.13 | 0.01 | **-15.70** |