# Neurofunctional basis underlying audiovisual integration of print and speech sound in Chinese children

Zhichao Xia [a, b * †], Ting Yang [c, d *], Xin Cui [a, e], Fumiko Hoeft [e, f, g, h], Hong Liu [c, d, f], Xianglin Zhang [a], Hua Shu [a †], Xiangping Liu [c, d †]

[a] State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern Institute for Brain Research, Beijing Normal University, China

[b] School of Systems Science, Beijing Normal University, China

[c] Faculty of Psychology, Beijing Normal University, China

[d] Beijing Key Laboratory of Applied Experimental Psychology, National Demonstration Center for Experimental Psychology Education, Faculty of Psychology, Beijing Normal University, China

[e] Haskins Laboratories, USA

[f] Department of Psychological Sciences and Brain Imaging Research Center, University of Connecticut, USA

[g] Department of Psychiatry and Weill Institute for Neurosciences and Dyslexia Center, University of California, San Francisco, USA

[h] Department of Neuropsychiatry, Keio University School of Medicine, Japan

* These authors contributed equally to this work

† **Corresponding authors**: Dr. Hua Shu, State Key Lab of Cognitive Neuroscience

and Learning, Beijing Normal University, China. Email: shuhua@bnu.edu.cn; Dr.

Xiangping Liu, Faculty of Psychology, Beijing Normal University, China. Email:

lxp599@163.com; Dr. Zhichao Xia, State Key Lab of Cognitive Neuroscience and

Learning, Beijing Normal University, China. Email: xiazc.psy@gmail.com

**Acknowledgments**: The authors would like to thank all the children and their

families for participating in this study and all the examiners for helping with data

collection. This work was supported by the Key Program of the National Social

Science Foundation of China (14ZDB157), National Key Basic Research Program of

China (2014CB846103), National Natural Science Foundation of China grants

(31271082, 31671126, 31611130107, 61374165, 81801782), Beijing Municipal

Science & Technology Commission (Z151100003915122), Fundamental Research

Fund for the Central Universities, China Postdoctoral Science Foundation

(2019T120062, 2018M641235), and Social Science Fund of Beijing (17YYA004).

**Conflict of Interest**: The authors declare no competing financial interests.

**Author Contributions**: Conceptualization: ZX, HS, XL; Investigation: ZX, TY, XC,

HL, XZ; Formal Analysis: ZX; Data Curation: ZX; Writing – Original Draft

Preparation: ZX; Writing – Review & Editing: ZX, TY, XC, FH, HL, XZ, HS, XL;

Funding Acquisition: ZX, HS, XL; Supervision: ZX, HS, XL

**Data Availability Statement**: The data that support the findings of this study are

available from the corresponding authors upon reasonable request. The data are not

publicly available due to privacy or ethical restrictions.

**Running Title**: Brain basis of audiovisual integration

## 48  Abstract

Effortless print-sound integration is essential to reading development, and the

superior temporal cortex (STC) is one most critical brain regions in alphabetic

languages. To examine the neural basis in a non-alphabetic language and its

relationship with reading ability, we conducted a functional magnetic resonance

imaging study in typically developing Chinese children. Audiovisual additive

enhancement and congruency effect in two scripts with different orthographic

depths—characters (opaque) and pinyin (a transparent alphabetic transcription of

characters)—were investigated. In terms of commonality, bilateral STCs displayed

additive enhancement in processing both character and pinyin stimulations.

Moreover, the magnitudes of congruency effect in the left STC for the two scripts

were highly correlated. On the other hand, differences were also observed. First,

areas in addition to the STC showed additive enhancement in processing pinyin-

sound associations. Second, while the bilateral superior temporal, left

inferior/middle frontal and parietal regions manifested a striking incongruency

effect (incongruent > congruent) for character-sound pairs, no significant clusters

were revealed for pinyin-sounds. Finally, the magnitude of congruency effect in the

left middle frontal gyrus for characters was specifically associated with silent

reading comprehension proficiency, indicating automatic semantic access and

integration during implicit character-sound processing. In contrast, the congruency

effect in the left STC for pinyin was specifically associated with oral reading fluency

that relies mainly on grapho-phonological mapping. To summarize, this study

revealed both universal and script-specific neurofunctional substrates underlying

audiovisual integration as well as their processing- and region-dependent

associations with reading abilities in typical Chinese children.

**Keywords**

Audiovisual integration, character, fMRI, pinyin, Chinese, reading development

**Highlights**

• The superior temporal cortex is involved in print-sound integration regardless of

orthographies.

• The neural manifestation of audiovisual integration is modulated by

orthographic transparency.

• The congruency effect in the left middle frontal gyrus for characters is correlated

to reading comprehension.

• The congruency effect in the left superior temporal gyrus for pinyin is correlated

with oral reading fluency.

• This study provides neural evidence on the role of pinyin in Chinese reading

acquisition.

## 86    1. Introduction

Efficient print-speech sound mapping is a prerequisite for and a hallmark of

successful reading acquisition (Blomert & Froyen, 2010; Fabio Richlan, 2019).

Behavioral and neuroimaging studies have revealed that the level of automatic

audiovisual integration uniquely accounts for reading outcomes in both shallow

(e.g., Dutch, German) and opaque orthographies (e.g., English) (Bakos, Landerl,

Bartling, Schulte-Körne, & Moll, 2017; Blau et al., 2010; Leo Blomert & Willems,

2010; Kronschnabel, Brem, Maurer, & Brandeis, 2014; McNorgan, Randazzo-

Wagner, & Booth, 2013).

Better knowledge about the neural basis underlying audiovisual integration

can further our understanding of how this process contributes to reading

development and provide objective training targets for children with reading

disorders (RD). In the past two decades, functional magnetic resonance imaging

(fMRI) has been widely used to answer the 'where' question. Two analytic

approaches are the most common methods to identify brain regions involved in

processing multimodal information. One is based on the additive model that

initially examined in animal electrophysiological studies where researchers found

audiovisual signals induced stronger (i.e., super-additive) or lower (i.e., sub-

additive) neural response than the summation of responses to unimodal auditory

and visual inputs in areas such as the STC (Stein & Stanford, 2008). This super-

additive enhancement criterion was then applied to fMRI data but proven too strict

about detecting multisensory integration regions. In contrast, with liberal variants

such as mean additivity enhancement criterion (i.e., congruent condition > mean of

unimodal auditory and visual conditions), areas including bilateral STCs can be

successfully identified (Beauchamp, 2005; Kronschnabel et al., 2014; McNorgan &

Booth, 2015). In terms of functional significance, the super- (and mean) additive

enhancement occurs early and reflects general aspects of processing (Kronschnabel

et al., 2014; Xu, Kolozsvari, Oostenveld, Leppanen, & Hamalainen, 2019). No

associations between additive enhancement with reading performance were found

(Kronschnabel et al., 2014). In this case, we expected that the bilateral STCs would

also show an additive enhancement in processing Chinese print-sound pairs.

Distinguishable brain response to congruent versus incongruent audiovisual

pairs is another reliable measure of multisensory integration (a.k.a. congruency

effect or incongruency effect if in the opposite direction) (Blau et al., 2010; Blau, van

Atteveldt, Ekkebus, Goebel, & Blomert, 2009; Holloway, van Atteveldt, Blomert, &

Ansari, 2015; van Atteveldt, Formisano, Goebel, & Blomert, 2004; van Atteveldt,

Formisano, Goebel, & Blomert, 2007). Compared with additive enhancement, the

congruency effect can reflect deep processes on the information carried by the

stimuli. In the pioneering study by van Atteveldt et al. (2004), the STC showed

stronger activation in the congruent condition compared with incongruent

condition. Although the involvement of STC in audiovisual integration of print and

speech sound has been replicated regardless of the level of processing (e.g.,

phoneme, syllable, or whole word) (Kronschnabel et al., 2014; McNorgan et al.,

2013; Wang, Karipidis, Pleisch, Fraga-Gonzalez, & Brem, 2020), the precise

manifestation of the effect (e.g., direction, strength) and whether other areas

display such an effect are associated with multiple factors, such as orthographic

depth (Holloway et al., 2015; Weiyong Xu, Kolozsvari, Monto, & Hämäläinen, 2018)

and reading ability (Blau et al., 2010; Blau et al., 2009; Kronschnabel et al., 2014).

In addition, task paradigm (McNorgan & Booth, 2015; van Atteveldt et al., 2007),

stimuli properties (Kronschnabel et al., 2014), and developmental stage (Plewko et

al., 2018; Wang et al., 2020) also affect the congruency effect but are beyond the

scope of this study.

Orthographic depth describes the consistency of mapping between

orthographic and phonological forms (e.g., grapheme-phoneme correspondence in

alphabetic languages, GPC hereafter), which significantly influences behavioral and

neural profiles of reading development and impairments in a given writing system

(Borleffs, Maassen, Lyytinen, & Zwarts, 2018; Richlan, 2014). The neural

processing accompanying audiovisual integration is also modulated by orthographic

transparency. For example, native speakers of English (deep) show an incongruency

effect in the STC during processing letter-sound pairs, which differs from the

congruency effect observed in Dutch (shallow) (Holloway et al., 2015). More

importantly, when the same group of English-speaking participants processed

numbers and number names, where the mapping was one-on-one, a strong

congruency effect appeared. However, the number of studies directly comparing

different orthographies is limited, and most existing evidence is restricted to

alphabetic languages. The question of whether the same neural substrate exists in

non-alphabetic languages with dramatically different linguistic characteristics, such

as having no GPC rules, remains largely open.

Chinese is particularly suited for answering cross-linguistic questions given

its logographic nature and extremely deep orthography (Perfetti, Cao, & Booth,

2013). Since that characters directly map to syllables and there are no GPC rules,

the print-sound mapping is quite arbitrary. Moreover, in many cases, different

characters share the same pronunciation (i.e., homophones), whereas a single

character may also have multiple sounds (i.e., polyphone). Therefore, the link

between print (character) and speech sound (syllable) is even more opaque. Based

on previous findings in alphabetic languages, an incongruency effect during

audiovisual integration was expected. In addition, grapho-semantic correspondence

is more systematic in Chinese characters, deeply involved in character recognition,

and plays a role equal to or even more important than grapho-phonological

processing in reading acquisition (Liu et al., 2017; Ruan, Georgiou, Song, Li, & Shu,

2018; Yang, Shu, McCandliss, & Zevin, 2013; Yang, Zevin, Shu, McCandliss, & Li,

2006). Thus, it is also reasonable to expect that additional regions, such as the

inferior and middle frontal cortex known to be involved in semantic processing,

would be recruited (Wu, Ho, & Chen, 2012; Zhao et al., 2014). Recently, Xu et al.

(2019) conducted a magnetoencephalography (MEG) study in Chinese adults with a

one-back task and identified an incongruency effect across the left frontal cortex

and STC. While the finding is interesting, some aspects need to be considered,

including the small sample size (n = 12), flat tone speech sounds that do not

correspond to any character as stimuli, and source reconstruction accuracy of MEG

data. Moreover, for experienced adult readers, audiovisual integration of characters

and corresponding sound is fully automated. Thus, in addition to cross-linguistic

issues, another important but unanswered question involves the neurofunctional

features of this process in the developing brain. Against this background, the

current study was to investigate the neural basis of integrating characters and

speech sounds in typically developing Chinese children.

In particular, we focused on the early-acquired single-element characters

given their fundamental role in learning-to-read in Chinese (Shu, Chen, Anderson,

Wu, & Xuan, 2003). At the very beginning stage, a small set of pictographic

characters are taught first. These simple characters will later be used as radicals in

complex compound characters (i.e., phonograms), which comprise greater than 80%

of Chinese characters (Chan & Siegel, 2001). Importantly, as a pictographic

character does not contain any components (i.e., radicals), there is no clue on its

pronunciation or meaning. Thus, the mappings between its visual form with sound

(syllable) and meaning (morpheme) are entirely arbitrary. In addition, Mandarin

Chinese offers a valuable opportunity to examine the influence of orthographic

depth since children learn pronunciations of characters with Hanyu pinyin (pinyin

hereafter). In brief, pinyin is an official Roman alphabetic system used to teach the

sounds of characters in Mainland China and Singapore (Yin et al., 2011). A child

learns pinyin first once entering school and then uses it to assist in conquering the

pronunciation of new characters (Guan, Liu, Chan, Ye, & Perfetti, 2011; Shu, Peng,

& McBride-Chang, 2008; Yan, Miller, Li, & Shu, 2008). In other words, pinyin acts

as a scaffold during reading acquisition. Given that pinyin borrows letters from

alphabets and the majority have consistent mappings with speech sounds

(phonemes), it is pretty transparent. Therefore, we included pinyin in this study to

directly examine the effect of orthographic depth and expected to observe a similar

activation pattern in processing pinyin-sounds as in shallow languages.

In addition to investigating the neural basis of audiovisual integration at the

group level, understanding how it correlates to individual heterogeneity in reading

abilities is important. On the one hand, children with RD showed a reduced

congruency effect compared with their typical peers (e.g., Blau et al., 2010; Blau et

al., 2009; Plewko et al., 2018). On the other hand, large individual differences in

reading ability are noted even within each group and can be accounted for by

variances at the brain level (Kast, Bezzola, Jancke, & Meyer, 2011; McNorgan,

Awati, Desroches, & Booth, 2014; McNorgan et al., 2013; Plewko et al., 2018; Wang

et al., 2020). Understanding such individual differences in linguistic processing,

which is often masked by averaging at the group level, is valuable theoretically and

practically (Kanai & Rees, 2011; Kidd, Donnelly, & Christiansen, 2018; Seghier &

Price, 2018). Therefore, we also adopted an individual differences approach to

investigate whether neural circuitries link congruency effects in processing

character/pinyin-sound pairs with typical reading acquisition.

## 2. Materials and Methods

### 2.1. Participants

The initial sample consisted of 54 typically developing children according to the following inclusion criteria: (a) native speaker of Mandarin Chinese, (b) right-handed, (c) normal audition, (d) normal or corrected-to-normal vision, (e) full-scale intelligence quotient (IQ) $\geq$ 80, (f) no history of neurological or psychiatric disorders, and (g) no reading difficulties (> 16th percentile on the standardized reading screening task *Character Recognition*) (Xue, Shu, Li, Li, & Tian, 2013). All the children received formal instruction on pinyin in the first half of the semester in the 1st grade. Before the experiment, the children and their parents/guardians were informed about the aim and procedure in detail and signed written consent. All behavioral and neuroimaging data were collected during 2018. This study received ethical approval from the Institutional Review Board of State Key Laboratory of Cognitive Neuroscience and Learning at Beijing Normal University.

The quality of each functional run was evaluated based on in-scanner task performance and the number of outliers (i.e., 'bad' data points) in blood-oxygen-level-dependent signal caused by severe head motion or spikes. Acceptance depended on the following criteria: (i) no more than 15% of the volumes (10 data points) were marked as outliers; (ii) no less than 75% in overall accuracy (combining all the conditions) and accuracy for each condition should be no less than 50%. The run failing to meet either criterion was removed from the analysis. As a result, 36

children had acceptable imaging data for the character experiment (23 girls; age

110-141 months; mean age = 127, grades 3-5). Eighteen children were excluded due

to incomplete data collection (n = 9), poor data quality (n = 8), or the combination of

poor data quality and task performance (n = 1). For pinyin, the final sample

consisted of 41 children (27 girls; age 110-141 months; mean age = 126, grades 3-5).

Thirteen children were excluded due to incomplete data collection (n = 6), poor data

quality (n = 6), or the combination of poor data quality and task performance (n =

1).

**2.2. Behavior measures**

Each child underwent testing individually in a silent booth. IQ was measured with

the abbreviated version of the *Chinese Wechsler Intelligence Scale for Children*

(*WISC-CR*) (Wechsler, 1974). Specifically, *Information*, *Similarities*, and *Digit Span*

subtests were used to estimate verbal IQ; *Picture Completion*, *Block Design*, and

*Coding* subtests were used to estimate performance IQ.

A set of neuropsychological tasks was used to measure children's reading and

reading-related cognitive-linguistic skills. In brief, *Character Recognition* that had

been used for identifying Chinese dyslexic children (Cui, Xia, Su, Shu, & Gong,

2016; Xia, Hoeft, Zhang, & Shu, 2016) was used here for the screening aim. *Word*

*List Reading* and *Silent Reading Comprehension* were used to measure oral reading

fluency and reading comprehension proficiency, respectively. The former relies more

on grapho-phonological mapping, whereas the latter relies more on grapho-semantic

processing (Xia et al., 2018). Finally, *Phoneme Deletion*, *RAN*, and *Morphological*

*Production* tasks were used to measure PA, RAN, and morphological awareness

(MA), the three critical cognitive-linguistic skills in Chinese reading development

(Lei et al., 2011).

*Character Recognition* is a standardized test for estimating the number of

characters children have learned. The test consisted of 150 characters selected from

grades 1-6 textbooks (Xue et al., 2013). The characters were arranged in the order of

difficulty on ten sheets of A4 paper. During the task, the participant was asked to

name the characters in sequence until failing in all 15 items on one page. There was

no time limit. Each correct answer is worth 1 point, and the full mark is 150.

*Word List Reading* is a standardized test for measuring the level of oral

reading fluency (Zhang et al., 2012). The test consisted of 180 high-frequency two-

character words arranged in a 9-column by 20-row matrix in a sheet of A4 paper.

The participant was required to read the words aloud in left-to-right, up-to-down

order as accurately and quickly as possible. The time used to complete the task was

recorded, and an index referring to the number of words the participant read

correctly per minute was calculated.

*Silent Reading Comprehension* is a standardized test for assessing children's

reading comprehension (Lei et al., 2011). The test included 100 items arranged

based on the number of characters (i.e., from a simple sentence to a short

paragraph). The participant was requested to read each sentence/paragraph silently

and decide the correctness of the meaning with a mark of ✓ or ✗ as fast as possible

within 3 minutes. The total number of characters in sentences with correct

responses was calculated and then transformed to the number of characters the

participant read per minute.

*Phoneme Deletion* was used to assess PA (Li, Shu, McBride-Chang, Liu, &

Peng, 2012). This task included three sections, and the participant was asked to

delete the initial, middle, or final sound from an orally presented syllable and

pronounce the remaining portion. Each section consisted of 3 practice trials and 8-

10 testing items. One correct response is worth 1 point for a total possible score of

28.

*Digit RAN* required the participant to name 50 one-digit numbers on a card

as accurately and rapidly as possible (Liu et al., 2017). Numbers 1, 2, 3, 5, and 8

were repeated ten times each and arranged randomly in a 5-column by 10-row

matrix. The task was administered twice, and the average time the participant

completed the task was used as the score.

*Morphological Production* was used to measure MA at the character level

(Shu, McBride-Chang, Wu, & Liu, 2006). In each trial, a character was orally

presented to the participant in a high-frequency word. The participant had to

respond with two new words. In one word, the given character retains the same

meaning as in the original case, whereas in the other word, the meaning of the

character should be different. This task contained 15 characters for a total possible

score of 30.

**2.3. Stimuli and experimental design**

In this study, we investigated neurofunctional correlates of audiovisual integration

in processing character-sound and pinyin-sound pairs separately. In the character

experiment, the materials were characters and corresponding speech sounds

(syllables) (**Figure 1A**). Fifty-six high frequency pictographic characters were

selected (Chinese Single-Character Word Database;

https://blclab.org/pyscholinguistic-norms-database/). All the characters are visually

simple, taught early, and used as radicals in complex compound characters. During

the experiment, visual stimuli were displayed in white, 'KaiTi' font, and 96 pt at the

center of a black background. A native Chinese male recorded the speech sounds

with a sampling rate of 44.1 kHz and 16-bit quantization. The audio files were

normalized to 85 dB and then edited with a bandpass (100-4000 Hz) filter with

Audacity (https://www.audacityteam.org/). The average duration of all sounds was

476.3 (±87.5) ms without pre- or post-silence. All the speech sounds can be

recognized easily. In the pinyin experiment, the visual stimuli were syllables

written in pinyin (e.g., sound /sheng 1/ can be written as 'shēng', **Figure 1B**), in

white, 'Century Schoolbook' font, and 90 pt. The auditory stimuli were the same as

those used in character conditions.

The fMRI procedure was adapted from Blau et al. (2010) (**Figure 1C**). A

block design was used with four conditions in each experiment: unimodal auditory

(Aud), unimodal visual (Vis), audiovisual congruent (avC), and audiovisual

incongruent (avI). The visual and auditory stimuli were presented simultaneously

in the avC and avI conditions. The entire study contained 4 functional runs, half of

which were for character and the other half for pinyin. To avoid priming from

character processing on pinyin processing, the pinyin experiment was always

conducted first. Each run included 8 experimental blocks (duration 20.8 s), with

each condition replicated twice and interleaved with 9 rest blocks (duration 20.8 s).

In each experimental block, there were 4 mini-blocks. Each mini-block consisted of a

1.5-s brain volume acquisition period and a 3.7-s silence period (i.e., sparse-

sampling) during which 4 trials of stimuli were presented. The order of the stimuli

and conditions were pseudorandomized. Considering the age of the participants, a

simple target detection task that does not require active monitoring of congruency

status was used to help maintain attention during scanning. Compared to the active

matching task, the passive paradigm will not change the automatic response

pattern during audiovisual integration (Blau et al., 2010; van Atteveldt et al., 2007).

The participant was asked to press the button with the right index finger accurately

and quickly to the auditory (440 Hz pure tone), visual (an unpronounceable symbol)

targets, or their combinations that appeared twice randomly per experimental block

(**Figure 1D**). To ensure that the child performed the task correctly, two practice

sessions were administered, including one outside the scanner and the other before

the first fMRI session inside the scanner.

### 2.4. Image acquisition

The 3-Tesla Siemens MAGNETOM Trio Tim scanner in Imaging Center for Brain

Research at Beijing Normal University was used to collect all the images with a 12-

channel head coil. For each child, 4 T2$^*$ sensitive gradient echo planar imaging

sequences were collected with the following parameters: repetition time (TR) = 5200

ms (1500 ms image acquisition plus 3700 ms delay), echo time = 32 ms, flip angle =

90 degrees, slice thickness = 4.5 mm, voxel size = 3.0 × 3.0 × 4.5 mm$^3$, interscan gap

= 0.675 mm, number of slices = 24, number of volumes = 68, and time of acquisition

= 5 minutes and 54 s. The sparse-sampling method provided a 3.7-s period of silence

for stimuli presentation. Such a design is recommended for fMRI experiments

investigating processing with auditory inputs (Talavage & Hall, 2012). Three

dummy scans were administered before image collection for each functional run to

avoid scanner equilibration effects. In addition, a high-resolution whole-brain T1-

weighted structural image (magnetization-prepared rapid acquisition with gradient

echo, TR = 2530 ms, echo time = 3.39 ms, inversion time = 1100 ms, flip angle = 7

degrees, slice thickness = 1.33 mm, voxel size = 1.33 × 1.0 × 1.33 mm$^3$, number of

axial slices = 144, and time of acquisition = 8 minutes and 7 s) was collected. Prior

to the formal scan, children were informed about the experimental procedure,

familiarized with the noise of MRI sequences, and trained to hold still. Earplugs

and noise-proof headphones were used to help reduce scanner noise, and foam pads

were used to help reduce head motion during scanning. A radiologist blinded to the

details of the study reviewed all the images to ascertain any pathological deviations.

**2.5. fMRI preprocessing**

Preprocessing of brain images was conducted with SPM12 (Wellcome Department of

Cognitive Neurology, London, UK, http://www.fil.ion.ucl). It included the following

steps: (a) head motion correction (realign), (b) ART-based outlier detection (the 'bad'

time point was defined as intensity > global mean ± 9 SD or frame-to-frame head

motion [framewise displacement] > 2 mm;

https://www.nitrc.org/projects/artifact_detect), (c) T1 image segmentation, (d)

normalization to the standard template in Montreal Neurological Institute (MNI)

space, and (e) smoothing with an 8-mm full-width at half-maximum Gaussian

kernel. No slice timing correction was performed given the discontinuous nature of

the signal in fMRI data collected using sparse-sampling (Perrachione & Ghosh,

2013). At the individual level, experimental conditions were modeled with a

generalized linear model. Parameters referring to outliers in the time course and

head motion (3 translation, 3 rotation, and 1 framewise displacement) were

included in the model to exclude the effect of nuisance covariates. Then, we

calculated two brain maps—avC vs. (Aud + Vis) / 2 and avC vs. avI, which were

used in the subsequent analyses on additive enhancement and congruency effect,

respectively.

**2.6. Statistical analyses**

Descriptive statistics of demographic and behavioral measures were conducted first,

followed by correlations between reading abilities and cognitive-linguistic skills.

The Bonferroni method was used to deal with multiple comparisons (corrected $p <$

0.05 / 15). Next, we evaluated in-scanner task performance. In this study, we

defined the accurate response as a button press to the target with a reaction time

(RT) between 200 and 2000 ms. The average reaction time was calculated for correct

responses.

The two aforementioned approaches were used in the whole brain analyses to

identify areas involved in audiovisual processing of character and pinyin

stimulations separately. The first approach was to detect mean additive

enhancement. It consisted of two criteria: (i) stronger activation in each unimodal

condition against the baseline, i.e., Aud > 0, Vis > 0. This criterion was used to

exclude areas that only responded to one input modality; (ii) stronger activation in

the audiovisual congruent condition compared with the mean of unimodal

conditions, i.e., avC > (Aud + Vis) / 2. For this, three one-sample $t$-tests were

administered, and intersections were identified. For each $t$-test, an uncorrected

threshold of $p$-voxel < 0.05 was applied, resulting in a conjoint $p < 0.05\textasciicircum 3 =$

0.000125 for the overlapping voxel. To be conservative, only clusters containing

more than 200 continuous voxels were considered to be significant. The second

approach was used to identify regions showing congruency effects. A one-sample $t$-

test was conducted with individual contrast maps of congruency (i.e., congruent

against incongruent). We focused on brain areas showing activation (uncorrected $p$-

voxel < 0.05) in at least one multimodal condition to avoid results produced by

differences in de-activation. A threshold of $p$-voxel < 0.005, FWE corrected $p$-cluster

< 0.05 was used to identify significant clusters while dealing with multiple

comparisons. Previous studies have repeatedly reported that the congruency effect

is associated with reading ability. To avoid missing the effect masked at the group

level, we conducted two regression analyses where the standard score of oral

reading fluency or silent reading comprehension proficiency was included as the

variate of interest, respectively. The same threshold of FWE corrected p < 0.05

(cluster-forming $p$-voxel < 0.005) was used to address multiple comparisons.

Nuisance variables of age and sex were controlled statistically in all the analyses.

With the results of whole brain analyses, we defined areas with different

combinations of additive enhancement, congruency effect, and congruency-reading

correlation as the regions-of-interest (ROIs). Such differences may indicate different

roles the areas play in audiovisual integration. Given that domain-specific (i.e.,

reading-related) processes were of particular interest, we did not include areas that

only showed additive enhancement. Those regions could be more involved in

domain-general processing. The aims of the ROI analysis were (1) to explicitly

present the effects and (2) to exclude the possibility that the absence of significance

on a specific effect was due to the strict whole brain threshold. For this, in the

region only showing congruency-related effects but no additive enhancement at the

whole brain level, we further examined its activation in unimodal conditions and

the contrast avC - (Aud + Vis) / 2. In the region showing additive enhancement and

congruency-behavior correlation but no congruency effect across the entire group at

the whole brain level, we examined the congruency effect with the average values

within the cluster. We also examined brain-behavior associations with the other

reading ability and cognitive-linguistic skills, while the nuisance variables of age

and sex were controlled statistically. Bonferroni correction was used to control the

number of behavior variables (corrected $p < 0.05 / 4 = 0.0125$). Finally, given that

the left STC was identified in both the character (showing additive enhancement

and incongruency effect) and pinyin experiments (showing additive enhancement

and congruency-reading correlation), we performed conjunction analysis to examine

the spatial relationship between the two clusters. We also calculated the correlation

between the magnitudes of congruency effect in processing character-sounds and

processing pinyin-sounds.

Significant clusters identified in the whole brain analyses were presented

over a FreeSurfer surface template with BrainNet Viewer (Xia, Wang, & He, 2013).

The peak location was reported in MNI coordinates, and corresponding brain

regions were localized and reported with the AAL atlas implemented in DPABI

(http://rfmri.org/dpabi). Behavioral analysis and ROI analysis were performed with

SPSS Statistics version 24 (SPSS Inc., Chicago, IL, USA).

**3. Results**

**3.1. Behavioral profiles and in-scanner task performance**

Descriptive statistics of demographics, cognitive-linguistic and reading skills for all

41 children are presented in **Table 1**. Significantly positive correlations were

observed between reading measures (all $p$'s < 0.05 / 15, Bonferroni correction; see

**Table 2** for the correlation coefficients and raw $p$-values). Regarding the

relationship with cognitive-linguistic skills, *Word List Reading* was significantly

associated with *Digit RAN* ($r$ = -0.597, $p$ < 0.001 < 0.05 / 15), but not with *Phoneme*

*Deletion* ($r$ = 0.288, $p$ = 0.068 > 0.05 / 15) or *Morphological Production* ($r$ = 0.344, $p$ =

0.028 > 0.05 / 15) with Bonferroni correction. In contrast, *Silent Reading*

*Comprehension* was significantly correlated with morphological production ($r$ =

0.487, $p$ = 0.001 < 0.05 / 15), but not with *Phoneme Deletion* ($r$ = 0.135, $p$ = 0.400 >

0.05 / 15) or *Digit RAN* ($r$ = -0.332, $p$ = 0.034 > 0.05 / 15) with Bonferroni correction.

This pattern indicates that in typically developing Chinese children who have

received formal reading instruction for 3-5 years, oral reading necessitates more

print-to-sound mapping, whereas reading comprehension relies more on semantic

processing.

Regarding the in-scanner task, all the children performed well (character

accuracy: $M$ [$SD$] = 96.5% [4.8], RT $M$ [$SD$] = 512 [81] ms; pinyin accuracy $M$ [$SD$] =

96.4% [3.7], RT $M$ [$SD$] = 530 [80] ms; **Table S1**), and no significant differences

were found between runs (all $p$'s > 0.1). These results indicate that participants

maintained attention to incoming stimuli during the entire scan session. The

results of ANOVA with condition as the repeated measure factor revealed no

significant effect on accuracy ($F$ = 1.00, $p$ = 0.395) and a significant effect on RT in

the character experiment ($F$ = 35.8, $p$ < 0.001). Post-hoc tests revealed the longest

RT for the unimodal visual condition ($p$'s ≤ 0.002 in all comparisons, Bonferroni

correction), followed by the auditory condition (Aud > avC: corrected $p$ < 0.001;

Aud > avI: corrected $p$ < 0.001). There was no difference between the two

crossmodal conditions. In the pinyin experiment, the effect of condition was

significant on both accuracy ($F$ = 10.1, $p$ < 0.001) and RT ($F$ = 54.1, $p$ < 0.001). Post-

hoc tests revealed that while children had the lowest accuracy in the visual

condition ($p$'s ≤ 0.005 in all comparison, Bonferroni correction), no differences were

found between any other pairs of conditions. In the unimodal visual condition,

participants also showed the longest RT (all corrected $p$'s ≤ 0.05), followed by the

auditory condition (Aud > avC: corrected $p$ < 0.001; Aud > avI: corrected $p$ < 0.001).

No differences were observed between the two crossmodal conditions.

**3.2. Brain results: the character experiment**

The bilateral STCs were identified as showing additive enhancement (**Figure 2A**;

**Table 3**; see **Figure S1** for activation in two unimodal conditions and the brain

map for avC > (Aud + Vis) / 2). On the other hand, a significantly incongruency

effect (i.e., avC < avI) was found in the left frontal (peak MNI -40, 14, 32), parietal

(peak MNI -36, -62, 48) and bilateral STCs (peak MNI, left: -54, -22, -2; right: 62, -

34, 10; **Figure 2B, Table 4**) at the FWE corrected threshold of $p$-cluster < 0.05. In

addition, whole-brain regression analysis identified a cluster in the left middle

frontal cortex (peak MNI -32, 10, 42) showing a positive correlation between the

congruency effect and silent reading comprehension proficiency at the FWE

corrected threshold of $p$-cluster < 0.05 (**Figure 2C**, **Table 4**). No regions were

identified as significant in whole-brain regression for oral reading fluency.

Based on the whole brain results, we defined three types of ROI and further

examined their response properties and relationships with reading abilities. The

first type of region (bilateral STCs) showed both additive enhancement and

incongruency effect (**Figure 2D**). No congruency-reading correlations were found

even at uncorrected $p$-voxel < 0.05. The second type of region (left IPL) showed only

incongruency effect in the whole brain analyses (**Figure 2E**). We found no

activation in either the unimodal condition or the audiovisual congruent condition.

No brain-behavior correlations were found in this area neither. The third type of

region (left MFG) showed a significant incongruency effect, of which the magnitude

was further associated with silent reading comprehension proficiency (**Figure 2F**).

In the ROI analysis, we found significant activation for unimodal auditory and

visual conditions in this area but observed no activation in the audiovisual

congruent condition. For the brain-behavior relationship, the magnitude of the

congruency also correlated with oral reading fluency ($r = 0.382$, $p = 0.026$) at an

uncorrected threshold, and showed correlated trends for digit RAN ($r = -0.319$, $p = $

0.066) and MA ($r = 0.300$, $p = 0.084$). Importantly, the positive correlation between

congruency and silent reading comprehension remained significant ($r = 0.494$, $p = $

0.004) even after additionally controlling for oral reading fluency. Finally, we found

that the positive correlation between congruency and reading comprehension was

driven by a negative correlation in the incongruent condition (avC: $r = 0.084$, $p =$

0.637; avI: $r = -0.487$, $p = 0.003$).

**3.3. Brain results: the pinyin experiment**

The audiovisual additive enhancement was detected in the left superior temporal,

inferior frontal, right superior, inferior temporal, and subcortical cortices (**Figure**

**3A**; **Table 3**; see **Figure S1** for activation in the unimodal conditions and brain

map for avC > (Aud + Vis) / 2). No regions showed a significant congruency effect in

either direction in the whole brain analysis, which is unexpected. However, voxel-

wise regression analysis revealed that oral reading fluency was positively correlated

with the magnitude of congruency effect in the left STC (peak MNI -56 -36 14)

(FWE corrected $p$-cluster < 0.05; **Figure 3B**, **Table 4**). No regions were identified

as significant in whole-brain regression for silent reading comprehension

proficiency. Therefore, we defined the area (left STC) showing both additive

enhancement and significant congruency-reading correlation as the ROI (**Figure**

**3C**). In the subsequent analyses, we found no congruency or incongruency effect in

this region even at the uncorrected threshold of $p < 0.05$. In terms of brain-behavior

correlation, the magnitude of congruency effect was correlated with silent reading

comprehension without multiple comparisons correction ($r = 0.371$, $p = 0.020$,

uncorrected). No correlations or correlated trends were found with PA ($r = -0.087$, $p$

$= 0.600$), digit RAN ($r = -0.248$, $p = 0.127$), or MA ($r = 0.195$, $p = 0.235$). Of

importance, the correlation between the congruency effect and oral reading fluency

remained significant ($r = 0.437$, $p = 0.006$) even after additionally controlling for

reading comprehension. Finally, different from the left MFG observed in the

character experiment, the positive congruency-reading correlation in the left STC

was driven by a positive correlation in the congruent condition (avC: $r = 0.381$, $p =$

0.017; avI: $r = -0.039$, $p = 0.815$), suggesting distinct mechanisms.

539           Given that the left STC was identified as functionally important (significant

in both additive enhancement and congruency effect related analyses) in

audiovisual processing in both experiments, we conducted follow-up conjunction

analysis and correlation analysis to examine their relationship. First, the two ROIs

spatially overlapped (**Figure 4A)**. Second, the magnitudes of the congruency effect

in processing character-sound pairs and pinyin-sound pairs were significantly

correlated ($r = 0.512$, $p = 0.002$; **Figure 4B**), while nuisance variables of age and sex

were statistically controlled.

**4. Discussion**

The present study investigated the brain bases underlying audiovisual integration

in characters (deep orthography) and pinyin (a transparent alphabetic coding

system) in typically developing Chinese children. The results revealed both script-

shared and script-unique neural substrates, supporting a universal principle

underlying the neural mechanism of reading-related processes across languages.

Furthermore, the evidence implies that the left MFG links audiovisual integration

of character-sounds with silent reading comprehension proficiency by supporting

automatic semantic access—reflecting a language-specific manifestation, while the

left STG links audiovisual integration of pinyin-sounds with oral reading fluency by

underpinning the shared grapho-phonological mapping component—providing a

possible mechanism with which pinyin helps with Chinese reading development.

**4.1. Neural processes accompanying print-sound integration in scripts**

**with contrasting transparencies**

Several neuroimaging studies have investigated the neural mechanism underlying

print-sound integration and its role in reading development with different

techniques. On the one hand, electroencephalogram (EEG) and MEG were used to

reveal the temporal properties of neural activities (Froyen, van Atteveldt, &

Blomert, 2010; Froyen, Van Atteveldt, Bonte, & Blomert, 2008; Froyen, Bonte, van

Atteveldt, & Blomert, 2009; Raij, Uutela, & Hari, 2000; Xu et al., 2019). These

studies demonstrated that integration occurs preconsciously in expert reading and

is reflected in the component of mismatch negativity/response. On the other hand,

fMRI research provided more precise and detailed information about the spatial

localization of the processing. Among others, the bilateral STCs have been identified

as the most relevant areas. Like other stimuli categories, processing audiovisual

information of scripts induces super-additivity (or mean additive) enhancement in

STC (Beauchamp, 2005; McNorgan & Booth, 2015; van Atteveldt et al., 2004).

Strikingly, differences in brain responses to audiovisual congruent against

incongruent print-sound pairs have been repeatedly reported in these areas (van

Atteveldt et al., 2004; van Atteveldt, Formisano, Blomert, & Goebel, 2007).

Importantly, such effects were observed in passive paradigms, pinpointing the

automatic nature of this effect (Blau et al., 2010; van Atteveldt et al., 2007).

Additive enhancement is more domain-general and has been reported in the

left STC in both transparent (van Atteveldt et al., 2004) and opaque languages

(McNorgan & Booth, 2015). Consistent with previous studies, we observed

audiovisual additive enhancement in the bilateral STCs in processing both

character-sound (deep) and pinyin-sound (shallow) associations. Compared with

additive enhancement, the congruency effect reflects more domain-specific

processing and is affected by factors, such as linguistic characteristics (Holloway et

al., 2015) and reading abilities (Blau et al., 2010; Blau et al., 2009; Karipidis et al.,

2018). Consistency of grapheme-phoneme mapping (orthographic depth or

transparency) is an essential feature in alphabetic languages. Differences in the

direction of the congruency effect and brain regions showing this effect have been

reported between languages with different orthographic depths (Blomert & Froyen,

2010; Holloway et al., 2015). In particular, greater activation in the congruent

condition than incongruent condition is more consistently reported in skilled

readers when processing letter-sound associations in transparent scripts. In

contrast, the opposite pattern is observed in opaque scripts (Holloway et al., 2015).

Chinese is a logographic language with an extremely deep orthography. Therefore,

we expected to observe an incongruency effect. In support of this hypothesis, the

results of the character experiment revealed that four regions (including the

bilateral STC) displayed significant incongruency effects. In contrast, in the absence

of a congruency or incongruency effect in processing pinyin-sound pairs at the group

level, we observed a positive correlation between the magnitude of congruency effect

and oral reading fluency. Higher skill readers showed a congruency effect, whereas

lower skill readers showed an incongruency effect. This pattern is consistent with

previous studies that revealed the lack of a congruency effect in dyslexia in

transparent languages (Blau et al., 2010; Blau et al., 2009), as well as these

investigated brain-behavior correlations at both the phoneme (Karipidis et al., 2016;

Karipidis et al., 2018) and syllable levels (McNorgan et al., 2014; Wang et al., 2020).

**4.2. Left frontal and parietal cortex in implicitly integrating audiovisual**

**information of Chinese characters**

Opaque orthography is not the only linguistic feature that affects the neural

processing of character-sound processing. Chinese characters are morpheme-

syllabic, which means that semantic information is deeply involved. Therefore, in

addition to grapho-phonological mapping, semantic access is an equally or even

more important aspect in reading (Bi, Han, Weekes, & Shu, 2007; Dang, Zhang,

Wang, & Yang, 2018; Liu et al., 2017; Ruan et al., 2018; Yang et al., 2013; Yang et

al., 2006). In this study, the left MFG displayed a significant incongruency effect

and showed a unique association between the magnitude of the effect and silent

reading comprehension proficiency. Considering that this region did not show

additive enhancement, we speculate that the observed incongruency effect is more

likely to be downstream and reflects semantic processing. In previous research,

frontal areas have also been reported along with a broad bilateral fronto-parietal

network. Its function was proposed as task-related top-down modulation (van

Atteveldt et al., 2007) or implicit domain-general conflict detection and resolution

(Holloway et al., 2015). Recently, the incongruency effect measured with MEG was

sourced to the frontal cortex in experienced Chinese adult readers (Xu et al., 2019).

Here, we replicated this finding by revealing the incongruency effect in the left

MFG in a developmental population in Chinese with fMRI and further

demonstrated that the role of this region in the implicit audiovisual integration of

characters is domain-specific and that the information automatically processed in

this region corresponds to semantics. This finding is also consistent with the opinion

that while the fundamental neural circuit for reading is universal, linguistic

features in a given language introduce different demands for specific cognitive

components (Perfetti et al., 2013; Rueckl et al., 2015).

In Chinese, neither grapho-semantic mapping nor phono-semantic mapping

is reliable at the character level. Given that no activation was found in the

audiovisual congruent condition for characters, the incongruency effect in the left

frontal cortex is more likely to reflect a resolution of the conflict between different

meanings accessed via visual and auditory inputs. Interestingly, the positive

correlation between the magnitude of congruency effect and reading comprehension

proficiency was driven by a negative correlation in the incongruent condition. In

other words, individuals with better comprehension performance showed lower

activation for non-matching print-sound pairs. This pattern could be caused by

higher neural efficiency in relevant processing, such as implicit suppression of

conflicting semantic information. It has previously been found that in the initial

stage of establishing a binding between grapheme and phoneme in Swiss-German,

typical readers show higher activation in incongruent conditions, and such

activation decreases as automation is achieved (Plewko et al., 2018).

In addition to the bilateral temporal and left frontal cortices, a left parietal

area also displayed a significant incongruency effect. Like the frontal region, this

region did not show additive enhancement, implying that higher activation in the

incongruent condition is more likely to be a downstream effect. Unlike the left MFG,

no activation was observed in either the unimodal auditory or visual condition, and

there was no congruency-reading correlation. Therefore, although previous studies

regarded the left parietal cortex as a part of the semantic network (Binder & Desai,

2011; Wang, Zhao, Zevin, & Yang, 2016), there was an absence of evidence

indicating that the incongruency effect in this area reflects semantic processing. In

another line of research, compared with English, the left parietal region was more

recruited in Chinese reading for visual orthographic analysis given the visual

complexity of characters (Perfetti et al., 2013). In this case, the incongruency effect

might reflect a conflict of orthographic information. However, here we found no

activation in the visual condition, going against this explanation. Third, although

the left partial cortex has not been associated with print-sound integration

previously in one's native language, a recent training study revealed that this

region was involved in acquiring novel grapho-phonological mapping (Xu,

Kolozsvari, Oostenveld, & Hamalainen, 2020). As its function in audiovisual

processing remains unclear, future studies are needed.

**4.3. The left STG links implicit integration of pinyin-sound pairs with oral**

**reading fluency**

While conquering the sounds of letters to an automatic level is one of the most

critical prerequisites in alphabetic languages (Blomert, 2011; Richlan, 2019),

learning pinyin—a transparent phonological coding system—plays a scaffolding role

at the initial stage of Chinese reading acquisition in Mainland China. Therefore, the

present study not only examined whether and how the neural features of

audiovisual integration are associated with orthographic transparency but also

provided information that helps illustrate the role of pinyin in developing fluent

reading ability in Chinese.

First, widespread brain regions, including the bilateral STCs, showed

additive enhancement in processing pinyin-sound associations, in line with previous

findings in alphabetic languages (Kronschnabel et al., 2014; McNorgan & Booth,

2015). For the congruency effect, we expected to observe a stronger activation in the

congruent condition than in the incongruent condition in the STC based on previous

studies in transparent scripts (Blau et al., 2010; van Atteveldt et al., 2004). At first

glance, no significant congruency or incongruency effect was observed. There could

be several reasons. First, the stimuli we used were at the syllable level. Differences

between the letter-phoneme integration and string-syllable integration have been

reported in previous studies (Kronschnabel et al., 2014). However, the authors also

found that neural response to word-like stimuli is more sensitive to reading

abilities. Thus, the alternative possibility is that the direction and strength of the

congruency effect are associated with reading abilities and masked by group

averaging. In line with the latter, we did observe a positive correlation between the

magnitude of the congruency effect in the left STG and oral reading fluency.

Specifically, children with higher oral reading scores are more likely to show

stronger congruency effects. In contrast, those with lower scores are more likely to

show weaker congruency effects or even incongruency effects. This finding is

consistent with studies, where a reduced congruency effect was found in individuals

with RD in shallow languages (Blau et al., 2010; Blau et al., 2009), studies exploring

brain-behavior correlation in transparent (Wang et al., 2020), and opaque languages

(McNorgan et al., 2014), as well as studies using a training paradigm (Karipidis et

al., 2016; Karipidis et al., 2018). The left STG is the central area representing

phonological information (Boets et al., 2013; Glezer et al., 2016) and is involved in

visual phonological processing in Chinese children and adults (Cao et al., 2010). Its

activation and connectivity change after learning visual-sound mappings (Dehaene

et al., 2010; Li, Xu, Luo, Zeng, & Han, 2020; Thiebaut de Schotten, Cohen,

Amemiya, Braga, & Dehaene, 2014), which can further scaffold later reading

development (Wang, Joanisse, & Booth, 2020). Noteworthily, the positive

correlation between the congruency effect in the left STG for pinyin-sound pairs and

oral reading fluency was driven by a positive correlation in the congruent condition.

This pattern is different from that in the left MFG in processing character-sound

pairs, indicating different mechanisms. It should also be noted that the current

findings were revealed with typical readers. Whether children with RD have deficits

or show other processing patterns requires further investigation.

The significant correlation between the magnitude of congruency effect in the

left STG to pinyin-sound pairs with oral reading fluency implies that learning

pinyin helps children develop the related cognitive components and underlying

neural circuits for fluent oral reading (though causal claims cannot be made from

this study). Compared with cognitive-linguistic skills such as PA and RAN, less

attention has been given to pinyin processing in understanding Chinese reading

development, probably because of its auxiliary role as a scaffold in reading

acquisition but not reading itself. However, although the number of studies is

limited, the pinyin-reading relationship has been demonstrated (Lin et al., 2010;

Pan et al., 2011; Siok & Fletcher, 2001; Zhang, Georgiou, Inoue, Zhong, & Shu,

2020). Pan et al. found that invented pinyin spelling at age 6 independently

predicted reading performance at ages 8 and 10 (Pan et al., 2011). Zhang and

colleagues' recent finding suggests a bidirectional relationship (Zhang et al., 2020).

Several explanations were proposed based on behavioral evidence. One is that

pinyin can help to establish better phonological awareness, the cognitive processing

shared by pinyin and character reading (Ding, Liu, McBride, & Zhang, 2015; Li et

al., 2020; McBride, Wang, & Cheang, 2018; Shu et al., 2008; Yin et al., 2011). Above

and beyond PA, there is also a unique contribution from pinyin processing skills to

character reading (Ju, Zhou, & delMas, 2021). It is reasonable to hypothesize that

pinyin may also help establish the correspondence between print and sound. In

support of this notion, a study revealed that training on pinyin uniquely improved

character-to-syllable mapping in Chinese as a second language learner (Guan et al.,

2011). In the present study, a spatially overlapping region also showed additive

enhancement and incongruency effect in processing characters. Furthermore, we

found that the magnitudes of congruency effect in processing two types of stimuli

were highly correlated. As the left posterior STC is involved in visual rhyming

judgment of Chinese characters and proposed to play a role in mapping between

orthography and phonology (Cao et al., 2010), these findings collectively suggest

that learning pinyin may shape the neural circuit for print-sound integration that

will later be recruited in learning characters.

**4.4. Caveats and future directions**

While this study investigated the neurofunctional basis of integrating audiovisual

information of characters and pinyin, representing one of the first efforts, caution

should be considered when interpreting the results. First, we focused on print-sound

integration at the syllable level, whereas previous studies mainly explored letter-

speech sound associations (but see Kronschnabel et al., 2014; Wang et al., 2020).

This is because Chinese has no GPC rules, and print-sound mapping occurs at the

syllable level. But similar to GPC rules in alphabetic languages, simple characters

are taught at the earliest stages and play fundamental roles in reading

development (Shu et al., 2003; Xing, Shu, & Li, 2004). Second, because our

participants were in grades 3-5, pinyin may play a different role than it does in

earlier stages of reading development. Siok and Fletcher (2001) unveiled significant

correlations between pinyin knowledge and reading in grades 2-5, but not in grade

1. Thus, it could be the particular age range that enabled us to observe the

relationship between print-sound processing in pinyin and oral reading fluency. To

further address this issue, it is necessary to conduct studies following preliterate

children until they achieve fluent reading. Finally, in this study, we uncovered the

brain bases of print-sound integrations in two contrasting scripts and their

relationship with reading performance in typically developing children. One next

question that warrants investigation in the future is whether neural processes are

altered in children with RD.

**5. Conclusion**

In the current study, we investigated regions that are involved in audiovisual

integration of character-sounds and pinyin-sounds. The results suggest the bilateral

STCs as the universal brain substrates underlying audiovisual integration across

scripts. At the same time, the precise manifestation of specific effects is influenced

by linguistic features in the given language. In particular, the left MFG is strongly

involved in the audiovisual integration of characters and correlated with reading

comprehension, suggesting that semantic representations were more efficiently

accessed and processed in readers with higher comprehension proficiency. In

contrast, the brain-behavior correlation in the left STC implies that pinyin may

help children develop better reading skills by shaping the shared component of

grapho-phonological mapping and shaping the underlying brain circuits.

**References**

Bakos, S., Landerl, K., Bartling, J., Schulte-Körne, G., & Moll, K. (2017). Deficits in

Letter-Speech Sound Associations but Intact Visual Conflict Processing in

Dyslexia: Results from a Novel ERP-Paradigm. *Front Hum Neurosci, 11*(116).

Beauchamp, M. S. (2005). Statistical criteria in FMRI studies of multisensory

integration. *Neuroinformatics, 3*(2), 93-113.

Bi, Y., Han, Z., Weekes, B., & Shu, H. (2007). The interaction between semantic and

the nonsemantic systems in reading: Evidence from Chinese.

*Neuropsychologia, 45*(12), 2660-2673.

Binder, J. R., & Desai, R. H. (2011). The neurobiology of semantic memory. *Trends

Cogn Sci, 15*(11), 527-536.

Blau, V., Reithler, J., van Atteveldt, N., Seitz, J., Gerretsen, P., Goebel, R., &

Blomert, L. (2010). Deviant processing of letters and speech sounds as

proximate cause of reading failure: a functional magnetic resonance imaging

study of dyslexic children. *Brain, 133*(Pt 3), 868-879.

Blau, V., van Atteveldt, N., Ekkebus, M., Goebel, R., & Blomert, L. (2009). Reduced

neural integration of letters and speech sounds links phonological and

reading deficits in adult dyslexia. *Current Biology, 19*(6), 503-508.

Blomert, L. (2011). The neural signature of orthographic-phonological binding in

successful and failing reading development. *Neuroimage, 57*(3), 695-703.

Blomert, L., & Froyen, D. (2010). Multi-sensory learning and learning to read. *Int J*

*Psychophysiol, 77*(3), 195-204.

Blomert, L., & Willems, G. (2010). Is there a causal link from a phonological

awareness deficit to reading failure in children at familial risk for dyslexia?

*Dyslexia, 16*(4), 300-317.

Boets, B., Op de Beeck, H. P., Vandermosten, M., Scott, S. K., Gillebert, C. R.,

Mantini, D., . . . Ghesquiere, P. (2013). Intact but less accessible phonetic

representations in adults with dyslexia. *Science, 342*(6163), 1251-1254.

Borleffs, E., Maassen, B. A. M., Lyytinen, H., & Zwarts, F. (2018). Cracking the

Code: The Impact of Orthographic Transparency and Morphological-Syllabic

Complexity on Reading and Developmental Dyslexia. *Front Psychol, 9*(2534),

2534.

Cao, F., Lee, R., Shu, H., Yang, Y. H., Xu, G. Q., Li, K. C., & Booth, J. R. (2010).

Cultural Constraints on Brain Development: Evidence from a Developmental

Study of Visual Word Processing in Mandarin Chinese. *Cereb Cortex, 20*(5),

1223-1233.

Chan, C. K. K., & Siegel, L. S. (2001). Phonological Processing in Reading Chinese

among Normally Achieving and Poor Readers. *J Exp Child Psychol, 80*(1), 23-

43.

Cui, Z., Xia, Z., Su, M., Shu, H., & Gong, G. (2016). Disrupted white matter

connectivity underlying developmental dyslexia: A machine learning

approach. *Human Brain Mapping, 37*(4), 1443-1458.

Dang, M., Zhang, R., Wang, X., & Yang, J. (2018). The Interaction Between

Phonological and Semantic Processing in Reading Chinese Characters. *Front

Psychol, 9*, 2748.

Dehaene, S., Pegado, F., Braga, L. W., Ventura, P., Nunes, G., Jobert, A., . . . Cohen,

L. (2010). How Learning to Read Changes the Cortical Networks for Vision

and Language. *Science, 330*(6009), 1359-1364.

Ding, Y., Liu, R. D., McBride, C., & Zhang, D. (2015). Pinyin Invented Spelling in

Mandarin Chinese-Speaking Children With and Without Reading

Difficulties. *J Learn Disabil, 48*(6), 635-645.

Froyen, D., van Atteveldt, N., & Blomert, L. (2010). Exploring the Role of Low Level

Visual Processing in Letter-Speech Sound Integration: A Visual MMN Study.

*Front Integr Neurosci, 4*, 9.

Froyen, D., Van Atteveldt, N., Bonte, M., & Blomert, L. (2008). Cross-modal

enhancement of the MMN to speech-sounds indicates early and automatic

integration of letters and speech-sounds. *Neurosci Lett, 430*(1), 23-28.

Froyen, D. J. W., Bonte, M. L., van Atteveldt, N., & Blomert, L. (2009). The Long

Road to Automation: Neurocognitive Development of Letter-Speech Sound

Processing. *J Cogn Neurosci, 21*(3), 567-580.

Glezer, L. S., Eden, G., Jiang, X., Luetje, M., Napoliello, E., Kim, J., & Riesenhuber,

M. (2016). Uncovering phonological and orthographic selectivity across the

reading network using fMRI-RA. *Neuroimage, 138*, 248-256.

Guan, C. Q., Liu, Y., Chan, D. H. L., Ye, F., & Perfetti, C. A. (2011). Writing

strengthens orthography and alphabetic-coding strengthens phonology in

learning to read Chinese. *Journal of Educational Psychology, 103*(3), 509-522.

Holloway, I. D., van Atteveldt, N., Blomert, L., & Ansari, D. (2015). Orthographic

Dependency in the Neural Correlates of Reading: Evidence from Audiovisual

Integration in English Readers. *Cereb Cortex, 25*(6), 1544-1553.

I Karipidis, I., Pleisch, G., Röthlisberger, M., Hofstetter, C., Dornbierer, D.,

Stämpfli, P., & Brem, S. (2016). Neural initialization of audiovisual

integration in prereaders at varying risk for developmental dyslexia. *Human*

*Brain Mapping*.

Ju, Z., Zhou, Y., & delMas, R. (2021). The contributions of separate pinyin skills and

oral vocabulary to Chinese word reading of U.S. Mandarin immersion third

graders. *Reading and Writing*.

Kanai, R., & Rees, G. (2011). The structural basis of inter-individual differences in

human behaviour and cognition. *Nature Reviews Neuroscience, 12*(4), 231-

242.

Karipidis, I. I., Pleisch, G., Brandeis, D., Roth, A., Röthlisberger, M., Schneebeli, M.,

. . . Brem, S. (2018). Simulating reading acquisition: The link between

reading outcome and multimodal brain signatures of letter–speech sound

learning in prereaders. *Scientific Reports, 8*(1), 7121.

Kast, M., Bezzola, L., Jancke, L., & Meyer, M. (2011). Multi- and unisensory

decoding of words and nonwords result in differential brain responses in

dyslexic and nondyslexic adults. *Brain and Language, 119*(3), 136-148.

Kidd, E., Donnelly, S., & Christiansen, M. H. (2018). Individual Differences in

Language Acquisition and Processing. *Trends Cogn Sci, 22*(2), 154-169.

Kronschnabel, J., Brem, S., Maurer, U., & Brandeis, D. (2014). The level of

audiovisual print-speech integration deficits in dyslexia. *Neuropsychologia,*

*62*, 245-261.

Lei, L., Pan, J., Liu, H., McBride-Chang, C., Li, H., Zhang, Y., . . . Shu, H. (2011).

Developmental trajectories of reading development and impairment from

ages 3 to 8 years in Chinese children. *J Child Psychol Psychiatry, 52*(2), 212-

220.

Li, H., Shu, H., McBride-Chang, C., Liu, H., & Peng, H. (2012). Chinese children's

character recognition: Visuo-orthographic, phonological processing and

morphological skills. *Journal of Research in Reading, 35*(3), 287-307.

Li, M., Xu, Y., Luo, X., Zeng, J., & Han, Z. (2020). Linguistic experience acquisition

for novel stimuli selectively activates the neural network of the visual word

form area. *Neuroimage, 215*, 116838.

Li, Y., Chen, X., Li, H., Sheng, X., Chen, L., Richardson, U., & Lyytinen, H. (2020).

A computer-based Pinyin intervention for disadvantaged children in China:

Effects on Pinyin skills, phonological awareness, and character reading.

*Dyslexia, 26*(4), 377-393.

Lin, D., McBride-Chang, C., Shu, H., Zhang, Y., Li, H., Zhang, J., . . . Levin, I.

(2010). Small Wins Big:Analytic Pinyin Skills Promote Chinese Word

Reading. *Psychol Sci, 21*(8), 1117-1122.

Liu, Y., Georgiou, G. K., Zhang, Y., Li, H., Liu, H., Song, S., . . . Shu, H. (2017).

Contribution of cognitive and linguistic skills to word-reading accuracy and

fluency in Chinese. *International Journal of Educational Research, 82*, 75-90.

McBride, C., Wang, Y., & Cheang, L. M.-L. (2018). Dyslexia in Chinese. *Current*

*Developmental Disorders Reports, 5*(4), 217-225.

McNorgan, C., Awati, N., Desroches, A. S., & Booth, J. R. (2014). Multimodal

Lexical Processing in Auditory Cortex Is Literacy Skill Dependent. *Cereb*

*Cortex, 24*(9), 2464-2475.

McNorgan, C., & Booth, J. R. (2015). Skill dependent audiovisual integration in the

fusiform induces repetition suppression. *Brain and Language, 141*, 110-123.

McNorgan, C., Randazzo-Wagner, M., & Booth, J. R. (2013). Cross-modal

integration in the brain is related to phonological awareness only in typical

readers, not in those with reading difficulty. *Front Hum Neurosci, 7*(388),

388.

Pan, J., McBride-Chang, C., Shu, H., Liu, H., Zhang, Y., & Li, H. (2011). What is in

the naming? A 5-year longitudinal study of early rapid naming and

phonological sensitivity in relation to subsequent reading skills in both native

Chinese and English as a second language. *Journal of Educational*

*Psychology, 103*(4), 897-908.

Perfetti, C., Cao, F., & Booth, J. (2013). Specialization and Universals in the

Development of Reading Skill: How Chinese Research Informs a Universal

Science of Reading. *Sci Stud Read, 17*(1), 5-21.

Perrachione, T. K., & Ghosh, S. S. (2013). Optimized Design and Analysis of Sparse-

Sampling fMRI Experiments. *Frontiers in Neuroscience, 7*(55).

Plewko, J., Chyl, K., Bola, Ł., Łuniewska, M., Dębska, A., Banaszkiewicz, A., . . .

Jednoróg, K. (2018). Letter and Speech Sound Association in Emerging

Readers With Familial Risk of Dyslexia. *Front Hum Neurosci, 12*(393).

Raij, T., Uutela, K., & Hari, R. (2000). Audiovisual integration of letters in the

human brain. *Neuron, 28*(2), 617-625.

Richlan, F. (2014). Functional neuroanatomy of developmental dyslexia: the role of

orthographic depth. *Front Hum Neurosci, 8*, 347.

Richlan, F. (2019). The Functional Neuroanatomy of Letter-Speech Sound

Integration and Its Relation to Brain Abnormalities in Developmental

Dyslexia. *Front Hum Neurosci, 13*(21).

Ruan, Y., Georgiou, G. K., Song, S., Li, Y., & Shu, H. (2018). Does writing system

influence the associations between phonological awareness, morphological

awareness, and reading? A meta-analysis. *Journal of Educational*

*Psychology, 110*(2), 180.

Rueckl, J. G., Paz-Alonso, P. M., Molfese, P. J., Kuo, W. J., Bick, A., Frost, S. J., . . .

921            Frost, R. (2015). Universal brain signature of proficient reading: Evidence

922            from four contrasting languages. *Proc Natl Acad Sci U S A, 112*(50), 15510-

923            15515.

924    Seghier, M. L., & Price, C. J. (2018). Interpreting and Utilising Intersubject

925            Variability in Brain Function. *Trends Cogn Sci, 22*(6), 517-530.

926    Shu, H., Chen, X., Anderson, R. C., Wu, N., & Xuan, Y. (2003). Properties of school

927            Chinese: Implications for learning to read. *Child development, 74*(1), 27-47.

928    Shu, H., McBride-Chang, C., Wu, S., & Liu, H. Y. (2006). Understanding Chinese

929            developmental dyslexia: Morphological awareness as a core cognitive

930            construct. *Journal of Educational Psychology, 98*(1), 122-133.

931    Shu, H., Peng, H., & McBride-Chang, C. (2008). Phonological awareness in young

932            Chinese children. *Dev Sci, 11*(1), 171-181.

933    Siok, W. T., & Fletcher, P. (2001). The role of phonological awareness and visual-

934            orthographic skills in Chinese reading acquisition. *Dev Psychol, 37*(6), 886-

935            899.

936    Stein, B. E., & Stanford, T. R. (2008). Multisensory integration: current issues from

937            the perspective of the single neuron. *Nat Rev Neurosci, 9*(4), 255-266.

938    Talavage, T. M., & Hall, D. A. (2012). How challenges in auditory fMRI led to

939            general advancements for the field. *Neuroimage, 62*(2), 641-647.

Thiebaut de Schotten, M., Cohen, L., Amemiya, E., Braga, L. W., & Dehaene, S.

(2014). Learning to read improves the structure of the arcuate fasciculus.

*Cereb Cortex, 24*(4), 989-995.

van Atteveldt, N., Formisano, E., Goebel, R., & Blomert, L. (2004). Integration of

letters and speech sounds in the human brain. *Neuron, 43*(2), 271-282.

van Atteveldt, N. M., Formisano, E., Blomert, L., & Goebel, R. (2007). The effect of

temporal asynchrony on the multisensory integration of letters and speech

sounds. *Cereb Cortex, 17*(4), 962-974.

van Atteveldt, N. M., Formisano, E., Goebel, R., & Blomert, L. (2007). Top–down

task effects overrule automatic multisensory responses to letter–sound pairs

in auditory association cortex. *Neuroimage, 36*(4), 1345-1360.

Wang, F., Karipidis, II, Pleisch, G., Fraga-Gonzalez, G., & Brem, S. (2020).

Development of Print-Speech Integration in the Brain of Beginning Readers

With Varying Reading Skills. *Front Hum Neurosci, 14*(289), 289.

Wang, J., Joanisse, M. F., & Booth, J. R. (2020). Neural representations of

phonology in temporal cortex scaffold longitudinal reading gains in 5- to 7-

year-old children. *Neuroimage, 207*, 116359.

Wang, X., Zhao, R., Zevin, J. D., & Yang, J. (2016). The Neural Correlates of the

Interaction between Semantic and Phonological Processing for Chinese

Character Reading. *Front Psychol, 7*.

Wechsler, D. (1974). *WISC-R, Wechsler intelligence scale for children, revised*:

Psychological Corporation.

Wu, C. Y., Ho, M. H., & Chen, S. H. (2012). A meta-analysis of fMRI studies on

Chinese orthographic, phonological, and semantic processing. *Neuroimage,*

*63*(1), 381-391.

Xia, M., Wang, J., & He, Y. (2013). BrainNet Viewer: a network visualization tool

for human brain connectomics. *PLoS One, 8*(7), e68910.

Xia, Z., Hoeft, F., Zhang, L., & Shu, H. (2016). Neuroanatomical anomalies of

dyslexia: Disambiguating the effects of disorder, performance, and

maturation. *Neuropsychologia, 81*, 68-78.

Xia, Z., Zhang, L., Hoeft, F., Gu, B., Gong, G., & Shu, H. (2018). Neural correlates of

oral word reading, silent reading comprehension, and cognitive

subcomponents. *International Journal of Behavioral Development, 0*(0),

0165025417727872.

Xing, H., Shu, H., & Li, P. (2004). The acquisition of Chinese characters: Corpus

analyses and connectionist simulations. *Journal of Cognitive Science, 5*(1), 1-

49.

Xu, W., Kolozsvari, O. B., Monto, S. P., & Hämäläinen, J. A. (2018). Brain

Responses to Letters and Speech Sounds and Their Correlations With

Cognitive Skills Related to Reading in Children. *Front Hum Neurosci,*

*12*(304).

Xu, W., Kolozsvari, O. B., Oostenveld, R., & Hamalainen, J. A. (2020). Rapid

changes in brain activity during learning of grapheme-phoneme associations

in adults. *Neuroimage, 220*, 117058.

Xu, W., Kolozsvari, O. B., Oostenveld, R., Leppanen, P. H. T., & Hamalainen, J. A.

(2019). Audiovisual Processing of Chinese Characters Elicits Suppression and

Congruency Effects in MEG. *Front Hum Neurosci, 13*(18), 18.

Xue, J., Shu, H., Li, H., Li, W., & Tian, X. J. J. o. P. R. (2013). The Stability of

Literacy-Related Cognitive Contributions to Chinese Character Naming and

Reading Fluency. *42*(5), 433-450.

Yan, M., Miller, K. F., Li, H., & Shu, H. (2008). What is the place for Pinyin in

beginning Chinese reading? Evidence from eye movements. In *Cognitive and*

*cultural influences on eye movements.* (pp. 343-360). Tianjin, China: Tianjin

People's Publishing House.

Yang, J., Shu, H., McCandliss, B. D., & Zevin, J. D. (2013). Orthographic influences

on division of labor in learning to read Chinese and English: Insights from

computational modeling. *Biling (Camb Engl), 16*(Spec Iss 2), 354-366.

Yang, J., Zevin, J. D., Shu, H., McCandliss, B. D., & Li, P. (2006). *A "triangle model"*

*of Chinese reading.* Paper presented at the 28th annual conference of the

cognitive science society, Mahwah, NJ.

Yin, L., Li, W., Chen, X., Anderson, R. C., Zhang, J., Shu, H., & Jiang, W. (2011).

The role of tone awareness and pinyin knowledge in Chinese reading. *Writing*

*Systems Research, 3*(1), 59-68.

Zhang, S., Georgiou, G. K., Inoue, T., Zhong, W., & Shu, H. (2020). Do pinyin and

character recognition help each other grow? *Early Childhood Research*

*Quarterly, 53*, 476-483.

Zhang, Y., Zhang, L., Shu, H., Xi, J., Wu, H., Zhang, Y., & Li, P. (2012). Universality

of categorical perception deficit in developmental dyslexia: an investigation of

Mandarin Chinese tones. *Journal of Child Psychology and Psychiatry, 53*(8),

874-882.

Zhao, J., Wang, X., Frost, S. J., Sun, W., Fang, S.-Y., Mencl, W. E., . . . Rueckl, J. G.

(2014). Neural division of labor in reading is constrained by culture:

A training study of reading Chinese characters. *Cortex, 53*, 90-106.

1013

**Table 1** Demographics and behavior profiles (n = 41)

| Measure | Mean | SD | Range |
|---|---|---|---|
| Age (month) | 126 | 8 | 110-141 |
| Grade (3rd / 4th / 5th) | 4 / 23 / 14 | | |
| Sex (female / male) | 27 / 14 | | |
| Verbal IQ (standard score) | 107 | 13 | 85-145 |
| Performance IQ (standard score) | 114 | 13 | 75-137 |
| Full-scale IQ (standard score) | 111 | 12 | 80-143 |
| Character Recognition (corrected item) | 119 | 12 | 86-146 |
| Word List Reading (word / minute) | 90 | 19 | 55-140 |
| Silent Reading Comprehension (character / minute) | 327 | 119 | 132-565 |
| Phoneme Deletion (corrected item) | 21 | 4 | 7-25 |
| Rapid Naming (second) | 18 | 3 | 11-24 |
| Morphological Production (item) | 24 | 4 | 13-30 |
| Number of 'bad' time points per fMRI run | | | |
| Character Experiment [a] | 2.4 | 3.37 | 0-10 |
| Pinyin Experiment | 0.74 | 1.54 | 0-9 |

*Note.* [a] n = 36

1014

**Table 2** Correlation between reading and cognitive-linguistic skills (n = 41)

| Measures | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|---|
| 1 Character Recognition | --- | | | | | |
| 2 Word List Reading | $r = 0.559$ | --- | | | | |
| | $p < 0.001$ | | | | | |
| 3 Silent Reading Comprehension | $r = 0.647$ | $r = 0.701$ | --- | | | |
| | $p < 0.001$ | $p < 0.001$ | | | | |
| 4 Phoneme Deletion | $r = 0.286$ | $r = 0.288$ | $r = 0.135$ | --- | | |
| | $p = 0.070$ | $p = 0.068$ | $p = 0.400$ | | | |
| 5 Rapid Naming | $r = -0.357$ | $r = -0.597$ | $r = -0.332$ | $r = -0.160$ | --- | |
| | $p = 0.022$ | $p < 0.001$ | $p = 0.034$ | $p = 0.319$ | | |
| 6 Morphological Production | $r = 0.566$ | $r = 0.344$ | $r = 0.487$ | $r = 0.122$ | $r = -0.117$ | --- |
| | $p < 0.001$ | $p = 0.028$ | $p = 0.001$ | $p = 0.446$ | $p = 0.466$ | |

1015

**Table 3** Cluster identified as showing audiovisual additive enhancement in each experiment

| Experiment | # Voxel | Brain area |
| --- | --- | --- |
| *Character* | 822 | Temporal_Sup_R, Temporal_Mid_R |
| | 354 | Temporal_Sup_L, Temporal_Mid_L |
| *Pinyin* | 2752 | Temporal_Mid_L, Temporal_Sup_L, Rolandic_Oper_L, Occipital_Mid_L, Temporal_Inf_L, SupraMarginal_L, Fusiform_L, Postcentral_L, Heschl_L, Temporal_Pole_Sup_L, Occipital_Inf_L |
| | 1532 | Temporal_Sup_R, Temporal_Mid_R, SupraMarginal_R |
| | 967 | Temporal_Mid_R, Temporal_Inf_R, Fusiform_R, Cerebelum_6_R, Occipital_Inf_R, Occipital_Mid_R |
| | 943 | Precentral_L, Frontal_Inf_Oper_L, Frontal_Inf_Tri_L, Rolandic_Oper_L |
| | 936 | Putamen_L, Thalamus_L, Insula_L, Vermis_3 |

*Note*. Brain area labeling is based on the AAL atlas (> 20 overlapping voxels).

1016

**Table 4** Results of the voxel-wise whole brain analyses on congruency effect and congruency-reading relationship

| Experiment and contrast | Brain area (AAL) | Direction | # Voxel | Peak $T$ | X | Y | Z |
|---|---|---|---|---|---|---|---|
| *Character* | | | | | | | |
| Congruency / Incongruency | Temporal_Mid_L, Temporal_Sup_L, Temporal_Pole_Sup_L, Rolandic_Oper_L, Hippocampus_L | avC < avI | 3310 | -7.92 | -54 | -22 | -2 |
| | Precuneus_R, Parietal_Inf_L, Parietal_Sup_L, Precuneus_L, Cuneus_R, Angular_L, Precentral_L, Occipital_Sup_L, Paracentral_Lobule_L, Occipital_Sup_R, occipital_Mid_L | avC < avI | 3308 | -5.79 | -36 | -62 | 48 |
| | Temporal_Sup_R, Temporal_Mid_R, Temporal_Pole_Sup_R, Temporal_Inf_R | avC < avI | 2319 | -6.68 | 62 | -34 | 10 |
| | Precentral_L, Frontal_Inf_Tri_L, Frontal_Mid_L, Frontal_Inf_Oper_L | avC < avI | 2894 | 5.83 | -40 | 14 | 32 |
| Congruency-ORF Correlation | | *n.s.* | | | | | |
| Congruency-SRC Correlation | Frontal_Mid_L, Precentral_L | positive | 558 | 4.07 | -32 | 10 | 42 |
| *Pinyin* | | | | | | | |
| Congruency / Incongruency | | *n.s.* | | | | | |
| Congruency-ORF correlation | Temporal_Sup_L, Temporal_Mid_L, Thalamus_L | positive | 468 | 4.32 | -40 | -42 | 6 |

| | |
|---|---|
| Congruency-SRC correlation | *n.s.* |

*Note*: Brain area labeling is based on the AAL atlas (> 20 overlapping voxels). Abbreviations: avC = audiovisual congruent, avI = audiovisual incongruent, ORF = oral reading fluency, SRC = silent reading comprehension.

1017

**Figure 1** Materials and experimental design. Example stimuli used in the character (A) and pinyin (B) experiments. (C) Schematic illustration of the fMRI procedure. (D) Auditory and visual targets in the in-scanner task.

**Figure 2** Results of the character experiment. (A) Clusters (yellow) showing

audiovisual additive enhancement are identified with the mean rule (Beauchamp,

2005) consisting of two criteria: (1) The region should activate in both unisensory

conditions (i.e., Aud > 0, Vis > 0); (2) The audiovisual congruent stimuli should induce

stronger activation than the mean activation of the unisensory conditions, i.e., avC >

(Aud + Vis) / 2. An uncorrected threshold of $p$-voxel < 0.05 is used for each contrast,

resulting in a conjoint probability of $p$-voxel < 0.05^3 = 0.000125. A cluster containing

no less than 200 continuous voxels was considered to show significant additive

enhancement. (B) Areas showing a significant congruency effect (blue = incongruency,

i.e., avC < avI). Threshold: cluster forming $p$-voxel < 0.005, Family-Wise Error (FWE)

corrected $p$-cluster < 0.05. (C) The region showing a significant correlation between

the magnitude of congruency effect and reading scores. Threshold: cluster forming $p$-

voxel < 0.005, FWE corrected $p$-cluster < 0.05. (D, E, F) The areas showing both

additive enhancement and congruency effect (red cluster), congruency effect only

(green cluster), and both congruency effect and congruency-reading correlation (violet

cluster) were defined as the regions-of-interest (ROIs). For each ROI, a bar plot

displaying activation for each condition and a boxplot displaying the contrast avC -

(Aud + Vis) / 2 are presented. A scatter plot is presented in the ROI showing a

significant correlation between the magnitude of congruency effect and reading

(residuals were obtained by regressing out nuisance variables of age and sex with a

general linear model). *Abbreviations*: Aud = unisensory auditory, Vis = unisensory

visual, avC = audiovisual congruent, avI = audiovisual incongruent, L = left

hemisphere, R = right hemisphere; *** $p$ < 0.001, *n.s.* = non-significant.

**Figure 3** Results of the pinyin experiment. (A) Clusters (yellow) showing audiovisual additive enhancement are identified with the mean rule (Beauchamp, 2005), which consists of two criteria: (1) The region should activate in both unisensory conditions (i.e., Aud > 0, Vis > 0); (2) The audiovisual congruent stimuli pairs should induce stronger activation than the average activation of the auditory and visual conditions, i.e., avC > (Aud + Vis) / 2. An uncorrected threshold $p$-voxel < 0.05 is used for each contrast, resulting in a conjoint probability of $p$-voxel < $0.05^3$ = 0.000125. A cluster consisting of no less than 200 continuous voxels under the threshold is considered to show additive enhancement. (B) A cluster displaying a significant correlation between the magnitude of congruency effect and oral reading fluency. Threshold: cluster forming $p$-voxel < 0.005, Family-Wise Error (FWE) corrected $p$-cluster < 0.05. (C) The intersection between areas showing mean additive audiovisual enhancement

and areas showing association with reading is defined as the region-of-interest (ROI).

The bar plot shows activation in each condition. There was no significant difference

between the audiovisual congruent condition (avC) and incongruent condition (avI).

Boxplot displays the contrast avC - (Aud, Vis) / 2. The scatter plot shows the

correlation between the magnitude of congruency effect and oral reading fluency

(residuals were obtained by regressing out nuisance variables of age and sex with a

general linear model). *Abbreviations*: Aud = unisensory auditory, Vis = unisensory

visual, avC = audiovisual congruent, avI = audiovisual incongruent, L = left

hemisphere, R = right hemisphere; *n.s.* = non-significant.

**Figure 4** (A) The regions-of-interest (ROIs) in the left superior temporal cortex

defined in the character experiment (green) and pinyin experiment (red), as well as

their intersection (yellow), are presented on a template brain. (B) The scatter plot

shows the significant correlation between the magnitude of congruency effect in the

character ROI and the pinyin ROI (residuals were obtained by regressing out

nuisance variables of age and sex with a general linear model).