1 Neurofunctional mechanisms underlying audiovisual

2 integration of characters and *pinyin* in Chinese

3 children

Zhichao Xia [a, b] [*], Ting Yang [c, d] [*], Xin Cui [a], Fumiko Hoeft [e, f, g, h], Hong Liu [c, d, e], Hua

Shu [a] [†], Xiangping Liu [c, d] [†]

[a] State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern

Institute for Brain Research, Beijing Normal University, China

[b] School of Systems Science, Beijing Normal University, China

[c] Faculty of Psychology, Beijing Normal University, China

[d] Beijing Key Laboratory of Applied Experimental Psychology, National

Demonstration Center for Experimental Psychology Education, Faculty of

Psychology, Beijing Normal University, China

[e] Department of Psychological Sciences and Brain Imaging Research Center,

University of Connecticut, USA

[f] Department of Psychiatry and Weill Institute for Neurosciences and Dyslexia

Center, University of California, San Francisco, USA

[g] Haskins Laboratories, USA

[h] Department of Neuropsychiatry, Keio University School of Medicine, Japan

[*] These authors contributed equally to this work

† Corresponding authors: Dr. Hua Shu, State Key Lab of Cognitive Neuroscience

and Learning, Beijing Normal University, China. Email: shuhua@bnu.edu.cn; Dr.

Xiangping Liu, Faculty of Psychology, Beijing Normal University, China. Email:

lxp599@163.com

Running Title: Brain basis of audiovisual integration

Number of pages: 46

Number of words in Abstract: 250, Introduction: 1053, Discussion: 1974.

Number of Tables: 2, Figures: 4

31 Abstract

Efficient integration of grapheme and phoneme is crucial for reading development

in alphabetic languages, and superior temporal cortex (STC) has been demonstrated

to be the most critical region for this process. To determine whether a similar

neural mechanism underlies such processing in non-alphabetic languages, and to

explore its relationship with reading, we conducted a functional magnetic resonance

imaging study in typically developing Chinese children. Highly frequent

pictographic characters and *pinyin*, a transparent alphabetic coding system that

assists individuals to learn the pronunciation of new characters were investigated.

In support of the orthographic depth hypothesis developed in alphabetic languages,

reverse congruency effect (i.e., higher activation in incongruent condition compared

to congruent condition) was identified in the left inferior frontal gyrus (IFG) and

bilateral STC for character (deep orthography) conditions. Furthermore, correlation

analysis revealed that the congruency contrast in the left IFG was associated with

proficiency in reading comprehension and morphological awareness, suggesting

reading-related semantic access during the implicit integration of multisensory

information regarding characters. For *pinyin* (shallow orthography) conditions,

while no regions showed a significant congruency effect at the group level in either

direction, the congruency contrast in the left superior temporal gyrus was positively

correlated with oral reading performance. This observation is consistent with

findings in transparent scripts on reading disorders. Taken together, these results

support and generalize the orthographic depth hypothesis to a logographic

language, highlight the role of the left IFG in character-syllable integration, and

underscore the role *pinyin* could play in developing fluent reading in Chinese

children.

Keywords

Highlights

• The left IFG and bilateral STC are involved in audiovisual integration of

characters.

• The congruency contrast in the left IFG in response to characters corresponds to

semantic access.

• The congruency contrast in the left STG in response to *pinyin* is correlated with

print-to-sound mapping.

• These findings support the orthographic depth hypothesis in a nonalphabetic

language.

• The results imply a close link between *pinyin* and Chinese reading acquisition.

1. Introduction

Audiovisual integration of print and speech sound is a prerequisite for successful reading acquisition (Blomert & Froyen, 2010; Richlan, 2019). Behavioral and neurofunctional evidence has revealed that the lack of automatic print-to-sound integration uniquely accounts for the reading failure in shallow orthographies beyond commonly reported reading-related cognitive-linguistic skills, such as phonological awareness (PA) (Bakos, Landerl, Bartling, Schulte-Körne, & Moll, 2017; Blau et al., 2010; Leo Blomert & Willems, 2010).

Better knowledge of the neural basis underlying audiovisual integration can further our understanding of how this process contributes to (a)typical reading development and provide objective training targets and evaluation indices for children with reading disorders (RD). To achieve this goal, neuroimaging techniques, especially functional magnetic resonance imaging (fMRI), have been widely applied in the past two decades. Across these studies, distinguishable brain responses to congruent versus incongruent audiovisual stimulus pairs have been demonstrated to be the most reliable measure of multisensory integration and have been named the 'congruency effect' (Blau et al., 2010; Blau, van Atteveldt, Ekkebus, Goebel, & Blomert, 2009; Holloway, van Atteveldt, Blomert, & Ansari, 2015; van Atteveldt, Formisano, Goebel, & Blomert, 2004; van Atteveldt, Formisano, Goebel, & Blomert, 2007). For example, in a pioneering study by van Atteveldt et al. (2004), superior temporal cortex (STC) showed a stronger activation in congruent than in

incongruent conditions. Furthermore, while the involvement of STC in the

audiovisual integration of print and speech sound has been replicated regardless of

the level of processing (e.g., phoneme, syllable, or whole word) (Kast, Bezzola,

Jancke, & Meyer, 2011; Mcnorgan, Randazzo-Wagner, & Booth, 2013), researchers

also found that the direction of the effect and the area displaying such effect are

influenced by multiple factors, where orthographic depth was an important one

(Holloway et al., 2015; Xu, Kolozsvari, Monto, & Hämäläinen, 2018).

Orthographic depth refers to the consistency of mapping between visual and

phonological forms in a writing system (e.g., grapheme-phoneme correspondence in

alphabetic languages, GPC hereafter). The orthographic depth hypothesis would

suggest that the congruency effect during audiovisual integration is in a direction

that is associated with the consistency of GPC. In support of this, researchers have

found that native speakers of English (deep orthography) showed a reverse

congruency effect (aka incongruency effect, higher activation for incongruent

condition), whereas there is a congruency effect (higher activation for congruent

condition) observed in Dutch (shallow orthography) (Holloway et al., 2015). More

importantly, when the same English-speaking participants processed digits, where

there is one-to-one mapping, the effect became positive. Although the conclusion is

clear, these findings are restricted to alphabetic languages. In order to broaden the

applicability of this hypothesis, research in non-alphabetic languages is necessary.

Chinese is particularly suited to for answer this question, given its

logographic nature and extremely deep orthography. In general, character is the

basic graphic unit. While different characters may share the same pronunciation

(i.e., homophones), a single character can also have multiple sounds (i.e.,

polyphone). Therefore, the link between print (character) and speech sound

(syllable) is quite opaque. According to the orthographic depth hypothesis, a reverse

congruency effect should be observed. Moreover, since grapho-semantic

correspondence is more systematic in Chinese and plays a role equal to or even

more important than phonological processing in Chinese reading acquisition (Liu et

al., 2017; Ruan, Georgiou, Song, Li, & Shu, 2018; Yang, Shu, McCandliss, & Zevin,

2012), additional regions such as the inferior and middle frontal cortex known to be

involved in semantic processing may be recruited (Wu, Ho, & Chen, 2012; Zhao et

al., 2014). Recently, Xu et al., (2019) conducted a magnetoencephalography (MEG)

study with an one-back task in Chinese adults and identified a reverse congruency

effect across the left frontal cortex and STC. However, some limitations need to be

considered, including small sample size (n = 12), using flat tone speech sounds that

do not exactly correspond to any character, and source reconstruction accuracy of

MEG. Moreover, in experienced readers, audiovisual integration of characters and

corresponding sound is fully automated. Thus, in addition to address these issues,

an important follow-up question should be neurofunctional features of this process

in developing populations. Against this background, the first aim of the current

study is to determine whether there is a cross-modal process supported by STC as

well as other brain areas associated with integrating characters and speech sounds

in typically developing Chinese children.

Of note, learning to read in Chinese is a longer process than learning most

alphabetic languages (Shu, Chen, Anderson, Wu, & Xuan, 2003). At the very

beginning stage, a small set of pictographic characters that are used as radicals in

composite characters are taught first, since more than 80% of Chinese characters

are phonograms that comprise semantic and phonetic radicals (Chan & Siegel,

2001). Given its important role, in this study, we specifically focused on these early-

acquired single-element characters. Importantly, because pictographic characters do

not contain radicals as phonological/semantic cues, the mapping between visual and

sound/meaning is direct and completely arbitrary. Moreover, Chinese offers a

valuable opportunity to examine the orthographic depth hypothesis using two

different scripts in the same brain. *Pinyin* is a Roman alphabetic system used to

represent the sound of characters in Mainland China and Singapore. A child will

learn *pinyin* first and then use it to assist in conquering the pronunciation of new

characters (Guan, Liu, Chan, Ye, & Perfetti, 2011; Yan, Miller, Li, & Shu, 2008). In

other words, it is more like a scaffold during reading acquisition. Since *pinyin*

borrows letters from English and has one-to-one mappings with speech sounds

(phonemes), it is reasonable to predict a similar activation pattern in response to

audiovisual stimulus pairs in *pinyin* and in transparent orthographies. Therefore,

we included *pinyin* in this study additionally to further examine the orthographic

depth hypothesis.

Finally, in addition to examining the congruency effect at the group level,

understanding multisensory integration in relation to individual heterogeneity in

reading is important. However, until now, the majority of research has been done by

comparing children with RD with their typical peers (e.g., Blau et al., 2010; Blau et

al., 2009; Plewko et al., 2018), leaving the brain-reading relationship in typically

developing children underexplored. Therefore, we additionally adopted an

individual differences approach to evaluate reading abilities and cognitive-linguistic

skills in an effect to shed light on whether and how neural circuitries link

multisensory integration of character/*pinyin* with typical reading acquisition.

2. Materials and Methods

2.1. Participants

Thirty children (20 girls; age 111 ~ 140 months, M = 127) were included in this

study. They were selected from the initial sample of 55 typical children according to

the following inclusion criteria: (a) native speaker of Chinese, (b) right-handed

(Oldfield, 1971), (c) normal audition and normal or corrected-to-normal vision, (d)

full-scale intelligence quotient (IQ) $\geq$ 85, (e) no history of neurological or psychiatric

disorders, (f) typical reading (no less than 1 SD below the norm average in the

reading screening task *Character Recognition*) (Xue, Shu, Li, Li, & Tian, 2013), (g)

completed all four task scans, (h) accurate response rate (ACC) for each run > 75%,

(i) no more than 17 (25%) 'bad' volumes in each run (see fMRI preprocessing). These

children learned *pinyin* formally in the first half of the semester in the 1st grade,

before learning characters. All the children and their parents/guardians were

informed about the aim and content of the study and signed written consent before

the experiment was conducted. All the behavioral and neuroimaging data were

collected in 2018. This study received ethical approval from the Institutional Review

Board of State Key Laboratory of Cognitive Neuroscience and Learning at Beijing

Normal University.

2.2. Behavior measures

Each child received testing individually in a silent booth. IQ was measured by the

abbreviated version of the *Chinese Wechsler Intelligence Scale for Children* (*WISC-*

CR) (Wechsler, 1974): *Information*, *Similarities*, and *Digit Span* subtests were used

to estimate verbal IQ and *Picture Completion*, *Block Design*, and *Coding* subtests

were used to evaluate performance IQ.

A set of neuropsychological tasks was used to measure children's reading and

reading-related cognitive-linguistic skills. In brief, as *Character Recognition* has

been used in previous studies for identifying Chinese dyslexic children (Cui, Xia, Su,

Shu, & Gong, 2016; Xia, Hoeft, Zhang, & Shu, 2016), we used it here for the

screening aim. *Word List Reading* and *Silent Reading Comprehension* were used to

measure oral reading fluency and reading comprehension proficiency, respectively.

The former task relies more on grapho-to-phonological access, while the latter relies

more on grapho-to-semantic processing (Xia et al., 2018). Lastly, tasks *Phoneme*

Deletion, *Digit RAN*, and *Morphological Production* tasks were used to measure PA,

RAN, and morphological awareness (MA), which are the most important cognitive-

linguistic skills for reading development in Chinese (Lei et al., 2011).

Character Recognition is a standardized test for estimating the number of

characters that the child has learned. It consists of 150 characters selected from

textbooks for grades 1-6 (Xue et al., 2013). The characters are arranged in the order

according to the grade on 10 sheets of A4 paper. During the task, the participant is

asked to name the characters in sequence until they fail in all 15 items on one page.

Each correct answer is worth 1 point, and the full mark is 150. No time limit was

applied.

Word List Reading consists of 180 high-frequency two-character words

(Zhang et al., 2012). These words are arranged in a 9-column by 20-row matrix in a

sheet of A4 paper. The participant is required to read words aloud in a left-to-right,

up-to-down order as accurately and quickly as possible. The time used to complete

the task is recorded, and an index referring to the number of words the participant

read correctly per minute is calculated as the final score.

Silent Reading Comprehension includes 100 sentences/paragraphs (Lei et al.,

2011). The items are arranged in order based on length. The participant is

requested to read each sentence/paragraph silently and decide the correctness of the

meaning with mark of ✓ or ✗ as fast as possible within 3 minutes. The sum of the

characters in sentences with correct responses is calculated. The final score

indicates the number of characters the participant read in one minute.

Phoneme Deletion is used to assess PA (Li, Shu, McBride-Chang, Liu, &

Peng, 2012). This task includes three sections, and the participant is asked to delete

the initial, middle, or final sound from orally presented syllables. Each section

consists of 3 practice trials and 8-10 testing items. In particular, the participant is

asked to pronounce what is left when a phoneme (consonant/vowel) is deleted from a

given syllable. One correct response is worth 1 point, and the total possible score is

28.

Digit RAN requires the participant to name 50 one-digit numbers on a card

as accurately and rapidly as possible (Liu et al., 2017). Numbers 1, 2, 3, 5, and 8

repeat five times and arrange randomly in a 5-column by 10-row matrix. The task is

administrated twice, and the average time the participant completes the task is

used as the final score.

Morphological Production is used to measure MA at the character level (Shu,

McBride-Chang, Wu, & Liu, 2006). In each trial, a character is orally presented in a

word to the participant. The participant has to respond with two new words. In one

word the given character retains the same meaning as that in the original case,

while in the other word the meaning of the character should be different. This task

contains 15 characters, with a full mark of 30.

2.3. Stimuli and experimental design

In this study, we investigated neurofunctional correlates underlying audiovisual

integration in both characters and *pinyin*. In the character conditions, the stimuli

were visually presented characters and their speech sounds (**Figure 1A**). Fifty-six

highly frequent pictographic characters were selected (Chinese Single-Character

Word Database; https://blclab.org/pyscholinguistic-norms-database/). All the

characters are simple and taught early. In the experiment, visual stimuli were

white characters displayed in 'KaiTi' font and 96 pt at the center of a black

background. The speech sounds were recorded from a native Chinese male with a

sampling rate of 44.1 kHz and 16-bit quantization. The audio files were normalized

to 85 dB and then edited with a bandpass (100-4000 Hz) filter with Audacity

(https://www.audacityteam.org/). The average duration of all sounds is 476.3 (±87.5)

ms, without pre- or post-silence. All the speech sounds can be correctly recognized

easily. For the *pinyin* conditions, the visual stimuli were the *pinyin* spelling of the

characters (in white, 'Century Schoolbook' font, and 90 pt), while the auditory

stimuli were the same as character conditions (**Figure 1B**).

The fMRI procedure was adapted from Blau et al. (2010) (**Figure 1C**). A

block design was used with four conditions in each type of material: auditory-only

(Aud), visual-only (Vis), audiovisual congruent (avC) and audiovisual incongruent

(avI). The visual and auditory stimuli were presented simultaneously in the

bimodal conditions. The entire experiment contained 4 functional runs, while the

character and *pinyin* stimuli were presented in separate ones. To prevent character

stimuli driving lexical access in *pinyin* processing, *pinyin* experiment was always

conducted first. Each run included 8 experimental blocks (duration 20.8 s) with each

condition replicated twice and interleaved with 9 rest blocks (duration 20.8 s). In

each experimental block, there were 4 miniblocks. Each miniblock consisted of a

1.5-s brain volume acquisition and a 3.7-s silence (i.e., delay, see image acquisition

part) during which 4 trials of stimuli were presented. The order of the stimuli and

conditions were pseudorandomized. Considering the age of participants, an

irrelevant perceptual cover task was used to have children maintaining their

attention on stimuli during scanning. Compared to the active matching task, a

simple task unrelated to the process of interest will not change the automatic

response pattern during the audiovisual integration (Blau et al., 2010; van

Atteveldt et al., 2007). The participant was asked to press the button with the index

finger of right hand accurately and quickly whenever the auditory (440 Hz pure

tone) and/or visual (an unpronounceable symbol) targets appeared (**Figure 1D**).

Only the one with an accurate rate equal to or above 75% in each run was included

in the present study. To ensure that the child understood the task correctly, two

practice sessions were administered, one outside the scanner and the other just

before the first fMRI session inside the scanner.

2.4. Image acquisition

A 3-Tesla Siemens MAGNETOM Trio Tim scanner in Imaging Center for Brain

Research at Beijing Normal University was used to collect all the images with a 12-

channel head coil. In each child, 4 T2* sensitive gradient echo planar imaging

sequences were collected with the following parameters: repetition time (TR) = 5200

ms, echo time = 32 ms, delay in TR = 3700 ms, flip angle = 90 degrees, slice

thickness = 4.5 mm, voxel size = 3.0 × 3.0 × 4.5 mm^3, interscan gap = 0.675 mm,

number of slices = 24, number of volumes = 68, and time of acquisition = 5 minutes

and 54 s. Of note, this sparse sampling method provides a 3.7-s period of silence for

stimuli presentation. Such a design is recommended for fMRI experiments on the

auditory cortex (Shah et al., 2000). In addition, a high-resolution whole-brain T1-

weighted structural image (magnetization-prepared rapid acquisition with gradient

echo, TR = 2530 ms, echo time = 3.39 ms, inversion time = 1100 ms, flip angle = 7

degrees, slice thickness = 1.33 mm, voxel size = $1.33 \times 1.0 \times 1.33$ mm^3, number of

axial slices = 144, and time of acquisition = 8 minutes and 7 s) was collected. Prior

to the formal scan, children were informed about the experimental procedure,

familiarized with the noise of MRI sequences, and trained to hold still. Foam pads

were used to help reduce head motion during scanning. A radiologist blind to the

details of the study reviewed all the images to ascertain any pathological deviations.

2.5. fMRI preprocessing

Preprocessing of images was conducted with SPM12 (Wellcome Department of

Cognitive Neurology, London, UK, http://www.fil.ion.ucl), including: (a) discarding

the first 4 volumes to avoid T1 equilibration effect, (b) head motion correction

(realign), (c) slice-time correction, (d) ART-based outlier detection (a 'bad' brain

volume was defined as intensity > 9 SD or frame-to-frame head motion > 2 mm;

https://www.nitrc.org/projects/artifact_detect), (e) T1 image segmentation, (f)

normalization to the standard template in MNI (Montreal Neurological Institute)

space, and (g) smoothing with an 8-mm full-width at half-maximum Gaussian

kernel. In the subject level, each condition was modeled with a generalized linear

model. Parameters referring to 'bad' volumes and head motion were included in the

model to remove the effect of nuisance covariates. Then, we calculated brain map of

the congruency contrast (avC-avI), which was used in the subsequent analyses.

2.6. Statistical analyses

Descriptive statistics of demographic and behavioral measures were conducted first,

followed by correlations between reading and cognitive-linguistic skills. Then, we

evaluated in-scanner task performance. In this study, we defined an accurate

response as a button press to the target(s) with a reaction time (RT) between 200

and 2000 ms. The average reaction time was calculated for correct responses. ACC

and RT were compared across runs.

In the brain analysis, to identify regions showing congruency effect in

character processing, one sample t-tests were conducted with individual contrast

maps of congruency (i.e., congruent minus incongruent conditions). To avoid results

produced by differences in de-activation, the neurobiological basis of which remains

unknown, we focused on brain areas showing activation in at least one multisensory

condition (p-voxel < 0.001, uncorrected). A threshold of p-voxel < 0.001, FWE

corrected p-cluster < 0.05 was used to identify significant clusters. These areas were

then used as regions-of-interest (ROIs) to explore the relationship between the

multisensory integration of characters with children's oral reading fluency and

reading comprehension proficiency. Nuisance variables of age, sex and performance

IQ were controlled statistically. Bonferroni corrected $p < 0.05$ was used to address

multiple comparison errors. For regions showing a significant relationship with

reading, we additionally investigated whether congruent or incongruent condition

significantly contributed to the correlation, as well as calculated the correlation

between the congruency contrast with PA, RAN and MA, to reveal the cognitive-

linguistic component that was most likely involved in the integration. Finally, we

conducted voxel-wise whole-brain correlation as the complementary approach. All

the analysis steps were replicated for *pinyin* conditions.

Brain results were presented over a surface template (smoothed ICBM152)

with BrainNet Viewer (Xia, Wang, & He, 2013). Significant clusters were reported

in MNI coordinates, and corresponding brain regions were localized and reported

with the AAL atlas implemented in DPABI (http://rfmri.org/dpabi). Behavioral

analysis and ROI analysis were performed with SPSS Statistics version 24 (SPSS

Inc., Chicago, IL, USA).

3. Results

3.1. Behavioral and in-scanner performance

Descriptive statistics of demographics, cognitive-linguistic and reading skills, and

in-scanner task performances are presented in **Table 1**. Positive correlations were

observed between all reading measures (p's < 0.05). Regarding the relation with

cognitive-linguistic skills, *Word List Reading* was associated with *Digit RAN* (r = -

0.425, p = 0.019), but not with *Phoneme Deletion* (r = 0.182, p = 0.336) or

Morphological Production (r = 0.341, p = 0.065). In contrast, *Silent Reading*

Comprehension was correlated with morphological production (r = 0.542, p = 0.002),

but not with *Phoneme Deletion* (r = -0.060, p = 0.754) or *Digit RAN* (r = -0.077, p =

0.686). This pattern indicates that in typically developing Chinese children who

have received formal reading instruction for approximately 4 years, fluent oral

reading necessitates print-to-sound mapping, while proficient reading

comprehension relies more heavily on semantic processing.

Regarding the in-scanner cover task, all children performed well (ACC: *M*

[*SD*] = 96.7% [2.8], RT: *M* [*SD*] = 505 [68] ms). There were no significant differences

in ACC (F = 0.29, p = 0.836) or RT (F = 2.59, p = 0.058) across runs, indicating that

participants focused their attention on incoming stimuli during the entire scan

session.

3.2. Brain results: character

In the voxel-wise whole-brain analysis, a significantly reverse congruency effect

(avC < avI) was found in left IFG (peak MNI -48 16 18) and bilateral STC (peak

MNI, left: -54 32 0, right: 58 -20 0; **Figure 2A and B, Table 2**) at a FWE corrected

threshold of p-cluster < 0.05. We defined these clusters as ROIs to further explore

their relationship with reading and found that the congruency contrast in the left

IFG was positively associated with reading comprehension proficiency (Bonferroni

corrected p < 0.05; **Figure 2C and D**), i.e., the greater the avC > avI, the better the

reading comprehension. The same positive correlation was also found between

congruency contrast and MA (Bonferroni corrected p < 0.05; **Figure 2E**), i.e., the

greater the avC > avI, the better the MA. Interestingly, while the congruency-

comprehension correlation was driven by individual differences in activation in the

incongruent condition (avC: $r = 0.164$, $p = 0.413$; avI: $r = -0.547$, $p = 0.003$), no

significant correlation was found between MA and activation in either avC ($r =$

0.326, $p = 0.097$) or avI condition ($r = -0.210$, $p = 0.292$). Finally, complementary

voxel-wise whole-brain correlation analysis revealed that reading comprehension

positively correlated with congruency contrast in the left IFG and inferior parietal

lobule (IPL) at a FWE corrected threshold of p-cluster < 0.05 (**Figure 3**, **Table 2**).

In line with the above analyses, such relationships were driven by the negative

correlation between reading and activation in the incongruent condition. Moreover,

congruency-MA correlations in both regions were significant (left IFG: $r = 0.593$, $p =$

0.001; left IPL: $r = 0.573$, $p = 0.002$). No additional regions were identified in whole-

brain regression for oral reading fluency.

3.3. Brain results: *pinyin*

Unexpectedly, no regions showed a significant congruency effect when the whole

brain analysis was performed across the entire group. However, voxel-wise

correlation analysis revealed that oral reading fluency positively correlated with the

congruency contrasts (avC-avI) in left STG (peak MNI -56 -36 14) (FWE corrected p-

cluster < 0.05; **Figure 4A and B**, **Table 2**). ROI analysis revealed that this

relationship was driven more by the avC condition ($r = 0.399$, $p = 0.039$), but not the

avI condition ($r = -0.277$, $p = 0.162$). No significant correlations were found between

congruency contrasts and cognitive-linguistic skills. Finally, complementary

analyses was performed by dividing participants into three groups based on their

oral word fluency (good [n = 8]: standardized score > 1, average [n = 10]: 0 <

standardized score < 1, and poor [n = 12]: standardized score < 0). Positive

congruency (avC > avI) were found in all eight children in the good group ($t = 5.753$,

$p < 0.001$; **Figure 4C**, **Table S1**), while negative patterns (avC < avI) were

observed in 10/12 children with poor reading fluency ($t = -3.635$, $p = 0.004$). No

significant effect existed in the average group ($t = 0.345$, $p = 0.738$).

4. Discussion

Establishing efficient print-to-sound mapping is essential for successful reading

acquisition (Blomert, 2011; Richlan, 2019). In alphabetic languages, the congruency

effect during audiovisual integration has repeatedly been observed in STC, while

the direction of effect is influenced by orthographic depth. Against the background

that the vast majority of research is restricted to alphabetic scripts, the present

study investigated the neurofunctional basis underlying audiovisual integration of

characters (deep orthography) and *pinyin* (transparent alphabetic coding system) in

typically developing Chinese children for the first time. The results in both scripts

are in line with predictions according to the orthographic depth hypothesis,

supporting a universal principle underlying the neural mechanism of audiovisual

integration of print and speech sound across languages. Moreover, we ascertained

that left IFG links multisensory integration of characters and reading

comprehension proficiency through automatic semantic access, while left STG links

multisensory integration of *pinyin* and oral reading fluency possibly through a

shared common mechanism of grapho-phonological mapping.

4.1. Left IFG and bilateral STC are involved in implicit integration of

multisensory information about Chinese characters

To date, various neuroimaging tools have been used to explore neural mechanisms

underlying audiovisual integration of print and speech sound. Techniques such as

electroencephalogram (EEG) and MEG have been used to focus on the temporal

dimension (Froyen, van Atteveldt, & Blomert, 2010; Froyen, Van Atteveldt, Bonte,

& Blomert, 2008; Froyen, Bonte, van Atteveldt, & Blomert, 2009; Raij, Uutela, &

Hari, 2000; Xu et al., 2019). These studies demonstrated that such integration is

preconscious in expert reading, which is reflected in a component of mismatch

negativity/response. On the other hand, fMRI research provides greater information

about spatial localization. These studies repeatedly observed difference in brain

responses to audiovisual congruent versus incongruent stimulus pairs in STC that

mainly consists of superior temporal gyrus and sulcus (van Atteveldt et al., 2004;

van Atteveldt, Formisano, Blomert, & Goebel, 2007). Importantly, such a

congruency effect has been confirmed in passive and irrelevant paradigms,

pinpointing the automatic nature of this effect (Blau et al., 2010; van Atteveldt et

al., 2007). Furthermore, this effect is modulated by linguistic features (Holloway et

al., 2015), and is associated with individual characteristics such as reading (Blau et

al., 2010; Blau et al., 2009; Karipidis et al., 2018).

Consistency of mapping between grapheme and phoneme is an important

feature in alphabetic languages. Differences in the directionality of the effect and

brain regions that show this effect have been found between languages with varying

orthographic depth (Blomert & Froyen, 2010; Holloway et al., 2015). In particular,

greater activation in congruent condition compared with incongruent condition is

more consistently reported in transparent scripts, while the opposite pattern is

observed in opaque scripts. Chinese is a logographic language with an extremely

deep orthography. Therefore, a reverse congruency effect can be expected based on

the orthographic depth hypothesis. In support of this idea, the present study

revealed three regions that showed a significantly reverse congruency effect,

including STC in both hemispheres, which have been repeatedly reported in

previous audiovisual integration studies. Moreover, deep orthography is not the

only linguistic feature that matters. Chinese is a morpheme-based language in

which semantic information is deeply involved, even at the basic character level.

That is, in addition to grapho-phonological mapping, semantic access is an equally

important aspect of reading (Bi, Han, Weekes, & Shu, 2007; Dang, Zhang, Wang, &

Yang, 2018; Liu et al., 2017; Ruan et al., 2018; Yang et al., 2012). In the current

study, the left IFG displayed a significantly incongruency effect, suggesting that

this region is associated with multisensory integration. More importantly, such an

effect is strongly associated with reading comprehension proficiency as well as

morphological processing. In previous research, frontal region has also been

reported, but along with a broad bilateral fronto-parietal network. Its functional

role was proposed as task related top-down modulation (van Atteveldt et al., 2007)

or implicit domain-general conflict detection and resolution (Holloway et al., 2015).

Recently, a reverse congruency effect was observed in frontal cortex in experienced

readers in Chinese, a non-alphabetic language, with MEG (Xu et al., 2019). Here,

we not only replicate such finding in a developmental population in Chinese with

fMRI, but further demonstrates the role of left IFG in the implicit audiovisual

integration of character is domain-specific and the information automatically

processed in this region corresponds to semantics.

In addition to left IFG, one more region displayed significant correlation

between congruency contrast and reading skills. In particular, congruency contrasts

in the left IPL were positively associated with reading comprehension proficiency

and morphological processing but not with oral reading fluency, phonological

processing or naming speed. This finding is in line with the role of this region in

semantic processing (Binder & Desai, 2011; Wang, Zhao, Zevin, & Yang, 2016). The

left IPL has not been identified to be associated with audiovisual integration

previously, however a recent training study revealed that this region was involved

in the acquisition of novel grapho-phonological mapping (Xu, Kolozsvari,

Oostenveld, & Hamalainen, 2020). Based on our and previous findings, it is

reasonable to suggest that the functional role of a region in audiovisual integration

is influenced by the linguistic features in the given language.

Taken together, these findings indicate that typically developing children in

upper elementary grades integrate high frequency characters and corresponding

speech sounds implicitly. Moreover, the linguistic features of Chinese writing

system drive the precise neural manifestation of the congruency effect, in line with

previous literature, supporting the notion that while the general neural circuit for

reading is universal, linguistic features in a given language introduce different

demands for specific cognitive components (Bolger, Perfetti, & Schneider, 2005;

Rueckl et al., 2015).

4.2. Left STG links implicit print-to-sound integration of *pinyin* with fluent

oral reading

Well-established print-to-sound mapping is essential for learning to read (Blomert,

2011; Richlan, 2019). While letter knowledge is one of the most important

prerequisites in alphabetic languages, *pinyin* – a transparent phonological coding

system plays a scaffolding role at the initial stage in Chinese reading acquisition.

Therefore, the present study, on the one hand utilizes a unique opportunity to

examine the orthographic depth hypothesis with both deep and shallow

orthographies in one language. On the other hand, it provides information that

helps illustrate the role of *pinyin* in developing fluent reading ability.

First, as proposed by the orthographic depth hypothesis, scripts with

transparent GPC should show a stronger brain activation in congruent condition

compared to incongruent condition in STC. At first glance, no significant

congruency effect was observed across the entire group. However, by grouping the

participants based on oral reading abilities, a clear pattern appeared in the left

STG. In particular, all the children in the better fluency group showed higher

activation in congruent versus incongruent conditions, similar to that observed in

shallow orthographies (van Atteveldt et al., 2004). Moreover, average readers

showed less congruency, while almost all below-average readers displayed a

negative pattern. This is in line with studies in individuals with RD in these

languages, where a less congruency effect was found (Blau et al., 2010; Blau et al.,

2009). In terms of the brain region, the left STG is the central area that represents

phonological information (Boets et al., 2013; Glezer et al., 2016). Its activation and

connection patterns are known to change when learning visual-sound mappings

(Dehaene et al., 2010; Li, Xu, Luo, Zeng, & Han, 2020; Thiebaut de Schotten, Cohen,

Amemiya, Braga, & Dehaene, 2014), which can further scaffold later reading

development (Wang, Joanisse, & Booth, 2019). It should be noted that the current

conclusion can only be made for typical readers since these below-average children

performed less well in the oral word reading task but were not classified as having

RD.

In addition to supporting the orthographic depth hypothesis, the significant

correlation between congruency contrasts in the left STG to audiovisual stimulus

pairs of *pinyin* with oral reading fluency may imply that learning *pinyin* help

typical children develop fluent oral reading ability (though causal claims cannot be

made from this study). In comparison to other cognitive-linguistic skills, such as PA

and RAN, less attention has been paid to the role of *pinyin* processing in Chinese

reading development, probably because it is a scaffold in reading acquisition but not

considered reading itself. However, although the number of studies is limited, the

pinyin-reading relationship has been demonstrated at the behavioral level (Lin et

al., 2010; Pan et al., 2011; Siok & Fletcher, 2001). In a longitudinal study, Pan et al.

demonstrated that invented *pinyin* spelling at age 6 independently predicted later

reading performance at ages 8 and 10 (Pan et al., 2011). Several explanations are

proposed based on behavioral evidences. One possibility is that *pinyin* can help to

establish better phonological representations (McBride, Wang, & Cheang, 2018).

Alternatively, training on *pinyin* can uniquely improve character-to-syllable

mapping, revealed in research focusing on second language learner of Chinese

(Guan et al., 2011). In this study, the congruency contrasts in the left STG to *pinyin*

were positively correlated with oral reading fluency, while a similar region also

showed multisensory integration of audiovisual pairs of characters. While

conclusion cannot be drawn from lack of evidence, our results that brain measures

were not correlated with auditory phonological processing (i.e., PA), may support

the second hypothesis that learning *pinyin* shapes the neural circuit that will later

be recruited in learning characters.

Overall, this study further supports the orthographic depth hypothesis by

extending it to *pinyin*, an alphabetic coding system used for representing

pronunciations of characters in Chinese. It also suggests that even though Chinese

is a logographic writing system, print-speech sound integration still plays a critical

role in reading development, in which the underlying grapho-phonological route

may be shaped by learning through *pinyin* prior to formal character reading

instruction.

4.3. Caveats and future directions

While the current findings are appealing, several caveats should be noted. First, in

this study, we investigated the neurofunctional basis underlying audiovisual

integration of *pinyin* for the first time. However, we did not observe a significant

congruency effect at the group level, as proposed by the orthographic depth

hypothesis. This may be because while all the participants had typical reading

skills, there were still individual differences in *pinyin* reading that we did not

assess directly. Nevertheless, a congruency effect was found in the fluent readers,

while a reverse pattern was observed in slow readers. Moreover, previous

longitudinal studies have revealed a significant role of *pinyin* processing in reading

acquisition (Lin et al., 2010; Pan et al., 2011; Siok & Fletcher, 2001). Taken

together, the results of this study propose the left STG as the possible neural basis

underlying such linkage, which should be examined in future studies. Second, since

the participants recruited in this study were in grades 3-5, *pinyin* may play a

different role in these individuals than it does in earlier stages of reading

development. In this case, it is important to conduct a study following preliterate

children until they achieve fluent reading in future. Finally, since we have

uncovered brain responses to audiovisual information and their relationship with

reading performance in typically developing children, such findings can be used as

the baseline for investigating whether neural processes are altered in children with

RD.

5. Conclusion

In the current study, we demonstrate a reverse congruency effect for the

audiovisual integration of characters and a congruency effect for *pinyin* in typically

developing Chinese children with fluent reading ability. This striking finding

supports and extends the orthographic depth hypothesis. Furthermore, we found

that the left IFG is strongly involved in the multisensory integration of characters

and is correlated with reading comprehension and morphological awareness,

suggesting that semantic representations were more automatically accessed in

readers with higher comprehension proficiency. Finally, this study revealed that

integrating multisensory information of *pinyin* in the left STG is correlated with

oral reading fluency, implying that the grapho-phonological mapping introduced by

pinyin learning may help children develop better reading skills.

Acknowledgements

The authors would like to thank all the children participating in this study, their

families, as well as all the examiners. This work was supported by the Key Program

of the National Social Science Foundation of China (14ZDB157), National Key Basic

Research Program of China (grant number 2014CB846103), National Natural

Science Foundation of China grants (grant number 31271082, 31671126,

31611130107, 61374165, 81801782), Beijing Municipal Science & Technology

Commission (grant number Z151100003915122), Fundamental Research Fund for

the Central Universities, China Postdoctoral Science Foundation (grant number

2019T120062, 2018M641235), and Social Science Fund of Beijing (grant number

17YYA004). No conflict declared.

References

Bakos, S., Landerl, K., Bartling, J., Schulte-Körne, G., & Moll, K. (2017). Deficits in

Letter-Speech Sound Associations but Intact Visual Conflict Processing in

Dyslexia: Results from a Novel ERP-Paradigm. *Front Hum Neurosci, 11*(116).

Bi, Y., Han, Z., Weekes, B., & Shu, H. (2007). The interaction between semantic and

the nonsemantic systems in reading: Evidence from Chinese.

Neuropsychologia, 45(12), 2660-2673.

Binder, J. R., & Desai, R. H. (2011). The neurobiology of semantic memory. *Trends

Cogn Sci, 15*(11), 527-536.

Blau, V., Reithler, J., van Atteveldt, N., Seitz, J., Gerretsen, P., Goebel, R., &

Blomert, L. (2010). Deviant processing of letters and speech sounds as

proximate cause of reading failure: a functional magnetic resonance imaging

study of dyslexic children. *Brain, 133*(Pt 3), 868-879.

Blau, V., van Atteveldt, N., Ekkebus, M., Goebel, R., & Blomert, L. (2009). Reduced

neural integration of letters and speech sounds links phonological and

reading deficits in adult dyslexia. *Current Biology, 19*(6), 503-508.

Blomert, L. (2011). The neural signature of orthographic-phonological binding in

successful and failing reading development. *Neuroimage, 57*(3), 695-703.

Blomert, L., & Froyen, D. (2010). Multi-sensory learning and learning to read. *Int J*

Psychophysiol, 77(3), 195-204.

Blomert, L., & Willems, G. (2010). Is there a causal link from a phonological

awareness deficit to reading failure in children at familial risk for dyslexia?

Dyslexia, 16(4), 300-317.

Boets, B., Op de Beeck, H. P., Vandermosten, M., Scott, S. K., Gillebert, C. R.,

Mantini, D., . . . Ghesquiere, P. (2013). Intact but less accessible phonetic

representations in adults with dyslexia. *Science, 342*(6163), 1251-1254.

Bolger, D. J., Perfetti, C. A., & Schneider, W. (2005). Cross-cultural effect on the

brain revisited: universal structures plus writing system variation. *Human*

Brain Mapping, 25(1), 92-104.

Chan, C. K. K., & Siegel, L. S. (2001). Phonological Processing in Reading Chinese

among Normally Achieving and Poor Readers. *J Exp Child Psychol, 80*(1), 23-

43.

Cui, Z., Xia, Z., Su, M., Shu, H., & Gong, G. (2016). Disrupted white matter

connectivity underlying developmental dyslexia: A machine learning

approach. *Human Brain Mapping, 37*(4), 1443-1458.

Dang, M., Zhang, R., Wang, X., & Yang, J. (2018). The Interaction Between

Phonological and Semantic Processing in Reading Chinese Characters. *Front*

Psychol, 9, 2748.

Dehaene, S., Pegado, F., Braga, L. W., Ventura, P., Nunes, G., Jobert, A., . . . Cohen,

L. (2010). How Learning to Read Changes the Cortical Networks for Vision

and Language. *Science, 330*(6009), 1359-1364.

Froyen, D., van Atteveldt, N., & Blomert, L. (2010). Exploring the Role of Low Level

Visual Processing in Letter-Speech Sound Integration: A Visual MMN Study.

Front Integr Neurosci, 4, 9.

Froyen, D., Van Atteveldt, N., Bonte, M., & Blomert, L. (2008). Cross-modal

enhancement of the MMN to speech-sounds indicates early and automatic

integration of letters and speech-sounds. *Neurosci Lett, 430*(1), 23-28.

Froyen, D. J. W., Bonte, M. L., van Atteveldt, N., & Blomert, L. (2009). The Long

Road to Automation: Neurocognitive Development of Letter-Speech Sound

Processing. *J Cogn Neurosci, 21*(3), 567-580.

Glezer, L. S., Eden, G., Jiang, X., Luetje, M., Napoliello, E., Kim, J., & Riesenhuber,

M. (2016). Uncovering phonological and orthographic selectivity across the

reading network using fMRI-RA. *Neuroimage, 138*, 248-256.

Guan, C. Q., Liu, Y., Chan, D. H. L., Ye, F., & Perfetti, C. A. (2011). Writing

strengthens orthography and alphabetic-coding strengthens phonology in

learning to read Chinese. *Journal of Educational Psychology, 103*(3), 509-522.

Holloway, I. D., van Atteveldt, N., Blomert, L., & Ansari, D. (2015). Orthographic

Dependency in the Neural Correlates of Reading: Evidence from Audiovisual

Integration in English Readers. *Cereb Cortex, 25*(6), 1544-1553.

Karipidis, I. I., Pleisch, G., Brandeis, D., Roth, A., Röthlisberger, M., Schneebeli, M.,

644   . . . Brem, S. (2018). Simulating reading acquisition: The link between

645   reading outcome and multimodal brain signatures of letter–speech sound

646   learning in prereaders. *Scientific Reports, 8*(1), 7121.

647   Kast, M., Bezzola, L., Jancke, L., & Meyer, M. (2011). Multi- and unisensory

648   decoding of words and nonwords result in differential brain responses in

649   dyslexic and nondyslexic adults. *Brain and Language, 119*(3), 136-148.

650   Lei, L., Pan, J., Liu, H., McBride-Chang, C., Li, H., Zhang, Y., . . . Zhang, Z. (2011).

651   Developmental trajectories of reading development and impairment from

652   ages 3 to 8 years in Chinese children. *Journal of Child Psychology and*

653   *Psychiatry, 52*(2), 212-220.

654   Li, H., Shu, H., McBride-Chang, C., Liu, H., & Peng, H. (2012). Chinese children's

655   character recognition: Visuo-orthographic, phonological processing and

656   morphological skills. *Journal of Research in Reading, 35*(3), 287-307.

657   Li, M., Xu, Y., Luo, X., Zeng, J., & Han, Z. (2020). Linguistic experience acquisition

658   for novel stimuli selectively activates the neural network of the visual word

659   form area. *Neuroimage*, 116838.

660   Lin, D., McBride-Chang, C., Shu, H., Zhang, Y., Li, H., Zhang, J., . . . Levin, I.

661   (2010). Small Wins Big:Analytic Pinyin Skills Promote Chinese Word

662   Reading. *Psychol Sci, 21*(8), 1117-1122.

Liu, Y., Georgiou, G. K., Zhang, Y., Li, H., Liu, H., Song, S., . . . Shu, H. (2017).

Contribution of cognitive and linguistic skills to word-reading accuracy and

fluency in Chinese. *International Journal of Educational Research, 82*, 75-90.

McBride, C., Wang, Y., & Cheang, L. M.-L. (2018). Dyslexia in Chinese. *Current*

Developmental Disorders Reports, 5(4), 217-225.

Mcnorgan, C., Randazzo-Wagner, M., & Booth, J. (2013). Cross-modal integration in

the brain is related to phonological awareness only in typical readers, not in

those with reading difficulty. *Front Hum Neurosci, 7*(388).

Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh

inventory. *Neuropsychologia, 9*(1), 97-113.

Pan, J., McBride-Chang, C., Shu, H., Liu, H., Zhang, Y., & Li, H. (2011). What is in

the naming? A 5-year longitudinal study of early rapid naming and

phonological sensitivity in relation to subsequent reading skills in both native

Chinese and English as a second language. *Journal of Educational*

Psychology, 103(4), 897-908.

Plewko, J., Chyl, K., Bola, Ł., Łuniewska, M., Dębska, A., Banaszkiewicz, A., . . .

Jednoróg, K. (2018). Letter and Speech Sound Association in Emerging

Readers With Familial Risk of Dyslexia. *Front Hum Neurosci, 12*(393).

Raij, T., Uutela, K., & Hari, R. (2000). Audiovisual integration of letters in the

human brain. *Neuron, 28*(2), 617-625.

Richlan, F. (2019). The Functional Neuroanatomy of Letter-Speech Sound

Integration and Its Relation to Brain Abnormalities in Developmental

Dyslexia. *Front Hum Neurosci, 13*(21).

Ruan, Y., Georgiou, G. K., Song, S., Li, Y., & Shu, H. (2018). Does writing system

influence the associations between phonological awareness, morphological

awareness, and reading? A meta-analysis. *Journal of Educational*

Psychology, 110(2), 180.

Rueckl, J. G., Paz-Alonso, P. M., Molfese, P. J., Kuo, W. J., Bick, A., Frost, S. J., . . .

Frost, R. (2015). Universal brain signature of proficient reading: Evidence

from four contrasting languages. *Proc Natl Acad Sci U S A, 112*(50), 15510-

15515.

Shah, N. J., Steinhoff, S., Mirzazade, S., Zafiris, O., Grosse-Ruyken, M. L., Jäncke,

L., & Zilles, K. (2000). The Effect of Sequence Repeat Time on Auditory

Cortex Stimulation During Phonetic Discrimination. *Neuroimage, 12*(1), 100-

108.

Shu, H., Chen, X., Anderson, R. C., Wu, N., & Xuan, Y. (2003). Properties of school

Chinese: Implications for learning to read. *Child development, 74*(1), 27-47.

Shu, H., McBride-Chang, C., Wu, S., & Liu, H. Y. (2006). Understanding Chinese

developmental dyslexia: Morphological awareness as a core cognitive

construct. *Journal of Educational Psychology, 98*(1), 122-133.

Siok, W. T., & Fletcher, P. (2001). The role of phonological awareness and visual-

orthographic skills in Chinese reading acquisition. *Dev Psychol, 37*(6), 886-

899.

Thiebaut de Schotten, M., Cohen, L., Amemiya, E., Braga, L. W., & Dehaene, S.

(2014). Learning to read improves the structure of the arcuate fasciculus.

Cereb Cortex, 24(4), 989-995.

van Atteveldt, N., Formisano, E., Goebel, R., & Blomert, L. (2004). Integration of

letters and speech sounds in the human brain. *Neuron, 43*(2), 271-282.

van Atteveldt, N. M., Formisano, E., Blomert, L., & Goebel, R. (2007). The effect of

temporal asynchrony on the multisensory integration of letters and speech

sounds. *Cereb Cortex, 17*(4), 962-974.

van Atteveldt, N. M., Formisano, E., Goebel, R., & Blomert, L. (2007). Top–down

task effects overrule automatic multisensory responses to letter–sound pairs

in auditory association cortex. *Neuroimage, 36*(4), 1345-1360.

Wang, J., Joanisse, M. F., & Booth, J. R. (2019). Neural representations of

phonology in temporal cortex scaffold longitudinal reading gains in 5- to 7-

year-old children. *Neuroimage, 207*, 116359.

Wang, X., Zhao, R., Zevin, J. D., & Yang, J. (2016). The Neural Correlates of the

Interaction between Semantic and Phonological Processing for Chinese

Character Reading. *Front Psychol, 7*.

Wechsler, D. (1974). *WISC-R, Wechsler intelligence scale for children, revised*:

Psychological Corporation.

Wu, C. Y., Ho, M. H., & Chen, S. H. (2012). A meta-analysis of fMRI studies on

Chinese orthographic, phonological, and semantic processing. *Neuroimage,*

63(1), 381-391.

Xia, M., Wang, J., & He, Y. (2013). BrainNet Viewer: a network visualization tool

for human brain connectomics. *PLoS One, 8*(7), e68910.

Xia, Z., Hoeft, F., Zhang, L., & Shu, H. (2016). Neuroanatomical anomalies of

dyslexia: Disambiguating the effects of disorder, performance, and

maturation. *Neuropsychologia, 81*, 68-78.

Xia, Z., Zhang, L., Hoeft, F., Gu, B., Gong, G., & Shu, H. (2018). Neural correlates of

oral word reading, silent reading comprehension, and cognitive

subcomponents. *International Journal of Behavioral Development, 42*(3), 342-

356.

Xu, W., Kolozsvari, O. B., Monto, S. P., & Hämäläinen, J. A. (2018). Brain

Responses to Letters and Speech Sounds and Their Correlations With

Cognitive Skills Related to Reading in Children. *Front Hum Neurosci,*

12(304).

Xu, W., Kolozsvari, O. B., Oostenveld, R., & Hamalainen, J. A. (2020). Rapid

changes in brain activity during learning of grapheme-phoneme associations

in adults. *bioRxiv*, 2020.2003.2012.988402.

Xu, W., Kolozsvári, O. B., Oostenveld, R., Leppänen, P. H. T., & Hämäläinen, J. A.

(2019). Audiovisual Processing of Chinese Characters Elicits Suppression and

Congruency Effects in MEG. *Front Hum Neurosci, 13*(18).

Xue, J., Shu, H., Li, H., Li, W., & Tian, X. J. J. o. P. R. (2013). The Stability of

Literacy-Related Cognitive Contributions to Chinese Character Naming and

Reading Fluency. *42*(5), 433-450.

Yan, M., Miller, K. F., Li, H., & Shu, H. (2008). What is the place for Pinyin in

beginning Chinese reading? Evidence from eye movements. In *Cognitive and*

cultural influences on eye movements. (pp. 343-360). Tianjin, China: Tianjin

People's Publishing House.

Yang, J., Shu, H. U. A., McCandliss, B. D., & Zevin, J. D. (2012). Orthographic

influences on division of labor in learning to read Chinese and English:

Insights from computational modeling. *Bilingualism: Language and*

Cognition, 16(2), 354-366.

Zhang, Y., Zhang, L., Shu, H., Xi, J., Wu, H., Zhang, Y., & Li, P. (2012). Universality

of categorical perception deficit in developmental dyslexia: an investigation of

Mandarin Chinese tones. *Journal of Child Psychology and Psychiatry, 53*(8),

874-882.

Zhao, J., Wang, X., Frost, S. J., Sun, W., Fang, S.-Y., Mencl, W. E., . . . Rueckl, J. G.

(2014). Neural division of labor in reading is constrained by culture:

A training study of reading Chinese characters. *Cortex, 53*, 90-106.

Table 1 Demographics and behavior measures of the participants

Measures	Mean	SD	Range
Age (month)	127	7	111 ~ 140
Sex (female/male)	20/10		
Verbal IQ (standard score)	110	13	85 ~ 145
Performance IQ (standard score)	116	13	82 ~ 137
Full scale IQ (standard score)	114	12	93 ~ 143
Character Recognition (item)	121	11	100 ~ 146
Standard score	0.62	0.74	-0.86 ~ 2.20
Word List Reading (word / minute)	95	17	71 ~ 140
Standard score	0.49	1.14	-0.91 ~ 3.53
Reading Fluency (character / minute)	359	107	176 ~ 565
Standard score	0.93	1.08	-0.57 ~ 3.08
Phoneme Deletion (item)	21	4	7 ~ 24
Rapid Naming (second)	17	2	11 ~ 23
Morphological Production (item)	24	4	13 ~ 30

Table 2 Significant clusters identified in brain analyses

Material and contrast	Label	Brain area (AAL)	$P_{\text{FWE-corrected}}$	Size	Peak T	X	Y	Z
Character								
Congruency (avC vs avI)	LMTG	Left middle temporal gyrus, superior temporal gyrus	0.009	450	-5.04	-54	-32	0
	LIFG	Left inferior frontal gyrus, triangular part and pars orbitalis, precentral gyrus	0.022	303	-4.09	-48	16	18
	RSTG	Right superior temporal gyrus, middle temporal gyrus	0.043	218	-5.98	58	-20	0
Congruency (avC-avI) correlated with reading comprehension	LPrCG	Left, precentral gyrus, inferior frontal gyrus, triangular part and pars orbitalis, middle frontal gyrus	0.008	471	5.44	-32	2	36
	LIPL	Left inferior parietal gyrus, superior parietal gyrus, middle occipital gyrus, angular gyrus	0.030	307	6.53	-32	-68	46
Pinyin								
Congruency (avC-avI) correlated with oral reading	LSTG	Left superior temporal gyrus, middle temporal gyrus	0.008	397	4.88	-56	-36	14

Note: The significant clusters were identified with an FWE corrected threshold of p-cluster < 0.05 (p-voxel < 0.001). Brain area labeling is

based on the AAL atlas. Cluster size refers to the number of voxels. avC = audiovisual congruent, avI = audiovisual incongruent.

Figure Legends

Figure 1 Materials and experimental design. Example stimuli in *pinyin* (A) and

character (B) conditions. (C) Schematic illustration of the fMRI procedure. (D)

Auditory and visual targets in the cover task.

Figure 2 Results of brain analysis comparing between character conditions. (A)

Areas showing a significant congruency effect (avC vs. avI). Threshold: p-voxel <

0.001, p-cluster < 0.05, Family-Wise Error (FWE) correction. (B) Bar plots showing

activation for each condition. (C) Regions showing significant correlations with

reading and related cognitive skills. (D) Scatter plot presenting a positive

correlation between the congruency effect in left IFG and reading comprehension

efficiency (Bonferroni corrected p < 0.05). (E) Scatter plot representing the positive

correlation between the congruency effect in the left IFG and morphological

awareness (Bonferroni corrected p < 0.05). Aud = auditory-only, Vis = visual-only,

avC = audiovisual congruent, avI = audiovisual incongruent, L = left, IFG = inferior

frontal gyrus, STC = superior temporal cortex, STG = superior temporal gyrus.

Figure 3 (A) Clusters displaying a significant correlation between the congruency

contrast and reading comprehension proficiency across the brain. Threshold: p-voxel

< 0.001, p-cluster < 0.05, Family-Wise Error (FWE) correction. Scatter plots

displaying the correlations in left IFG (B) and the left IPL (C). IFG = inferior frontal

gyrus, IPL = inferior parietal lobular.

Figure 4 (A) Clusters displaying a significant correlation between the congruency

effect in *pinyin* and oral reading efficiency across the brain. Threshold: p-voxel <

0.001, p-cluster < 0.05, Family-Wise Error (FWE) correction. (B) Scatter plot

displaying the correlation. (C) Bar plots showing the activation in each condition in

individuals with different levels of oral reading fluency.

Figure S1 Brain activations for each condition in (A) *pinyin* and (B) character

tasks. A threshold of uncorrected p-voxel < 0.001 is used for presentation purposes.

Figure 1 Materials and experimental design. Example stimuli in *pinyin* (A) and

character (B) conditions. (C) Schematic illustration of the fMRI procedure. (D)

Auditory and visual targets in the cover task

Figure 2 Results of brain analysis comparing between character conditions. (A) Areas showing a significant congruency effect (avC vs. avI). Threshold: *p*-voxel < 0.001, *p*-cluster < 0.05, Family-Wise Error (FWE) correction. (B) Bar plots showing the activation in each condition. (C) Regions showing significant correlations with reading and related cognitive skills. (D) Scatter plot presenting a positive correlation between the congruency effect in left IFG and reading comprehension efficiency (Bonferroni corrected p < 0.05). (E) Scatter plot representing the positive correlation between the congruency effect in the left IFG and morphological awareness (Bonferroni corrected p < 0.05). Aud = auditory-only, Vis = visual-only, avC = audiovisual congruent, avI = audiovisual incongruent, L = left, IFG = inferior frontal gyrus, STC = superior temporal cortex, STG = superior temporal gyrus.

Figure 3 (A) Clusters displaying a significant correlation between the congruency contrast and reading comprehension proficiency across the brain. Threshold: *p*-voxel < 0.001, *p*-cluster < 0.05, Family-Wise Error (FWE) correction. Scatter plots displaying the correlations in the left IFG (B) and left IPL (C). IFG = inferior frontal gyrus, IPL = inferior parietal lobular.

818   **Figure 4** (A) Clusters displaying a significant correlation between the congruency

819   effect in *pinyin* and oral reading efficiency across the brain. Threshold: *p*-voxel <

820   0.001, *p*-cluster < 0.05, Family-Wise Error (FWE) correction. (B) Scatter plot

821   displaying the correlation. (C) Bar plots showing the activation in each condition in

822   individuals with different levels of oral reading fluency.