Bacterial influence on the maintenance of symbiotic yeast through *Drosophila* metamorphosis

Running title: Bacterial influence on yeast maintenance

Robin Guilhot[1*], Antoine Rombaut[1], Anne Xuéreb[1], Kate Howell[2], Simon Fellous[1]

[1] CBGP, INRAE, CIRAD, IRD, Montpellier SupAgro, Univ Montpellier, Montpellier, France

[2] Faculty of Veterinary and Agricultural Sciences, University of Melbourne, Parkville, Vic 3010, Australia

*corresponding author's email address: guilhoro@gmail.com

Key words: maintenance through metamorphosis, symbiosis, microbial interactions, *Drosophila*, *Saccharomyces cerevisiae*, extracellular bacteria

Summary statement

Bacterial symbionts of *Drosophila* influence yeast maintenance through fly metamorphosis, a novel observation that may have consequences for the evolution of insect-yeast-bacteria interactions.

Abstract

Interactions between microbial symbionts of metazoan hosts are emerging as key features of symbiotic systems. Little is known about the role of such interactions on the maintenance of symbiosis through host's life cycle. We studied the influence of symbiotic bacteria on the maintenance of symbiotic yeast through metamorphosis of the fly *Drosophila melanogaster*. To this end we mimicked the development of larvae in natural fruit. In absence of bacteria yeast was never found in young adults. However, yeast could maintain through metamorphosis when larvae were inoculated with symbiotic bacteria isolated from *D. melanogaster* faeces. Furthermore, an Enterobacteriaceae favoured yeast transstadial maintenance. Because yeast is a critical symbiont of *D. melanogaster* flies, bacterial influence on host-yeast association may have consequences for the evolution of insect-yeast-bacteria tripartite symbiosis and their cooperation.

Introduction

Metazoans often form associations with non-obligatory symbiotic microorganisms. Microbial symbiont can influence host phenotype, and hosts determine symbiont multiplication and dispersal (Ferrari and Vavre, 2011). The importance of interactions between microorganisms is relatively better understood in the context of parasitism than in the context of beneficial symbionts (Alizon et al., 2013; Tollenaere et al., 2016; Zélé et al., 2018). Beneficial microbial symbionts can nonetheless interact in a wide variety of manners (Comolli, 2014; Seth and Taga, 2014; Hassani et al., 2018) and affect both the dynamics of each microorganism and the phenotype of their host (Wargo and Hogan, 2006; Newell and Douglas, 2014; Callens et al., 2018; Gould et al., 2018; Sommer and Newell, 2019).

Fungi-bacteria interactions are well described microbial interactions due to their importance in human health, industry and domestic life (Kobayashi and Crouch, 2009; Jouhten et al., 2016; Carbonetto et al., 2018). In the wild, yeasts and other fungi associate with extracellular bacteria in a wide variety of habitats, including decaying plant materials where they interact with the larvae and adults of saprophagous insects such as *Drosophila* flies. Symbioses between *Drosophila* and either yeast and bacteria have however been largely studied separately. Each type of microorganism affects *Drosophila* physiology, nutrition, reproduction and behavior (Ryu et al., 2008; Anagnostou et al., 2010; Shin et al., 2011; Storelli et al., 2011; Becher et al., 2012; Broderick and Lemaitre, 2012; Wong et al., 2014) and may maintain through - a variable part of - *Drosophila* life cycle (Bakula, 1969; Starmer et al., 1988; Hoang et al., 2015; Pais et al., 2018). In natural fruit, *D. melanogaster* larval development can be impossible, or largely compromised, in absence of yeast (Becher et al., 2012), suggesting symbiosis with yeast may not be dispensable for larvae. A handful of studies considering both fungi and bacteria showed that direct interaction between yeast and bacteria can modulate fly behavior (Fischer et al., 2017) and that bacteria can affect fly attraction to yeast (Leitão-Gonçalves et al., 2017). There is to date little data on how interactions between microbial symbionts affect their transmission from the host or maintenance among host life stages, which would have wide consequences for the ecological dynamics of the microorganisms, and consequently, the evolution of their symbiosis with flies.

It is established that *Drosophila* flies contribute to bacteria and yeast dynamics through effects on local multiplication and dispersal (Gilbert, 1980; Ganter, 1988; Starmer et al.,

1988; Chandler et al., 2012; Stamps et al., 2012; Buser et al., 2014). *Saccharomyces* yeast, for example, attracts adult flies with volatiles, these adults then acquire the microorganism and later inoculate it in fruit where larvae develop. An alternative mean of insect vectoring for symbiotic bacteria and yeast of larvae would rely on the transstadial maintenance, from the larval to the adult stage. It would enable the colonization of freshly emerged adults by larval symbionts and, provided they maintain and are shed later in life, their possible dispersal to new patches of resources. The maintenance of yeast or bacteria throughout the *Drosophila* life cycle have been investigated (Rohlfs and Hoffmeister, 2005; Pais et al., 2018), including maintenance from larvae to adults through metamorphosis (i.e. transstadial maintenance or transstadial transmission) (Bakula, 1969; Starmer et al., 1988; Ridley et al., 2012; Duneau and Lazzaro, 2018). To our knowledge, how interactions between microbial symbionts affect symbiont maintenance through *Drosophila* life stages remains however unknown. Here, we investigated the maintenance of a wild isolate of *Saccharomyces cerevisiae* yeast from larvae to young adults *Drosophila melanogaster*. We found that yeast presence in adults only occurred when larvae were associated with symbiotic bacteria and its frequency depended on the identity of these bacteria.

Material and methods

Biological material

We used a *Drosophila melanogaster* (Meigen, 1830) Oregon-R strain usually maintained on banana medium (233 g.L^{-1} banana, 62 g.L^{-1} sugar, 62 g.L^{-1} dead yeast, 25 g.L^{-1} ethanol, 10 g.L^{-1} agar and 5 g.L^{-1} nipagin) at 21°C with a 14 h/10 h day-night cycle. The four bacterial strains used had been isolated from feces of adult Oregon-R flies and had been described in Guilhot et al. (2019). Bacterial strains were identified as *Staphylococcus* sp. (accession number MK461976 in the NCBI database), *Enterococcus* sp. (MK461977), an Enterobacteriaceae (MK461978) and an Actinobacteria (MK461979). The taxonomic resolution of our analyses is unfortunately modest and these bacteria do not usually dominate the *Drosophila* microbiota. However similar strains had already been identified as associated with Drosophilids (Chandler et al., 2011; Staubach et al., 2013). More importantly, microbial species and strains can evolve rapidly and with large consequences on their effects on host phenotype (Winans et al., 2017; Martino et al., 2018). For this reason, the proper description of symbiont effects on host phenotype in relevant experimental conditions (Guilhot et al. 2019) may be more important than high taxonomical resolution to understand symbiosis.

The yeast *Saccharomyces cerevisiae* (Meyen ex Hansen, 1883) strain was isolated from a wild Drosophilid in the *'Le Domaine de l'Hortus'* vineyard, near Montpellier in Southern France.

Experimental design

The main experiment was conducted on sterile vials. Each experimental unit consisted of twenty *D. melanogaster* eggs free of cultivable bacteria and yeast deposited on an artificial wound of a surface-sterilized grape berry that was inoculated, or not, with specific microorganisms. Eggs were laid by conventionally-reared *D. melanogaster* Oregon-R females on solidified grape juice, which contained the antibiotic streptomycin (1 mg.L^{-1}, from a standard streptomycin solution of 1 mg.ml^{-1} in 1 mM EDTA (Sigma-Aldrich ref. 85886)) in order to remove parental bacteria. Grape berries were surface-sterilized; they were hence dipped in a 2% bleach solution, rinsed with sterile water and dried before use. We ensure these procedures were efficient: no cultivable bacteria or yeast were found in surface-sterilized berries and eggs homogenates. Every grape berry received 10^4 *S. cerevisiae* yeast cells suspended in 10 µl. Berries inoculated with *S. cerevisiae* were then allocated to six

different bacterial treatments: no bacteria (control, n = 18); one of the four bacterial strains

described above (10^4 cells of the same type: n (*Staphylococcus*) = 13, n (*Enterococcus*) = 13,

n (Enterobacteriaceae) = 11, n (Actinobacteria) = 13); and a mixture of the four bacteria

(2.5×10^3 cells of each type, n = 9). Replicates were organized in eleven blocks launched over

four days.

Newly formed pupae were aseptically removed daily from their larval container and placed in

a sterile new vial until adult emergence. This procedure mimicked natural insect behavior as

D. melanogaster larvae usually crawl out of their substrate before pupation (Sokolowski et al.,

1986; Woltz and Lee, 2017), which incidentally prevent the exposure of young adults to the

microorganisms present in the larval substrate. It was not logistically feasible to process every

adult independently. We therefore elected to assess microbial content in groups of adults that

emerged on the same day. For each grape berry, we randomly selected a single pupa and

pooled all the adults, females as males, that emerged on the day as this pupa. This protocol

allowed the detection of yeast (and bacterial) cells associated with freshly emerged adults that

may have been present externally or internally. Sampled adults were homogenized in sterile

PBS using a Tissue Lyser II (Qiagen) and Ø3 mm glass balls. Serially diluted fly samples

were plated on Lysogeny Broth (LB) plates to count microbial cells after incubation at 24°C.

Colonies of the five microbial symbionts (yeast and bacteria) were distinguished according to

their morphology (see Guilhot et al. 2019).

Effects of bacteria on yeast development in the larval fruit may explain some of our results.

We hence collected the remaining juice from grape berries two days after the formation of the

last pupa. Although quantifying yeast in fruit when larvae were feeding would have been

informative too, we elected to not disturb the development of the larvae and waited the larvae

left their substrate. Serially diluted fruit samples were plated on LB plates to count yeast

colonies at 24°C. As above, microbial colonies were distinguished based on their

morphology.

In parallel to the main experiment, bacteria were also inoculated without yeast on cubes of

laboratory medium (which contains dead yeast) following the procedure above to assess

bacteria transstadial maintenance in their environment of origin (i.e. the banana medium used

to rear the fly colony and described above). The six different bacterial treatments were: no

bacteria (control, n = 12); one of the four bacterial strains described above (n

(*Staphylococcus*) = 12, n (*Enterococcus*) = 7, n (Enterobacteriaceae) = 11, n (Actinobacteria)

= 11); and a mixture of the four bacteria (n = 14). Replicates were organized in fifteen blocks

launched over four days.

Statistical analyses

We hypothesized larval bacteria would affect yeast transstadial maintenance. We estimated

this phenomenon in groups of 1 to 11 freshly emerged adults (median = 5, IQR = 4). Yeast-

positive samples contained 1 to 150 cells per adult fly (Fig. S1). This variation was not

investigated statistically due to low statistical power. Whether live yeast cells were present or

not was analyzed using a generalized linear model with binomial distribution and logit link

function. Tested factors comprised bacterial treatment, number of adults in the groups, yeast

concentration in the fruit, age of the adults and experimental block. Statistical power did not

enable testing the interaction between bacterial treatment and number of adults in the groups

(but see Fig. S2). Because we used groups of adult flies it was mandatory to take into account

the number of individuals per pool. The biological material employed (i.e. wild and laboratory

strains and populations) informs on the factors that can influence transstadial symbiont

maintenance in a qualitative fashion, but could not indicate their quantitative occurrence in

the field. Backward model selection allowed to eliminate non-significant terms (i.e. yeast

concentration in the fruit and age of the pooled flies) from the initial complete model.

Contrasts were used to detect significant differences between bacterial treatment levels.

Numbers of replicates varied among bacterial treatments due to differential larval mortality.

However, the analysis of larval survival revealed no significant effect of the bacteria on this

trait (Guilhot et al., 2019).

To study the effect of the bacterial treatment on the yeast concentration (log-transformed) in

larval fruit substrate, we used a linear mixed model with Restricted Maximum Estimate

Likelihood. Experimental block was defined as a random factor.

Analyzes were performed with JMP (SAS, 14.1).

Results

Bacterial treatment significantly affected *S. cerevisiae* presence in freshly emerged adult flies ($\chi^2 = 20.30$, df = 5, p = 0.001). Yeast was never observed in adult flies that emerged from control treatments, unlike treatments with bacteria at the larval stage (contrast 'All treatments with bacteria' vs 'Control': $\chi^2 = 11.2$, df = 1, p < 0.0001) (Fig. 1). Young adult flies that had developed with the Enterobacteriaceae alone or in mixture with the other bacteria were more likely to harbor live yeast cells than the other treatments with bacteria at the larval stage (contrast 'With Enterobacteriaceae' vs 'All other treatments with bacteria': $\chi^2 = 4.52$, df = 1, p = 0.03) (Fig. 1). As expected, the number of individuals in the assayed pool significantly and positively affected the likelihood of yeast observation ($\chi^2 = 7.54$, df = 1, p = 0.01) (Fig. S2) – supporting the need to include this factor in all the analyses. The age of freshly emerged adult flies ($\chi^2 = 0.65$, df = 1, p = 0.42) and the yeast concentration in the larval medium ($\chi^2 = 0$, df = 1, p = 1) had not significant influence on yeast presence in adults.

The bacterial treatment had no significant effect on the yeast concentration in the medium two days after the formation of the last pupa ($F_{5,49} = 1.18$, p = 0.33) (Fig. 2). Yeast presence in fruit flesh was detected in all replicates but one.

Bacteria could be observed in young adults that emerged from most combinations of larval environment and bacterial treatment (Fig. S3). However, the proportion of bacteria-positive groups never exceeded 25%. When bacteria were detected, load varied from 1 to 33 bacterial cells per adult fly (data not analyzed statistically). The observation of the inoculated bacteria in emerged adults shows these bacteria sampled in laboratory adults reared on artificial medium could establish symbiosis with larvae, even in fruit substrate.

Discussion

Our most important result is that larval bacteria influenced yeast transstadial maintenance (Fig. 1). In control treatments that were not inoculated by bacteria, yeast was never found in freshly emerged adult flies. On the contrary, the presence of bacteria at the larval stage favored yeast maintenance through host metamorphosis. In particular, inoculation by the Enterobacteriaceae bacterium (alone or in mixture) led to greater *S. cerevisiae* transstadial maintenance than the other bacterial treatments (Fig. 1). The propensity to favor yeast maintenance hence seemed to vary among bacteria.

It is well known that coinfecting symbionts (mutualistic as parasitic) often affect each other's horizontal transmission to new hosts in holometabolous insects (Azambuja et al., 2005; Fellous and Koella, 2009; Gendrin and Christophides, 2013; Hegde et al., 2015) and other multicellular organisms (Azambuja et al., 2005; Lass et al., 2013; Barret et al., 2016; Bonnet et al., 2017; Zélé et al., 2018). However, we know a single other case of microbial interactions affecting symbiont maintenance through complete metamorphosis: in *Galleria mellonella* butterflies, the bacterium *Enterococcus mundtii* interacts with host immunity during the pupal stage to shape adult bacterial microbiota (Johnston and Rolff, 2015). Although we used fresh fruit and a wild yeast strain, flies and bacteria were laboratory sourced. Our experiment hence shows bacteria affect yeast transstadial maintenance in *D. melanogaster*, but further work will be necessary to unveil the pervasiveness of this phenomenon in the field.

What mechanisms may underlie symbiont transstadial maintenance, and how did bacteria affect it? The maintenance of *S. cerevisiae* yeast and several bacterial strains through *Drosophila* metamorphosis are congruent with previous reports of the transstadial maintenance of extracellular microbial symbionts in Drosophilids (Bakula, 1969; Starmer et al., 1988; Ridley et al., 2012; Duneau and Lazzaro, 2018; Téfit et al., 2018), other Dipterans (Radvan, 1960; Capuzzo et al., 2005; Rochon et al., 2005; Damiani et al., 2008; Lauzon et al., 2009; Gendrin and Christophides, 2013; Nayduch and Burrus, 2017; Majumder et al., 2020) and other holometabolous insects (Hammer and Moran, 2019). Microbial symbionts could maintain on inner or outer walls of the pupal chamber (Kaltenpoth et al., 2010; Wang and Rozen, 2017). In *Drosophila melanogaster*, bacterial cells of *Escherichia coli* were found associated with the inner pupal membrane (Bakula, 1969). Alternatively, adults might retrieve symbionts by consuming their own meconium - the remaining of larval midgut that is excreted right after adult emergence (Moll et al., 2001; Broderick and Lemaitre, 2012;

Gendrin and Christophides, 2013). The mechanism of bacterial influence on yeast
maintenance through metamorphosis is not trivial either. The Enterobacteriaceae that favored
yeast maintenance, despite presenting a wide metabolic spectrum (Guilhot et al., 2019),
probably not improved fruit quality by concentrating or synthetizing nutrients (Ramiro et al.,
2016) as had no significant effect on fly phenotype in this context (Guilhot et al., 2019).
Besides, the concentration of yeast cells in fruit did not correlated with the presence of yeast
in the freshly emerged adults and was not affected by the bacterial treatment (Fig. 2). This
lack of quantitative relationships suggests yeast maintenance through metamorphosis may be
determined by qualitative processes rather than mere cell numbers. Several bacteria are
known to interact with *Drosophila* host signaling (e.g. Shin et al., 2011; Storelli et al., 2011).
Symbiotic bacteria could therefore elicit host or yeast physiological responses in a way that
would affect the likelihood of transstadial maintenance.

Yeast transstadial maintenance in *D. melanogaster* may have consequences for the spatial
spread of the yeast and the evolution of the symbiosis. Yeast needs active transport by insect
vectors to disperse among the ephemeral patch of resources formed by fruits (Starmer and
Lachance, 2011). *Drosophila* adults contribute to such yeast dispersal through two non-
excluding mechanisms. First, it is well established that yeasts produce chemical volatiles that
attract adult flies (Palanca et al., 2013; Buser et al., 2014; Scheidler et al., 2015; Anagnostou
et al., 2016; Bellutti et al., 2018; Günther et al., 2019; Lewis and Hamby, 2019), which favors
their acquisition and vectoring by insects to new resource patches (Buser et al., 2014).
Whether this phenomenon reflects yeast adaptation to insect vectoring is however debated
(Günther and Goddard, 2019). Second, yeast maintenance through *Drosophila* metamorphosis
- as demonstrated here - would enable the dispersal to new resource patches of larval
symbionts (e.g. fruit, possibly infested with insect larvae) by colonized emerging adults. Such
continuity in symbiosis over the life-cycle selects larval symbionts for beneficial effects on
host fitness (Ebert, 2013). The microbial strains the most beneficial to larval development (for
example in terms of larval survival) would be the ones best dispersed to new resources
patches by the vigorous or numerous adult hosts they favored the development of.
Furthermore, the maintenance of larval microbial symbionts until adult emergence may also
benefit the host as freshly emerged adults could be less susceptible to opportunistic pathogens
due to symbiont prior presence (Blum et al., 2013; Johnston and Rolff, 2015; Obadia et al.,
2017). To conclude, transstadial maintenance of larval symbionts has implication for the
dynamics and evolution of both hosts and microorganisms, the effects of bacteria on yeast we

report may therefore affect important aspects of symbiosis. These are new and anticipated consequences of insect association with bacteria.

Symbiont-symbiont interactions are emerging as key features of numerous taxa (Ferrari and Vavre, 2011; Álvarez-Pérez et al., 2019, Mathé-Hubert et al., 2019), including *Drosophila* flies (Fischer et al., 2017; Gould et al., 2018). Studying microbial symbionts one by one may be more tractable, however our experiment illustrates understanding the nature and diversity of host-symbiont relationships necessitates encompassing the complexity of natural communities.

Acknowledgements

We thank Edouard Jurkevitch, Natacha Kremer and Elodie Vercken for critical reading of an earlier version of this manuscript and Laure Benoit, Marie-Pierre Chapuis, Romain Gallet and Philippe Gautier for methodological help.

Competing interests

No competing interests declared.

Author contributions

R.G., A.R. and S.F. designed the experiment. R.G., A.R., A.X. and S.F. ran the experiment. K.H. collected the yeast isolate. R.G. and S.F. analyzed the data and wrote the manuscript.

Funding

This work was supported by French National Research Agency through the 'SWING' project (ANR-16-CE02-0015) and by Agropolis Fondation under the reference ID 1505-002 through the 'Investissements d'avenir' program (Labex Agro: ANR-10-LABX-0001-01).

Data availability

The dataset is available in the open data repository Zenodo (doi: 10.5281/zenodo.3546129) (Guilhot et al., 2020).

References

Alizon, S., De Roode, J.C. and Michalakis, Y. (2013). Multiple infections and the evolution of virulence. *Ecology letters* **16**, 556-567.

Álvarez-Pérez, S., Lievens, B. and Fukami, T. (2019). Yeast–bacterium interactions: the next frontier in nectar research. *Trends in plant science* **24**, 393-401.

Anagnostou, C., Dorsch, M. and Rohlfs, M. (2010). Influence of dietary yeasts on *Drosophila melanogaster* life-history traits. *Entomologia Experimentalis et Applicata* **136**, 1-11.

Azambuja, P., Garcia, E.S. and Ratcliffe, N.A. (2005). Gut microbiota and parasite transmission by insect vectors. *Trends in parasitology* **21**, 568-572.

Bakula, M. (1969). The persistence of a microbial flora during postembryogenesis of *Drosophila melanogaster*. *Journal of invertebrate pathology* **14**, 365-374.

Barret, M., Guimbaud, J.F., Darrasse, A. and Jacques M.A. (2016). Plant microbiota affects seed transmission of phytopathogenic micro-organisms. *Mol. Plant. Pathol.* **17**, 791-795.

Becher, P.G., Flick, G., Rozpędowska, E., Schmidt, A., Hagman, A., Lebreton, S., Larsson, M.C., Hansson, B.S., Piškur, J. and Bengtsson, M. (2012). Yeast, not fruit volatiles mediate *Drosophila melanogaster* attraction, oviposition and development. *Functional Ecology* **26**, 822-828.

Bellutti, N., Gallmetzer, A., Innerebner, G., Schmidt, S., Zelger, R. and Koschier, E.H. (2018). Dietary yeast affects preference and performance in *Drosophila suzukii*. *Journal of pest science* **91**, 651-660.

Blum, J.E., Fischer, C.N., Miles, J. and Handelsman, J. (2013). Frequent replenishment sustains the beneficial microbiome of *Drosophila melanogaster*. *MBio* **4**, e00860-13.

Bonnet, S.I., Binetruy, F., Hernández-Jarguín, A.M. and Duron, O. (2017). The tick microbiome: why non-pathogenic microorganisms matter in tick biology and pathogen transmission. *Frontiers in Cellular and Infection Microbiology* **7**, 236.

Broderick, N.A. and Lemaitre, B. (2012). Gut-associated microbes of *Drosophila melanogaster*. *Gut microbes* **3**, 307-321.

Buser, C.C., Newcomb, R.D., Gaskett, A.C. and Goddard, M.R. (2014). Niche construction initiates the evolution of mutualistic interactions. *Ecology Letters* **17**, 1257-1264.

Callens, M., Watanabe, H., Kato, Y., Miura, J. and Decaestecker, E. (2018). Microbiota inoculum composition affects holobiont assembly and host growth in *Daphnia*. *Microbiome* **6**, 56.

Capuzzo, C., Firrao, G., Mazzon, L., Squartini, A. and Girolami, V. (2005). '*Candidatus* Erwinia dacicola', a coevolved symbiotic bacterium of the olive fly *Bactrocera oleae* (Gmelin). *International journal of systematic and evolutionary microbiology* **55**, 1641-1647.

Carbonetto, B., Ramsayer, J., Nidelet, T., Legrand, J. and Sicard, D. (2018). Bakery yeasts, a new model for studies in ecology and evolution. *Yeast* **35**, 591-603.

Chandler, J. A., Lang, J. M., Bhatnagar, S., Eisen, J. A., and Kopp A. (2011). Bacterial communities of diverse *Drosophila* species: ecological context of a host–microbe model system. *PLoS Genetics* **7**, e1002272.

Chandler, J.A., Eisen, J.A. and Kopp, A. (2012). Yeast communities of diverse *Drosophila* species: comparison of two symbiont groups in the same hosts. *Appl. Environ. Microbiol.* **78**, 7327-7336.

Comolli, L.R. (2014). Intra-and inter-species interactions in microbial communities. *Frontiers in microbiology* **5**, 629.

Damiani, C., Ricci, I., Crotti, E., Rossi, P., Rizzi, A., Scuppa, P., Esposito, F., Bandi, C., Daffonchio, D. and Favia, G. (2008). Paternal transmission of symbiotic bacteria in malaria vectors. *Current Biology* **18**, R1087-R1088.

Duneau, D.F. and Lazzaro, B.P. (2018). Persistence of an extracellular systemic infection across metamorphosis in a holometabolous insect. *Biology letters* **14**, 20170771.

Ebert, D. (2013). The epidemiology and evolution of symbionts with mixed-mode transmission. *Annual Review of Ecology, Evolution, and Systematics* **44**, 623-643.

Fellous, S. and Koella, J.C. (2009). Infectious dose affects the outcome of the within-host competition between parasites. *The American Naturalist* **173**, E177-E184.

Ferrari, J. and Vavre, F. (2011). Bacterial symbionts in insects or the story of communities affecting communities. *Philosophical Transactions of the Royal Society B: Biological Sciences* **366**, 1389-1400.

Fischer, C.N., Trautman, E.P., Crawford, J.M., Stabb, E.V., Handelsman, J. and Broderick, N.A. (2017). Metabolite exchange between microbiome members produces compounds that influence *Drosophila* behavior. *Elife* **6**, e18855.

Ganter, P.F. (1988). The vectoring of cactophilic yeasts by *Drosophila*. *Oecologia* **75**, 400-404.

Gendrin, M. and Christophides, G.K. (2013). The *Anopheles* mosquito microbiota and their impact on pathogen transmission. In *Anopheles mosquitoes-New insights into Malar. vectors*. InTech.

Gilbert, D.G. (1980). Dispersal of yeasts and bacteria by *Drosophila* in a temperate forest. *Oecologia* **46**, 135-137.

Gould, A.L., Zhang, V., Lamberti, L., Jones, E.W., Obadia, B., Korasidis, N., Gavryushkin, A., Carlson, J.M., Beerenwinkel, N. and Ludington, W.B. (2018). Microbiome interactions shape host fitness. *Proceedings of the National Academy of Sciences* **115**, E11951-E11960.

Guilhot, R., Rombaut, A., Xuéreb, A., Howell, K. and Fellous, S. (2019). Environmental specificity in *Drosophila*-bacteria symbiosis affects host developmental plasticity. *bioRxiv* doi:10.1101/717702, version 3: peer-reviewed and recommended by *PCI Evolutionary Biology* doi:10.24072/pci.evolbiol.100085

Guilhot, R., Rombaut, A., Xuéreb, A., Howell, K. and Fellous, S. (2020). Data from: Bacterial influence on the maintenance of symbiotic yeast through *Drosophila* metamorphosis. *Zenodo*. https://zenodo.org/record/3546129#.Xpjl7XvgrIV

Günther, C.S. and Goddard, M.R. (2019) Do yeasts and *Drosophila* interact just by chance? *Fungal Ecology* **38**, 37-43.

Günther, C.S., Knight, S.J., Jones, R. and Goddard, M.R. (2019). Are *Drosophila* preferences for yeasts stable or contextual? *Ecology and evolution* **9**, 8075-8086.

Hammer, T.J. and Moran, N.A. (2019). Links between metamorphosis and symbiosis in holometabolous insects. *Philosophical Transactions of the Royal Society B* **374**, 20190068.

Hassani, M.A., Durán, P. and Hacquard, S. (2018). Microbial interactions within the plant holobiont. *Microbiome* **6**, 58.

Hegde, S., Rasgon, J.L. and Hughes, G.L. (2015). The microbiome modulates arbovirus transmission in mosquitoes. *Current opinion in virology* **15**, 97-102.

Hoang, D., Kopp, A., and Chandler, J. A. (2015). Interactions between *Drosophila* and its natural yeast symbionts - Is *Saccharomyces cerevisiae* a good model for studying the fly-yeast relationship? *PeerJ* **3**, e1116.

Johnston, P.R. and Rolff, J. (2015). Host and symbiont jointly control gut microbiota during complete metamorphosis. *PLoS Pathogens* **11**, e1005246.

Jouhten, P., Ponomarova, O., Gonzalez, R. and Patil, K.R. (2016). *Saccharomyces cerevisiae* metabolism in ecological context. *FEMS yeast research* **16**, fow080.

Kaltenpoth, M., Goettler, W., Koehler, S. and Strohm, E. (2010). Life cycle and population dynamics of a protective insect symbiont reveal severe bottlenecks during vertical transmission. *Evolutionary ecology* **24**, 463-477.

Kobayashi, D.Y. and Crouch, J.A. (2009). Bacterial/fungal interactions: from pathogens to mutualistic endosymbionts. *Annual review of phytopathology* **47**, 63-82.

Lass, S., Hudson, P.J., Thakar, J., Saric, J., Harvill, E., Albert, R. and Perkins, S.E. (2013). Generating super-shedders: co-infection increases bacterial load and egg production of a gastrointestinal helminth. *Journal of the Royal Society Interface* **10**, 20120588.

Lauzon, C.R., McCombs, S.D., Potter, S.E. and Peabody, N.C. (2009). Establishment and vertical passage of *Enterobacter (Pantoea) agglomerans* and *Klebsiella pneumoniae* through all life stages of the Mediterranean fruit fly (Diptera: Tephritidae). *Annals of the Entomological Society of America* **102**, 85 – 95.

Leitão-Gonçalves, R., Carvalho-Santos, Z., Francisco, A.P., Fioreze, G.T., Anjos, M., Baltazar, C., Elias, A.P., Itskov, P.M., Piper, M.D.W. and Ribeiro, C. (2017). Commensal

bacteria and essential amino acids control food choice behavior and reproduction. *PLoS Biology* **15**, e2000862.

Lewis, M.T. and Hamby, K.A. (2019). Differential impacts of yeasts on feeding behavior and development in larval *Drosophila suzukii* (Diptera: Drosophilidae). *Scientific reports* **9**, 1-12.

Majumder, R., Sutcliffe, B., Taylor, P.W. and Chapman, T.A. (2020). Microbiome of the Queensland fruit fly through metamorphosis. *Microorganisms* **8**, 795.

Martino, M.E., Joncour, P., Leenay, R., Gervais, H., Shah, M., Hughes, S., Gillet, B., Beisel, C. and Leulier, F. (2018). *Cell host & microbe* **24**, 109-119.

Mathé-Hubert, H., Kaech, H., Hertaeg, C., Jaenike, J. and Vorburger, C. (2019). Nonrandom associations of maternally transmitted symbionts in insects: The roles of drift versus biased cotransmission and selection. *Molecular ecology* **28**, 5330-5346.

Moll, R.M., Romoser, W.S., Modrakowski, M.C., Moncayo, A.C. and Lerdthusnee, K. (2001). Meconial peritrophic membranes and the fate of midgut bacteria during mosquito (Diptera: Culicidae) metamorphosis. *Journal of medical entomology* **38**, 29-32.

Nayduch, D. and Burrus, R.G. (2017). Flourishing in filth: house fly–microbe interactions across life history. *Annals of the Entomological Society of America* **110**, 6-18.

Newell, P.D. and Douglas, A.E. (2014). Interspecies interactions determine the impact of the gut microbiota on nutrient allocation in *Drosophila melanogaster*. *Appl. Environ. Microbiol.* **80**, 788-796.

Obadia, B., Güvener, Z.T., Zhang, V., Ceja-Navarro, J.A., Brodie, E.L., William, W.J. and Ludington, W.B. (2017). Probabilistic invasion underlies natural gut microbiome stability. *Current Biology* **27**, 1999-2006.

Pais, I.S., Valente, R.S., Sporniak, M. and Teixeira, L. (2018). *Drosophila melanogaster* establishes a species-specific mutualistic interaction with stable gut-colonizing bacteria. *PLoS Biology* **16**, e2005710.

Palanca, L., Gaskett, A.C., Günther, C.S., Newcomb, R.D. and Goddard, M.R. (2013). Quantifying variation in the ability of yeasts to attract *Drosophila melanogaster*. *PLoS One* **8**, e75332.

Ramiro, R.S., Pollitt, L.C., Mideo, N. and Reece, S.E. (2016). Facilitation through altered resource availability in a mixed-species rodent malaria infection. *Ecology letters* **19**, 1041-1050.

Radvan, R. (1960). Persistence of bacteria during development in flies. *Folia Microbiologica* **5**, 50.

Ridley, E.V., Wong, A.C., Westmiller, S. and Douglas, A.E. (2012). Impact of the resident microbiota on the nutritional phenotype of *Drosophila melanogaster*. *PloS One* **7**, e36765.

Rochon, K., Lysyk, T.J. and Selinger, L.B. (2005). Retention of *Escherichia coli* by house fly and stable fly (Diptera: Muscidae) during pupal metamorphosis and eclosion. *Journal of medical entomology* **42**, 397 – 403.

Rohlfs, M. and Hoffmeister, T.S. (2005). Maternal effects increase survival probability in *Drosophila subobscura* larvae. *Entomologia Experimentalis et Applicata* **117**, 51-58.

Ryu, J.H., Kim, S.H., Lee, H.Y., Bai, J.Y., Nam, Y.D., Bae, J.W., Lee, D.G., Shin, S.C., Ha, E.M. and Lee, W.J. (2008). Innate immune homeostasis by the homeobox gene caudal and commensal-gut mutualism in *Drosophila*. *Science* **319**, 777-782.

Scheidler, N.H., Liu, C., Hamby, K.A., Zalom, F.G. and Syed, Z. (2015). Volatile codes: correlation of olfactory signals and reception in *Drosophila*-yeast chemical communication. *Scientific reports* **5**, 14059.

Seth, E.C. and Taga, M.E. (2014). Nutrient cross-feeding in the microbial world. *Frontiers in microbiology* **5**, 350.

Shin, S.C., Kim, S.H., You, H., Kim, B., Kim, A.C., Lee, K.A., Yoon, J.H., Ryu, J.H. and Lee, W.J. (2011). *Drosophila* microbiome modulates host developmental and metabolic homeostasis via insulin signaling. *Science* **334**, 670-674.

Sokolowski, M.B., Bauer, S.J., Wai-Ping, V., Rodriguez, L., Wong, J.L. and Kent, C. (1986). Ecological genetics and behaviour of *Drosophila melanogaster* larvae in nature. *Animal Behaviour* **34**, 403-408.

Sommer, A.J. and Newell, P.D. (2019). Metabolic basis for mutualism between gut bacteria and its impact on the *Drosophila melanogaster* host. *Appl. Environ. Microbiol.* **85**, e01882-18.

Stamps, J.A., Yang, L.H., Morales, V.M. and Boundy-Mills, K.L. (2012). *Drosophila* regulate yeast density and increase yeast community similarity in a natural substrate. *PLoS One* **7**, e42238.

Starmer, W.T., Peris, F. and Fontdevila, A. (1988). The transmission of yeasts by *Drosophila buzzatii* during courtship and mating. *Animal behaviour* **36**, 1691-1695.

Starmer, W.T. and Lachance, M.A. (2011). Yeast ecology. In *The Yeasts: a Taxonomic Study*, pp. 65-83. Elsevier, Waltham, MA.

Staubach, F., Baines, J. F., Künzel, S., Bik, E. M. and Petrov, D. A. (2013). Host species and environmental effects on bacterial communities associated with *Drosophila* in the laboratory and in the natural environment. *PloS One* **8**, e70749.

Storelli, G., Defaye, A., Erkosar, B., Hols, P., Royet, J. and Leulier, F. (2011). *Lactobacillus plantarum* promotes *Drosophila* systemic growth by modulating hormonal signals through TOR-dependent nutrient sensing. *Cell metabolism* **14**, 403-414.

Téfit, M.A., Gillet, B., Joncour, P., Hughes, S. and Leulier, F. (2018). Stable association of a *Drosophila*-derived microbiota with its animal partner and the nutritional environment throughout a fly population's life cycle. *Journal of insect physiology* **106**, 2-12.

Tollenaere, C., Susi, H. and Laine, A.L. (2016). Evolutionary and epidemiological implications of multiple infection in plants. *Trends in plant science* **21**, 80-90.

Wang, Y. and Rozen, D.E. (2017). Gut microbiota colonization and transmission in the burying beetle *Nicrophorus vespilloides* throughout development. *Appl. Environ. Microbiol.* **83**, e03250-16.

Wargo, M.J. and Hogan, D.A. (2006). Fungal - bacterial interactions: a mixed bag of mingling microbes. *Current opinion in microbiology* **9**, 359-364.

Winans, N.J., Walter, A., Chouaia, B., Chaston, J.M., Douglas, A.E. and Newell, P.D. (2017). A genomic investigation of ecological differentiation between free-living and *Drosophila*-associated bacteria. *Molecular ecology* **26**, 4536-4550.

Woltz, J.M. and Lee, J.C. (2017). Pupation behavior and larval and pupal biocontrol of *Drosophila* suzukii in the field. *Biological control* **110**, 62-69.

Wong, A.C.N., Dobson, A.J. and Douglas, A.E. (2014). Gut microbiota dictates the metabolic response of *Drosophila* to diet. *Journal of Experimental Biology* **217**, 1894-1901.

Zélé, F., Magalhães, S., Kéfi, S. and Duncan, A.B. (2018). Ecology and evolution of facilitation among symbionts. *Nature communications* **9**, 4869.

Figures

Fig. 1. Transstadial maintenance of *Saccharomyces cerevisiae* in response to bacterial treatment. Symbols indicate the proportion of groups of freshly emerged adult flies containing yeast per bacterial treatment (n = number of adult groups per bacterial treatment). 95% binomial confidence intervals were calculated using normal approximation method. These results are qualitative as we used groups of adult flies to estimate yeast transstadial maintenance (Fig S2).

Fig. 2. Yeast concentration in grape berry flesh after the formation of the last pupa. Concentration is expressed in number of yeast cells per 200 µl of fruit flesh. Symbols indicate mean ± s.e.m.