1   Title

2   **Instantaneous movement-unrelated midbrain activity**
3   **modifies ongoing eye movements**
4

Authors

Antimo Buonocore[1,2], Matthias P. Baumann[1,2], Ziad M. Hafed[1,2]

1. Werner Reichardt Centre for Integrative Neuroscience, Tübingen University,
Tübingen, BW, 72076, Germany
2. Hertie Institute for Clinical Brain Research, Tübingen University,
Tübingen, BW, 72076, Germany
12

Corresponding author:

Antimo Buonocore
Werner Reichardt Centre for Integrative Neuroscience
Otfried-Müller Str. 25
Tübingen, 72076, Germany
Tel: +49 7071 29 88821
antimo.buonocore@cin.uni-tuebingen.de
21

Abstract

At any moment in time, new information is sampled from the environment and interacts with ongoing brain state. Often, such interaction takes place within individual circuits that are capable of both mediating the internally ongoing plan as well as representing exogenous sensory events. Here we investigated how sensory-driven neural activity can be integrated, very often in the same neuron types, into ongoing oculomotor commands for saccades. Despite the ballistic nature of saccades, visually-induced action potentials in the superior colliculus (SC), a structure known to drive eye movements, not only occurred intra-saccadically, but they were also associated with highly predictable modifications of the ongoing eye movements. Such modifications were also possible by peri-saccadically injecting single, double, or triple electrical microstimulation pulses into the SC. Our results suggest instantaneous readout of the SC map during movement generation, irrespective of activity source, and explain a significant component of kinematic variability of motor outputs.

Introduction

A hallmark of the central nervous system is its ability to process an incredibly complex amount of incoming information from the environment in parallel. This is achieved through multiplexing of functions, either at the level of individual brain areas or even at the level of individual neurons themselves. For example, in different motor modalities like arm (Alexander and Crutcher, 1990; Shen and Alexander, 1997; Breveglieri et al., 2016) or eye (Goldberg and Wurtz, 1972a, b; Wurtz and Goldberg, 1972; Mohler and Wurtz, 1976; Bruce and Goldberg, 1985) movements, a large fraction of the neurons contributing to the motor command are also intrinsically sensory in nature, hence being described as sensory-motor neurons. In this study, we aimed to investigate the implications of such sensory and motor multiplexing using vision and the oculomotor system as our model of choice.

A number of brain areas implicated in eye movement control, such as the midbrain superior colliculus (SC) (Wurtz and Albano, 1980; Munoz and Wurtz, 1995), frontal eye fields (FEF) (Bruce and Goldberg, 1985; Schall and Hanes, 1993; Schall et al., 1995; Tehovnik et al., 2000), and lateral intra-parietal area (LIP) (Mazzoni et al., 1996), contain many so-called visual-motor neurons. These neurons burst both in reaction to visual stimuli entering into their response fields (RF's) as well as in association with triggering eye movements towards these RF's. In some neurons, for example in the SC (Mohler and Wurtz, 1976; Mays and Sparks, 1980; Edelman and Goldberg, 2001; Willeke et al., 2019), even the motor bursts themselves are contingent on the presence of a visual target at the movement endpoint. In the laboratory, the properties of visual and motor bursts are frequently studied in isolation, by dissociating the time of visual onsets (evoking "visual" bursts) from the time of saccade triggering (evoking "motor" bursts). However, in real life, exogenous sensory events can happen at any time in relation to our own ongoing internal state. Thus, "visual" spikes at one visual field location may, in principle, be present at the same time as "motor" spikes for a saccade to another location. What are the implications of such simultaneity? Answering this

question is important to clarify mechanisms of readout from circuits in which functional

multiplexing is prevalent.

In the SC, our focus here, there have been many debates about how this structure

contributes to saccade control (Waitzman et al., 1991; Ivan et al., 2018). In recent proposals

(Goossens and Van Opstal, 2006; Van Opstal and Goossens, 2008; Goossens and Van

Opstal, 2012), it was suggested that every spike emitted by SC neurons during their "motor"

bursts contributes a mini-vector of movement tendency, such that the aggregate sum of all

output spikes is read out by downstream structures to result in a given movement trajectory.

However, implicit in these models is the assumption that only action potentials within a

narrow time window around movement triggering (the "motor" burst) matter. Any other

spiking, by the same or other neurons, before or after the eye movement is irrelevant. This

causes a significant readout problem, since downstream neurons do not necessarily have

the privilege of knowing which spikes should now count for a given eye movement

implementation and which not.

Indeed, from an ecological perspective, an important reason for multiplexing could be

exactly to maintain flexibility to rapidly react to the outside world, even in a late motor control

structure. In that sense, rather than invoking mechanisms that allow actively ignoring "other

spiking" activity outside of the currently triggered eye movement (whether spatially or

temporally), one would predict that SC readout, at any one moment, should be quite

sensitive to any spiking activity regardless of its source. We experimentally tested this

hypothesis. We "injected" SC spiking activity around the time of saccade generation, but at

a spatially dissociated location. We uncovered causal evidence that the entire landscape of

SC activity can instantaneously contribute to individual saccade metrics, explaining a

component of behavioral variability previously unaccounted for.

Results

Stimulus-driven SC "visual" bursts can occur intra-saccadically

We first tested the hypothesis that visually-induced action potentials can occur in the SC intra-saccadically; that is, simultaneously with motor-related bursts. We exploited the topographic nature of the SC in representing visual and motor space (Cynader and Berman, 1972; Robinson, 1972; Chen et al., 2019). We asked two monkeys (P and N) to maintain steady fixation on a central spot. Prior work has shown that this condition gives rise to frequent microsaccades, which are associated with movement-related bursts in the rostral region of the SC representing small visual eccentricities and movement vectors (Hafed et al., 2009; Hafed and Krauzlis, 2012a; Chen et al., 2019; Willeke et al., 2019). We then presented a visual stimulus at a more eccentric location, and we recorded neural activity at this location (Fig. 1A, B). Depending on the timing of the visual stimulus relative to a given microsaccade, we could measure visual burst strength (in both visual and visual-motor neurons; Methods) either in isolation of microsaccades or when a microsaccade was in-flight. If SC visual bursts could still occur intra-saccadically, then one would expect that visual burst strength should be similar whether the burst timing happened when a microsaccade was being triggered or not. We ensured that all sites did not simultaneously burst for microsaccade generation (Figs. 1B, S1), to ensure that we were only measuring visual bursts and not concurrent movement-related activity. Such movement-related activity was expectedly in more rostral SC sites, representing foveal visual eccentricities, as we also confirmed in the same two monkeys (and sometimes in the very same sessions) in an earlier report (Chen et al., 2019).

Figure 1 Injecting arbitrary, movement-unrelated spiking activity into the SC map around the time of saccade generation. (A) A monkey steadily fixated while we presented an eccentric stimulus in a recorded neuron's RF (purple). The stimulus location was spatially dissociated from the motor range of microsaccades being generated (orange circles). This allowed us to experimentally inject movement-unrelated "visual" spikes into the SC map around the time of microsaccade generation. **(B)** We injected "visual" spikes at eccentric retinotopic locations (purple dots) distinct from the neurons that would normally exhibit motor bursts for microsaccades. The orange line and shaded area denote the mean and 95% confidence interval, respectively, of all microsaccade amplitudes that we observed. The neurons in which we injected "visual" spikes (purple dots) were not involved in generating these microsaccades (Fig. S1). The origin of the shown log-polar plot corresponds to 0.03 deg eccentricity (Hafed and Krauzlis, 2012b). **(C)** In a second experiment, a monkey generated a saccade to a given target location. We injected single, double, or triple "electrical" spikes into the SC map, again at a site unrelated to the currently generated saccade (purple RF and yellow symbol for the microstimulated site, versus orange for the saccade vector). The relative relationship between saccade target eccentricity and microstimulated SC site was similar to the relative relationship between movement sizes and "visual" spikes in **A**: in both cases, we injected movement-unrelated SC activity at a site more eccentric than the internally generated movement.

Regardless of microsaccade direction, "visual" bursts could still occur in the SC even if there was an ongoing eye movement. To illustrate this, Fig. 2A shows the stimulus-driven visual burst of an example neuron with and without concurrent microsaccades. The stimulus in this case consisted of a vertical sine wave grating of 40% or 80% contrast (Methods). The top panel of the figure shows the times of microsaccade onsets (crosses) and ends (squares) for all movements occurring in the session that overlapped with the interval of visual burst occurrence, which we defined to be 30-100 ms. This latter interval was chosen based on the bottom panel of the figure, showing the actual visual bursts. The gray curve shows average firing rate when there were no microsaccades from -100 ms to +150 ms relative to stimulus onset, and the orange curve shows average firing rate when the visual burst (shaded interval) coincided with at least a part of an ongoing microsaccade (top panel).

As can be seen, intra-saccadic "visual" bursts could still occur, and they were similar in

strength to saccade-free visual bursts. This was true across our population; for each neuron,

we plotted in Fig. 2B peak firing rate after stimulus onset when there was a concurrent

microsaccade being generated (inset schematic near the y-axis) as a function of peak firing

rate after stimulus onset when there was no concurrent microsaccade (inset schematic near

the x-axis). For this analysis, we pooled trials from the highest 3 contrasts (20%, 40%, and

80%) that we used in our study for simplicity (Methods), but similar conclusions could also

be reached for individual stimulus contrasts. There was no difference in visual burst strength

as a function of coincident microsaccades ($t(84) = 1.6905$, $p = 0.09$). Moreover, SC visual

bursts could still occur intra-saccadically whether the stimulus was activating the same SC

side generating a given movement or the opposite SC side (Fig. S2).

Therefore, at the time of movement execution (that is, at the time of a movement-

related burst in one part of the SC map), it is possible to have spatially dissociated visual

bursts in another part of the map. We next investigated how such additional "visual" spikes

(at an unrelated spatial location relative to the movements) affected the eye movements that

they were coincident with.

158

Figure 2 SC visual bursts can still occur intra-saccadically. **(A)** We measured the firing rate of an example neuron when a stimulus appeared inside its RF without any nearby microsaccades (gray firing rate curve) or when the same stimulus appeared while microsaccades were being executed around the time of visual burst occurrence (orange firing rate curve). The raster above the firing rate curves shows the individual microsaccades that occurred in the session for the orange firing rate curve (cross means movement onset, and square means movement end). For all of the movements, the visual burst overlapped with at least parts of the movements. Error bars denote 95% confidence intervals, and the shaded violet region denotes our estimate of visual burst interval (30-100 ms after stimulus onset). **(B)** At the population level, there was no difference in peak firing rate with saccades detected during a visual burst (y-axis and schematic near it) or without saccades around the visual burst (x-axis and schematic near it). The y-axis data correspond to the top left schematic view of the SC map (Hafed and Chen, 2016), in which extra-foveal visual spikes (purple) occurred in the SC concurrently with foveal motor spikes (orange) triggering microsaccades. The x-axis data correspond to the situation in the bottom right schematic, with no movement-related motor bursts at the time of the visually-induced spikes.

Peri-saccadic stimulus-driven "visual" bursts systematically influence eye movement metrics

If "visual" bursts can be present somewhere on the SC map at a time when "motor" bursts elsewhere on the map are to be read out by downstream neurons, then one might expect that each additional "visual" spike on the map should contribute to the executed movement metrics and cause a change in saccades. This would suggest a highly lawful relationship between the strength of the peri-saccadic "visual" burst and the amount of eye movement alteration that is observed. We explored this by relating the behavioral properties of the saccades in our task to the temporal relationship between their onset and the presence of "visual" spikes in the SC map caused by an unrelated stimulus onset.

We first confirmed a clear correlation between microsaccade amplitudes and eccentric stimulus onsets (Hafed and Ignashchenkova, 2013; Buonocore et al., 2017b; Malevich et al., 2020a). Our stimuli consisted of vertical sine wave gratings having different luminance contrasts (Methods). We plotted the time course of microsaccade amplitudes relative to grating onset for microsaccades that were spatially congruent with grating location (that is, having directions towards grating location; Methods). For the present analysis, we only focused on stimulus eccentricities of ≤4.5 deg because these had the strongest effects on microsaccades (Fig. S3). As expected (Hafed and Ignashchenkova, 2013; Buonocore et al., 2017b; Malevich et al., 2020a), there was a transient increase in microsaccade amplitude approximately 80-90 ms after grating onset (Fig. 3A). Critically, the increase reflected the stimulus properties, because it was stronger with higher stimulus contrast (main effect of contrast: $F(2,713) = 81.55$, $p < 1.27427*10^{-32}$), and there were also qualitatively different temporal dynamics: amplitude increases occurred earlier for higher (~75 ms) than lower (~85 ms) contrasts. Because we had simultaneously recorded neural data, we then analyzed, for the same trials, the SC visual bursts that were associated with the appearing gratings in these sessions. Visual bursts started earlier, and were stronger, for higher stimulus contrasts (Fig. 3B) (Li and Basso, 2008; Chen et al., 2015), similar to the amplitude changes in microsaccades. Moreover, the timing of the microsaccadic effects (Fig. 3A) was similar to the timing of the SC visual bursts (Fig. 3B), showing a short lag of ~20 ms relative to the bursts that is consistent with an efferent processing delay from SC neurons to the final extraocular muscle drive.

Therefore, as we hypothesized in previous reports (Hafed and Ignashchenkova, 2013; Buonocore et al., 2017b; Malevich et al., 2020a), not only is it possible for SC visual bursts to occur intra-saccadically (Fig. 2), but such bursts are temporally aligned with concurrent changes in microsaccade amplitudes (Fig. 3). We next uncovered a highly lawful impact of each injected extra "spike" per recorded neuron on saccade metrics.

Figure 3 Saccade metrics are altered when the movements coincide with SC visual bursts, and the alteration is related to SC visual burst strength. (A) Time course of saccade amplitude in our task relative to stimulus onset (vertical blue line). The data were subdivided according to stimulus contrast (three different colors representing the three highest contrasts in our task). Movements amplitudes were small (microsaccades) in the baseline pre-stimulus interval, but they sharply increased after stimulus onset, reaching a peak at around 70-80 ms. Moreover, the metric alteration clearly depended on stimulus contrast. **(B)** Normalized firing rates relative to stimulus onset (vertical blue line) for all extra-foveal neurons that we recorded simultaneously with the eye movement data in **A**. The alterations in movement metrics in **A** were strongly correlated in both time and amplitude with the properties of SC visual bursts. Error bars denote 95% confidence intervals. Figs. S3, S4 shows results with more eccentric neurons and stimuli (>4.5 deg).

221   The number of extra "visual" spikes per recorded neuron occurring intra-saccadically

222   was linearly related to metric alterations in microsaccades. For each eye movement towards

223   the recently appearing stimulus (that is, congruent with stimulus location), we counted how

224   many "visual" spikes by the concurrently recorded neuron occurred in the interval between

225   eye movement onset and saccade peak velocity. That is, we tested for the impact of the

number of extra "visual" spikes by a given recorded neuron as the SC population was being

read out by downstream pre-motor and motor structures to execute the triggered movement.

This per-neuron spike count was a proxy for how adding additional "visual" spikes in the SC

population at a site unrelated to the movement vector can "leak" downstream when the

saccade gate is opened. Moreover, since the extra spikes were more eccentric than the

sizes of the congruent microsaccades, we expected that the contribution would act to

increase microsaccade amplitudes (as in Fig. 3A). We focused, for now, on neurons at

eccentricities ≤4.5 deg (but still more eccentric than microsaccade amplitude; Fig. 1B)

because our earlier analyses showed that the clearest metric changes to tiny microsaccades

occurred under these circumstances (Figs. 3, S3, S4). We found a clear, lawful relationship

between the amount of "extra" spikes that occurred intra-saccadically on movement metrics.

These spikes were unrelated to the originally planned "motor" burst; they were spatially

dissociated but temporally coincident with saccade triggering, and they were also driven by

an exogenous visual stimulus onset. To demonstrate this observation, we plotted in Fig. 4A

the average microsaccadic eye movement trajectory in the absence of any additional SC

"visual" bursts (dark red; the curve labeled 0 spikes; Methods). We then plotted average

microsaccade size whenever any given recorded eccentric neuron had a visual burst such

that 1 spike of this visual burst happened to occur between movement onset and movement

peak velocity (red; 1 spike). The amplitude of the microsaccade was significantly larger than

with 0 spikes. We then progressively looked for movements with 2, 3, 4, 5, or 6 "visual"

spikes per recorded neuron that happened to occur intra-saccadically; there were

progressively larger and larger microsaccades (Fig. 4A). Across all data, the number of

"visual" spikes (per recorded neuron) that occurred intra-saccadically was lawfully and

linearly driving the amplitude increase of the (smaller) saccades (Fig. 4B) (Towards

condition, F-statistic vs. constant model: $F = 799$, $p<0.0001$; estimated coefficients :

intercept $= 0.1412$, $t = 30.0298$, $p<0.0001$; slope $= 0.0914$, $t = 28.26$, $p<0.0001$). On the

other hand, microsaccades directed opposite to the RF did not show any systematic modulation by the number of injected intra-saccadic spikes (Opposite condition, F-statistic vs. constant model: F = 3.83, p = 0.05; estimated coefficients: intercept = 0.11553, t = 57.13.0298, p<0.0001; slope = 0.0048, t = 1.9584, p = 0.05), suggesting that it is difficult to reduce microsaccade size below the already small amplitude of these tiny eye movements (Hafed, 2011).

These results suggest that there is an instantaneous specification of saccade metrics described by the overall activity present on the SC map, and they provide a much more nuanced view of the correlations between SC visual bursts and microsaccade amplitudes shown in Fig. 3. Every SC spike matters: all activity happening intra-saccadically and at locations of the SC map different from the saccade endpoint goal is interpreted as part of the motor command by downstream neurons.

For completeness, we also considered the same analyses as in Fig. 4 (that is, with congruent movements) but for more eccentric SC "visual" bursts (Fig. S5). The effects were present but in a significantly weaker manner than in Fig. 4, suggesting that the distance of the "extra" spiking activity on the SC map from the planned movement vector matters (Towards condition, F-statistic vs. constant model: F = 77.1, $p<2.13*10^{-18}$; estimated coefficients : intercept = 0.12254, t = 60.889, p<0.0001; slope = 0.012496, t = 8.7813, $p<2.13*10^{-18}$). This observation, while still showing that every spike matters, is inconsistent with recent models of saccade generation by the SC (Goossens and Van Opstal, 2006; Van Opstal and Goossens, 2008; Goossens and Van Opstal, 2012), which do not necessarily implement any kind of local versus remote interactions in how the SC influences saccade trajectories through individual spike effects.

Figure 4 The number of exogenous, movement-unrelated "visual" spikes to occur intra-saccadically
linearly adds to the executed movement's amplitude. (A) For every recorded neuron from Fig. 3 and every
microsaccade to occur near the visual burst interval (Fig. 1), we counted the number of spikes recorded from
the neuron that occurred intra-saccadically (between movement onset and movement peak velocity). We did
this for movements directed towards the RF location (Fig. 1B; Methods). We then plotted radial eye position
(aligned to zero in both the x- and y-axes) relative to saccade onset after categorizing the movements by the
number of intra-saccadic spikes. When no spikes were recorded during the eye movement, saccade
amplitudes were small (darkest curve), consistent with microsaccades during steady fixation. Adding "visual"
spikes in the SC map during the ongoing movement systematically increased movement amplitudes. Error
bars denote s.e.m. **(B)** To summarize the results in **A**, we plotted mean saccade amplitude against the number
of intra-saccadic "visual" spikes for movements directed towards the RF locations (faint red dots). There was
a linear increase in amplitude with each additional spike per recorded neuron (orange line representing the
best linear fit of the underlying raw data). For movements opposite the RF locations (faint green dots and green
line), there was no impact of intra-saccadic "visual" spikes on movement amplitudes. Error bars denote one
(**A**) and two (**B**) standard errors of the mean. Fig. S5 shows results for intra-saccadic spikes from more
eccentric neurons (>4.5 deg).

To further investigate the results of Fig. 4, we next explored more detailed temporal

interactions between SC visual bursts and saccade metric changes. Across all trials from all

neurons analyzed in Fig. 4, we measured the time of any given trial's visual burst peak

relative to either microsaccade onset (Fig. 5A), microsaccade peak velocity (Fig. 5B), or

microsaccade end (Fig. 5C), and we sorted the trials based on burst peak time relative to

microsaccade onset (i.e. the trial sorting in all panels in Fig. 5 was always based on the data

from panel A). We then plotted individual trial spike rasters with the bottom set of rasters

representing trials with the SC "visual" burst happening much earlier than microsaccade

onset and the top set being trials with the SC "visual" burst occurring after microsaccade

end. The rasters were plotted in gray in Fig. 5, except that during a putative visual burst

interval (30-100 ms from stimulus onset), we color-coded the rasters by the microsaccade

amplitude observed in the same trials (same color coding scheme as in Fig. 4A). The

marginal plot in Fig. 5D shows microsaccade amplitudes for the sorted trials (with a moving

average window of 30 trials in steps of 30 trials; Methods). We used this marginal plot to

estimate which sorted trials were associated with the beginning of microsaccade amplitude

increases (from the bottom of the raster and moving upward) and which trials were

associated with the end of the microsaccade amplitude increases. As can be seen,

whenever SC "visual" bursts occurred pre- and intra-saccadically, microsaccade amplitudes

were dramatically increased by two- to three-fold relative to baseline microsaccade

amplitudes (blue horizontal lines). For visual bursts after peak velocity (Fig. 5C), the effect

was diminished, consistent with efferent delays from SC activity to extraocular muscle

activation (Miyashita and Hikosaka, 1996; Munoz et al., 1996; Stanford et al., 1996; Gandhi

and Keller, 1999b; Katnani and Gandhi, 2012).

Therefore, at the time in which SC activity is to be read out by downstream neurons

to implement a saccadic eye movement (right before movement onset to right before

movement end, e.g. Miyashita and Hikosaka, 1996; Munoz et al., 1996; Stanford et al., 1996;

Gandhi and Keller, 1999b; Katnani and Gandhi, 2012), additional movement-unrelated SC

spiking activity is also read out and has a direct impact on eye movement metrics.

Figure 5 Exogenous, movement-unrelated SC spikes have the greatest impact on movement metrics when they occur peri-saccadically. (A) Individual trial spike rasters across all neurons ≤4.5 deg eccentricity and all movements towards RF locations. The spike rasters are sorted based on the time of the visual burst (peak firing rate after stimulus onset) relative to saccade onset (bottom left: trials with visual bursts earlier than microsaccades; top right: trials with visual bursts later than microsaccades). The spike rasters are plotted in gray except during the interval 30-100 ms after stimulus onset (our visual burst interval; Fig. 2) to highlight the relative timing of the visual burst to movement onset. Spikes in the visual burst interval are color-coded according to the observed movement amplitude on a given trial (legend on the left). As can be seen, microsaccades were enlarged when extra-foveal SC spiking (stimulus-driven visual bursts) occurred right before and during the microsaccades (see marginal plot of movement amplitudes in **D**). **(B)** Same as **A**, and with the same trial sorting, but with burst timing now aligned to movement peak velocity. **(C)** Same as **A**, **B**, and with the same trial sorting, but with burst timing now aligned to movement end. The biggest amplitude effects occurred when the exogenous "visual" spikes occurred pre- and intra-saccadically, but not post-saccadically. **(D)** Microsaccade amplitudes (30-trial moving average) on all sorted trials in **A**-**C**. Blue horizontal lines denote the range of trials for which there was a significant increase in movement amplitudes (Methods).

To obtain even more precise knowledge of the time needed for any injected "visual" spikes to start influencing saccade metrics, we then selected all individual trial spike rasters from Fig. 5A, and we counted the number of spikes occurring within any given 5 ms time bin relative to eye movement onset. We did this for all time bins between -100 ms and +100 ms from movement onset, and we also binned the movements by their amplitude ranges (Fig. 6A). The two smallest microsaccade amplitude bins reflected baseline movement amplitudes (see Fig. 3A), and they expectedly occurred when there was no "extra" spiking activity in the SC around their onset (Fig. 6A, two darkest reds). For all other amplitude bins, the larger movements were always associated with the presence of extra "visual" spikes on the SC map (more eccentric than the normal microsaccade amplitudes) occurring between -30 ms and +30 ms from saccade onset (Fig. 6A). Note how the timing of the effect was constant across amplitude bins, suggesting that it is the relative timing of extra "visual"

spikes and movement onset that mattered; the amplitude effect (that is, the different colored

curves) simply reflected the total number of spikes that occurred during the critical time

window of movement triggering (consistent with Fig. 4). Therefore, additional "visual" spikes

in the SC at a time consistent with saccade-related readout by downstream neurons

essentially "leak" into the saccade being generated. On the other hand, the pattern of Fig.

6A was not present for movements going opposite to the recorded neuron's RF's, for which,

if anything, there was a lower number of spikes happening during the peri-saccadic interval

(Fig. 6B). This suggests that it was easier to trigger microsaccades in one direction when

no activity was present in the opposite SC.

The strongest evidence that any "extra" spiking activity present on the SC map can

systematically alter the amplitude of the eye movements, irrespective of our experimental

manipulation of visual bursts, can be seen in Fig 6C. Here, we selected microsaccades

launched 100 ms after grating onset, temporally distant from the time of the earlier visual

bursts. This was to demonstrate that there is actually no need for a stimulus-driven visual

burst to be present for us to observe effects of extraneous spiking activity on triggered eye

movements. We counted the number of spikes occurring within the first 25 ms of the

selected movements. Because sustained firing rates were relatively low, which reduced the

total amount of data available to us for this analysis, we restricted the analysis to

microsaccades in which we could count either 2, 3, or 4 "extra" spikes emitted by a given

recorded neuron during the microsaccades. Even when the spikes were no longer strongly

associated with the stimulus-induced visual burst, their presence on the SC map at a site

more eccentric than microsaccade amplitudes was enough to modulate eye movement

amplitudes in a systematic manner, increasing the amplitude above baseline levels (blue

line) when more spikes were present. Such modulation was again not visible for movements

going in the opposite direction from the recorded neuron's RF's (Fig. 6D), again suggesting

 that there is a lower limit to how small microsaccades can become with opposite drive from

 the other SC.

Figure 6 Exogenous, movement-unrelated "visual" spikes affect movement metrics when they occur within approximately +/- 30 ms from movement onset, and they need not be part of a stimulus-driven visual burst to be effective. (A) For different microsaccade amplitude ranges from Fig. 5 (color-coded curves), we counted the number of exogenous spikes occurring from a recorded extra-foveal SC neuron within any given 5 ms time bin around movement onset (range of times tested: -100 ms to +100 ms from movement onset). The lowest two microsaccade amplitude ranges (0.1-0.2 and 0.2-0.3 deg) reflected baseline amplitudes during steady-state fixation (e.g. Fig. 3), and they were not correlated with additional extra-foveal spiking activity around their onset (two darkest red curves). For all other larger microsaccades, they were clearly associated with precise timing of extra-foveal "visual" spikes occurring within approximately +/- 30 ms from movement onset, regardless of movement size. **(B)** Same as **A** but for movements opposite the recorded neuron's RF locations. There were fewer spikes during the peri-saccadic interval, suggesting that it was easier to trigger eye movements when there was no activity present in the opposite SC. **(C)** Relation between the number of intra-saccadic spikes and saccade amplitudes for eye movements triggered 100 ms after grating onset, temporally distant from the time of the visual burst. Each dot represents a saccade amplitude with the associated number of spikes detected within 25 ms after movement onset. Solid squares represent the averages. The presence of intra-saccadic spikes on the SC map was enough to modulate movement amplitudes even outside of our usual visual burst interval of earlier analyses. Blue indicates baseline microsaccade amplitude. **(D)** Same as **C** but for movements opposite the recorded RF location.

Overall, these results demonstrate that there is a tight time window around saccade

onset (Fig. 6) in which any movement-unrelated spikes in sites other than the saccade goal

representation can induce a systematic variation in the motor program.

Peri-saccadic single, double, or triple pulse electrical microstimulation alters eye movement

metrics

Our results so far demonstrate that as little as one single extra action potential by each

visually-activated neuron was sufficient to alter ongoing microsaccades (Figs. 4-6). To

establish a causal role for such exogenous SC spiking activity, we next replaced visual

stimulus onsets with single, double, or triple electrical microstimulation pulses that were

strategically timed to occur peri-saccadically. We did this in a third monkey (M) for which we

know that amplitude effects like those in Fig. 3 still took place (Buonocore et al., 2019;

Malevich et al., 2020a). We performed an experiment analogous to peri-saccadic perceptual

experiments in which brief visual stimuli occur near the time of saccade onset (e.g. Honda,

1989; Morrone et al., 1997; Ross et al., 1997; Lappe et al., 2000; Krekelberg et al., 2003;

Buonocore et al., 2017a; Grujic et al., 2018), but we replaced visual stimuli with "electrical"

stimuli. The monkey generated an instructed visually-guided saccade, and we introduced

electrical microstimulation pulses at random times (Methods). To simulate the effects of

Figs. 4-6 in this "electrical" spiking paradigm, we had the monkey generate visually-guided

saccades to a parafoveal location (2.2–5.5 deg across sessions), and the microstimulating

electrode was always at a site representing a larger eye movement vector along the same

direction (Methods). This makes the experiment similar in principle to the conditions of Figs.

4-6 above: additional spiking activity in the same SC but at a location more eccentric than

that of the planned eye movement. Prior to microstimulation, we confirmed that the visually-

guided saccade target location was outside of the visual and movement RF's at the

microstimulated SC site. That is, the microstimulated SC site was not directly involved in

generating the visually-guided saccade; it was just injecting extraneous spikes analogous to

the "visual" spikes in Figs. 2-6. More importantly, we confirmed that triple pulse

microstimulation, the strongest of our electrical manipulations, did not itself evoke any

systematic saccadic eye movements (even microsaccades) during a control fixation

condition (Fig. S6). We then proceeded to the main experiment.

Figure 7 shows results from an example microstimulation session. In Fig. 7A, we

plotted the trajectories of saccades to a location approximately 1.9 deg to the right and 3

deg above fixation without coincident microstimulation. In Fig. 7B (red traces), the monkey

generated the same visually-guided saccades, but these saccades were now paired with 3

microstimulation pulses (inter-pulse interval: 3.3 ms; 30 μA current amplitude per pulse;

Methods) starting within +/-10 ms of movement onset. The saccades were now elongated

relative to the baseline saccades (also shown in faint color in Fig. 7B to facilitate comparison

between the two conditions), consistent with the results of Figs. 4-6 above. Thus, the

saccades were pulled towards the site in which additional SC spiking was injected, such that

the overall displacement of the eye during the saccades was slightly longer. Such elongation

was further clarified when we plotted radial eye velocity. Injecting 3 microstimulation pulses

at saccade onset introduced a secondary velocity transient, shortly afterwards, to elongate

the movements relative to baseline (Fig. 7C, D). Therefore, exogenous movement-unrelated

"electrical" activity had a similar effect to the "visual" activity described in our earlier

experiments.

Across sessions, we tested single, double, or triple pulse microstimulation in peri-

saccadic intervals. To summarize all results, we first defined a baseline saccadic

displacement amplitude as the overall displacement caused by saccades in which no

microstimulation pulse occurred near saccade onset (Methods). We then obtained a time

course of saccadic displacement as a function of peri-saccadic microstimulation pulses by

measuring, for any time window of pulse times relative to saccade onset, the saccadic

displacement observed and normalizing it to the baseline displacement. This gave us a

percentage change of saccadic displacements as a function of microstimulation pulse times

relative to saccade onset, and it also allowed us to summarize all sessions together when

they individually each had slightly different SC sites being microstimulated. As shown in Fig.

8, even single and double pulse microstimulation affected ongoing eye movement metrics.

The increase in eye displacement started about 25 ms before saccade onset, and it lasted

for 25 ms thereafter, consistent with the timing described in Fig. 6. Moreover, consistent with

Figs. 4-6, the amplitude changes were sensitive to the number of injected microstimulation

pulses. With one injected "electrical" spike, the alteration in the displacement was modest

(~6%) and loosely centered around the saccade onset. With two and three injected spikes,

the alteration was significantly stronger (~8%) and in a more precise time window around

the saccade. Overall, the trajectories were clearly pulled towards the RF locations of the

stimulated sites.

Figure 7 Replacing "visual" with "electrical" exogenous spikes similarly alters ongoing eye movement metrics. Example session from the experiment of Fig. 1C. A monkey generated a saccade to a target at 1.9 deg and 3 deg horizontally and vertically, respectively. In a control manipulation (**A**), 3 pulses of electrical microstimulation were injected into the SC 40-50 ms before saccade onset, and at a site more eccentric than the saccade target (but along a similar vector direction; Methods). In peri-saccadic microstimulation (**B**), the same 3 pulses were applied to the same SC site, but now within +/- 10 ms from saccade onset. Each curve in **A**, **B** shows the total saccadic displacement of the eye on individual trials (from 50 ms before saccade onset to 100 ms afterwards), and the curves were all aligned together at the origin (eye position at saccade onset). The gray curves in **B** are the same as those in **A**, and they demonstrate how the red curves in **B** were associated with longer saccadic eye displacements than in control. **(C, D)** Radial eye velocity traces for the same eye movements in **A**, **B** (but plotting data only from saccade onset this time). Faint curves show individual trials, and thick curves denote the across-trial averages. Electrical pulses at saccade onset clearly resulted in a subsequent prolongation of ongoing saccades, evident as a secondary velocity pulse approximately 30 ms later (**D**). Figure S6 shows that 3 pulses alone (without an ongoing plan) were not sufficient to evoke a saccadic eye movement.

Figure 8 Peri-saccadic electrical microstimulation with 1, 2, or 3 spikes alters saccade metrics. Time courses of saccadic displacement as a function of peri-saccadic microstimulation. We measured the total saccadic displacement (0-100 ms from saccade onset) as a function of the time of electrical microstimulation that was applied peri-saccadically. Amplitude displacements were normalized to the measured displacement when microstimulation was far from saccade onset (i.e. averaged across ±25 ms centered on 50 ms before saccade onset). Each panel shows results with 1 (**A**), 2 (**B**), or 3 (**C**) peri-saccadic electrical microstimulation pulses applied to the SC at a site more eccentric than the saccade target location (but along a similar vector direction). Single pulse microstimulation had a modest impact on the eye displacement, and it was loosely centered around saccade onset. With double and triple pulse microstimulation variants, the alteration was significantly stronger and clearly centered in a precise time window around the saccade. This time window was highly consistent with the time window of Fig. 6A with "visual" injected spikes that altered microsaccade metrics. Error bars denote 95% confidence intervals.

Discussion

We experimentally injected movement-unrelated spiking activity into the SC map at the time of saccade generation. We found that such activity was sufficient to significantly alter the metrics of the generated saccadic eye movement, suggesting an instantaneous readout of the entire SC map for implementing any individual movement.

Our results reveal a component of motor variability that we believe has been previously unaccounted for, namely, the fact that ever-present spiking activity in the entire SC map (whether due to sustained firing rates for a stimulus presented in the RF, like in our first experiment, or otherwise) can "leak" into the readout performed by downstream motor structures when executing a movement. In fact, our results from Fig. 6C showed that any intra-saccadic spikes on the SC map (far from the location of the motor burst) were sufficient to modulate microsaccade metrics, meaning that there was no need for a stimulus onset or even a stimulus-driven visual burst, like in Figs. 3-6, to give rise to our observations. Indeed, saccades during natural viewing show an immense amount of kinematic variability when compared to simplified laboratory tasks with only a single saccade target (Berg et al., 2009). In such natural viewing, natural images with plenty of low spatial frequency image power are expected to strongly activate a large number of SC neurons, which prefer low spatial frequencies, around the time of saccades (Chen et al., 2018; Khademi et al., 2020).

From an ecological perspective, our results demonstrate a remarkable flexibility of the oculomotor system during eye movement generation. Historically, saccades were thought to be controlled by an open-loop control system due to their apparent ballistic nature. However, other evidence, including our current results, clearly showed that individual saccades are actually malleable brain processes. In our case, we experimentally tried to generate a movement-unrelated "visual" or "electrical" burst of activity that precisely coincided with the time of saccade triggering. We uncovered an instantaneous readout of the entire SC map that includes all the activity related to the ongoing motor program as well

as the "extra" activity. In real life, this extra activity might happen due to external sensory

stimulation, such as the presence of a new object in the visual scene.

Such integration of sensory signals into the motor plan was also not merely a "loose"

leakage phenomenon; rather, it exhibited a lawful additive process between the "visual"

spikes injected into the SC population and the altered microsaccade amplitudes (Figs. 4, 6).

The more "visual" spikes that occurred intra-saccadically, the larger the microsaccades

became, following a linear relationship. We discount the possibility that this effect was due

to movement-related bursts per se, because we ensured that the neurons that we recorded

from were not exhibiting movement bursts for the ranges of eye movements that we

analyzed (Fig. S1). We suggest that this additive vector mechanism might underlie many of

the effects commonly seen in experimental psychophysics, in which saccade kinematics are

systematically altered by the presence of sudden irrelevant visual information available as

close as 40 ms to movement onset (Edelman and Xu, 2009; Buonocore and McIntosh, 2012;

Guillaume, 2012; Buonocore et al., 2016; Buonocore et al., 2017b; Malevich et al., 2020a).

Our hypothesis is that these modulations are a behavioral manifestation of the

instantaneous readout of the activity on the SC map, as we also previously hypothesized

(Hafed and Ignashchenkova, 2013; Buonocore et al., 2017b). Moreover, similar specification

mechanisms can be seen in the instantaneous alteration of eye velocity during smooth

pursuit when small flashes are presented (Buonocore et al., 2019), and even with ocular

position drift during fixation (Malevich et al., 2020b). These observations extend the

mechanisms uncovered in our study to the pursuit system and beyond, and they also relate

to sequential activation of SC neurons during curved saccades associated with planning

sequences of movements (Port and Wurtz, 2003).

Most intriguingly, our results motivate similar neurophysiological studies on sensory-

motor integration in other oculomotor structures. For example, our own ongoing experiments

in the lower oculomotor brainstem, at the very final stage for saccade control (Keller, 1974;

Büttner-Ennever et al., 1988; Gandhi and Keller, 1999a; Missal and Keller, 2002), are revealing highly thought provoking visual pattern analysis capabilities of intrinsically motor neurons (Buonocore et al., 2020). These and other experiments will, in the future, clarify the mechanisms behind multiplexing of visual and motor processing in general, across other subcortical areas, like pulvinar, and also cortical areas, like FEF and LIP. Moreover, these sensory-motor integration processes can have direct repercussions on commonly used behavioral paradigms in which microsaccades and saccades happen around the time of attentional cues/probes and can alter performance (Hafed, 2013; Hafed et al., 2015; Tian et al., 2016; Buonocore et al., 2017b).

Our electrical microstimulation manipulations are also interesting because they add to what we believe is already strong causal evidence from our visual burst manipulations. While previous studies have demonstrated the possibility to terminate the current saccadic motor plan in favor of a new one by electrical microstimulation of the SC (Gandhi and Keller, 1999b), our experiments rather focused on the flexibility of "leaking" extra subtle activity into the ongoing oculomotor plan. As such, we used low currents, and we microstimulated peri-saccadically at more eccentric locations than the generated saccades, unlike in the earlier studies. Our experiments therefore complement these earlier reports performing SC electrical microstimulation.

Consistent with the above sentiment, our study generally illuminates emerging and classic models of the role of the SC in saccade control. In a recent model by Goossens and Van Opstal (2006), it was suggested that every SC spike during a motor burst contributes a mini-vector of eye movement tendency, such that the aggregate sum of movement tendencies comprises the overall trajectory. Our results are consistent with this model, and related ones also invoking a role of SC activity levels in instantaneous trajectory control (Waitzman et al., 1991; Ivan et al., 2018), in the sense that we did observe linear contributions of additional SC spikes on eye movement metrics (Figs. 4, 6). However, as

stated in Introduction, our results add to this model the notion that there need not be a

"classifier" identifying particular SC spikes as being the movement-related spikes of the

current movement and other spikes as being irrelevant. More importantly, we found

diminishing returns of relative eccentricity between the "extra" spikes and the current motor

burst (Figs. 3, S3, 4, S5). According to their model, the more eccentric spikes that we

introduced from more eccentric neurons should have each contributed "mini-vectors" that

were actually larger than the "mini-vectors" contributed by the less eccentric spikes from the

less eccentric neurons. So, if anything, we should have expected larger effects for the more

eccentric neurons. This was clearly not the case. Therefore, this model needs to consider

local and remote interactions more explicitly. The model also needs to consider other factors

like input from other areas. Indeed, Peel et al. (2019) reported that the SC generates fewer

saccade-related spikes during FEF inactivation, even for matched saccade amplitudes.

Similarly, we recently found that microsaccades without visual guidance can be associated

with fewer active SC neurons than similarly-sized microsaccades with visual guidance,

because of so-called visually-dependent saccade-related neurons (Willeke et al., 2019).

Finally, SC motor bursts themselves are very different for saccades directed to upper versus

lower visual field locations, without an apparent difference in saccade kinematics (Hafed

and Chen, 2016). All of these observations suggest that further research on the SC

contributions to saccade trajectory control is strongly needed.

In all, we believe that our results expose highly plausible neural mechanisms

associated with robust behavioral effects on saccades accompanied by nearby visual

flashes in a variety of paradigms, and they also motivate revisiting a classic

neurophysiological problem, the role of the SC in saccade control, from the perspective of

visual-motor multiplexing within individual brain circuits, and even individual neurons

themselves.

Acknowledgements

We were funded by the Deutsche Forschungsgemeinschaft (DFG) through the Research Unit: FOR1847 (project: HA6749/2-1). We were also funded by the Werner Reichardt Centre for Integrative Neuroscience (CIN; DFG EXC307) and the Hertie Institute for Clinical Brain Research.

Declaration of interests

The authors declare no competing interests.

Author contributions

AB, MPB, ZMH performed the microstimulation experiments. AB, ZMH analyzed the recording data. AB, MPB, ZMH analyzed the microstimulation data. AB, MPB, ZMH wrote and edited the manuscript.

Methods

Animal preparation

We collected data from three (M, N, and P) adult, male rhesus monkeys (Macaca mulatta) that were 6-8 years of age and weighed 6-8 kg. The experiments were approved (licenses: CIN3/13; CIN4/19G) by ethics committees at the regional governmental offices of the city of Tuebingen and were in accordance with European Union guidelines on animal research and the associated implementations of these guidelines in German law. The monkeys were prepared using standard surgical procedures necessary for behavioral training and intracranial recordings. In short, monkeys N and P had a chamber centered on the midline and aiming at the superior colliculus (SC) with an angle of 35 and 38 degrees posterior of vertical in the sagittal plane, respectively. Monkey M was implanted with a similar chamber having an angle of 38 degrees posterior of vertical in the sagittal plane. The details of the surgical procedures were described in previous reports (monkeys N and P: Chen and Hafed, 2013; Monkey M: Willeke et al., 2019). To record eye movements with high temporal and spatial precision, all monkeys were also implanted with a scleral search coil. This allowed eye tracking using the magnetic induction technique (Fuchs and Robinson, 1966; Judge et al., 1980). Monkeys N and P were each implanted in the right and left eye, respectively. Monkey M was implanted in both eyes.

Experimental control system and monkey setup

We used a custom-built real-time experimental control system that drove stimulus presentation and ensured monkey behavioral monitoring and reward delivery. It also controlled an electrical microstimulation device (Stimpulse; FHC, Inc.) to trigger microstimulation. The details of the system are reported in recent publications (Chen and Hafed, 2013; Tian et al., 2016).

During the testing sessions, the animals were head fixed and seated in a standard primate chair placed at a distance of 74 cm from a CRT monitor. The eye height was aligned with the center of the screen. The room was completely dark with the only light source being the monitor. All stimuli were presented over a uniform gray background (Monkey M: 29.7 Cd/m^2; Monkeys N and P: 21 Cd/m^2). In all the experiments, the fixation spot consisted in a small square made of 3 by 3 pixels (about 8.5 by 8.5 min arc) colored in white (Monkey M: 86 Cd/m^2; Monkeys N and P: 72 Cd/m^2). The central pixel had the same color as the background.

Behavioral tasks and electrophysiology

Experiment 1: injecting visual spikes at the time of saccade generation

We performed a novel analysis of SC data reported on earlier; our behavioral task is therefore described in detail in (Chen et al., 2015). Briefly, we used a fixation paradigm during which we introduced a peripheral transient visual event at random intervals (see Fig. 1A). Each trial started with a white fixation spot presented at the center of the display over a uniform gray background. The monkey was required to align its gaze with the fixation spot. Because fixation is an active process, this steady-state fixation paradigm allowed us to have a scenario in which microsaccades were periodically generated (Hafed and Ignashchenkova, 2013). After a random interval, we presented a stimulus consisting of a vertical sine wave grating of 2.2 cycles/deg spatial frequency and filling the visual response field (RF) of the recorded neuron. The stimulus onset allowed experimentally injecting visual spikes into the SC at retinotopic locations dissociated from the neurons involved in microsaccade generation (Hafed et al., 2009; Willeke et al., 2019). Therefore, we could investigate the influence of such injected spiking activity if it happened to occur in the middle of an ongoing microsaccade (see Results). We varied the contrast of the grating across trials in order to vary the amount of injected SC spiking activity around the time of microsaccade

generation. Specifically, grating contrast could be one of 5%, 10%, 20%, 40%, or 80% (Chen

et al., 2015). For the current study, we only analyzed trials with the highest 3 contrasts. We

related microsaccade kinematics to injected "visual" spiking activity. Overall, we analyzed

84 SC visual (44) and visual-motor (40) neurons in two monkeys (N and P).

Experiment 2: injecting electrical spikes at the time of saccade generation

In monkey M, we performed peri-saccadic electrical microstimulation of the SC during a

visually-guided saccade task (Fig. 1C). That is, rather than peri-saccadic spiking activity

induced in the SC by visual onsets, we electrically microstimulated the SC around the time

of saccade generation.

Before each session, we first identified the direction and amplitude of the visual field

location associated with the electrode position in the SC topographic map. We did so by

mapping the visual and movement RF's encountered by the electrode using a standard

delayed visually-guided saccade task (Chen et al., 2015). In a second step, we confirmed

the electrode location by assessing the size and direction of the saccade vector evoked

through electrical microstimulation (our electrodes had relatively low impedances of <1

MOhm, and they were placed in the intermediate layers of the SC). We programmed the

microstimulation parameters of a given test on the Stimpulse (FHC, Inc.) system's user

interface, and we then triggered the microstimulation train using our real-time experiment

control system. While the monkey was maintaining fixation on a continuously visible fixation

spot, we microstimulated the SC with symmetric biphasic current pulses having an amplitude

of 30-45 μA and inter-pulse interval of 3.3 ms (i.e. pulse train frequency of 300Hz). The pulse

train duration was relatively short (70-150 ms), and each individual pulse consisted of a 150

μs cathodal phase followed immediately by a 150 μs anodal phase. Such electrical

microstimulation reliably elicited short-latency eye movements of a particular size and

direction (Fig. S6A).

Once the RF's of the neurons at the SC site were established using both recording and microstimulation, as per the above procedures, we started the main experiment, which consisted of a simple visually-guided saccade task. The monkey fixated, and the fixation spot was then removed with a simultaneous presentation of a visual target in the periphery (white circle of 0.45 deg radius). We chose the location of the visual target such that it was not overlapping with the RF's of the neurons that we microstimulated earlier. Specifically, the location of the target stimulus was chosen to be about halfway between the fixation spot and the RF locations being microstimulated on the SC map. Thus, when we injected "electrical" stimuli into the SC, we were injecting activity at a site that was spatially dissociated from the saccade target location; this is conceptually the same as the situation in the "visual" spike experiments above (Fig. 1A, B). In different blocks, we injected either 1, 2, or 3 electrical pulses into the SC peri-saccadically. Pulse amplitude currents were similar to above, and inter-pulse intervals (for double and triple pulse microstimulation) were 3.3 ms. To achieve peri-saccadic microstimulation, as in peri-saccadic visual presentation experiments (e.g. Buonocore et al., 2017a; Grujic et al., 2018), we randomized the time of microstimulation from trial to trial. Specifically, each trial started with an initial fixation interval of approximately 300-700 ms. When the saccade target appeared, we introduced a random delay between approximately 70 ms and 170 ms (uniform distribution) before injecting electrical spikes. Given typical visually-guided saccade reaction times, this resulted in electrical microstimulation pulses appearing either well before, during, or well after saccade onset. This, in turn, allowed us to assess the effects of peri-saccadic single, double, or triple pulse microstimulation in the SC (Fig. 6 in Results). In a separate control block, we confirmed that 3 pulses electrically injected into the SC (with the same parameters as in the main experiment) during simple fixation did not elicit any systematic eye movements (Fig. S6B, C). Thus, whatever impacts that we had on visually-guided saccade metrics were not due to an explicitly evoked saccade caused by electrical microstimulation.

Data analysis

All analyses were performed with custom scripts in Matlab (MathWorks, Inc.). Most of the analyses involving grouping the eye movement data into groups of movements going either towards or opposite a recorded neuron's RF. To make this classification, we first calculated the angle of the RF relative to the fixation spot. Then, all eye movements with an angle ±90 degrees around the RF direction were classified as being "towards" the RF. All remaining eye movements were classified as being directed "opposite" to the RF. Movement angles were defined as the arctangent subtended by the horizontal and vertical component between movement onset and end. RF angles were defined as the arctangent subtended by the horizontal and vertical coordinates of the RF locations relative to the fixation spot.

To analyze peak firing rates "without saccades" (Figs. 2, 3B, S2, S3B), we selected all trials in which there were no microsaccades btween -100 ms and 200 ms relative to stimulus onset. We then averaged all the firing rates across trials, and we determined the peak firing rate for each neuron from the across-trial average curve. For the peak firing rate "with saccades", we took all the trials in which a microsaccade was either starting or ending during the so-called visual burst interval, which we defined to be the interval 30-100 ms after stimulus onset. Paired-sample t-tests were performed to test the influence of saccades on the peak visual burst with an α level of 0.05 unless otherwise stated (Figs. 2, S2).

To summarize the time courses of microsaccade amplitudes after stimulus onset (Figs. 3A, S3A), we selected the first saccade of each trial that was triggered within the interval from -100 ms to +150 ms relative to stimulus onset. All the microsaccade amplitudes were then pooled together across monkeys and sessions. The microsaccade amplitude time course was obtained by filtering the data with a running average window of 50 ms with a step size of 10 ms. To statistically test the effect of grating contrasts on these time courses, we performed a one-way ANOVA on saccade amplitudes for all saccades occurring between 50 ms and 100 ms after stimulus onset. To compare the effect of grating contrast on SC

visual bursts (Figs. 3B, S3B), for each neuron, we normalized the firing rate based on the

maximum firing rate elicited by the strongest contrast. Subsequently, we calculated the

mean firing rate of the population and the 95% confidence interval for the different contrast

levels. Both the amplitude and the firing rate analyses focused on the three highest contrasts

because the first two contrast levels did not have a visible impact on eye movement

behavior.

For some analyses (Figs. 4-6, S5), we explored the relationship between the number

of spikes emitted by a recorded neuron and saccade amplitude. This allowed us to directly

investigate the effect of each single additional spike per recorded neuron in the SC map on

an ongoing saccade, irrespective of other variables. We selected all the trials in which an

eye movement was performed in the direction of the grating soon after its presentation. The

analysis was restricted to the three highest contrasts, since they had a clear effect on the

eye movement behavior (Fig.3A). For each selected saccade, we counted the number of

spikes happening from a given recorded neuron between movement onset movement peak

velocity. For Figs. 4A, S5A, we calculated the "radial eye position" from saccade start as the

Euclidian distance of any eye position sample (i.e. at any millisecond) recorded during an

eye movement relative to the eye position at movement onset (see for similar procedures:

Hafed et al., 2009; Buonocore et al., 2017b). To make statistical inferences on the effects

of the number of spikes on saccade amplitude (Figs. 4B, S5B), we proceeded by fitting a

generalized linear model to the raw data with equation: $y = \beta_0 + \beta_1 \cdot x$ where 'x' was our

predictor variable, the number of spikes, and 'y' was the predicted amplitude. The

parameters fitted were: β_0 the intercept, β_1 the slope. We imposed a cutoff of at least 15

trials for each level of the predictor, leading to exclusion of spike counts bigger than six.

To study the time window of influence of each added spike on saccade amplitudes

(Figs. 5, 6), we generated raster plots from all trials of all sessions by aligning each spike

raster trace to the saccade occurring on the same trial (either saccade start, peak velocity,

or end). The saccades were chosen as those that happened after stimulus onset, and the

alignment was based on the time of peak visual burst after stimulus onset on a given trial

relative to the time of the movement. We selected data from the three highest stimulus

contrasts and also with eye movements directed towards the RF, since the modulation in

behavior was most pronounced in these cases (e.g. Fig. 4). To identify the point at which

the amplitude diverged from a baseline level in Fig. 5D, we first sorted all the amplitudes

based on burst time relative to saccade onset (Fig. 5A). Then, we made bins of 30 trials

each from which we derived the mean amplitude values (Fig. 5D). We also tested the

amplitudes of each bin against the first one (baseline amplitude) to determine when the

amplitude increase was significant. To do so, we performed two-sample independent t-tests

between each pair, and we adjusted the alpha level with Bonferroni correction. We chose

as an index for a significant increase in amplitude the point at which three consecutive bins

were significantly different from the baseline (first horizontal blue line in the trial sorting of

Fig. 5D). The next three consecutive bins that did not differ anymore from the baseline

indicated that the amplitude increase was not significant anymore (second horizontal blue

line in the trial sorting of Fig. 5D).

For the microstimulation experiment, we first ensured that the monkey was

maintaining stable fixation in an interval between -50 ms to 300 ms around target onset and

that the eyes were within a fixation window of one degree radius. We then selected as

response saccades all the eye movements made 75 ms to 400 ms after target appearance

and with an amplitude larger than halfway between fixation and target location. Such strict

criteria were used because the target could have been placed in relatively rostral regions,

where microsaccades happen frequently and might therefore interfere with the goal directed

saccade. Subsequently, on the selected saccade, we calculated the "eye displacement" as

the radial position of the eyes from saccade start to 100 ms thereafter. This value was then

normalized by the average displacement obtained with microstimulation in a time window of

±25 ms centered on 50 ms before movement onset (i.e. well before the onset of our

behavioral effect) to allow the pooling of different sessions that had different target

eccentricities. The eye displacement curve (Fig. 8) was calculated by binning the eye

displacement values in a time interval of 50 ms before to 100 ms after saccade onset, with

a bin size of 50 ms and a step of 1 ms. This procedure was repeated independently for the

1, 2, or 3 spike blocks.

References

Alexander GE, Crutcher MD (1990) Neural representations of the target (goal) of visually guided arm movements in three motor areas of the monkey. J Neurophysiol 64:164-178.

Berg DJ, Boehnke SE, Marino RA, Munoz DP, Itti L (2009) Free viewing of dynamic stimuli by humans and monkeys. J Vis 9:19.11-15.

Breveglieri R, Bosco A, Galletti C, Passarelli L, Fattori P (2016) Neural activity in the medial parietal area V6A while grasping with or without visual feedback. Sci Rep 6:28893.

Bruce CJ, Goldberg ME (1985) Primate frontal eye fields. I. Single neurons discharging before saccades. Journal of Neurophysiology 53:603-635.

Buonocore A, McIntosh RD (2012) Modulation of saccadic inhibition by distractor size and location. Vision Research 69:32-41.

Buonocore A, McIntosh RD, Melcher D (2016) Beyond the point of no return: effects of visual distractors on saccade amplitude and velocity. Journal of Neurophysiology 115:752-762.

Buonocore A, Fracasso A, Melcher D (2017a) Pre-saccadic perception: Separate time courses for enhancement and spatial pooling at the saccade target. PLoS ONE 12:e0178902-0178923.

Buonocore A, Skinner J, Hafed ZM (2019) Eye Position Error Influence over "Open-Loop" Smooth Pursuit Initiation. The Journal of Neuroscience 39:2709-2721.

Buonocore A, Baumann MP, Hafed ZM (2020) Visual pattern analysis by motor neurons (Abstract). Computational and Systems Neuroscience (Cosyne) 147.

Buonocore A, Chen C-Y, Tian X, Idrees S, Münch TA, Hafed ZM (2017b) Alteration of the microsaccadic velocity-amplitude main sequence relationship after visual transients: implications for models of saccade control. Journal of Neurophysiology 117:1894-1910.

Büttner-Ennever JA, Cohen B, Pause M, Fries W (1988) Raphe nucleus of the pons containing omnipause neurons of the oculomotor system in the monkey, and its homologue in man. Journal of Comparative Neurology 267:307-321.

Chen CY, Hafed ZM (2013) Postmicrosaccadic Enhancement of Slow Eye Movements. Journal of Neuroscience 33:5375-5386.

Chen CY, Ignashchenkova A, Thier P, Hafed ZM (2015) Neuronal Response Gain Enhancement prior to Microsaccades. Current Biology 25:2065-2074.

Chen CY, Hoffmann KP, Distler C, Hafed ZM (2019) The Foveal Visual Representation of the Primate Superior Colliculus. Curr Biol 29:2109-2119 e2107.

Chen CY, Sonnenberg L, Weller S, Witschel T, Hafed ZM (2018) Spatial frequency sensitivity in macaque midbrain. Nature Communications 9:2852.

Cynader M, Berman N (1972) Receptive-field organization of monkey superior colliculus. J Neurophysiol 35:187-201.

Edelman JA, Goldberg ME (2001) Dependence of Saccade-Related Activity in the Primate Superior Colliculus on Visual Target Presence. Journal of Neurophysiology 86:676-691.

Edelman JA, Xu KZ (2009) Inhibition of Voluntary Saccadic Eye Movement Commands by Abrupt Visual Onsets. Journal of Neurophysiology 101:1222-1234.

Fuchs AF, Robinson DA (1966) A method for measuring horizontal and vertical eye movement chronically in the monkey. J Appl Physiol 21:1068-1070.

Gandhi NJ, Keller EL (1999a) Activity of the brain stem omnipause neurons during saccades perturbed by stimulation of the primate superior colliculus. Journal of Neurophysiology 82:3254-3267.

Gandhi NJ, Keller EL (1999b) Comparison of Saccades Perturbed by Stimulation of the
Rostral Superior Colliculus, the Caudal Superior Colliculus, and the Omnipause
Neuron Region. Journal of Neurophysiology 82:3236-3253.
Goldberg ME, Wurtz RH (1972a) Activity of superior colliculus in behaving monkey. I.
Visual receptive fields of single neurons. J Neurophysiol 35:542-559.
Goldberg ME, Wurtz RH (1972b) Activity of superior colliculus in behaving monkey. II.
Effect of attention on neuronal responses. J Neurophysiol 35:560-574.
Goossens HH, Van Opstal AJ (2006) Dynamic ensemble coding of saccades in the
monkey superior colliculus. J Neurophysiol 95:2326-2341.
Goossens HHLM, Van Opstal AJ (2012) Optimal Control of Saccades by Spatial-Temporal
Activity Patterns in the Monkey Superior Colliculus. PLoS Computational Biology
8:e1002508.
Grujic N, Brehm N, Gloge C, Zhuo W, Hafed ZM (2018) Perisaccadic perceptual
mislocalization is different for upward saccades. J Neurophysiol 120:3198-3216.
Guillaume A (2012) Saccadic inhibition is accompanied by large and complex amplitude
modulations when induced by visual backward masking. Journal of Vision 12:5-5.
Hafed ZM (2011) Mechanisms for generating and compensating for the smallest possible
saccades. European Journal of Neuroscience 33:2101-2113.
Hafed ZM (2013) Alteration of Visual Perception prior to Microsaccades. Neuron 77:775-
786.
Hafed ZM, Krauzlis RJ (2012a) Similarity of superior colliculus involvement in
microsaccade and saccade generation. Journal of Neurophysiology 107:1904-1916.
Hafed ZM, Krauzlis RJ (2012b) Similarity of superior colliculus involvement in
microsaccade and saccade generation. J Neurophysiol 107:1904-1916.
Hafed ZM, Ignashchenkova A (2013) On the Dissociation between Microsaccade Rate and
Direction after Peripheral Cues: Microsaccadic Inhibition Revisited. Journal of
Neuroscience 33:16220-16235.
Hafed ZM, Chen CY (2016) Sharper, Stronger, Faster Upper Visual Field Representation
in Primate Superior Colliculus. Curr Biol 26:1647-1658.
Hafed ZM, Goffart L, Krauzlis RJ (2009) A Neural Mechanism for Microsaccade
Generation in the Primate Superior Colliculus. Science (New York, NY) 323:940-
943.
Hafed ZM, Chen C-Y, Tian X (2015) Vision, Perception, and Attention through the Lens of
Microsaccades: Mechanisms and Implications. Frontiers in Systems Neuroscience
9:779.
Honda H (1989) Perceptual localization of visual stimuli flashed during saccades.
Perception & Psychophysics 45:162-174.
Ivan S, Uday KJ, Neeraj JG (2018) Instantaneous Midbrain Control of Saccade Velocity.
The Journal of Neuroscience 38:10156-10167.
Judge SJ, Richmond BJ, Chu FC (1980) Implantation of magnetic search coils for
measurement of eye position: an improved method. Vision Res 20:535-538.
Katnani HA, Gandhi NJ (2012) The relative impact of microstimulation parameters on
movement generation. Journal of Neurophysiology 108:528-538.
Keller EL (1974) Participation of Medial Pontine Reticular-Formation in Eye-Movement
Generation in Monkey. Journal of Neurophysiology 37:316-332.
Khademi F, Chen CY, Hafed ZM (2020) Visual feature tuning of superior colliculus neural
reafferent responses after fixational microsaccades. J Neurophysiol 123:2136-2153.
Krekelberg B, Kubischik M, Hoffmann KP, Bremmer F (2003) Neural correlates of visual
localization and perisaccadic mislocalization. Neuron 37:537-545.
Lappe M, Awater H, Krekelberg B (2000) Postsaccadic visual references generate
presaccadic compression of space. Nature 403:892-895.

Li X, Basso MA (2008) Preparing to move increases the sensitivity of superior colliculus
neurons. J Neurosci 28:4561-4577.
Malevich T, Buonocore A, Hafed ZM (2020a) Dependence of the stimulus-driven
microsaccade rate signature on visual stimulus polarity.
bioRxiv:2020.2005.2023.112417.
Malevich T, Buonocore A, Hafed ZM (2020b) Rapid stimulus-driven modulation of slow
ocular position drifts. bioRxiv:2020.2004.2002.022335.
Mays LE, Sparks DL (1980) Dissociation of visual and saccade-related responses in
superior colliculus neurons. J Neurophysiol 43:207-232.
Mazzoni P, Bracewell RM, Barash S, Andersen RA (1996) Motor intention activity in the
macaque's lateral intraparietal area. I. Dissociation of motor plan from sensory
memory. J Neurophysiol 76:1439-1456.
Missal M, Keller EL (2002) Common inhibitory mechanism for saccades and smooth-
pursuit eye movements. Journal of Neurophysiology.
Miyashita N, Hikosaka O (1996) Minimal synaptic delay in the saccadic output pathway of
the superior colliculus studied in awake monkey. Exp Brain Res 112:187-196.
Mohler CW, Wurtz RH (1976) Organization of monkey superior colliculus: intermediate
layer cells discharging before eye movements. J Neurophysiol 39:722-744.
Morrone MC, Ross J, Burr DC (1997) Apparent position of visual targets during real and
simulated saccadic eye movements. Journal of Neuroscience 17:7941-7953.
Munoz DP, Wurtz RH (1995) Saccade-related activity in monkey superior colliculus. I.
Characteristics of burst and buildup cells. Journal of Neurophysiology 73:2313-
2333.
Munoz DP, Waitzman DM, Wurtz RH (1996) Activity of neurons in monkey superior
colliculus during interrupted saccades. Journal of Neurophysiology 75:2562-2580.
Peel TR, Dash S, Lomber SG, Corneil BD (2019) Frontal eye field inactivation alters the
readout of superior colliculus activity for saccade generation in a task-dependent
manner. bioRxiv:646604.
Port NL, Wurtz RH (2003) Sequential activity of simultaneously recorded neurons in the
superior colliculus during curved saccades. J Neurophysiol 90:1887-1903.
Robinson DA (1972) Eye movements evoked by collicular stimulation in the alert monkey.
Vision Res 12:1795-1808.
Ross J, Morrone MC, Burr DC (1997) Compression of visual space before saccades.
Nature 386:598-601.
Schall J, Hanes D, Thompson K, King D (1995) Saccade target selection in frontal eye
field of macaque. I. Visual and premovement activation. The Journal of
Neuroscience 15:6905-6918.
Schall JD, Hanes DP (1993) Neural basis of saccade target selection in frontal eye field
during visual search. Nature 366:467-469.
Shen L, Alexander GE (1997) Neural correlates of a spatial sensory-to-motor
transformation in primary motor cortex. J Neurophysiol 77:1171-1194.
Stanford TR, Freedman EG, Sparks DL (1996) Site and parameters of microstimulation:
evidence for independent effects on the properties of saccades evoked from the
primate superior colliculus. Journal of Neurophysiology 76:3360-3381.
Tehovnik EJ, Sommer MA, Chou IH, Slocum WM, Schiller PH (2000) Eye fields in the
frontal lobes of primates. Brain Res Brain Res Rev 32:413-448.
Tian X, Yoshida M, Hafed ZM (2016) A Microsaccadic Account of Attentional Capture and
Inhibition of Return in Posner Cueing. Frontiers in Systems Neuroscience 10.
Van Opstal AJ, Goossens HHLM (2008) Linear ensemble-coding in midbrain superior
colliculus specifies the saccade kinematics. Biological cybernetics 98:561-577.

Waitzman DM, Ma TP, Optican LM, Wurtz RH (1991) Superior colliculus neurons mediate
the dynamic characteristics of saccades. Journal of Neurophysiology 66:1716-1737.
Willeke KF, Tian X, Buonocore A, Bellet J, Ramirez-Cardenas A, Hafed ZM (2019)
Memory-guided microsaccades. Nature Communications 10:3710.
Wurtz RH, Goldberg ME (1972) Activity of superior colliculus in behaving monkey. 3. Cells
discharging before eye movements. J Neurophysiol 35:575-586.
Wurtz RH, Albano JE (1980) Visual-motor function of the primate superior colliculus. Annu
Rev Neurosci 3:189-226.
966

Supplementary figures

Figure S1 Injected "visua" spikes in our experiments were in neurons that were not directly involved in generating the saccades that were being altered in our main analyses of Figs. 2-6. Mean firing rate of all of our neurons aligned to saccade onset (blue line) for eye movements made towards the RF (**A**, orange) or opposite it (**B**, green). The movements were all selected during a baseline pre-stimulus period (-100 to -25 ms from stimulus onset) in the absence of any visual stimuli inside the RF's. Error bars denote 95% confidence intervals. The neurons did not show any bursts for microsaccade generation, confirming that we were only measuring visual bursts and not concurrent movement-related activity in Figs. 2-6.

Figure S2 Visual bursts in the SC could happen intra-saccadically whether the movement being
generated was towards the recorded neurons' RF locations or opposite them. Peak firing rate for all the
recorded neurons when the visual burst was happening with or without microsaccades (similar analyses to
Fig. 2, but now separating movement directions relative to RF locations; Methods). **(A)** When microsaccades
were going towards the neurons' RF locations (orange), there was no difference between peak firing rate
compared to the peak firing rate without microsaccades (t(84) = -0.3867, p = 0.6999). **(B)** When microsaccades
were going opposite to the RF locations (green), the peak firing rate was slightly reduced compared to the
peak firing rate without microsaccades (t(84) = 3.6223, p < $4.9876*10^{-4}$), but it was still clearly present.
Therefore, regardless of movement directions, intra-saccadic SC visual bursts can still occur.
991
992

Figure S3 More eccentric stimuli relative to the generated movement amplitudes had weaker effects on metric alterations than the less eccentric stimuli of Fig. 3. (A) Time courses of microsaccade amplitudes relative to stimulus onset when the visual stimuli were presented at eccentricities >4.5 deg (and <20 deg; Fig. 1B). The figure is otherwise formatted identically to Fig. 3. As can be seen, there were weaker effects of more eccentric stimuli on microsaccades, even though the stimuli were made bigger to fill the RF's (Methods) and also even though the raw visual bursts showed similar properties to the more central neurons' visual bursts (**B** and Fig. S4). **(B)** Normalized firing rates of the more eccentric neurons relative to stimulus onset (compared to Fig. 3B). The raw firing rates are shown in Fig. S4, and, together with the current figure, they demonstrate that there was a weaker impact of more eccentric spiking activity on microsaccades; that is, the more eccentric bursts were similar in strength to the more central bursts (Fig. S4), but they still had a smaller impact on microsaccade amplitudes in **A**. Note that, consistent with Hafed and Ignashchenkova (2013); Buonocore et al. (2017b); Malevich et al. (2020a), microsaccade amplitudes at the time of SC visual bursts were decreased relative to baseline (by a small amount) for movements that were opposite the stimulus locations (see Fig. 6). This suggests that visual bursts opposite a planned movement might hamper the movement's execution (Buonocore et al., 2017b).

L014 **Figure S4 Despite smaller effects on microsaccade amplitudes (Fig. S3), more eccentric visual bursts**
L015 **were not weaker than more central ones.** Peak firing rate measurements for neurons with an RF location
L016 ≤4.5 deg (Near, orange) or >4.5 deg (Far, blue) from fixation. Each dot represents one neuron. Solid squares
L017 represent the averages for each group. Error bars represents two standards error of the mean. In this example,
L018 the visual stimuli presented to the neurons were gratings with the second highest contrast.
L019

L020

Figure S5 Intra-saccadic "visual" spikes from more eccentric neurons still linearly increased microsaccade amplitudes, but with a much weaker effect size. (A) Radial eye position relative to saccade onset grouped by the number of "visual" spikes counted between movement onset and saccade peak velocity, for eye movements going towards the recorded neuron's RF location (similar to Fig. 4A). In this analysis, the RF was always located at an eccentricity >4.5 deg (Fig. 1B). We used the same grouping and color conventions as in Fig. 4A but in gradients of blue instead of red. When no spikes were recorded during the eye movement, saccade amplitudes were relatively small (darkest blue curve). Adding visual spikes in the SC map during the ongoing movements slightly increased their amplitudes (1 to 6, color-coded from dark to light blue). However, the effect was much milder than for neurons closer in eccentricity to the foveal movement endpoints (Fig. 4A). **(B)** Mean saccade amplitude against the number of intra-saccadic visual spikes (faint blue dots), similar in formatting to Fig. 4B. There was a linear increase in amplitude relative to the number of injected visual spikes (blue line), similar to Fig. 4B for the more central neurons. However, the slop of the effect was significantly lower. The solid lines represent the best linear fit of the underlying raw data. Error bars represent two standard errors of the mean.

Time relative to microstimulation onset (ms) (A)

Time relative to microstimulation onset (ms) (B)

Time relative to microstimulation onset (ms) (C)

Figure S6 Triple pulse microstimulation, our strongest experimental manipulation in Figs. 6, 7, did not by itself evoke saccades. (A) One example microstimulation session with a short spike train of 70 ms duration at 300Hz and with a current amplitude of 30µA. Horizontal (orange) and vertical (purple) eye positions are shown from individual trials aligned on microstimulation onset (blue line). Soon after the microstimulation pulse train onset, saccadic eye movements were evoked in a direction congruent with the microstimulated SC site (up to the right). About 100 ms later, the eyes made a saccade back to the fixation point to recover, especially since the fixation spot was still visible during the microstimulation train. The red vertical lines on the x-axis indicate each microstimulation pulse with a spacing of 3.3 ms. **(B)** Same as **A**, but with only a train of 3 microstimulation pulses (still at 300Hz and current amplitude of 30µA). No eye movements were systematically evoked after the microstimulation train was applied. The figure shows ten example trials. **(C)** Histogram of saccadic reaction times relative to microstimulation onset during the injection of 3 electrical spikes (the condition in **B**). The frequency of saccades was the same before and after microstimulation, supporting the conclusion that 3 electrical spikes were insufficient, on their own, to evoke eye movements.