A miR-124-mediated post-transcriptional mechanism controlling the cell fate switch of astrocytes to induced-neurons

Elsa Papadimitriou[1], Paraskevi N. Koutsoudaki[1,&], Timokratis Karamitros[2,#], Dimitra Karagkouni[3,#], Dafni Chroni-Tzartou[4], Christos Gkemisis[1], Maria Margariti[1], Evangelia Xingi[5], Irini Thanou[1], Socrates J. Tzartos[4], Artemis G. Hatzigeorgiou[3], Dimitra Thomaidou[1,5,*]

Affiliations

[1]Neural Stem Cells and Neuro-imaging Group, Department of Neurobiology, Hellenic Pasteur Institute

[2]Bioinformatics and Applied Genomics Unit, Department of Microbiology, Hellenic Pasteur Institute

[3]DIANA-Lab, Hellenic Pasteur Institute & Dept. of Computer Science and Biomedical Informatics, Univ. of Thessaly

[4]Laboratory of Molecular Neurobiology and Immunology, Department of Neurobiology, Hellenic Pasteur Institute

[5]Light Microscopy Unit, Hellenic Pasteur Institute

[#]equally contributing authors

[&]Present address: Molecular Carcinogenesis Group, Department of Histology and Embryology, Medical School, National and Kapodistrian University of Athens, Athens, Greece

[*] Corresponding author

Email: thomaidou@pasteur.gr

Abstract

miRNA miR-124 has been employed supplementary to neurogenic TFs and other miRNAs to enhance direct neurogenic conversion by suppressing multiple non-neuronal targets. Aim of the study was to investigate whether miR-124 is sufficient to drive direct reprogramming of astrocytes to induced-neurons (iNs) on its own and elucidate its independent mechanism of reprogramming action. Our data show that miR-124 is a potent driver of the reprogramming switch of astrocytes towards an immature neuronal fate by directly targeting the RNA-binding protein Zfp36l1 and subsequently de-repressing Zfp36l1 neurogenic interactome. To this end miR-124 contribution in iNs production largely recapitulates endogenous neurogenesis pathways, being further enhanced upon addition of the neurogenic compound ISX9, which greatly improves both miR-124-induced reprogramming efficiency and iNs' functional maturation. Importantly, miR-124 is potent to guide direct neuronal reprogramming of reactive astrocytes to iNs of cortical identity *in vivo* following cortical trauma, a novel finding confirming its robust reprogramming action under neuroinflammatory conditions.

Introduction

Direct astrocytic reprogramming to induced-neurons (iNs) is a powerful approach for manipulating cell fate, as it takes advantage of the intrinsic neural stem cell (NSC) potential of reactive astrocytes (Magnusson et al. 2014), while it offers the possibility of reprogramming resident brain cells. To this end astrocytic cell fate conversion to iNs has been well-established in vitro (Berninger et al. 2007; Heinrich et al. 2010; Aravantinou-Fatorou et al. 2015) and in vivo (Guo et al. 2014; Torper et al. 2013; Mattugini et al. 2019) using combinations of transcription factors (TFs) or chemical cocktails (L. Zhang et al. 2015; Xiang Li et al. 2015; Gao et al. 2017). Challenging the expression of lineage-specific TFs during reprogramming is accompanied by changes in the expression of regulatory RNAs, mostly miRNAs, that post-transcriptionally modulate high numbers of neurogenesis-promoting factors, indicating that miRNAs act as key regulators of cell fate conversion (Xuekun Li and Jin 2010). To this end, miRNAs have been introduced, supplementary or alternatively to TFs, to instruct direct neuronal reprogramming (Yoo et al. 2011).

Among neurogenic miRNAs, miR-124 has been shown to contribute to efficient neurogenic conversion of fibroblasts when coupled with certain TFs or other miRNAs, in particular miR-9/9*,and the strong neuronal reprogramming potential of this cocktail has been elaborately studied at the transcriptomic and epigenetic level (Ambasudhan et al. 2011; Abernathy et al. 2017; Wohl and Reh 2016; Victor et al. 2014). miR-124 is the most abundant miRNA in the CNS, participating both in embryonic (Maiorano and Mallamaci 2009) and adult neurogenesis (Cheng et al. 2009; Åkerblom et al. 2012), while its expression persists in mature neurons (Deo et al. 2006). miR-124 acts globally to increase the expression levels of neuronal genes, by repressing components of major neuronal gene repressor complexes, such as the anti-neural transcriptional repressor REST complex (Visvanathan et al. 2007; Baudet et al. 2012; Volvert et al. 2014) and the Polycomb Repressive Complex 2 (PRC2) (Neo et al. 2014; S. W. Lee et al. 2018), while it also participates at the post-transcriptional regulation of neuronal transcripts by targeting the neuron-specific splicing global repressor Ptbp1 (Makeyev et al. 2007). Besides its roles in transcriptional and post-transcriptional regulation, miR-124 neurogenic action also relates to chromatin dynamics, as it acts as a key mediator of a chromatin permissive environment for neuronal reprogramming through its involvement in the formation of the neuron specific chromatin remodeling complex nBAF (Yoo et al. 2009).

However, although miR-124 has been lately utilized in many reprogramming cocktails for the neurogenic conversion of fibroblasts (Ambasudhan et al. 2011; Birtele et al. 2019; Jiang et al.

2015; Victor et al. 2018; Yoo et al. 2011), neither its potential to induce fate conversion of
astrocytes to iNs *in vitro* or *in vivo*, nor its mechanism of action in instructing direct
reprogramming on its own have been investigated. In this study we show that miR-124 is
sufficient to instruct reprogramming of cortical astrocytes to immature induced-neurons (iNs)
in vitro controlling the reprogramming "switch" of astrocytes towards the neuronal fate by
down-regulating genes with important regulatory roles in astrocytic function. Among these
we identified for the first time the RNA binding protein Zfp36l1, implicated in ARE-mediated
mRNA decay (Lai et al. 2000), as a direct target of miR-124 and further found certain neuronal-
specific Zfp36l1targets that participate in cortical development being de-repressed in miR-
124-iNs, implying that miR-124-induced reprogramming recapitulates pathways of
endogenous cortical neurogenesis. Importantly, by blocking miR-124 specific binding in
Zfp36l1 3'UTR, we revealed that miR-124/Zfp36l1 interaction is one of the drivers of miR-124-
induced astrocytic cell fate switch and induction of neuronal identity. To enhance the
neuronal differentiation of reprogrammed immature iNs, we combined miR-124 with
isoexasole-9 (ISX9) chemical compound known to possess neurogenesis-promoting properties
(Schneider et al. 2008; Xiang Li et al. 2015). Our functional and mechanistic analysis of the two
molecules combination revealed that in vitro addition of ISX9 promoted both the neurogenic
conversion and greatly enhanced the functional maturation of miR-124+ISX9-iNs reinforcing
their passage through a Tbr2+ intermediate stage. Importantly, *in vivo* miR-124 was also
potent either alone or along with ISX9, to guide neuronal reprogramming of reactive
astrocytes to iNs following cortical trauma, with the vast majority of iNs exhibiting deep-layer
cortical identity, implying the strong miR-124 reprogramming capacity within the injured brain
micro-environment.

Results

miR-124 is sufficient to instruct reprogramming of postnatal cortical astrocytes to immature induced- neurons (immature-iNs)

In order to study the potential of miR-124 to instruct neuronal reprogramming of astrocytes on its own, cultured postnatal day3-5 (P3-5) mouse cortical astrocytes were transfected with the miR-124-3p mimic 3 times every two days and were supplemented with the antioxidants vitamin E and ascorbic acid (**Fig.1A**). After 1 week (day7) nearly 35% of miR-124 over-expressing cells in culture were Tuj1+ exhibiting multipolar morphology **(Fig.1B,C)**, as compared to control astrocytes that received scrambled miRNA (sc-miRNA) (**Suppl.Fig.1A**), where no Tuj1positivity was detected. Still, miR-124-iNs exhibited low differentiation potential and only 19% of the cells remained Tuj1+ at day14 of the reprogramming protocol (**Fig.1B, C**). The ability of miR-124 to instruct neurogenic reprogramming was further supported by RT-qPCR expression analysis of several neurogenic transcription factors (TFs) at day7, where miR-124 overexpression induced the up-regulation of the mRNA levels of the proneural bHLH TFs *Mash1* and to a lesser extend *Neurog2* (**Fig.1D**), while it additionally up-regulated TFs related to both dorsal (*Tbr2, Tbr1, Fezf2* and *Cux1*) (**Fig.1E**) and ventral telencephalon development (*Gsx2, Dlx1*) (**Fig.1F**). We also observed up-regulation of TFs related to neuronal differentiation (*Sox4, Sox11, Hes6*) (**Fig.1G**), however we failed to detect an up-regulation of *NeuroD1* (**Fig.1G**), which is known to play a crucial role in neuronal reprogramming (Pataskar et al. 2016; Guo et al. 2014; Matsuda et al. 2019). Instead, miR-124 significantly reduced *NeuroD1* mRNA levels, implying that this reduction may contribute to the low differentiation capacity of miR-124-iNs. Further, immunofluorescence analysis indicated that the majority of miR-124-iNs (nearly 80%) were Mash1+ and also exhibited low Tbr2 expression (nearly 70%), while only a small percentage of reprogrammed cells (15%) were positive for the ventral TF Gsx2 (**Fig.1H, I**), indicating that the Mash1-Tbr2 trajectory is most prominently activated in the majority of miR-124-iNs.

The neurogenic compound ISX9 greatly enhances the miR-124-induced reprogramming efficiency and differentiation state of iNs

The observed down-regulation of NeuroD1 by miR-124 prompted us to supplement the reprogramming medium from day2 to day10 with the chemical compound ISX9, known to up-regulate NeuroD1 levels and enhance neuronal differentiation (Schneider et al. 2008). Indeed, the addition of ISX9 led to the acquisition of a more differentiated neuronal-like phenotype

with a smaller soma and longer projections (**Fig.2A**) and significantly increased the percentage

of Tuj1+ reprogrammed cells to 62% at day7, with an average of 38% of them being detected

at day14 (**Fig.2B**). Importantly, ISX9 was potent to reverse the miR-124-induced reduction of

NeuroD1 mRNA levels (**Fig.2C**) and to greatly elevate the transcriptional levels of *Neurog2* and

Tbr2 peaking at day7 (**Fig.2D**), while it also induced a moderate reduction in the mRNA levels

of *Mash1* (**Fig.2E**). Interestingly, ISX9 was not able to induce reprogramming of astrocytes on

its own (sc-miRNA+ISX9) (**Suppl.Fig.1B**), despite evoking robust up-regulation of the mRNA

levels of *NeuroD1* (**Fig.2D**) and other neurogenic TFs (**Suppl.Fig.1C**) and to a small, but

significant, extend the protein levels of Mash1 and Tbr2 (**Suppl.Fig.1D, E**). However, addition

of ISX9 in the reprogramming medium along with miR-124 significantly increased the

percentage of Tbr2+/Tuj1+ iNs, without significantly affecting either the percentage of

Mash1+/Tuj1+ iNs or Gsx2+/Tuj1+ iNs relative to miR-124 alone (**Fig.2F, G**). Quantification of

the protein levels of Mash1 and Tbr2 by measuring their mean nuclear fluorescence intensity

in miR-124-iNs and miR-124+ISX9-iNs at day7, revealed a significant enhancement of Tbr2

protein levels (**Fig.2H**) and a down-regulation of Mash1 levels (**Fig.2I**) following ISX9 addition.

These results led us to the inference that addition of ISX9 in the reprogramming medium

reinforces the passage of miR-124+ISX9-iNs through a Tbr2+ intermediate stage, which seems

to be the main reprogramming route these iNs follow. Interestingly, the addition of ISX9 also

significantly enhanced the expression of *Insm1* (**Suppl.Fig.1F**), a key transcriptional regulator

of intermediate progenitors (IPs) (Elsen et al. 2018), further supporting the notion that iNs

pass through an intermediate stage bearing molecular characteristics of endogenous IPs.

miR-124+ISX9-iNs exhibit characteristics of mature, electrophysiologically active neurons

The majority of miR-124-iNs and miR-124+ISX9-iNs were positive for the cortical deep-layer

TF Tbr1 at day14 (**Suppl.Fig.2A, B**). We mainly observed a moderate nuclear Tbr1 expression

in miR-124-iNs, whereas miR-124+ISX9-iNs exhibited strong cytoplasmic Tbr1 expression,

besides a moderate nuclear one (**Suppl.Fig.2A**). After 21 days in culture, nearly 80% of Tuj1+

miR-124+ISX9-iNs were also positive for the mature neuronal markers MAP2 and Synapsin1,

exhibiting a differentiated neuronal morphology (**Fig3.A,B**), while miR-124-iNs at day21 did

not exhibit signs of further maturation and only a small percentage of them were positive for

MAP2 and Synapsin1 (**Fig.3B**). Additionally, at day28 the majority of miR-124+ISX9-iNs (nearly

90%) were positive for the glutamatergic marker vGlut1 (**Fig.3C, D**), while only 12% of them

were found positive for GABA (**Fig.3D, Suppl.Fig.2C**).

In order to further establish the functional maturation state of miR-124-iNs and miR-124+ISX9-iNs we performed electrophysiological analysis with whole-cell voltage-clamp and current-clamp recordings at different time points from day15 to day27. Rapidly inactivating inward Na^+ currents and persistent outward K^+ currents were recorded in miR-124+ISX9-iNs after 22 days in culture (n=47 out of 80 analyzed cells) in response to depolarizing voltage steps (**Fig.3E, left panel, Fig. 3F**), while further application of TTX and TEA – selective Na^+ channels' (Na_V) and K^+ channels' (K_V) blockers respectively – confirmed that the Na_V were responsible for the inward currents and K_V for the outward currents (**Fig.3E, middle and right panels**). A small amount of K_V channels started being present in miR-124+ISX9-iNs before day21, while they became more evident in days 23-27 **(Fig.3F)**, in accordance with the ability of almost all recorded miR-124+ISX9-iNs (n=21 out of 30 recorded cells) to generate repetitive action potentials (APs) upon membrane depolarization (**Fig.3G**). Finally, rare spontaneous post-synaptic current activity was detected in few mature miR-124+ISX9-iNs (day27) (**Fig.3H**). On the other hand, miR-124-iNs exhibited lower amounts of Na_V and K_V channels (not shown) and thus were not capable of firing action potentials (APs) (n=15 cells).

The majority (80%) of miR-124+ISX9-iNs (**Fig.3I**) and a few miR-124-iNs (not shown) were capable of responding to different concentrations of GABA early in the course of their maturation (day22), even before the appearance of APs, which is in compliance with the expression of GABA receptors in early stages of neuronal development (Luján, Shigemoto, and López-Bendito 2005). Additionally, miR-124+ISX9-iNs were capable of responding to L-glutamate in a concentration dependent manner (**Fig.3J**), while L-glutamate-sensitive inward current was completely blocked after co-application of 100μM L-glutamate and 20μM CNQX, indicating the presence of AMPA/kainite receptors (**Fig.3K**).

miR-124 and ISX9 exhibit both independent and cooperative transcriptional contributions in the reprogramming process to iNs

To in-depth analyze the molecular mechanism through which miR-124 contributes to the reprogramming process either alone or following ISX9 supplementation, we performed RNA-sequencing of miR-124-iNs and miR-124+ISX9-iNs at day7, using as controls astrocytes obtained the initial day of the reprogramming (day1) and sc-miRNA transfected astrocytes at day7. The differential expression analysis was performed between day7 miR-124-iNs or miR-124+ISX9-iNs and day1 astrocytes (astro) (miR-124-iNs vs astro and miR-124+ISX9-iNs vs astro respectively), whereas the day7 sc-miRNA-transfected astrocytes (sc-miRNA astro) were used as the ultimate control for the analysis of miR-124 target genes (see Fig.5 below). We

identified 4,233 differentially expressed genes (DEGs) in miR-124-iNs vs astro and 6,652 DEGs

in miR-124+ISX9-iNs vs astro (1≤log$_2$(fold change)≤-1, FDR<0.05) (**Suppl.Fig.3A**).

Heat map analysis of DEGs (miR-124-iNs vs astro and miR-124+ISX9-iNs vs astro) that belonged

to the GO terms: Glial cell differentiation, Gliogenesis, Astrocyte development, Generation of

neurons, Neuron differentiation, Regulation of neuron differentiation, Neurotransmitter

transport and Synaptic signaling (**Fig.4A**) indicated that miR-124 alone efficiently down-

regulated a gene cluster enriched in astrocytic genes (**Cluster I**), with a small additional

contribution from ISX9. At the same time miR-124 up-regulated a gene cluster of neuronal

specific genes (**Cluster III**) that was to a large extend further up-regulated by ISX9

supplementation, while ISX9 highly up-regulated a neuronal specific gene cluster (**Cluster II**)

that was most exclusively expressed in miR-124+ISX9-iNs.

GO enrichment analysis of biological processes for the up-regulated DEGs of both miR-124-

iNs vs astro and miR-124+ISX9-iNs vs astro, further revealed that miR-124 alone up-regulated

genes related to generation of neurons and neuron differentiation as well as to more specific

neuronal functions mostly associated to synaptic transmission (**Fig.4B in orange**), while the

addition of ISX9 greatly enhanced the number of up-regulated genes related not only to

similar GO terms up-regulated by miR-124, but also to more mature neuronal functions such

as action potential, axon development and subtype specific synaptic transmission (**Fig.4B in**

red). Furthermore, enrichment analysis of the down-regulated DEGs of miR-124-iNs vs astro

indicated that many of them were related to cell cycle, gliogenesis, and astrocyte

differentiation (**Fig.4C**). Interestingly, this analysis revealed a strong effect of miR-124 in

down-regulating components of many signaling pathways, including MAPK, PKB, canonical

Wnt, TGF-β, BMP, Notch and JAK/Stat signaling pathways (**Fig.4C**), known to play important

roles in astrocytic identity and function (Gross et al. 1996; Acaz-Fonseca et al. 2019; Kang and

Hébert 2011; Yang et al. 2012).

Since reprogramming is a process that implicates great changes in the transcriptomic, post-

transcriptomic and epigenetic landscape of trans-differentiating cells, we sought to identify

the differentially expressed transcription factors (TFs), epigenetic factors (EFs) and RNA

binding proteins (RBPs) in our datasets that could possibly drive the reprogramming process.

Heat map analysis of astrocytic TFs (**Suppl.Fig.3B**) indicated that miR-124 alone potently

down-regulated TFs related to astrocytic function such as *Id1, Id3, Tcf4, Tcf7l1, Rbpj, Nfic,*

Zcchc24, Pparg, Nr3c1 and *Tead1*, while the addition of ISX9 exhibited only a small

contribution to their down-regulation (**Suppl.Fig.3B**). Importantly, validation of many of those

genes with RT-qPCR verified the observed trend and also indicated that ISX9 alone (sc-

miRNA+ISX9) failed to down-regulate their mRNA levels (**Suppl.Fig.3C**). In parallel, heat map
analysis of up-regulated neuronal-specific TFs revealed that miR-124 led to the up-regulation
of TFs related to telencephalon development such as *Tox, Foxo6, Scrt1, Scrt2, Rcor2, Rarb,*
Rxrg, Dlx5 and *Sox21* (**Fig.4D**), along with TFs that we had already identified through prior RT-
qPCR analysis (*Mash1, Insm1, Hes6, Sox4, Fezf2, Gsx2, Dlx1*). Additionally, the
supplementation with ISX9 increased the number of TFs implicated in telencephalon
development, among which *Eomes (Tbr2), Prdm8, Ovol2, Tfap2c, Tshz2, Lhx6* and *Cux2*
(**Fig.4D**). Surprisingly, we identified a rather large set of TFs highly up-regulated only following
ISX9 addition, related to more ventral/caudal brain regions than the telencephalon, such as
the retina, midbrain, hindbrain/spinal cord (**Fig.4D**), a finding posing the possibility that ISX9
expands region-specific neuronal identity at least at the transcriptional level. Validation of
selected TFs/EFs expressed either in telencephalon (**Suppl.Fig.3D**) or in midbrain
(**Suppl.Fig.3E**) and hindbrain/spinal cord (**Suppl.Fig.3F**) by RT-qPCR verified their observed up-
regulation by the addition of ISX9.

Interestingly, heat map analysis of differentially expressed RNA-binding proteins (RBPs)
revealed that miR-124 was sufficient to down-regulate many RBPs expressed in astrocytes and
other non-neuronal cells, such as the splicing factors *Ptbp1, Snrpa1, Lgals3, Isy1, Ddx39* and
Syf2, as well as the mRNA decay proteins *Zfp36, Zfp36l1, Zfp36l2* (**Fig.4E**). In addition, miR-124
moderately up-regulated several neuronal specific RBPs, mostly relevant to mRNA splicing and
stabilization, among which *Elavl2, Elavl4, Nova1, Rbfox1, Rbfox2, Celf3, Nol3, Nol4* and
Adarb1, while the addition of ISX9 induced further up-regulation of their mRNA levels and
significantly increased the number of genes implicated in neuron specific splicing, editing or
translation such as *Aplp1, Celf4, Celf5, Celf6, Elavl3, Ern2, Esrp2, Rbfox3, Rbm11, Ssrm4* and
Tdrd6 (**Fig.4E**).

The analysis of differentially expressed EFs revealed that miR-124 increased on its own the
levels of several EFs related to epigenetic transcriptional activation, including the H3K4
methyltransferase *Kmt2c*, the DNA demethylase *Tet1*, the chromatin remodeling factors
Smarca1, Smarca2, Chd7, as well as the neuronal BAF (nBAF) complex component *Ss18l1*
(Crest) (**Suppl.Fig.4G**). On the other hand ISX9 further contributed in up-regulating the brain
specific chromatin remodeling factor Chd5, the nBAF complex components *Actl6b* and *Dpf3*
and the H3K4 methyltransferase *Smyd3*, while interestingly it majorly contributed in down-
regulating EFs related to epigenetic transcriptional repression, such as the H3K9
methyltransferases *Suv39h1* and *Suv39h2*, the histone deacetylases *Hdac5* and *Hdac7*, the

components of repressor complexes *Cbx5, Sap18* and *Rbbp4*, and the component of non-neuronal BAF complexes *Dpf2* (**Suppl.Fig.4G**).

The above observations led us to the conclusion that miR-124 over-expression is sufficient to induce the astrocytic reprogramming switch towards an immature cortical neuronal fate through down-regulation of many glial specific genes, many of which are implicated in different regulatory levels, such as the transcriptional, post-transcriptional, epigenetic and signaling pathway level. The addition of ISX9, on the other hand, acts auxiliary to miR-124 reprogramming action enhancing neuronal-specific gene transcription contributing to the maturation of miR-124 immature-iNs.

The non-neuronal RBP Zfp36l1 is a novel direct target of miR-124

To get a closer insight into the post-transcriptional effects of miR-124 on the astrocytic transcriptome, we sought to determine the direct targets of miR-124 that could act as drivers of the reprogramming switch. Argonaute (AGO) HITS-CLIP experiments are considered the avant-garde of high-throughput methodologies for the direct detection of miRNA targets on a transcriptome-wide scale. Therefore, we utilized publicly available AGO-HITS-CLIP data, performed in mouse brain cortex (Chi et al. 2009) in order to detect miR-124 binding sites. The analysis revealed 171 miR-124 direct targets that were also defined as down-regulated in the miR-124-iNs vs sc-miRNA astro RNA-Seq analysis ($\log_2$(fold change)≤-1, FDR<0.01). miR-124 targets were subsequently filtered to examine genes expressed in astrocytes, utilizing a published reference list for expressed transcripts in distinct cell types of the mouse brain (Y. Zhang et al. 2014), ending up with 130 miR-124 direct target genes (**Fig.5A**).

Among these genes, a prominent target was the RBP Zfp36l1, which is implicated in the mRNA decay (Lai et al. 2000) and is highly expressed in cortical glial cells (Weng et al. 2019), cortical radial glial precursors (Yuzwa et al. 2017; Weng et al. 2019) and other non-neuronal cells (Carrick and Blackshear 2007). miR-124 directly binds to the 3' UTR of Zfp36l1 transcript with perfect seed complementarity (7mer-M8 site) (**Fig.5B**), while in re-analysis of publicly available AGO-HITS-CLIP data from human motor cortex and cingulate gyrus tissues, the miR-124 binding site on the 3' UTR of ZFP36L1 human transcript was found to be conserved (**Fig.5B**).

The efficient down-regulation of *Zfp36l1* by miR-124 was validated by RT-qPCR, where it was shown to be a very early event in the course of reprogramming, with a greater than 50% reduction of the control *Zfp36l1* mRNA levels only 24h after the first transfection (day2) (**Suppl.Fig.4A**). Interestingly, ISX9 alone was not potent to down-regulate *Zfp36l1* mRNA levels

(**Fig.5C**), further supporting our initial hypothesis that ISX9 alone cannot instruct reprogramming of astrocytes to iNs, possibly by failing to down-regulate astrocytic fate genes.

Since Zfp36l1 acts by mediating degradation of its mRNA targets, we were interested in identifying Zfp36l1 mRNA targets, being up-regulated in our analysis. For this purpose, we combined two publicly available Zfp36l1 individual-nucleotide resolution CLIP-Seq data (iCLIP-Seq) from thymocytes (Vogel et al. 2016) and B lymphocytes (Galloway et al. 2016) and ended up with 621 Zfp36l1 direct targets that are up-regulated in miR-124-iNs vs sc-miRNA astro ($\log_2$(fold change)≥1, FDR<0.05), which importantly correspond to 47% of miR-124 up-regulated DEGs. GO enrichment analysis revealed that many of these genes are implicated in neurogenesis, neuron projection development, synaptic transmission, axonogenesis, dendritic morphogenesis and telencephalon development (**Fig.5D**). Interestingly, many of them were also found to regulate transcription and RNA processing (**Fig.5D**), highlighting an important regulatory role for many Zfp36l1 targets, which could possibly have great impact on the reprogramming process. Among these targets we found neurogenic TFs, such as *Tox, Tox3, Rcor2, Cux1, Hes6, Lzts1* and *Mllt11* as well as EFs also related to neurogenesis such as *Chd3, Chd7, Kmt2c* and *Tet1* (**Fig.5E**). Notably, we also identified as Zfp36l1 direct targets the neuronal RBPs *Elavl4, Nova1* and *Rbfox1* (**Fig.5E**). This constitutes a significant finding that delineates the neuronal RBPs' repression directly by Zfp36l1, being relieved upon miR-124-mediated Zfp36l1 down-regulation.

We subsequently examined the mRNA levels of the Zfp36l1 targets, *Elavl4, Nova1, Rbfox1, Rcor2* and *Tox* upon miR-124 overexpression with RT-qPCR and verified that miR-124 was potent to induce their up-regulation (**Fig.5F**). We also verified the increase of the protein levels of the neurogenic TF and target of Zfp36l1, Tox, in miR-124-iNs at day5 by immunofluorescence analysis (**Suppl.Fig.4B, C**).

Notably, we observed that the combination of miR-124 with ISX9 greatly further enhanced the mRNA levels of the RBP targets of Zfp36l1, *Elavl4* and *Rbfox1* (**Suppl.Fig.4D**), as well as the other neuronal RBPs of the nElavl family, *Elavl2* (**Suppl.Fig.4E**) and *Elavl3* (**Suppl.Fig.4F**), highlighting the importance of ISX9 in reinforcing the mRNA levels of significant neuronal specific RBPs during the course of reprogramming.

Collectively, these data show that the targeting of Zfp36l1 by miR-124 de-represses the Zfp36l1 neurogenic interactome, which corresponds to half of the miR-124-up-regulated genes and includes neuronal-specific genes with important regulatory functions. Thus the

miR-124/Zfp36l1 interaction bears the potential to play a key regulatory role in the control of the neurogenic reprogramming switch of astrocytes by miR-124.

Targeting of Zfp36l1 by miR-124 plays a key role in the miR-124-induced cell fate switch of astrocytes to iNs

To investigate the impact of the miR-124/Zfp36l1 interaction on the miR-124-mediated reprogramming of astrocytes, we used a target site blocker (TSB) oligonucleotide, which competitively binds to the miR-124 binding site on the 3'UTR of Zfp36l1 mRNA and blocks its down-regulation by the miR-124-RISC complex (**Fig.6A**). We verified the inhibition of the miR-124-induced down-regulation of Zfp36l1 protein levels by the addition of TSB in a dose dependent manner (miR-124:TSB molecular ratio 4:1, 2:1) at day5 of the reprogramming protocol by western blot (**Fig.6B**). Next, we tested the effect of TSB on the mRNA levels of the neuronal targets of Zfp36l1, *Tox, Rbfox1, Nova1, Rcor2* and *Elavl4* by RT-qPCR initially at day3 and observed a dose-dependent reduction of the miR-124-induced up-regulation of *Tox, Rbfox1, Nova1* and *Rcor2*, but not of *Elavl4*, in the presence of increasing concentrations of TSB (miR-124:TSB molecular ratio 4:1, 2:1 and 1:1) (**Fig.6C**). The mRNA levels of *Tox* and *Rbfox1* retained the same response to TSB at a later time point, at day5 (**Suppl.Fig.5A, B**), an effect that was not observed for the other Zfp36l1 targets *Nova1*, *Rcor2* and *Elavl4* (**Suppl.Fig.5C**), implying that their post-transcriptional regulation becomes more complex as the neuronal conversion is gradually established. Importantly, TSB resulted in a statistically significant and dose-dependent reduction in the percentage of Tuj1+ miR-124-iNs at day5 (**Fig.6D, E**), which was accompanied by an evident alteration of their characteristic iN morphology (**Fig.6D, F**). More specifically, a morphological analysis of the number of processes extending from the soma and the size of the soma of Tuj1+ cells indicated that TSB addition resulted in gradual abolishment of their multipolar neuronal morphology with fine processes, characteristic of miR-124-iNs (**Fig.6D left panel, F**), and instead largely led to the retention of a premature astrocyte-like morphology with bigger soma (**Suppl.Fig.5D**) and none or very few processes (**Fig.6D right panel, F**).

These observations strongly indicate that the targeting of Zfp36l1 by miR-124 plays a crucial role in the miR-124-induced neuronal reprogramming action by unlocking neurogenic genes with important regulatory activity and by controlling the necessary morphological changes.

miR-124 induces reprogramming of resident reactive astrocytes to iNs with deep-layer cortical identity *in vivo* following cortical trauma

To evaluate the *in vivo* reprogramming potential of miR-124 to drive cell fate conversion of resident reactive cortical astrocytes to iNs alone or supplemented by ISX9, we overexpressed miR-124 via lentiviral transfer into the mechanically injured cortex of 3-4 months old mice. To this end we stereotaxically injected either one of the two lentiviruses, control LV-GFP and LV-miR-124-GFP, in order to transduce reactive astrocytes surrounding the cortical injury site, whileISX9 was systemically administered beginning 2 days afterviral injection for 5 consecutive days in a group of mice (**Fig.7A**). We initially sacrificed a group of mice 6 days after viral administration (10 days after cortical trauma induction) to evaluate the original phenotype of transduced cells surrounding the injury site (**Suppl.Fig.6A**) and observed that the majority of them (approximately 68%)in both conditions were GFAP+ reactive astrocytes, while an average of 10% were NeuN+ neurons. The rest of the transduced cells were equally either Iba-1+ microglial cells or Olig2+oligodendrocytes (**Suppl.Fig.6B**). For all markers examined at this time point, there were no statistically significant differences between LV-124-GFP and control LV-GFP, indicating that the two viruses are expected to transduce the same percentage of astrocytes and neurons upon injection into the cortical parenchyma. Further analysis of the percentage of NeuN+/GFP+ transduced cells 3 weeks after the stereotaxic injection revealed that it was significantly increased in the animals that received the miR-124-overexpressing virus, but not the control virus (**Fig.7B, C**). In particular, while approximately 10% of the transduced cells were NeuN+ in LV-GFP-injected animals, showing no significant difference between the 6 days and 3 week time points, 74% of the LV-miR-124-GFP-transduced cells were NeuN+ 3 weeks after the injection, showing significant difference between the two time points, indicating a strong potential of miR-124 to direct conversion of reactive astrocytes into iNs *in vivo*. The same effect was observed when the injection of LV-miR-124-GFPvirus was accompanied by systemic administration of ISX9, where 71% of transduced cells were found to be NeuN+, while no significant difference was observed at 3 weeks after co-treatment of ISX9 in comparison to LV-miR-124-GFP injection alone (**Fig.7C**).

Overexpression of miR-124either alone or along with ISX9 administration, appeared to drive iNs towards a deep layer cortical identity, as the vast majority of the NeuN+ transduced cells after 3 weeks in both groups were also positive for the deep layer cortical marker Tbr1, with percentages reaching 98% in LV-miR-124-iNs and 88% in LV-miR-124-GFP+ISX9-iNs, with no significant difference observed between the two groups (**Fig.7D, E**).

401   The above *in vivo* data indicate that miR-124 is sufficient to drive on its own neurogenic
402   reprogramming of reactive astrocytes within the injured cortex micro-environment that
403   render it a potent candidate molecule for further in vivo application.

Discussion

In this study we attempted to isolate miR-124 mechanism of action from that of other reprogramming co-factors and we provide evidence that miR-124 drives the trans-differentiation switch of cortical astrocytes to an immature iN phenotype of cortical identity, recapitulating its endogenous role in embryonic cortical development (Maiorano and Mallamaci 2009) and SVZ adult neurogenesis (Cheng et al. 2009; Åkerblom et al. 2012). Indeed, miR-124 directed astrocytes through a multipolar intermediate stage expressing TFs such as *Mash1, Tbr2, Insm1, Rcor2, Fezf2* and *Tbr1* all associated with neuronal commitment and differentiation during embryonic (Kwan, Šestan, and Anton 2012; Elsen et al. 2018)and adult neurogenesis (Díaz-Guerra et al. 2013). Intriguingly, miR-124 over-expression led to a significant down-regulation of the neuronal differentiation bHLH TF NeuroD1, which has been reported as a temporal, stage-specific event in early stages of eye development in Xenopus (Liu et al. 2011). To this end to reinforce the differentiation of immature miR-124-iNs, we supplemented miR-124 with the neurogenic compound ISX9, known to up-regulate neuronal specific genes, among which NeuroD1. ISX9 acts by increasing intracellular Ca^{2+} signaling leading to HDAC5 nuclear exit and MEF2 de-repression (Schneider et al. 2008) and has been already used in chemical reprogramming protocols for the conversion of fibroblasts and astrocytes to iNs (Gao et al. 2017; Xiang Li et al. 2015). Additionally, ISX9 enhances the differentiation of iPSC-derived neurons from Huntington's disease patients, indicating that it also possesses a neuronal maturation-promoting activity (Lim et al. 2017). Indeed, the addition of ISX9 greatly improved both the reprogramming efficiency and differentiation status of miR-124+ISX9-iNs, by enhancing *Tbr2, Neurog2* and *NeuroD1* transcriptional levels reinforcing their passage through a Tbr2+ intermediate stage and subsequently promoting their differentiation leading to the acquisition of electrophysiologically active iNs. Interestingly, despite a robust up-regulation of the levels of NeuroD1 and many other neuronal TFs by ISX9 alone, control sc-miRNA+ISX9-treated astrocytes failed to undergo reprogramming. NeuroD1 has been extensively reported to possess a strong 'pioneer factor' reprogramming capacity towards the neuronal fate upon its viral-mediated overexpression (Pataskar et al. 2016; Matsuda et al. 2019; Guo et al. 2014; Rivetti Di Val Cervo et al. 2017), however we hypothesize that very high levels of *NeuroD1* – not being triggered by ISX9 supplementation in our system – are required to inflict these effects.

The RNA-Seq analysis we performed, highlighted the importance of miR-124 over-expression in the down-regulation of astrocytic TFs, RBPs, EFs and components of signaling pathways with significant regulatory role in astrocytic identity and function, which is in accordance with the

documented role of miR-124 in favoring neuronal fate at the expense of astrocytic (Neo et al.

2014) and implies that astrocytic identity barriers need to be repressed before the induction

of a neuronal cell fate during the reprogramming process. Conversely, ISX9 had a small

contribution in the repression of astrocytic genes, thus this might be the reason for its inability

to confer the reprogramming switch on its own.

Our analysis of the direct miR-124 targets utilizing AGO-HITS-CLIP data from mouse cortex (Chi

et al. 2009) revealed the RBP *Zfp36l1* as a novel target of miR-124. Of note, we also verified

the miR-124/Zfp36l1 interaction in human bearing the same binding site, by analyzing AGO-

HITS-CLIP data from human motor cortex and cingulate gyrus (Boudreau et al. 2014), a finding

highlighting the importance of this conserved interaction during mammalian brain evolution.

Many studies have identified several direct targets of miR-124 with important regulatory role

in neurogenesis, acting at the transcriptional level, such as the TFs *Sox9* (Cheng et al. 2009),

Lhx2 (Sanuki et al. 2011) and the components of the REST repressor complex, *Scp1*

(Visvanathan et al. 2007) and *Rcor1* (Baudet et al. 2012; Volvert et al. 2014); or at the

epigenetic level, such as the component of the PRC2 complex *Ezh2* (Neo et al. 2014) and the

component of the BAF complex BAF53a (Yoo et al. 2009); as well as at the post-transcriptional

level, such as the RBP involved in alternative splicing Ptbp1 (Makeyev et al. 2007).

Here, we report for the first time that miR-124 is directly implicated in the regulation of

another process mediated by RBPs, the mRNA decay, apart from the well-characterized miR-

124/Ptbp1 circuitry (Makeyev et al. 2007; Yeom et al. 2018; Xue et al. 2013). Zfp36l1 is a

member of the Zfp36 family of proteins along with Zfp36 and Zfp36l2, which act by binding to

AU rich elements (AREs) in the 3'UTR of their mRNA targets mediating their destabilization(Lai

et al. 2000). Zfp36l1 is expressed in cortical radial glial precursors of the VZ (Yuzwa et al. 2017;

Weng et al. 2019; DeBoer et al. 2013), in cortical glial cells (Weng et al. 2019) and in non-

neuronal cells (Carrick and Blackshear 2007; Chen et al. 2015). Interestingly, the ARE-

dependent mRNA decay is regulated by other neurogenic miRNAs as well, since the close

paralog of Zfp36l1, Zfp36, is targeted by miR-9 (Dai et al. 2015), suggesting a combined

regulation of Zfp36 family members by the two miRNAs to counteract the destabilization of

neuronal mRNAs during neurogenesis.

In order to identify targets of Zfp36l1 being up-regulated in our system we examined data

from Zfp36l1-iCLIP-Seq experiments in thymocytes (Vogel et al. 2016) and B lymphocytes

(Galloway et al. 2016). Although our analysis was restricted to a not relevant to the astrocytic

transcriptome, we identified a rather large number of Zfp36l1 targets being up-regulated in

miR-124-iNs that interestingly correspond to nearly half of the up-regulated genes by miR-

124. Importantly, many Zfp36l1 targets exhibit significant regulatory role in neurogenesis and neuronal differentiation, such as the studied here TFs *Tox* and *Rcor2* and the RBPs *Rbfox1*, *Elavl4,* and *Nova1*. Further on, to experimentally validate our hypothesis that Zfp36l1 is a direct miR-124 target implicated in astrocytes' cell fate switch, we used a target site blocker (TSB) that efficiently antagonizes the binding of miR-124 in the 3'UTR of *Zfp36l1* mRNA and showed that the disruption of miR-124/Zfp36l1 interaction has a negative impact on the reprogramming process, reducing the number of Tuj1+ miR-124-iNs and abolishing their multipolar morphology. Of note, our experiments with the TSB blocker revealed that among the neuron-specific RBPs that are targets of Zfp36l1, *Rbfox1* and *Nova1* exhibit a pattern of direct post-transcriptional regulation by contrast to *Elavl4*, implying the stronger involvement of other RBPs in its post-transcriptional regulation. In accordance with this, Zfp36 has been reported to target *Elavl2, Elavl3, Elavl4* and *Nova1* (Dai et al. 2015), uncovering a complementary and synergistic role of Zfp36l1 and Zfp36 in repressing neuron-specific RBPs in non-neuronal cells.

Importantly, the here identified Zfp36l1 direct target, *Rbfox1*, has been reported to be a critical post-transcriptional regulator of gene expression during cortical development (J. A. Lee et al. 2016; Weyn-Vanhentenryck et al. 2014). To the same direction, we show here that Zfp36l1 also directly targets the TFs *Tox* and *Rcor2*, both known to be involved in the control of mammalian corticogenesis, with Tox regulating among others the levels of dorsal telenchephalon TFs and marker of IPs Tbr2 (Artegiani et al. 2015) and Rcor2 being expressed along with Insm1 in cortical IPs promoting dorsal telencephalon patterning (Wang et al. 2016; Elsen et al. 2018). Additionally, the repression of the multipolar morphology of miR-124-iNs by the TSB blocker could imply a role for the miR-124/Zfp36l1 interaction in the acquisition of a multipolar morphology characteristic of endogenous VZ/SVZ IPs (Mizutani et al.,2018), a fact corroborated by the GO analysis of Zfp36l1 target genes, which presented the GO terms neuronal projection development and dendrite morphogenesis with high statistical significance. Thus, this analysis revealed certain transcriptional and post-transcriptional cortical neurogenesis regulators being de-repressed by the miR124/Zfp36l1 interaction **(Fig.6G)**, strengthening the notion that miR-124-mediated reprogramming of astrocytes recapitulates molecular aspects of endogenous neurogenesis.

Furthermore, the addition of ISX9 significantly contributed to the up-regulation of the mRNA levels of the neuronal RBPs *Elavl2, Elavl3, Elavl4* (Ince-Dunn et al. 2012; Scheckel et al. 2016) and *Rbfox1* (Weyn-Vanhentenryck et al. 2014), reinforcing the action of miR-124 in inducing the switch from the neuronal transcripts' destabilizing RBPs to the stabilizing neuronal RBPs

(**Fig.6G**). In parallel, ISX9 supplementation greatly enhanced the transcriptional levels and the
number of TFs related to telencephalon development and/or adult neurogenesis, further
promoting the cortical identity of iNs already initiated by miR-124. Surprisingly, ISX9 also up-
regulated a large set of TFs related to the development of other non-telencephalic brain
regions such as the midbrain and the hindbrain/spinal cord. However, the mechanism of the
observed here ISX9-mediated up-regulation of ventral/caudal TFs still remains elusive. Of
note, ISX9 has been shown to affect the epigenetic landscape leading to an open chromatin
state by increasing H3/H4 acetylation in pancreatic β-cells (Dioum et al. 2011), suggesting that
an epigenetic mechanism may be responsible for the observed transcriptional regional
expansion, a hypothesis that needs to be further explored.

Here we also demonstrate that miR-124 is capable of directly converting reactive cortical
astrocytes surrounding a mechanically-induced trauma to iNs of deep-layer cortical identity.
Brain injury is known to facilitate in vivo reprogramming, rendering astrocytes more plastic by
activating NSC genes' expression (Götz et al. 2015). *In vivo* reprogramming of reactive
astrocytes to neuronal precursors and iNs has been achieved following forced expression of
TFs among which Sox2 (Niu et al. 2013), Neurog2 (Grande et al. 2013), NeuroD1 (Guo et al.
2014; Rivetti Di Val Cervo et al. 2017) and Nurr1/Neurog2 (Mattugini et al. 2019), in some
cases combined with anti-apoptotic and/or anti-oxidant treatment to enhance survival
(Gascón et al. 2016). However, it is the first time that miR-124 is shown to be sufficient to
reprogram reactive astrocytes to NeuN+ iNs with a high efficiency of 74%. Unlike to the *in*
vitro situation, co-administration of ISX9 didn't significantly further enhance miR-124-induced
reprogramming efficiency *in vivo*, possibly reflecting the strong effect of the surrounding
micro-environment in supporting miR-124-induced neuronal conversion. Indeed, pro-
inflammatory cytokines like TNF, and growth factors such as EGF, FGF2 and SHH present in
the injured cortex have been reported to facilitate the neurogenic conversion of astrocytes
(Gabel et al. 2016; Grande et al. 2013; Sirko et al. 2013).

Interestingly, following either miR-124 or miR-124+ISX9 treatment all iNs possess a deep-layer
cortical identity expressing Tbr1 protein. Taking into account the fact that the injury has been
performed in a deep cortical area, this observation indicates that, besides the *in vitro* observed
endogenous potency of reprogrammed iNs to up-regulate Tbr1 levels, region-specific
neuronal identity barriers are imposed to newly produced iNs by the surrounding cortical
microenvironment, in accordance with recent findings unravelling the existence of layer-
driven neuronal identity during *in vivo* astrocytic reprogramming to iNs (Mattugini et al. 2019).

Taken together this study highlighted the strong potency of miR-124 to instruct the cell fate

switch of astrocytes, post-transcriptionally triggering cortical neurogenesis pathways being

unlocked by the direct targeting of *Zfp36l1*. Additionally, our *in vitro* results give mechanistic

insight from a developmental perspective into the combined action of miR-124 and ISX9 in

driving direct reprogramming of astrocytes to mature iNs. Importantly, our findings point to

an even stronger *in vivo* reprogramming capacity of miR-124 than the one observed *in vitro*,

opening the possibility for its use in therapeutic protocols with minimal support by other

neurogenic or anti-inflammatory/anti-oxidant factors.

Materials and Methods

Primary cultures of postnatal cortical astrocytes

Primary postnatal astrocytic cultures from P3-P5 mice were prepared as previously described (Aravantinou-Fatorou et al. 2015). Briefly, the cerebral cortexes from 2-3 P3-P5 C57BL/6 mice were collected in ice cold HBSS (Invitrogen), the tissue was washed three times with HBSS and digested with 0.04% trypsin (Sigma) and 10µg/ml DNAse (Sigma) for 5 min at 37°C. After digestion cells were mechanically dissociated, centrifuged for 5 min at 850 rpm (120 g), re-suspended in DMEM 4.5g/lt glucose (Invitrogen) containing 10% FBS (Invitrogen), 1% Penicillin/Streptomycin (Pen/Strep) (Sigma) and placed in a T75 flask pre-coated with poly-D-lysine (PDL) (Sigma). When culture reached confluence (usually after 7 days), the flask was shaken in a horizontal shaker at 200-250 rpm for 20h, in order to obtain a pure astrocytic culture, free from neurons, oligodendrocytes and microglia. The remaining cells were digested with 0.5% trypsin-EDTA (Invitrogen) for 5 min at 37°C, centrifuged at 850 rpm, re-suspended in fresh DMEM 4.5g/lt glucose 10% FBS, 1% Pen/Strep and divided in two new T75 flasks pre-coated with PDL. Half of the medium was changed every two days.

In vitro reprogramming protocol

For the reprogramming of astrocytes to induced-neurons, 40,000 astrocytes were seeded in 10mm coverslips coated with 20µg/ml poly-L-ornithine (PLO) (Sigma) overnight and 5µg/ml laminin for 3 h at 37°C (Sigma). Once cells reached>90% confluence (usually after 1-2 days) they became transfected with 80nM miR-124-3p mimics or sc-miRNA mimics (negative control) (Thermo) using Lipofectamine 2000 (Invitrogen) according to manufacturer's instructions (day1). The next day the astrocytic medium (DMEM 4.5g/lt glucose, 10% FBS, 1% Pen/Strep) was replaced with the reprogramming medium: Neurobasal (Invitrogen) supplemented with 1X B-27 (Invitrogen), 1X GlutaMAX (Invitrogen), 20µM vitamin E (a-tocopherol) (Sigma) and 200mM ascorbic acid (Sigma). The same process of transfection was repeated twice at day3 and day5. Vitamin E was added to the medium until day4, while ascorbic acid was added throughout the reprogramming protocol. At day7 the reprogramming medium was changed to the neuronal differentiation medium: Neurobasal supplemented with 1X B-27, 1X GlutaMAX, 20ng/ml BDNF (R&D Systems), 0.5mM cAMP (Sigma) and 200mM ascorbic acid. In the miR-124+ISX9-reprogrammed cells, 10µM of ISX9 chemical compound (Tocris) were added from day2 to day10. All the mediums added to the reprogrammed cells were pre-conditioned for 24 h in a confluent astrocytic culture.

RT-qPCR analysis

For the RT-qPCR analysis experiments, total RNA was extracted using the Nucleospin miRNA kit (Macherey-Nagel) and 500-800ng of total RNA were used for cDNA synthesis with the Superscript II reverse transcriptase (Invitrogen) according to manufacturer's instructions. Quantitative real time PCR was performed using SYBR Select Master Mix (Applied Biosystems) and samples were run in the ViiA 7 Real-Time PCR System (Applied Biosystems). The primers used are listed in **Table 1**. Each sample was analyzed in triplicates, gene expression was calculated using the ΔΔCt method and all the results were normalized to β-actin expression. Relative expression was estimated setting the values of sc-miRNA transfected astrocytes to 1. All experiments were performed at least in triplicates.

Immunocytochemistry

Cells were washed once with PBS and then fixed with 4% paraformaldehyde for 20 min at room temperature. Afterwards, cells were washed three times with PBS and blocked with 5% normal donkey serum (NDS) (Merck-Millipore), 0.1% Triton X-100 in PBS for 1 h at room temperature. For nuclear staining, cells were permeabilized with 0.25% Triton X-100 in PBS for 10 min at room temperature and washed three times with PBS prior to blocking. Next, cells were incubated with primary antibodies, diluted in 1% NDS, 0.05% Triton X-100 in PBS overnight at 4°C. The next day, cells were washed three times with PBS and incubated with secondary antibodies diluted in 1% NDS, 0.05% Triton X-100 in PBS for 2h at room temperature. The nuclei of the cells were stained with ProLong Gold Antifade Reagent with DAPI (Cell Signalling). The following primary antibodies were used in this study: mouse anti-Tuj1 (Covance, 1:600), chicken anti-Tuj1 (Millipore, 1:1000), mouse anti-MAP2 (Millipore, 1:200), rabbit anti-Synapsin1 (Abcam, 1:200), rat anti-Mash1 (R&D Systems, 1:100), rabbit anti-Tbr2 (Abcam, 1:200), rabbit anti-Gsx2 (Millipore, 1:400), rabbit anti-Tox (Atlas antibodies, 1:200), mouse anti-vGlut1 (Millipore, 1:1000) and rabbit anti-GABA (Sigma, 1:10,000). The secondary antibodies used in this study were Alexa Fluor 546-, Alexa Fluor 488- and Alexa Fluor 647-conjugated secondary antibodies (Life Technologies). Images were acquired with a 20x or 40x objective (1024x1024 pixels, 1μm Z-step) using a Leica TCS SP8 confocal microscope (LEICA Microsystems). For each experiment measurements from 20-25 fields per coverslip were obtained for each condition.

Electrophysiology

For whole-cell recordings iNs plated in PLO-laminin coated coverslips were used for electrophysiological experiments beginning at day15 up to day27 of the reprogramming protocol. The coverslips with the cells were placed onto a recording chamber and viewed using an Olympus CKX41 microscope with a 40x lens. The cells were bathed in a solution containing: 140mM NaCl, 2.8mM KCl, 2mM $CaCl_2$, 4mM $MgCl_2$, 10mM Glucose, and 20mM HEPES. For whole-cell recordings we used a capillary glass with Filament (Sutter instrument) to fabricate low resistance recording pipettes (~5 MΩ) and filled with: 140mM KCl, 2mM $CaCl_2$, 2mM $MgCl_2$, 2mM Mg-ATP, 5mM EGTA and 10mM HEPES. Osmolarity and pH of all solutions were adjusted appropriately before experiments. Data were acquired at room temperature (22–24°C) using an EPC9 HEKA amplifier and an ITC-16 acquisition system with a patchmaster software (HEKA). Data analysis was carried out using OriginPro8. **Voltage protocols:** The membrane of the cells was held at a holding potential of -70 mV and step depolarizing pulses were applied. Depolarization steps were applied for 50 msec in 10 mV increments from -80 mV to +50 mV with a sweep interval time of 1 sec and sweep duration of 500 ms. Each depolarizing pulse was proceeded by a hyperpolarizing step to -120 mV. **Current protocols:** Cells we held at their resting membrane potential (0pA) and depolarizing current steps from -20 pA to 200 pA from a holding current of 0pA were applied.

RNA-Seq experiment and bioinformatic analysis

For the RNA-Seq experiment, the following samples were prepared in 3 biological replicates: astrocytes (day1), sc-miRNA-transfected astrocytes (day7), miR-124-iNs (day7) and miR-124+ISX9-iNs (day7). Total RNA was extracted using the Nucleospin miRNA kit (Macherey-Nagel) according to manufacturer's instructions. Libraries were prepared with TruSeq RNA Library Prep Kit v2 (Illumina) and 75c single-end sequencing in an Illumina NextSeq 550 sequencer. Raw libraries were quality checked and preprocessed using FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Mapping of reads against the Mouse Transcriptome (v.GRCm38.rel79) and transcript abundance estimation on Transcripts Per Million (TPM) values was performed using kallisto (Bray et al. 2016). Analysis of differential expression, interpretation and visualization was subsequently performed using kallisto-compatible Sleuth tool (Pimentel et al. 2017) and R-base functions. Gene ontology (GO) enrichment analysis was performed using the Gene Ontology Panther Classification System (http://pantherdb.org/).

Analysis of AGO-CLIP-Seq data

AGO-HITS-CLIP datasets, performed in mouse brain cortex tissue (P13 neocortex, 5 replicates)

and human brain tissues (motor cortex, cingulate gyrus) from 2 individuals, were retrieved

from the publications (Chi et al. 2009) and (Boudreau et al. 2014) respectively. Raw libraries

were quality checked using FastQC (www.bioinformatics.babraham.ac.uk/projects/fastqc/),

while adapters/ contaminants were detected utilizing an in-house developed pipeline and the

Kraken suite (Davis et al. 2013). Pre-processing was performed with TrimGalore (Krueger

2015) and Cutadapt (Martin 2011). CLIP-Seq libraries were aligned against the reference

genomes, i.e. GRCh38 and mm10 assemblies for human and mouse respectively, with

GMAP/GSNAP (Wu and Nacu 2010) spliced aligner, allowing up to 2 mismatches. microCLIP

CLIP-Seq-guided model (Paraskevopoulou et al. 2018) was utilized to identify binding events

for the expressed miRNAs. In case of multiple replicates (i.e. mouse brain cortex) a miRNA

binding event had to be present in at least two replicates to be considered as valid. Top

expressed miRNAs were retrieved from the relevant publications. Human and mouse

transcriptomes were compiled from ENSEMBL v96 (Cunningham et al. 2019) to annotate the

retrieved miRNA binding events. Identified miRNA binding sites residing on 3' UTR regions

were retained and subsequently filtered to preserve only genes expressed in astrocytes. A list

of ~10,000 genes, expressed in astrocytes, with FPKM ≥ 2, was retrieved from a reference

publication and retained for analysis (Y. Zhang et al. 2014).

Target Site Blocker (TSB) experiment

For the functional validation of the miR-124/Zfp36l1 interaction a custom made miRCURY

locked nucleic acid (LNA) miRNA Power Target Site Blocker (TSB) (Qiagen) was used with the

following sequence: TTACAAGGCACTAAGTTGCTT. TSB was transfected in astrocytes along

with sc-miRNA or miR-124-3p mimics using Lipofectamine 2000 (Invitrogen) in different

molecular ratios: miR-124:TSB, 4:1 (80nM:20nM), 2:1 (80nM:40nM) and 1:1 (80nM:80nM).

Western blot

Cells were washed once with ice-cold PBS and lysed for 15 min in ice-cold lysis buffer (150mM

NaCl, 50mM Tris (pH 7.5), 1%v/v Triton X-100, 1mM EDTA, 1mM EGTA, 0.1% SDS, 0.5% sodium

deoxycholate) containing PhosSTOP phosphatase inhibitors and a complete protease inhibitor

mixture (Roche Life Science), then centrifuged at 20,000 g for 15 min, followed by collection

of the supernatant and measurement of the protein concentration by Bradford assay

(Applichem). Proteins were separated by SDS-polyacrylamide gel electrophoresis (PAGE) and

transferred onto nitrocellulose membranes (Maine Manufacturing). Nonspecific binding sites

were blocked in TBS/ 0.1% Tween 20/5% skimmed milk for 1 hour at 20°C followed by
overnight incubation with primary antibodies diluted in TBS/0.1% Tween20/5% BSA. Primary
antibodies used were rabbit anti-Zfp36l1 (Abcam, 1:500) and mouse anti-βactin (Millipore,
1:1000). Incubation with HRP-conjugated secondary antibodies, anti-mouse-HRP (Thermo,
1:10,000) and anti-rabbit-HRP (Thermo, 1:5,000) was performed for 2 hours at room
temperature and protein bands were visualized using the Clarity Western ECL Substrate (BIO-
RAD).

Lentiviral production

For lentiviral in vivo transduction, VSV-G (Vesicular Stomatitis Virus–Glycoprotein)–
pseudotyped lentiviruses were used either for the over-expression of miR-124 along with GFP
or as control expressing only GFP. More specifically, for lentiviral particles' production, HEK
293T cells cultured in 10-cm Petri dishes at a 50-60% confluence were co-transfected with 10
μg lentiviral plasmid expressing miR-124-1 precursor under the CMV promoter and GFP under
the EF1 promoter (SBI System Biosciences) or 10 μg lentiviral plasmid expressing GFP under
the CMV promoter and the packaging plasmids pVSV-G (3.5 μg), MDL (6.5 μg), and RSV-REV
(2.5 μg) (all kindly provided by Dr. Matsas lab) with calcium phosphate. The following day the
culture medium was replaced with fresh one, the supernatant containing the lentiviral
particles was collected 48 h and 72 h (second harvest) after transfection and concentrated by
ultracentrifugation at 25,000 rpm (80,000 x g) for 2 h at 4°C using a sucrose gradient.

Cortical trauma, viral injection and ISX9 administration

This study was carried out in strict compliance with the European Directive 2010/63/EU and
the Greek National Law 161/91 for Use of Laboratory Animals, according to FELASA
recommendations for euthanasia and the National Institutes of Health Guide for Care and Use
of Laboratory Animals. All protocols were approved by the Animal Care and Use Committee
of the Hellenic Pasteur Institute (Animal House Establishment Code: EL 25 BIO 013). License
No 2585/29-5-18 for the experiments was issued by the Greek authorities of the Veterinary
Department of the Athens Prefecture. The manuscript was prepared in compliance with the
ARRIVE guidelines for reporting animal research.
Adult male and female FVB mice (8-16 weeks old) were deeply anaesthetized using inhalable
isoflurane, and positioned in a stereotaxic apparatus. The dorsal surface of the skull was
exposed through a midline incision and a burr hole was drilled at the following coordinates:
antero-posterior (AP) -1.0 mm, caudal to Bregma; lateral (L) 1.0 mm to the midline (Franklin
and Paxinos, 2001). A 26-gauge needle was inserted into the brain parenchyma in a depth of
0.9 mm from the surface of the brain to create a trauma in the cortex, avoiding the corpus

callosum and hippocampus. The inserted needle was moved along the anterior-posterior axis between positions (AP) -1.1 and -0.9 to widen the trauma. The skin was sutured, a local analgesic cream containing 2.5% lidocain and 2.5% prilocain was applied and the animals were kept warm until they were fully awake. Viral injection took place 4 days after the cortical trauma. A 10µl Hamilton syringe (Hamilton) with a 26-gauge needle was slowly inserted into the brain tissue at coordinates (AP) -1.1 mm, (L) 1.0 mm, and (V): 1.0 mm, from the same burr hole on the skull and 2µl of lentiviral concentrate was injected at a rate of 0.5 µl/min. The needle was left in position for 5 min after each injection and then withdrawn gently. A second viral injection was repeated at coordinates (AP) -0.9 mm, (L) 1.0 mm, and (V): 1.0 mm with similar procedures, and surgery was completed as described above. A group of animals was injected with the lentivirus LV-miR-124-GFP (group LV-124) and another one with the control lentivirus LV-GFP (group LV-GFP). A subgroup of the LV-124 group received intraperitoneally 20 mg/kg of ISX9 (Tocris) diluted in (2-Hydroxypropyl)-β-cyclodextrin (Sigma) (ISX9 concentration: 2 mg/ml in 30% (2-Hydroxypropyl)-β-cyclodextrin (Sigma) diluted in sterile ddH2O) once a day, for 5 consecutive days, beginning 48 h after lentiviral injection. Animals were sacrificed 6 days or 3 weeks after viral injection.

Tissue Preparation, Histology, and Immunohistochemistry

For histology, mice were deeply anaesthetized by inhaling isoflurane, and perfused with 4% paraformaldehyde (PFA) via left cardiac ventricle. The brains were removed, post-fixed in 4% PFA overnight, and then cryo-protected in 20% sucrose overnight. Tissues were then frozen in -20°C isopentane and cut into 20 µm-thick coronal sections on a cryostat (Leica CM1900), collected on silane-coated slides and stored at -20°C. For detection of specific antigens with immunofluorescence, sections were left for 15 min in room temperature, washed in PBS, and blocked with 5% normal goat or donkey serum (Merck-Millipore) in PBT (0.1% Triton X-100/PBS) for 1 h. Incubation with primary antibodies took place overnight at 4°C. Primary antibodies to assess transduced cell identity used were: chicken polyclonal anti-GFP (Abcam, 1:1000) to detect the transplanted cells, mouse monoclonal anti-neuronal nuclei (NeuN) (Merck-Millipore, 1:300) to identify mature neurons, rabbit polyclonal anti-glial fibrillary acidic protein (GFAP) (Dako,1:600) to detect astrocytes; rabbit polyclonal anti-oligodendrocyte transcription factor 2 (Olig2) (Merck-Millipore, 1:200) to detect oligodendrodytes, rabbit polyclonal anti-ionized calcium-binding adapter 1 (Iba-1) (Wako, 1:600) for detection of microglia and rabbit polyclonal anti-Tbr1 (Abcam,1:250) for detection of deep layer cortical neurons. Following incubation with primary antibodies, sections were washed with PBS and incubated for 2 h with the appropriate secondary antibodies conjugated with AlexaFluor 488

(green), 546 (red), or 647 (blue) and Hoechst (Molecular Probes) for nuclei staining. Finally,

sections were washed and coverslipped with Mowiol (Calbiochem). Images were acquired

with a 40x objective using Leica TCS SP8 and Leica TCS-SP5II confocal microscopes (LEICA

Microsystems).

Image analysis

Images were analyzed using Fiji/ImageJ software (National Institutes of Health).

In vitro analysis: mean fluorescence intensity of Tbr2, Tox and Mash1 staining inside the cell

nuclei was quantified using a custom-written macro implemented in Fiji. Initially automatic

detection of nuclei was performed using the Otsu method and Tuj1+ cells were selected

based on their mean intensity value above a user-defined threshold of 40, followed by a

manual validation according to cell morphology (cells with an astrocyte-like morphology with

a big, rectangular soma and none or few processes were excluded). Mean fluorescence

intensity of Tbr2, Tox and Mash1 inside the cell nuclei ROIs was measured both for Tuj1- and

Tuj1+ cells. Quantification was performed in maximum intensity projections. For each

experiment measurements from at least 200-300 cells were obtained for each condition.

Morphological characterization of Tuj1+ cells in the TSB blocker experiments was also

conducted using Fiji. More specifically, morphological characterization included quantification

of the cell body area and the number of processes extending from the soma of Tuj1+ cells.

Tuj1+ cells were selected based on the mean intensity value of their soma above a user-

defined threshold and were sorted in 3 groups: cells with a multipolar morphology bearing 3

or more processes extending from the soma, cells with 1-2 processes and cells exhibiting an

astrocyte-like morphology with a rectangular soma possessing none or 1-2 processes.

Quantification was performed in maximum intensity projections. For each experiment

measurements from at least 100-150 cells were obtained for each condition.

In vivo analysis: for each animal and each immunofluorescence staining, cell counting was

performed on brain sections collected at 240 lm intervals across the whole antero-posterior

extent of the hippocampus (bregma-0.5mm up to -2.5mm) in a total number of 3-4 mice for

each experimental condition. For estimation of the ratio of transduced cells that have a

specific phenotype, images of all GFP+ cells found in each set of sections were acquired and

double-positivity with cell type-specific markers was evaluated by an observer "blind" to

treatment group and time point. All GFP+ cells found in the cortex within these sections were

imaged and analyzed. Representative confocal images shown in **Fig.7** and **Suppl.Fig.6** are

obtained from coronal sections, antero-posterior positions between 21.3 and 22.2 relative to

Bregma.

Statistical analysis

All in vitro quantified data are presented as average ± SD, unless otherwise indicated. Two-tailed Student t-test was used to calculate statistical significance with p values for all the data obtained from the experiments with the TSB blocker, while for the rest of the data a one-tailed Student t-test was used. p values less than 0.05 ($p < 0.05$) were considered indicative of significance. In vivo data were assessed using a one-way analysis of variance (ANOVA). When interactions were detected, group comparisons were performed using a two-sample assuming unequal variances test.

Data Availability

High throughput sequencing data are deposited in the European Nucleotide Archive under Study accession PRJEB38603.

Acknowledgments

This work was financially supported by: 'BIOIMAGING-GR: A Greek Research Infrastructure for Visualizing and Monitoring Fundamental Biological Processes (MIS 5002755)', funded by the Operational Program "Competitiveness, Entrepreneurship and Innovation" (NSRF 2014-2020), co-financed by Greece and the European Union (European Regional Development Fund); ARISTEIA-II 'Astro-Rep' 3713 Excellence Grant of the Greek Ministry of Education and Fondation Santé Grant 2017-2018, awarded to DT. We also acknowledge funding from the Stavros Niarhos Foundation (SFN) Grant to the Hellenic Pasteur Institute, as part of the Foundation's initiative to support the Greek Research Center ecosystem; Greek General Secretariat of Research and Technology 'Action for the Study of Neurodegenerative Diseases on the Basis of Precision Medicine' and 'KRIPIS-II' Action (MIS 5002486) under the Operational Strategic Reference Framework 2014–2020. We would like to thank Dr. Era Taoufik for critical comments on the manuscript.

Author Contributions

EP and DT conceived whole project, designed experiments and analyzed the data; EP, CG and MM conducted in vitro experiments; DCT and SJT designed and performed electrophysiology experiments; TK, DK and AGH designed and performed bioinformatics analysis and EP contributed in analyzing the data; PNK designed and performed in vivo experiments, analyzed relevant data and contributed in paper writing; IT contributed to in vivo experiments and relevant data analysis; EX developed the Fiji macro and helped with image analysis workflow; EP and DT wrote the manuscript; DT supervised the project and acquired funding.

Conflict of Interest

The authors report no conflict of interest.

References

Abernathy, Daniel G., Woo Kyung Kim, Matthew J. McCoy, Allison M. Lake, Rebecca Ouwenga, Seong Won Lee, Xiaoyun Xing, et al. 2017. "MicroRNAs Induce a Permissive Chromatin Environment That Enables Neuronal Subtype-Specific Reprogramming of Adult Human Fibroblasts." *Cell Stem Cell* 21 (3): 332-348.e9. https://doi.org/10.1016/j.stem.2017.08.002.

Acaz-Fonseca, Estefania, Ana Ortiz-Rodriguez, Iñigo Azcoitia, Luis M. Garcia-Segura, and Maria Angeles Arevalo. 2019. "Notch Signaling in Astrocytes Mediates Their Morphological Response to an Inflammatory Challenge." *Cell Death Discovery*. https://doi.org/10.1038/s41420-019-0166-6.

Åkerblom, Malin, Rohit Sachdeva, Isabelle Barde, Sonia Verp, Bernhard Gentner, Didier Trono, and Johan Jakobsson. 2012. "MicroRNA-124 Is a Subventricular Zone Neuronal Fate Determinant." *Journal of Neuroscience*. https://doi.org/10.1523/JNEUROSCI.0558-12.2012.

Ambasudhan, Rajesh, Maria Talantova, Ronald Coleman, Xu Yuan, Saiyong Zhu, Stuart A. Lipton, and Sheng Ding. 2011. "Direct Reprogramming of Adult Human Fibroblasts to Functional Neurons under Defined Conditions." *Cell Stem Cell*. https://doi.org/10.1016/j.stem.2011.07.002.

Aravantinou-Fatorou, Katerina, Felipe Ortega, Dafni Chroni-Tzartou, Nasia Antoniou, Cornelia Poulopoulou, Panagiotis K. Politis, Benedikt Berninger, Rebecca Matsas, and Dimitra Thomaidou. 2015. "CEND1 and NEUROGENIN2 Reprogram Mouse Astrocytes and Embryonic Fibroblasts to Induced Neural Precursors and Differentiated Neurons." *Stem Cell Reports* 5 (3): 405–18. https://doi.org/10.1016/j.stemcr.2015.07.012.

Artegiani, Benedetta, Antonio M Jesus Domingues, Sara Bragado Alonso, Elisabeth Brandl, Simone Massalini, Andreas Dahl, and Federico Calegari. 2015. "Tox: A Multifunctional Transcription Factor and Novel Regulator of Mammalian Corticogenesis." *The EMBO Journal* 34 (7): 896–910. https://doi.org/10.15252/embj.201490061.

Baudet, Marie Laure, Krishna H. Zivraj, Cei Abreu-Goodger, Alistair Muldal, Javier Armisen, Cherie Blenkiron, Leonard D. Goldstein, Eric A. Miska, and Christine E. Holt. 2012. "MiR-124 Acts through CoREST to Control Onset of Sema3A Sensitivity in Navigating Retinal Growth Cones." *Nature Neuroscience*. https://doi.org/10.1038/nn.2979.

Berninger, Benedikt, Marcos R. Costa, Ursula Koch, Timm Schroeder, Bernd Sutor, Benedikt Grothe, and Magdalena Götz. 2007. "Functional Properties of Neurons Derived from in Vitro Reprogrammed Postnatal Astroglia." *Journal of Neuroscience*. https://doi.org/10.1523/JNEUROSCI.1615-07.2007.

Birtele, Marcella, Yogita Sharma, Srisaiyini Kidnapillai, Shong Lau, Thomas B. Stoker, Roger A. Barker, Daniella Rylander Ottosson, Janelle Drouin-Ouellet, and Malin Parmar. 2019. "Dual Modulation of Neuron-Specific MicroRNAs and the REST Complex Promotes Functional Maturation of Human Adult Induced Neurons." *FEBS Letters*. https://doi.org/10.1002/1873-3468.13612.

Boudreau, Ryan L., Peng Jiang, Brian L. Gilmore, Ryan M. Spengler, Rebecca Tirabassi, Jay A. Nelson, Christopher A. Ross, Yi Xing, and Beverly L. Davidson. 2014. "Transcriptome-Wide Discovery of MicroRNA Binding Sites in Human Brain." *Neuron*. https://doi.org/10.1016/j.neuron.2013.10.062.

Bray, Nicolas L., Harold Pimentel, Páll Melsted, and Lior Pachter. 2016. "Near-Optimal Probabilistic RNA-Seq Quantification." *Nature Biotechnology*. https://doi.org/10.1038/nbt.3519.

Carrick, Danielle M., and Perry J. Blackshear. 2007. "Comparative Expression of Tristetraprolin (TTP) Family Member Transcripts in Normal Human Tissues and Cancer Cell Lines." *Archives of Biochemistry and Biophysics*. https://doi.org/10.1016/j.abb.2007.04.011.

Chen, Ming Tai, Lei Dong, Xin Hua Zhang, Xiao Lin Yin, Hong Mei Ning, Chao Shen, Rui Su, et al. 2015. "ZFP36L1 Promotes Monocyte/Macrophage Differentiation by Repressing CDK6." *Scientific Reports*. https://doi.org/10.1038/srep16229.

Cheng, Li Chun, Erika Pastrana, Masoud Tavazoie, and Fiona Doetsch. 2009. "MiR-124 Regulates Adult Neurogenesis in the Subventricular Zone Stem Cell Niche." *Nature Neuroscience*. https://doi.org/10.1038/nn.2294.

Chi, Sung Wook, Julie B. Zang, Aldo Mele, and Robert B. Darnell. 2009. "Argonaute HITS-CLIP Decodes MicroRNA-MRNA Interaction Maps." *Nature*. https://doi.org/10.1038/nature08170.

Cunningham, Fiona, Premanand Achuthan, Wasiu Akanni, James Allen, M. Ridwan Amode, Irina M. Armean, Ruth Bennett, et al. 2019. "Ensembl 2019." *Nucleic Acids Research*.

https://doi.org/10.1093/nar/gky1113.

Dai, Weijun, Wencheng Li, Mainul Hoque, Zhuyun Li, Bin Tian, and Eugene V. Makeyev. 2015. "A Post-
Transcriptional Mechanism Pacing Expression of Neural Genes with Precursor Cell
Differentiation Status." *Nature Communications*. https://doi.org/10.1038/ncomms8576.

Davis, Matthew P.A., Stijn van Dongen, Cei Abreu-Goodger, Nenad Bartonicek, and Anton J. Enright.
2013. "Kraken: A Set of Tools for Quality Control and Analysis of High-Throughput Sequence
Data." *Methods*. https://doi.org/10.1016/j.ymeth.2013.06.027.

DeBoer, E. M., M. L. Kraushar, R. P. Hart, and M. R. Rasin. 2013. "Post-Transcriptional Regulatory
Elements and Spatiotemporal Specification of Neocortical Stem Cells and Projection Neurons."
Neuroscience. https://doi.org/10.1016/j.neuroscience.2013.05.042.

Deo, Monika, Jenn Yah Yu, Kwan Ho Chung, Melissa Tippens, and David L. Turner. 2006. "Detection of
Mammalian MicroRNA Expression by in Situ Hybridization with RNA Oligonucleotides."
Developmental Dynamics. https://doi.org/10.1002/dvdy.20847.

Díaz-Guerra, Eva, Jaime Pignatelli, Vanesa Nieto-Estévez, and Carlos Vicario-Abejón. 2013.
"Transcriptional Regulation of Olfactory Bulb Neurogenesis." *Anatomical Record*.
https://doi.org/10.1002/ar.22733.

Dioum, Elhadji M., Jihan K. Osborne, Sean Goetsch, Jamie Russell, Jay W. Schneider, and Melanie H.
Cobb. 2011. "A Small Molecule Differentiation Inducer Increases Insulin Production by
Pancreatic β Cells." *Proceedings of the National Academy of Sciences of the United States of
America*. https://doi.org/10.1073/pnas.1118526109.

Elsen, Gina E., Francesco Bedogni, Rebecca D. Hodge, Theo K. Bammler, James W. MacDonald, Susan
Lindtner, John L.R. Rubenstein, and Robert F. Hevner. 2018. "The Epigenetic Factor Landscape of
Developing Neocortex Is Regulated by Transcription Factors Pax6→ Tbr2→ Tbr1." *Frontiers in
Neuroscience*. https://doi.org/10.3389/fnins.2018.00571.

Gabel, Sebastien, Eric Koncina, Gauthier Dorban, Tony Heurtaux, Cindy Birck, Enrico Glaab, Alessandro
Michelucci, Paul Heuschling, and Luc Grandbarbe. 2016. "Inflammation Promotes a Conversion
of Astrocytes into Neural Progenitor Cells via NF-KB Activation." *Molecular Neurobiology*.
https://doi.org/10.1007/s12035-015-9428-3.

Galloway, Alison, Alexander Saveliev, Sebastian Łukasiak, Daniel J. Hodson, Daniel Bolland, Kathryn
Balmanno, Helena Ahlfors, et al. 2016. "RNA-Binding Proteins ZFP36L1 and ZFP36L2 Promote
Cell Quiescence." *Science*. https://doi.org/10.1126/science.aad5978.

Gao, Longfei, Wuqiang Guan, Min Wang, Huihan Wang, Jiali Yu, Qing Liu, Binlong Qiu, et al. 2017.
"Direct Generation of Human Neuronal Cells from Adult Astrocytes by Small Molecules." *Stem
Cell Reports*. https://doi.org/10.1016/j.stemcr.2017.01.014.

Gascón, Sergio, Elisa Murenu, Giacomo Masserdotti, Felipe Ortega, Gianluca L. Russo, David Petrik,
Aditi Deshpande, et al. 2016. "Identification and Successful Negotiation of a Metabolic
Checkpoint in Direct Neuronal Reprogramming." *Cell Stem Cell* 18 (3): 396–409.
https://doi.org/10.1016/j.stem.2015.12.003.

Götz, Magdalena, Swetlana Sirko, Johannes Beckers, and Martin Irmler. 2015. "Reactive Astrocytes as
Neural Stem or Progenitor Cells: In Vivo Lineage, In Vitro Potential, and Genome-Wide
Expression Analysis." *GLIA*. https://doi.org/10.1002/glia.22850.

Grande, Andrew, Kyoko Sumiyoshi, Alejandro López-Juárez, Jennifer Howard, Bhuvaneswari Sakthivel,
Bruce Aronow, Kenneth Campbell, and Masato Nakafuku. 2013. "Environmental Impact on
Direct Neuronal Reprogramming in Vivo in the Adult Brain." *Nature Communications* 4.
https://doi.org/10.1038/ncomms3373.

Gross, Robert E., Mark F. Mehler, Peter C. Mabie, Ziying Zang, Linda Santschi, and John A. Kessler.
1996. "Bone Morphogenetic Proteins Promote Astroglial Lineage Commitment by Mammalian
Subventricular Zone Progenitor Cells." *Neuron*. https://doi.org/10.1016/S0896-6273(00)80193-
2.

Guo, Ziyuan, Lei Zhang, Zheng Wu, Yuchen Chen, Fan Wang, and Gong Chen. 2014. "In Vivo Direct
Reprogramming of Reactive Glial Cells into Functional Neurons after Brain Injury and in an
Alzheimer's Disease Model." *Cell Stem Cell* 14 (2): 188–202.
https://doi.org/10.1016/j.stem.2013.12.001.

Heinrich, Christophe, Robert Blum, Sergio Gascón, Giacomo Masserdotti, Pratibha Tripathi, Rodrigo
Sánchez, Steffen Tiedt, Timm Schroeder, Magdalena Götz, and Benedikt Berninger. 2010.
"Directing Astroglia from the Cerebral Cortex into Subtype Specific Functional Neurons." *PLoS
Biology*. https://doi.org/10.1371/journal.pbio.1000373.

Ince-Dunn, Gulayse, Hirotaka J. Okano, Kirk B. Jensen, Woong Yang Park, Ru Zhong, Jernej Ule, Aldo Mele, et al. 2012. "Neuronal Elav-like (Hu) Proteins Regulate RNA Splicing and Abundance to Control Glutamate Levels and Neuronal Excitability." *Neuron*. https://doi.org/10.1016/j.neuron.2012.07.009.

Jiang, Houbo, Zhimin Xu, Ping Zhong, Yong Ren, Gaoyang Liang, Haley A. Schilling, Zihua Hu, et al. 2015. "Cell Cycle and P53 Gate the Direct Conversion of Human Fibroblasts to Dopaminergic Neurons." *Nature Communications*. https://doi.org/10.1038/ncomms10100.

Kang, Wenfei, and Jean M. Hébert. 2011. "Signaling Pathways in Reactive Astrocytes, a Genetic Perspective." *Molecular Neurobiology*. https://doi.org/10.1007/s12035-011-8163-7.

Krueger, Felix. 2015. "Trim Galore!: A Wrapper Tool around Cutadapt and FastQC to Consistently Apply Quality and Adapter Trimming to FastQ Files." *Babraham Institute*.

Kwan, Kenneth Y., Nenad Šestan, and E. S. Anton. 2012. "Transcriptional Co-Regulation of Neuronal Migration and Laminar Identity in the Neocortex." *Development*. https://doi.org/10.1242/dev.069963.

Lai, Wi S., Ester Carballo, Judith M. Thorn, Elizabeth A. Kennington, and Perry J. Blackshear. 2000. "Interactions of CCCH Zinc Finger Proteins with MRNA. Binding of Tristetraprolin-Related Zinc Finger Proteins to AU-Rich Elements and Destabilization of MRNA." *Journal of Biological Chemistry*. https://doi.org/10.1074/jbc.M001696200.

Lee, Ji Ann, Andrey Damianov, Chia Ho Lin, Mariana Fontes, Neelroop N. Parikshak, Erik S. Anderson, Daniel H. Geschwind, Douglas L. Black, and Kelsey C. Martin. 2016. "Cytoplasmic Rbfox1 Regulates the Expression of Synaptic and Autism-Related Genes." *Neuron*. https://doi.org/10.1016/j.neuron.2015.11.025.

Lee, Seong Won, Young Mi Oh, Ya Lin Lu, Woo Kyung Kim, and Andrew S. Yoo. 2018. "MicroRNAs Overcome Cell Fate Barrier by Reducing EZH2-Controlled REST Stability during Neuronal Conversion of Human Adult Fibroblasts." *Developmental Cell*. https://doi.org/10.1016/j.devcel.2018.06.007.

Li, Xiang, Xiaohan Zuo, Junzhan Jing, Yantao Ma, Jinlin Jiaming Wang, Defang Liu, Jialiang Zhu, et al. 2015. "Small-Molecule-Driven Direct Reprogramming of Mouse Fibroblasts into Functional Neurons." *Cell Stem Cell* 17 (2): 195–203. https://doi.org/10.1016/j.stem.2015.06.003.

Li, Xuekun, and Peng Jin. 2010. "Roles of Small Regulatory RNAs in Determining Neuronal Identity." *Nature Reviews Neuroscience*. https://doi.org/10.1038/nrn2739.

Lim, Ryan G., Lisa L. Salazar, Daniel K. Wilton, Alvin R. King, Jennifer T. Stocksdale, Delaram Sharifabad, Alice L. Lau, et al. 2017. "Developmental Alterations in Huntington's Disease Neural Cells and Pharmacological Rescue in Cells and Mice." *Nature Neuroscience* 20 (5): 648–60. https://doi.org/10.1038/nn.4532.

Liu, Kaili, Ying Liu, Weichuan Mo, Rong Qiu, Xiumei Wang, Jane Y. Wu, and Rongqiao He. 2011. "MiR-124 Regulates Early Neurogenesis in the Optic Vesicle and Forebrain, Targeting NeuroD1." *Nucleic Acids Research*. https://doi.org/10.1093/nar/gkq904.

Luján, R., R. Shigemoto, and G. López-Bendito. 2005. "Glutamate and GABA Receptor Signalling in the Developing Brain." *Neuroscience*. https://doi.org/10.1016/j.neuroscience.2004.09.042.

Magnusson, Jens P., Christian Göritz, Jemal Tatarishvili, David O. Dias, Emma M.K. Smith, Olle Lindvall, Zaal Kokaia, and Jonas Frisén. 2014. "A Latent Neurogenic Program in Astrocytes Regulated by Notch Signaling in the Mouse." *Science*. https://doi.org/10.1126/science.346.6206.237.

Maiorano, Nicola Antonio, and Antonello Mallamaci. 2009. "Promotion of Embryonic Cortico-Cerebral Neuronogenesis by MiR-124." *Neural Development*. https://doi.org/10.1186/1749-8104-4-40.

Makeyev, Eugene V., Jiangwen Zhang, Monica A. Carrasco, and Tom Maniatis. 2007. "The MicroRNA MiR-124 Promotes Neuronal Differentiation by Triggering Brain-Specific Alternative Pre-MRNA Splicing." *Molecular Cell*. https://doi.org/10.1016/j.molcel.2007.07.015.

Martin, Marcel. 2011. "Cutadapt Removes Adapter Sequences from High-Throughput Sequencing Reads." *EMBnet.Journal*. https://doi.org/10.14806/ej.17.1.200.

Matsuda, Taito, Takashi Irie, Shutaro Katsurabayashi, Yoshinori Hayashi, Tatsuya Nagai, Nobuhiko Hamazaki, Aliya Mari D. Adefuin, et al. 2019. "Pioneer Factor NeuroD1 Rearranges Transcriptional and Epigenetic Profiles to Execute Microglia-Neuron Conversion." *Neuron*. https://doi.org/10.1016/j.neuron.2018.12.010.

Mattugini, Nicola, Riccardo Bocchi, Volker Scheuss, Gianluca Luigi Russo, Olof Torper, Chu Lan Lao, and Magdalena Götz. 2019. "Inducing Different Neuronal Subtypes from Astrocytes in the Injured Mouse Cerebral Cortex." *Neuron* 103 (6): 1086-1095.e5.

https://doi.org/10.1016/j.neuron.2019.08.009.

Mizutani, Ken Ichi. 2018. "Physiological Significance of Multipolar Cells Generated from Neural Stem
Cells and Progenitors for the Establishment of Neocortical Cytoarchitecture." *Genes to Cells*.
https://doi.org/10.1111/gtc.12546.

Neo, Wen Hao, Karen Yap, Suet Hoay Lee, Liang Sheng Looi, Piyush Khandelia, Sheng Xiong Neo,
Eugene V. Makeyev, and I. Hsin Su. 2014. "MicroRNA MiR-124 Controls the Choice between
Neuronal and Astrocyte Differentiation by Fine-Tuning Ezh2 Expression." *Journal of Biological*
Chemistry. https://doi.org/10.1074/jbc.M113.525493.

Niu, Wenze, Tong Zang, Yuhua Zou, Sanhua Fang, Derek K. Smith, Robert Bachoo, and Chun Li Zhang.
2013. "In Vivo Reprogramming of Astrocytes to Neuroblasts in the Adult Brain." *Nature Cell*
Biology 15 (10): 1164–75. https://doi.org/10.1038/ncb2843.

Paraskevopoulou, Maria D., Dimitra Karagkouni, Ioannis S. Vlachos, Spyros Tastsoglou, and Artemis G.
Hatzigeorgiou. 2018. "MicroCLIP Super Learning Framework Uncovers Functional
Transcriptome-Wide MiRNA Interactions." *Nature Communications*.
https://doi.org/10.1038/s41467-018-06046-y.

Pataskar, Abhijeet, Johannes Jung, Pawel Smialowski, Florian Noack, Federico Calegari, Tobias Straub,
and Vijay K Tiwari. 2016. "NeuroD1 Reprograms Chromatin and Transcription Factor Landscapes
to Induce the Neuronal Program." *The EMBO Journal*.
https://doi.org/10.15252/embj.201591206.

Pimentel, Harold, Nicolas L. Bray, Suzette Puente, Páll Melsted, and Lior Pachter. 2017. "Differential
Analysis of RNA-Seq Incorporating Quantification Uncertainty." *Nature Methods*.
https://doi.org/10.1038/nmeth.4324.

Rivetti Di Val Cervo, Pia, Roman A. Romanov, Giada Spigolon, Débora Masini, Elisa Martín-Montañez,
Enrique M. Toledo, Gioele La Manno, et al. 2017. "Induction of Functional Dopamine Neurons
from Human Astrocytes in Vitro and Mouse Astrocytes in a Parkinson's Disease Model." *Nature*
Biotechnology. https://doi.org/10.1038/nbt.3835.

Sanuki, Rikako, Akishi Onishi, Chieko Koike, Rieko Muramatsu, Satoshi Watanabe, Yuki Muranishi,
Shoichi Irie, et al. 2011. "MiR-124a Is Required for Hippocampal Axogenesis and Retinal Cone
Survival through Lhx2 Suppression." *Nature Neuroscience*. https://doi.org/10.1038/nn.2897.

Scheckel, Claudia, Elodie Drapeau, Maria A. Frias, Christopher Y. Park, John Fak, Ilana Zucker-Scharff,
Yan Kou, et al. 2016. "Regulatory Consequences of Neuronal ELAV-like Protein Binding to Coding
and Non-Coding RNAs in Human Brain." *ELife*. https://doi.org/10.7554/eLife.10421.

Schneider, Jay W., Zhengliang Gao, Shijie Li, Midhat Farooqi, Tie Shan Tang, Ilya Bezprozvanny, Doug
E. Frantz, and Jenny Hsieh. 2008. "Small-Molecule Activation of Neuronal Cell Fate." *Nature*
Chemical Biology 4 (7): 408–10. https://doi.org/10.1038/nchembio.95.

Sirko, Swetlana, Gwendolyn Behrendt, Pia Annette Johansson, Pratibha Tripathi, Marcos Costa, Sarah
Bek, Christophe Heinrich, et al. 2013. "Reactive Glia in the Injured Brain Acquire Stem Cell
Properties in Response to Sonic Hedgehog Glia." *Cell Stem Cell*.
https://doi.org/10.1016/j.stem.2013.01.019.

Torper, Olof, Ulrich Pfisterer, Daniel A. Wolf, Maria Pereira, Shong Lau, Johan Jakobsson, Anders
Björklund, Shane Grealish, and Malin Parmar. 2013. "Generation of Induced Neurons via Direct
Conversion in Vivo." *Proceedings of the National Academy of Sciences of the United States of*
America 110 (17): 7038–43. https://doi.org/10.1073/pnas.1303829110.

Victor, Matheus B., Michelle Richner, Tracey O. Hermanstyne, Joseph L. Ransdell, Courtney Sobieski,
Pan Yue Deng, Vitaly A. Klyachko, Jeanne M. Nerbonne, and Andrew S. Yoo. 2014. "Generation
of Human Striatal Neurons by MicroRNA-Dependent Direct Conversion of Fibroblasts." *Neuron*.
https://doi.org/10.1016/j.neuron.2014.10.016.

Victor, Matheus B., Michelle Richner, Hannah E. Olsen, Seong Won Lee, Alejandro M. Monteys,
Chunyu Ma, Christine J. Huh, et al. 2018. "Striatal Neurons Directly Converted from
Huntington's Disease Patient Fibroblasts Recapitulate Age-Associated Disease Phenotypes."
Nature Neuroscience. https://doi.org/10.1038/s41593-018-0075-7.

Visvanathan, Jaya, Soo Kyung Seunghee Lee, Bora Lee, Jae W. Lee, and Soo Kyung Seunghee Lee.
2007. "The MicroRNA MiR-124 Antagonizes the Anti-Neural REST/SCP1 Pathway during
Embryonic CNS Development." *Genes and Development*. https://doi.org/10.1101/gad.1519107.

Vogel, Katharina U., Lewis S. Bell, Alison Galloway, Helena Ahlfors, and Martin Turner. 2016. "The
RNA-Binding Proteins Zfp36l1 and Zfp36l2 Enforce the Thymic β-Selection Checkpoint by
Limiting DNA Damage Response Signaling and Cell Cycle Progression." *The Journal of*

Immunology. https://doi.org/10.4049/jimmunol.1600854.

Volvert, Marie Laure, Pierre Paul Prévot, Pierre Close, Sophie Laguesse, Sophie Pirotte, James Hemphill, Florence Rogister, et al. 2014. "MicroRNA Targeting of CoREST Controls Polarization of Migrating Cortical Neurons." *Cell Reports*. https://doi.org/10.1016/j.celrep.2014.03.075.

Wang, Yixuan, Qian Wu, Peng Yang, Chenfei Wang, Jing Liu, Wenyu Ding, Wensu Liu, et al. 2016. "LSD1 Co-Repressor Rcor2 Orchestrates Neurogenesis in the Developing Mouse Brain." *Nature Communications*. https://doi.org/10.1038/ncomms10481.

Weng, Qinjie, Jincheng Wang, Jiajia Wang, Danyang He, Zuolin Cheng, Feng Zhang, Ravinder Verma, et al. 2019. "Single-Cell Transcriptomics Uncovers Glial Progenitor Diversity and Cell Fate Determinants during Development and Gliomagenesis." *Cell Stem Cell*. https://doi.org/10.1016/j.stem.2019.03.006.

Weyn-Vanhentenryck, Sebastien M., Aldo Mele, Qinghong Yan, Shuying Sun, Natalie Farny, Zuo Zhang, Chenghai Xue, et al. 2014. "HITS-CLIP and Integrative Modeling Define the Rbfox Splicing-Regulatory Network Linked to Brain Development and Autism." *Cell Reports*. https://doi.org/10.1016/j.celrep.2014.02.005.

Wohl, Stefanie Gabriele, and Thomas Andrew Reh. 2016. "MiR-124-9-9* Potentiates Ascl1-Induced Reprogramming of Cultured Müller Glia." *GLIA*. https://doi.org/10.1002/glia.22958.

Wu, Thomas D., and Serban Nacu. 2010. "Fast and SNP-Tolerant Detection of Complex Variants and Splicing in Short Reads." *Bioinformatics*. https://doi.org/10.1093/bioinformatics/btq057.

Xue, Yuanchao, Kunfu Ouyang, Jie Huang, Yu Zhou, Hong Ouyang, Hairi Li, Gang Wang, et al. 2013. "Direct Conversion of Fibroblasts to Neurons by Reprogramming PTB-Regulated MicroRNA Circuits." *Cell* 152 (1–2): 82–96. https://doi.org/10.1016/j.cell.2012.11.045.

Yang, Chunzhang, Rajiv R. Iyer, Albert C.H. Yu, Raymund L. Yong, Deric M. Park, Robert J. Weil, Barbara Ikejiri, Roscoe O. Brady, Russell R. Lonser, and Zhengping Zhuang. 2012. "β-Catenin Signaling Initiates the Activation of Astrocytes and Its Dysregulation Contributes to the Pathogenesis of Astrocytomas." *Proceedings of the National Academy of Sciences of the United States of America*. https://doi.org/10.1073/pnas.1118754109.

Yeom, Kyu Hyeon, Simon Mitchell, Anthony J. Linares, Sika Zheng, Chia Ho Lin, Xiao Jun Wang, Alexander Hoffmann, and Douglas L. Black. 2018. "Polypyrimidine Tract-Binding Protein Blocks MiRNA-124 Biogenesis to Enforce Its Neuronal-Specific Expression in the Mouse." *Proceedings of the National Academy of Sciences of the United States of America*. https://doi.org/10.1073/pnas.1809609115.

Yoo, Andrew S., Brett T. Staahl, Lei Chen, and Gerald R. Crabtree. 2009. "MicroRNA-Mediated Switching of Chromatin-Remodelling Complexes in Neural Development." *Nature*. https://doi.org/10.1038/nature08139.

Yoo, Andrew S., Alfred X. Sun, Li Li, Aleksandr Shcheglovitov, Thomas Portmann, Yulong Li, Chris Lee-Messer, Ricardo E. Dolmetsch, Richard W. Tsien, and Gerald R. Crabtree. 2011. "MicroRNA-Mediated Conversion of Human Fibroblasts to Neurons." *Nature*. https://doi.org/10.1038/nature10323.

Yuzwa, Scott A., Michael J. Borrett, Brendan T. Innes, Anastassia Voronova, Troy Ketela, David R. Kaplan, Gary D. Bader, and Freda D. Miller. 2017. "Developmental Emergence of Adult Neural Stem Cells as Revealed by Single-Cell Transcriptional Profiling." *Cell Reports*. https://doi.org/10.1016/j.celrep.2017.12.017.

Zhang, Lei, Jiu Chao Yin, Hana Yeh, Ning Xin Ma, Grace Lee, Xiangyun Amy Chen, Yanming Wang, et al. 2015. "Small Molecules Efficiently Reprogram Human Astroglial Cells into Functional Neurons." *Cell Stem Cell*. https://doi.org/10.1016/j.stem.2015.09.012.

Zhang, Ye, Kenian Chen, Steven A. Sloan, Mariko L. Bennett, Anja R. Scholze, Sean O'Keeffe, Hemali P. Phatnani, et al. 2014. "An RNA-Sequencing Transcriptome and Splicing Database of Glia, Neurons, and Vascular Cells of the Cerebral Cortex." *Journal of Neuroscience*. https://doi.org/10.1523/JNEUROSCI.1860-14.2014.

 Figure legends

Figure 1: miR-124 is sufficient to instruct reprogramming of postnatal cortical astrocytes to
iNs

(A) Overview of the miR-124-mediated reprogramming protocol. **(B)** Immunostaining of
astrocytes reprogrammed with miR-124 at day7 and day14 of the reprogramming protocol
with anti-Tuj1 antibody. **(C)** Quantification of the percentage of Tuj1+ reprogrammed cells
(average ± SD, n=4 independent experiments, **p<0.01 and ***p<0.001 versus sc-miRNA).
RT-qPCR analysis of the mRNA levels of the proneural TFs, *Mash1* and *Neurog2* **(D)**, the dorsal
telencephalon TFs, *Tbr2, Tbr1, Fezf2, Cux1* **(E)**, the ventral telencephalon *TFs, Gsx2* and *Dlx1*
(F), and the neuronal differentiation TFs, *Sox4, Sox11, Hes6* and *NeuroD1* **(G)**. Data are
presented as fold change versus sc-miRNA (average ± SD, n=3 independent experiments,
p<0.01 and ***p<0.001 versus sc-miRNA). **(H) Co-immunostaining of astrocytes
reprogrammed with miR-124 at day7 of the reprogramming protocol with anti-Mash1/Tuj1,
anti-Tbr2/Tuj1 and anti-Gsx2/Tuj1 antibodies. **(I)** Quantification of the percentage of Mash1+,
Tbr2+ and Gsx2+ in Tuj1+ reprogrammed cells (average ± SD, n=3 independent experiments,
**p<0.01 and ***p<0.001 versus sc-miRNA).

Figure 2: The neurogenic compound ISX9 greatly enhances the miR-124-induced
reprogramming efficiency and differentiation state of iNs

(A) Immunostaining of astrocytes reprogrammed with miR-124 or miR-124+ISX9 at day7 and
day14 of the reprogramming protocol with anti-Tuj1 antibody. **(B)** Quantification of the
percentage of Tuj1+ reprogrammed cells at the timepoints day1, day5, day7, day10 and day14
of the reprogramming protocol (average ± SD, n=3 independent experiments for day5 and
day10, n=8 for day7 and n=4 for day14, *p<0.05 and ***p<0.001 versus miR-124). **(C)**RT-qPCR
analysis of the mRNA levels of NeuroD1 at day7 of the reprogramming protocol. Data are
presented as log$_2$(fold change) versus sc-miRNA (average ± SD, n=3 independent experiments,
**p<0.01 and ***p<0.001 versus sc-miRNA). RT-qPCR analysis of the mRNA levels of the TFs,
Neurog2 and Tbr2 (**D**) and Mash1 (**E**) at the time points day1, day5, day7, day10 and day14 of
the reprogramming protocol. Data are presented as log$_2$(fold change) (**D**) and fold change (**E**)
versus astrocytes (day1) (average ± SD, n=3 independent experiments, *p<0.05, **p<0.01 and
***p<0.001 versus miR-124). **(F)** Co-immunostaining of astrocytes reprogrammed with miR-
124+ISX9 at day7 of the reprogramming protocol with anti-Mash1/Tuj1, anti-Tbr2/Tuj1 and
anti-Gsx2/Tuj1 antibodies. **(G)** Quantification of the percentage of Mash1+, Tbr2+ and Gsx2+
in Tuj1+ reprogrammed cells either with miR-124 or miR-124+ISX9 at day7 (average ± SD, n=3

independent experiments, **p<0.01 versus miR-124). Measurement of the mean nuclear

fluorescence intensity of Tbr2 **(H)** and Mash1 **(I)** in Tuj1+ reprogrammed cells either with miR-

124 or miR-124+ISX9 at day7 (co-immunostaining with anti-Tbr2/Mash1/Tuj1 antibodies). A

representative experiment is shown of n=3 independent experiments (mean ± SD, n=326 cells

for miR-124 and n=540 cells for miR-124+ISX9, ***p<0.001 versus miR-124).

Figure 3: miR-124+ISX9-iNs exhibit characteristics of mature, electrophysiologically active

neurons

(A) Co-immunostaining of miR-124+ISX9-iNs at day21 of the reprogramming protocol with

anti-MAP2/Tuj1 and anti-MAP2/Synapsin1antibodies. Inset area indicated in white frame. **(B)**

Quantification of the percentage of Tuj1+ miR-124-iNs and miR-124+ISX9-iNs at day21 of the

reprogramming protocol. The percentage of MAP2/Syn1 double positive (DP) Tuj1+ iNs is

shown in blue (average ± SD, n=3 independent experiments, ***p<0.001 refers to

%MAP2/Syn1 DP in Tuj1+ miR-124+ISX9-iNs versus miR-124-iNs and $^{\#\#\#}$p<0.001 refers to

%Tuj1 miR-124+ISX9-iNs versus miR-124-iNs). **(C)** Co-immunostaining of miR-124+ISX9-iNs at

day28 of the reprogramming protocol with anti-vGlut1/Tuj1 antibodies. **(D)** Quantification of

the percentage of vGlut1+/Tuj1+ and GABA+/Tuj1+ miR-124+ISX9-iNs at day28 (average ± SD,

n=3 independent experiments). **(E)** Superimposed traces of inward Na$^+$ currents and outward

K$^+$ currents evoked by depolarizing voltage steps obtained from a miR-124+ISX9-iN (day22)

(left panel). Superimposed traces of inward and outward currents evoked by the same

protocol after 1 min pre-application of 1μM TTX + 10mM TEA, showing the inhibition of both

inward and outward currents (middle panel), followed by full recovery of the current traces

after 3 min wash of the cell (right panel). **(F)** Superimposed traces of inward Na$^+$ currents and

outward K$^+$ currents evoked by depolarizing voltage steps obtained from a miR-124+ISX9-iN

(day27). **(G)** Example of a repetitive action potential induced from a mature miR-124+ISX9-iN

(day27) by different current steps (injections) at the current clamp mode (the protocol of

current clamp is shown upper right). (**H**) Example of a mature miR-124+ISX9-iN (day27) that

exhibits spontaneous post-synaptic activity at -70 mV holding potential. Representative

traces of ionic currents induced by application of the neurotransmitter GABA in two different

concentrations (300μM and 1mM) obtained from a miR-124+ISX9-iN in the early stage of

neuronal maturation (day22) **(I)** and the neurotransmitter glutamate in two different

concentrations (100μM and 300μM) obtained from a miR-124+ISX9-In in the late stage of

maturation (day27) **(J)**.The cell membrane potential was held at -70 mV and the time of

agonist application is indicated in bars above the traces. **(K)** Superimposed traces obtained

from a mature miR-124+ISX9-iN (day 26) with application of 100 μM glutamate (Glut) or co-

application of 100μM Glut+CNQX indicated that the antagonist CNQX inhibits the AMPA/

kainate glutamate receptor.

Figure 4: RNA-Seq analysis revealed both independent and cooperative transcriptional

contributions of miR-124 and ISX9 in the reprogramming process

(A) Heat map analysis of 300 up- and down-regulated DEGs that belong to the GO terms: Glial

cell differentiation, Gliogenesis, Astrocyte development, Generation of neurons, Neuron

differentiation, Regulation of neuron differentiation, Neurotransmitter transport and Synaptic

signaling. **(B)** GO analysis of biological processes for the up-regulated DEGs in miR-124-iNs vs

astro (in orange) and miR-124+ISX9-iNs vs astro (in red). **(C)** GO analysis of biological processes

for the down-regulated DEGs in miR-124-iNs vs astro. GO terms are ranked according to

$\log_{10}$FDR and the intensity legends indicate the fold enrichment of each GO term. **(D)** Heat

map analysis of 54 up-regulated differentially expressed TFs clustered according to the brain

region they are developmentally expressed (telencephalon, retina, midbrain and hindbrain).

(E) Heat map analysis of 40 up- and down-regulated RBPs.

Figure 5: The RNA-binding protein Zfp36l1 is a novel direct target of miR-124

(A) Venn diagram of the miR-124 targets, derived from AGO-HITS-CLIP and the down-

regulated DEGs of miR-124 vs sc-miRNA astro RNA-Seq data ($\log_2$(fold change)≤-1, FDR<0.01).

Identified miR-124 targets were combined with a public reference list of genes, expressed in

astrocytes, resulting in a set of 130 genes. **(B)** miR-124 direct binding to the 3' UTR of *Zfp36l1*

and *ZFP36L1*, in mouse and human species respectively. miR-124 binds with perfect seed

complementarity (7mer-M8 site) in both species. **(C)** RT-qPCR validation of the mRNA levels

of miR-124 direct target *Zfp36l1* at day7 (average ± SD, n=3 independent experiments,

p<0.01 and ***p<0.001 vs sc-miRNA). **(D) GO analysis of biological processes for Zfp36l1

direct targets that are also significantly up-regulated in miR-124-iNs vs sc-miRNA astro. GO

terms are ranked according to $\log_{10}$FDR and the intensity legend shows the fold enrichment

of each GO term. **(E)** Volcano plot comparing the $\log_2$(fold change) of TPM values in the miR-

124-iNs vs sc-miRNA astro condition versus the $\log_{10}$(FDR) values. Significantly up-regulated

($\log_2$(fold change)≥1, FDR<0.05) and down-regulated ($\log_2$(fold change)≤-1, FDR<0.05) genes

are shown in green and orange respectively. Labels of Zfp36l1 and neuronal-specific up-

regulated genes that are also Zfp36l1 direct targets are portrayed. **(F)** RT-qPCR validation of

the mRNA levels of the Zfp36l1 targets *Elavl4, Nova1, Rbofox1, Rcor2* and *Tox* at day7 (average

± SD, n=3 independent experiments, *p<0.05, **p<0.01 and ***p<0.001 versus sc-miRNA).

Figure 6: Targeting of Zfp36l1 by miR-124 plays a key role in the miR-124-induced cell fate switch of astrocytes to iNs

(A) Schematic representation of the TSB binding region in the 3'UTR of *Zfp36l1* mRNA. **(B)** Western blot analysis of the protein levels of Zfp36l1 in the absence or presence of increasing concentrations of TSB (miR-124:TSB molecular ratio 4:1 and 2:1) at day5 of the reprogramming protocol. Actin protein levels have been used as loading control for normalization of Zfp36l1 protein levels. Normalized Zfp36l1 protein levels in sc-miRNA transfected astrocytes have been set to 1 and the fold change of normalized Zfp36l1 protein levels of each condition vs sc-miRNA is presented in the box below. **(C)** RT-qPCR analysis of the mRNA levels of the Zfp36l1 targets *Tox, Rbfox1, Nova1, Rcor2* and *Elavl4* in the presence or absence of increasing concentrations of TSB (miR-124:TSB molecular ratio 4:1, 2:1 and 1:1) at day3 of the reprogramming protocol (average ± SD, n=3 independent experiments,*p<0.05, **p<0.01 and ***p<0.001 versus sc-miRNA and [#]p<0.05, [##]p<0.01 versus miR-124). (**D**) Immunostaining of astrocytes reprogrammed with miR-124 or miR-124 along with increasing concentrations of TSB (4:1 and 2:1) with anti-Tuj1 antibody at day5 of the reprogramming protocol. **(E)** Quantification of the percentage of Tuj1+ cells at day5 (average ±SD, n=3 independent experiments *p<0.05 and **p<0.01 versus miR-124). **(F)** Morphological characterization of Tuj1+ cells at day5. Quantification of the percentages of multipolar Tuj1+ cells (3 or more processes extending from soma), Tuj1+ cells with 1-2 processes and Tuj1+ cells with an astrocyte-like morphology (rectangular soma with none or 1-2 processes) (average ± SD, n=3 independent experiments, **p<0.01 versus miR-124). **(G)** Proposed model of independent and cooperative transcriptional and post-transcriptional contributions of miR-124 and ISX9 in miR-124/ISX9–induced direct neuronal conversion of cortical astrocytes: miR-124 represses astrocytic identity by down-regulating many astrocytic genes and directly targeting the RBP Zfp36l1, leading to the up-regulation of the direct Zfp36l1 targets and TFs *Tox* and *Rcor2*. Tox is known to up-regulate the cortical IP marker and TF *Tbr2* (Artegiani et al. 2015), which is also further up-regulated by ISX9, along with the neurogenic TFs *Neurog2* and *NeuroD1*. Additionally, miR-124-induced down-regulation of *Zfp36l1* results in the up-regulation of the neuronal-specific RBPs *Nova1* and *Rbfox1*, the latter being further strongly up-regulated by ISX9. These RBPs along with the strongly up-regulated nElavl RBPs *Elavl2, Elavl3* and *Elavl4* by ISX9 contribute to the stabilization and efficient translation of many neuronal mRNAs (J. A. Lee et al. 2016; Scheckel et al. 2016; Weyn-Vanhentenryck et al. 2014). Finally ISX9 transcriptionally up-regulates many neuronal maturation genes leading to the functional maturation of miR-124+ISX9-iNs. Post-

transcriptional events and the RBPs are highlighted in blue, transcriptional events along with

the TFs are shown in green, while down-regulated genes and (blocked) inhibitory mechanisms

are shown in grey, (dashed lines present knowledge from the literature).

Figure 7: miR-124 induces reprogramming of resident reactive astrocytes to iNs with deep-

layer cortical identity *in vivo* following cortical trauma

(A) Experimental setup. **(B)** LV-124-transduced cells in the peritraumatic cortical parenchyma

expressing the mature neuronal marker NeuN, 3 weeks after viral transduction. Inset area

indicated in white frame. Scale bar 50μm. **(C)** Percentage of control LV-GFP and LV-124

transduced cells expressing the mature neuronal marker NeuN, 6 days and 3 weeks post-

transduction with or without treatment with ISX9. LV-GFP 6 days vs 3 weeks: p=0.104, LV-124

6 days vs 3 weeks: p**=0.00116, LV-124 6 days vs LV-124-ISX9 3 weeks: p**=0.00104, LV-GFP

3 weeks vs LV-124 3 weeks: p**=0.0036, LV-GFP 3 weeks vs LV-124-ISX9 3 weeks:

p**=0.00337, LV-124 3 weeks vs LV-124-ISX9 3 weeks: p=0.8427, n=4 for all groups and time

points. **(D)** LV-miR-124-transduced cells in the peritraumatic cortical parenchyma expressing

the deep layer cortical neuronal marker Tbr1 3 weeks after viral transduction. Image from an

animal transduced with LV-miR-124 and co-treated with ISX9. Inset area indicated in white

frame. White arrows indicate NeuN+/Tbr1+ transduced cells. Scale bar 50μm. **(E)** Percentage

of LV-124-transduced cells expressing Tbr1, 3 weeks post-transduction with or without

treatment with ISX9. LV-124 3 weeks vs LV-124-ISX9 3 weeks: p=0.1503, n=4 for both groups.

1248

1249

1250

1251

1252

1253

1254

1255

1256

 Table 1: Sequences of primers for real-time PCR used in this study

Gene	Forward primer	Reverse primer
Mash1	TCTCCGGTCTCGTCCTACTC	CAAAGTCCATTCCCAGGAGA
Neurog2	GTCCCCATACAGCTGCACTT	CAGGTGAGGTGCATAACGGT
Tbr2	CACCCAGAATCTCCTAACACTG	AGCCTCGGTTGGTATTTGTG
Tbr1	TCCAGACGTTCACTTTTCCG	CCCGTGTAGATCGTGTCATAG
Cux1	AGCAGAGACTTTAAGGGAACAG	GCAGCCAACTCTACTTCTAGG
Fezf2	TTTGTGGCAAAGGCTTTCAC	TCTTGTCGTTGTGGGTGTG
Gsx2	GATTCCACTGCCTCTCCATG	CGGGACAGGTACATATTGGAAG
Dlx1	CAGTTCCGTGCAGTCCTAC	ATTGTCCTGGGTTTACGGATC
Sox4	GAACGCCTTTATGGTGTGGT	GAACGGAATCTTGTCGCTGT
Sox11	CCCTGTCGCTGGTGGATAAG	GGTCGGAGAAGTTCGCCTC
Hes6	TACCGAGGTGCAGGCCAA	AGTTCAGCTGAGACAGTGGC
NeuroD1	TTGAAGCCATGAATGCAGAG	TCTTGGGCTTTTGATCATCC
Scrt1	AATCATGCCCAGGTCCTTC	CCACGTAGTCACTGAGGTATC
Lhx6	GACACCATGATCGAGAACCTC	CAATTGCTCTGCGGTGAAG
Chd5	ATCTACGAAATCTGGCACCG	CCCTTGTGGATCTCAGACTTG
En1	CTACTCATGGGTTCGGCTAAC	TCTTTAGCTTCCTGGTGCG
Foxa1	AGGGTTGGATGGTTGTGTC	AGGCCGGAGTTCATGTTG
Lmx1b	CGGGATCGGAAACTGTACTG	AGCAGAAACAGCCCAAGTG
Hoxc4	CAAGCAACCCATAGTCTACCC	AACTCTTTCTCTAATTCCAGGACC
Phox2a	TCCCTTCTCTGGAGTTCTGTC	GATAATGCCAGGTCCAGAAGG
Elavl2	AATAACAGGGCAGAGCTTGG	TCTGATTGAGGCTGAGCTTG
Elavl3	GGTTCGGGATAAGATCACAGG	CAGAACTGGGACGTGCATAG
Elavl4	AGAATCCTGCAAACTCGTGAG	ATGGTTTTGGTCTGGAGTCTG
Nova1	GCTGGCTACCTCTGGATCAT	TGGGATGCCATTTAGCTTGC
Zfp36l1	CACACCAGATCCTAGTCCTTG	CTGGGAGTGCTGTAGTTGAG
Zfp36	TCTCTTCACCAAGGCCATTC	GAGTCCGAGTTTATGTTCCAAAG
Rbpj	TCCCAAAACCCGGATAACC	TTTCGCATAGCTTCCCTAGTG
Tcf7l1	CTACAGCAACGACCACTTCTC	GGTAATACGGTGACAGCTCAG

Tcf4	CACAAACCATTACAGCACCTC	GTGTGGTCAGGAGAATAGATCG
Nfic	GGACGGAAGACATAGAAGGAG	GGGCTGTTGAATGGAGATTTG
Tox	TGCTCTCCAATTCCATCTCTG	CTGTCTGATGTCTGTAGGCTG
Rcor2	GCATGTACCTGAGTCCTGAAG	CTGCTATTGGTCTGCTTCATG
Rbfox1	AGTTACGGACGAGTTTATGCTG	AGAGAACGAGACCCACATCA

1258

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

AGO−CLIP−Seq miRNA targets

RNA−Seq downregulated genes

1426

8547

Astrocyte genes

B

Mouse
chr12:80109221-80109245

mmu-miR-124-3p

5'- U   G   C G G U G A A U G C C- 3'
AAGGCAC
| | | | | | |
UUCCGUG
3'- G   A   U U C A A C G A A G A C A U U U -5'

Zfp36l1

Human
chr14:68789157-68789181

hsa-miR-124-3p

5'- U   G   C G G U G A A U G C C A A- 3'
AAGGCAC
| | | | | | |
UUCCGUG
3'- G   A   U U C A A C G A A G A C A U U U -5'

ZFP36L1

C

Zfp36l1

D

GO biological processes for Zfp36l1 mRNA targets

neurogenesis
neuron projection development
regulation of trans−synaptic signaling
neuron differentiation
dendrite morphogenesis
axonogenesis
neurotransmitter transport
telencephalon development
glutamate receptor signaling pathway
pallium development
regulation of transcription. DNA−templated
regulation of RNA metabolic process
cerebral cortex cell migration

−log10(FDR)

2 4 6 8 10

Fold Enrichment

2 4 6

E

F

Zfp36l1 mRNA targets

Figure 6

Figure 7

A

Cortical trauma   LV-124/ LV-GFP injection   Sacrifice

ISX9 20 mg/kg

Day 0   Day 4   Day 6   Day 10   Day 25

B   GFP/NeuN/Hoechst

50 μm

C   NeuN

% NeuN+ GFP+/GFP+ cells

LV-GFP   LV-miR-124   LV-GFP   LV-miR-124   LV-miR-124 +ISX9

6 days   3 weeks

D   GFP/Tbr1/NeuN

50 μm

E   Tbr1

% Tbr1+NeuN+GFP+/NeuN+GFP+cells

LV-miR-124   LV-miR-124 +ISX9