Mechanisms of antibiotic action shape the fitness landscapes of resistance mutations

Colin Hemez[1,2]*, Fabrizio Clarelli[3,4], Adam C. Palmer[5], Leonid Chindelevitch[6], Theodore Cohen[7], and Pia Abel zur Wiesch[3,4,8]*

[1] Systems Biology Institute, Yale University, West Haven, CT USA 06516
[2] Department of Molecular, Cellular, and Developmental Biology, Yale University, New Haven, CT USA 06520
[3] Institute of Pharmacy, Arctic University of Norway, Tromsø, Norway 9017
[4] Center for Infectious Disease Dynamics, Department of Biology, Pennsylvania State University, University Park, PA USA 16802
[5] Department of Pharmacology, Computational Medicine Program, Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel Hill, NC USA 27599
[6] School of Computing Science, Simon Fraser University, Burnaby, BC, Canada V5A 1S6
[7] Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven, CT USA 06520
[8] Centre for Molecular Medicine Norway, Nordic EMBL Partnership, Oslo, Norway 0318
* Correspondence to colin.hemez@yale.edu, pia.z.wiesch@uit.no

Abstract

Antibiotic-resistant pathogens are a major public health threat. Understanding how an antibiotic's mechanism of action influences the emergence of resistance could help to improve the design of new drugs and to preserve the effectiveness of existing ones. To this end, we developed a model that links bacterial population dynamics with antibiotic-target binding kinetics. Our approach allows us to derive mechanistic insights on drug activity from population-scale experimental data and to quantify the interplay between drug mechanism and resistance selection. We find that whether a drug acts as a bacteriostatic or bactericidal agent has little influence on resistance selection. We also show that heterogeneous drug-target binding within a population enables antibiotic-resistant bacteria to evolve secondary mutations, even when drug doses remain above the resistant strain's minimum inhibitory concentration. Our work suggests that antibiotic doses beyond this "secondary mutation selection window" could safeguard against the emergence of high-fitness resistant strains during treatment.

(word count: 150)

Introduction

The emergence and spread of antibiotic-resistant bacterial pathogens is an urgent global problem that threatens to undermine one of the most essential components of modern medicine (WHO, 2012). Antibiotic resistance is also expensive, adding an average of US $1400 to the costs of treatment for each of the 2.8 million patients who become infected with a drug-resistant bacterium in the United States annually (CDC, 2019; Thorpe, Joski, & Johnston, 2018; WHO, 2014). The scarcity of promising new antimicrobial drugs with novel mechanisms of action further exacerbates the challenges associated with managing the spread of drug resistance (Roberts, Kruger, Paterson, & Lipman, 2008; Silver, 2011). Given the increasing incidence of resistant bacterial infections and the lack of new drugs on the horizon, clinicians, researchers, and global leaders must act to preserve the effectiveness of the world's existing antibiotic drug arsenal (WHO, 2012).

Antibiotic treatment induces a strong selective pressure on bacterial populations to evolve resistance (Hughes, 2014; Rao, 1998). Resistance mutations raise the minimum inhibitory concentration (MIC) of an antibiotic, the amount of drug needed to suppress the growth of a bacterial culture (Andrews, 2001). However, alleles that confer drug resistance also frequently carry fitness costs (Andersson & Hughes, 2010; Melnyk, Wong, & Kassen, 2015; Vogwill & MacLean, 2015), predominantly because antibiotics target vital cellular functions (such as DNA replication and protein synthesis). Resistance mechanisms reduce the ability of a drug to disrupt its target, but do so at the expense of optimal physiological function (Lovmar et al., 2009).

With few exceptions (Engelberg & Artman, 1964), resistance-causing alleles induce physiological impairments in both drug-free and drug-containing environments, though resistant strains may only suffer a strict competitive disadvantage (i.e. a slower growth rate)

against sensitive strains in drug-free conditions. A range of antibiotic concentrations therefore

exists within which drug-resistant strains have a selective advantage over their drug-

susceptible counterparts. Dosing drugs within this "resistance selection window" (also called

the "mutant selection window") can allow for the proliferation of drug-resistant

subpopulations (Karl Drlica & Zhao, 2007; Gullberg et al., 2011; Yu, Baeder, Regoes, & Rolff,

2018). Recent advances in antimicrobial pharmacodynamics have leveraged the characteristics

of resistance selection windows to design dosing strategies that minimize the selection of

resistant pathogens without sacrificing treatment efficacy (Cui et al., 2006; Mohamed, Cars, &

Friberg, 2014; Yu et al., 2018).

The existence of resistance mutations that confer physiological impairments in both

drug-free and drug-containing environments implies that resistant strains face selective

pressures to evolve secondary mutations that alleviate these impairments, and that these

selective pressures exist even under continuous drug exposure (Loftie-Eaton et al., 2017;

Maisnier-Patin, Berg, Liljas, & Andersson, 2002). Secondary mutations can increase bacterial

fitness (through faster growth rates) in the absence of drugs, or they can confer elevated levels

of drug tolerance to preexisting resistant subpopulations (through attenuated drug-target

interactions, faster growth rates in the presence of drugs, or both). In the case of increased

bacterial fitness, secondary mutations enable drug-resistant mutants to compete against drug-

susceptible strains in resource-limited, antibiotic-free environments (Andersson & Hughes,

2010; Durão, Balbontín, & Gordo, 2018; Levin, Perrot, & Walker, 2000), and are implicated in

the spread of drug resistance across a wide range of timescales and clinical settings (Handel,

Regoes, & Antia, 2006). In the case of increased drug tolerance, secondary mutations can be

the underlying cause of treatment failure (Ahn et al., 2015; Merker et al., 2018). Elucidating the

dynamics of secondary mutation emergence during treatment is thus crucial for managing the

spread of resistance.

Since resistance mutations are frequently associated with fitness costs (Melnyk et al.,

2015; Vogwill & MacLean, 2015) both *in vivo* (Majcherczyk, Barblan, Moreillon, & Entenza,

2008) and *in vitro* (Zhang, Sahin, McDermott, & Payot, 2006), studies on the resistance

selection window and on secondary adaptation have yielded valuable insights into the

emergence of drug-resistant bacteria during treatment. However, the design of optimal

resistance-mitigating drug dosing strategies remains challenging for two reasons. One

obstacle is that bacteria may acquire resistance through a multitude of mechanisms that

reduce antibiotic efficacy (Blair, Webber, Baylay, Ogbolu, & Piddock, 2015). These molecular

mechanisms may themselves influence the fitness landscape of resistance mutations (that is,

the relationship between the fitness cost of resistance and the selective advantage conferred

by the resistance mutation in drug-containing environments). A second challenge is that an

antibiotic's mechanism of action may affect the strength of selection for resistant strains over

drug-susceptible strains during treatment. One important feature of an antibiotic's cellular-

level mechanism of action is whether the drug controls bacterial populations by increasing

the rate of bacterial killing (i.e. bactericidal action) or by decreasing the rate of bacterial

replication (i.e. bacteriostatic action). Clinicians and researchers alike have argued that these

modes of antimicrobial action influence the dynamics of resistance selection (Frenoy &

Bonhoeffer, 2018; Stratton, 2003).

The design of resistance-mitigating antibiotic usage therefore depends on an

understanding of how a drug's mechanism of action, a pathogen's mechanism of resistance,

and the fitness landscape of resistance affect selection pressures during treatment. Tractable

and quantitative strategies for systematically exploring all of these factors have so far been

lacking. To address this gap, we developed a dynamical model that simulates the growth and

death of bacterial populations under antibiotic exposure using molecular-scale descriptions of

drug-target binding kinetics and cellular-scale descriptions of a drug's mechanism of action.

In our model, higher numbers of inactivated drug-target complexes within a cell lead to

increases in antibiotic effect (either bacteriostatic, bactericidal, or a combination of the two).

The relationship between drug-target inactivation and antibiotic effect can take the shape of a

linear (i.e. gradual) or stepwise (i.e. sudden) function, as well as other intermediate forms

(**Supplementary Figure S1**). The model enables us to estimate critical pharmacodynamic

parameters from experimental datasets as effectively as with classical approaches (Regoes et

al., 2004), to simulate the fitness landscapes of resistance mutations against drugs with diverse

mechanisms of action, and to quantify the probability of secondary mutation emergence

within resistant subpopulations of bacteria during treatment.

We find that bacteria with resistance mechanisms that confer even modest reductions

in drug-target binding affinity can incur strikingly high (80-99%) fitness costs while still

maintaining higher drug tolerances than their susceptible counterparts, regardless of the

antibiotic's mechanism of action. We also find that drugs with stepwise effects on bacterial

growth and death have narrower resistance selection windows than do drugs with linear

effects. However, our model suggests that whether a drug acts primarily through bactericidal

or bacteriostatic action has comparatively little influence on the strength of resistance

selection during treatment. We further demonstrate that, even with aggressive treatment

regimens, heterogeneous drug-target occupancy within a population enables fitness-impaired

resistant strains to develop secondary mutations that can lead to treatment failure. Our work

cautions that fitness costs may not limit the emergence of resistant strains that evolve through

reductions in drug-target binding affinity. We propose the "secondary mutant selection

window" as a novel pharmacodynamic characteristic of a drug that should be assessed

alongside other classic parameters such as the MIC and the resistance selection window when

designing robust resistance-mitigating antibiotic dosing strategies.

Results

A model that links bacterial population dynamics with molecular mechanisms of antibiotic drug action

We developed a linear dynamical model to describe the effect of a constant concentration of

drug on the growth and death rates of a bacterial population (**Figure 1A**) (see **Methods**, *Model*

formulation and analysis for a mathematical description of the model). We assume that each

bacterial cell in the population carries an identical number N of intracellular proteins that the

drug targets for inactivation. Drug molecules inactivate target proteins by binding to them

with a rate k_F and can dissociate from the target with a rate k_R. The affinity K_D of the drug is

thus the ratio of off-rate to on-rate, $K_D = k_R/k_F$. The model assumes that the growth and death

rates of a bacterial cell depend on its drug-target occupancy (that is, the number of inactivated

drug-target complexes it contains) (Clarelli et al., 2019; Wiesch et al., 2015). We denote drug-

target occupancy with the index i, which ranges from 0 to N. Cells harboring successively

larger numbers of inactivated drug-target complexes have successively faster death rates

and/or slower growth rates, depending on the mechanism of action of the drug (see **Results**,

Classification of drug action). We thus define the growth rate ($G[i]$) and death rate ($D[i]$) of each

subpopulation as discrete monotonic functions of drug-target occupancy. In practice, $G[i]$ and

$D[i]$ take the form of constrained logistic functions each controlled by a steepness and

inflection point parameter, allowing us to define quasi-linear, quasi-stepwise, quasi-

exponential, and sigmoid curves (**Supplementary Figure S1**).

Figure 1 – Features of a model that links bacterial population dynamics with the cellular mechanisms of antibiotic drug action. (**A**) Illustration of the model. We consider a population B_i of bacterial cells harboring i inactive drug-target complexes. The change in the size of B_i is a function of cellular growth and death rates (each of which is determined by the value of i), and of the molecular kinetics of the drug binding and unbinding to its protein target. The total bacterial population is given by the sum $B_0 + B_1 + ... + B_{N-1} + B_N$, where N is the number of drug targets per cell. (**B**) Dynamics of a bacterial population exposed to a drug dose above the minimum inhibitory concentration (MIC). The black line represents the total bacterial population; shaded lines represent subpopulations with x and fewer inactivated drug-target complexes. (**C**) Proportion of the bacterial subpopulation B_i as a share of total

population for the first three hours of the curve shown in panel (B). (**D**) Pharmacodynamic curves derived from the model for a wild-type (light green) and drug-resistant (dark green) bacterial strain. The MIC is denoted as the drug concentration at which the net bacterial growth rate is zero. Inset: the resistance selection window (green shading) is given by the drug concentration range within which the drug-resistant strain exhibits a higher—but still positive—net growth rate compared to the wild-type strain. G_o denotes the growth rate of the wild-type strain in the absence of antibiotic (i.e. the growth rate for subpopulation B_o). D_N denotes the maximum death rate of bacterial strains when all N cellular targets are inactivated (i.e. the death rate of subpopulation B_N).

____

The model tracks the growth and death of all $N+1$ bacterial subpopulations, each denoted B_i, over time (**Figure 1B**). Drug concentration determines the net growth rate of the entire bacterial population (**Supplementary Figure S2**). In the absence of drug, the population grows exponentially at a rate equal to the difference between the drug-free growth and death rates (G_o and D_o, respectively). When drug is present, the composition of bacterial subpopulations asymptotes towards a steady state after a transient phase during which drug molecules bind to their targets (**Figure 1C**). At steady state, the relative composition of bacterial subpopulations does not depend on the total size of the population.

We can calculate the MIC of a drug directly from model parameters (see **Methods**, *Calculation of the minimum inhibitory concentration*), and we can simulate clinically observed drug resistance mutations by modulating the parameters of the model that influence the value of the MIC. Changes in the binding kinetics of the drug (i.e. k_F and k_R) simulate target modification mutations that decrease the affinity of an antibiotic molecule to a cellular protein (Billal, Feng, Leprohon, Légaré, & Ouellette, 2011; Everett, Jin, Ricci, & Piddock, 1996; Gao et al., 2010). Changes to the value of N represent changes in the number of protein targets per cell, equivalent to target up- or downregulation (Brochet, Couvé, Zouine, Poyart, & Glaser, 2008; Palmer, Chait, & Kishony, 2018; Palmer & Kishony, 2014). We assume that fitness costs associated with resistance alleles take the form of reduced growth rates, and we simulate this cost by reducing the drug-free growth rate of resistant strains by a factor c_R such that the

maximum growth rate of a resistant strain ($G_{o,RES}$) relative to that of a wild-type strain is $G_{o,RES}$

$= G_o(1-c_R)$. When c_R ranges from 0 (no cost) to 1 (no growth whatsoever), the resistant strain

exhibits a slower growth rate relative to that of the wild-type. If c_R is negative, the resistant

strain exhibits a faster drug-free growth rate than does the wild-type strain, as has been

observed in rare cases with some fluoroquinolone-resistant *Escherichia coli* isolates (Lindgren,

Marcusson, Sandvang, Frimodt-Møller, & Hughes, 2005). The model also enables us to

generate pharmacodynamic curves by calculating the net growth rates of simulated bacterial

populations over a range of drug concentrations (**Figure 1D**). The resistance selection window

constitutes the range of drug concentrations over which a drug-resistant mutant strain has a

higher but strictly positive net growth rate relative to that of its wild-type counterpart (**Figure

1D**, inset).

Inferring cellular mechanisms of antibiotic action from population-scale data

To test the utility of our biochemical model for gaining cellular-scale insights into

antimicrobial drug mechanisms from population-scale experiments, we calibrated our model

to a family of time-kill curves of the gram-negative bacterium *Escherichia coli* challenged to

ciprofloxacin, a fluoroquinolone first brought to market in 1987. Ciprofloxacin has two known

molecular targets in bacteria, both of which are heterotetrameric type-II topoisomerases: the

DNA gyrase complex ($GyrA_2B_2$) and DNA topoisomerase IV ($ParC_2E_2$). However, ciprofloxacin

preferentially binds to the $GyrA_2B_2$ complex in gram-negative bacteria (Karl Drlica, Malik,

Kerns, & Zhao, 2008). We used a mass-spectrometry based estimate for the number of

$GyrA_2B_2$ complexes per *E. coli* cell ($N \sim 183$) as the number of drug targets within each

bacterium (Wiśniewski & Rakus, 2014).

We implemented an adaptive simulated annealing algorithm to calibrate the

parameters of our model to an experimental dataset of ciprofloxacin time-kill curves

(**Methods,** *Model calibration via simulated annealing*). We performed 249 independent

parameterizations using the algorithm and selected the parameter set that yielded the lowest

objective function value (**Figure 2A, Table 1, Supplementary Figure S3**). Bacterial persistence

(Dörr, Lewis, & Vulić, 2009; Harms, Maisonneuve, & Gerdes, 2016) likely plays a role in the

slower-than-expected population decline that we observe experimentally at high drug

concentrations. At antibiotic doses below those that elicit persistence, the calibrated model

accurately recapitulates the pharmacodynamic curve derived from experimental data

(**Supplementary Figure S4**).

214
215
216   **Figure 2 – Calibrating the model to experimental data reveals underlying mechanisms of**
217   **drug action.** (**A**) Comparison between calibrated biochemical model (solid lines) and
218   experimental data (shaded points). The experimental data (**Supporting Data File 1**) represent

time-kill curves of *Escherichia coli* exposed to ciprofloxacin. (**B**) Comparison of calibrated biochemical model the E_{MAX} pharmacodynamic model (Regoes et al., 2004). We fit the E_{MAX} model to the same experimental dataset shown in panel (A) and compared correlation coefficients and MICs. Red points in the MIC panel denote experimentally-measured ciprofloxacin MICs for *E. coli* strains isolated prior to the widespread emergence of quinolone resistance (see **Supporting Data File 2**). The solid horizontal line represents the mean of experimental measurements, and the dashed lines indicate the 95% confidence interval. (**C**) Cellular growth and death rates as a function of ciprofloxacin-GyrA$_2$B$_2$ complex number (*i*) for the model calibrated to the experimental data shown in panel (A). (**D**) Four extreme schemes of drug action resulting from two characteristics (activity and steepness) of a drug's effect on growth and death rates as a function of drug-target occupancy. **Supplementary Figure S5** shows the simulated bacterial kill curves for these schemes at 4x MIC.

____

Model parameters				
Name	**Description**	**Value**	**Units**	**Source**
N	Number of target proteins per cell (i.e. GyrA$_2$B$_2$ copy number)	183	cell^{-1}	(Wiśniewski & Rakus, 2014)
G_0	Bacterial growth rate in the absence of drug	0.526	hr^{-1}	Model calibration
D_0	Bacterial death rate in the absence of drug	5.40×10^{-3}	hr^{-1}	(Wang et al., 2010)
D_N	Bacterial death rate in saturating concentrations of drug	7.53	hr^{-1}	Model calibration
k_F	Drug-target binding rate	5.23×10^3	M^{-1} sec^{-1}	Model calibration
k_R	Drug-target unbinding rate	3.17×10^{-4}	sec^{-1}	Model calibration
α_G	Steepness of growth rate function G[*i*]	16.8	# drug-target complexes^{-1}	Model calibration
α_D	Steepness of death rate function D[*i*]	7.29	# drug-target complexes^{-1}	Model calibration
γ_G	Inflection point of growth rate function G[*i*]	24.9	# drug-target complexes	Model calibration
γ_D	Inflection point of death rate function D[*i*]	359	# drug-target complexes	Model calibration
B_0	Initial size of bacterial population at the start of drug treatment	6.88×10^9	cell ml^{-1}	Model calibration
μ_R	Mutation rate for drug resistance emergence	2.00×10^{-7}	cell^{-1} division^{-1}	(Martinez & Baquero, 2000; Schulz zur Wiesch, Engelstädter, & Bonhoeffer, 2010)
μ_C	Mutation rate for emergence of secondary mutations in resistant strains	2.00×10^{-6}	cell^{-1} division^{-1}	(Martinez & Baquero, 2000; Schulz zur Wiesch et al., 2010)
c_R	Cost of resistance mutation, such that the antibiotic-free growth rate of a resistant mutant is $G_0(1 - c_R)$	0.25	Non-dimensional	(Gagneux et al., 2006)

Table 1 – Model parameters. We obtained the values of k_F, k_R, α_G, α_D, γ_G, γ_D, and B_o by calibrating the model to experimental data (**Figure 2**). We inferred antibiotic-free growth rate and antibiotic-saturated death rate (G_o and D_N) by fitting an exponential curve to ciprofloxacin kill curves using 0 and 2.19 μg/ml of drug, respectively (**Supplementary Figure S11**). We use a constrained logistic function to model the growth and death rates of bacterial

cells as a function of bound target number, where α controls the steepness of the logistic
function and γ controls the inflection point of the logistic function (**Supplementary Figure**
S1). Parameters not obtained from model calibrations to experimental data were retrieved
from the literature. For the bacterial death rate in the absence of drug (D_o), we used the mean
of death rates reported in Wang et al., 2010.
____

We compared our biochemical model's ability to capture critical pharmacodynamic

characteristics of a drug against that of an E_{MAX} model (Regoes et al., 2004). The E_{MAX} approach

describes net bacterial growth rate directly as a function of drug concentration and does not

accommodate molecular descriptions of drug-target interactions. Such models have been

used extensively to estimate pharmacodynamic parameters, to design drug dosing regimens,

and to predict the strength of resistance selection at nonzero drug concentrations. Our

formulation delivers performance comparable to that of the E_{MAX} model for fitting

experimental time-kill curves (**Figure 2B**, left panel) and more accurately estimates MIC

(which we determined to be 8.9×10^{-3} μg/ml for ciprofloxacin) from these data (**Figure 2B**, right

panel). This demonstrates the validity of our approach for deriving pharmacodynamic

insights similar to what an E_{MAX} model provides.

Our model furthermore offers capabilities that the E_{MAX} approach lacks, including the

ability to estimate molecular kinetic parameters of drug-target binding from population-scale

data. To test the robustness of these estimates, we analyzed the K_D values for ciprofloxacin

binding to *E. coli* GyrA$_2$B$_2$ generated for the 249 independent parameterizations described

above. The best 90% of all calibrations (that is, the 224 fits with the lowest objective function

values) consistently converged upon a narrow range of affinity values (95% confidence

interval: 7.2×10^{-8} to 1.6×10^{-7} M) (**Supporting Data File 3**). Our estimates lie within the range of

K_D values of ciprofloxacin for *E. coli* GyrA$_2$B$_2$ reported from experimental measurements,

which span from 3.2×10^{-8} to 3.0×10^{-6} M (Jungkind & American Society for Microbiology

Eastern Pennsylvania, 1995; Kampranis & Maxwell, 1998; Shen & Pernet, 1985; Siporin, Heifetz,

& Domagala, 1990).

Classification of antibiotic action

Another unique feature of our approach is the ability to describe bacterial growth and death

rates as a function of drug-target occupancy. For ciprofloxacin, the calibrated model predicts

three regimes of bacterial subpopulation dynamics in relation to $GyrA_2B_2$ inactivation: a

growth regime in which bacterial replication dominates among subpopulations with low

numbers of inactivated targets, a stalling regime for intermediate numbers of drug-target

complexes in which neither growth nor death is appreciable, and a killing regime at high

numbers of inactivated targets in which bacterial death increases quasi-exponentially (**Figure**

2C). The forms of G[*i*] and D[*i*] that we obtain here suggest that ciprofloxacin has a

multimodal mechanism of action, a result consistent with prior experimental studies (K.

Drlica, 1999; Karl Drlica et al., 2008; Silva, Lourenço, Queiroz, & Domingues, 2011). The drug

stalls cellular replication at intermediate target occupancies and induces killing only at higher

doses. Like many antibiotics, ciprofloxacin thus exhibits both bactericidal and bacteriostatic

effects on microbial populations (Pankey & Sabath, 2004; Silva et al., 2011). Our biochemical

model represents this explicitly.

Most drugs nonetheless demonstrate a greater degree of bactericidal or bacteriostatic

activity at clinically relevant doses (Nemeth, Oesch, & Kuster, 2015), and we hypothesized that

the ability of a drug to stall growth or to accelerate death may affect the selection for resistant

strains and the emergence of secondary mutations. We also suspected that the relationship

between drug-target occupancy and antibiotic effect—reflected in the steepness of the *G*[*i*]

and *D*[*i*] functions—could further shape the dynamics of resistance selection.

These two characteristics (bactericidal versus bacteriostatic activity and drug effect steepness) represent two general dimensions along which a drug's mechanism of action can affect the growth and death of bacterial populations. Four extreme cases of drug action thus exist (**Figure 2D**). In the case of a purely bacteriostatic antibiotic, death rates are a constant function of inactivated drug-target complex number (that is, $D[i] = D_o$ for all values of i). For a purely bactericidal antibiotic, the growth rate of all bacterial subpopulations remains constant ($G[i] = G_o$ for all values of i). The steepness of the drug effect is reflected in the form of the function $D[i]$ for bactericidal antibiotics and $G[i]$ for bacteriostatic antibiotics (**Supplementary Figure S1**). We defined linear and stepwise onset of action as our two extremes, as other monotonic forms are intermediate cases of these curves.

The opposing effects of increased drug resistance and decreased cellular fitness

Mutations that confer resistance against antibiotics often come at the cost of reduced growth rates compared to those of drug-susceptible strains (Andersson & Hughes, 2010; Melnyk et al., 2015). The balance of replication and death determines bacterial net growth both in the absence and in the presence of antibiotics, and very high fitness costs associated with resistance can prevent bacterial viability at any drug concentration (Björkman, Nagaev, Berg, Hughes, & Andersson, 2000). We sought to investigate the quantitative basis for the trade-off between drug resistance and cellular growth and to investigate how the drug mechanisms defined above influence the range of permissible fitness costs that a drug-resistant mutant can incur while still maintaining a drug susceptibility that is lower than that of a wild-type strain. In the simplest case of the model, where the number of target molecules per cell is 1, the expression for the MIC captures the opposing effects of drug resistance and cellular growth (see **Methods**, *Calculation of minimum inhibitory concentration* for derivation):

[**Equation 1**]

$$\text{MIC} = \frac{(k_R + D_N)}{k_F D_N} G_0$$

The MIC increases with reductions of the on-rate kinetics of drug-target binding (k_F) and with

increases in the off-rate kinetics of drug-target binding (k_R), but decreases with fitness costs

that manifest as reductions in the drug-free growth rate (G_o). These proportionalities hold for

any number N of drug targets.

We modeled the opposing effects of biochemical changes that reduce drug

susceptibility (i.e. altered drug-target binding kinetics or target upregulation) and the fitness

costs of these biochemical changes. We considered a set of five antibiotics with an identical

protein target and identical molecular kinetic parameters (that is, the target number N, the

drug-target on-rate k_F, and the drug-target off-rate k_R are constant for the wild-type strain)

(**Supplementary Table S1**, **Supplementary Figure S5**). One antibiotic in the set features

growth and death dynamics derived from the model calibration to ciprofloxacin time-kill

curve data (**Figure 2C**). The other four antibiotics are hypothetical and feature growth and

death dynamics that represent four extremes of antibiotic action (**Figure 2D**). We simulated

mutant strains of *E. coli* that acquire drug resistance phenotypes either through changes in the

molecular kinetics of drug binding (k_F or k_R) or by increasing the copy number N of the drug's

cellular protein target. Each of these resistance mechanisms has been observed in clinical

isolates of drug-resistant, gram-negative bacteria (Blair et al., 2015; Melnyk et al., 2015;

Redgrave, Sutton, Webber, & Piddock, 2014). We then simulated fitness costs associated with

the resistance mutation and calculated the mutant strain's MIC relative to that of the wild-

type strain.

For resistance acquired through changes in the kinetics of drug-target binding (k_F and

k_R), we found that mutants can tolerate strikingly high (80-99%) fitness costs while still

maintaining an MIC that is greater than that of the drug-susceptible wild-type (**Figure 3**, top

and middle rows). This permissibility of fitness costs exists for all five of the drug mechanisms

we simulated, although drugs that act with linear effects (Bacteriostatic/Linear and

Bactericidal/Linear) have a narrower range of permissible fitness costs than do drugs that act

with stepwise effects. For all drug mechanisms, mutant strains make larger gains in MIC by

decreasing the on-rate kinetics of drug-target binding (k_F) than they do by increasing the off-

rate kinetics of drug-target binding (k_R) by the same amount (**Supplementary Figure S6**). That

is, mutations that lead to the same change in drug-target affinity (as quantified by the

dissociation constant $K_D = k_R/k_F$) through different changes in the on- and off-rate binding

kinetics do not necessarily have the same range of permissible fitness costs. This has

biological significance—limiting the opportunity for a drug to bind to its target, thereby

preventing the drug from actuating its effects on cellular growth and death, should lead to

lower drug susceptibilities than would accelerating the rate at which an already-formed drug-

target complex disassociates. The difference in the fitness effects of mutations that modify k_F

and k_R is especially pronounced for bactericidal drugs that elicit linear increases in cellular

death (Bactericidal/Linear).

Figure 3 – Drug mechanism influences the fitness landscapes of resistance mutations. We calculated the MIC, expressed as a fold-change relative to the MIC of the wild-type, for mutant strains carrying (top row) drug targets with reduced binding kinetics (k_F), (middle row) drug targets with accelerated unbinding kinetics (k_R), or (bottom row) increased numbers of drug target molecules (N). Mutant strains also carry fitness costs, expressed as a fractional reduction in drug-free growth rate relative to wild-type. When modulating the number of drug target molecules N (bottom row), we assumed that cells require a fixed number of active protein targets to grow at a normal rate and that cellular killing is induced when a fixed number of inactive drug-target complexes form within a cell. Thus, the inflection point for the growth rate function (γ_G) changes concomitantly with N such that N-γ_G remains constant, while the inflection point for the death rate function (γ_D) remains constant (see **Supplementary Figure S1** for illustrations of the effects of γ_G and γ_D on bacterial growth and death rates).

____

Ciprofloxacin exhibits a bactericidal effect by permitting GyrA$_2$B$_2$-mediated cleavage

of DNA but preventing DNA re-ligation, resulting in widespread and eventually

insurmountable chromosome fragmentation (Karl Drlica et al., 2008; Pan, Yague, & Fisher,

2001). When simulating the overexpression of target proteins in resistant cells (**Figure 3,**

bottom row) we therefore assumed that bacterial killing is induced when a fixed number of

inactivated drug-target molecules form within a cell (that is, we assume a toxicity threshold

whereby γ_D remains constant with changing N). Conversely, we assumed that a resistant cell

requires a fixed number of active, non-complexed target proteins in order to maintain its

maximum growth rate (that is, a survival threshold). γ_G thus changes in step with N such that

N-γ_G remains constant. We made these same assumptions for the four hypothetical

antibiotics.

We found that target overexpression has a diversity of effects on resistance that

depend on the mechanism of action of the drug. For ciprofloxacin and its multimodal effects

on growth and death, small increases in target number can lead to modest increases in MIC,

even when the resistant cell faces large fitness costs as a result of $GyrA_2B_2$ overexpression.

However, larger increases in target number lead to reductions in MIC. This result is consistent

with experimental studies on target amplification, in which the overexpression of *gyrAB* in *E.*

coli resulted in increased susceptibility to ciprofloxacin (Palmer & Kishony, 2014). Target

overexpression leads to substantial gains in resistance against bacteriostatic drugs that exhibit

stepwise effects, even at very high fitness costs. The effect of target overexpression on drug

resistance is negligible for bactericidal drugs and for bacteriostatic drugs with a linear effect

on growth stalling.

Drug mechanism shapes the resistance selection window

To understand how a drug's mechanism of action affects the propensity to select for resistance

during treatment, we simulated the pharmacodynamics of wild-type and drug-resistant

strains challenged to each of the five drugs in the set outlined above. MICs for clinical isolates

of ciprofloxacin-resistant *E. coli* strains with single point mutations in GyrA, which may

reduce the affinity of ciprofloxacin to $GyrA_2B_2$, range from 10 to 16 times greater than the MIC

of a drug-susceptible wild-type (Everett et al., 1996; Morgan-Linnell & Zechiedrich, 2007;

Piddock, 1999; Redgrave et al., 2014). Data on the fitness costs associated with mutant GyrA-

mediated ciprofloxacin resistance in *E. coli* are sparse, but studies of rifampicin-resistant

clinical isolates of *Mycobacterium tuberculosis* with point mutations in the *rpoB* gene have

suggested that a 20-30% reduction in growth rate is approximately the maximum fitness cost

that drug-resistant mutants can incur before facing extinction in competitive drug-free

environments (Gagneux et al., 2006). To model drug-resistant strains, we therefore scaled k_F

and k_R such that the MIC of the resistant strain is 12 times that of its drug-susceptible

counterpart given a 25% fitness cost (c_R = 0.25) (**Figure 4A**).

409
410
Figure 4 – The propensity to select for resistant mutants depends on drug mechanism. (**A**)
We modeled wild-type strains using the parameters obtained from the calibration detailed in
Figure 2. (**B**) Relationship between MICs of resistant strains (expressed as multiples of
MIC_{WT}) and fitness cost of resistance. Horizontal dashed lines indicate the MICs of the wild-
type and resistant strains described in panel (A); the vertical dashed line indicates the fitness
cost at which all resistant strains have the same fold-increase in MIC relative to that of wild-

type (c_R = 0.25). (**C**) Pharmacodynamic curves for the wild-type and resistant strains described in panel (**A**). (**D**) Resistance selection windows for drug-resistant strains. The fitness advantage of resistant strains over wild-type strains is shown within the drug concentration range in which the resistant strain has a positive net growth rate that is larger than that of the wild-type. The fitness advantage is expressed as a proportion of the resistant strain's growth rate in the absence of drug ($G_{o,RES}$).

____

A nearly linear relationship exists between drug resistance and fitness cost for strains resistant to drugs with a linear effect on growth or death (**Figure 4B**, Bacteriostatic/Linear and Bactericidal/Linear). By contrast, drugs with stepwise effects on growth and killing (Bacteriostatic/Stepwise and Bactericidal/Stepwise) exhibit only modest reductions in MIC until they incur very high (>90%) fitness costs. We determined resistance selection windows for strains resistant to the five drugs in our set by simulating pharmacodynamic curves for wild-type and resistant strains (**Figure 4C**). To quantify the magnitudes of selection for resistant strains, we calculated the difference in net growth rates between wild-type and susceptible strains over the concentration range that defines the resistance selection window for each drug (**Figure 4D**). For linear-effect bacteriostatic drugs (Bacteriostatic/Linear), we found that the resistance selection window begins at drug concentrations as low as 200x below the MIC of the susceptible strain. Drugs with stepwise effects on growth or killing (Bacteriostatic/Stepwise and Bactericidal/Stepwise) have narrower resistance selection windows than their counterparts with more linear activity profiles.

Consistent with prior studies on the pharmacodynamic profiles of antimicrobial agents (Mohamed et al., 2014; Nielsen & Friberg, 2013; Yu et al., 2018), we find that the size of the resistance selection window is associated with the steepness of a drug's pharmacodynamic curve. Given a cellular effect (i.e. bacteriostatic or bactericidal), drugs with steeper pharmacodynamic curves tend to have narrower selection windows (**Supplementary Figure S7**). However, we also find that strains resistant to drugs with narrower resistance selection

windows have higher net growth rates within the resistance selection regime than do strains

resistant to drugs with wider resistance selection windows (**Figure 4D**). This finding has clear

clinical significance: drugs with steeper pharmacodynamic profiles feature relatively small

concentration ranges that select for resistance, but the negative consequences of dosing within

the resistance selection window are higher for these drugs.

The secondary mutant selection window is narrower for antibiotics with stepwise effects on growth

and death

The genotypic space for mutations that confer resistance to antibiotics by modifying the

binding kinetics of a drug to its target, such as those described in **Figure 4**, is typically highly

constrained (Levin et al., 2000; Palmer & Kishony, 2013). This is because a return to a drug-

susceptible state requires reversion of the specific genetic changes that conferred resistance in

a bacterial population, whereas secondary mutation accumulation can involve a wider range

of genetic changes throughout the cell's metabolic network. Therefore, the probability that a

bacterial population evolves secondary mutations that compensate for the fitness costs of a

resistance mutation is often higher than the probability that a bacterial population will revert

to susceptibility in drug-free environments (Isalan et al., 2008; Maisnier-Patin et al., 2002).

During treatment, resistant bacterial populations may also accumulate secondary mutations

that further raise MIC. In order to understand how drug mechanism influences such

secondary adaptation, we simulated the emergence of secondary mutants from drug-resistant

subpopulations of a bacterial population faced with antibiotic challenge (**Supplementary**

Figure S8; **Methods**, *Simulating the emergence of secondary mutations*).

The probability of secondary mutation emergence is substantially higher for drugs

with linear effects on cellular growth and death than it is for drugs with stepwise effects

(**Figure 5A**). This holds true for both bactericidal and bacteriostatic agents. Counterintuitively,

then, the suppression of secondary mutation emergence is not necessarily guaranteed by

rapid killing as suggested by earlier studies (Lipsitch & Levin, 1997). Likewise, rapid

attenuation of cell division does not halt the emergence of secondary mutations. We studied

the basis for this result by investigating the steady-state target occupancy distributions of cells

under antibiotic exposure. By accounting for the kinetics of drug-target binding, our

biochemical model shows that target occupancy among cells follows a distribution and is not

a single value even in otherwise clonal bacterial subpopulations (**Figure 5B**). This results in

heterogeneous replication rates within the drug-resistant subpopulation (**Supplementary**

Figure S9) that allow some bacteria to mutate. Classical population-dynamic models of

antibiotic action (Lipsitch & Levin, 1997; Regoes et al., 2004), which assume that a drug affects

the net growth rate of all cells equally, overlook this phenomenon.

481
482

Figure 5 – Emergence of secondary mutations among resistant subpopulations of infecting
bacteria. (**A**) Probability of a drug-resistant strain with secondary mutations emerging from
an infecting bacterial population before the infection is cleared (i.e. before the total bacterial

population decreases to less than 1). The initial population size for this simulation is 10^9 cells.
Inset shows probabilities of secondary mutation emergence before infection clearance when
the drug concentration used is 2x MIC_{RES}. (**B**) Frequency distributions of inactive drug-target
complexes for drug-resistant subpopulations undergoing steady-state exponential decline at
2x MIC_{RES}. (**C**) Probability of secondary mutant emergence from bacterial subpopulations
with *i* inactivated drug-target complexes, shown for ciprofloxacin dosed at 2x MIC_{RES}. (**D**)
Probability of secondary mutant emergence from bacterial subpopulations as a function of
drug dose, shown for ciprofloxacin dosed at 2x MIC_{RES}. Probabilities are shown as absolute
values (left panel) and as values normalized to the total probability of compensation for the
entire bacterial population over the course of treatment (right panel). (**E**) Resistance and
secondary mutant selection windows for different drug action mechanisms. The resistance
selection window (middle green) is defined as the drug concentration range over which a
drug-resistant strain has a growth advantage over wild-type. The secondary mutant selection
window (dark green) is defined as the drug concentration range over which the probability of
a resistant strain with secondary mutations emerging before infection clearance exceeds 10^{-4}
(see **Supplementary Figure S10** and **Methods**, *Simulating the emergence of secondary mutations*).
Dashed lines indicate the MICs of the wild-type and resistant strains. CIP: ciprofloxacin; S/S:
bacteriostatic/stepwise effect; S/L: bacteriostatic/linear effect; C/S: bactericidal/stepwise effect;
C/L: bactericidal/linear effect; MIC_{WT}: MIC of the wild-type strain; MIC_{RES}: MIC of the
resistant strain.
____

For ciprofloxacin doses only slightly above the MIC of the resistant strain ([Drug] = 2x

MIC_{RES}), we found that secondary mutations are most likely to emerge once the bacterial

population has reached a steady-state target occupancy distribution (**Figure 5C**). A

considerable probability of secondary mutation emergence nonetheless exists among

bacterial subpopulations with low numbers of inactivated drug-target complexes. These low-

occupancy subpopulations have faster growth rates and thus higher mutation rates. They are

also present in very large numbers during the initial stages of treatment, when drug molecules

are binding to their cellular targets and before the overall population begins to decline

(**Figure 1C**). We found that drug concentration influences the likelihood of a secondary

mutant arising from a steady-state or a low-occupancy subpopulation (**Figure 5D**). While the

overall probability of secondary mutation emergence decreases with higher drug dose (**Figure**

5D, left panel), the relative probability that a secondary mutation arises from a low-occupancy

population is greater for higher drug doses (**Figure 5D**, right panel). This implies that

secondary mutations are more likely to emerge very early during treatment when high drug

doses are used.

Prior studies have estimated that the probability of the existence of a fitness cost-free

bacterial pathogen prior to treatment ranges from 3×10^{-4} to 5×10^{-5} per infection (Colijn,

Cohen, Ganesh, & Murray, 2011). We sought to determine the range of drug concentrations

over which the likelihood of secondary mutation emergence during treatment is at least as

high as the likelihood for preexisting secondary resistance. We therefore determined the drug

concentration at which the probability of secondary mutation emergence before population

extinction equals 10^{-4} (that is, each treatment course has a 1 in 10,000 chance of giving rise to a

resistant strain bearing secondary mutations). We used this value as an upper boundary for

the "secondary mutant selection window," the range of drug concentrations over which the

probability of the emergence of a drug-resistant bacterial strain with secondary mutations is

substantial (**Supplementary Figure S10**). The secondary mutant selection window extends the

range of drug concentrations defined by the resistance selection window over which drug-

resistant strains may be selected (**Figure 5E**).

As with the resistance selection window, we found that the size of the secondary

mutant selection window varies dramatically depending on a drug's mechanism of action.

Drugs with linear effects on cellular growth and death have larger secondary mutant selection

windows than do drugs with stepwise effects on cellular growth and death. This is because for

drugs with stepwise effects, it is possible to shift the entire distribution of target occupancy to

a range where bacterial replication is virtually eliminated (or where bacterial death far

outweighs replication) across the entire population. With linear action, replication can still

occur even at high target occupancy, enabling the emergence of mutants. Drugs that fully

suppress cellular replication above MIC (i.e. Bacteriostatic/Stepwise) have small secondary

mutant selection windows, as the probability that additional mutations emerge over the

course of treatment is equal to the probability that a resistant strain with secondary mutations

emerges during the transient phase of drug-target binding immediately after treatment onset,

which lasts on the order of a few hours (**Figure 1C**).

Discussion

The increasing prevalence of first line- and multi-drug resistant bacteria (WHO, 2012, 2014)

signals the need for new antibiotics and robust drug dosing strategies that minimize the

emergence and spread of resistance (CDC, 2019). Despite this need, little is known about the

role that a drug's mechanism of action plays on the evolution of antibiotic resistance. We

studied the relationship between drug mechanism and drug resistance using a mathematical

model that connects bacterial population dynamics with molecular-scale descriptions of drug-

target binding kinetics (**Figure 1A**). Our biochemical model allows us to describe bacterial

replication and death as functions of drug-target occupancy, enables us to estimate molecular

kinetic parameters from population-scale data, and delivers performance on par with that of

classical pharmacodynamic models (**Figure 2B**).

We calibrate the model to an experimental dataset of ciprofloxacin time-kill curves

(**Figure 2A, Table 1**), and we show that drug-resistant strains can incur strikingly high fitness

costs associated with mutations that reduce drug-target binding kinetics (**Figure 3**). We find

that the relationship between drug-target inactivation and antibiotic effect (i.e. bacterial

killing, growth stalling, or both) exerts a strong influence on the strength of selection for

resistant strains during treatment, regardless of whether the drug is bactericidal or

bacteriostatic (**Figure 4D**). We also show that the molecular kinetics of drug-target binding

within cells results in heterogeneous replication rates among members of an otherwise

homogenous population (**Figure 5B**). This enables some drug-resistant strains to develop

secondary mutations that can further reduce drug susceptibility, increase resilience in drug-

free environments, and ultimately lead to treatment failure.

The clinical consequence of the frequently-observed trade-off between bacterial

fitness and drug resistance (Andersson & Hughes, 2010) is the existence of a resistance

selection window—a range of drug concentrations that selects for the propagation of drug-

resistant strains over their drug-susceptible counterparts (Karl Drlica & Zhao, 2007; Roberts et

al., 2008). It is important to note that numerous factors not captured by the resistance

selection window can contribute to resistance selection in clinical settings, most notably

ecological interactions between drug-susceptible strains, drug-resistant strains, and host

physiology (Day, Huijben, & Read, 2015). Our approach nonetheless enables us to isolate the

roles that a drug's mechanism of action play in driving the emergence of resistance.

We show that the resistance selection window is narrower for drugs that exert their

effects on growth or death in a stepwise (i.e. sudden) manner, resulting in a steeper

pharmacodynamic curve (**Figure 4C-4D, Supplementary Figure S7**). This result is consistent

with other studies on the pharmacodynamics of antimicrobial agents, which have found that

the size of the resistance selection window is associated with the steepness of the

pharmacodynamic curve (Mohamed et al., 2014; Nielsen & Friberg, 2013; Yu et al., 2018). The

characteristics of antimicrobial agents that enable steeper pharmacodynamic curves

nonetheless remain poorly described. Models that capture the effects of antibiotic drugs on

multiple scales, such as that described in this study and elsewhere (Clarelli et al., 2019; Wiesch

et al., 2015), could serve as helpful tools for studying the interplay between a drug's molecular

mechanism and its effect on bacterial population dynamics, enabling the design of new

antimicrobial agents with narrow resistance selection windows.

Mutations that alleviate the fitness costs associated with drug resistance and/or that

further raise a strain's MIC play a major role in driving the spread of antimicrobial resistance

across bacterial populations and clinical settings (Handel et al., 2006). Our study sheds

quantitative light on the mechanistic factors that govern the emergence of these secondary

mutations during treatment. We propose the use of the secondary mutant selection window

(**Supplementary Figure S10**) as a tool for illustrating the likelihood of further mutation

acquisition at nonzero drug concentrations. As with the size of the resistance selection

window, the size of the secondary mutant selection window varies greatly depending on the

mechanism of action of the antibiotic (**Figure 5E**). We stress that the secondary mutant

selection window does not necessarily indicate a region on the pharmacodynamic profile of a

drug over which the selection of a resistant strain with secondary mutations is favored. The

strength of selection depends on the physiological effect of the secondary mutation itself—

that is, whether the mutation accelerates growth rate, slows drug-target binding, or exerts a

multitude of other possible effects. Indeed, secondary mutations that act strictly by restoring

growth rates to wild-type levels lead only to modest (usually sublinear) increases in MIC

(**Figure 4B**), implying that strains with cost-free resistance phenotypes would still have MICs

well below the upper boundary for the secondary mutant selection windows shown in **Figure**

5E. Rather, the secondary mutant selection window defines the drug concentration range

within which the accumulation of secondary mutations is substantial and therefore clinically

significant.

Suppressing secondary mutation is crucial for reducing the survival of drug-resistant

mutants in antibiotic-free environments, where drug-resistant strains enter into direct

competition with other microbial organisms for limited resources (Andersson & Hughes, 2010;

Durão et al., 2018). We demonstrate that dosing drugs at or slightly above the MIC of a

resistant strain may not be sufficient for preventing the spread of resistance, and that—for

drugs with linear effects on bacterial growth and death as a function of drug-target

occupancy—there exist appreciable risks of selecting for secondary mutations even at doses

substantially above the MIC of the resistant strain. Reassessing the range of drug

concentrations that selects for resistant mutants as a composite of the resistance selection

window and the secondary mutant selection window (**Figure 5E, Supplementary Figure S10**)

could facilitate the design of drug dosing strategies that holistically mitigate the emergence

and spread of resistance.

Our study shows that both bactericidal and bacteriostatic drugs are capable of

exhibiting narrow resistance selection windows and low probabilities of secondary mutation

emergence in bacterial populations subjected to antibiotic treatment. This finding challenges

the long-accepted notion that bactericidal agents are superior to bacteriostatic agents in

suppressing the emergence of resistance during treatment (Stratton, 2003), and signals the

need to look beyond a drug's ability to kill or stall bacterial replication to assess the risks of

resistance emergence. The relationship between drug-target inactivation and overall

antibiotic effect has a much stronger influence on the strength of resistance selection than

does the drug's bacteriostatic or bactericidal activity (**Figure 4D**). The processes that may

dictate such a relationship for any given antibiotic nonetheless remain enigmatic. This

underscores the need for deeper experimental and theoretical research on the molecular

processes that govern the pharmacodynamics of antibiotic drugs.

The proper use of antibiotics in clinical and non-clinical settings constitutes a core

action for addressing the worldwide threat of antibiotic resistance (CDC, 2019). The

quantitative approach we present in this study may prove useful for identifying strategies that

manage the emergence of resistance to existing and future antimicrobial agents. We argue

that dosing regimens should account for a drug's resistance and secondary mutant selection

windows if they are to minimize the selection of resistance phenotypes during treatment. Our

findings suggest that even drugs with seemingly straightforward pharmacodynamic

classifications (i.e. bacteriostatic versus bactericidal action) can set bacterial populations on

complex and sometimes counterintuitive evolutionary trajectories with respect to resistance

selection. In the clinic, there exists little evidence that bactericidal antibiotics lead to more

favorable outcomes than do bacteriostatic antibiotics, especially for combatting

uncomplicated infections (Leekha, Terrell, & Edson, 2011; Pankey & Sabath, 2004). Yet it is

precisely in the treatment of uncomplicated, drug-susceptible infections that the greatest

gains are to be made in mitigating the emergence of resistance. Mechanistic models such as

that presented in this study can help to uncover clinically useful drug characteristics that

classical models may overlook. We envision a coupling of our quantitative approach with

high-throughput experimental platforms (Kulesa, Kehe, Hurtado, Tawde, & Blainey, 2018;

Schoepp et al., 2017) to aid in the development of new drugs with optimal pharmacodynamic

profiles and to accelerate the discovery of drug- and pathogen-specific dosing regimens that

minimize resistance emergence.

Methods

Bacterial time-kill curve experiment

We conducted time-kill curve experiments using *Escherichia coli* strain BW25113 (Coli Genetic

Stock Center #7636) (Datsenko & Wanner, 2000). We diluted overnight cultures of BW25113

1:1000 into pre-warmed lysogeny broth (LB) and grew cells to an optical density at 600nm

(OD_{600}) of 0.50. We then prepared a 1:3 dilution series of ciprofloxacin (highest concentration:

2.19 μg/ml) and added the antibiotics to bacterial cultures. We quantified bacterial population

sizes at regular (20-30 min) time intervals by plating a 1:10 dilution series of liquid culture onto

LB agar plates and counting colony forming units in technical triplicate. The bacterial kill

curve obtained at the highest ciprofloxacin concentration was used to determine the

maximum death rate (D_N) of bacterial cells, and a growth curve obtained using the same

protocol with the omission of ciprofloxacin was used to determine the maximum growth rate

(G_o) of cells in an antibiotic free environment (**Supplementary Figure S11**).

Model formulation and analysis

Our biochemical model constitutes a system of linear ordinary differential equations that

describe how successive numbers of inactivated drug-target complexes affect bacterial

replication and death. We consider a population of initial size B_o of phenotypically

homogenous and clonal bacteria exposed to a constant concentration C_o of drug. When no

drug is present, bacterial cells replicate at a rate G_o and die at a rate D_o. All cells have an

identical number N of proteins that drug molecules target for inactivation. We assume first-

order kinetics for drug-target binding: drug molecules bind to cellular protein targets within

cells, thereby inactivating the protein, at a rate k_F. Inactivated drug-protein targets dissociate

at a rate k_R. The first-order affinity of the drug to its protein target (K_D) is therefore the ratio of

the molecular dissociation rate to the molecular on-rate ($K_D = k_R/k_F$).

We stratify the entire bacterial population into $N+1$ subpopulations according to the

number i of inactivated drug-target complexes within each cell (i.e. the drug-target

occupancy), and we assume that growth and death rates of each bacterial cell depend on the

drug-target occupancy. That is, bacterial subpopulations with a larger drug-target occupancy

have slower growth rates and/or higher death rates than do bacterial subpopulations with a

smaller drug-target occupancy. Growth rate is therefore a monotonically decreasing discrete

function $G[i]$, and death rate is a monotonically increasing discrete function $D[i]$. We use

generalized logistic equations (**Supplementary Figure S1**) to describe overall growth and

death rates as a function of drug-target occupancy, allowing these functions to take the form

of a line, a sigmoidal curve, an exponential curve, or a step function. We assume that when a

drug inactivates all N protein targets in a cell, growth rate falls to zero (for bacteriostatic

drugs), death rate attains a maximal value D_N (for bactericidal drugs), or growth and death

rates are both affected (for drugs with mixed bactericidal and bacteriostatic action). In all of

these cases, the maximal rate of killing or growth attenuation can occur before all N target

proteins are inactivated if, for instance, $G[i]$ and/or $D[i]$ are step functions with inflection

points between 0 and N. During replication, a bacterial cell partitions its inactivated drug-

target complexes to two daughter cells according to a binomial distribution.

The change over time in the number of bacterial cells with exactly i inactivated drug-

target complexes (B_i) thus depends on the growth rate G_i, the death rate D_i, and the binding

kinetics of the drug to its protein target:

[**Equation 2**]

$$\frac{dB_i(t)}{dt} = (i+1)k_R B_{i+1} + \big(N-(i-1)\big)k_F C_0 B_{i-1} - ik_R B_i - (N-i)k_F C_0 B_i - D_i B_i - G_i B_i + \sum_{j=i}^{N} 2\frac{\binom{j}{i}}{2^j} G_j B_j$$

The first four terms on the right side of this equation describe changes in B_i due to drug-target

binding and unbinding. The fifth term describes bacterial death, and the sixth and seventh

terms describe bacterial growth. We can then define $B(t)$ as a vector whose elements comprise

the set of all bacterial subpopulations (B_0, B_1, ..., B_i, ..., B_{N-1}, B_N) at a given time. Because

Equation 2 is linear, we can describe the temporal dynamics of the entire bacterial population

with this vector, whose representation is a system of linear differential equations:

[**Equation 3**]

711
$$\frac{d\vec{B}(t)}{dt} = A\vec{B}$$

In the equation above, *A* denotes the matrix of coefficients describing the system of equations

for the vector *B*(*t*). The values for the coefficients in *A* depend on the concentration C_o of drug,

on the drug's binding kinetics, and on the growth and death rate functions *G*[*i*] and *D*[*i*].

Equation 3 represents an initial value problem. This system of linear differential

equations with a constant coefficient matrix has a unique solution given by

[**Equation 4**]

718
$$\vec{B}(t) = e^{At}\vec{B}_0$$

where the vector $\vec{B}_0$ denotes the initial composition of bacterial subpopulations at *t* = 0. The

solution can also be written as a linear superposition of a product of complex exponentials

(with arguments determined by eigenvalues) and polynomials (whose degree is determined by

the geometric multiplicity of these eigenvalues and whose coefficients are uniquely

determined by the initial conditions). In practice, *B*(*t*) describes a family of exponential growth

and decay curves that represent the replication and death of all *N*+1 bacterial subpopulations

over time (**Figure 1B**). We solve for *B*(*t*) numerically by calculating the matrix exponential of *A*

using a scaling and squaring algorithm implemented in MATLAB (MathWorks, Newton, MA)

(Al-Mohy & Higham, 2009).

Calculation of minimum inhibitory concentration

We define the MIC as the concentration C_o of an antibiotic such that any concentration of

drug at or above C_o is guaranteed to cause the eventual extinction of the bacterial population.

This occurs precisely when one eigenvalue of matrix *A* (from **Equation 3**) is zero and all other

eigenvalues have a negative real component. We thus express the MIC as

[**Equation 5**]

735
$$MIC = \inf\left\{ C_0 > 0 \mid \max\left(\mathcal{R}e(eig(A)) \right) = 0 \right\}.$$

With this formulation, finding the MIC amounts to finding the value of C_o such that the

greatest real component of the eigenvalues of A is zero. Deriving the expression for the MIC in

the simplest case of the model, when $N = 1$, serves to illustrate this approach. For the purposes

of this derivation, we consider a drug that elicits both a bactericidal and a bacteriostatic effect,

so $G[i = 1] = 0$ and $D[i = 1] = D_N$. However, the approach for finding the MIC is identical for any

mechanism of drug action. The matrix A describing all bacterial subpopulations ($B_{i=o}$ and $B_{i=1}$)

in this simple case is

[**Equation 6**]

744
$$A = \begin{bmatrix} G_0 - k_F C_0 & k_R \\ k_F C_0 & -(k_R + D_N) \end{bmatrix}.$$

We wish to find the concentration C_{MIC} of antibiotic that yields negative real components of

all but one eigenvalues λ of matrix A. For the 2-by-2 matrix given by **Equation 6**, the

characteristic polynomial is given by $\lambda^2 - \text{tr}(A)\lambda + \det(A)$, and the Routh-Hurwitz stability

criterion needed to satisfy the negative value constraints on λ is $\text{tr}(A) \leq 0$ and $\det(A) \geq 0$. For the

matrix described in **Equation 6**, these expressions correspond to

[**Equation 7**]

751
$$G_0 - k_F C_0 - k_R - D_N \leq 0$$

752   and

753   [**Equation 8**]

754
$$(G_0 - k_F C_0)(-k_R - D_N) - k_F k_R C_0 \geq 0.$$

755   Solving for the concentration C_o in both of these cases yields

756   [**Equation 9**]

757
$$C_0 \geq \frac{G_0 - k_R - D_N}{k_F}$$

758   in the case of **Equation 7** and

759 [**Equation 10**]

760
$$C_0 \geq \frac{(k_R + D_N)G_0}{k_F D_N}$$

in the case of **Equation 8**. We expect the value of k_R to be greater than that of G_o (that is, we

expect the rate of drug-target unbinding to be greater than the rate of bacterial replication).

We also expect the value of the death rate at saturating drug concentrations (D_N) to be

nonzero and positive. Therefore, **Equation 9** is guaranteed to be satisfied if **Equation 10** is also

satisfied. We thus find the expression for the MIC to be

[**Equation 11**]

767
$$C_{MIC} = \frac{(k_R + D_N)G_0}{k_F D_N}.$$

768 From this expression, we can infer the following proportionalities for the value of the MIC

769 relative to the values of other model parameters:

770 [**Equation 12**]

771
$$C_{MIC} \propto G_0$$

772
$$C_{MIC} \propto 1/k_F$$

773
$$C_{MIC} \propto k_R.$$

Polynomial expressions for the MIC, as shown in **Equation 11**, become exceedingly

complex beyond $N = 3$. However, we conjecture (although we have not been able to prove)

that the structure of the linear system shown in **Equation 3** guarantees the existence of the

MIC. For larger values of N, we leverage numerical schemes to calculate the eigenvalues of

matrix *A*. We use MATLAB's *eig()* function, which calculates eigenvalues using the QZ

algorithm (Moler & Stewart, 1973).

Model calibration via simulated annealing

Numerical values for the model parameters N, D_o, μ_R, and μ_C were obtained from the literature

(**Table 1**). The values for G_o and D_N were obtained by fitting experimental kill curves at drug

concentrations of zero and 2.19 μg/ml, respectively, to exponential functions (**Supplementary**

Figure S11). We leveraged an adaptive simulated annealing algorithm coupled with local

gradient descent to obtain the remaining parameters (k_F, k_R, α_G, α_D, γ_G, and γ_D). Detailed

descriptions of the adaptive simulated annealing algorithm are available elsewhere

(Henderson, Jacobson, & Johnson, 2003; Ingber, 1995); in brief, simulated annealing is a global

optimization algorithm capable of escaping local minima. It is therefore well suited to

applications involving the optimization of many parameters. Adaptive simulated annealing is

a variant on the classical simulated annealing algorithm that probes global parameter space

with greater efficiency by accounting for each parameter's magnitude when formulating a

new parameter set at every iteration of the algorithm. We used adaptive simulated annealing

to minimize the difference between experimental time-kill curves and model simulations of

bacterial populations challenged to the same antibiotic doses. The difference between

experimental observation and simulation is expressed through the objective function, whose

value ψ the adaptive simulated annealing algorithm seeks to minimize:

[**Equation 13**]

$$\psi = \sum_i \sum_j (W \, |E - B|)^2.$$

E denotes an m-by-n matrix of experimentally-measured population sizes at m drug

concentrations and n timepoints, B denotes simulated population sizes at the same drug

concentrations and timepoints, and W denotes an m-by-n weighting matrix (for our

application, simply a matrix of ones). B is a function of the parameters being optimized (that

is, $B = f(k_F, k_R, \alpha_G, \alpha_D, \gamma_G, \gamma_D)$).

Coupling the adaptive simulated annealing optimization with a local gradient descent

assures that our calibration procedure always converges on a local minimum. We used an

exponential cooling schedule for the simulated annealing algorithm, which assures that the

optimization runs ergodically (Ingber, 1995). That is, repeating the optimization many times

from random initial starting conditions in parallel yields roughly the same results as running

the optimization once for a very long time. This allowed us to parallelize the optimization

procedure by running the algorithm repeatedly across several cores of a computer and to

characterize the distributions of parameter values obtained from these calibrations

(**Supplementary Figure S3**). After performing 249 independent model calibrations, we

selected the parameter set with the lowest objective function value to use in subsequent

simulations. The parameter values for this set are shown in **Table 1**. Parameter sets for all

model optimizations performed are available in **Supporting Data File 3**.

Simulating the emergence of secondary mutations

We assumed that drug-resistant bacterial strains with secondary mutations that compensate

for fitness costs and/or that further increase MIC emerge from preexisting drug-resistant

subpopulations present in the initial population at the start of treatment (**Supplementary**

Figure S8). The size of the drug-resistant subpopulation at treatment onset ($B_{0,R}$) is given by

the mutation-selection balance, which expresses the size of the drug-resistant subpopulation

at which the rate of emergence of drug resistance alleles by spontaneous mutation equals the

rate of elimination of those alleles due to competitive fitness costs (Johnson, 1999):

[**Equation 14**]

827
$$B_{0,R} = \frac{B_0 \mu_R}{c_R}$$

Here, μ_R denotes the mutation rate for drug resistance emergence per unit time.

In order to quantify the probability of secondary mutation emergence from this drug-

resistant subpopulation, we adapted a formulation that Lipsitch and Levin developed to study

the evolution of drug-resistant bacterial strains during antibiotic treatment (Lipsitch & Levin,

1997). We assumed that secondary mutations emerge exclusively due to errors in DNA

replication during bacterial growth. The expected number of resistant cells with secondary

mutations that emerge from a bacterial population with i inactivated drug-target complexes

($E(M_{RC,i})$) is proportional to the total number of replications that the subpopulation undergoes

before extinction and the rate of secondary mutation emergence:

[**Equation 15**]

$$E\left(M_{RC,i}\right) = \mu_C \int_0^{t_{EXT,i}} G_{R,i}\, B_{R,i}(t)\, dt$$

In this equation, μ_C denotes the secondary mutation rate, $G_{R,i}$ represents the growth rate of a

resistant strain with exactly i inactivated drug-target complexes, $B_{R,i}(t)$ describes the

population dynamics of the ith drug-resistant bacterial subpopulation, and $t_{EXT,i}$ describes the

amount of time elapsed from treatment onset until the bacterial subpopulation is eliminated

($B_{i,R} = 1$ when $t = t_{EXT}$). The total number $E(M_{RC})$ of resistant mutants with secondary mutations

that we expect to observe over the course of treatment is thus the sum of **Equation 15** over all

values of i, and the probability P_{RC} that a compensated resistant mutant will emerge over the

course of treatment follows from the Poisson assumption that secondary mutations arise

stochastically and independently of other mutations:

[**Equation 16**]

$$P_{RC} = 1 - e^{-\left(\sum_{i=0}^{N} E(M_{RC,i})\right)}.$$

The summation term in **Equation 16** describes the total number of resistant strains with

secondary mutations expected to emerge before extinction. This equation thus quantifies the

Poisson probability that at least one resistant strain with a secondary mutation will emerge

over the course of treatment.

Code and data

We wrote all code in MATLAB. All of the code written for this study is available as a software

package in **Supplementary File 1**. Experimental data represented in **Figures 2A and 2B** and in

Supplementary Figure S4 are available within **Supporting Data Files 1, 2, and 4**, and the

parameter values for all iterations of model optimization are available in **Supporting Data**

File 3.

Acknowledgements
We extend sincere thanks to Sören Abel, Benjamin Akhuetie-Oni, and Laura Quinto for helpful feedback on the manuscript.

Author Contributions
Conceptualization: P.AzW.
Data curation: C.H.
Formal analysis: L.C., C.H., F.C., P.AzW.
Funding acquisition: P.AzW., T.C.
Investigation: C.H., F.C., A.P.
Methodology: L.C., P.AzW., C.H., A.P.
Project administration: P.AzW., T.C.
Resources: A.P., T.C., P.AzW.
Software: C.H.
Validation: C.H., L.C.
Visualization: C.H.
Writing – Original draft preparation: C.H.
Writing – Review & editing: C.H., P.AzW., F.C., L.C., T.C.

Competing interests
The authors declare no competing interests.

References

Ahn, S. H., Kim, D. H., Lee, A. R., Kim, B. K., Park, Y. K., Park, E.-S., . . . Kim, K.-H. (2015). Substitution at rt269 in Hepatitis B Virus Polymerase Is a Compensatory Mutation Associated with Multi-Drug Resistance. *PloS one, 10*(8). doi:10.1371/journal.pone.0136728

Al-Mohy, A. H., & Higham, N. J. (2009). A New Scaling and Squaring Algorithm for the Matrix Exponential. *SIAM Journal on Matrix Analysis and Applications, 31*(3), 970-989. doi:10.1137/09074721X

Andersson, D. I., & Hughes, D. (2010). Antibiotic resistance and its cost: is it possible to reverse resistance? *Nature Reviews Microbiology, 8*(4), 260-271. doi:10.1038/nrmicro2319

Andrews, J. M. (2001). Determination of minimum inhibitory concentrations. *The Journal of Antimicrobial Chemotherapy, 48 Suppl 1*, 5-16. doi:10.1093/jac/48.suppl_1.5

Billal, D. S., Feng, J., Leprohon, P., Légaré, D., & Ouellette, M. (2011). Whole genome analysis of linezolid resistance in Streptococcus pneumoniae reveals resistance and compensatory mutations. *BMC genomics, 12*, 512. doi:10.1186/1471-2164-12-512

Björkman, J., Nagaev, I., Berg, O. G., Hughes, D., & Andersson, D. I. (2000). Effects of Environment on Compensatory Mutations to Ameliorate Costs of Antibiotic Resistance. *Science, 287*(5457), 1479-1482. doi:10.1126/science.287.5457.1479

Blair, J. M. A., Webber, M. A., Baylay, A. J., Ogbolu, D. O., & Piddock, L. J. V. (2015). Molecular mechanisms of antibiotic resistance. *Nature Reviews Microbiology, 13*(1), 42-51. doi:10.1038/nrmicro3380

Brochet, M., Couvé, E., Zouine, M., Poyart, C., & Glaser, P. (2008). A Naturally Occurring Gene Amplification Leading to Sulfonamide and Trimethoprim Resistance in Streptococcus agalactiae. *Journal of Bacteriology, 190*(2), 672-680. doi:10.1128/JB.01357-07

CDC. (2019). *Antibiotic Resistance Threats in the United States, 2019*. Retrieved from Atlanta,
GA:
Clarelli, F., Palmer, A., Singh, B., Storflor, M., Lauksund, S., Cohen, T., . . . Wiesch, P. A. z.
(2019). Drug-target binding quantitatively predicts optimal antibiotic dose levels.
bioRxiv, 369975. doi:10.1101/369975
Colijn, C., Cohen, T., Ganesh, A., & Murray, M. (2011). Spontaneous Emergence of Multiple
Drug Resistance in Tuberculosis before and during Therapy. *PloS one, 6*(3), e18327.
doi:10.1371/journal.pone.0018327
Cui, J., Liu, Y., Wang, R., Tong, W., Drlica, K., & Zhao, X. (2006). The mutant selection
window in rabbits infected with Staphylococcus aureus. *The Journal of Infectious
Diseases, 194*(11), 1601-1608. doi:10.1086/508752
Datsenko, K. A., & Wanner, B. L. (2000). One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. *Proceedings of the National Academy of
Sciences of the United States of America, 97*(12), 6640-6645.
doi:10.1073/pnas.120163297
Day, T., Huijben, S., & Read, A. F. (2015). Is selection relevant in the evolutionary emergence
of drug resistance? *Trends in Microbiology, 23*(3), 126-133.
doi:10.1016/j.tim.2015.01.005
Dörr, T., Lewis, K., & Vulić, M. (2009). SOS Response Induces Persistence to Fluoroquinolones
in Escherichia coli. *PLOS Genetics, 5*(12), e1000760. doi:10.1371/journal.pgen.1000760
Drlica, K. (1999). Mechanism of fluoroquinolone action. *Current Opinion in Microbiology, 2*(5),
504-508.
Drlica, K., Malik, M., Kerns, R. J., & Zhao, X. (2008). Quinolone-mediated bacterial death.
Antimicrobial Agents and Chemotherapy, 52(2), 385-392. doi:10.1128/AAC.01617-06
Drlica, K., & Zhao, X. (2007). Mutant selection window hypothesis updated. *Clinical Infectious
Diseases: An Official Publication of the Infectious Diseases Society of America, 44*(5),
681-688. doi:10.1086/511642
Durão, P., Balbontín, R., & Gordo, I. (2018). Evolutionary Mechanisms Shaping the
Maintenance of Antibiotic Resistance. *Trends in Microbiology, 26*(8), 677-691.
doi:10.1016/j.tim.2018.01.005
Engelberg, H., & Artman, M. (1964). Studies on streptomycin-dependent bacteria: Effect of
streptomycin on protein synthesis by streptomycin-sensitive, streptomycin-resistant and
streptomycin-dependent, mutants of Escherichia coli. *Biochimica et Biophysica Acta
(BBA) - Specialized Section on Nucleic Acids and Related Subjects, 80*(2), 256-268.
doi:10.1016/0926-6550(64)90098-2
Everett, M. J., Jin, Y. F., Ricci, V., & Piddock, L. J. (1996). Contributions of individual
mechanisms to fluoroquinolone resistance in 36 Escherichia coli strains isolated from
humans and animals. *Antimicrobial Agents and Chemotherapy, 40*(10), 2380-2386.
Frenoy, A., & Bonhoeffer, S. (2018). Death and population dynamics affect mutation rate
estimates and evolvability under stress in bacteria. *PLOS Biology, 16*(5), e2005056.
doi:10.1371/journal.pbio.2005056
Gagneux, S., Long, C. D., Small, P. M., Van, T., Schoolnik, G. K., & Bohannan, B. J. M. (2006).
The Competitive Cost of Antibiotic Resistance in Mycobacterium tuberculosis. *Science,
312*(5782), 1944-1946. doi:10.1126/science.1124410
Gao, W., Chua, K., Davies, J. K., Newton, H. J., Seemann, T., Harrison, P. F., . . . Howden, B. P.
(2010). Two Novel Point Mutations in Clinical Staphylococcus aureus Reduce Linezolid

Susceptibility and Switch on the Stringent Response to Promote Persistent Infection.
PLOS Pathogens, 6(6), e1000944. doi:10.1371/journal.ppat.1000944

Gullberg, E., Cao, S., Berg, O. G., Ilbäck, C., Sandegren, L., Hughes, D., & Andersson, D. I.
(2011). Selection of Resistant Bacteria at Very Low Antibiotic Concentrations. *PLOS*
Pathogens, 7(7), e1002158. doi:10.1371/journal.ppat.1002158

Handel, A., Regoes, R. R., & Antia, R. (2006). The role of compensatory mutations in the
emergence of drug resistance. *PLoS computational biology, 2*(10), e137.
doi:10.1371/journal.pcbi.0020137

Harms, A., Maisonneuve, E., & Gerdes, K. (2016). Mechanisms of bacterial persistence during
stress and antibiotic exposure. *Science (New York, N.Y.), 354*(6318).
doi:10.1126/science.aaf4268

Henderson, D., Jacobson, S. H., & Johnson, A. W. (2003). The Theory and Practice of Simulated
Annealing. In F. Glover & G. A. Kochenberger (Eds.), *Handbook of Metaheuristics* (pp.
287-319). Boston, MA: Springer US.

Hughes, D. (2014). Selection and evolution of resistance to antimicrobial drugs. *IUBMB life,*
66(8), 521-529. doi:10.1002/iub.1278

Ingber, L. (1995). *Adaptive Simulated Annealing: Lessons Learned*. Retrieved from McLean,
VA:

Isalan, M., Lemerle, C., Michalodimitrakis, K., Horn, C., Beltrao, P., Raineri, E., . . . Serrano, L.
(2008). Evolvability and hierarchy in rewired bacterial gene networks. *Nature,*
452(7189), 840-845. doi:10.1038/nature06847

Johnson, T. (1999). The approach to mutation-selection balance in an infinite asexual population,
and the evolution of mutation rates. *Proceedings. Biological Sciences, 266*(1436), 2389-
2397. doi:10.1098/rspb.1999.0936

Jungkind, D. L., & American Society for Microbiology Eastern Pennsylvania, B. (1995).
Antimicrobial resistance: A crisis in healthcare: Plenum Press.

Kampranis, S. C., & Maxwell, A. (1998). The DNA gyrase-quinolone complex. ATP hydrolysis
and the mechanism of DNA cleavage. *The Journal of Biological Chemistry, 273*(35),
22615-22626. doi:10.1074/jbc.273.35.22615

Kulesa, A., Kehe, J., Hurtado, J. E., Tawde, P., & Blainey, P. C. (2018). Combinatorial drug
discovery in nanoliter droplets. *Proceedings of the National Academy of Sciences,*
115(26), 6685-6690. doi:10.1073/pnas.1802233115

Leekha, S., Terrell, C. L., & Edson, R. S. (2011). General Principles of Antimicrobial Therapy.
Mayo Clinic Proceedings, 86(2), 156-167. doi:10.4065/mcp.2010.0639

Levin, B. R., Perrot, V., & Walker, N. (2000). Compensatory mutations, antibiotic resistance and
the population genetics of adaptive evolution in bacteria. *Genetics, 154*(3), 985-997.

Lindgren, P. K., Marcusson, L. L., Sandvang, D., Frimodt-Møller, N., & Hughes, D. (2005).
Biological Cost of Single and Multiple Norfloxacin Resistance Mutations in Escherichia
coli Implicated in Urinary Tract Infections. *Antimicrobial Agents and Chemotherapy,*
49(6), 2343-2351. doi:10.1128/AAC.49.6.2343-2351.2005

Lipsitch, M., & Levin, B. R. (1997). The population dynamics of antimicrobial chemotherapy.
Antimicrobial Agents and Chemotherapy, 41(2), 363-373.

Loftie-Eaton, W., Bashford, K., Quinn, H., Dong, K., Millstein, J., Hunter, S., . . . Top, E. M.
(2017). Compensatory mutations improve general permissiveness to antibiotic resistance
plasmids. *Nature Ecology & Evolution, 1*(9), 1354. doi:10.1038/s41559-017-0243-2

Lovmar, M., Nilsson, K., Lukk, E., Vimberg, V., Tenson, T., & Ehrenberg, M. (2009). Erythromycin resistance by L4/L22 mutations and resistance masking by drug efflux pump deficiency. *The EMBO Journal, 28*(6), 736-744. doi:10.1038/emboj.2009.17

Maisnier-Patin, S., Berg, O. G., Liljas, L., & Andersson, D. I. (2002). Compensatory adaptation to the deleterious effect of antibiotic resistance in Salmonella typhimurium. *Molecular Microbiology, 46*(2), 355-366.

Majcherczyk, P. A., Barblan, J.-L., Moreillon, P., & Entenza, J. M. (2008). Development of glycopeptide-intermediate resistance by Staphylococcus aureus leads to attenuated infectivity in a rat model of endocarditis. *Microbial Pathogenesis, 45*(5-6), 408-414. doi:10.1016/j.micpath.2008.09.003

Martinez, J. L., & Baquero, F. (2000). Mutation frequencies and antibiotic resistance. *Antimicrobial Agents and Chemotherapy, 44*(7), 1771-1777. doi:10.1128/aac.44.7.1771-1777.2000

Melnyk, A. H., Wong, A., & Kassen, R. (2015). The fitness costs of antibiotic resistance mutations. *Evolutionary Applications, 8*(3), 273-283. doi:10.1111/eva.12196

Merker, M., Barbier, M., Cox, H., Rasigade, J.-P., Feuerriegel, S., Kohl, T. A., . . . Niemann, S. (2018). Compensatory evolution drives multidrug-resistant tuberculosis in Central Asia. *eLife, 7*, e38200. doi:10.7554/eLife.38200

Mohamed, A. F., Cars, O., & Friberg, L. E. (2014). A pharmacokinetic/pharmacodynamic model developed for the effect of colistin on Pseudomonas aeruginosa in vitro with evaluation of population pharmacokinetic variability on simulated bacterial killing. *Journal of Antimicrobial Chemotherapy, 69*(5), 1350-1361. doi:10.1093/jac/dkt520

Moler, C., & Stewart, G. (1973). An Algorithm for Generalized Matrix Eigenvalue Problems. *SIAM Journal on Numerical Analysis, 10*(2), 241-256. doi:10.1137/0710024

Morgan-Linnell, S. K., & Zechiedrich, L. (2007). Contributions of the combined effects of topoisomerase mutations toward fluoroquinolone resistance in Escherichia coli. *Antimicrobial Agents and Chemotherapy, 51*(11), 4205-4208. doi:10.1128/AAC.00647-07

Nemeth, J., Oesch, G., & Kuster, S. P. (2015). Bacteriostatic versus bactericidal antibiotics for patients with serious bacterial infections: systematic review and meta-analysis. *Journal of Antimicrobial Chemotherapy, 70*(2), 382-395. doi:10.1093/jac/dku379

Nielsen, E. I., & Friberg, L. E. (2013). Pharmacokinetic-Pharmacodynamic Modeling of Antibacterial Drugs. *Pharmacological Reviews, 65*(3), 1053-1090. doi:10.1124/pr.111.005769

Palmer, A. C., Chait, R., & Kishony, R. (2018). Nonoptimal Gene Expression Creates Latent Potential for Antibiotic Resistance. *Molecular Biology and Evolution, 35*(11), 2669-2684. doi:10.1093/molbev/msy163

Palmer, A. C., & Kishony, R. (2013). Understanding, predicting and manipulating the genotypic evolution of antibiotic resistance. *Nature Reviews. Genetics, 14*(4), 243-248. doi:10.1038/nrg3351

Palmer, A. C., & Kishony, R. (2014). Opposing effects of target overexpression reveal drug mechanisms. *Nature Communications, 5*, 4296. doi:10.1038/ncomms5296

Pan, X. S., Yague, G., & Fisher, L. M. (2001). Quinolone resistance mutations in Streptococcus pneumoniae GyrA and ParC proteins: mechanistic insights into quinolone action from enzymatic analysis, intracellular levels, and phenotypes of wild-type and mutant proteins.

Antimicrobial Agents and Chemotherapy, 45(11), 3140-3147. doi:10.1128/AAC.45.11.3140-3147.2001

Pankey, G. A., & Sabath, L. D. (2004). Clinical Relevance of Bacteriostatic versus Bactericidal Mechanisms of Action in the Treatment of Gram-Positive Bacterial Infections. *Clinical Infectious Diseases, 38*(6), 864-870. doi:10.1086/381972

Piddock, L. J. (1999). Mechanisms of fluoroquinolone resistance: an update 1994-1998. *Drugs, 58 Suppl 2*, 11-18. doi:10.2165/00003495-199958002-00003

Rao, G. G. (1998). Risk Factors for the Spread of Antibiotic-Resistant Bacteria. *Drugs, 55*(3), 323-330. doi:10.2165/00003495-199855030-00001

Redgrave, L. S., Sutton, S. B., Webber, M. A., & Piddock, L. J. V. (2014). Fluoroquinolone resistance: mechanisms, impact on bacteria, and role in evolutionary success. *Trends in Microbiology, 22*(8), 438-445. doi:10.1016/j.tim.2014.04.007

Regoes, R. R., Wiuff, C., Zappala, R. M., Garner, K. N., Baquero, F., & Levin, B. R. (2004). Pharmacodynamic functions: a multiparameter approach to the design of antibiotic treatment regimens. *Antimicrobial Agents and Chemotherapy, 48*(10), 3670-3676. doi:10.1128/AAC.48.10.3670-3676.2004

Roberts, J. A., Kruger, P., Paterson, D. L., & Lipman, J. (2008). Antibiotic resistance--what's dosing got to do with it? *Critical Care Medicine, 36*(8), 2433-2440. doi:10.1097/CCM.0b013e318180fe62

Schoepp, N. G., Schlappi, T. S., Curtis, M. S., Butkovich, S. S., Miller, S., Humphries, R. M., & Ismagilov, R. F. (2017). Rapid pathogen-specific phenotypic antibiotic susceptibility testing using digital LAMP quantification in clinical samples. *Science Translational Medicine, 9*(410). doi:10.1126/scitranslmed.aal3693

Schulz zur Wiesch, P., Engelstädter, J., & Bonhoeffer, S. (2010). Compensation of fitness costs and reversibility of antibiotic resistance mutations. *Antimicrobial Agents and Chemotherapy, 54*(5), 2085-2095. doi:10.1128/AAC.01460-09

Shen, L. L., & Pernet, A. G. (1985). Mechanism of inhibition of DNA gyrase by analogues of nalidixic acid: The target of the drugs is DNA. *PNAS, 82*, 307-311.

Silva, F., Lourenço, O., Queiroz, J. A., & Domingues, F. C. (2011). Bacteriostatic versus bactericidal activity of ciprofloxacin in Escherichia coli assessed by flow cytometry using a novel far-red dye. *The Journal of Antibiotics, 64*(4), 321-325. doi:10.1038/ja.2011.5

Silver, L. L. (2011). Challenges of Antibacterial Discovery. *Clinical Microbiology Reviews, 24*(1), 71-109. doi:10.1128/CMR.00030-10

Siporin, C., Heifetz, C. L., & Domagala, J. M. (1990). *The new generation of quinolones*: M. Dekker.

Stratton, C. W. (2003). Dead bugs don't mutate: susceptibility issues in the emergence of bacterial resistance. *Emerging Infectious Diseases, 9*(1), 10-16. doi:10.3201/eid0901.020172

Thorpe, K. E., Joski, P., & Johnston, K. J. (2018). Antibiotic-Resistant Infection Treatment Costs Have Doubled Since 2002, Now Exceeding $2 Billion Annually. *Health Affairs, 37*(4), 662-669. doi:10.1377/hlthaff.2017.1153

Vogwill, T., & MacLean, R. C. (2015). The genetic basis of the fitness costs of antimicrobial resistance: a meta-analysis approach. *Evolutionary Applications, 8*(3), 284-295. doi:10.1111/eva.12202

Wang, P., Robert, L., Pelletier, J., Dang, W. L., Taddei, F., Wright, A., & Jun, S. (2010). Robust
growth of Escherichia coli. *Current biology: CB, 20*(12), 1099-1103.
doi:10.1016/j.cub.2010.04.045
WHO. (2012). *The evolving threat of antimicrobial resistance: options for action* (978-92-4-
450318-8). Retrieved from https://apps.who.int/iris/handle/10665/44812
WHO. (2014). *Antimicrobial resistance: Global report on surveillance*. Retrieved from Geneva:
Wiesch, P. A. z., Abel, S., Gkotzis, S., Ocampo, P., Engelstädter, J., Hinkley, T., . . . Cohen, T.
(2015). Classic reaction kinetics can explain complex patterns of antibiotic action.
Science Translational Medicine, 7(287), 287ra273-287ra273.
doi:10.1126/scitranslmed.aaa8760
Wiśniewski, J. R., & Rakus, D. (2014). Quantitative analysis of the Escherichia coli proteome.
Data in Brief, 1, 7-11. doi:10.1016/j.dib.2014.08.004
Yu, G., Baeder, D. Y., Regoes, R. R., & Rolff, J. (2018). Predicting drug resistance evolution:
insights from antimicrobial peptides and antibiotics. *Proceedings. Biological Sciences,*
285(1874). doi:10.1098/rspb.2017.2687
Zhang, Q., Sahin, O., McDermott, P. F., & Payot, S. (2006). Fitness of antimicrobial-resistant
Campylobacter and Salmonella. *Microbes and Infection, 8*(7), 1972-1978.
doi:10.1016/j.micinf.2005.12.031
1108
1109

Supplementary Figure S1 – Bacterial growth and death rates as a function of drug-target occupancy. We define the functions G[i] and D[i] as constrained logistic curves such that G[i = 0] = G_o, G[$i = N$] = 0, D[i = 0] = D_o, and D[$i = N$] = D_N. The parameters α_G and α_D define the steepness of the logistic curves for the growth and death rate function, respectively. α_G and α_D are unitless and range from 1 to 500; 1 yields a quasi-linear function, while 500 yields a quasi-step function. The parameters γ_G and γ_D define the inflection point of the logistic curves for the growth and death rate function, respectively. γ_G ranges from $-N$ to N and γ_D ranges from 0 to $2N$; the curve is quasi-sigmoidal if γ_G and γ_D are in between 0 and N and is quasi-exponential if γ_G and γ_D are outside of these bounds.

1123
1124
Supplementary Figure S2 – Simulated time-kill curves of *Escherichia coli* **exposed to a range**
of drug concentrations. We used the parameter set outlined in **Table 1** to model the growth
and death of bacterial populations subjected to drug concentrations up to 16x minimum
inhibitory concentration (MIC). Drug concentrations are expressed as factors of the MIC. The
net growth rate of the entire bacterial population over the time course of the simulation
decreases with increasing drug concentration.
1131
1132

1133
1134
Supplementary Figure S3 – Results from 249 independent model calibrations to
experimental data. We used adaptive simulated annealing coupled with gradient descent (see
Methods, *Model calibration via simulated annealing*) to fit the model to experimental kill curve
data of *E. coli* exposed to ciprofloxacin (**Supporting Data File 1**). Shown in this figure are the
results for 249 independent model fits, each beginning with randomly-chosen values for the
parameters describing drug-target binding rate k_F, drug-target unbinding rate k_R, steepness of
the growth rate function α_G, steepness of the death rate function α_D, inflection point of the
growth rate function γ_G, and inflection point of the death rate function γ_D. (**A**) Frequency
distribution of objective function values obtained from independent model calibrations. The
objective function value describes the goodness of the fit between experimental data and
simulation; smaller values indicate higher goodness of fit. (**B-H**) Optimization plots showing
randomly chosen initial parameter values (x-axis) and calibrated parameter values (y-axis) for
all independent model calibrations. The optimized parameters are k_F (**B**), k_R (**C**), K_D (the ratio
of k_R to k_F) (**D**), α_G (**E**), α_D (**F**), γ_G (**G**), and γ_D (**H**). The final objective function value of each
model fit is colored according to the color bar above panel (**A**).
1150
1151

1152
1153
Supplementary Figure S4 – Pharmacodynamic curves generated from experimental data
and from the calibrated model. The experimental pharmacodynamic curve was generated
by calculating the net growth rates of *E. coli* exposed to a set of ciprofloxacin drug
concentrations (**Supporting Data File 1**). The time-kill curves of this same experimental
dataset are shown in **Figure 2A**; see **Supporting Data File 4** for experimental data on net
growth rate as a function of drug concentration. The model-calibrated pharmacodynamic
curve was generated by simulating bacterial time-kill curves over the same range of drug
concentrations used in the experiment and calculating associated net growth rates.
1162
1163

Supplementary Figure S5 – Simulated population curves for ciprofloxacin and for four extreme modes of antibiotic drug mechanism. We simulated a bacterial population of 10^8 cells exposed to antibiotic drug at 4x MIC. The ciprofloxacin curve corresponds to the drug mechanism obtained from the model calibration to experimental data and detailed in **Figure 2C**, and the remaining curves correspond to the extreme schemes of drug mechanism shown in **Figure 2D**.

Supplementary Figure S6 – MIC as a function of drug-target binding and unbinding kinetics. The MIC of a mutant (normalized to the MIC of the wild-type) is plotted against the fold-change in (**A**) drug-target binding (k_F) or (**B**) drug-target complex disassociation (k_R). For this simulation, mutants have no fitness costs associated with changes in k_F and k_R ($c_R = 0$). For drug-target binding (k_F), fold increase in MIC is directly proportional to fold decrease in k_F for all drug mechanisms. In both panels, the dashed line indicates the line of direct proportionality. MIC_{WT}: MIC of the wild-type strain; MIC_{RES}: MIC of the resistant strain.

1185
1186
Supplementary Figure S7 – Drugs with steeper pharmacodynamic curves tend to have
narrower resistance selection windows. To quantify the steepness of pharmacodynamic
curves, we fit the curves for drug-resistant strains shown in **Figure 4C** to the
pharmacodynamic function formulated by Regoes et al. (Regoes et al., 2004). The equation
describes the net growth rate G_{net} of a bacterial population as a function of drug concentration
C_o and other parameters (MIC, G_o, D_N) derived from the model:

1193
$$G_{net} = G_0 - \frac{(G_0 - D_N)(C_0/\text{MIC})^\kappa}{(C_0/\text{MIC})^\kappa - (D_N/G_0)}$$

In this equation, κ describes the Hill coefficient, which serves as a measure of the steepness of
the pharmacodynamic curve. Larger values of κ indicate steeper curves. For each of the drug
mechanisms described in this study (**Supplementary Table 1**), we generated
pharmacodynamic curves for drug-resistant mutants (**Figure 4C**, solid lines), determined the
value of κ that best fits the curve, and plotted κ against the range of drug concentrations that
represents the resistance selection window (**Figure 4D**). MIC_WT: MIC of the wild-type strain;
MIC_RES: MIC of the resistant strain.
1201

1202
1203
Supplementary Figure S8 – Emergence of secondary mutations within subpopulations of
drug-resistant bacteria during antibiotic treatment. When simulating the emergence of
secondary mutations, we assume that a drug-resistant subpopulation (middle green) of
bacteria is present at the start of treatment; the size of this subpopulation is given by the
mutation selection balance of the allele that confers the drug-resistance phenotype (Johnson,
1999). We calculate the probability that a drug-resistant strain with secondary mutations (dark
green) emerges from this subpopulation before the elimination of the drug-resistant strain (at
time t_{EXT}).
1212
1213

1214
1215
1216 **Supplementary Figure S9 – Distributions of growth and death rates for drug-resistant**
1217 **bacterial subpopulations undergoing steady-state exponential decline at 2x MIC_{RES}.** Boxes
1218 denote the central 50% of the growth and death rate distributions, and whiskers denote the
1219 central 95% of the growth and death rate distributions.
1220
1221

1222
1223
Supplementary Figure S10 – The secondary mutant selection window. The secondary
mutant selection window comprises the drug concentration range over which the net growth
of the drug-resistant strain is negative but the probability of secondary resistance emergence
before the end of treatment exceeds a defined threshold (in our simulations, 10^{-4}, or a 1 in
10,000 chance). Four regimes of selection exist: the null selection window in which the wild-
type strain dominates, the resistance selection window, the secondary mutant selection
window, and the complete killing window. We simplify these four regimes by disregarding
the relative strengths of selection for each strain in each regime and we instead illustrate the
boundaries of each region along a drug concentration axis (top bar); these simplified selection
regimes are shown for all five drug mechanisms studied in **Figure 5E.**
1234
1235

1236
1237
Supplementary Figure S11 – Obtaining G_o and D_N from experimental data. (**A**) To obtain the
value of G_o (growth rate in the absence of antibiotic) used in simulations, we fit an exponential
growth curve to experimental data for *E. coli* cells grown in the absence of antibiotic. (**B**) To
determine the value of D_N (maximum death rate in saturating conditions of antibiotic), we fit
an exponential decay curve to experimental data for *E. coli* cells exposed to 2.19 µg/ml of
ciprofloxacin (~200 x MIC). The population size deviates from exponential decay at later
timepoints (dashed and shaded) likely because of the emergence of persistent subpopulations
of bacteria (Dörr et al., 2009). The R^2 values shown are the linear correlation coefficients for
the model fit, and are not the correlation coefficients for the log-transform of the data.
1247
1248

1249

Drug mechanisms					
Activity	Ciprofloxacin	Bacteriostatic	Bacteriostatic	Bactericidal	Bactericidal
Steepness		Stepwise	Linear	Stepwise	Linear
Abbreviation	**CIP**	**S/S**	**S/L**	**C/S**	**C/L**
α_G	16.8	500	1	1	1
α_D	7.28	1	1	500	1
γ_G	24.8	35	35	35	35
γ_D	364	150	150	150	150
Bacteriostatic Potency	1	1	1	0	0
Bactericidal potency	1	0	0	1	1

1250

Supplementary Table S1 – Parameters for a set of five drugs with different mechanisms of
action. The parameters α_G and α_D describe the steepness of the growth and death rate
functions, respectively, around the inflection point. The parameters γ_G and γ_D describe the
inflection points of the growth and death rate functions (see **Supplementary Figure S1**).
Bacteriostatic potency refers to the magnitude of growth rate decline at saturating
concentrations of drug; a value of 1 indicates that that growth rate declines to zero in
saturating concentrations of drug ($G[i = N] = 0$), and a value of 0 indicates that growth rate is
unaffected by drug concentration ($G[i] = G_o$ for all i). Bactericidal potency refers to the
magnitude of death rate increase at saturating conditions of drug; a value of 1 indicates that
death rate increases to maximum in saturating concentrations of drug ($D[i = N] = D_N > D_o$), and
a value of 0 indicates that death rate is unaffected by drug concentration ($D[i] = D_o$ for all i).
All other parameters (including drug-target binding rate k_F, drug-target unbinding rate k_R, and
target number N) are identical for all drugs in the set.
1264
1265

1266 **Supplementary File 1 – MATLAB code package containing the code written for this study.**
1267 This file contains scripts that we used to implement our model, to analyze data, and to
1268 generate the numeric values for all main text and supplementary figures.
1269
1270 **Supporting Data File 1 – Experimental data for the ciprofloxacin time-kill curve**
1271 **experiment represented in Figure 2A and Supplementary Figure S11.**
1272
1273 **Supporting Data File 2 – Experimentally-measured minimum inhibitory concentrations**
1274 **(MICs) for ciprofloxacin against *Escherichia coli* represented in Figure 2B.** We collated this
1275 list of experimentally-measured MICs from the literature; study sources are given in the file.
1276
1277 **Supporting Data File 3 – Model calibrations obtained via simulated annealing.** Starting and
1278 ending values for all model parameters are given for each iteration of the model fitting
1279 procedure described in **Methods,** *Model calibration via simulated annealing*.
1280
1281 **Supporting Data File 4 – Experimental pharmacodynamic curve data represented in**
1282 **Supplementary Figure S4.** We generated these data by calculating the net growth rates of
1283 bacterial populations at each drug concentration listed in **Supporting Data File 1**.
1284