A microRNA Signature of Metastatic Colorectal Cancer

Running title: miRNA in metastatic colorectal cancer

Eirik Høye[1,2,*], Bastian Fromm[1,3,*], Paul Heinrich Michael Böttger[4], Diana Domanska[5], Annette Torgunrud[1], Christin Lund-Andersen[1,2], Torveig Weum Abrahamsen[1], Åsmund Avdem Fretland[2,6,7], Vegar Johansen Dagenborg[1,2], Susanne Lorenz[8], Bjørn Edwin[6,7,2], Eivind Hovig[1,9], Kjersti Flatmark [10,1,2**]

[1]   Department of Tumor Biology, Institute for Cancer Research, The Norwegian Radium Hospital, Oslo University Hospital, Oslo, Norway

[2]   Institute of Clinical Medicine, Medical Faculty, University of Oslo, Oslo, Norway

[3]   Science for Life Laboratory, Department of Molecular Biosciences, The Wenner-Gren Institute, Stockholm University, Stockholm, Sweden

[4]   Independent researcher, Linz, Austria

[5]   Department of Pathology, University of Oslo, Oslo, Norway

[6]   The Intervention Centre, Rikshospitalet, Oslo University Hospital, Oslo, Norway

[7]   Department of Hepato-Pancreato-Biliary Surgery, Rikshospitalet, Oslo University Hospital, Oslo, Norway

[8]   Department of Core Facilities, Institute for Cancer Research, The Norwegian Radium Hospital, Oslo University Hospital, Oslo, Norway

[9]   Center for Bioinformatics, Department of Informatics, University of Oslo, Oslo, Norway

[10] Department of Gastroenterological Surgery, The Norwegian Radium Hospital, Oslo University Hospital, Nydalen, Oslo, Norway

[*] These authors contributed equally to this work

[**]Corresponding author, email: kjersti.flatmark@rr-research.no

Abstract

Although microRNAs (miRNA) are involved in all hallmarks of cancer, miRNA dysregulation in metastasis remains poorly understood and contradictory results have been published. The aim of this work was to identify miRNAs associated with metastatic progression of colorectal cancer (CRC). Novel and previously published next generation sequencing (NGS) datasets generated from 268 samples with primary (pCRC) and metastatic CRC (mCRC; liver, lung and peritoneal metastases) and tumor adjacent tissues were analyzed. Differential expression analysis was performed using a meticulous bioinformatics pipeline, including only bona fide miRNAs, utilizing miRNA-tailored quality control and processing, and applying a physiologically meaningful cut-off value (100 reads per million). The results were adjusted for host tissue background expression and samples from the different metastatic sites were independently analyzed. A metastatic signature containing five miRNAs up-regulated at multiple metastatic sites was identified (Mir-210_3p, Mir-191_5p, Mir-8-P1b_3p *(mir-141-3p)*, Mir-1307_5p, and Mir-155_5p) along with a number of miRNAs that were differentially expressed at individual metastatic sites. Several of these have previously been implicated in metastasis through involvement in epithelial-to-mesenchymal transition and hypoxia, while other identified miRNAs represent novel findings. The identified differentially expressed miRNAs confirm known associations and contribute novel insights into miRNA involvement in the metastatic process. The use of open science practices facilitates reproducibility, and new datasets may easily be added to the publicly available pipeline to continuously improve the knowledge in the field. The identified set of miRNAs provides a reliable starting-point for further research into the role of miRNAs in metastatic progression.

Introduction

Colorectal cancer (CRC) is a heterogeneous disease and a leading cause of cancer-related deaths worldwide[1], and metastatic progression to the liver, lungs and peritoneal surface remains the primary cause of CRC-related mortality. Metastasis is a complex process where cancer cells undergo adaptation to enable survival and establishment of tumors in organs with very different microenvironments[2,3]. The genomic and transcriptomic changes in metastatic CRC (mCRC) remain incompletely understood, particularly in the context of organ-specific metastasis[4–6].

MicroRNAs (miRNAs) are evolutionary ancient post-transcriptional gene regulators that are involved in numerous biological processes and are molecular players in human disease, including cancer[7,8]. MiRNAs can be extracted from tissues and body fluids, and because of their remarkable chemical stability and the availability of sensitive detection methods, miRNAs have been suggested as cancer biomarkers[9–11]. However, to date, no miRNAs have been clinically implemented as biomarkers of CRC[5,12–18]. In mCRC, in particular, there is little consensus regarding which miRNAs are up- and down-regulated, limiting our understanding of their role in metastatic progression[5].

The lack of consensus likely reflects some well-known caveats related to analysis of miRNAs in human disease. Non-miRNA sequences have been incorrectly annotated as miRNA genes[19], and a bioinformatics workflow specifically tailored for analysis of miRNAs in bulk tissue samples has not been available. Furthermore, differential expression analysis has been performed without ensuring the presence of physiologically relevant tissue expression levels[20]. Also, accounting for differences in cellular composition is important, since many miRNAs are exclusively expressed in particular cell types or at specific developmental time points[21–24], likely confounding analysis of bulk tissue samples. Finally, to elucidate the role of miRNAs in mCRC, failure to consider differences related to metastatic location, and differences in normal background expression, may have contributed to inconsistent results[12,25].

To overcome these challenges, the publicly available, manually curated miRNA gene database MirGeneDB (mirgenedb.org), was used as miRNA reference[26], and a novel bioinformatics pipeline was developed. The bioinformatics work-flow included use of the miRTrace software as a universal quality control pipeline specifically for miRNA next-generation sequencing (NGS) data[27] with subsequent processing using miRge3.0[28]. A strict cut-off of 100 reads per million (RPM) was applied as the minimum expression level for physiological

relevance[12]. Taking cell-type specific miRNA expression into account, metastatic samples from different sites

were independently analyzed, which allowed correction for different background expression levels at the

individual sites. Existing publicly available miRNA-sequencing (miRNA-seq) patient derived datasets,

combined with novel miRNA-seq datasets from pCRC and mCRC with normal adjacent tissues, were analyzed,

after quality control totaling 268 datasets. Using this unbiased analytical approach, a novel mCRC miRNA

signature was identified, which partially overlaps with previous reports, but which also includes several miRNAs

previously not identified in this context.

Results

NGS data collection and processing

New NGS datasets were successfully generated from 85 samples: pCRC (n=3), tumor adjacent colorectum (nCR; n=3), liver metastases (mLi; n=19), tumor adjacent liver (nLi; n=9), lung metastases (mLu; n=25), tumor adjacent lung (nLu; n=7), and peritoneal metastases (PM; n=20). From five studies, previously published datasets containing NGS analyses of miRNA in pCRC and mCRC were also included[12,14,15,29] while data from four published studies were not accessible despite repeated requests[30–33]. In total, 350 NGS datasets were subjected to quality assessment using the miRTrace pipeline. The majority of the new NGS datasets and datasets from three previously published studies[12,15,34] fulfilled the quality control criteria. Two studies were excluded, one because of low quality reads in the majority of samples[29], the other because of contamination of reads from other organisms[14] (Supplementary file 1). After quality control, a total of 268 NGS datasets remained for further analysis, including pCRC (n=120), nCR (n=25), mLi (n=35), nLi (n=20), mLu (n=28), nLu (n=10) and PM (n=30). These datasets were then successfully processed and mapped to MirGeneDB[26] using miRge3.0[28].

Global miRNA expression

When analyzing global miRNA expression using the dimensionality reducing algorithm uniform manifold approximation and projection for dimension reduction (UMAP), the datasets clustered according to the tissue of origin (Fig. 1). The normal tissues (nLi, nLu and nCR) formed distinct clusters reflecting the unique transcriptional profiles of these organs. nLi and nCR clustered separately from the corresponding tumor tissues, while nLu clustered close to the mLu tissue datasets. In general, the pCRC and mCRC dataset clusters were less homogeneous than the normal tissue counterparts, but with exception of the PM datasets, which exhibited considerable overlap with the other malignant datasets, the pCRC, mLi, and mLu datasets all formed distinct clusters.

Differential miRNA expression between pCRC and nCR

Thirty-two miRNAs were up-regulated and 35 miRNAs were down-regulated when comparing pCRC to nCR. Among the differentially expressed miRNAs were many well-known oncomiRs, including Mir-21_5p, multiple MIR-17 family members, Mir-31, Mir-221 (up-regulated) and Mir-8-P1b_3p (miR-141) (down-regulated). Several cell-type specific miRNAs were also detected at different levels in the two tissues, including higher

104   levels of Mir-17-P1a/P1b_5p (mir-17; CD14+ monocytes) and Mir-223_3p (dendritic cells), and lower levels of

105   Mir-486_5p and Mir-451_5P (red blood cells), the Mir-143_5p and Mir-145_5p (mesenchymal cells), Mir-

106   150_5p (lymphocytes), Mir-375_3p and Mir-192-P1_5p (epithelial cells), and Mir-342_3p (dendritic cells,

107   lymphocytes and macrophages)[22]. For a complete overview of differentially expressed miRNAs between pCRC

108   and nCR, see Supplementary file 3.

109   Differential expression analysis identifies miRNA signature of mCRC

After performing site specific differential expression analysis and correcting for background expression, a total

of 26 miRNAs were identified as differentially expressed in one or more of the metastatic tissues compared to

pCRC (Fig. 2 and Table 1). Two miRNAs, Mir-210_3p and Mir-191_5p, were up-regulated at all three

metastatic sites. In addition, three miRNAs were up-regulated at two of the three sites; Mir-8-P1b_3p in mLi and

mLu, Mir-1307_5p in mLi and PM, and Mir-155 in mLu and PM (Fig. 3). All these miRNAs were expressed

well above the threshold for biological significance, with Mir-191_5p and Mir-8-P1b_3p being expressed at

particularly high levels (>1000 RPM).

In addition, several miRNAs were up- or down-regulated at individual metastatic sites. In mLi, four miRNAs

were identified as up-regulated, Mir-10-P1a_5p, Mir-592_5p, Mir-1247_5p and Mir-425_5p, while Mir-486_5p

was down-regulated. Of these, Mir-10-P1a_5p was expressed at exceptionally high levels, with 97 123 RPM in

pCRC and 164 925 RPM in mLi. In mLu Mir-19-P1_3p, Mir-19-P2a/P2b_3p, Mir-374-P1_5p and Mir-142_5p

were up-regulated while Mir-423_5p, Let-7-P1b_5p, Mir-197_3p, Mir-92-P1c_3p, Mir-362-P2/P4_3p and Mir-

221_3p were down-regulated. In PM Mir-506-P3_3p, Mir-506-P4a1/P4a2/P4b_3p, Mir-154-P9_3p and Mir-

127_3p, Mir-154-P36_3p were up-regulated and Mir-223_3p were down-regulated.

124   qPCR validation

125   qPCR analysis of randomly selected mLi (n=11) and pCRC (n=11) samples not included in the previous NGS

126   analysis replicated the findings from the NGS data, validating up-regulation of Mir-210_3p in mLi compared to

127   pCRC. Welch two-sided t-test on dCq values, t-statistic = -2.25, degrees of freedom = 19.95, p-value = 0.036.

128   Analysis of cell-type specific miRNAs

Of the 45 previously validated cell-type specific miRNAs, only 25 had a mean expression greater than 100 RPM

in at least one of the tissues and formed the basis for the following analysis. The relative expression levels of

cell-type specific miRNAs showed clear differences between the tissues, particularly prominent for the tumor

adjacent tissues (Fig. 4A). For instance, hepatocyte specific Mir-122_5p was detected at high and moderate

levels in nLi and mLi, respectively, and at extremely low levels in the other tissues. Also, Mir-143_3p and Mir-

145_5p was detected at higher levels in nCR and nLu relative to the other datasets (p=4.38E-02 and p=3.42E-04,

respectively). The epithelial cell specific Mir-8-P2a_3p and Mir-8-P2b_3p, were detected at higher levels in the

intestinal epithelial-derived tissues compared to nLi and nLu (p=2.20E-16 and p=2.24E-16, respectively). In line

with these findings, principal component analysis (PCA) (Fig. 4B) showed distinct clusters for nLi and nLu,

whereas nCR clustered closer to pCRC and the metastatic tissues. The correlation circle (Fig. 4C) shows the

loading of the PCA, indicating the direction and relative contribution of the 15 cell-type specific miRNAs that

contributed most to the clustering of each tissue.

Gene set enrichment (GSE) analysis

RBiomirGS analysis revealed multiple GO terms and KEGG pathways that were predicted to be either more or

less repressed because of differential expression of miRNAs between pCRC and mCRC. Among terms predicted

to be less repressed, several were common to mLi and mLu. These included the five Molecular Function GO

terms: Transcription Coregulator Activity, Transcription Coactivator Activity, Transcription Regulator Activity,

mRNA Binding and Poly Purine Tract Binding. Also estimated to be less repressed were the eight Cellular

Component GO terms, including Transcription Repressor Complex, Caveola, Transferase Complex, Golgi

Aparatus, Glutamatergic Synapse, Catalytic Complex, Nuclear Envelope and Transcription Regulator Complex.

Furthermore, 92 Biological Process GO terms, many of which are related to transcriptional activity, were also

estimated to have reduced miRNA repression in both mLi and mLu. Among the KEGG pathways, only mLu had

pathways predicted to be less repressed, including Long Term Potentiation, Axon Guidance, Long Term

Depression, Ubiquitin Mediated Proteolysis, Neurotrophin Signaling Pathway, TGF Beta Signaling Pathway,

Adherens Junction and Phosphatidylinositol Signaling System. One KEGG pathway, Systemic Lupus

Erythematosus, was estimated to be more repressed in both mLi and mLu, while the Parkinsons Disease KEGG

pathway was repressed only in mLu. Notably, no differentially repressed GO terms or KEGG pathways were

shared between the PM datasets and the other two sites. There were also many GO terms and KEGG pathways

that were differentially repressed specifically to individual sites. All results for the GSE are shown in

Supplementary file 6.

Discussion

The initial differential expression analysis comparing pCRC and nCR revealed 68 differentially expressed miRNAs, reflecting major and genome-wide changes of malignant transformation. Among the up-regulated miRNAs, a subset representing previously reported "oncoMirs" was identified, including Mir-21, and members of the three clusters Mir-17-92, Mir-31, and Mir-221. These miRNAs have validated mRNA targets in CRC, such as phosphatase and tensin homolog (PTEN), transforming growth factor beta receptor (TGFBR), and Smad[16,35,36]. A number of other miRNAs previously connected to CRC were also up-regulated, such as MIR-15 family member Mir-29-P2a/P2b (*miR-29b*), two MIR-96 family members, Mir-135-P3, Mir-130-P2a (*miR-301a*)[37,38], Mir-181-P1c[39], and Mir-224[16,35]. One up-regulated miRNA, Mir-95-P2 (*miR-421*), has not been reported for CRC previously, but its deregulation has been demonstrated in other cancers[40–42]. The remaining down-regulated miRNAs were all previously reported to be associated with CRC; for instance, the tumor suppressor MIR-10 family (4 genes), MIR-15, MIR-192 and MIR-194[16,35] have been implicated in CRC progression. The identification of miRNAs previously reported to be deregulated in CRC and associated with relevant signaling pathways provides confidence in our analytical approach.

Using the same analytical strategy, a metastatic signature was identified, which included five miRNAs that were differentially expressed in at least two metastatic sites compared to pCRC. Of these, Mir-210_3p and Mir-191_5p were up-regulated at all three metastatic sites, which would suggest a strong metastasis-related significance. Mir-210_3p is in the literature known as the "hypoxamiR" because of its key involvement in the cellular response to hypoxia. The Mir-210_3p promoter has a binding site for hypoxia inducible transcription factors HIF-1α and HIF-2α[43] which coordinate cellular responses to hypoxic stress, including regulation of key pathways in metastasis, such as angiogenesis, cell proliferation, differentiation and apoptosis[44]. Increased Mir-210_3p expression, which was additionally validated by qPCR in a separate set of samples, therefore suggests hypoxic stress to be a common feature of CRC metastasis, irrespective of metastatic site. In contrast, the role of Mir-191_5p in metastasis is less well established, and this miRNA is therefore an interesting candidate for further follow-up studies. Additionally, three miRNAs, Mir-8-P1b_3p, Mir-1307_5p, and Mir-155_5p, were up-regulated at two of three metastatic sites. Mir-8-P1b_3p (*miR-141*) is a member of the MIR-8 (*miR-200*) family, which is strongly involved in epithelial to mesenchymal transition (EMT) and targets transcription factors ZEB1 and ZEB2, which in turn suppress E-cadherin expression[45,46]. In our data, Mir-8-P1b_3p was down-regulated in pCRC relative to nCR, while up-regulated in mLi and mLu (Fig. 3). This fits well with the concept that tumors

undergo EMT as part of tumorigenesis at the primary site, while in the established metastasis the inverse process,

mesenchymal to epithelial transition, is necessary to establish growth in the new metastatic microenvironment.

Mir-155_5p, which was preferentially expressed in mLu and PM, is reported to be specifically expressed in

lymphocytes and macrophages[22], possibly indicating higher abundance of these cell types in the metastases

compared to pCRC. Less is known about the biological activity of Mir-1307_5p, although reports have

suggested a role in lung adenocarcinoma proliferation[47] and as a predictor of hepatocellular carcinoma

metastasis[48]. The identification of this miRNA as highly expressed in mCRC points to its involvement in

metastasis, and Mir-1307_5p therefore represents another target for further studies. Several of the signature

miRNAs have previously been implicated in the metastatic process, providing evidence that their observed up-

regulation represents adaptations to survival metastatic site.

To further investigate potential biological implications of miRNA expression, GSE analyses were performed

to explore predicted effects on mRNA expression based on miRNAs that were differentially expressed at the

metastatic sites. Through these analyses, GO terms related to transcription were estimated to be less repressed at

the metastatic sites, and KEGG pathways "Axon Guidance", "Long-Term Potentiation" and "Long-Term

Depression" were estimated to be less repressed in mLu, possibly representing adaptations to challenges in this

microenvironment. However, the biology of mRNA repression by miRNAs is complex. The seed sequence of

any miRNA can usually target a large number of mRNAs, and an mRNA can potentially be targeted for

repression by multiple miRNAs[5]. The functional role of these miRNAs will therefore be dependent on the local

cellular contexture, and because the net effect on mRNA expression is highly unpredictable, GSE analysis

cannot replace experimental validation. Furthermore, a miRNA may potentially have abnormal expression in

multiple diseases, such as Mir-21, which has been shown to be up-regulated in 29 different diseases in addition

to CRC[49]. Therefore, the functional role of the differentially expressed miRNAs is still not clear, but this work

still represents an excellent starting point for further studies, as well as a basis for interpreting the, often

contradictory, existing literature.

In order to obtain results that are physiologically relevant, the absolute expression level of a miRNA is an

important consideration. A cut-off level of 100 RPM was therefore applied as a minimum expression level to be

included in these analyses[20]. Several of the differentially expressed miRNAs were expressed at very high

absolute levels, for instance with Mir-191_5p, Mir-8-P1b_3p and Mir-10-P1a_5p all being expressed at >1000

RPM in both pCRC and mCRC. Expression levels of this magnitude strongly suggest that alterations in

expression of these miRNAs will influence target mRNA levels. Mir-10-P1a_5p is of particular interest, being

highly and differentially expressed in mLi compared to pCRC (164 925 RPM and 97 123 RPM, respectively),

suggesting specific adaptations to the unique conditions of this microenvironment. Previous experimental

evidence suggests that the Mir-10-P1a_5p paralogue, Mir-10-P1b_5p, is associated with enhanced metastatic

capability by down-regulation of metastasis suppressor Hoxd10[50–52]. Since Mir-10-P1a_5p and Mir-10-P1b_5p

share the same seed sequence (ACCCUGU), the range of mRNA targets and functional roles would be expected to

be similar.

In clinical cancer studies, the most commonly available material will be bulk tissue samples, and a tumor

biopsy will therefore contain a variable amount of cancer cells of epithelial origin together with a mixture of cell

types from the "host tissue" (such as fibroblasts, endothelial cells, myocytes, blood cells and immune cells). The

importance of keeping cell specificity in mind when interpreting miRNA expression levels is illustrated by the

high levels of hepatocyte specific Mir-122_5p in liver metastases compared to pCRC, which is likely due to the

presence of hepatocytes in the metastatic tissues, and do not indicate that this is a metastasis biomarker.

Similarly, Mir-143_3p/Mir-145_5p were previously suggested to be tumor suppressors due to apparent "down-

regulation" in pCRC relative to nCR[53]. In our datasets, both miRNAs were detected at much lower levels in

pCRC relative to nCR, but since these miRNAs are exclusively expressed in mesenchymal cells, the low

expression in pCRC can only be explained by mesenchymal cells being less frequently present in the colorectal

tumors than in the normal colorectal wall[54,55]. Expression of known cell-type specific miRNAs could also inform

on the relative composition of cell types in the different tissues, and as expected, the three tumor adjacent tissues

analyzed clustered separately and had distinct miRNA expression patterns. For instance, the epithelial specific

miRNAs, such as Mir-8-P2a_3p (miR-200b) and Mir-8-P2b_3p (miR-200c), had similar levels in the malignant

tissues, which is consistent with the intestinal epithelial origin of the cancer cells. This means that considering

background expression was a necessary step to ensure that cell composition effects would not confound the

differential expression analysis, and failure to do so would likely have led to incorrect identification of miRNAs

not associated with metastasis. Taken together, when analyzing bulk tissue samples, it is therefore necessary to

consider the composition of the "host tissue", which in this study was handled by defining and correcting for

background expression and by analyzing tumor samples from different metastatic sites as separate entities.

In addition to applying a 100 RPM cut-off level, correcting for background expression, and analyzing

metastatic locations as separate entities, a number of other methodological improvements were implemented in

this work. A systematic effort was made to ensure the quality of the analyses and making the data and the

analytical pipeline transparent and available to other researchers in the interest of reproducibility. A novel

quality assessment step was introduced in the analysis, using the recently developed miRTrace algorithm,

resulting in the exclusion of two previous studies[14,29]. Furthermore, our recently established database

MirGeneDB, containing only *bona fide* miRNA genes, was used as a reference for mapping miRNA reads to the

human genome, representing a further level of precision compared to the previously more commonly used

repository, miRBase[56]. Regretfully, miRBase has been shown to contain a proportion of incorrect miRNA

annotations[19], and including such sequences in the analyses could potentially lead to weakening the power of

statistical analyses and result in misleading conclusions. Furthermore, to facilitate reproducibility, a common set

of bioinformatics tools were used, and raw NGS datasets from all studies were run through an identical pipeline,

from postprocessing and read alignment to miRNA gene counting, using the latest version of miRge[28], a

bioinformatics pipeline specifically developed with miRNAs in mind. Combining all these measures and

specialized tools, the resulting set of differentially expressed miRNAs between pCRC and mCRC therefore

highly likely represents a reliable mCRC signature.

In summary, this work represents the most comprehensive attempt to characterize miRNA expression in

mCRC. An unbiased, stringent and transparent bioinformatics approach was developed and applied to a large

compilation of new and previously published NGS datasets to identify miRNAs associated with metastatic

progression in CRC. Comparison of pCRC and nCR replicated many previous findings of up- and down-

regulation of well-known oncomiRs and tumor-suppressor miRNAs, supporting the analytical strategy.

Correction for background expression was performed, and tumor samples from different metastatic sites were

analyzed as separate entities using a 100 RPM expression cut-off level to ensure biological relevance. A

metastatic signature containing five miRNAs that were up-regulated at multiple metastatic sites was identified

along with a number of miRNAs differentially expressed at individual metastatic sites. Many of these miRNAs

have previously been implicated as key players in the metastatic process, while for others, the involvement in

tumor cell adaptations at the distant site represent novel findings. The use of open science practices and the

biological relevance of the findings lend confidence in the resulting mCRC signature, which provides a starting-

point for further elucidation of the role of miRNAs in metastatic progression. The pipeline developed for this

analysis is freely available to other researchers to expand on the results presented in this work, as well for

exploring other cancer entities and disease settings. This work could therefore represent a first step to generate a

miRNA tissue expression "atlas" for researchers interested in miRNA biomarkers, for instance to identify

miRNAs that are specifically deregulated in a disease of interest.

Materials and Methods

Patient samples

Tissue samples were obtained from study specific biobanks: pCRC and nCR samples were from the LARC-EX study (NCT02113384); mLi and nLi samples from the OSLO-COMET trial (NCT01516710); mLu and nLu samples from our lung metastasis biobank (S-06402b) and PM samples from the Peritoneal Surface Malignancies biobank (NCT02073500). The studies were approved by the Regional Ethics Committee of South-East Norway, and patients were included following written informed consent. Patient samples were collected at the time of surgery and were snap frozen in liquid nitrogen at the time of collection and stored at -80°C. Samples were prepared and processed as described in[57].

RNA extraction and NGS

RNA was extracted using Qiagen Allprep DNA/RNA/miRNA universal kit, which simultaneously isolates genomic DNA and total RNA. RNA concentration was evaluated using a NanoDrop spectrophotometer (ThermoFisher, Waltham, Massachusetts, USA) and RNA integrity was evaluated using the Bioanalyzer RNA 6000 Nano kit (Agilent Technologies, Santa Clara, California, USA). MiRNA NGS libraries were then prepared using TruSeq Small RNA Library protocol and sequences using HiSeq 2500 High Throughput Sequencer (all from Illumina, San Diego, California, USA).

Identification of published NGS datasets

A literature search was conducted for the terms "microRNA + CRC + next generation sequencing" in different variations, and reviews were studied[16,35]. Publicly available datasets were downloaded from European Genome-phenome Archive (EGA), the Sequence Read Archive (SRA) and the Gene Expression Omnibus (GEO)[12–15,34].

Data processing, read alignment and gene counting

All datasets were processed using the same pipeline. miRTrace[27] was used for preprocessing and quality control (QC) of raw data (FASTQ files). Briefly, low-quality reads, defined as reads where less than 50% of nucleotides had a Phread quality score greater than 20 were discarded. 3p adapter sequences were trimmed, and reads made up of repetitive elements and reads shorter than 18nt were removed. After miRTrace QC, samples were excluded if < 25% of reads were between 20 and 25 nt, if > 75% of reads were discarded, or if < 10% of reads were

303 identified as miRNA. If > 50% of the datasets in a study failed the QC criteria, or if significant contamination

304 was detected, it was excluded. After QC, raw data processing, read alignment and gene counting was performed

305 using miRge3.0[28], with MirGeneDB2.0[26] as reference. To account for cross mapping of miRNA genes, miRge3.0

306 merges miRNA genes with very similar sequences into one annotation, reducing 537 human miRNA annotations

307 in MirGeneDB2.0 to a total of 389 unique annotations.

Analysis of global miRNA expression

309 For data visualization of global miRNA expression, read counts were normalized using the variance Stabilizing

310 Transformation() (VST) function from the DESeq2 package[58]. UMAP was used to visualize the similarity of

311 datasets on the global miRNA expression level. The UMAP algorithm reduces the 389 dimensions of unique

312 miRNA genes into two dimensions for visualization[59]. The umap R package was used, and datasets were

313 annotated by tissue.

Differential expression analysis

The differential expression analysis can be viewed in the supplementary R-markdown file, (Supplementary file

3).

Differential expression analysis was performed using DESeq2 (version 1.26)[58], which estimates LFC and its

standard error (SE), using raw, non-normalized read counts as input. Hypothesis testing was performed by a

Wald test against the null hypothesis LFC = 0, followed by the Benjamini-Hochberg procedure to correct for

multiple hypothesis testing, applying a false discovery rate (FDR) threshold of < 0.05. The LFC shrinkage

function in DESeq2, lfcShrink(), was enabled, to shrink fold changes for miRNAs with higher variance.

Differentially expressed miRNAs were filtered for relevance by requiring an LFC > 0.58 or LFC < -0.58, and

also requiring that at least one of the compared tissues had a mean expression > 100 RPM, a level previously

suggested as minimal cut-off for physiological activity[20]. MiRNAs known to be cell-type specific[22] were also

labelled for each miRNA. Information regarding normal tissue background expression levels was obtained by

analyzing differential expression between nCR versus nLi and nLu datasets, and pCRC versus nLi and nLu

datasets. Then, in the mCRC versus pCRC differential expression analysis, miRNAs differentially expressed in

the same direction in the corresponding normal tissue were not considered differentially expressed in the

respective metastatic site. For the PM tissue datasets, where no tumor adjacent tissue was available, the union of

nLi and nLu background expression was used.

qPCR validation

Eleven additional randomly chosen mLi and 11 pCRC tissue samples were selected for qPCR validation of increased expression of Mir-210_3p in mLi compared to pCRC. Synthetic RNA Spike-Ins UniSp2, UniSp4 and UniSp5 (Qiagen, Düsseldorf, Germany Cat. No. 339390) were added pre-isolation. RT-PCR was done with miRCURY LNA RT Kit (Qiagen Cat. No. 339340), adding UniSp6 and cel-miR-39-3p RNA Spike-Ins (Qiagen Cat. No 339390). qPCR was done using Qiagen miRCURY SYBR Green Kit (Qiagen Cat. No. 339345) and miRCURY LNA miRNA PCR Assays (Qiagen Cat. No. 339306). Mir-103 (Qiagen primer Cat. No. YP00204306) was used as reference miRNA, while the Mir-210_3p primer was Qiagen Cat. No. YP00204333. Two PCR replicates were run per primer assay. Mir-210_3p Cq values were normalized to the reference miRNA, to obtain the dCq value. Welch two-sided t-test was used to compare the dCq values, to test if the expression levels were different between mLi and pCRC. (Supplementary file 5).

Analysis of cell-type specific miRNAs

Two analyses were performed to assess differences in expression levels of 45 previously validated cell-type specific miRNAs[22], and thereby infer differences in cell composition in the tissues. A heatmap using z-scores of RPM values per miRNA was made to illustrate the relative expression of each miRNA in the tissues (Fig. 4A). Principle component analysis (PCA) plots were made with the FactoMineR PCA() function to illustrate which miRNAs were more or less prevalent in each tissue. Welch two-sided t-test comparing mean VST values between two groups of tissues was done to assess if the observed differences in cell-type specific miRNA levels were statistically significant (Supplementary file 4).

Gene set enrichment analysis

Gene set enrichment analysis (GSEA) was performed using RBiomirGS[60] on GO Molecular Function, Cellular Component and Biological Process, and KEGG pathways, downloaded from https://www.gsea-msigdb.org/gsea/msigdb/collections.jsp. miRNA to mRNA predicted interactions were combined with data from the miRNA differential expression analysis (DESeq2 estimated LFC and FDR values for each mCRC site versus pCRC). The input values to RBiomirGS were corrected for normal background expression, reducing LFC towards 0 and increasing FDR towards 1, depending on the magnitude of the normal background. All miRNAs were then given an S_{miRNA} score -log10 p-value * sign($\log_2$FC)., and for each mRNA, a S_{mRNA} score was calculated by summing up the S_{miRNA} scores of all the predicted miRNA to mRNA interactions. The S_{mRNA}

359   scores were then used to perform logistic regression, which provides likelihoods of gene sets being more or less

360   suppressed due to differential expression of miRNAs.

Acknowledgements

The authors thank Michael Hackenberg for help with data acquisition and Marc Halushka for discussions.

This work was supported by the South-Eastern Norway Regional Health Authority [grant numbers 2014041 to KF, 2018014 to EH], the Norwegian Cancer Society [grant number 4499184 to CLA] and the Research Council of Norway [grant number 218325].

Competing interests

The authors have declared no competing interests.

References

Ferlay J *et al.* Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015; **136**: E359-86.

Nguyen DX, Bos PD, Massagué J. Metastasis: from dissemination to organ-specific colonization. *Nat Rev Cancer* 2009; **9**: 274–284.

Riihimäki M, Hemminki A, Sundquist J, Hemminki K. Patterns of metastasis in colon and rectal cancer. *Sci Rep* 2016; **6**: 29765.

Massague J, Obenauf AC. Metastatic colonization by circulating tumour cells. *Nature* 2016; **529**: 298–306.

Flatmark K, Høye E, Fromm B. microRNAs as cancer biomarkers. Scandinavian Journal of Clinical and Laboratory Investigation. 2016; **76**: S80–S83.

Arnold M, Sierra MS, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global patterns and trends in colorectal cancer incidence and mortality. *Gut* 2017; **66**: 683–691.

Bartel DP. Metazoan MicroRNAs. *Cell* 2018; **173**: 20–51.

Dalmay T, Edwards DR. MicroRNAs and the hallmarks of cancer. *Oncogene* 2006; **25**: 6170–6175.

Jung M *et al.* Robust microRNA stability in degraded RNA preparations from human tissue and cell samples. *Clin Chem* 2010; **56**: 998–1006.

Volinia S *et al.* A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Natl Acad Sci U S A* 2006; **103**: 2257–2261.

Baffa R *et al.* MicroRNA expression profiling of human metastatic cancers identifies cancer gene targets. *J Pathol* 2009; **219**: 214–221.

Neerincx M *et al.* MiR expression profiles of paired primary colorectal cancer and metastases by next-generation sequencing. *Oncogenesis* 2015; **4**: e170.

Röhr C *et al.* High-throughput miRNA and mRNA sequencing of paired colorectal normal, tumor and metastasis tissues and bioinformatic modeling of miRNA-1 therapeutic applications. *PLoS One* 2013; **8**: e67461.

Goossens-Beumer IJ *et al.* MicroRNA classifier and nomogram for metastasis prediction in colon cancer. *Cancer Epidemiol Biomarkers Prev* 2015; **24**: 187–197.

Schee K *et al.* Deep Sequencing the MicroRNA Transcriptome in Colorectal Cancer. PLoS ONE. 2013; **8**: e66165.

Cekaite L, Eide PW, Lind GE, Skotheim RI, Lothe RA. MicroRNAs as growth regulators, their function and biomarker status in colorectal cancer. *Oncotarget* 2016; **7**: 6476–6505.

To KK, Tong CW, Wu M, Cho WC. MicroRNAs in the prognosis and therapy of colorectal cancer: From bench to bedside. *World J Gastroenterol* 2018; **24**: 2949–2973.

Git A *et al.* Systematic comparison of microarray profiling, real-time PCR, and next-generation sequencing technologies for measuring differential microRNA expression. *RNA* 2010; **16**: 991–1006.

Fromm B *et al.* A Uniform System for the Annotation of Vertebrate microRNA Genes and the Evolution of the Human microRNAome. *Annu Rev Genet* 2015; **49**: 213–242.

Mullokandov G *et al.* High-throughput assessment of microRNA activity and function using microRNA sensor and decoy libraries. *Nat Methods* 2012; **9**: 840–846.

21   Witwer KW, Halushka MK. Toward the promise of microRNAs – Enhancing reproducibility and rigor in
microRNA research. *RNA Biol* 2016; **13**: 1103–1116.

22   McCall MN *et al.* Toward the human cellular microRNAome. *Genome Res* 2017; **27**: 1769–1781.

23   Rie D de *et al.* An integrated expression atlas of miRNAs and their promoters in human and mouse. *Nat*
Biotechnol 2017. doi:10.1038/nbt.3947.

24   Juzenas S *et al.* A comprehensive, cell specific microRNA catalogue of human peripheral blood. Nucleic
Acids Research. 2017; **45**: 9290–9301.

25   Mudduluru G *et al.* A Systematic Approach to Defining the microRNA Landscape in Metastasis. *Cancer*
Res 2015; **75**: 3010–3019.

26   Fromm B *et al.* MirGeneDB 2.0: the metazoan microRNA complement. *Nucleic Acids Res* 2020.
doi:10.1093/nar/gkz885.

27   Kang W, Eldfjell Y, Fromm B, Estivill X, Biryukova I, Friedländer MR. miRTrace reveals the organismal
origins of microRNA sequencing data. *Genome Biol* 2018; **19**: 213.

28   Patil AH, Halushka MK. miRge3.0: a comprehensive microRNA and tRF sequencing analysis pipeline.
Cold Spring Harbor Laboratory. 2021; : 2021.01.18.427129.

29   Rohr C *et al.* High-throughput miRNA and mRNA sequencing of paired colorectal normal, tumor and
metastasis tissues and bioinformatic modeling of miRNA-1 therapeutic applications. *PLoS One* 2013; **8**:
e67461.

30   Sun Y, Wang L, Guo S-C, Wu X-B, Xu X-H. High-throughput sequencing to identify miRNA biomarkers
in colorectal cancer patients. Oncology Letters. 2014; **8**: 711–713.

31   Sun G, Cheng Y-W, Lai L, Huang T-C, Wang J, Wu X *et al.* Signature miRNAs in colorectal cancers were
revealed using a bias reduction small RNA deep sequencing protocol. *Oncotarget* 2016; **7**: 3857–3872.

32   Liang G *et al.* Deep sequencing reveals complex mechanisms of microRNA deregulation in colorectal
cancer. *Int J Oncol* 2014; **45**: 603–610.

33   Hamfjord J *et al.* Differential expression of miRNAs in colorectal cancer: comparison of paired tumor
tissue and adjacent normal mucosa using high-throughput sequencing. *PLoS One* 2012; **7**: e34150.

34   Selitsky SR *et al.* Transcriptomic Analysis of Chronic Hepatitis B and C and Liver Cancer Reveals
MicroRNA-Mediated Control of Cholesterol Synthesis Programs. mBio. 2015; **6**. doi:10.1128/mbio.01500-
15.

35   Strubberg AM, Madison BB. MicroRNAs in the etiology of colorectal cancer: pathways and clinical
implications. *Dis Model Mech* 2017; **10**: 197–214.

36   Schee K, Boye K, Abrahamsen TW, Fodstad O, Flatmark K. Clinical relevance of microRNA miR-21,
miR-31, miR-92a, miR-101, miR-106a and miR-145 in colorectal cancer. *BMC Cancer* 2012; **12**: 505.

37   Zhang W *et al.* MicroRNA-301a promotes migration and invasion by targeting TGFBR2 in human
colorectal cancer. *J Exp Clin Cancer Res* 2014; **33**: 113.

38   Ma X *et al.* Modulation of tumorigenesis by the pro-inflammatory microRNA miR-301a in mouse models
of lung cancer and colorectal cancer. *Cell Discov* 2015; **1**: 15005.

39   Guo X *et al.* miR-181d and c-myc-mediated inhibition of CRY2 and FBXL3 reprograms metabolism in
colorectal cancer. *Cell Death Dis* 2017; **8**: e2958.

40   Scaravilli M *et al.* MiR-1247-5p is overexpressed in castration resistant prostate cancer and targets
MYCBP2. The Prostate. 2015; **75**: 798–805.

41 Chen L, Tang Y, Wang J, Yan Z, Xu R. miR-421 induces cell proliferation and apoptosis resistance in
human nasopharyngeal carcinoma via downregulation of FOXO4. *Biochem Biophys Res Commun* 2013;
435: 745–750.

42 Zhao J *et al.* MicroRNA-7: a promising new target in cancer therapy. *Cancer Cell Int* 2015; **15**: 103.

43 Chan YC, Banerjee J, Choi SY, Sen CK. miR-210: the master hypoxamir. *Microcirculation* 2012; **19**: 215–
223.

44 Rankin EB, Giaccia AJ. Hypoxic control of metastasis. *Science* 2016; **352**: 175–180.

45 Pencheva N, Tavazoie SF. Control of metastatic progression by microRNA regulatory networks. *Nat Cell
Biol* 2013; **15**: 546–554.

46 Gregory PA *et al.* The miR-200 family and miR-205 regulate epithelial to mesenchymal transition by
targeting ZEB1 and SIP1. *Nat Cell Biol* 2008; **10**: 593–601.

47 Du X *et al.* MiR-1307-5p targeting TRAF3 upregulates the MAPK/NF-κB pathway and promotes lung
adenocarcinoma proliferation. *Cancer Cell Int* 2020; **20**: 502.

48 Eun JW *et al.* Circulating Exosomal MicroRNA-1307-5p as a Predictor for Metastasis in Patients with
Hepatocellular Carcinoma. *Cancers* 2020; **12**. doi:10.3390/cancers12123819.

49 Jenike AE, Halushka MK. miR-21: a non-specific biomarker of all maladies. *Biomark Res* 2021; **9**: 18.

50 Ma L, Reinhardt F, Pan E, Soutschek J, Bhat B, Marcusson EG *et al.* Therapeutic silencing of miR-10b
inhibits metastasis in a mouse mammary tumor model. *Nat Biotechnol* 2010; **28**: 341–347.

51 Ma L, Teruya-Feldstein J, Weinberg RA. Tumour invasion and metastasis initiated by microRNA-10b in
breast cancer. *Nature* 2007; **449**: 682–688.

52 Wang Y, Li Z, Zhao X, Zuo X, Peng Z. miR-10b promotes invasion by targeting HOXD10 in colorectal
cancer. *Oncol Lett* 2016; **12**: 488–494.

53 Michael MZ, O' Connor SM, van Holst Pellekaan NG, Young GP, James RJ. Reduced accumulation of
specific microRNAs in colorectal neoplasia. *Mol Cancer Res* 2003; **1**: 882–891.

54 Chivukula RR *et al.* An essential mesenchymal function for miR-143/145 in intestinal epithelial
regeneration. *Cell* 2014; **157**: 1104–1116.

55 Kent OA, McCall MN, Cornish TC, Halushka MK. Lessons from miR-143/145: the importance of cell-
type localization of miRNAs. *Nucleic Acids Res* 2014; **42**: 7528–7538.

56 Griffiths-Jones S, Grocock RJ, van Dongen S, Bateman A, Enright AJ. miRBase: microRNA sequences,
targets and gene nomenclature. *Nucleic Acids Res* 2006; **34**: D140-4.

57 Østrup O *et al.* Molecular signatures reflecting microenvironmental metabolism and chemotherapy-induced
immunogenic cell death in colorectal liver metastases. Oncotarget. 2017; **8**.
doi:10.18632/oncotarget.19350.

58 Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with
DESeq2. Genome Biology. 2014; **15**. doi:10.1186/s13059-014-0550-8.

59 McInnes L, Healy J, Melville J. UMAP: Uniform Manifold Approximation and Projection for Dimension
Reduction. arXiv [stat.ML]. 2018.http://arxiv.org/abs/1802.03426.

60 Zhang J, Storey KB. RBiomirGS: an all-in-one miRNA gene set analysis solution featuring target mRNA
mapping and expression profile integration. *PeerJ* 2018; **6**: e4262.

Figure legends

Fig. 1 Global miRNA expression according to tissue of origin

UMAP cluster plot based on global miRNA expression, normalized by VST, from the DESeq2 bioconductor package, annotated by tissue of origin.

Uniform approximation and projection for dimension reduction (UMAP); varianceStabilizingTransformation (VST); primary colorectal cancer (pCRC); normal colorectal tissue (nCR); CRC liver metastasis (mLi); normal adjacent liver tissue (nLi); CRC lung metastasis (mLu); normal adjacent lung tissue (nLu); CRC peritoneal metastasis (PM).

Fig. 2 Differential expression analysis identifies miRNAs associated with metastatic colorectal cancer

a Volcano plots showing differentially expressed miRNAs in mLi, mLu and PM compared to pCRC. The horizontal axis shows LFC relative to pCRC, while the vertical axis shows -log10 FDR. The identified miRNAs were differentially expressed between pCRC and mCRC, with expression levels greater than 100 RPM in one of the tissues, and results were corrected for tumor adjacent background expression. **b** Bar plots of LFC of differentially expressed miRNAs in pCRC compared to the individual metastatic sites.

Log2 fold change (LFC); false discovery rate (FDR); reads per million (RPM); primary colorectal cancer (pCRC); normal colorectal tissue (nCR); CRC liver metastasis (mLi); normal adjacent liver tissue (nLi); CRC lung metastasis (mLu); normal adjacent lung tissue (nLu); CRC peritoneal metastasis (PM).

Fig. 3 Differential expression analysis identifies a five-gene miRNA signature up-regulated at multiple metastatic sites

a Bar plots showing LFC of the five signature miRNAs that were differentially expressed between pCRC and at least two mCRC sites. Mir-210_3p and Mir-191_5p were up-regulated at all three metastatic sites, while, Mir-8-P1b_3p was up-regulated in mLi and mLu, Mir-1307_5p in mLi and PM, and Mir-155 in mLu and PM, after

correcting for multiple hypothesis testing. Error bars represent 95% confidence intervals from DESeq2 estimated

SE of the LFC. **b** Scatter plots of Log10 RPM in each tissue.

Log2 fold change (LFC); reads per million (RPM); standard error (SE) primary colorectal cancer (pCRC);

normal colorectal tissue (nCR); CRC liver metastasis (mLi); normal adjacent liver tissue (nLi); CRC lung

metastasis (mLu); normal adjacent lung tissue (nLu); CRC peritoneal metastasis (PM).

Fig. 4 Cell-type specific miRNAs provide insight into the cellular composition of tissues

a Heatmap illustrating the expression of 25 cell-specific miRNAs for each tissue that had mean expression > 100

RPM in at least one tissue. The color scale indicates the z-score of RPM for each miRNA. **b** PCA plot based on

analysis of 25 cell-specific miRNA with expression>100 reads per million, VST normalized, colored by tissue.

Transparent points are individual samples, solid points represent the statistical mean and shaded ellipses is 95%

confidence of the mean of each tissue. **c** The correlation circle shows PCA loadings, which are the direction and

relative contribution the top 15 cell-type specific miRNAs had on clustering of each tissue.

Reads per million (RPM); principal component analysis (PCA); varianceStabilizingTransformation (VST);

primary colorectal cancer (pCRC); normal colorectal tissue (nCR); CRC liver metastasis (mLi); normal adjacent

liver tissue (nLi); CRC lung metastasis (mLu); normal adjacent lung tissue (nLu); CRC peritoneal metastasis

(PM).

Table 1 Differentially expressed miRNAs in mCRC compared to pCRC according to
metastatic site

Reads per million (RPM); log 2 fold change (LFC) estimated using DESeq2; standard error (SE); primary

colorectal cancer (pCRC); normal colorectal tissue (nCR); CRC liver metastasis (mLi); normal adjacent liver

tissue (nLi); CRC lung metastasis (mLu); normal adjacent lung tissue (nLu); CRC peritoneal metastasis (PM);

false discovery rate (FDR).

Availability

Data from current and previous studies can be found at EGA accession number EGAS00001001127 (14), GEO accession numbers: GSE57381 (31), GSE46622 (15) and GSE63119 (16), and SRA accession PRJNA397121 for datasets prepared in this study and (17). Bioinformatics pipeline can be found at (https://github.com/eirikhoye/mirna_pipeline)

Supplementary material

Supplementary file 1: miRTrace QC reports (.html)

Supplementary file 2: miRge3.0 count matrix and sample metadata (.csv)

Supplementary file 3: R-markdown: DESeq2 differential expression analysis (.html)

Supplementary file 4: R-markdown: code for figures (.html)

Supplementary file 5: R-markdown qPCR analysis (.html)

Supplementary file 6: R-markdown GSE results (.html)

Fig 1

pCRC
(n=120)

PM
(n=30)

mLu
(n=28)

nCR
(n=25)

nLu
(n=10)

mLi
(n=35)

UMAP2

nLi
(n=20)

UMAP1

Fig 2

a

b

mLi vs pCRC   mLu vs pCRC   PM vs pCRC

Signature miRNA   Differentially Expressed   Background Expression   Not Significant

Fig 3

Fig 4

a

Myocyte | Hepatocyte | Mesenchymal | Red Blood Cell | Lymphocyte | Macrophage | Neural | Epithelial | Endothelial | Melanocyte | RPE | Islet Cell | CD14+ Monocyte

Rows: PM, mLu, mLi, pCRC, nLu, nLi, nCR

Columns: Mir–133–P1_3p, Mir–122_5p, Mir–143_3p, Mir–145_5p, Mir–144_5p, Mir–451_5p, Mir–486_5p, Mir–146–P2_5p, Mir–342_3p, Mir–155_5p, Mir–150_5p, Mir–223_3p, Mir–24–P1_3p, Mir–128–P1_3p, Mir–192–P1_5p, Mir–375_3p, Mir–8–P2a_3p, Mir–8–P2b_3p, Mir–126_5p, Mir–204–P1_5p, Mir–335_5p, Mir–7–P1_5p, Mir–17–P1a_5p, Mir–15–P1a_5p, Mir–15–P1b_5p

VST Z-score: −1 0 1 2

b

Dim2 (13.4 %)

nLu, mLu, nCR, PM, pCRC, mLi, nLi

c

Mir-146-P2_5p, Mir-145_5p, Mir-24-P1_3p, Mir-342_3p, Mir-143_3p, Mir-126_5p, Mir-8-P2b_3p, Mir-451_5p, Mir-486_5p, Mir-7-P1_5p, Mir-8-P2a_3p, Mir-144_5p, Mir-17-P1a_5p, Mir-122_5p, Mir-192-P1_5p

nLu, mLu, nCR, pCRC, PM, mLi, nLi

Dim1 (23.3 %)

Table 1 Differentially expressed miRNAs in mCRC compared to pCRC according to metastatic site

MirGeneDB ID (miRBase ID)	pCRC RPM	mLi vs pCRC			mLu vs pCRC			PM vs pCRC		
		RPM	LFC (SE)	FDR	RPM	LFC (SE)	FDR	RPM	LFC (SE)	FDR
Mir-191_5p	13 630	23 253	0.74 (0.15)	3.67E-06	26 989	0.74 (0.16)	9.81E-06	20 429	0.61 (0.15)	2.42E-04
Mir-210_3p	291	685	1.26 (0.18)	4.73E-11	693	1.15 (0.19)	1.55E-08	491	0.71 (0.18)	3.37E-04
Mir-1307_5p	492	887	0.85 (0.17)	2.99E-06				797	0.62 (0.17)	8.29E-04
Mir-8-P1b_3p (mir-141-3p)	7518	11 957	0.64 (0.19)	4.07E-03	15 111	0.75 (0.21)	1.12E-03			
Mir-155_5p	387				733	0.76 (0.18)	7.24E-05	654	0.70 (0.17)	1.59E-04
Mir-10-P1a_5p (mir-10a-5p)	97 123	164 925	0.86 (0.19)	8.04E-05						
Mir-592_5p	75	137	0.98 (0.27)	4.88E-03						
Mir-1247_5p	69	127	0.85 (0.28)	2.58E-02						
Mir-425_5p	700	1117	0.73 (0.12)	2.90E-08						
Mir-486_5p	2013	1489	-0.66 (0.23)	3.47E-02						
Mir-142_5p	3453				7117	0.84 (0.18)	2.01E-05			
Mir-19-P2a/P2b_3p	1297				2345	0.80 (0.15)	9.38E-07			
Mir-374-P1_5p	115				217	0.72 (0.12)	9.07E-09			
Mir-19-P1_3p	379				597	0.65 (0.16)	3.42E-04			
Mir-423_5p	433				202	-1.17 (0.17)	1.35E-10			
Let-7-P1b_5p	1057				650	-0.86 (0.16)	2.55E-07			
Mir-197_3p	181				113	-0.74 (0.13)	1.76E-07			
Mir-92-P1c_3p	1398				981	-0.72 (0.18)	2.22E-04			
Mir-362-P2/P4_3p	441				302	-0.65 (0.14)	2.36E-05			
Mir-221-P2_3p	1633				1101	-0.60 (0.14)	8.50E-05			
Mir-506-P3_3p	6							112	2.69 (0.29)	6.35E-09
Mir-506 P4a1/P4a2/P4b_3p	14							191	2.72 (0.29)	5.75E-11
Mir-154-P9_3p	253							455	0.85 (0.14)	1.43E-08
Mir-127_3p	2263							2888	0.61 (0.20)	1.42E-02
Mir-154-P36_3p	195							330	0.74 (0.13)	7.13E-07
Mir-223_3p	619							280	-0.94 (0.21)	4.62E-05

Reads per million (RPM); log 2 fold change (LFC) estimated using DESeq2; standard error (SE); primary colorectal cancer (pCRC); normal colorectal tissue (nCR); CRC liver metastasis (mLi); normal adjacent liver tissue (nLi); CRC lung metastasis (mLu); normal adjacent lung tissue (nLu); CRC peritoneal metastasis (PM); false discovery rate (FDR)