Neural population dynamics underlying expected value computation

Hiroshi Yamada*[123], Yuri Imaizumi[4], Masayuki Matsumoto[123]

[1]Division of Biomedical Science, Faculty of Medicine, University of Tsukuba, 1-1-1 Tenno-dai, Tsukuba, Ibaraki 305-8577, Japan.

[2]Graduate School of Comprehensive Human Sciences, University of Tsukuba, 1-1-1 Tenno-dai, Tsukuba, Ibaraki 305-8577, Japan

[3]Transborder Medical Research Center, University of Tsukuba, 1-1-1 Tenno-dai, Tsukuba, Ibaraki 305-8577, Japan

[4]Medical Sciences, University of Tsukuba, 1-1-1 Tenno-dai, Tsukuba, Ibaraki 305-8577, Japan.

*Address correspondence to

Hiroshi Yamada, Ph.D.

Division of Biomedical Science, Faculty of Medicine, University of Tsukuba

1-1-1 Tennodai, Tsukuba, Ibaraki, 305-8577 Japan.

TEL/FAX: 81-29-853-6013, e-mail: h-yamada@md.tsukuba.ac.jp

Abstract

Computation of expected values, i.e., probability times magnitude, seems to be a dynamic integrative process performed in the brain for efficient economic behavior. However, neural dynamics underlying this computation remain largely unknown. We examined (1) whether four core reward-related regions detect and integrate the probability and magnitude cued by numerical symbols and (2) whether these regions have distinct dynamics in the integrative process. Extractions of mechanistic structure of neural population signal demonstrated that expected-value signals simultaneously arose in central part of orbitofrontal cortex (cOFC, area 13m) and ventral striatum (VS). These expected-value signals were incredibly stable in contrast to weak and/or fluctuated signals in dorsal striatum and medial OFC. Notably, temporal dynamics of these stable expected-value signals were unambiguously distinct: sharp and gradual signal evolutions in cOFC and VS, respectively. These intimate dynamics suggest that cOFC and VS compute the expected-values with unique time constants, as distinct, partially overlapping processes.

Keywords: Monkey, Neural population dynamics, expected values, integration, rewards

Introduction

Economic behavior requires a reliable perception of the world for maximizing benefit (Houthakker, 1950; Samuelson, 1950; Savage, 1954; Von Neumann & Morgenstern, 1944). Such maximization is achieved in the first place by computing expected values (i.e. probability times magnitude) (Glimcher et al., 2008) in the brain, which seems to be a dynamic process to detect and integrate the probability and magnitude to yield expected value signals. Indeed, humans and animals behave as if they compute the expected values (Glimcher et al., 2008; Kahneman & Tversky, 1979; Stephens & Krebs, 1986). One salient example, discovered over a century ago and repeatedly measured, is human economic behavior, in which a series of models originating from the standard theory of economics (Von Neumann & Morgenstern, 1944) has been developed to describe efficient economic behavior. Despite the ubiquity of this phenomenon, a dynamic integrative process to compute the expected values from probability and magnitude remain largely unknown.

Substantial previous researches in the past two decades have shown that many regions of the brain process rewards in terms of signaling the probability and/or magnitude, mostly during economic choice behavior (Barraclough et al., 2004; Eshel et al., 2016; Hirokawa et al., 2019; Lopatina et al., 2016; Ma & Jazayeri, 2014; Platt & Glimcher, 1999; Roesch et al., 2009; Rudebeck & Murray, 2014; Tobler et al., 2005; Xie & Padoa-Schioppa, 2016; Yamada et al., 2018; Zhou et al., 2019). In these previous literature, the computation of expected values are assumed to be processed by these regions without their neural dynamics, in line

with an expected value theory shared across multiple disciplines (Glimcher et al.,

2008; Stephens & Krebs, 1986; Sutton & Barto, 1998; Von Neumann &

Morgenstern, 1944). Neuroimaging studies in humans and non-human primates

have also suggested that multiple brain regions in the reward circuitry (Haber &

Knutson, 2010) are involved in this computational process (Fouragnan et al., 2019;

Howard et al., 2015; Howard & Kahnt, 2017; Hsu et al., 2009; Levy & Glimcher,

2012; J. O'Doherty et al., 2004; Papageorgiou et al., 2017; Tom et al., 2007),

although the underlying neural mechanism has not been elucidated because of the

limited time resolution of current neuroimaging techniques (Goense & Logothetis,

2008; Milham et al., 2018). These reward-related brain regions may employ the

expected-value computation, however, none have captured and compared the

temporal aspects of neural activities in the multiple candidate brain regions. These

observations have led us to test the hypothesis that neural population dynamics

within subsecond-order time resolutions (Chen & Stuphorn, 2015; Churchland et

al., 2012; Mante et al., 2013; Murray et al., 2017; Takei et al., 2017) play key roles

in the expected-value computation, that is the detection and integration of the

probability and magnitude on multiple neural population ensembles.

We targeted reward-related cortical and subcortical structures of non-human

primates (Haber & Knutson, 2010) including the central orbitofrontal cortex (cOFC,

area 13M), the medial orbitofrontal cortex, (mOFC, area 14O), the dorsal striatum

(DS, caudate nucleus), and ventral striatum (VS), all of which are known to

represent the neural correlate with stimulus values during economic choice

behavior. To dissociate the integrative process to compute expected values from

a choice process employed during economic choices (Chen & Stuphorn, 2015;

Gardner et al., 2019; Yoo & Hayden, 2020), we recorded the neural activity in a

non-choice situation; monkeys perceive and compute the expected values from

probability and magnitude symbols. We used a recently developing mathematical

approach, called state space analysis (Chen & Stuphorn, 2015; Churchland et al.,

2012; Mante et al., 2013; Murray et al., 2017), to test how the expected-values

computation is processed within each of the four neural population ensembles in

an order of 10^{-2}-second time resolution. We found that the cOFC and VS neural

populations coincidently detect and integrate the probability and magnitude

information to yield the expected value signals on neural population assembles.

These signals are maintained in a stable state, though the cOFC signal sharply

develops in contrast to gradual evolution of the VS signal. Contrastingly, the

signals in the mOFC and DS were weak and/or fluctuated . Thus, we conclude that

the cOFC and VS neural populations employ a common integrative computation

of expected values from probability and magnitude.

Results

Task and behavior in monkeys

Based on the vast literature on human behavioral economics and by harnessing

the well-developed visual and cognitive abilities in non-human primates, we

designed a behavioral task in which monkeys estimated the expected values of

rewards from numerical symbols, mimicking events performed by humans. The

task involved a visual pie-chart that included two numerical symbols associated

with the probability and magnitude of fluid rewards with great precision. After

monkeys fixated the central grey target, a visual pie-chart comprising green and

blue pies was presented (Figure 1A). The number of the green pies indicated the

magnitude of fluid rewards in 0.1 mL increments (0.1-1.0 mL). Simultaneously, the

number of the blue pies indicated the probability of reward in 0.1 increments (0.1-

1 where 1 indicates a 100% chance). After a 2.5 second delay, the visual pie-chart

disappeared, and a reward outcome was provided to the monkeys with the

indicated amount and probability of reward, unless no reward was given. Under

this experimental condition, the expected values of rewards are defined as the

probability multiplied by the magnitude cued by the numerical symbols (Table. S1).

To examine whether the monkeys accurately perceived the expected values

from the numerical symbols for probability and magnitude, we applied a choice

task to the monkeys (Figure 1B). The two monkeys exhibited a near-efficient

performance in selecting a larger expected value option among two alternatives

during choice trials (Figure 1C). We examined which of the following three

behavioral models best described the monkey's behavior: model 1 (M1), monkeys

make choices based on the number of pies; model 2 (M2), monkeys make choices

based on the probability and magnitude; and model 3 (M3), monkeys make choices

based on the expected values (Table. S1). Comparisons of the model

performances based on Akaike's Information Criterion (AIC) and Bayesian

Information Criterion (BIC) (Burnham & Anderson, 2004) revealed that the model

3 best explained the monkey's behavior as indicated by the smallest AIC and BIC

values (Figure S1). The model 3 consistently showed the highest pseudo r-

squared values in each monkey (Figure 1D). These results indicated that the

monkeys utilize the expected values estimated from the numerical symbols for

probability and magnitude.

Neural population data

We constructed four pseudo-populations of neurons by recording single-neuron

activity during the single cue task (Figure 1A) from the DS (194 neurons), VS (144

neurons), cOFC (190 neurons) and mOFC (158 neurons) (Figure 1E). The four

constructed neural populations exhibited changes in their activities at different

times in the task trials (Figure S2A). Approximately 40 to 50 % of neurons in the

four neural populations demonstrated peak activity during the cue presentation

(Figure S2B, Chi-squared test, n = 686, P = 0.32, X^2 = 3.55, df = 3), with several

basic firing properties (Figure S2C-F). Strong peak activities with short-latency

were observed in the cOFC (Kruskal-Wallis test, Latency: Figure S2C, n = 314, P

= 0.013, H = 10.9, df = 3, Peak firing rate: Figure S2D, n = 314, P < 0.001, H =

32.1, df = 3). Activity changes were slow in the mOFC (Figure S2E, Kruskal-Wallis

test, n = 314, P = 0.003, H = 13.4, df = 3). Baseline firing rates were the highest in

the cOFC (Figure S2F, Kruskal-Wallis test, n = 686, P < 0.001, H = 60.3, df = 3).

In short, the strong activity with short-latency frequently occurred in the cOFC in

contrast to the phasic activity at various latencies in the DS and VS and relatively

tonic and gradual activity changes in the mOFC.

Conventional analyses for detecting the expected-value signals

We first applied common conventional analyses (linear regression, model selection

based on AIC, and model selection based on BIC) to the four neural populations

to examine neural modulations by the probability, magnitude, and expected values

at a single neuron level (see Methods). During a fixed one second time window

after the cue onset, these analyses showed that neurons in all the four brain

regions signal the probability, magnitude, and expected values to some extent

(Figure 2). For example, neurons signaling the expected values were found in each

brain region (Figure 2A-H, Neural modulation pattern was defined as the Expected

value type based on all the three analyses. Regression coefficients for probability

and magnitude, A-B, DS, 6.17, $P < 0.001$ and 2.54, $P = 0.007$, C-D, VS: 7.14, $P <$

0.001 and 6.71, $P < 0.001$, E-F, cOFC, 8.55, $P < 0.001$ and 11.1, $P < 0.001$), while

the classification of neural modulation types was dependent on the methods in

some neurons (Regression coefficients for probability and magnitude, G-H, mOFC,

1.76, $P = 0.032$, and 0.50, $P = 0.54$, AIC-based model selection: Expected value

type, linear regression: Probability type, and BIC-based model selection: non-

modulated type). In addition, neurons signaling probability or magnitude were also

observed in each brain region (Figure 2I-L, blue and green). Moreover, a subset of

neurons in the cOFC and VS signaled high-risk high-return or low-risk low-return

(Figure S3). These neurons were characterized as a strong activity elicited when

the cue indicated a low probability and large magnitude (hence, high-risk high-

return, Figure 2J and K, brown). Indeed, each neural population was composed of

a mixture of these signals (Figure 2I-L), indicating that signals for the expected

values and its components (i.e., probability and magnitude) appeared in each

neural population during one second after the cue onset. If we analyzed these neural modulation patterns through a task trial (Figure 2M-P), there was no significant difference in the proportions of neural modulation types, except the VS (chi-squared test: DS, n = 104, df = 75, X^2 = 91.4, P = 0.096; VS, n = 104, df = 75, X^2 = 98.2, P = 0.037; cOFC, n = 104, df = 75, X^2 = 83.2, P = 0.242; mOFC, n = 104, df = 75, X^2 = 79.0, P = 0.353), indicating that these conventional analyses did not clearly provide evidence of temporal dynamics in the modulation patterns of neural populations. Thus, we developed an analytic tool to examine how the detection and integration of probability and magnitude are processed within these neural population ensembles.

State space analysis for detecting neural population dynamics

The state space analysis can provide temporal dynamics of neural population signal related to cognitive and motor performances (Churchland et al., 2012; Mante et al., 2013). In our lottery task, such population dynamics can describe how the expected values are evolved within the neural population ensembles. To describe how each neural population detects and integrates the probability and magnitude into the expected values, we represented each neural population signal as a time-series of vectors in the space of probability and magnitude in two steps. First, we used a linear regression to project a time series of each neural activity into the regression subspace composed of the probability and magnitude in each neural population. This step captures the across-trial variance caused by probability and magnitude moment-by-moment at a population level. Second, we applied a

principal component analysis (PCA) to the time series of neural activities in the

regression subspace in each neural population. This step determines the main

feature of the neural population signal moment-by-moment in the space of

probability and magnitude. Because activations are dynamic and changing over

time, the analysis identified whether and how signal transformations occurred to

convert the probability and magnitude into the expected values as a time-series of

eigenvectors (Figure 3A). The directions of these eigenvectors capture the

expected values as an angle moment-by-moment at a population level (Figure 3B).

We evaluated properties of the eigenvectors for the first and second principal

components (PC1 and PC2) in each neural population in terms of the vector angle,

size, and deviance (Figure 3C). A stable population signal is described as a small

variation in the eigenvector properties through a trial, whereas an unstable

population signal is described as a large variation in the eigenvector's properties.

It must be noted that our procedure is a variant of the state space analysis in line

with the use of linear regression to identify dimensions of a neural population signal

(Chen & Stuphorn, 2015; Mante et al., 2013). It was not, however, aimed at

projecting the population activity as trajectories in multi-dimensional space.

Stable and unstable neural population signals

The eigenvector analyses yielded clear differences in neural population signals

among the four populations (Figure 4A-D). We first confirmed adequate

performance of the state space analysis indicated by the percentages of variance

explained in each population (Figure 4A). The VS population exhibited the highest

performance among the four neural populations, which was followed by the cOFC

and DS populations, with the lowest performance exhibited by the mOFC

population. Thus, performances to process the probability and magnitude

information was distinct among the four neural populations.

To characterize the whole structure of each neural population signal, we

analyzed the aggregated properties of the eigenvectors without a temporal order

of the vectors through a task trial. We first examined the properties of the

eigenvectors for PC1. The aggregated eigenvectors revealed both stable and

unstable neural population signals during the cue presentation (Figure 4B, green).

The VS population exhibited the highest performance (37%) and the eigenvectors

for PC1 were stable throughout cue presentation, with directions close to 45º, i.e.,

the expected value (Figure 4B, VS, vector angle, PC1, mean ± s.e.m., 37.5º ± 0.98,

7.5º difference from 45º). The cOFC population also exhibited a stable expected-

value signal with the second-best performance (31%), but they were deviated more

from the ideal expected-value signal (Figure 4B, cOFC, vector angle, PC1, mean

± s.e.m., 59.4º ± 1.16, 14.4º difference from 45º, Wilcoxon rank-sum test, n = 52,

W = 122, P < 0.001). Stability of the vectors was the best in the VS and cOFC, as

indicated by the smallest deviation from its mean vector among the four neural

populations (Figure 4C, left, PC1). Thus, the VS and cOFC populations signaled

the expected values in a stable manner.

In contrast, unstable population signals were observed in the DS and mOFC

(Figure 4B, green). The DS population showed a considerable variability in their

eigenvectors (Figure 4C, left, PC1) compared to those in the VS and cOFC neural

populations. The signal carried by the DS neural population was close to 0º, i.e.,

the probability (Figure 4B, DS, vector angle, PC1, mean ± s.e.m., DS, 11.4º ± 1.72)

with a performance closer to that of the cOFC (29%). The mOFC population

exhibited a large variability in the eigenvectors (Figure 4B, mOFC, PC1, vector

angle, mean ± s.e.m., 38.1º ± 5.80, Figure 4C, left, PC1) due to the poorest

performance of PCA (14%), indicating a weak and fluctuating population signal.

Thus, neural populations in the DS and mOFC did not signal the expected values

due to the dynamic changes and weakness of the signals, respectively.

Second, we examined the properties of the eigenvectors for PC2. The

eigenvectors for PC2 revealed another feature of the neural population signals,

that reflected risk-return in the VS and cOFC (Figure S3, Figure 4B, blue, vector

angle, PC2, mean ± s.e.m., VS, 306.7º ± 1.07, 8.3º difference from 315º, cOFC,

322.4º ± 1.94, 7.4º difference from 315º). The deviations from the ideal risk-return

signal were not significantly different between the VS and cOFC populations

(Wilcoxon rank-sum test, n = 52, W = 319, P = 0.737). These signals were equally

stable in the VS and cOFC (Figure 4C, right, PC2). In clear contrast, the DS and

mOFC signals were unstable and fluctuated more (Figure 4C, right, vector angle,

PC2, mean ± s.e.m., DS, 64.8 ± 19.0, mOFC, 320.2 ± 8.77), similar to those

observed for PC1 (Figure 4C, left, PC1). Thus, the VS and cOFC were key brain

regions to signal risk-return as well as the expected values within their neural

population ensembles, suggesting that integrated information of the probability and

magnitude could be signaled in these neural populations.

To further examine the significance of these findings, we used a shuffle control

procedure in two ways (see Methods). First, we randomly shuffled the allocation of

probability and magnitude conditions to neural activity in each trial in each neuron

(shuffled condition 1). If we shuffled the linear projection of neural activity into the

regression subspace by this way, the neural population structure disappeared in

all the four brain regions (Figure 4F). The performance of PCA for PC1 and PC2

were all below 20% (Figure 4E) and were significantly reduced from the observed

data in all the four brain regions, even in the mOFC (Figure 5A, variance explained,

$P < 0.001$ for all populations in PC1 and PC2). The deviance of eigenvectors under

the shuffle control clearly increased in the cOFC and VS, which resulted in no

significant differences among the four neural populations (Figure 4G, Kruskal-

Wallis test, PC1, n = 104, df = 3, H = 4.99, P = 0.18, PC2, n = 104, df = 3, H = 2.78,

P = 0.43). We also tested another shuffle control, in which the trial conditions were

shuffled in each analysis window throughout a trial (shuffled condition 2). Under

this full shuffled control, the PCA performances decreased further, albeit slightly

(Figure 4I and Figure 5B), without significant differences among the four

populations (Figure 4J-K, Deviance, Kruskal-Wallis test, PC1, n = 104, df = 3, H =

1.38, P = 0.71, PC2, n = 104, df = 3, H = 0.53, P = 0.91). Thus, these shuffle

procedures appropriately evaluated the significance of our population findings.

Next, we examined whether the size of eigenvectors is different among the

four neural populations, which represents the extent of neural modulation due to

the probability and magnitude in each neural population as an arbitrary unit. The

size of the eigenvectors was not significantly different (Figure 4D, left, PC1,

Kruskal-Wallis test, n = 104, df = 3, H = 2.62, P = 0.45, right, PC2, n = 104, df = 3,

H = 4.76, P = 0.19), but it strongly depended on a temporal resolution for the

analysis (Figure S4). Size of the eigenvector decreased as the analysis window

size decreased (Figure S4B, E, and F), although all results and conclusions

described above were maintained across the window's sizes (Figure S4A-D). The

decrease in eigenvector size would be because signal-to-noise ratios generally

decrease if the window size decreases. These effects were observed as the

decrease in PCA performance (Figure S4A) and percentages of neural

modulations in the conventional analyses (Figure S5).

Collectively, these observations suggested that the probability and magnitude

of rewards could be detected and integrated within the activity of cOFC and VS

neural populations as the expected value and risk-return signals in a stable state.

305 Temporal structure of the neural population signals

To characterize temporal aspects of the VS and cOFC neural populations that yield

the expected values signals, we compared temporal dynamics of all the four neural

population signals. Specifically, we compared temporal patterns of the vector

change exhibited by each neural population (Figure 6). At the time point after the

cue onset when the monkeys initiated the expected-value computation, all the four

neural populations developed eigenvectors (Figure 6A). Size of the eigenvectors

increased and then decreased within a second, however temporal patterns of this

change in size were different among the four neural populations. The onset

latencies, detected by comparing to the vector size during the baseline period,

seemed to be coincident for the cOFC, VS, and DS population, which was followed

by a late noisy signal in the mOFC (Figure 6B). In contrast, the detected peak of

vector size for each neural population seemed to appear at different times. To

statistically examine these temporal dynamics at the population level, we used a

bootstrap re-sampling technique (see Methods).

The analysis revealed no significant difference in the onset latencies among

cOFC, VS, and DS populations (Figure 6C, bootstrap re-sampling, onset latency,

mean ± s.d., cOFC, 107.1 ± 26.0 ms, VS, 138,7 ± 61.3 ms, DS, 155.0 ± 52.4 ms)

while these signals were followed by a late noisy signal in the mOFC (mOFC, 287

± 98.8 ms). In contrast, if we compared the peak latencies (Figure 6D), the cOFC

exhibited the earliest peak (292 ± 37.5 ms), followed by the DS (371 ± 43.0 ms),

mOFC (444 ± 113.5 ms), and VS (508 ± 76.7), which exhibited the latest peak.

Thus, the expected-value signal sharply developed in the cOFC in contrast to the

gradual development in the VS. Note that mOFC signals were very noisy as

indicated by the large variation in vector size during the baseline period (Figure 6B,

bottom, see horizontal line).

Lastly, we examined temporal changes in vector angle, which indicate how

fast the stable expected value signals evoked in the cOFC and VS (Figure 6E). As

seen in the time series of the vector angle after detected onsets, the signals carried

by the VS and cOFC neural populations at the early time period were almost 45º

(i.e., expected values), indicating that these two neural populations integrate the

probability and magnitude information into the expected values just after the

numerical symbol appeared (see intercepts of the regression lines). Moreover,

these two expected value signals were not the same, but rather idiosyncratic in

each neural population: a gradual and slight shift of the vector angle directed to

90º (i.e., magnitude, cOFC, Figure 6E, regression coefficient, r = 5.31, n = 129, t =

6.04, df = 126, P < 0.0001) or 0º (i.e., probability, Figure 6E, VS, r = -3.91, n = 127,

t = -4.16, df = 124, P < 0.0001) was observed toward the end of cue presentation.

Similar to the VS, the DS population showed the same direction of the angle shift

(Figure 6E, DS, r = -5.38, n = 127, t = -3.31, df = 124, P = 0.001). In contrast, such

significant shift of the vector angle was not observed in the mOFC population (r =

-4.30, n = 120, t = -0.94, df = 117, P = 0.351). These results indicated that the

neural populations in both the VS and cOFC integrate the probability and

magnitude information into the expected values immediately after the cue

presentation, whereas their temporal dynamics were idiosyncratic in each of the

two stable population signals.

Discussion

Extraction of neural population dynamics is a recently developing approach for

understanding computational processes implemented in the domain of cognitive

and motor processing (Chen & Stuphorn, 2015; Churchland et al., 2012; Mante et

al., 2013; Murray et al., 2017; Takei et al., 2017). This approach provides a

mechanistic structure of neural population signals regarding temporal aspects,

such as oscillatory activities during reaching (Churchland et al., 2012), co-

activation patterns of spinal neurons and muscles (Takei et al., 2017), and a

dynamic unfolding of task-related activity for perceptual decisions (Mante et al.,

2013). Here, we found that the VS and cOFC neural populations maintain the

stable expected-value signals at a population level (Figure 4). This is the first

mechanistic demonstration of expected value signals embedded in multiple neural

populations when monkeys compute the expected values from the numerical

symbols cueing the probability and magnitude of rewards. The temporal dynamics

of these two stable neural populations are unique in the aspect of time constants

(Figure 6B-D) and gradual shifts of their structures (Figure 6E). These results

indicate that the cOFC and VS compute the expected values as distinct, partially

overlapping processes. These expected-value signals could be followed by the

transition of neural population signals from evaluation to choices during economic

decision makings (Chen & Stuphorn, 2015; Gardner et al., 2019; Yoo & Hayden,

2020).

Two idiosyncratic expected-value signals in the cOFC and VS

State space analysis can detect both stable (Murray et al., 2017) and flexible

(Mante et al., 2013) neural signals at a population level. In the present study,

expected-value signals observed in the VS and cOFC were similarly stable in terms

of the fluctuation of vector angles but were significantly different in terms of the

temporal aspects (Figure 6). These signal properties indicate that information

processing in these two brain regions were not same. For example, the fast cOFC

signal may reflect the calculation of expected values from the probability and

magnitude symbols like mental arithmetic, while the slow VS signal may reflect the

secondary process to maintain the calculated expected-value information. It is

known that the fronto-striatal projection plays a large role in a variety of cognitive functions (Haber & Knutson, 2010). Since the cOFC project to the VS anatomically, these two processes must act cooperatively through the cortico-basal ganglia loop. Indeed, both population signals were similar in terms of the heterogeneous signals carried by each individual neuron (Figure 2J and K) throughout the task trial (Figure 2N and O). However, these two expected-value signals were unambiguously distinctive in terms of their time course (Figure 6B-D) and gradual shift (Figure 6E). Therefore, the cOFC and VS may compute the expected values within each of these cortical and striatal local circuits in a co-operative manner.

Our results were consistent with human imaging studies, in which activity in the VS and cOFC represents value-related signals (Noonan et al., 2017; J. O'Doherty et al., 2004; Yan et al., 2016), but were not consistent with the evidences that value signals exist in human ventromedial prefrontal cortex (vmPFC) (Levy & Glimcher, 2012; Tom et al., 2007), which includes mOFC. The reasons for why the mOFC showed very weak signals related to all aspects of the expected values in the present study are unclear (Figure 2L and 4B). One possibility of this inconsistency may come from the species differences between human and non-human primates in the orbitofrontal network (Wallis, 2011). The mOFC is a part of the vmPFC, but the comparability between human and macaque monkeys remains elusive. Another possibility is that the vmPFC is not involved in simple information processing, such as association between cues and outcomes, but is involved in more complicated behavioral context for making economic decisions (Yamada et al., 2018) and setting of mood (Ongur & Price, 2000).

Fluctuating signals in the DS and mOFC

Fluctuating signals were observed in the DS and mOFC because of the instability

or weakness of the signals (Figure 4). The mOFC signal would not be completely

meaningless, since the PCA performance in mOFC population was better than that

in shuffle controls (Figure 5). However, the signal carried by mOFC population was

weak (Figure 2L), indicating that the eigenvector fluctuation in mOFC population

reflects weak signal modulations by the probability and magnitude. In contrast,

PCA performance in the fluctuated DS population was equivalent to that in the

cOFC population (Figure 4A), where the stable expected-value signal appeared.

Moreover, considerable modulation of the neural activity in DS was observed in

the conventional analyses (Figure 2I and M). Thus, the fluctuating DS signal must

reflect some functional role employed by the DS neural population in detecting and

integrating the probability and magnitude information. The DS signal fluctuated

with a significant shift from relatively close to the expected values to probability

signal (Figure 6E, top). The consistent direction of the shift between VS and DS

populations implies that striatal neural populations may prefer probabilistic

phenomena (Ma & Jazayeri, 2014; Pouget et al., 2013), whereas cOFC neural

population may prefer magnitude, that is continuous variable.

The expected value signals and economic choices

Economic choices employed in the brain seems to be dynamic sequential

processes, such as the expected value computation, their comparison among

alternatives, and making a choice. Recent findings suggest that these may or may

not be discrete/continuous and overlapping processes until choices are made by

the subject (Chen & Stuphorn, 2015; Yoo & Hayden, 2020). Neural correlates of

the probability, magnitude, and the expected values have been extensively

reported in a diverse set of brain regions (J. P. O'Doherty, 2014) without their

dynamics. They may support a possibility that the expected value signals were

computed in wide brain regions as a network as in the present study, although

those signals were represented for the economic choice (Enomoto et al., 2020) or

an array of alternative non-value related processes (J. P. O'Doherty, 2014), just as

motor response for example. Although signals in the cOFC and VS were fluctuated

(Figure 4B), these populations also signals expected values at the beginning of

cue presentation (Figure 6A and E), suggesting the widespread evolution of the

expected value signals might occur at very early time period through a reward

circuitry.

Significance of the population signals revealed by our state space analysis

State space analysis reveals temporal structures of neural population in multi-

dimensional space for both cognitive (Murray et al., 2017) and motor tasks

(Churchland et al., 2012; Takei et al., 2017). However, the interpretation of the

extracted population structure depends on the way to use this procedure (Elsayed

& Cunningham, 2017). In the present study, we did not seek to determine the

population structure as a trajectory in neural state space, as was performed in

previous studies. Instead, we aimed to detect the main features underscoring the

population structure in the space of the probability and magnitude that compose the expected values. For this purpose, stability of the regression subspace is critical. We elaborately projected the neural firing rates into the regression subspace by preparing a completely orthogonal data matrix in our task design. Thus, our state space analysis is informative on whether and how the expected-value signals are composed of the probability and magnitude moment-by-moment as the series of eigenvectors.

Conclusions

A dynamic integrative process of probability and magnitude underlies computation of expected values in restricted brain regions, i.e., cOFC-VS, which could occur before subjects make economic choices. Since the computation of expected values are followed by the choice process during economic choices, single cue task enables us to extract the integrative process for expected value computation. The existence of the neural population signals for expected-values is consistent with the expected values theory, whereas the co-existence of risk-return signals may reflect a behavioral bias for risk-preferences, a phenomenon observed across species (Stephens & Krebs, 1986; Yamada, Tymula, et al., 2013). The sharp and slow evolutions of expected-value signals in the cOFC and VS, respectively, indicate that each brain region has a unique time constant in the expected-value computation. When the monkeys perceive the probability and magnitude from numerical symbols, learned expected-values may be computed and recalled through the OFC-striatum circuit (Hirokawa et al., 2019). Our results indicate that

476   the expected-value signals as observed in population ensemble activities are

477   compatible with the framework of dynamical systems (Churchland et al., 2012;

478   Mante et al., 2013).

Acknowledgments

The authors appreciate the valuable comments of Tomomichi Oya, Tomohiko Takei, Tsuyoshi Setogawa, Jun Kunimatsu, Masafumi Nejime, Narihisa Matsumoto, Hiroshi Abe, and Takashi Kawai. In addition, the authors thank Takashi Kawai, Ryo Tajiri, Yoshiko Yabana, and Yuki Suwa for their technical assistance. Monkey FU was provided by NBRP "Japanese Monkeys" through the National Bio Resource Project of the MEXT, Japan.

Funding: This research was supported by JSPS KAKENHI Grant Number JP:15H05374, 18K19492, 19H05007, Takeda Science Foundation, Council for Addiction Behavior Studies, Narishige Neuroscience Research Foundation, The Ichiro Kanehara Foundation (H.Y.); JSPS KAKENHI Grant Number JP:26710001, MEXT KAKENHI Grant Number JP:16H06567 (M.M.).

Competing interests: The authors declare no competing of interests.

 Figures and legends

Figure 1. Task and behavior for recording neural activity.

(**A**) Sequence of events during the single cue task. A single visual pie-chart comprising

green and blue pies was presented to the monkeys. (**B**) Choice task. Two visually

displayed pie-charts were presented to the monkeys left and right to the center. After visual

fixation on the re-appeared central target, the central fixation target disappeared, and

monkeys chose either one of the targets by fixating on the target. A block of choice trials

was interleaved between single cue trial blocks. During the choice trials, neural activity

was not recorded. (**C**) Percentages of right target choice during the choice task plotted

against the expected values of left and right target options. (**D**) Pseudo r-squared

estimated in the three behavioral models. M1: number of pies. M2: probability and

magnitude. M3: the expected values. (**E**) An illustration of neural recording areas based

on sagittal magnetic resonance images. Neurons were recorded from medial (mOFC, 14O,

orbital part of area 14) and central parts of the orbitofrontal cortex (cOFC, 13M, medial

part of area 13) at A31-A34 anterior-posterior (A-P) level. Neurons were also recorded

from the dorsal (DS) and ventral striatum (VS) at A21-A27. The white scale bar indicates

5 mm.

Figure 2. Expected-value signals detected by conventional analyses.

(**A**) An example activity histogram of a DS neuron modulated by the expected values during the single cue task. The activity aligned to the cue onset is represented for three different levels of probability (0.1-0.3, 0.4-0.7, 0.8-1.0) and magnitude (0.1-0.3 mL, 0.4-0.7 mL, 0.8-1.0 mL) of rewards. Gray hatched time windows indicate a 1 s time window used to estimate the neural firing rates shown in **B**. (**B**) An activity plot of the DS neuron during the 1 s time window shown in **A** against the probability and magnitude of rewards. (**C-D**) same as **A-B**, but for a VS neuron. (**E-F**) same as **A-B**, but for a cOFC neuron. (**G-H**) same as **A-B**, but for a mOFC neuron. (**I-L**) Plots of the regression coefficients for the probability and magnitude of rewards estimated for all neurons in the DS (**I**), VS (**J**), cOFC (**K**), and mOFC (**L**). Filled colors indicate the neural modulation pattern classified by the

BIC-based model selection. P: Probability type, M: Magnitude type, EV: Expected value

type, and R-R: Risk-Return type. The non-modulated type is indicated by the small open

circle. (**M-P**) Percentages of neural modulation types defined based on the BIC-based

model selection through the cue presentation in the DS (**M**), VS (**N**), cOFC (**O**), and mOFC

(**P**). The results using 0.1 s time window (without overlap) were shown. The classification

of neural modulation types was dependent on the methods, although an overall tendency

for the differences in neural modulations among the neural populations was consistent

among the three analyses.

Figure 3. Schematic depictions for the analysis of neural population dynamics using PCA.

(**A**) Time series of a neural population activity projected into a regression subspace

composed of the probability and magnitude. A series of eigenvectors was obtained by

applying PCA once to each of the four neural populations. PC1 and PC2 indicate the first

and second principal components, respectively. The number of eigenvectors obtained by

the PCA was 2.7 s divided by the analysis window size for the probability and magnitude;

27, 54, and 135 eigenvectors in 0.1, 0.05, or 0.02 s time window, respectively. (**B**)

Examples of eigenvectors at time of *i* th analysis window for the probability and magnitude,

whose direction indicates a signal characteristic at that time represented on the population

ensemble activity. EV: expected values (45º, 225º), M: magnitude (90º, 270º), P:

probability (0º,180º), R-R: risk-return (135º, 315º). (**C**) Characteristics of the eigenvectors

evaluated quantitatively. Angle: vector angle from horizontal axis from 0º to 360º. Size:

length of the eigenvector. Deviance: difference between vectors.

Figure 4. Neural populations provide stable expected-value signals in the VS and cOFC.

(**A**) Cumulative variance explained by PCA in the four neural populations. Dashed line

indicates percentages of variances explained by PC1 and PC2 in each neural population.

(**B**) Overlay plots of the series of eigenvectors for PC1 and PC2 in the four neural

populations. a.u. indicates arbitrary unit. (**C**) Box plots of the deviance from the mean

vector estimated in each neural population for the PC1 (left) and PC2 (right). (**D**) Box plots

of the vector size estimated in each neural population for the PC1 (left) and PC2 (right).

(**E-H**) Same as **A-D**, but for the PCA under the shuffled condition 1. See Methods for

details. (**I-L**) Same as **A-D**, but for the PCA under the shuffled condition 2. In **C-D**, **G-H**,

and **K-L**, asterisks indicate statistical significance between two populations using

Wilcoxon rank-sum test with Bonferroni correction for multiple comparisons (**,*, and §

indicate statistical significance at $P < 0.01$, $P < 0.05$, and $0.05 < P < 0.06$ (close to

significance), respectively). The results are shown by using 0.1 s analysis window.

Figure 5. Probability density of the explained variances by PCA in the shuffled controls.

(**A**) Probability density of variances explained by PCA for PC1 to PC4 under the shuffled condition 1 (see Methods for details). The probability density was estimated with 1,000 repeats of the shuffles in each neural population. (**B**) Probability density of variance explained by the PCA for PC1 to PC4 under the shuffled condition 2 (see Methods for details). The probability density was estimated with 1,000 repeats of the shuffles in each neural population. In **A** and **B**, dashed lines indicate the variances explained by PCA in each of the four neural populations without the shuffle.

Figure 6. Gradual and sharp evolutions of neural population signals in the VS and cOFC.

(**A**) Plots of the time series of eigenvectors for PC1 in 0.02 s analysis windows shown in

a sequential order during 1 s after the cue onset. Horizontal and vertical scale bars indicate

the eigenvectors for the probability and magnitude in arbitrary units, respectively. (**B**) Plots

of the time series of vector size during 1 s after the cue onset. Horizontal dashed lines

indicate three standard deviations of the mean vector size during the baseline period, a

0.3 s time period before the cue onset. Solid colored lines indicate interpolated lines using

a cubic spline function to provide a resolution of 0.005 s. Vertical dashed lines indicate the

onset (left) and peak (right) latencies for the changes in vector sizes. (**C**) Probability

densities of the onset latencies of the four neural population signals. The probability

densities were estimated using bootstrap re-samplings. Vertical dashed lines indicate the

means. Horizontal solid lines indicate the bootstrap standard errors. (**D**) same as **C**, but

for the peak latencies of the four neural population signals. (**E**) Plots of the time series of

vector angle from the detected onset to the onset of outcome feedback. Solid black lines

indicate regression slopes. In **C** and **D**, asterisks indicate statistical significance estimated

using bootstrap re-samplings (*** and * indicate statistical significance at $P < 0.001$ and P

< 0.05, respectively). In **E**, triple asterisks indicate a statistical significance of the

regression slope at $P < 0.001$. Data for the PC2 is not shown.

Materials and Methods

Subjects and experimental procedures

Two rhesus monkeys were used (*Macaca mulatta,* SUN, 7.1 kg, male; *Macaca fuscata*, FU, 6.7 kg, female). All experimental procedures were approved by the Animal Care and Use Committee of University of Tsukuba (protocol#. H30.336) and performed in compliance with the US Public Health Service's Guide for the Care and Use of Laboratory Animals. Each animal was implanted with a head-restraint prosthesis. Eye movements were measured using a video camera system at 120 Hz. Visual stimuli were generated by a liquid-crystal display at 60 Hz that was placed 38 cm from the monkey's face when seated. The subjects performed the cued lottery task five days a week. The subjects practiced the cued lottery task for 10 months, after which they became proficient at making choices of lottery options.

Behavioral task

Cued lottery tasks. Animals performed one of two visually cued lottery tasks: *single cue task* or *choice task*. Activity of neurons were recorded only during the single cue task.

Single cue task: At the start of each trial, the monkeys had 2 s to align their gaze to within 3° of a 1° diameter grey central fixation target. After fixating for 1 s, an 8° pie-chart providing information about the probability and magnitude of rewards was presented for 2.5 s at the same location as the central fixation target. The pie-chart was then removed and 0.2 s later a 1 kHz and 0.1 kHz tone (0.15 s duration)

indicated reward and no-reward outcomes, respectively. The high tone preceded a reward by 0.2 s. The low tone indicated that no reward would be delivered. The animals received a fluid reward, for which magnitude and probability were indicated by the green and blue pie-charts, respectively; otherwise, no-reward was delivered. An inter-trial interval of 4 to 6 s followed.

Choice task: At the start of each trial, the monkeys had 2 s to align their gaze to within $3°$ of a $1°$ diameter grey central fixation target. After fixating for 1 s, two peripheral $8°$ pie-charts providing information about the probability and magnitude of rewards for each of the two target options were presented for 2.5 s, $8°$ to the left and right of the central fixation location. Grey $1°$ choice targets appeared at these same locations. After a 0.5 s delay, the fixation target disappeared, cueing saccade initiation. The animals were free to choose for 2 s by shifting their gaze to either target, within $3°$ of the choice target. A 1 kHz and 0.1 kHz tone of 0.15 s duration indicated reward and no-reward outcomes, respectively. The animals received the fluid reward indicated by the green pie-chart of the chosen target, with the probability indicated by the blue pie-chart of the chosen target; otherwise, no-reward was delivered. An inter-trial interval of 4 to 6 s followed.

Pay-off and block structure. Green and blue pie-charts indicated reward magnitudes from 0.1 to 1.0 mL in 0.1 mL increments and reward probabilities from 0.1 to 1.0 in 0.1 increments, respectively (Table S1). A total of 100 pie-charts was used. In the single cue task, each of the 100 pie-charts was presented once in random order. In the choice task, two out of the 100 pie-charts were randomly

allocated to the two options. During one session of electrophysiological recordings,

20 to 50 trial blocks of the choice task were interleaved with 100 to 120 trial blocks

of the single cue task.

Calibration of reward supply system. The precise amount of liquid reward was

controlled and delivered to the monkeys with a solenoid valve. An 18-gauge tube

(0.9 mm inner diameter) was attached to the tip of the delivery tube to reduce the

trial-by-trial variability of reward supply. The amount of reward in each payoff

condition was calibrated by measuring the weight of water with 0.002 g precision

(hence, 2μL) on a single trial basis. This calibration method was the same as that

in a previous study (Yamada et al., 2018).

Electrophysiological recordings

We used conventional techniques for recording single neuron activity from the DS,

VS, cOFC, and mOFC. The monkeys were implanted with recording chambers (28

× 32 mm) targeting the OFC and striatum, centered 28 mm anterior to the

stereotaxic coordinates. The locations of the chambers were verified using

anatomical magnetic resonance imaging. In each recording session, a stainless-

steel guide tube was placed within a 1-mm spacing grid, and a tungsten

microelectrode (1-3 MΩ, FHC) was passed through the guide tube. For recording

of neurons in the mOFC and cOFC, the electrode was lowered until it approximated

the bottom of the brain after passing through the cingulate cortex or dorsolateral

prefrontal cortex or between them, respectively. For recording of neurons in the

DS, the electrode was lowered until low spontaneous activity was observed after

passing through the cortex and white matter. For recording of neurons in the VS,

the electrode was lowered further until it passed through the internal capsule. At

the end of the VS recording session, the electrode was occasionally lowered close

to the bottom of the brain to confirm the recording depth relative to the bottom.

Electrophysiological signals were amplified, band pass filtered, and monitored.

Single neuron activity was isolated on the basis of spike waveforms. We recorded

from the four brain regions of single hemisphere of the two monkeys: 194 DS

neurons (98 and 96 from monkeys SUN and FU, respectively), 144 VS neurons

(89, SUN and 55, FU), 190 cOFC neurons (98, SUN and 92, FU), and 158 mOFC

neurons (64, SUN and 94, FU). The activity of all single neurons was sampled

when the activity of an isolated neuron demonstrated a good signal-to-noise ratio

(>2.5). Blinding was not performed. The sample sizes required to detect effect

sizes (number of recorded neurons, number of recorded trials in a single neuron,

and number of monkeys) were estimated with reference to previous studies (Chen

& Stuphorn, 2015; Yamada, Inokawa, et al., 2013; Yamada et al., 2018). Neural

activity was recorded during 100-120 trials of the single cue task. Note that

presumed projection neurons (PANs, phasically active neurons) (Yamada et al.,

2016) were recorded from the DS and VS.

Statistical analysis

For statistical analysis, we used the statistical software package R (http://www.r-

project.org/). All statistical tests for behavioral and neural analyses were two-tailed.

Effects of units on statistical analysis. In the present study, we used two variables

for analyses; probability and magnitude. We defined the probability of reward from

0.1 to 1.0, and the magnitude of reward from 0.1 to 1.0 mL. Under this definition of

units, the effects of probability and magnitude on the data were equivalent. Thus,

the data were not standardized in the analyses.

Behavioral analysis

We examined whether the monkey's choice behavior depended on the expected

values of the two options located left and right side to the center. We pooled choice

data across all recording sessions, yielding 44,883 and 19,292 choice trials for

monkeys SUN and FU, respectively. The percentage of right target choices was

estimated for all combinations of the expected values of left and right target options.

Model Fitting. The percentage of choosing the right-side option was analyzed using

a general linear model with binominal distribution,

$$Pchooses_R = 1 / (1 + e^{-z}) \tag{1}$$

where the relation between $Pchooses_R$ and Z was given by the logistic function in

each of the following three models: number of pies displayed (M1), probability and

magnitude (M2), and the expected values (M3).

The first model, M1, assumed that the monkeys chose a target by comparing

the number of pies for two targets,

$$Z = b_0 + b_1 Npie_L + b_2 Npie_R \tag{2}$$

where b_0 is the intercept, and $Npie_L$ and $Npie_R$ are the number of pies contained in left and right pie-chart stimuli, respectively. Values of b_0 to b_2 were free parameters and estimated by maximizing log likelihood.

The second model, M2, assumed that the monkeys chose a target by comparing the probability and magnitude of two targets,

$$Z = b_0 + b_1 P_L + b_2 P_R + b_3 M_L + b_4 M_R \qquad (3)$$

where b_0 is the intercept; P_L and P_R are the probability of rewards for left and right pie-chart stimuli, respectively; and M_L and M_R are the magnitude of rewards for left and right pie-chart stimuli, respectively. Values of b_0 to b_4 were free parameters and estimated by maximizing log likelihood.

The third model, M3, assumed that the monkeys chose a target by comparing the expected values of rewards for two targets,

$$Z = b_0 + b_1 EV_L + b_2 EV_R \qquad (4)$$

where b_0 is the intercept, EV_L and EV_R are the expected values of rewards as probability times magnitude for left and right pie-chart stimuli, respectively. Values of b_0 to b_2 were free parameters and estimated by maximizing log likelihood.

Model comparisons. To identify the best structural model to describe the monkeys' behavior, we compared the three models described above. In each model, we estimated a combination of the best-fit parameters to explain monkeys' choice behavior. We compared their goodness-of-fits based on Akaike's Information Criterions (AIC) and Bayesian Information Criterion (BIC) (Burnham & Anderson, 2004),

AIC (Model) = −2L + 2k (5)

BIC (Model) = −2L + k log n (6)

where L is the maximum log-likelihood of the model, k is the number of free

parameters, and n is the sample size. After estimating the best-fit parameters in

each model, we selected one model that exhibited the smallest AIC and BIC. To

evaluate the model fits, we estimated a McFadden's pseudo r-squared statistic

using the following equation:

Pseudo r-squared = (L_0 - L_{Model}) / L_0 (7)

where L_{Model} is the maximum log likelihood for the model given the data, and L_0 is

the log likelihood under the assumption that all free parameters were zero in the

model.

Neural analysis

Basic firing properties. Peri-stimulus time histograms (PSTH) were drawn for each

single neuron activity aligned at the visual cue onset. To display the histogram of

neurons, the peak activity (maximum firing rate in each histogram) was detected

for each neuron. The average activity curves were smoothed using a 50 ms

Gaussian kernel (σ = 50 ms) and normalized by the peak firing rates. The

percentage of neurons that showed an activity peak during cue presentation was

compared among the four brain regions using a chi-squared test at $P < 0.05$. Basic

firing properties, such as peak firing rates, latency of the peak, and duration of the

peak activity (half peak width) were compared among the four brain regions using

parametric and non-parametric tests, respectively, with a statistical significance

level of $P < 0.05$. Baseline firing rates during 1 s before the appearance of central

fixation targets were also compared with a statistical significance level of $P < 0.05$.

Estimation of neural firing rates through task trials. We analyzed neural activity

during a 2.7 s time period from the onset of pie-chart stimuli to the onset of outcome

feedbacks during the single cue task. To obtain a time series of neural firing rates

through a trial, we estimated the firing rates of each neuron for every 0.1, 0.05, or

0.02 s time window (without overlap) during the 2.7 s period. No gaussian kernel

was used.

Estimation of neural firing rates in a fixed time window. We analyzed neural activity

during a 1 s time window after the onset of pie-chart stimuli during the single cue

task. The 1 s activity was used for the conventional analyses below. No gaussian

kernel was used.

Conventional analyses to detect neural modulations in each individual

neuron

Linear regression and model selection. For conventional and standard analyses of

neural modulation by the values of probability and magnitude indicated by pie-chart

stimuli, we used *linear regression* and *model selection analyses.* As mentioned

above, we estimated the firing rate of each neuron during the 1 s time period after

the onset of pie-chart stimulus during the single cue task. No gaussian kernel was

used.

Linear regression. Neural discharge rates (F) were fitted by a linear combination of the following variables:

$$F = b_0 + b_p \text{Probability} + b_m \text{Magnitude} \qquad (8)$$

where Probability and Magnitude are the probability and magnitude of rewards indicated by the pie-chart, respectively. b_0 is the intercept. If b_p and b_m were not 0 at $P < 0.05$, the discharge rates were regarded as being significantly modulated by that variable.

On the basis of the linear regression, activity modulation patterns were categorized into several types: "Probability" type with a significant b_p and without a significant b_m; "Magnitude" type without a significant b_p and with a significant b_m; "Expected value" type with significant b_p and b_m with both the same signs (i.e., positive b_p and positive b_m or negative b_p and negative b_m); "Risk-Return" type with significant b_p and b_m with both having opposite signs (i.e., negative b_p and positive b_m or positive b_p and negative b_m) and "non-modulated" type without significant b_p and b_m. Note that risk-return types reflect high-risk high-return (prefer low probability and large magnitude) or low-risk low-return (prefer high probability and low magnitude).

Model selection. Neural discharge rates (F) were fitted by the following five models:

$$\text{M1:} \quad F = b_0 \qquad (9)$$

$$\text{M2:} \quad F = b_0 + b_p \text{Probability} \qquad (10)$$

$$\text{M3:} \quad F = b_0 + b_m \text{Magnitude} \qquad (11)$$

M4: $F = b_0 + b_p$ Probability $+ b_m$ Magnitude   (12)

M5: $F = b_0 + b_{ev}$ Expected value   (13)

where Expected value is the expected value indicated by the visual pie-chart as

the probability multiplied by magnitude (Table S1). b_0 is the intercept. Probability

and Magnitude are the probability and magnitude of reward indicated by the pie-

chart, respectively. Among the five models, we selected one model that exhibited

the smallest AIC or BIC.

If the selected model was M1, neurons were defined as the "non-modulated"

type. If the selected model was M2, neurons were defined as the "Probability" type.

If the selected model was M3, neurons were defined as the "Magnitude" type. If

the selected model was M4 with the same signs of b_p and b_m, neurons were defined

as the "Expected value" type. If the selected model was M4 with opposite signs of

b_p and b_m, neurons were defined as the "Risk-Return" type. If the selected model

was M5, neurons were defined as the "Expected value" type.

Application of conventional analyses to neural activity through task trials. We

applied the three conventional analyses above (linear regression, AIC-based

model selection, and BIC-based model selection) for the activity of neurons

estimated at every time window in the four brain regions. As mentioned above, we

estimated the firing rate of each neuron for every 0.1, 0.05, or 0.02 s time window

(without overlap) during the 2.7 s period. No gaussian kernel was used. The activity

modulation type was defined in each time window during the 2.7 s period. The

analyses described percentages of the neural modulation types throughout cue

presentation (Figure S5).

Population dynamics using principal component analysis

Estimation of neuron firing rates through task trials. As mentioned above, we

estimated the firing rate in each neuron every 0.1, 0.05, or 0.02 s time window

(without overlap) during the 2.7 s period. No gaussian kernel was used.

Regression subspace. We used a linear regression to determine how the

probability and magnitudes of rewards affected the activities of each neuron in the

four neural populations. Each neural population was composed of all the recorded

neurons in each brain region. We first prepared the probability and magnitude as

0.1 to 1.0 and 0.1 to 1.0 mL, respectively. We then described average firing rates

of the neuron *i* at time *t* as a linear combination of the probability and magnitude in

each neural population:

$$F_{(i,t,k)} = b_{0(i,t)} + b_{1(i,t)}\text{Probability}_{(k)} + b_{2(i,t)}\text{Magnitude}_{(k)} \qquad (14)$$

where $F_{(i,t,k)}$ is the average firing rates of neuron *i* at time t on trial *k,* Probability$_{(k)}$

is the probability of rewards cued to the monkeys on trial *k;* Magnitude$_{(k)}$ *is* the

magnitude of rewards cued to the monkeys on trial *k*. The regression coefficients

$b_{0(i,t)}$ to $b_{2(i,t)}$ describe the degree to which the firing rates of neuron *i* depend on the

mean firing rates (hence, firing rates independent of task variables), the probability

of rewards, and the magnitude of rewards, respectively, at a given time *t* during

the trials.

We used the regression coefficients described in Eq. 14, above, to identify

how the dimensions of neural population signals were composed from the

probability and magnitudes as aggregated properties of individual neural activity.

This step corresponds to the fundamental conceptual step of viewing the

regression coefficients as a temporal structure of neural modulations by the

probability and magnitude at the population level. Our procedures are analogous

to the state-space analysis performed by Mante et al (Mante et al., 2013), in which

the regression coefficients were used to provide an axis (or dimension) of the

interested variables in multi-dimensional state space obtained by principal

component analysis (PCA). In the present study, our orthogonalized task design

allowed us to project the neural firing rates into the regression subspace reliably.

It must be noted that our analyses were not aimed at describing population

dynamics of the neural signals as a trajectory in the multi-dimensional task space,

which is the standard goal of state space analysis.

Principal component analysis. We used a PCA to identify dimensions of the neural

population signal in the orthogonal spaces composed of the probability and

magnitude of rewards in each of the four neural populations. In each neural

population, we first prepared a two dimensional data matrix X of size $N_{(neuron)} \times N$

$_{(C \times T)}$; the regression coefficient vectors, $b_{1(i,t)}$ and $b_{2(i,t)}$ in Eq.14, whose rows

correspond to the total number of neurons in each neural population and whose

columns correspond to C, the total number of conditions (i.e., two: probability and

magnitude); and T, the total number of analysis windows (i.e., 2.7 s divided by the

window size). The PCs of this data matrix are vectors $v_{(a)}$ of length $N_{(neuron)}$, total

number of recorded neurons; if $N_{(C \times T)}$ is larger than $N_{(neuron)}$, otherwise the length

is $N_{(C \times T)}$. The PCs were indexed from the principal component explaining the most

variance to the least variance. The eigenvectors were obtained by prcomp ()

function in the R software. It must be noted that we did not perform de-noising in

the PCA (Mante et al., 2013), since we did not aim to project firing rates into state

space. Instead, we intended to use the principal components to identify main

features of the neural modulation signals at the population level through task trials.

Eigenvectors. When we applied the PCA to the data matrix *X*, we could deconstruct

the set into eigenvectors and eigenvalues. The eigenvectors and eigenvalues exist

as pairs with every eigenvector having a corresponding eigenvalue. In our analysis,

the eigenvectors at time t represent a vector in the space of probability and

magnitude. The eigenvalues at time t for the probability and magnitude are scalars,

indicating how much variance is in the data in that vector. The first principal

component is thus the eigenvectors with the highest eigenvalues. We analyzed

eigenvectors for the first (PC1) and second principal components (PC2) in the

following analyses. It must be noted here that we applied the PCA once in each

neural population and thus the total variances contained in the data were different

among the four populations.

Analysis of eigenvectors. We evaluated characteristics of the eigenvectors for PC1

and PC2 in each of the four neural populations in terms of the vector angle, size,

and deviance in the space of probability and magnitude (Figure 3). The angle is

the vector angle from the horizontal axis from 0º to 360º. The size is the length of

the eigenvector. The deviance is the difference between vectors, and we estimated

the deviance from the mean vector in each neural population. These three

characteristics of the eigenvectors were compared among the four neural

populations at $P < 0.05$ using the Kruskal-Wallis test and Wilcoxon rank-sum test

with Bonferroni correction for multiple comparisons. The vector during the first 0.1

s was extracted from these analyses.

Shuffle control for PCA. To examine the significance of the population structure

described by the PCA, we performed two shuffle controls. When we projected

neural activity into the regression subspace, we randomized the data by shuffling

in two ways. In the shuffled condition 1, $b_{1(i,t)}$ and $b_{2(i,t)}$ in Eq.14 were estimated with

the randomly shuffled allocation of trial number k to the Probability$_{(k)}$ and

Magnitude$_{(k)}$ only once for all time t in each neuron. This shuffle provided a data

matrix X of size $N_{(neuron)} \times N_{(C \times T)}$, eliminating the modulation of probability and

magnitude observed in the condition C, but retaining the temporal structure of

these modulations across time. In the shuffled condition 2, $b_{1(i,t)}$ and $b_{2(i,t)}$ in Eq.14

were estimated with the randomly shuffled allocation of trial number k to the

Probability$_{(k)}$ and Magnitude$_{(k)}$ at each time t in each neuron. This shuffle provided

a data matrix X of size $N_{(neuron)} \times N_{(C \times T)}$, eliminating the structure across conditions

and times. In these two shuffle controls, matrix X was estimated 1,000 times. The

PCA performance was evaluated by constructing distributions of the explained

variances for PC1 to PC4 (Figure 5). The statistical significance of the explained

variances by PC1 and PC2 was estimated based on the bootstrap standard errors

(i.e., standard deviation of the reconstructed distribution).

Bootstrap re-samplings for onset and peak latencies of the neural population

signals. To detect onset and peak latencies of the population signal, we analyzed

dynamic changes in population structure with the size of the eigenvectors in each

of the four populations. We used a time-series of eigenvectors in 0.02 s analysis

windows and estimated sizes of the time-series of vectors for PC1. To obtain

smooth changes in vector size, a cubic spline function was applied with a resolution

of 0.005 s (Figure 6B). Vector sizes during a baseline period were obtained by

applying PCA to the matrix data X with time t from 0.3 s before cue onset to the

onset of feedback (i.e., 3.0 s time period). A standard deviation of the vector sizes

during the baseline period was obtained in each of the four neural populations. An

onset latency of the population signal was defined as a time when the spline curve

was above a value of more than three s.d. during the baseline period. Peak latency

of the population signal was defined as the time from cue onset to the time when

the maximum size of the vectors was obtained.

We estimated mean latencies of the onset and peak using a parametric

bootstrap re-sampling method. In each neural population, the neurons were

randomly re-sampled with a duplicate, and a data matrix X of size $N_{(neuron)} \times N_{(C \times T)}$

was obtained. The PCA was applied to the data matrix X. The time-series of the

eigenvectors was obtained, and their sizes were estimated. The onset and peak

latencies were estimated as above. This resampling was conducted 1,000 times,

and distributions of the onset and peak latencies were obtained (Figure 6C and D).

The statistical significance of the onset and peak latencies was estimated based

on the bootstrap standard errors (i.e., standard deviation of the reconstructed

distribution) (Efron. & Tibshirani., 1993).

References
942

Barraclough, D. J., Conroy, M. L., & Lee, D. (2004). Prefrontal cortex and decision
making in a mixed-strategy game. *Nat Neurosci, 7*(4), 404-410.

Burnham, K., & Anderson, D. (2004). Multimodel inference: understanding AIC and BIC
in model selection. *Sociol. Method Res., 33*, 261–304.

Chen, X., & Stuphorn, V. (2015). Sequential selection of economic good and action in
medial frontal cortex of macaques during value-based decisions. *Elife, 4*.
doi:10.7554/eLife.09418

Churchland, M. M., Cunningham, J. P., Kaufman, M. T., Foster, J. D., Nuyujukian, P.,
Ryu, S. I., & Shenoy, K. V. (2012). Neural population dynamics during reaching.
Nature, 487(7405), 51-56. doi:10.1038/nature11129

Efron., B., & Tibshirani., R. (1993). *An Introduction to the Bootstrap*: Chapman &
Hall/CRC.

Elsayed, G. F., & Cunningham, J. P. (2017). Structure in neural population recordings: an
expected byproduct of simpler phenomena? *Nat Neurosci, 20*(9), 1310-1318.
doi:10.1038/nn.4617

Enomoto, K., Matsumoto, N., Inokawa, H., Kimura, M., & Yamada, H. (2020).
Topographic distinction in long-term value signals between presumed dopamine
neurons and presumed striatal projection neurons in behaving monkeys. *Sci Rep,*
10(1), 8912. doi:10.1038/s41598-020-65914-0

Eshel, N., Tian, J., Bukwich, M., & Uchida, N. (2016). Dopamine neurons share common
response function for reward prediction error. *Nat Neurosci, 19*(3), 479-486.
doi:10.1038/nn.4239

Fouragnan, E. F., Chau, B. K. H., Folloni, D., Kolling, N., Verhagen, L., Klein-Flugge,
M., . . . Rushworth, M. F. S. (2019). The macaque anterior cingulate cortex
translates counterfactual choice value into actual behavioral change. *Nat*
Neurosci, 22(5), 797-808. doi:10.1038/s41593-019-0375-6

Gardner, M. P. H., Conroy, J. C., Sanchez, D. C., Zhou, J., & Schoenbaum, G. (2019).
Real-Time Value Integration during Economic Choice Is Regulated by
Orbitofrontal Cortex. *Curr Biol, 29*(24), 4315-4322 e4314.
doi:10.1016/j.cub.2019.10.058

Glimcher, P. W., Camerer, C. F., Fehr, E., & Poldrack, R. A. (2008). *Neuroeconomics:*
Decision Making and the Brain. New York: Elsevier.

Goense, J. B., & Logothetis, N. K. (2008). Neurophysiology of the BOLD fMRI signal in
awake monkeys. *Curr Biol, 18*(9), 631-640. doi:10.1016/j.cub.2008.03.054

Haber, S. N., & Knutson, B. (2010). The reward circuit: linking primate anatomy and
human imaging. *Neuropsychopharmacology, 35*(1), 4-26.
doi:10.1038/npp.2009.129

Hirokawa, J., Vaughan, A., Masset, P., Ott, T., & Kepecs, A. (2019). Frontal cortex
neuron types categorically encode single decision variables. *Nature, 576*(7787),
446-451. doi:10.1038/s41586-019-1816-9

Houthakker, H. S. (1950). Revealed Preference and the Utility Function. *Economica,*
17(66), 159-174.

Howard, J. D., Gottfried, J. A., Tobler, P. N., & Kahnt, T. (2015). Identity-specific
coding of future rewards in the human orbitofrontal cortex. *Proc Natl Acad Sci U*
S A, 112(16), 5195-5200. doi:1503550112 [pii]
10.1073/pnas.1503550112

Howard, J. D., & Kahnt, T. (2017). Identity-Specific Reward Representations in
Orbitofrontal Cortex Are Modulated by Selective Devaluation. *J Neurosci,*
37(10), 2627-2638. doi:JNEUROSCI.3473-16.2017 [pii]

10.1523/JNEUROSCI.3473-16.2017

Hsu, M., Krajbich, I., Zhao, C., & Camerer, C. F. (2009). Neural response to reward

anticipation under risk is nonlinear in probabilities. *J Neurosci, 29*(7), 2231-2237.

doi:10.1523/JNEUROSCI.5296-08.2009

Kahneman, D., & Tversky, A. (1979). Prospect theory: An analysis of decisions under

risk. *Econometrica, 47*, 313–327.

Levy, D. J., & Glimcher, P. W. (2012). Comparing apples and oranges: using reward-

specific and reward-general subjective value representation in the brain. *J*

Neurosci, 31(41), 14693-14707. doi:10.1523/JNEUROSCI.2218-11.2011

Lopatina, N., McDannald, M. A., Styer, C. V., Peterson, J. F., Sadacca, B. F., Cheer, J.

F., & Schoenbaum, G. (2016). Medial Orbitofrontal Neurons Preferentially Signal

Cues Predicting Changes in Reward during Unblocking. *J Neurosci, 36*(32),

8416-8424. doi:10.1523/JNEUROSCI.1101-16.2016

Ma, W. J., & Jazayeri, M. (2014). Neural coding of uncertainty and probability. *Annu Rev*

Neurosci, 37, 205-220. doi:10.1146/annurev-neuro-071013-014017

Mante, V., Sussillo, D., Shenoy, K. V., & Newsome, W. T. (2013). Context-dependent

computation by recurrent dynamics in prefrontal cortex. *Nature, 503*(7474), 78-

84. doi:10.1038/nature12742

Milham, M. P., Ai, L., Koo, B., Xu, T., Amiez, C., Balezeau, F., . . . Schroeder, C. E.

(2018). An Open Resource for Non-human Primate Imaging. *Neuron, 100*(1), 61-

74 e62. doi:10.1016/j.neuron.2018.08.039

Murray, J. D., Bernacchia, A., Roy, N. A., Constantinidis, C., Romo, R., & Wang, X. J.

(2017). Stable population coding for working memory coexists with

heterogeneous neural dynamics in prefrontal cortex. *Proc Natl Acad Sci U S A,*

114(2), 394-399. doi:10.1073/pnas.1619449114

Noonan, M. P., Chau, B. K. H., Rushworth, M. F. S., & Fellows, L. K. (2017).

Contrasting Effects of Medial and Lateral Orbitofrontal Cortex Lesions on Credit

Assignment and Decision-Making in Humans. *J Neurosci, 37*(29), 7023-7035.

doi:10.1523/JNEUROSCI.0692-17.2017

O'Doherty, J., Dayan, P., Schultz, J., Deichmann, R., Friston, K., & Dolan, R. J. (2004).

Dissociable roles of ventral and dorsal striatum in instrumental conditioning.

Science, 304(5669), 452-454. doi:10.1126/science.1094285

O'Doherty, J. P. (2014). The problem with value. *Neurosci Biobehav Rev, 43*, 259-268.

doi:S0149-7634(14)00084-0 [pii]

10.1016/j.neubiorev.2014.03.027

Ongur, D., & Price, J. L. (2000). The organization of networks within the orbital and

medial prefrontal cortex of rats, monkeys and humans. *Cereb Cortex, 10*(3), 206-

219.

Papageorgiou, G. K., Sallet, J., Wittmann, M. K., Chau, B. K. H., Schuffelgen, U.,

Buckley, M. J., & Rushworth, M. F. S. (2017). Inverted activity patterns in

ventromedial prefrontal cortex during value-guided decision-making in a less-is-

more task. *Nat Commun, 8*(1), 1886. doi:10.1038/s41467-017-01833-5

Platt, M. L., & Glimcher, P. W. (1999). Neural correlates of decision variables in parietal

cortex. *Nature, 400*(6741), 233-238.

Pouget, A., Beck, J. M., Ma, W. J., & Latham, P. E. (2013). Probabilistic brains: knowns

and unknowns. *Nat Neurosci, 16*(9), 1170-1178. doi:10.1038/nn.3495

Roesch, M. R., Singh, T., Brown, P. L., Mullins, S. E., & Schoenbaum, G. (2009).

Ventral striatal neurons encode the value of the chosen action in rats deciding

between differently delayed or sized rewards. *J Neurosci, 29*(42), 13365-13376.

doi:29/42/13365 [pii]

10.1523/JNEUROSCI.2572-09.2009

Rudebeck, P. H., & Murray, E. A. (2014). The orbitofrontal oracle: cortical mechanisms

for the prediction and evaluation of specific behavioral outcomes. *Neuron, 84*(6),

1143-1156. doi:10.1016/j.neuron.2014.10.049

Samuelson, P. A. (1950). The Problem of Integrability in Utility Theory. *Economica,*

17(68), 355–385.

Savage, L. J. (1954). *The Foundations of Statistics*. New. York: John Wiley and Sons.

Stephens, D., & Krebs, J. (1986). *Foraging Theory*. New Jersey: Princeton Univ. Press.

Sutton, R. S., & Barto, A. G. (1998). *Reinforcement Learning*. Cambridge: The MIT
press.
Takei, T., Confais, J., Tomatsu, S., Oya, T., & Seki, K. (2017). Neural basis for hand
muscle synergies in the primate spinal cord. *Proc Natl Acad Sci U S A, 114*(32),
8643-8648. doi:10.1073/pnas.1704328114
Tobler, P. N., Fiorillo, C. D., & Schultz, W. (2005). Adaptive coding of reward value by
dopamine neurons. *Science, 307*(5715), 1642-1645.
Tom, S. M., Fox, C. R., Trepel, C., & Poldrack, R. A. (2007). The neural basis of loss
aversion in decision-making under risk. *Science, 315*(5811), 515-518.
doi:315/5811/515 [pii]
10.1126/science.1134239
Von Neumann, J., & Morgenstern, O. (1944). *Theory of Games and Economic Behavior*.
New Jersey: Princeton Univ. Press.
Wallis, J. D. (2011). Cross-species studies of orbitofrontal cortex and value-based
decision-making. *Nat Neurosci, 15*(1), 13-19. doi:10.1038/nn.2956
Xie, J., & Padoa-Schioppa, C. (2016). Neuronal remapping and circuit persistence in
economic decisions. *Nat Neurosci, 19*(6), 855-861. doi:10.1038/nn.4300
Yamada, H., Inokawa, H., Hori, Y., Pan, X., Matsuzaki, R., Nakamura, K., . . .
Minamimoto, T. (2016). Characteristics of fast-spiking neurons in the striatum of
behaving monkeys. *Neurosci Res, 105*, 2-18. doi:10.1016/j.neures.2015.10.003
Yamada, H., Inokawa, H., Matsumoto, N., Ueda, Y., Enomoto, K., & Kimura, M. (2013).
Coding of the long-term value of multiple future rewards in the primate striatum.
J Neurophysiol, 109(4), 1140-1151. doi:10.1152/jn.00289.2012
Yamada, H., Louie, K., Tymula, A., & Glimcher, P. W. (2018). Free choice shapes
normalized value signals in medial orbitofrontal cortex. *Nat Commun, 9*(1), 162.
doi:10.1038/s41467-017-02614-w
Yamada, H., Tymula, A., Louie, K., & Glimcher, P. W. (2013). Thirst-dependent risk
preferences in monkeys identify a primitive form of wealth. *Proc Natl Acad Sci U
S A, 110*(39), 15788-15793. doi:1308718110 [pii]
10.1073/pnas.1308718110

Yan, C., Su, L., Wang, Y., Xu, T., Yin, D. Z., Fan, M. X., . . . Chan, R. C. (2016).

Multivariate Neural Representations of Value during Reward Anticipation and

Consummation in the Human Orbitofrontal Cortex. *Sci Rep, 6*, 29079.

doi:10.1038/srep29079

Yoo, S. B. M., & Hayden, B. Y. (2020). The Transition from Evaluation to Selection

Involves Neural Subspace Reorganization in Core Reward Regions. *Neuron,*

105(4), 712-724 e714. doi:10.1016/j.neuron.2019.11.013

Zhou, J., Gardner, M. P. H., Stalnaker, T. A., Ramus, S. J., Wikenheiser, A. M., Niv, Y.,

& Schoenbaum, G. (2019). Rat Orbitofrontal Ensemble Activity Contains

Multiplexed but Dissociable Representations of Value and Task Structure in an

Odor Sequence Task. *Curr Biol, 29*(6), 897-907 e893.

doi:10.1016/j.cub.2019.01.048

1092

Supplemental Information

Neural population dynamics underlying expected value

computation

Hiroshi Yamada*, Yuri Imaizumi, Masayuki Matsumoto

Supplemental Figures

Figure S1. Model fits to the choice behavior of two monkeys.

(**A**) Plots of the AIC values in monkey SUN and FU estimated in the three behavioral

models, M1: number of pie. M2: probability and magnitude. M3: the expected values. (**B**)

Same as **A**, but for the BIC values.

Figure S2. Basic firing properties of neurons in the DS, VS, cOFC, and mOFC.

(**A**) Color map histograms of neuronal activities recorded from the four brain regions. Each horizontal line indicates the neural activity aligned to the cue onset, that was averaged for all lottery conditions. Firing rates of a neuron were normalized to the peak activity. (**B**) The percentages of neurons that showed an activity peak during the cue presentation. (**C**) Latency plots of the peak activity after the cue presentation. (**D**) Firing rates of the peak activity observed during the cue presentation. (**E**) Plots of the half-peak width, which indicates the phasic nature of activity changes. (**F**) Plots of the baseline firing rates during 1 s time period before the onset of the central fixation target. In **C-F**, asterisks indicate statistical significance among two neural populations using Wilcoxon rank-sum test with Bonferroni correction for multiple comparisons (**,*, and § indicate statistical significance at $P < 0.01$, $P < 0.05$, and $0.05 < P < 0.06$ (close to significance), respectively).

Figure S3. Risk-return signals detected by the conventional analyses.

(**A**) An example activity histogram of a VS neuron modulated by both the probability and

magnitude of reward with opposite signs (i.e., negative b_p and positive b_m). The activity

aligned to cue onset is represented for three different levels of the probability (0.1-0.3, 0.4-

0.7, 0.8-1.0) and magnitude (0.1-0.3 mL, 0.4-0.7 mL, 0.8-1.0 mL) of rewards. Gray

hatched areas indicate a 1 s time window to estimate the neural firing rates shown in **B**.

The neural modulation pattern was defined as the Risk-Return type based on the linear

regression and AIC-based model selection, and as Magnitude type based on the BIC-

based model selection. Regression coefficients were -2.44 (*P* = 0.039) and 4.86 (*P* <

0.001) for the probability and magnitude, respectively. (**B**) Activity plots of the VS neuron

during the 1 s time window shown in **A** against the probability and magnitude of rewards.

(**C-D**) same as **A-B**, but for a cOFC neuron. The neural modulation type was defined as

the Risk-Return type based on all the three analyses. Regression coefficients were -6.65

(*P* < 0.001) and 3.82 (*P* < 0.001) for the probability and magnitude, respectively.

Figure S4. Effects of the analysis window size on the PCA.

(**A**) Cumulative variances explained by PCA in the four neural populations. Dashed lines indicate the percentages of variance explained by PC1 and PC2 in each neural population. The size of the analysis windows was 0.1, 0.05, and 0.02 s, respectively. (**B**) Overlay plots of the series of eigenvectors in the four neural populations. The eigenvectors for PC1 and PC2 are shown. The size of the analysis windows was 0.1, 0.05, and 0.02 s, respectively. a.u. indicates arbitrary units. (**C**) Box plots of the deviance from the mean vector estimated in each neural population are shown for the PC1. (**D**) same as (**C**), but for the PC2. (**E**) Box plots of the vector size estimated in each neural population are shown for the PC1. (**F**) same as (**E**), but for the PC2. In **C-F**, asterisks indicate statistical significance between two neural populations using Wilcoxon rank-sum test with Bonferroni correction for multiple comparisons (**, *, and § indicate statistical significance at $P < 0.01$, $P < 0.05$, and $0.05 < P < 0.06$ (close to significance), respectively).

Figure S5. Effects of the analysis window size on the conventional analyses.

(**A**) Percentage plots of the neural modulation type defined by using the model selection based on BIC values. The data for 0.1 s analysis windows is shown for the four neural populations. M: Magnitude type, P: Probability type, EV: Expected value type, R-R: Risk-Return type. (**B**) same as **A**, but for the data in 0.05 s analysis window. (**C**) same as **A**, but for the data in 0.02 s analysis window.

68 **Supplemental table**

EV	Probability									
	.10	.20	.30	.40	.50	.60	.70	.80	.90	1.0
.10	.01	.02	.03	.04	.05	.06	.07	.08	.09	.10
.20	.02	.04	.06	.08	.10	.12	.14	.16	.18	.20
.30	.03	.06	.09	.12	.15	.18	.21	.24	.27	.30
.40	.04	.08	.12	.16	.20	.24	.28	.32	.36	.40
.50	.05	.10	.15	.20	.25	.30	.35	.40	.45	.50
.60	.06	.12	.18	.24	.30	.36	.42	.48	.54	.60
.70	.07	.14	.21	.28	.35	.42	.49	.56	.63	.70
.80	.08	.16	.24	.32	.40	.48	.56	.64	.72	.80
.90	.09	.18	.27	.36	.45	.54	.63	.72	.81	.90
1.0	.10	.20	.30	.40	.50	.60	.70	.80	.90	1.0

(Magnitude (ml))

72 **Table S1. Payoff matrix regarding probability and magnitude of rewards.**

73 Payoff matrix of the probability and magnitude of rewards indicated by the blue and green

74 pie chart. The expected values [EV] defined as probability times magnitude are shown for

75 the 100 possible combinations. All the 100 possible combinations of the reward probability

76 and magnitude were used during the single cue and choice task.