Cyanobacterial growth and morphology are influenced by carboxysome positioning and temperature

Rees Rillema[a*], Joshua S. MacCready[a], Anthony G. Vecchiarelli[a#]

[a] Department of Molecular, Cellular, and Developmental Biology, University of Michigan, Ann Arbor, MI, USA.

* Present Address: BioMolecular Science Program, Michigan State University, East Lansing, MI, USA.

Running Title: Carboxysome mispositioning alters cell physiology

[#]**Address correspondence to Anthony G. Vecchiarelli:** ave@umich.edu

ABSTRACT: Cyanobacteria are the prokaryotic group of phytoplankton responsible for a

significant fraction of global CO_2 fixation. Like plants, cyanobacteria use the enzyme Ribulose

1,5-bisphosphate Carboxylase/Oxidase (RuBisCO) to fix CO_2 into organic carbon molecules via

the Calvin-Benson-Bassham cycle. Unlike plants, cyanobacteria evolved a carbon concentrating

organelle called the carboxysome - a proteinaceous compartment that encapsulates and

concentrates RuBisCO along with its CO_2 substrate. In the rod-shaped cyanobacterium

Synechococcus elongatus PCC7942, we recently identified the McdAB system responsible for

uniformly distributing carboxysomes along the cell length. It remains unknown what role

carboxysome positioning plays with respect to cellular physiology. Here, we show for the first

time that a failure to distribute carboxysomes leads to a temperature-dependent decrease in cell

growth rate, cell division arrest, cell elongation, asymmetric cell division, and a significant

reduction in cellular levels of RuBisCO. Unexpectedly, we also report that even wild-type *S.*

elongatus undergoes filamentous growth at the cool, but environmentally-relevant, growth

temperature of $20^{\circ}C$. The findings suggest that carboxysome positioning by the McdAB system

functions to maintain the carbon-fixation efficiency of RuBisCO by preventing carboxysome

aggregation, which is particularly important at temperatures where rod-shaped cyanobacteria

adopt a filamentous morphology.

IMPORTANCE: Photosynthetic cyanobacteria are responsible for almost half of global CO_2

fixation. Due to eutrophication, rising temperatures, and increasing atmospheric CO_2

concentrations, cyanobacteria have recently gained notoriety for their ability to form massive

blooms in both freshwater and marine ecosystems across the globe. Like plants, cyanobacteria

use the most abundant enzyme on Earth, RuBisCO, to provide the sole source of organic carbon

required for its photosynthetic growth. Unlike plants, cyanobacteria have evolved a carbon-

concentrating organelle called the carboxysome that encapsulates and concentrates RuBisCO

with its CO_2 substrate to significantly increase carbon-fixation efficiency and cell growth. We

recently identified the positioning system that distributes carboxysomes in cyanobacteria.

However, the physiological consequence of carboxysome mispositioning in the absence of this

distribution system remains unknown. Here we find that carboxysome mispositioning triggers

temperature-dependent changes in cell growth and morphology as well as a significant reduction

in cellular levels of RuBisCO.

INTRODUCTION

Cyanobacteria represent a phylum of diverse prokaryotic organisms where many

fundamental biological processes have remained largely understudied. As the evolutionary

ancestor of algae and plant chloroplasts, all cyanobacteria perform oxygenic photosynthesis and

fix carbon dioxide through the Calvin-Benson-Bassham cycle. Unlike chloroplasts,

cyanobacteria encapsulate their RuBisCO and carbonic anhydrase within large (~150 nm)

selectively permeable protein-based organelles called carboxysomes. This mechanism generates

an environment around RuBisCO that is significantly enriched in CO_2, which increases the

carboxylation activity of RuBisCO, while simultaneously reducing photorespiration (1).

In the model rod-shaped cyanobacterium *Synechococcus elongatus* PCC 7942

(hereafter *S. elongatus*), carboxysomes were found to be uniformly distributed down the length

of individual cells (2). This equidistant positioning supports equal inheritance of carboxysomes

following cell division and maximum diffusion of substrates and products across the

carboxysome shell. Carboxysomes are essential for the growth and survival of all cyanobacteria

and are responsible for ~35% of global carbon-fixation through atmospheric CO_2 assimilation

(Dworkin, 2006; Kerfeld and Melnicki, 2016). However, it remains unknown how the

subcellular organization of carboxysomes influences cyanobacterial physiology; a question of

considerable ecological, evolutionary, and biotechnological importance.

Savage *et. al.* were first to report that a ParA-type ATPase (hereafter McdA

- Maintenance of carboxysome distribution A) is required for positioning carboxysomes in *S.*

elongatus (2). ParA family members have well established roles in the segregation of bacterial

chromosomes and plasmids (4, 5). Less studied are ParA family members shown to be required

in the positioning of diverse protein complexes, such as those involved in secretion (6, 7),

chemotaxis (8–10), conjugation (11), cell division (12, 13), and cell motility (14, 15), as well as

bacterial microcompartments (BMCs), such as the carboxysome (2, 16). We recently identified a

small novel protein, McdB, responsible for generating dynamic McdA gradients on the nucleoid

(16). McdB colocalizes and directly interacts with carboxysomes and removes McdA from the

nucleoid in their vicinity. We found that carboxysomes use a Brownian-ratchet mechanism

whereby McdB-bound carboxysome motion occurs in a directed and persistent manner toward

increased concentrations of McdA on the nucleoid. We also recently found that the McdAB

system is widespread among β-cyanobacteria (17); an incredibly diverse and widely distributed

phylum of bacteria that display complex morphologies (18). Together, the data suggest that the

equidistant positioning of carbon-fixing carboxysomes in cyanobacteria is important, but the

physiological consequences of carboxysome mispositioning in the absence of the McdAB system

remain unclear.

When Savage *et. al.* first identified the McdA requirement for carboxysome positioning,

only a minor decrease in CO_2 fixation was found in a Δ*mcdA* strain, but growth at 30ºC under

ambient CO_2 was the only condition studied (2). Rising sea surface temperatures and coastal

eutrophication in marine ecosystems are pervasive effects of climate change (19). Rising

temperatures increase enzyme kinetics (20), leading to higher rates of protein synthesis,

RuBisCO activity, and photosynthesis (21). Temperature therefore directly influences

cyanobacterial growth and metabolic rates (22, 23).

Here, we show for the first time that a failure to distribute carboxysomes leads to

temperature-dependent decreases in growth rate, cell division arrest, cell elongation, asymmetric

cell division, and a significant reduction in cellular levels of RuBisCO. We also report that,

unexpectedly, wild-type *S. elongatus* undergoes filamentous growth at 20ºC; an environmentally

relevant growth temperature for *S. elongatus* not commonly used in the lab. We propose that

carboxysome positioning by the McdAB system functions as part of an autotrophic growth

strategy that maintains the carbon-fixation efficiency of RuBisCO by preventing carboxysome

aggregation. In the absence of carboxysome positioning or when cells are grown at cooler

temperatures, we propose that cell elongation and asymmetric cell division are responses to

organic carbon limitation, due to the decreased enzymatic activity of RuBisCO under these

conditions.

RESULTS

We performed *in vivo* microscopy to determine how carboxysome organization was altered

in *mcdA*, *mcdB*, and *mcdAB* deletion strains compared to wild-type *S. elongatus*. In *S. elongatus*,

each RuBisCO enzyme is composed of eight Large (Rbc<u>L</u>) and eight Small (Rbc<u>S</u>) subunits to

form $RbcL_8S_8$ (24). Therefore, to image carboxysomes, the fluorescent protein mTQ (monomeric

Turquoise2) was fused to the C-terminus of RbcS to make RbcS-mTQ (see Methods for strain

construction details). *RbcS-mTQ* was expressed using a second copy of its native promoter

(inserted at neutral site 1) in addition to wild-type *rbcS* at its native locus in wild-type cells as

well as in the three deletion strains – Δ*mcdA*, Δ*mcdB*, and Δ*mcdAB*. The presence of RbcS-mTQ

did not alter growth rate **(Supplemental Figure S1)**. Carboxysomes of *S. elongatus* encapsulate

between 800 to 1,500 RuBisCO enzymes (25). Therefore, RbcS-mTQ provided a bright and

high-contrast marker for quantifying the subcellular organization of carboxysomes, as previously

shown (16). We also performed Phase Contrast imaging to monitor for potential changes in cell

morphology, and Chlorophyll fluorescence imaging to verify that cells treated in our analyses

were photosynthetically active and therefore viable.

In the lab, *S. elongatus* is typically grown at 32°C; with a doubling time on the order of

several hours depending on growth conditions (temperature, light, and CO_2 availability). This

was the temperature used in our previous study that identified the McdAB carboxysome

positioning system (16). However, *S. elongatus* was recently shown to grow faster at 40°C (26,

27). Therefore, in anticipation of a growth defect in the absence of carboxysome positioning in

our Mcd system mutants, we began our studies by growing cultures at 40°C in constant light

under ambient (0.04%) or high (2%) CO_2 concentrations.

Carboxysome mispositioning correlates with minor defects in cellular physiology at 40°C

We first verified that carboxysomes are organized by the McdAB system at 40°C, as we

found previously at 32°C (16). We examined carboxysome positioning under high CO_2

conditions (2% CO_2) to maximize growth. We found that RbcS-mTQ labeled carboxysomes

were equally spaced down the long axis of wild-type *S. elongatus* cells **(Figure 1A)**. The Δ*mcdA*,

Δ*mcdB*, and Δ*mcdAB* mutants lost this uniform positioning of carboxysomes **(Figure 1B-D)**; as

observed previously at 32°C (16).

We compared the nearest-neighbor spacing of carboxysome foci as a function of cell

length **(Figure 1E)**. Wild-type cells showed uniform carboxysome spacing (0.55 ± 0.20 μm)

regardless of cell length. All three mutants, on the other hand, displayed a gradual increase in the

nearest-neighbor spacing of carboxysome foci as cell length increased. Both the median and

variability in spacing were greater in the mutants compared to wild-type **(Figure 1F)**. The

increased spacing resulted in fewer carboxysome foci per cell **(Figure 1G)**, and within single

cells, there were fewer carboxysome foci per unit cell length **(Figure 1H)**. When quantifying the

fluorescence intensity **(Figure 1I)** and size of carboxysome foci **(Figure 1J)**, it became apparent

that the increased spacing in all three mutant populations was not likely a result of fewer

carboxysomes being assembled, but rather, carboxysomes were coalescing into massive

aggregates. Indeed, when comparing carboxysome foci number across cell populations, we find

that ~ 95 % of wild-type cells (n = 440) had three or more foci, whereas ~ 80 % of all mutant

populations had three or fewer foci **(Figure 1K)**. Roughly 20% of cells from all three mutant

populations had a single carboxysome aggregate, whereas wild-type cells never had less than two

foci. The 40°C data shows that the McdAB system not only equally spaces carboxysomes down

the cell length, it also serves as an anti-aggregator; preventing carboxysomes from coalescing.

We then asked if carboxysome mispositioning affected cell physiology at 40°C. Wild-

type *S. elongatus* cells were 3.1 ± 0.5 microns in length **(Figure 2A)** and 1.25 ± 0.05 microns in

width **(Figure 2B)**. We found that Δ*mcdA* cells were of a similar length and width compared to

that of wild-type **(Figure 2AB)**. The Δ*mcdB* and Δ*mcdAB* cells had a slightly longer median cell

length, but a significantly wider distribution **(Figure 2A)**. These mutants were also thinner

compared to wild type **(Figure 2B)**. The presence of both longer and shorter cells suggested asymmetric cell division events. We quantified the frequency of symmetric (mid-cell) versus asymmetric (non-mid-cell) division events and found that asymmetric division was exclusive to the Δ*mcdB* and Δ*mcdAB* populations **(Figure 2C)**. The data show that while both McdA and McdB are required for positioning carboxysomes, the loss of McdB elicits an asymmetric division phenotype at 40°C.

The changes in cell morphology suggested that, even with high CO_2 levels, carboxysome mispositioning may alter cell growth. However, we found only modest reductions in growth rate for all three mutants **(Figure 2D-E)**. Even when grown slowly in ambient CO_2 (0.04%), we did not observe significant differences in growth rate compared to wild-type **(Figure 2F-G)**. Overall, at 40°C, carboxysomes are mispositioned in all three Mcd mutants, with moderate increases in carboxysome spacing due to aggregation, which ultimately results in fewer carboxysome foci per cell. We found an asymmetric cell division phenotype exclusive to cell without McdB. But in all deletion strains, growth rate was not significantly slower than wild type, with or without CO_2, at this temperature.

Carboxysome mispositioning correlates with cell division arrest, cell elongation, and asymmetric cell division at 30°C

We continued our study at 30°C, a growth temperature closer to what we used to first identify the McdAB carboxysome positioning system (16). As we showed previously, and similar to our 40°C data, wild-type *S. elongatus* cells have equally spaced carboxysomes **(Figure 3A)**, while in all three mutants, carboxysomes were mispositioned **(Figure 3B-D)**. Wild-type cells displayed the same carboxysome spacing distance (0.50 ± 0.20 mm) regardless of cell

length, while all three mutants had increased carboxysome spacing, and variability in spacing, as

cell length increased **(Figure 3E-F)**. Intriguingly, we found that mutant cell lengths were

significantly longer compared to wild-type. This cell elongation phenotype was more extreme in

the Δ*mcdB* and Δ*mcdAB* mutants, resulting in more distantly spaced carboxysome foci **(Figure**

3F). The data once again shows that McdB plays a currently unknown role in carboxysome

function, separate from its role in positioning with McdA.

The increased spacing resulted in fewer carboxysome foci per cell **(Figure 3G)**, and per

unit cell length **(Figure 3H)**. Carboxysome foci in all three mutants were significantly larger

than that of wild-type, suggesting aggregation **(Figure 3I-J)**. While ~ 90 % of wild-type cells (n

= 486) had three or more foci, ~ 70 % of all mutant populations had three or fewer foci **(Figure**

3K). Again, ~ 20 % of cells in all three mutant populations contained a single carboxysome

aggregate. This is a striking reduction in carboxysome foci when considering the cell elongation

phenotype exclusive to the mutant cell lines.

As the carboxysome spacing data suggested, all three mutants had significantly longer

cell lengths compared to wild-type when grown at 30°C **(Figure 4A)**. Median cell width was

similar across all strains, however Δ*mcdB* and Δ*mcdAB* cell populations displayed significantly

wider distributions in width **(Figure 4B)**. Once again, the Δ*mcdB* and Δ*mcdAB* mutant

populations displayed a significant number of asymmetrical division events (~ 70 % of dividing

cells) compared to wild-type or Δ*mcdA* **(Figure 4C)**. We propose that carboxysome

mispositioning and aggregation at 30°C elicits cell division arrest, cell elongation and, when

McdB is absent, asymmetric cell division; phenotypes that were largely masked when cells were

grown at 40°C.

183   The changes in cell morphology suggested that the mutants may display slower growth

184 rates, even when grown at high CO_2. However, that was not the case. Under high CO_2

185 conditions, no significant reduction in growth rate was observed for all three mutants when

186 compared to wild-type **(Figure 4D-E)**. Even when grown in ambient CO_2, reductions in growth

187 rate of the mutants were minor **(Figure 4F-G)**.

188   Overall, we find that at 30°C, carboxysomes are mispositioned in the $\Delta mcdA$, $\Delta mcdB$,

189 and $\Delta mcdAB$ mutants; with drastic increases in carboxysome spacing due to aggregation,

190 resulting in fewer carboxysome foci per cell. We unveiled a cell elongation phenotype for all

191 three mutant populations, and strikingly, we also found an asymmetric cell division phenotype

192 that was exclusive to cells lacking McdB; a phenotype also found at 40°C but exacerbated at

193 30°C **(compare Panel C in Figures 2 and 4)**. Despite these carboxysome aggregation and cell-

194 morphology phenotypes, only minor decreases in growth rate were observed. We propose that

195 carboxysome aggregation decreases the carbon-fixing activity of encapsulated RuBisCO, which

196 triggers an organic carbon-limitation response - cell division arrest, elongation, and asymmetric

197 cell division.

199 ***S. elongatus* elongates at 20°C and this mode of growth is slower in Mcd system mutants**

200   A ten degree drop in growth temperature unveiled the physiological consequences of

201 carboxysome mispositioning at 30°C, which were largely masked at 40°C. The catalytic rate of

202 carboxysome-encapsulated RuBisCO is often considered the bottle neck of photosynthesis

203 because the enzyme is inefficient (28) and temperature dependent (20). The light reactions of

204 photosynthesis, however, are considered temperature independent (20, 29). We therefore decided

205 to study the effects of carboxysome mispositioning when cells were grown at 20°C, which is

206   within the environmentally relevant temperature range - *S. elongatus* PCC 7942 was originally

207   isolated from a fresh water source in the San Francisco bay area, with an annual temperature

208   range of 8°C to 25°C (30).

209   Unexpectedly, we found that even wild-type *S. elongatus* undergoes extreme cell

210   elongation when grown at 20°C **(Figure 5A)**. *S. elongatus* has been previously studied at this

211   temperature (31), but to our knowledge, this is the first time cell morphology has been directly

212   observed. Despite the dramatically long cell lengths, carboxysomes were still robustly aligned

213   down the entire longitudinal axis of the cell **(Figure 5A)**. All three mutants were also elongated,

214   but to a lesser extent, and carboxysomes were clearly mispositioned **(Figure 5B-D)**.

215   Wild-type showed the same uniform carboxysome spacing distance (0.50 ± 0.30 μm)

even in cells as along as 20 microns **(Figure 5E)**. All three mutants, on the other hand, displayed

dramatically increased carboxysome spacing, and variability in spacing, as cell length increased

(Figure 5F). Since all cell types dramatically elongated at 20°C, the loss of carboxysome

positioning resulted in a massive reduction in the number of carboxysome foci per cell **(Figure**

5G), and per unit cell length **(Figure 5H)**. Carboxysome foci in all three mutants were

significantly larger than that of wild-type, suggesting carboxysome aggregation **(Figure 5I-J)**.

While ~ 90% of wild-type cells (n=388) had six or more foci, ~ 80% of all mutant populations

had less than five **(Figure 5K)**. The data implicates carboxysome positioning by the McdAB

system as a requirement for maximizing carbon-fixation in cells that are elongated when grown

at cooler but environmentally relevant temperatures.

226   We quantified cell length at 20°C and found that the median length of wild-type cells was

significantly longer than all three mutant populations **(Figure 6A)**. The Δ*mcdA* cells were

similar in width to that of wild-type, while Δ*mcdB* and Δ*mcdAB* cell populations were notably

thinner **(Figure 6B)**. Consistent with the wide distributions in cell length **(Figure 6A)**, a

significant fraction of division events at this growth temperature were asymmetric across all cell

populations. But the frequency of asymmetric division events was still highest in the strains

lacking McdB **(Figure 6C)**.

When growth rates were assayed at high CO_2 **(Figure 6D-E)** or ambient CO_2 **(Figure 6F-**

G), we finally found statistically significant reductions in growth rate for all three mutants when

compared to wild-type. However, the reductions in growth rate depended on whether McdA or

McdB was absent, and whether the cells were grown with high or ambient CO_2. First, the $\Delta mcdA$

growth rate was 2-times slower than wild-type at high CO_2 **(Figure 6E)**. But when grown in

ambient CO_2, the $\Delta mcdA$ growth rate was similar to wild-type **(Figure 6G)**. It is known that

carboxysome quantity significantly increases when *S. elongatus* is grown in ambient CO_2 (32).

Therefore, our data suggest that with ambient CO_2, the greater number of carboxysomes

compensates for a loss in their positioning. Alternatively, a decrease in growth rate for $\Delta mcdB$

and $\Delta mcdAB$ was found with high- or ambient-levels of CO_2 **(Figure 6E and G)**. The data once

again shows that McdB plays a key role in carboxysome function outside of its role in

positioning with McdA.

Overall, we identified a cell elongation phenotype in wild-type *S. elongatus* cells that

occurs when grown at colder, but environmentally relevant temperatures. All three Mcd system

mutants elongated to a lesser extent and have aggregated carboxysomes, which resulted in

drastically fewer carboxysome foci per cell compared to wild-type. Significant decreases in

growth rate were observed, even with high CO_2 present. The findings suggest that in Mcd mutant

or wild-type cells, colder growth temperatures decrease the enzyme kinetics of RuBisCO to the

point where cell division arrest, elongation, and asymmetric cell division is triggered in response

to carbon-limitation. At 30ºC, this response was only found in the Mcd mutants likely due to

reduced carbon-fixation efficiency resulting from carboxysome aggregation. At 40ºC, the

enzyme kinetics of RuBisCO are fast enough to compensate for carboxysome aggregation, which

would explain why Mcd mutants largely displayed cell morphologies and growth rates similar to

that of wild-type at higher growth temperatures.

McdAB mutants have significantly decreased levels of intracellular RuBisCO

RuBisCO is the sole enzyme providing organic carbon to cellular biomass production and

the phototrophic growth of *S. elongatus*. Therefore, we set out to determine if carboxysome

mispositioning and the changes in cell morphology of McdAB mutants at 30ºC correlated with

altered cellular levels of RuBisCO. Immunoblot analysis against the Large subunit of RuBisCO

(RbcL) showed that RuBisCO content is roughly half in all three mutants compared to wild-type

cells **(Figure 7A-B)**. RbcL content from cell cultures was normalized based on Atpβ quantity

from immunoblot analysis as shown previously (33, 34). The data suggest that although the

mutants house massive carboxysome aggregates that are much brighter than those uniformly

positioned in wild-type cells **(Supplemental Figure S2)**, the loss of a functioning McdAB

system results in a reduction in the total cellular content of RuBisCO.

We continued our quantification of cellular RuBisCO content using the RbcS-mTQ

fluorescence signal. When quantifying the average RbcS-mTQ intensity per cell, only wild-type

seemed to increase RuBisCO content in response to decreasing growth temperature **(Figure 7C)**.

This finding is consistent with previous studies showing that RuBisCO abundance in

phytoplankton populations (35), including cyanobacterial species (44), inversely correlates with

growth temperature. We then normalized the RbcS-mTQ signal to cell length to account for the

differences in cell morphology with Mcd system mutants and varying growth temperatures. At

40°C, RuBisCO content was similar across all cell populations, albeit more variable in the

mutants **(Figure 7D)**. At 30°C, all three mutants had roughly 2-fold less RuBisCO compared to

that of wild type **(Figure 7E)**; a decrease that is consistent with our immunoblot quantification at

32°C **(see Figure 7B)**. The data confirm that the immunoblot analysis using Atpβ standardization

provided a valid measure of cellular RuBisCO abundance, and that the RbcS-mTQ intensity

measurements *in vivo* are a reasonable proxy for RuBisCO content. Finally, at 20°C, all cell

populations including wild-type had the lowest levels of RuBisCO content when normalized to

cell length **(Figure 7F)** compared to that at higher temperatures **(Figure 7D-E)**. Also, cells

without McdB had almost 2-fold less RuBisCO compare to wild-type or Δ*mcdA* cells **(Figure**

7F). Overall, our findings suggest that carboxysome mispositioning results in decreased levels of

cellular RuBisCO, and unveiled a critical but currently unknown role for McdB in carboxysome

integrity and carbon-fixing function.

DISCUSSION

We recently identified the McdAB system responsible for the equidistant positioning of

carboxysomes in the rod-shaped cyanobacterium *S. elongatus* PCC 7942 (16). We also recently

found that McdAB systems are widespread among β-cyanobacteria (17). These findings suggest

important and widespread, but currently unknown, physiological roles for carboxysome

positioning. Here, we show for the first time that a failure to distribute carboxysomes leads to a

number of temperature-dependent changes in cell physiology: decrease in cell growth rate, cell

division arrest, cell elongation, asymmetric cell division, and a significant reduction in cellular

levels of RuBisCO.

At the three temperatures tested (20, 30, and 40°C), all three mutant cell-types housed

few and irregularly spaced carboxysome aggregates, compared to wild-type cells with uniformly-

spaced and -sized carboxysome foci. The data here provides quantitative support for our previous

findings that the McdAB system equally distributes carboxysomes to opposite sides of the cell to

ensure their inheritance following cell division (2, 16), akin to ParA-based plasmid partition

systems in bacteria (4). But in addition, and particularly important for protein-based cargoes, the

McdAB system serves to prevent carboxysome aggregation. This 'anti-aggregation' activity

serves as a homeostasis mechanism that regulates carboxysome size, number, composition,

positioning, and ultimately, its carbon-fixing function in the cell. It has recently been found that

the pyrenoid, the functional analog of the carboxysome in the chloroplasts of the model alga

Chlamydomonas, is also spatially regulated and this activity affects carbon-fixation (36). It

remains to be seen if other BMCs also use spatial regulation mechanisms to avoid aggregation

during their self-assembly to optimize enzymatic efficiency.

Physiological defects associated with carboxysome mispositioning are temperature-

dependent.

Despite the drastic mispositioning and aggregation of carboxysomes in the *mcdA*, *mcdB*,

and *mcdAB* deletion strains of *S. elongatus*, only minor changes in cell morphology and growth

rate were found when compared to wild-type at the most optimal growth temperature used in this

study (40°C) **(Figures 2)**. At 30°C however, all three mutants displayed filamentous growth, and

a significant fraction of cell division events in the mutant populations lacking McdB were

asymmetric **(Figure 4)**. Despite these changes to cell morphology, the growth rates were similar

to that of wild-type; even in ambient CO_2. At our coldest but environmentally relevant growth

temperature of 20ºC, we found that all cell populations, even wild-type, underwent filamentous

growth and asymmetric cell division (Figure 6). Wild-type cells at 20ºC were as much as 5 to

10-times longer than the median length when grown at 30ºC or 40ºC. Even in these extremely

elongated cells, the McdAB system robustly distributed carboxysomes down the entire cell

length (Figure 5A). This mode of filamentous growth was slower in all Mcd system mutants,

resulting in significantly shorter filaments compared to wild-type (Figure 6). Overall, we find

temperature-dependent physiological defects associated with carboxysome mispositioning and

aggregation; phenotypes that were masked at high growth rates and temperatures typically used

in the lab, and unveiled at lower but environmentally relevant temperatures.

Cell division arrest, filamentation, and asymmetric cell division – A stress response to

carbon-limitation?

Many bacteria can change shape in response to growth conditions (37). *E. coli* and *B.*

subtilis produce larger cells under nutrient-rich conditions and smaller cells under nutrient-

limited conditions (38, 39). *Pseudomonas aeruginosa* elongates to enhance nutrient uptake

during carbon and nitrogen starvation (40); *Caulobacter crescentus* cells increase cell area in

response to phosphate starvation (41); and abundant human gut species such as *Bacteriodes*

thetaiotaomicron elongate under sugar-limited conditions (42). Our findings suggest that *S.*

elongatus filamentation and asymmetric cell division are responses triggered by organic carbon-

limitation, resulting from reduced carbon-fixation activity of RuBisCO in this obligate

photoautotroph. Cell filamentation (i) prevents the birth of cells devoid of carboxysomes, (ii)

gives the cell an opportunity to increase carbon-fixation by producing more RuBisCO as a larger

aggregate, or by nucleating the formation of additional carboxysomes *de novo* elsewhere along

the elongated cell and (iii) the increased cell-surface area would increase the light-harvesting

capability for photochemistry.

A carbon-limitation response parsimoniously explains the two triggers identified here: (1)

carboxysome aggregation and (2) colder growth temperatures. What is the mechanism behind the

carboxysome aggregation trigger? All Mcd system mutants displayed significant reductions in

their total cellular RuBisCO content (Figure 7). Therefore, carboxysome aggregation could

trigger the carbon-limitation response simply due to lower levels of RuBisCO as a result of

losses in carboxysome integrity and leakage, enzyme stability, and/or degradation. Alternatively,

wild-type cells have carboxysomes of a homogenous size, and as cells grow, so do the number of

carboxysomes and their net surface area. The net surface area of the carboxysome shell, where

RuBisCO substrates and products interface with the cytoplasm, significantly decreases due to

carboxysome aggregation. As a result, substrate and product permeability would suffer, which

could lead to decreased carbon-fixation rates by encapsulated RuBisCO. It is therefore also

possible that the carbon-limitation response is triggered by decreased RuBisCO activity due to

decreases in the effective surface area of aggregated carboxysomes. These two proposals are not

necessarily mutually exclusive from one another. Consistent with these proposals, it was recently

found in Synechococcus sp. PCC 7002 that mispositioning of inactivated carboxysomes towards

the cell poles is strongly correlated with their rapid degradation (35).

We propose this adaptive response to carbon-limitation is also triggered by low

temperature growth, even in wild-type *S. elongatus*. At high temperature (i.e. 40ºC), RuBisCO

activity is high (43) and sufficient to compensate for carboxysome mispositioning and RuBisCO

aggregation in Mcd system mutants. Therefore, the carbon-limitation response is not triggered.

At 30ºC, RuBisCO activity is still high enough to prevent a carbon-limitation response in wild-

type cells, but not in Mcd mutants with few and aggregated carboxysomes. As a result, cell

filamentation is triggered only in the Mcd mutants. At 20°C, RuBisCO activity has reduced to a

level that, even in wild-type cells, triggers the carbon-limitation response. Mcd mutants are

starved for carbon to the point where growth is slowed, resulting in shorter filaments compared

to wild-type.

Spatial regulation of carboxysomes influences RuBisCO activity and abundance

Psychrophilic phytoplankton species deal with low temperatures, well below the thermal

optimum of most enzymes, by increasing the abundance of RuBisCO (35). It has recently been

found that low temperature growth (18 to 22°C) results in increased RuBisCO levels in the

marine cyanobacterium *Synechococcus* PCC 7002, compared to growth at 26°C or 30°C (44).

This trend was also observed in natural phytoplankton assemblages across a wide latitudinal

range (35). Consistently, we found here that the RbcS-mTQ signal within carboxysome foci, and

within the entire cell, inversely correlated with decreasing growth temperatures.

Previous studies have also shown positive relationships between growth rate and

RuBisCO abundance (35, 44–48). Elevated carbon fixation rates during blooms in polar regions

have been shown to be associated with several-fold increases in RuBisCO. Twenty degrees

Celsius is well below the thermal optimum for RuBisCO activity (49), and well below the

optimal growth temperature of *S. elongatus*. We therefore conclude that increasing RuBisCO

abundance and triggering cell elongation both represent acclimation responses that compensate

for the decreased catalytic rate of RuBisCO at low-temperature growth. This response is further

compounded when carboxysomes are mispositioned and aggregated in our Mcd mutants. It is

important to note that filamentation occurred here under high CO_2 – a condition whereby *S.*

elongatus does not require the carbon-concentrating activity of carboxysomes for growth (50).

Therefore, the evidence further supports that it is not CO_2 substrate limitation causing

filamentation, but rather RuBisCO activity that is significantly compromised when complexed in

carboxysome aggregates or when cells are grown at colder temperatures. These findings shed

light on the importance of the McdAB system in maximizing the carbon-fixation activity of

RuBisCO by preventing aggregation during carboxysome self-assembly.

Overall, we show that carboxysomes are organized by the McdAB system in

cyanobacteria and can appropriately respond to variability in RuBisCO abundance and activity

along a wide temperature gradient. Our results suggest that carboxysome homeostasis provided

by the McdAB system is part of an autotrophic growth strategy in elongated cells - RuBisCO

abundance increases and is distributed into homogeneously-sized carboxysomes at low

temperature to overcome the lower catalytic rates of this temperature-dependent enzyme (35).

McdB plays a role in carboxysome function outside of its positioning with McdA

Regardless of growth temperature, Δ*mcdB* and Δ*mcdAB* mutants elicited stronger cell

elongation and asymmetric division phenotypes compared to Δ*mcdA*. Our findings suggest that

McdB plays a currently unknown but critical role in carboxysome function, outside of its role in

positioning with McdA. Consistently, our recent bioinformatics analysis of McdAB systems

across cyanobacteria identified numerous species with orphan McdBs, once again suggesting

functional roles independent of McdA (17). We have also shown previously via bacterial two-

hybrid analysis that McdA does not physically associate with any carboxysome component,

while McdB directly interacts with a number of shell proteins (16). We have also recently found

that purified McdB undergoes Liquid-Liquid Phase-Separation (LLPS) *in vitro* (17). This activity

is intriguing given recent studies showing that both α- and β-carboxysomes, as well as the algal

pyrenoid, have intrinsically-disordered proteins that from liquid-like condensates with RuBisCO

(51–53). Collectively, these studies suggest that LLPS is a common feature underlying

carboxysome biogenesis. It is intriguing to speculate that the LLPS activity of McdB and other

carboxysome components are related and potentially influence each other.

Cold growth temperature triggers elongation and asymmetric cell division

Across the bacterial world, complex systems maintain cell-size homeostasis. We find

here that *S. elongatus* maintains cell-size homeostasis when grown at 30°C or 40°C, but

undergoes filamentation when grown at the environmentally-relevant temperature of 20°C. A

growing number of bacterial species are known to elongate into a filamentous morphology

amidst environmental changes to promote survival. For example, *E. coli* cells become

filamentous during infection (54, 55), and during DNA damage, the SOS response blocks cell

division until damage has been repaired, resulting in elongation (56). Several other forms of

stress have been shown to induce elongation, including host environment, antibiotics, nutrient

access, pH, heat shock, and osmotic fluctuations (57–60). Bacteria can clearly exist in diverse

morphological states, in part dictated by their environment (37, 61).

To our knowledge, we find here the first example of bacterial filamentation caused by

cold-growth temperature. We propose colder temperatures trigger filamentation indirectly as a

carbon-limitation response due to the reduced carbon-fixing activity of RuBisCO. Alternatively,

transition into a filamentous morphology has been proposed to confer advantages, such as

avoiding phagocytosis via the host immune response (54, 55, 57, 61) or to protect against

predation in aquatic environments (58, 62–64). It is attractive to speculate that the growth

temperature of 20ºC coincides with the seasonal temperature during which predation occurs (63).

That is, *S. elongatus* may use temperature as a cue to elongate so as to avoid planktivorous

protists. Consistent with this possibility, previous studies have demonstrated that several

freshwater bacteria, including *Caulobacter crescentus* (58), exhibit high phenotypic plasticity

and can transition to a filamentous morphology – a transition that may be specifically triggered

in the presence of a size-selective protistan predator (63, 65, 66). It is possible that some aquatic

bacteria have evolved to undergo filamentation at temperatures that coincide with grazing

season.

Several studies have characterized the mechanisms by which bacterial filamentation is

induced, particularly under DNA damage or exposure to antibiotics (56). But how bacterial cells

enter and exit these filamentous states to ensure survival during changes in environment remains

poorly characterized and is a recent question of interest. We find here that *S. elongatus* forms

filaments when grown at low temperature or when carboxysomes are mispositioned. When these

filaments divide, it is asymmetric – forming a daughter cell of 'normal' length. It has been

recently shown that *S. elongatus* can also be induced to filament under dim-light stress, and then

divides asymmetrically to form daughter cells of the correct size when brought back into well-lit

conditions (67). Somewhat related, chlorophyll fluorescence along the cell length of our Mcd

mutants was much more heterogeneous, particularly at 20ºC, compared to wild-type **(see Figure**

5A-D). It should be noted that in addition to sensing light and inorganic carbon substrates, *S.*

elongatus responds to lowering temperatures by slowing and even turning off circadian rhythms;

thereby down regulating cell growth, division, and metabolic flux (68, 69).

The Min system has been shown to play a role in the asymmetric cell division and

daughter-cell sizing of filamentous *E. coli* (60), *V. parahaemolyticus* (70, 71), and *S. elongatus*

(67). Recent findings suggest that division restoration at the poles of these filaments is regulated

by a combination of Min oscillations, FtsZ levels and *terminus* segregation, resulting in daughter

cells of the right length (72, 73). The mechanism by which filamentation and asymmetric

division occurs in *S. elongatus* is an area of future research. Taken together, the conserved ability

for various bacterial species to undergo filamentation and morphological recovery, some of

which showing direct selective benefits, strongly suggests that this differentiation plays an

important role in survival and proliferation.

MATERIALS AND METHODS

Construct designs. All constructs in this study were generated using Gibson Assembly (74)

from synthetized dsDNA and verified by sequencing. Constructs contained flanking DNA that

ranged from 500 to 1500 bp in length upstream and downstream of the targeted insertion site to

promote homologous recombination into target genomic loci (75).

Generation of Bacterial Strains. All *Synechococcus elongatus* PCC7942 transformations were

performed as previously described (75). Plasmid constructs of *mcdA*, *mcdB*, and *mcdAB*

deletions were created by replacing the respective coding sequences with a kanamycin resistance

cassette. All fluorescent strains were transformed using plasmid pAH40, which contains a

chloramphenicol resistance cassette and a second copy of the *rbcS* promoter and gene, attached

at the 3' end with the gene encoding for fluorescent protein mTurquoise2 (mTQ) and separated

with a GSGSGS linker, inserted into neutral site 1. Transformed cells were plated on BG-11

agar. Single colonies were picked into 96-well plates containing 300 μL of BG-11 with 6.35 ug

ml^{-1} kanamycin or 3.125 ug ml^{-1} chloramphenicol. Concentrations of both respective antibiotics

were gradually increased to 12.5 μg ml^{-1}. Cultures were verified for complete insertion via PCR

and removed from antibiotics for experiments.

Growth Conditions. Both wild-type and mutant *Synechococcus elongatus* PCC7942 strains

were grown in 125 mL culture flasks (Corning) in 50 mL BG-11 (Sigma) medium buffered with

1 g L^{-1} HEPES to pH 8.3 shaken at 130 rpm or on BG-11 plates containing 1.5% (w/v) agar. All

strains were maintained and grown under constant LED illumination of 100 μE m^{-2} s^{-1} at either

20°C, 30°C, or 40°C in 2% or 0.04% CO_2 as specified. Cultures grown in 0.04% CO_2 were

grown in air with no additional CO_2. Cultures were regularly diluted with fresh medium to

maintain exponential growth phase for subsequent imaging and immunoblot analyses. For

cloning, One Shot™ TOP10 Chemically Competent *E. coli* (ThermoFisher) were grown

aerobically at 37° in Luria-Broth medium.

Fluorescence microscopy. Microscopy was performed using exponentially growing cells at an

OD of 0.4. Two milliliters of culture were spun down at 15,000 x g for 60 s, resuspended in 100

μL of BG-11. Five microliters were transferred to a square 1.5% agarose + BG-11 pad, which

was then flipped onto a 35 mm cell culture dish with a #1.5 glass coverslip bottom (ManTek).

All images were captured using a Nikon Eclipse Ti2 inverted microscope with a PlanApo

Objective lens (100x, 1.45NA, oil immersion), with phase contrast transillumination, and with a

SOLA LED light source for imaging chlorophyll fluorescence (Excitation: 560/40nm (540-

580nm), Emission: 630/70nm (593-668nm), Dichroic Mirror: 585nm) and RbcS-mTQ

fluorescence (Excitation: 436/20nm (426-446nm), Emission: 480/40nm (460-500nm), Dichroic

Mirror: 455nm). Images were acquired using a Photometrics Prime 95B Back-illuminated

sCMOS Camera. Image analysis was performed using Fiji (76) and the MicrobeJ plugin (77).

Growth Curve Measurements. Cultures were inoculated at a starting OD_{750} of 0.1-0.2 with

fresh BG-11. Cell growth was monitored at OD_{750} using a DS-11 Spectrophotometer (DENOVI)

at the specified time points. Growth rates were calculated using the linear portion of the

exponential phase of growth. Error bars represent the standard deviation from three biological

replicates that were recorded from different culture flasks.

MicrobeJ quantification. Multiple fields of view were taken for each cell strain using three

channels: Phase contrast provided cell perimeters, and fluorescence microscopy provided

chlorophyll autofluorescence and RbcS-mTQ intensities. These data were analyzed using

MicrobeJ (77). In each cell-line, cell length detection was performed using the rod-shaped

descriptor and thresholding set to 0.4µm < area < max, 0.71µm < width range < 2 µm, 0 µm <

width variation < 0.2 µm, and 0 µm < angularity amplitude < 0.35 µm. Carboxysome detection

was performed using the point function with a tolerance of 20 and an intensity minimum of 500.

Associations, shape descriptors, profiles, and distances were recorded for each strain. All

MicrobeJ quantification was also verified manually. Graphs and statistical analyses were

generated with Graph Pad Prism.

Immunoblot Analysis. Cells were lysed with a Qsonica sonication system (20 cycles - 30 s on,

10s off at 30% power) in 0.5 mL RuBisCO extraction buffer (50 mM EPPS at pH 8.1, 1% PVPP,

1 mM EDTA, 10 mM DTT, 0.1% Triton, and sigma protease inhibitor). Laemmli sample buffer

(0.1 mL of 4x) was added to lysate prior to loading 10 µL on a 4%–12% Bis-Tris NuPAGE gel

(Invitrogen). Gels were transferred onto a Mini-size polyvinylidene difluoride membrane (Bio-

Rad) using a Trans-Blot Turbo system (Bio-Rad). The membrane was immunoprobed using

rabbit polyclonal antisera against RbcL and the beta subunit of ATP synthase, AtpB (Agrisera)

and then goat anti-rabbit IgG secondary antibody (LI-COR). Membrane signals were visualized

and quantified at 600 nm using LI-COR Image Studio. For each sample, there were three

replicates from the same cell lysate. We normalized RbcL signal using AtpB; a method

previously used in Zhang *et al*., 2012 and specifically with *S. elongatus* PCC7942 in Sun *et al*.,

2016 (33, 34).

ACKNOWLEDGMENTS

We would like to thank Drs. Lyle Simmons, Vincent Denef, and Daniel Ducat for providing

critical feedback in experimental direction and data interpretation. We would also like to thank

Pusparanee Anne Hakim for experimental assistance and critical reading of the manuscript.

REFERENCES:

1.   Rae BD, Long BM, Badger MR, Price GD. 2013. Functions, compositions, and evolution

of the two types of carboxysomes: Polyhedral microcompartments that facilitate CO2

fixation in cyanobacteria and some proteobacteria. Microbiol Mol Biol Rev.

2.   Savage DF, Afonso B, Chen AH, Silver PA. 2010. Spatially Ordered Dynamics of the

Bacterial Carbon Fixation Machinery. Science (80-) 327:1258–1261.

3.   Kerfeld CA, Melnicki MR. 2016. Assembly, function and evolution of cyanobacterial

carboxysomes. Curr Opin Plant Biol 31:66–75.

4.   Baxter JC, Funnell BE. 2014. Plasmid Partition Mechanisms. Microbiol Spectr 2.

5.   Badrinarayanan A, Le TBK, Laub MT. 2015. Bacterial Chromosome Organization and

Segregation. Annu Rev Cell Dev Biol 31:171–199.

6.   Viollier PH, Sternheim N, Shapiro L. 2002. Identification of a localization factor for the

polar positioning of bacterial structural and regulatory proteins. Proc Natl Acad Sci USA

99:13831–13836.

7.   Perez-Cheeks BA, Planet PJ, Sarkar IN, Clock SA, Xu Q, Figurski DH. 2012. The product

of tadZ, a new member of the parA/minD superfamily, localizes to a pole in

Aggregatibacter actinomycetemcomitans. Mol Microbiol 83:694–711.

8.   Thompson SR, Wadhams GH, Armitage JP. 2006. The positioning of cytoplasmic protein

clusters in bacteria. Proc Natl Acad Sci USA 103:8209–8214.

9.   Ringgaard S, Schirner K, Davis BM, Waldor MK. 2011. A family of ParA-like ATPases

promotes cell pole maturation by facilitating polar localization of chemotaxis proteins.

Genes Dev 25:1544–1555.

10.   Alvarado A, Kjær A, Yang W, Mann P, Briegel A, Waldor MK, Ringgaard S. 2017.

Coupling chemosensory array formation and localization. Elife 6.

11.   Atmakuri K, Cascales E, Burton OT, Banta LM, Christie PJ. 2007. Agrobacterium

ParA/MinD-like VirC1 spatially coordinates early conjugative DNA transfer reactions.

EMBO J 26:2540–2551.

12.   Raskin DM, De Boer PAJ. 1999. Rapid pole-to-pole oscillation of a protein required for

directing division to the middle of Escherichia coli. Proc Natl Acad Sci U S A.

13.   Jordan A, Chandler J, MacCready JS, Huang J, Osteryoung KW, Ducat DC. 2017.

Engineering Cyanobacterial Cell Morphology for Enhanced Recovery and Processing of

574   Biomass. Appl Env Microbiol 83.

575   14.   Youderian P, Burke N, White DJ, Hartzell PL. 2003. Identification of genes required for

576   adventurous gliding motility in Myxococcus xanthus with the transposable element

577   mariner. Mol Microbiol 49:555.

578   15.   Kusumoto A, Shinohara A, Terashima H, Kojima S, Yakushi T, Homma M. 2008.

579   Collaboration of FlhF and FlhG to regulate polarflagella number and localization in Vibrio

580   alginolyticus. Microbiology.

581   16.   MacCready JS, Hakim P, Young EJ, Hu L, Liu J, Osteryoung KW, Vecchiarelli AG,

582   Ducat DC. 2018. Protein gradients on the nucleoid position the carbon-fixing organelles of

583   cyanobacteria. Elife 7.

584   17.   MacCready JS, Basalla JL, Vecchiarelli AG. 2020. Origin and Evolution of Carboxysome

585   Positioning Systems in Cyanobacteria. Mol Biol Evol 37:1434–1451.

586   18.   Shih PM, Wu D, Latifi A, Axen SD, Fewer DP, Talla E, Calteau A, Cai F, Tandeau De

587   Marsac N, Rippka R, Herdman M, Sivonen K, Coursin T, Laurent T, Goodwin L, Nolan

588   M, Davenport KW, Han CS, Rubin EM, Eisen JA, Woyke T, Gugger M, Kerfeld CA.

589   2013. Improving the coverage of the cyanobacterial phylum using diversity-driven

590   genome sequencing. Proc Natl Acad Sci U S A.

591   19.   Doney SC, Ruckelshaus M, Emmett Duffy J, Barry JP, Chan F, English CA, Galindo HM,

592   Grebmeier JM, Hollowed AB, Knowlton N, Polovina J, Rabalais NN, Sydeman WJ,

593   Talley LD. 2012. Climate Change Impacts on Marine Ecosystems. Ann Rev Mar Sci.

594   20.   Geider RJ. 1987. Light and Temperature Dependence of the Carbon to Chlorophyll a

595   Ratio in Microalgae and Cyanobacteria: Implications for physiology and growth of

596   phytoplankton. New Phytol.

21.   Raven JA, Geider RJ. 1988. Temperature and algal growth. New Phytol.

22.   Eppley R. 1972. Temperature and phytoplankton growth in the sea. Fish Bull.

23.   Chen B, Liu H, Huang B, Wang J. 2014. Temperature effects on the growth rate of marine

picoplankton. Mar Ecol Prog Ser.

24.   Tabita FR, Hanson TE, Satagopan S, Witte BH, Kreel NE. 2008. Phylogenetic and

evolutionary relationships of RubisCO and the RubisCO-like proteins and the functional

lessons provided by diverse molecular formsPhilosophical Transactions of the Royal

Society B: Biological Sciences.

25.   Sun Y, M Wollman AJ, Huang F, Leake MC, Liu L-N. 2019. Single-Organelle

Quantification Reveals Stoichiometric and Structural Variability of Carboxysomes

Dependent on the Environment. Plant Cell Adv Publ.

26.   Ohbayashi R, Nakamachi A, Hatakeyama TS, Watanabe S, Kanesaki Y, Chibazakura T,

Yoshikawa H, Miyagishima S. 2019. Coordination of Polyploid Chromosome Replication

with Cell Size and Growth in a Cyanobacterium. MBio 10.

27.   Yu J, Liberton M, Cliften PF, Head RD, Jacobs JM, Smith RD, Koppenaal DW, Brand JJ,

Pakrasi HB. 2015. Synechococcus elongatus UTEX 2973, a fast growing cyanobacterial

chassis for biosynthesis using light and CO_2. Sci Rep.

28.   Erb TJ, Zarzycki J. 2018. A short history of RubisCO: the rise and fall (?) of Nature's

predominant CO2 fixing enzyme. Curr Opin Biotechnol.

29.   Ensminger I, Busch F, Huner NPA. 2006. Photostasis and cold acclimation: Sensing low

temperature through photosynthesis. Physiol Plant.

30.   Welkie DG, Rubin BE, Diamond S, Hood RD, Savage DF, Golden SS. 2019. A Hard

Day's Night: Cyanobacteria in Diel Cycles. Trends Microbiol.

31.   Burns RA, MacDonald CD, McGinn PJ, Campbell DA. 2005. Inorganic carbon repletion

621   disrupts photosynthetic acclimation to low temperature in the cyanobacterium

622   Synechococcus elongatus. J Phycol.

623   32.   Sun Y, Wollman AJM, Huang F, Leake MC, Liu LN. 2019. Single-organelle

624   quantification reveals stoichiometric and structural variability of carboxysomes dependent

625   on the environment. Plant Cell.

626   33.   Sun Y, Casella S, Fang Y, Huang F, Faulkner M, Barrett S, Liu LN. 2016. Light

627   modulates the biosynthesis and organization of cyanobacterial carbon fixation machinery

628   through photosynthetic electron flow. Plant Physiol 171:530–541.

629   34.   Zhang P, Eisenhut M, Brandt AM, Carmel D, Silén HM, Vass I, Allahverdiyeva Y,

630   Salminen TA, Aro EM. 2012. Operon flv4-flv2 provides cyanobacterial photosystem ii

631   with flexibility of electron transfer. Plant Cell.

632   35.   Young JN, Goldman JAL, Kranz SA, Tortell PD, Morel FMM. 2015. Slow carboxylation

633   of Rubisco constrains the rate of carbon fixation during Antarctic phytoplankton blooms.

634   New Phytol.

635   36.   Itakura AK, Chan KX, Atkinson N, Pallesen L, Wang L, Reeves G, Patena W, Caspari O,

636   Roth R, Goodenough U, McCormick AJ, Griffiths H, Jonikas MC. 2019. A Rubisco-

637   binding protein is required for normal pyrenoid number and starch sheath morphology in

638   Chlamydomonas reinhardtii Proc Natl Acad Sci 116:18445 LP –

639   18454.

640   37.   Kysela DT, Randich AM, Caccamo PD, Brun Y V. 2016. Diversity Takes Shape:

641   Understanding the Mechanistic and Adaptive Basis of Bacterial Morphology. PLOS Biol

642   14:e1002565.

643 38. Jun S, Si F, Pugatch R, Scott M. 2018. Fundamental principles in bacterial physiology—

644   history, recent progress, and the future with focus on cell size control: a review. Reports

645   Prog Phys 81:056601.

646 39. Sargent MG. 1975. Control of cell length in Bacillus subtilis. J Bacteriol 123.

647 40. Steinberger RE, Allen AR, Hansma HG, Holden PA. 2002. Elongation correlates with

648   nutrient deprivation in Pseudomonas aeruginosa - Unsaturated biofilms. Microb Ecol

649   43:416–423.

650 41. Gonin M, Quardokus EM, O'Donnol D, Maddock J, Brun Y V. 2000. Regulation of stalk

651   elongation by phosphate in Caulobacter crescentus. J Bacteriol 182:337–347.

652 42. Rangarajan AA, Koropatkin NM, Biteen JS. 2020. Nutrient-dependent morphological

653   variability of Bacteroides thetaiotaomicron. Microbiology 166:624–628.

654 43. Galmés J, Hermida-Carrera C, Laanisto L, Niinemets Ü. 2016. A compendium of

655   temperature responses of Rubisco kinetic traits: variability among and within

656   photosynthetic groups and impacts on photosynthesis modeling. J Exp Bot 67:5067–5091.

657 44. Fernández-González C, Pérez-Lorenzo M, Pratt N, Moore CM, Bibby TS, Marañón E.

658   2020. Effects of Temperature and Nutrient Supply on Resource Allocation, Photosynthetic

659   Strategy, and Metabolic Rates of Synechococcus sp. J Phycol.

660 45. Falkowski PG, Sukenik A, Herzig R. 1989. Nitrogen limitation in Isochrysis Galbana

661   (Haptophyceae). II. Relative abundance of chloroplast proteins. J Phycol.

662 46. Raven JA, Johnston AM. 1991. Mechanisms of inorganic-carbon acquisition in marine

663   phytoplankton and their implications for the use of other resources. Limnol Oceanogr.

664 47. Losh JL, Morel FMM, Hopkinson BM. 2012. Modest increase in the C:N ratio of N-

665   limited phytoplankton in the California Current in response to high CO_2. Mar Ecol Prog

666   Ser.

667   48.   Losh JL, Young JN, Morel FMM. 2013. Rubisco is a small fraction of total protein in

668   marine phytoplankton. New Phytol.

669   49.   Galmés J, Aranjuelo I, Medrano H, Flexas J. 2013. Variation in Rubisco content and

670   activity under variable climatic factors. Photosynth Res.

671   50.   Cameron JC, Wilson SC, Bernstein SL, Kerfeld CA. 2013. Biogenesis of a Bacterial

672   Organelle: The Carboxysome Assembly Pathway. Cell 155:1131–1140.

673   51.   Oltrogge LM, Chaijarasphong T, Chen AW, Bolin ER, Marqusee S, Savage DF. 2020.

674   Multivalent interactions between CsoS2 and Rubisco mediate α-carboxysome formation.

675   Nat Struct Mol Biol 27:281–287.

676   52.   Wang H, Yan X, Aigner H, Bracher A, Nguyen ND, Hee WY, Long BM, Price GD, Hartl

677   FU, Hayer-Hartl M. 2019. Rubisco condensate formation by CcmM in beta-carboxysome

678   biogenesis. Nature2019/01/25. 566:131–135.

679   53.   Freeman Rosenzweig ES, Xu B, Kuhn Cuellar L, Martinez-Sanchez A, Schaffer M,

680   Strauss M, Cartwright HN, Ronceray P, Plitzko JM, Förster F, Wingreen NS, Engel BD,

681   Mackinder LCM, Jonikas MC. 2017. The Eukaryotic CO2-Concentrating Organelle Is

682   Liquid-like and Exhibits Dynamic Reorganization. Cell 171:148-162.e19.

683   54.   Horvath DJ, Li B, Casper T, Partida-Sanchez S, Hunstad DA, Hultgren SJ, Justice SS.

684   2011. Morphological plasticity promotes resistance to phagocyte killing of uropathogenic

685   Escherichia coli. Microbes Infect 13:426–437.

686   55.   Justice SS, Hung C, Theriot JA, Fletcher DA, Anderson GG, Footer MJ, Hultgren SJ.

687   2004. Differentiation and developmental pathways of uropathogenic Escherichia coli in

688   urinary tract pathogenesis. Proc Natl Acad Sci 101:1333–1338.

56. Kreuzer KN. 2013. DNA Damage Responses in Prokaryotes: Regulating Gene

Expression, Modulating Growth Patterns, and Manipulating Replication Forks. Cold

Spring Harb Perspect Biol 5.

57. Justice SS, Hunstad DA, Cegelski L, Hultgren SJ. 2008. Morphological plasticity as a

bacterial survival strategy. Nat Rev Microbiol 6:162–168.

58. Heinrich K, Leslie DJ, Morlock M, Bertilsson S, Jonas K. 2019. Molecular basis and

ecological relevance of Caulobacter cell filamentation in freshwater habitats. MBio 10.

59. Bos J, Zhang Q, Vyawahare S, Rogers E, Rosenberg SM, Austin RH. 2015. Emergence of

antibiotic resistance from multinucleated bacterial filaments. Proc Natl Acad Sci 112:178

LP – 183.

60. Wehrens M, Ershov D, Rozendaal R, Walker N, Schultz D, Kishony R, Levin PA, Tans

SJ. 2018. Size Laws and Division Ring Dynamics in Filamentous Escherichia coli cells.

Curr Biol 28:972-979.e5.

61. Yang DC, Blair KM, Salama NR. 2016. Staying in Shape: the Impact of Cell Shape on

Bacterial Survival in Diverse Environments. Microbiol Mol Biol Rev 80:187 LP – 203.

62. Wortinger MA, Quardokus EM, Brun Y V. 1998. Morphological adaptation and inhibition

of cell division during stationary phase in Caulobacter crescentus. Mol Microbiol 29:963–

973.

63. Pernthaler J, Zöllner E, Warnecke F, Jürgens K. 2004. Bloom of filamentous bacteria in a

mesotrophic lake: Identity and potential controlling mechanism. Appl Environ Microbiol

70:6272–6281.

64. Pernthaler J. 2005. Predation on prokaryotes in the water column and its ecological

implications. Nat Rev Microbiol 3:537–546.

65.   Hahn MW, Moore ERB, Höfle MG. 1999. Bacterial Filament Formation, a Defense

Mechanism against Flagellate Grazing, Is Growth Rate Controlled in Bacteria of Different

Phyla. Appl Environ Microbiol 65:25 LP – 35.

66.   Simek K, Vrba J, Pernthaler J, Posch T, Hartman P, Nedoma J, Psenner R. 1997.

Morphological and compositional shifts in an experimental bacterial community

influenced by protists with contrasting feeding modes. Appl Environ Microbiol 63:587 LP

– 595.

67.   Liao Y, Rust MJ. 2018. The Min Oscillator Defines Sites of Asymmetric Cell Division in

Cyanobacteria during Stress Recovery. Cell Syst.

68.   Broddrick JT, Welkie DG, Jallet D, Golden SS, Peers G, Palsson BO. 2019. Predicting the

metabolic capabilities of Synechococcus elongatus PCC 7942 adapted to different light

regimes. Metab Eng.

69.   Murayama Y, Kori H, Oshima C, Kondo T, Iwasaki H, Ito H. 2017. Low temperature

nullifies the circadian clock in cyanobacteria through Hopf bifurcation. Proc Natl Acad

Sci U S A.

70.   Muraleedharan S, Freitas C, Mann P, Glatter T, Ringgaard S. 2018. A cell length-

dependent transition in MinD-dynamics promotes a switch in division-site placement and

preservation of proliferating elongated Vibrio parahaemolyticus swarmer cells. Mol

Microbiol 109:365–384.

71.   MacCready JS, Vecchiarelli AG. 2018. In long bacterial cells, the Min system can act off-

center. Mol Microbiol 109:268–272.

72.   Raghunathan S, Chimthanawala A, Krishna S, Vecchiarelli AG, Badrinarayanan A. 2020.

Asymmetric chromosome segregation and cell division in DNA damage-induced bacterial

filaments. bioRxiv 2020.03.16.993485.

73.   Cayron J, Dedieu A, Lesterlin C. 2020. Bacterial filament division dynamics allows rapid

post-stress cell proliferation. bioRxiv 2020.03.16.993345.

74.   Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO. 2009.

Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat Methods.

75.   Clerico EM, Ditty JL, Golden SS. 2007. Specialized Techniques for Site-Directed

Mutagenesis in Cyanobacteria, p. 155–171. *In* . Humana Press.

76.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,

Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,

Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image

analysis. Nat Methods 9:676–682.

77.   Ducret A, Quardokus EM, Brun Y V. 2016. MicrobeJ, a tool for high throughput bacterial

cell detection and quantitative analysis. Nat Microbiol 1:1–7.

758   **Table 1**: Strains used in this study

Strain	Description	Source
Synechococcus elongatus PCC7942	Wild Type	-
Synechococcus elongatus PCC7942	ΔmcdA	(16)
Synechococcus elongatus PCC7942	ΔmcdB	(16)
Synechococcus elongatus PCC7942	ΔmcdAB	(16)
Synechococcus elongatus PCC7942	*rbcS-mTQ*	Wild Type transformed with pAH40
Synechococcus elongatus PCC7942	Δ*mcdA* + *rbcS-mTQ*	Δ*mcdA* transformed with pAH40
Synechococcus elongatus PCC7942	Δ*mcdB* + *rbcS-mTQ*	Δ*mcdB* transformed with pAH40
Synechococcus elongatus PCC7942	Δ*mcdAB* + *rbcS-mTQ*	Δ*mcdA*B transformed with pAH40

762
Figure 1. McdAB system mutants display fewer and mispositioned carboxysome aggregates at
40°C. (**A-D**) Microscopy images of the specified cell strains grown at 40°C in 2% CO_2. Phase
contrast (black), chlorophyll (red), and carboxysomes (green). Phase Contrast (PC) channel is
blue in the merge for better contrast. (**E**) Spacing between carboxysome foci as a function of cell
length. (**F**) Spacing between carboxysome foci in the same cell. (**G**) Carboxysome foci number
per cell. (**H**) Number of carboxysome foci per unit cell length for each strain. (**I**) Carboxysome
peak foci intensity for each cell strain. AU = Arbitrary Units. (**J**) Quantification of the maximum
length across carboxysome foci. (**F-J**) Solid bars represent the median and dashed lines
demarcate the 95% confidence interval. Statistical significance was based on a nonparametric
Mann-Whitney test. *** = P < 0.001, ** = P < 0.01, * = P < 0.05, n.s. = not significant. (**K**)
Population percentages of cells with the specified number of carboxysome foci. n ≥ 1000
carboxysomes from 440 cells of each strain.

775
776

777
Figure 2. Carboxysome mispositioning correlates with subtle changes in cell physiology at
40°C. (**A**) Cell lengths and (**B**) widths of WT population compared to McdAB system mutants.
Cells were grown at in 2% CO_2. n = 440 cells of each strain. Statistical significance was based on
a nonparametric Mann-Whitney test. *** = P < 0.001, ** = P < 0.01, * = P < 0.05, n.s. = not
significant. (**C**) Percentage of cells undergoing symmetric (mid-cell; light grey bar) or
asymmetric (non-mid-cell; dark grey bar) division. (**D**) Growth curves of all strains grown at
40°C in 2% CO_2. (**E**) Quantification of mean exponential growth rate at 40°C in 2% CO_2. (**F**)
Growth curves of all strains grown at 40°C in 0.4% CO_2. (**G**) Quantification of mean exponential
growth rate at 40°C in 0.4% CO_2. (**D-G**) Error bars represent the standard deviation from three
independent biological replicates. Statistical significance was based on an unpaired t-test. *** =
P < 0.001, ** = P < 0.01, * = P < 0.05, n.s. = not significant.

Figure 3. McdAB system mutants house few and mispositioned carboxysome aggregates at
30°C. (**A-D**) Microscopy images of the specified cell strains grown in 2% CO_2. Phase contrast
(black), chlorophyll (red), and carboxysomes (green). Phase Contrast (PC) channel is blue in the
merge for better contrast. (**E**) Spacing between carboxysome foci as a function of cell length. (**F**)
Spacing between carboxysome foci in the same cell. (**G**) Carboxysome foci number per cell. (**H**)
Number of carboxysome foci per unit cell length for each strain. (**I**) Carboxysome peak foci
intensity for each cell strain. Arbitrary Units (AU). (**J**) Quantification of the maximum length
across carboxysome foci. (**F-J**) Solid bars represent the median and dashed lines demarcate the
95% confidence interval. Statistical significance was based on a nonparametric Mann-Whitney
test. *** = $P < 0.001$, ** = $P < 0.01$, * = $P < 0.05$, n.s. = not significant. (**K**) Population
percentages of cells with the specified number of carboxysome foci. $n \geq 1000$ carboxysomes
from 440 cells of each strain.

Figure 4. McdAB mutants display cell elongation and asymmetric cell division phenotypes at
30°C. (**A**) Cell lengths and (**B**) widths of WT population compared to McdAB system mutants.
Cells were grown in 2% CO_2. n = 440 cells of each strain. Statistical significance was based on a
nonparametric Mann-Whitney test. *** = P < 0.001, ** = P < 0.01, * = P < 0.05, n.s. = not
significant. (**C**) Percentage of cells undergoing symmetric (mid-cell; light grey bar) or
asymmetric (non-mid-cell; dark grey bar) division. (**D**) Growth curves of all strains grown at
30°C in 2% CO_2. (**E**) Quantification of mean exponential growth rate at 30°C in 2% CO_2. (**F**)
Growth curves of all strains grown at 30°C in 0.4% CO_2. (**G**) Quantification of mean exponential
growth rate at 30°C in 0.4% CO_2. (**D-G**) Error bars represent the standard deviation from three
independent biological replicates. Statistical significance was based on an unpaired t-test. *** =
P < 0.001, ** = P < 0.01, * = P < 0.05, n.s. = not significant.

819
Figure 5. At 20°C wild-type and McdAB system mutants elongate, but only the mutants display
few and mispositioned carboxysome aggregates. (**A-D**) Microscopy images of the specified cell
strains grown in 2% CO_2. Phase contrast (black), chlorophyll (red), and carboxysomes (green).
Phase Contrast (PC) channel is blue in the merge for better contrast. (**E**) Spacing between
carboxysome foci as a function of cell length. (**F**) Spacing between carboxysome foci in the
same cell. (**G**) Carboxysome foci number per cell. (**H**) Number of carboxysome foci per unit cell
length for each strain. (**I**) Carboxysome peak foci intensity for each cell strain. AU = Arbitrary
Units. (**J**) Quantification of the maximum length across carboxysome foci. (**F-J**) Solid bars
represent the median and dashed lines demarcate the 95% confidence interval. Statistical
significance was based on a nonparametric Mann-Whitney test. *** = $P < 0.001$, ** = $P < 0.01$, *
= $P < 0.05$, n.s. = not significant. (**K**) Population percentages of cells with the specified number
of carboxysome foci. n ≥ 1000 carboxysomes from 440 cells of each strain.

837
Figure 6. At 20°C wild-type and McdAB mutants elongate and undergo asymmetric cell
division, but only mutants show slowed growth rates. (**A**) Cell lengths and (**B**) widths of WT
population compared to McdAB system mutants. Cells were grown in 2% CO_2. n = 440 cells of
each strain. Statistical significance was based on a nonparametric Mann-Whitney test. *** = P <
0.001, ** = P < 0.01, * = P < 0.05, n.s. = not significant. (**C**) Percentage of cells undergoing
symmetric (mid-cell; light grey bar) or asymmetric (non-mid-cell; dark grey bar) division. (**D**)
Growth curves of all strains grown at 20°C in 2% CO_2. (**E**) Quantification of mean exponential
growth rate at 20°C in 2% CO_2. (**F**) Growth curves of all strains grown at 20°C in 0.4% CO_2. (**G**)
Quantification of mean exponential growth rate at 20°C in 0.4% CO_2. (**D-G**) Error bars represent
the standard deviation from three independent biological replicates. Statistical significance was
based on an unpaired t-test. *** = P < 0.001, ** = P < 0.01, * = P < 0.05, n.s. = not significant.

Figure 7. McdAB mutants have decreased cellular levels of RuBisCO. (**A**) Western Blot
analysis of RbcL signal (top) and Atpß (bottom). Wild-type (lanes 1-3), Δ*mcdA* (lanes 4-6),
Δ*mcdB* (lanes 7-9), and Δ*mcdAB* (lanes 10-12), purified RbcL control (lane 13). (**B**)
Quantification of immunoblot signal for the cell type specified normalized against Atpβ signal.
Cells were grown at 30°C in 2% CO_2. Error bars represent the standard deviation from three
independent biological replicates. (**C**) Quantification of the average RbcS-mTQ intensity per
cell in each strain at the specified temperatures. (**D-F**) Quantification of the average RbcS-mTQ
intensity normalized to cell length in each strain at the specified temperatures. n ≥ 338 per cell
population. Statistical significance was based on an unpaired t-test. *** = P < 0.001, ** = P <
0.01, * = P < 0.05, n.s. = not significant.

879   **Supplemental Figures**
880

881
882   **Figure S1.** Presence of RbsC-mTQ carboxysome marker does not influence growth. (**A**) Growth
883   curves of wild-type *S. elongatus* with and without RbcS-mTQ carboxysome marker. Cells were
884   grown at 30°C in 2% CO_2. (**B**) Comparison of growth rates. Error bars represent the standard
885   deviation from three independent biological replicates. Unpaired t-test shows growth rates were
886   not statistically different.

887
888   **Figure S2.** Comparison of the peak RbcS-mTQ foci intensities for each strain across the three
889   growth temperatures tested in this study. AU = Arbitrary Units AU. n > 1000 carboxysomes per
890   cell-line at each temperature. Solid bars represent the median and dashed lines demarcate the
891   95% confidence interval. Statistical significance was based on a nonparametric Mann-Whitney
892   test. *** = P < 0.001, ** = P < 0.01, * = P < 0.05, n.s. = not significant.
893