Spike-frequency dependent inhibition and excitation of neural activity by high-frequency ultrasound

Martin Loynaz Prieto[1], Kamyar Firouzi[2], Butrus T. Khuri-Yakub[2], Daniel V. Madison[1], and Merritt Maduke[1]

[1]Department of Molecular and Cellular Physiology, and [2]E. L. Ginzton Laboratory, Stanford University

Condensed title: Spike-Rate-Dependent Neuromodulation by Ultrasound [50 characters or less]

SUMMARY [40 words or less] Prieto et al. describe how ultrasound can either inhibit or potentiate action potential firing in hippocampal pyramidal neurons and demonstrate that these effects can be explained by increased potassium conductance.

1 **ABSTRACT [300 words or less]**
2
Ultrasound can modulate action-potential firing *in vivo* and *in vitro*, but the mechanistic basis of

this phenomenon is not well understood. To address this problem, we used patch-clamp

recording to quantify the effects of focused, high-frequency (43 MHz) ultrasound on evoked

action potential firing in CA1 pyramidal neurons in acute rodent hippocampal brain slices. We

find that ultrasound can either inhibit or potentiate firing in a spike-frequency-dependent

manner: at low (near-threshold) input currents and low firing frequencies, ultrasound inhibits

firing, while at higher input currents and higher firing frequencies, ultrasound potentiates firing.

The net result of these two competing effects is that ultrasound increases the threshold current

for action potential firing, the slope of frequency-input curves, and the maximum firing

frequency. In addition, ultrasound slightly hyperpolarizes the resting membrane potential,

decreases action potential width, and increases the depth of the afterhyperpolarization. All of

these results can be explained by the hypothesis that ultrasound activates a sustained

potassium conductance. According to this hypothesis, increased outward potassium currents

hyperpolarize the resting membrane potential and inhibit firing at near-threshold input

currents, but potentiate firing in response to higher input currents by limiting inactivation of

voltage-dependent sodium channels during the action potential. This latter effect is a

consequence of faster action-potential repolarization, which limits inactivation of voltage-

dependent sodium channels, and deeper (more negative) afterhyperpolarization, which

increases the rate of recovery from inactivation. Based on these results we propose that

ultrasound activates thermosensitive and mechanosensitive two-pore-domain potassium (K2P)

channels, through heating or mechanical effects of acoustic radiation force. Finite-element

modelling of the effects of ultrasound on brain tissue suggests that the effects of ultrasound on

firing frequency are caused by a small (less than 2°C) increase in temperature, with possible

additional contributions from mechanical effects

INTRODUCTION

Ultrasound can non-invasively modulate action potential activity in neurons *in vivo* and *in vitro*,

with improved depth penetration and spatial resolution relative to other non-invasive

neuromodulation modalities, and it may therefore become an important new technology in

basic and clinical neuroscience (Fry et al., 1958; Gavrilov et al., 1996; Tufail et al., 2010;

Bystritsky et al., 2011; Fomenko et al., 2018; Blackmore et al., 2019). Investigation of this

phenomenon has predominantly focused on low-frequency ultrasound (defined here as less

than 3 MHz, although there is no firmly defined boundary between "high" and "low" frequency

in the neuromodulation field), but higher ultrasound frequencies have also been shown to

modulate action potential firing *in vitro* (Menz et al., 2013; Menz et al., 2019) and to directly

modulate ion channel activity in heterologous systems (Kubanek et al., 2016; Prieto et al.,

2018). The focus on lower frequency ultrasound is understandable, since envisioned clinical

applications involving transcranial focused ultrasound have been a primary motivation for

research on ultrasound neuromodulation, and loss of ultrasound power due to attenuation in

the skull limits these applications to low frequency ultrasound. For applications in which

transmission through the skull does not impose limits on frequency, such as *in vitro* studies,

neuromodulation in the peripheral nervous system (Downs et al., 2018; Cotero et al., 2019;

Zachs et al., 2019), neuromodulation using subcranial implants, or neuromodulation in

experimental animal model systems involving craniotomies or acoustically transparent cranial

windows, high frequencies have a distinct advantage in terms of the greater spatial resolution

that can be achieved. Even for *in vivo* applications in human subjects, however, the spatial

resolutions that can be achieved with low-frequency, transcranial ultrasound neuromodulation

are on the order of millimeters, making ultrasound neuromodulation superior in this respect to

other, more established forms of non-invasive brain stimulation (Tyler et al., 2018).

These applications motivate investigation of the fundamental physical, cellular, and

molecular mechanisms underlying neuromodulation, which are all not well understood for

either high- or low-frequency ultrasound. It remains an open question to what extent these

mechanisms overlap for high- and low-frequency ultrasound neuromodulation. In terms of the

basic physical mechanism by which acoustic energy is transduced into effects on biological

tissue, most proposed mechanisms for ultrasound neuromodulation involve either heating due

to absorption of acoustic energy (Hand, 1998), mechanical effects of acoustic radiation force

(Duck, 1998; Sarvazyan et al., 2010), or effects of cavitation (the nucleation, growth, oscillation,

and sometimes collapse, of microscopic gas bubbles) (Leighton, 1998; Wu and Nyborg, 2008;

Krasovitski et al., 2011; Plaksin et al., 2016). Of these, the first two increase with acoustic

frequency, while the probability of cavitation decreases with acoustic frequency. There are also

many unanswered questions at the cellular level. Both excitatory and inhibitory effects of

ultrasound have been observed using direct or indirect measures of neural activity at the

population level (Bystritsky et al., 2011; Blackmore et al., 2019), but it is unclear whether the

direct effect of ultrasound at the cellular level is excitatory or inhibitory. Of course, the answer

to this question could depend on any number of possible relevant biological and experimental

variables, such as species, tissue, specific neural subtype, ultrasound stimulus parameters, or

whether effects on intrinsic or evoked activity are measured. For example, a cellular-level

excitatory effect, specific to GABAergic interneurons, could produce an inhibitory effect at the

population level. This leads to the question of whether certain subpopulations of neurons are

more sensitive to ultrasound than others, and if so, what are the specific molecular

mechanisms underlying the differences in sensitivity. Do certain ion channels respond directly

to ultrasound? What biophysical properties might account for the sensitivity of these channels

to ultrasound, and how might cell-type specific differences in the density and localization of

these channels, and the way in which they interact with other ion channels to regulate action

potential firing, produce differences in the response to ultrasound?

One reason there are so many outstanding questions regarding ultrasound

neuromodulation is that patch-clamp recordings of the effects of ultrasound on action-potential

firing in neurons have been unavailable. At low ultrasound frequencies, we (Prieto et al., 2018)

and others (Tyler et al., 2008) have found that patch-clamp seals are extremely unstable in the

presence of ultrasound at low frequencies, precluding detailed, mechanistic studies of

ultrasound neuromodulation with this technique, which provides quantitative information on

action-potential timing and dynamics unobtainable with other techniques. However, we have

previously shown that stable patch-clamp recordings can be achieved using ultrasound at the

frequency of 43 MHz (Prieto et al., 2018). Here, we use patch-clamp recording to measure the

effects of ultrasound at 43 MHz and 50 W/cm^2 on action-potential firing in response to injected

current in pyramidal neurons of the CA1 layer of the hippocampus in acute rodent brain slices.

We find that ultrasound has a bidirectional, spike-frequency dependent effect on excitability,

and that this and other observed neurophysiological effects of ultrasound can be explained by

activation by ultrasound of a steady K$^+$ current, resembling that of two pore domain potassium

channels (K2P channels).

MATERIALS AND METHODS

Slice preparation. Brain slices were prepared from male Sprague-Dawley rats, 35-50 days old.

Rats were anesthetized with isofluorane and decapitated, and the brain was immediately

removed and placed in ice-cold artificial cerebral spinal fluid (ACSF), bubbled with 95% O_2 and

5% CO_2. The hippocampus was dissected out and placed on the slicing apparatus, consisting of a

manual micrometer and a gravity-driven vertical slicing mechanism, with the CA1 layer oriented

approximately parallel to the slicing blade. Slices (500 microns thick) were prepared and then

stored in a humidified chamber with an atmosphere of 95% O_2/5% CO_2, resting on a square of

filter paper placed on a dish of ACSF. Slices were used within 1-6 hours of slice preparation.

Animals were handled in accordance with protocols approved by Stanford University's

Institutional Animal Care and Use Committee.

Ultrasound. Continuous wave ultrasound at 43 MHz and 50 W/cm^2 was applied to brain slices

using a set-up similar to that we previously used for our experiments on cultured cells (Prieto et

al., 2018), except that the tissue was visualized with a dissecting microscope at low

magnification. The bottom of the experimental chamber was a 25-micron film of polystyrene,

plasma-treated with a Harrick plasma cleaner (Harrick Plasma, Ithaca, NY). Ultrasound was

transmitted from below (through the polystyrene film) with the sound beam perpendicular to

the bottom of the chamber. The 43 MHz transducer was a custom-built device, calibrated as

described previously (Prieto et al., 2013), excited using an ENI 403LA (37 dB) amplifier (ENI,

Rochester, NY). The focal volume of the transducer is approximately a cylinder 90 microns in

diameter by 500 microns long, and the focal distance is approximately 4.2 mm. The set-up was

based on the stage from a Zeiss Axioskop-2 microscope (Zeiss Microscopes, Jena, Germany),

with the housing for the sub-stage condenser modified to accommodate the transducer, such

that the position of the transducer could be adjusted using the controls for alignment of the

condenser, and the position of the tissue sample relative to the transducer could be adjusted

with the microscope stage. The transducer was coupled to the polystyrene film at the bottom

of the experimental chamber using a small volume of distilled water held in place by a rubber

O-ring attached to the tip of the transducer with silicone grease. The focal volume of the

transducer was aligned along the z-axis using a pulse-echo protocol, adjusting the height of the

transducer to maximize the echo signal from the bottom of the empty chamber. The focus was

aligned in the x-y plane by adding to the chamber a small volume of ACSF, barely sufficient to

cover the bottom of the chamber, such that a thin layer of solution was spread over the bottom

of the chamber. Ultrasound pulses, one second in length, were then applied, raising a mound of

fluid at the focus of the transducer (due to the radiation force produced by reflection of the

acoustic wave at the interface between the solution and the air above it (Duck, 1998)). The

mound of fluid was then aligned in the x-y plane to the center of a reticle in one eyepiece of the

dissecting microscope, and, after adding additional ACSF and the tissue sample to the chamber,

the center of the reticle was aligned with the region of the tissue targeted for patch-clamp

recording. The ultrasound intensity (50 W/cm^2) is the spatial peak, pulse average intensity for

the free field. The interval between ultrasound applications was at least 12 seconds.

Electrophysiology. Current clamp recordings were performed using an Axon Instruments

Axoclamp-2B amplifier operating in "Bridge" mode and Digidata 1330A digitizer with pClamp

software (Molecular Devices, Sunnyvale, CA), except for the preliminary experiments in

Supplemental Figure 1, which were performed with an Axon Instruments Axopatch 200B

amplifier (Molecular Devices, Sunnyvale, CA). Patch-clamp recording was performed using a

"blind-patch" approach (Blanton et al., 1989; Malinow and Tsien, 1990; Castaneda-Castellanos

et al., 2006), in which the recording pipette was positioned above the CA1 layer of the

hippocampus, as identified visually at low magnification, and then slowly lowered into the

tissue while applying positive pressure to the pipette and monitoring the pipette tip resistance

in voltage-clamp mode. In the blind-patch approach, a small decrease in tip resistance is used to

indicate possible contact of the pipette tip with a neuron in the absence of the usual visual

cues. Typically, the first two instances of possible cell contact were not used, to avoid patching

on cells at the surface of the tissue that may have been damaged during the slicing procedure.

Gigaseals and the subsequent whole-cell recording configuration were obtained following the

standard procedure in voltage-clamp mode before switching to current-clamp mode. In most

experiments, slices were held in place with a Warner Instruments RC-22 slice anchor (Harvard

Bioscience, Hamden, CT); the experiments in Supplemental Figure 1 and some of the

experiments in Figure 1E were performed without a slice anchor. No obvious effects of the slice

anchor on the ultrasound response were noted. Series resistance, monitored and compensated

throughout the recording, was between 30 and 100 MΩ. All of the neurons used for

experiments could be unambiguously identified as pyramidal cells by their distinct adaptive

firing patterns in response to 2-s current steps. Current records were low-pass filtered at 10 kHz

and sampled at 100 kHz. Brain slices were continuously perfused with ACSF (in mM: 119 NaCl,

2.5 KCl, 1.3 MgSO$_4$, 2.5 CaCl$_2$, 1 NaH$_2$PO$_4$, 26.2 NaHCO$_3$, 11 glucose), bubbled with 95% O$_2$/5%

CO_2, at ~100-250 mL/hour. The internal solution was (in mM): 126 K-gluconate, 4 KCl, 10 HEPES,

4 Mg-ATP, 0.3 Na_2GTP, 10 Na-phosphocreatine, 10 sucrose, and 50 U/mL creatine

phosphokinase (porcine), pH 7.2 (KOH). This internal solution contains an ATP-regenerating

system (phosphocreatine and creatine phosphokinase) because we found that the strength of

the response to ultrasound was unstable, gradually declining over the course of a recording

unless the ATP-regenerating system was included (Supplemental Figure 2). Na-phosphocreatine

was obtained from Abcam and creatine phosphokinase was obtained from EMD Millipore. All

other salts and chemicals were obtained from either Sigma-Aldrich or Fisher Scientific. (For

some of the preliminary recordings shown in Figure 1E, the creatine phosphokinase was

omitted from the internal solution, or a different internal solution, containing 120 K-gluconate,

40 HEPES, 5 $MgCl_2$, 0.3 Na_2GTP, 2 Na_2ATP, pH 7.2 (KOH) was used. Creatine phosphokinase was

also omitted for the experiments in Supplemental Figure 4). Other than the reduction of the

ultrasound response over time in the absence of creatine phosphokinase, no obvious

differences in recordings with different internal solutions were noted.) Because creatine

phosphokinase increases the viscosity of the solution, making it difficult to obtain gigaseals, a

small volume of internal solution without the enzyme was added to the tip of the pipette

(enough to fill approximately the first 3 mm of the tip) before back-filling the pipette with the

enzyme-containing solution. Pipettes were pulled from thick walled glass and had resistances

between 5 and 10 MΩ. Recordings were performed at room temperature (21-23°C), except for

the experiments in Supplemental Figure 4, which were performed at near physiological

temperature (30°C). For these experiments, the temperature of the external solution was

regulated and monitored with a Warner Instruments CL-100 bipolar temperature controller

equipped with a SC-20 in-line heater/cooler and a thermistor (Warner Instruments, Hamden,

CT). The external solution was heated to 35-37°C while being bubbled with 95% O_2/5% CO_2 and

then passed through the heater/cooler and cooled to achieve the target temperature of 30°C in

the bath solution. (The external solution was cooled rather than heated to avoid loss of oxygen

tension and formation of gas bubbles due to heating of oxygen-saturated solution.)

Data analysis. Current records were analyzed in Igor Pro (Wavemetrics, Lake Oswego, OR) with

user-written procedures. Action-potential threshold was defined as the point at which the first

derivative of the voltage reached 4% of its peak value during the rising phase of the action

potential. This quantitative criterion was previously found to correspond with action-potential

thresholds as identified visually (Khaliq and Bean, 2010; Yamada-Hanff and Bean, 2015), and we

found that it also works well with our data, using phase plots to visually confirm the threshold

value. Action-potential height was defined as the difference between the action-potential peak

and the action-potential threshold voltage. Action-potential width was measured at 50% of

action potential height defined in this manner. Threshold current levels for action-potential

firing were estimated based on a series of current steps in 10-pA increments. Frequency-input

plots and action-potential parameters (height, width, latency, and interspike intervals) were

determined from the average values of at least 3 trials each for the control and ultrasound

conditions. Frequency-input trials were performed alternatingly for the control and ultrasound

conditions, with the first condition tested varying randomly on a cell-by-cell basis.

Average traces for analysis of the effects of ultrasound on membrane resting potential and

membrane capacitance were derived from at least 3 voltage traces.

Statistical significance was assessed using paired or unpaired two-tailed Student's t-tests, with P

< 0.05 defined as significant. Statistical analysis was performed in Microsoft Excel.

Finite-element simulations. Finite-element were performed in COMSOL (COMSOL Inc., Palo

Alto, CA, USA). The simulation domain had radially symmetric geometry and was 6 mm in the

axial direction. The simulation domain contained four layers of different materials: a lower layer

of water (4.2 mm thick in the axial direction), followed by a layer of polystyrene (25 microns

thick), followed by a layer of brain tissue (500 microns thick), followed by an upper layer of

water (1.275 mm thick). The width of the simulation domain in the axial direction was 1 mm

(for simulation for acoustic pressure and heating) or 5 mm (for simulation of mechanical

deformation) in the axial direction. A 940-micron diameter by 100-micron height arc on the

lower axial boundary of the simulation domain represented the quartz lens of the transducer.

Simulations of acoustic pressure, heating, and static displacement in response to

radiaton force were performed as described previously (Prieto et al., 2018). For simulation of

dynamic tissue displacement in response to radiation force, the brain slice was modeled as a

incompressible, linear viscoelastic material (Calhoun et al., 2019), characterized by Young's

modulus, Poisson's ratio, and shear viscosity, loaded by the fluid layer above it. The polystyrene

was modeled as a linear elastic material, because we determined in a series of simulations that

including viscosity of the polystyrene had no effect on the tissue displacement. A time step of

0.1 ms was used for simulation of the dynamic tissue displacement. Material properties used

for water, polystyrene, and brain tissue used in the simulation and sources for these values are

given in Table 1. Additional details on mesh size, boundary conditions, and solver configurations

are available in Prieto et al. (2018).

SUMMARY OF SUPPLEMENTAL MATERIAL

Supplemental materials include four figures showing: the effect of ultrasound at different

intensities on action potential firing frequency (Supplemental Figure 1); stabilization of the

response to ultrasound by an ATP-regenerating system in the internal solution (Supplemental

Figure 2); effect of ultrasound on action potential height (Supplemental Figure 3); and the

effects of ultrasound on action potential firing and waveform at near physiological temperature

(30°C) (Supplemental Figure 4).

RESULTS

We measured the effects of ultrasound on action-potential firing in hippocampal CA1 pyramidal

cells using the set-up shown in Figure 1A (described in *Materials and Methods*). Throughout the

experiments reported here, ultrasound was applied at 43 MHz and 50 W/cm^2 as a 1-s,

continuous-wave pulse. In an initial exploration of the effects of intensity on the ultrasound

response, we determined that 50 W/cm^2 had a sufficiently robust effect on firing frequency to

permit quantitative investigation of this effect (Supplemental Figure 1), but we did not perform

a detailed investigation of the intensity dependence. We chose to use continuous-wave

ultrasound (without additional low-frequency modulation within the pulse) because

continuous-wave ultrasound was previously found to be optimal for ultrasound

neuromodulation of retinal ganglion cells at 43 MHz (Menz et al., 2013). With these ultrasound

parameters, we found robust, reproducible inhibition of action-potential firing by ultrasound

using the protocol illustrated in Figure 1B. In these experiments, a current-injection amplitude

sufficient to induce firing at an average frequency of ~4-12 Hz during the first 500 ms of the

current step (corresponding to the overlap between the ultrasound stimulus and current step)

was used. This range of firing frequencies is physiologically relevant and sufficient to detect

either inhibition or potentiation of firing. With these experimental conditions, we established

that the response to ultrasound is highly reproducible, both on a trial-by-trial basis within the

same cell (Figure 1C-D), and between cells, with similar effects seen in over fifty cells (Figure

1E). In addition, the response to ultrasound was stable over the course of recordings lasting

over 30 minutes (Figure 1F, Supplemental Figure 2); in a few cases where the patch seal lasted

for over 90 minutes, the ultrasound response remained stable.

Effects of ultrasound on frequency-input curves. To explore the effects of ultrasound on

excitability over a wider range of firing frequencies, we generated frequency-input (f-i) curves

comparing average firing frequencies as a function of input current in the presence and

absence of ultrasound. An example f-i curve generated with the protocol illustrated in Figure 1B

is shown in Figure 2A, along with example voltage traces in Figure 2B. The average spike

frequency during the first 500 ms of the current step was compared to the spike frequency in

the same time window in the absence of ultrasound. To compare the effects of ultrasound

across neurons, we converted the f-i curves into plots of the relative increase or decrease in

firing frequency as a function of the input current (Figure 2C-D). These data reveal two distinct

regimes with contrasting inhibitory and excitatory ultrasound effects. At relatively low input

currents, near the threshold for action-potential firing under this current stimulation protocol,

ultrasound decreases the average firing frequency; while at relatively high input currents, well

above the action-potential threshold, ultrasound increases the average firing frequency.

Between these two regimes, there is a transitional region where there is little or no effect on

average firing frequency, presumably due to the balance between competing inhibitory and

excitatory effects. Other notable effects of ultrasound on the f-i curves are an increase in the

threshold current for action-potential firing, an increase in the slope of the f-i relationship in

the approximately linear region of the f-i curve, and an increase in the maximum firing

frequency in the sublinear "plateau" region of the curve (Figure 2A). The mean (± SEM) slope of

linear region of the f-i curve increased from 0.108 ± 0.007 Hz/pA in the control condition to

0.145 ± 0.012 Hz/pA in the ultrasound condition; and the mean maximum firing frequency

increased from 23 ± 1 Hz in the control condition to 30 ± 2 Hz/pA in the ultrasound condition (N

= 9).

254   Action-potential firing behavior is determined by the interaction between numerous K^+,

Na^+, and other ionic currents (Madison and Nicoll, 1984; Bean, 2007). Some of these currents

are clearly identified with specific ion channel subtypes, while the molecular identity of others

is still uncertain. Thus, f-i curves are a complicated function of the density, localization,

conductance, and kinetic properties of these channels/currents. Some currents inactivate

relatively rapidly and only influence firing frequency during the initial response to a sustained

depolarizing current step, while others show slow, voltage-dependent activation, and only

influence firing frequency late in a current step; still other currents can influence firing

frequency throughout a sustained depolarization. To explore the molecular basis of the

response to ultrasound, we therefore generated a second set of f-i curves with ultrasound

applied 1 s after the start of a 3-s current step (Figure 3A).

Ultrasound also had a bidirectional, spike-frequency-dependent effect on excitability

when it was applied 1 second after the start of a current step (Figure 3). Again, ultrasound

decreased firing frequency in the low-firing-frequency, near-threshold region of the f-i curve,

and increased spike rate in the high-firing-frequency, suprathreshold region of the curve (Figure

3B-D). However, the excitatory effect was more pronounced than we observed when the

ultrasound pulse started 500 ms before the start of the current step. Here, ultrasound

potentiated firing frequency by several hundred percent for high input currents (Figure 3D), as

compared with a maximum potentiation of 49 ± 16% at 450 pA seen with the earlier ultrasound

application (Figure 2C). This reflects the fact that in response to prolonged injection of high

amplitude currents, accumulation of voltage-gated Na$^+$ channel (Na$_V$ channel) inactivation can

drive pyramidal cells into a refractory state where spiking is infrequent and irregular or entirely

absent (for example, the voltage trace for the control condition at +450 pA in Figure 3C shows

an initial steep decline in action potential height, followed by a gradual partial recovery of

action potential height, followed by a period of no action potential activity). If neurons are in

this refractory state during the ultrasound application, ultrasound can "rescue" firing (as in the

example voltage trace for the ultrasound condition at +450 pA in Figure 3C). This refractory

state probably does not occur under normal physiological conditions, but the ability of

ultrasound to rescue action-potential firing under these conditions still provides an important

clue as to the molecular mechanisms underlying the effects of ultrasound on firing frequency,

as discussed further below. A hint of this rescue phenomenon is also seen when the ultrasound

application starts before the current step, as seen in the abrupt increase in the potentiation

effect at +450 pA (Figure 2C).

Effects of ultrasound on interspike intervals. To examine the effects of ultrasound on action-

potential firing in more detail we compared the latency to the first spike, and the intervals

between subsequent spikes (interspike intervals), in the control and ultrasound conditions

(Figure 4). To summarize these results, and to account for the variability in intrinsic excitability

between cells, we averaged *instantaneous* firing frequencies (latency and interspike intervals)

across cells firing at approximately the same *average* firing frequency (either 5, 10, or 20 Hz) in

the control condition at whatever input current was necessary to achieve these average firing

frequencies, and at the same input current in the ultrasound condition (Figure 4A-C). We note,

however, that this averaging procedure can obscure some of the details of ultrasound's effects.

At 5 Hz the effect of ultrasound applied 500 ms before the start of the current step is

predominantly inhibitory (as seen in the longer average latency and interspike intervals in the

ultrasound as compared with the control condition (Figure 4A), but the interval between the

first and second spikes was actually shorter in the ultrasound condition than the control

condition in some cells (6 out of 13 cells in this data set). This effect occurs because, even at

relatively low average firing frequencies, pyramidal cells will occasionally fire "doublets" or

high-frequency bursts of two action potentials, in which a second action potential is triggered

by the after-depolarization of the initial action potential. When this occurs, ultrasound

decreases the interval between spikes (Figure 4D). This result indicates that the mechanism by

which ultrasound potentiates firing at high average firing frequencies is also active under

conditions of low overall average firing frequency, during localized periods of high-frequency

firing. A similar combination of inhibitory and excitatory effects can be observed at 10 Hz and

even 20 Hz (Figure 4B-C). At 20 Hz, ultrasound still increased the latency to the first spike (16 ±

2 ms versus 12 ± 1 ms in the control condition; N = 12, P = 5.2 x 10^{-4}, paired, two-tailed

Student's t-test), despite decreasing the interspike interval for all subsequent spikes (Figure 4C).

The effect of ultrasound was also mixed when ultrasound was applied 1 s after the start of the

current step (Figure 4E).

Effects of ultrasound on resting membrane potential. Ultrasound also has effects on resting

membrane potential, which can be observed by averaging several voltage traces aligned to the

onset of the ultrasound pulse, in the absence of injected current. As shown in Figure 5A,

ultrasound has a slight hyperpolarizing effect on resting membrane potential. The average

voltage traces also show another interesting effect of ultrasound. In addition to the relatively

constant hyperpolarization, there is a transient *depolarization* of the resting membrane

potential, preceding the hyperpolarization effect and acting on a faster time scale, at the onset

of the ultrasound pulse; this transient depolarization is matched by a roughly symmetrical

transient hyperpolarization at the offset of the pulse (Figure 5A, *arrows*). The symmetrical,

on/off nature of these transients suggests that they are caused by changes in membrane

capacitance, as does the fact that they occur much faster than the steady-state changes in

resting membrane potential (Figure 5A-C). (Changes in membrane potential due to changes in

capacitance can occur much faster than those due to ionic currents because they do not involve

actual redistribution of charges across the membrane and are therefore not limited by

membrane conductance). Thus, the results in Figure 5 can be described by three distinct steps:

1) ultrasound rapidly increases membrane capacitance (C) at the onset of the ultrasound pulse,

which causes the membrane voltage (V) to become less negative (due to an increase in the

denominator in the equation $V=Q/C$, where Q is the negative total charge on the membrane);

2) ionic currents then slowly change the membrane voltage to a steady-state value determined

by the total ionic current (one or more ion channels having been affected by ultrasound)

resulting in membrane hyperpolarization; 3) at the offset of the ultrasound pulse, capacitance

rapidly relaxes back to its initial value, producing a transient decrease in membrane voltage,

through essentially the same mechanism as in step 1. We investigate the physical basis of these

capacitance changes and their relationship to the effects of ultrasound on excitability further

below, but one point worth mentioning here is that the change in capacitance is too small for

its effect on the rate of membrane charging (less than 1% change in membrane time constant)

to have a significant effect on excitability in and of itself.

The K2P channel hypothesis. What ion channels might be responsible for the effects of

ultrasound on excitability and resting membrane potential? A compelling hypothesis—able to

explain all of our data—is that ultrasound activates a fast-activating, non-inactivating potassium

channel, such as members of the K2P potassium channel family. While commonly described as

"voltage-independent", K2P channels (with the exception of TWIK-1 channels) actually have an

outwardly-rectifying, voltage-dependent open probability under physiological K^+ gradients due

to the interaction of permeant ions in the selectivity filter with an activation gate (Schewe et

al., 2016). Nonetheless, their rate of activation is fast (millisecond time-scale) and voltage-

independent, such that they are functionally similar to truly voltage-independent channels with

outwardly rectifying single-channel conductance. A primary reason for suspecting K2P channels

is that the effects of ultrasound are similar regardless of whether ultrasound is presented 500

ms before or 1 s after the start of the current step (compare Figures 2 and 3), consistent with

the idea that ultrasound affects a channel that does not undergo prolonged voltage-dependent

inactivation during sustained depolarizations. Related to this point, the effects of ultrasound

are not diminished by repetitive, high-frequency action potential firing (for example, in Figure

4C effects of ultrasound are clearly present throughout the entire 20-Hz, 10-spike train).

Further, a striking feature of the effects of ultrasound on spike intervals is that ultrasound

always increases the latency to the first spike (Figure 4A-C), regardless of the input current and

the rate of approach to the initial action potential threshold, consistent with the idea that

ultrasound affects a fast-activating, non-inactivating channel.

CA1 pyramidal neurons express a variety of K2P channel subunits. Expression of TASK-1

and TASK-3, TREK-1 and TREK-2, TRAAK, and TWIK has been shown at the mRNA level (Talley et

al., 2001), while expression at the protein level has been shown has been shown for TASK-3 in

CA1 pyramidal neurons specifically (Marinc et al., 2014), and for TRAAK throughout the central

nervous system (Brohawn et al., 2019). In addition, functional expression of TASK-like currents

has been shown in CA1 pyramidal neurons using patch-clamp recording (Taverna et al., 2005).

TREK and TRAAK channels are particularly interesting in the present context since they are

exceptionally sensitive to mechanical force and to increases in temperature between

approximately 20 and 40°C (Maingret et al., 2000; Kang et al., 2005). Thus, TREK and TRAAK

channels are responsive to the two leading candidate mechanisms by which ultrasound at 43

MHz could modulate ion channel activity.

Activation of K2P channels by ultrasound could account for all of the neurophysiological

effects of ultrasound described so far: hyperpolarization of resting membrane potential;

inhibition of action-potential firing in response to near-threshold current injections; and—

although this last point may seem counter-intuitive—potentiation of action potential firing at

high firing frequencies. Hyperpolarization of resting membrane potential by increased outward

K^+ current is straightforward, as is the idea that K^+ current can inhibit firing, but K^+ current can

also potentiate firing by its effects on action-potential repolarization and afterhyperpolarization

(AHP). By accelerating the rate of membrane repolarization following the peak of an action

potential (thereby reducing action potential width) K^+ current can reduce inactivation of

voltage-dependent Na^+ channels during the action potential, and by increasing the depth of the

AHP, it can accelerate the voltage-dependent recovery of Na_V channels from inactivation. Both

of these effects would tend to increase the population of Na_V channels available for activation

in response to depolarizing current and would increase the maximum action potential firing

frequency, as we in fact see in response to ultrasound (Figures 2 and 3). This mechanism is well-

known and widespread in neurophysiology, with several K^+ channels, including both K2P

channels and voltage-dependent K^+ channels (K_V channels), having been shown to facilitate

high-frequency firing (Lien and Jonas, 2003; Brickley et al., 2007; Gu et al., 2007; Gonzalez et al.,

2009; Liu and Bean, 2014; Kanda et al., 2019). The idea that the potentiation of firing by

ultrasound is due to effects on action-potential repolarization and AHP is supported by the

results in Figure 4C. For neurons firing at an average firing frequency of 20 Hz in the absence of

ultrasound, ultrasound increases the latency to the first spike, while it decreases the intervals

between all subsequent spikes. The lack of a potentiating effect on the first spike, despite the

otherwise strongly potentiating effects of ultrasound, indicates that the potentiating effect acts

through a process (such as action-potential repolarization and AHP) that occurs *after* the

initiation of the first action potential. The idea that ultrasound can potentiate firing by

activating K^+ current makes specific predictions about the effects of ultrasound on action

potential waveform: ultrasound should accelerate repolarization, decrease action potential

width, and increase the depth of the AHP.

Effects of ultrasound on action-potential waveform. To test the idea that potentiation of firing

by ultrasound is due to activation of K^+ channels, we examined the effects of ultrasound on

action-potential waveform in our recordings. The effects of ultrasound on action-potential

waveform are consistent with the idea that ultrasound facilitates high-frequency firing by

accelerating action-potential repolarization. Figure 6A-F shows the effect of ultrasound applied

500 ms before the start of the current step on action-potential width for cells firing at average

frequencies of 5, 10, and 20 Hz. (Ultrasound also had effects on action-potential height,

although these were less pronounced than the effects on width; effects on height are detailed

in Supplemental Figure 3.) Ultrasound decreased action-potential width for every action

potential at all firing frequencies. As shown in Figure 6G-H, ultrasound also decreased action-

potential width when applied 1 s after the start of a current step, again indicating that the

channels responsible for these effects continue to influence firing frequency and remain

responsive to ultrasound throughout sustained depolarizing current steps.

The effects of ultrasound on action-potential width tended to counteract the

broadening of action-potential width that occurs during high-frequency firing. Figure 7 plots

action potential width as a function of action potential number and input current for the

control and ultrasound conditions. In the control condition, there are dramatic differences in

width between the first action potential and subsequent action potentials at high input

currents, while in the ultrasound condition these differences are much less pronounced. To

quantify this effect, we measured the difference in width between the first and third, first and

fifth, and first and last action potentials during the ultrasound stimulus, in response to a +450

pA current step for the control and ultrasound conditions. These differences (mean ± SEM,

control vs. ultrasound, N = 9) were 2.6 ± 0.9 vs. 0.7 ± 0.1 ms, 1.5 ± 0.1 vs. 0.5 ± 0.1 ms, and 1.2 ±

0.2 vs 0.5 ± 0.1 ms for the third, fifth, and last action potentials (P = 0.064, 1.4×10^{-5}, and

0.0013, paired, two-tailed Student's t-test). One plausible interpretation of this result is that

K2P channels activated by ultrasound cause the membrane voltage during the action potential

to repolarize before slower-activating K_V channels, which would otherwise contribute to action-

potential repolarization, are activated. Since time-dependent activation and inactivation of K_V

channels causes action-potential broadening during repetitive firing (Giese et al., 1998; Shao et

al., 1999; Yue and Yaari, 2004; Kim et al., 2005; Gu et al., 2007), an increase in the contribution

of K2P channels lacking time-dependent inactivation with a concomitant decrease in the

contribution of K_V channels to action-potential repolarization would reduce time- and

frequency-dependent action potential broadening.

Although the effects of ultrasound on action-potential widths are predominantly due to

acceleration of the repolarization phase, we also noted effects on the rising phase of the action

potential. These effects are readily apparent in the first derivative of membrane voltage during

the action potential (Figure 8A) or in action potential phase plots (plots of the first derivative of

voltage versus voltage, Figure 8B). In fact, the maximum rates of voltage rise and fall during the

action potential were both consistently increased by ultrasound throughout a spike train

(Figure 8C-D). The effect on the falling phase is to be expected based on the observed decrease

in spike width and the hypothesis that ultrasound activates K2P channels. The effect on the

rising phase is also consistent with this hypothesis, as activation of K2P leading to reduced Na_V

channel inactivation would increase the number of Na_V channels available to activate during

the rising phase of the action potential. Alternatively, increased K2P conductance could

increase the rate of action potential rise by decreasing the membrane time constant (Brickley et

al., 2007). Consistent with these results, decreases in the rates of action-potential rise and fall

were seen with knock-outs of K2P channels in cerebellar granule neurons (Brickley et al., 2007)

and hypothalamic hypocretin/orestin neurons (Gonzalez et al., 2009), while in a heterologous

action-potential firing model higher levels of K2P expression increased the rate of action-

potential rise (MacKenzie et al., 2015).

In addition to effects of ultrasound on the rising and falling phases of the action

potential, we also found effects on the AHP. To quantify these effects, we measured the voltage

minimum between action potentials during repetitive firing. Because measurements of this

parameter are very sensitive to changes in resting membrane voltage and series resistance that

can occur over long recording times, we compared voltage minimums before, during, and after

ultrasound application within the same voltage trace (Figure 9A-B) and made a similar

comparison for control recordings. We performed these comparisons for ultrasound applied 1 s

after the start of a current, for cells firing at an average frequency of 5 Hz in the control

condition. This firing frequency is near the transition region between the inhibitory and

potentiating effects of ultrasound on spike frequency, such that the spike frequency is similar

for the control and ultrasound conditions, allowing us to compare a similar number of

interspike voltage minima for the control and ultrasound conditions (Figure 9C). This analysis

demonstrates that the depth of the AHP is greater during the ultrasound application than

before or after it, or during the same time windows for the control condition. Together with the

effects of ultrasound on spike waveform (Figure 6), this result supports the idea that ultrasound

activates a sustained outward current, which limits Na_V channel inactivation and thereby

potentiates high-frequency firing. Removal of Na_V channel inactivation by membrane

hyperpolarization also explains how ultrasound can rescue spiking in neurons that have entered

a refractory state due to accumulation of Na_V channel inactivation (Figure 3C, *bottom right*).

Physical mechanism of neuromodulation by high-frequency ultrasound. The idea that

ultrasound acts on K2P channels is also consistent with the physical effects of ultrasound on

biological tissue. At 43 MHz, two plausible mechanisms through which ultrasound might

modulate the activity of ion channels are heating and mechanical stress due to acoustic

radiation force. Absorption of acoustic energy by biological tissue as heat can increase its

temperature, with effects on ion channel gating and all other biological reactions. Absorption

also results in attenuation of ultrasound intensity as the wave propagates, creating spatial

gradients in intensity that give rise to radiation force, which in turn produces tissue

displacement and strain. At the microscopic scale, this displacement and strain may involve

increased tension in the cell membrane, cytoskeleton, and extracellular matrix, all of which may

affect excitability through mechanical effects on ion channel proteins. Among the K2P channels

that may be expressed by CA1 pyramidal cells, TREK and TRAAK channels are especially

sensitive to thermal and mechanical stimuli.

To gain further insight into these physical mechanisms, we performed finite-element

simulations of the effects of ultrasound on brain slices in the context of our experimental set-

up. The simulated spatial profiles of ultrasound-induced heating and macroscopic tissue

displacement in response to radiation force are shown in Figure 10A-B. Notably, the spatial

profiles of heating and displacement effects are significantly wider than the 90-micron diameter

of the focal volume of the ultrasound beam, with significant heating and displacement

occurring several hundreds of microns away from the beam axis (Figure 10C). This is an

important result because the thermo- and mechanosensitive K2P channels TREK-1 and TRAAK

are expressed at high density at the nodes of Ranvier of vertebrate neurons (Brohawn et al.,

2019; Kanda et al., 2019). The first node of Ranvier is located approximately 100 microns from

the axon initial segment (Kole, 2011). In our experiments, the soma of the patched neuron is

located approximately on the axis of the ultrasound beam (see *Materials and Methods*) so the

first node of Ranvier is probably within the region of the tissue exposed to thermal and

mechanical effects of ultrasound. Axonal K^+ channels play important roles in regulating

excitability (Shah et al., 2008; Kole, 2011; Kanda et al., 2019). It is therefore plausible that a

subpopulation of TREK-1 or TRAAK channels at the nodes of Ranvier could contribute to the

neurophysiological effects of ultrasound. The magnitude of the temperature change is also

consistent with a thermal mechanism for the effects of ultrasound. The maximum temperature

change in the simulation is 1.3 C; temperature changes of this size have previously been shown

to affect neural excitability (Owen et al., 2019). The maximum value of the simulated

displacement (1.7 microns) is similar to the displacement measured in the retina during

ultrasound neuromodulation with stimulus parameters similar to those used here (Menz et al.,

2019).

Since the activity of TREK and TRAAK channels is highly temperature-sensitive, we

considered whether our results might represent an artefact due to the experiments being

performed at room temperature (21-23°C). Room temperature is near the threshold for

temperature activation of these channels, such that they are mainly inactive in the absence of

other stimuli such as membrane tension, lipid agonists, or acidic pH (Maingret et al., 2000; Kang

et al., 2005). Thus at room temperature in the absence of additional gating stimuli, the relative

increase in thermosensitive K2P current would be greater in our experiments than at

physiological temperatures, which might lead us to overestimate the importance of K2P

channels in the response to ultrasound. On the other hand, the midpoints of the temperature-

activation curves for TREK and TRAAK channels are near 37°C (in other words, near body

temperature in mammals) so that the absolute increase in K2P current in response to increased

temperature is near maximal at physiological temperatures, so we might instead be

underestimating the thermal effects of ultrasound on K2P channels that would occur in

physiological contexts. To cut through this speculation and address these issues, we repeated

our experiments measuring the neurophysiological effects of ultrasound in cells firing at a spike

frequency of approximately 5 Hz at near-physiological temperature (30°C) (Supplemental Figure

4). Similar to what we observed at room temperature, ultrasound inhibited action potential

firing at this relatively low spike frequency and decreased action potential width, indicating that

these effects are not especially sensitive to the ambient temperature. However, the

hyperpolarizing effect of ultrasound on the resting membrane voltage that we observed at

room temperature was no longer apparent at 30°C, possibly due to increased noise in the

baseline voltage or more hyperpolarized (in other words, closer to the K^+ reversal potential)

resting voltage at higher temperature.

It is instructive to consider the amplitude and time course of the membrane capacitance

change in response to ultrasound (Figure 5) in the context of possible thermal and mechanical

mechanisms. As described above (under *Effects of ultrasound on resting membrane potential*),

the capacitance change in response to ultrasound is fast relative to the membrane time

constant, so we can assume that the total charge on the membrane is constant during the

initial transient depolarization in response to ultrasound (Figure 5A, *left arrow*). In other words,

the numerator in the equation V = Q/C is constant, so for small changes in voltage the relative

change in voltage is approximately inversely proportional to the relative change in capacitance.

Empirically, membrane capacitance increases by approximately 1% per degree C (Taylor, 1965).

This is consistent with the size of the simulated temperature rise (peak simulated temperature

rise of 1.3°C compared with the measured amplitude of the initial decrease in voltage of 0.7 ±

0.1%). However, the time course of the temperature rise (Figure 10D) is much slower than the

time course of the voltage transient (which again, assuming constant Q, is identical to the time

course of the capacitance change). The time course of the change in resting membrane

potential (173 ± 18 ms, Figure 5), however, parallels that of the temperature rise. (A

capacitance change on the time course of the simulated temperature rise would not have a

significant effect on the membrane voltage, as it would be counteracted by ionic currents.)

Thus, the simulated ultrasound heating results strongly suggest that ultrasound affects action-

potential firing in our experiments at least in part through a thermal effect on ion channels, but

do not explain the presence of the capacitive transients.

We considered whether the time course of the capacitive transients could instead be

explained by the dynamics of the tissue mechanical response to acoustic radiation force. We

sought to determine whether, having already modeled the static displacement of the tissue, we

could, without retroactively changing any of the tissue material properties, obtain a time course

for tissue displacement similar to that of the capacitive transients using a simple viscoelastic

model with reasonable tissue viscous properties (see *Materials and Methods*). We found that

this could be achieved using a shear viscosity (μ) of 1 Pa·s (Figure 10E). Since the tissue is

essentially incompressible in our model (Poisson's ratio (ν) = 0.4998), this is equivalent to a

relaxation time of $2\mu\cdot(1+\nu)/E = 6$ ms (where E is Young's modulus). Biological tissue is a highly

heterogeneous material that displays a variety of active and passive mechanical responses to

force, spanning time scales from milliseconds to hours (Ricca et al., 2013), and as a result its

viscous properties are highly sensitive to the time scale of the measurement and even complex

viscoelastic models encompassing multiple relaxation times may not fully describe the

viscoelastic behavior of tissue. Nonetheless, the shear viscosity/relaxation time in our model is

reasonable for a soft, gel-like material, and is comparable to fast relaxation times observed

experimentally in brain tissue (Arbogast and Margulies, 1999; Abolfathi et al., 2009; Rashid et

al., 2012, 2013). Moreover, the simulated time course of displacement is consistent with

experimental measurements of the tissue displacement in response to ultrasound at 43 MHz

and 40 W/cm^2 in the salamander retina, which was found to be complete in less than 10 ms

(Menz et al., 2019).

We can therefore make the reasonable assumption that the capacitive transients are

due to a mechanical effect on membrane properties, and we can estimate the size of the

potential ion-channel gating effects that would occur as a result of this mechanical effect. The

capacitance of a lipid bilayer membrane is given by $C = \varepsilon \cdot \varepsilon_0 \cdot A/L$, where ε is the dielectric

constant of the hydrophobic core of the lipid bilayer, ε_0 is the permittivity (polarizability) of free

space, A is membrane area, and L is the thickness of the hydrophobic core of the membrane.

For small strains like those under consideration here, lipid bilayer membranes can be

considered incompressible, such that a 1% increase in capacitance corresponds to a 0.5%

increase in area and a 0.5% decrease in thickness (White and Thompson, 1973; Alvarez and

Latorre, 1978). An increase in membrane area can be converted to an increase in membrane

tension (γ) according to $\gamma = \Delta A \cdot K_A$, where ΔA is the relative change in area and K_A is the area

elastic constant of the membrane. Area elastic constants measured for lipid membranes are on

the order of 100's of mN/m (Evans et al., 1976; Kwok and Evans, 1981; Needham and Nunn,

1990). If the capacitance transients are due to membrane strain, the resulting membrane

tension is on the order of a few 0.1 mN/n to a few mN/m. These values are similar to the

tension thresholds for activation of mechanosensitive K2P channels (estimated as 0.5-4 mN/m

for activation of TREK-1 and TRAAK (Brohawn et al., 2014)), which are low relative to other

known mammalian mechanosensitive channels. Notably, a recent *in vivo* ultrasound

neuromodulation study of the murine sciatic nerve at 4 MHz, found that tissue displacement *in*

vivo was highly correlated with the neuromodulation effects (Lee et al., 2020). Nonetheless,

additional data or theoretical advances would be required to firmly associate these capacitive

transients with changes in membrane tension. If such an association could be made, it would

provide strong evidence that ultrasound modulates action potential firing through mechanical

effects of radiation force in our experiments. At present, our results do not rule out this idea,

but the case for mechanical effects remains speculative, while the role of thermal effects seems

highly plausible. Nonetheless, our simulation results support the conclusion that both inhibitory

and excitatory effects of high-frequency ultrasound on action-potential firing are due to

activation of thermo- and mechanosensitive K2P channels.

DISCUSSION

To summarize, the neurophysiological effects of ultrasound that we have described here can all

be explained by activation of a sustained outward current. We argue that the molecular basis of

this outward current is most likely one or more of the K2P channels expressed by CA1

pyramidal neurons. Although a variety of voltage-dependent K^+ currents shape the action-

potential waveform and regulate excitability in these neurons, several arguments suggest that

K2P channels are the molecular basis of the ultrasound-activated outward conductance. First,

the K2P channels TREK and TRAAK, being strongly mechanosensitive and thermosensitive, have

biophysical properties that make them especially sensitive to physical effects of ultrasound.

Second, the fact that ultrasound has similar effects on firing frequency whether it is applied 500

ms before or 1 s after the start of a current step suggests that ultrasound affects firing through

a channel that does not undergo prolonged voltage-dependent inactivation during sustained

depolarizations. Finally, the neurophysiological effects of ultrasound are, strikingly, essentially

the opposite of those caused by knock-out of K2P channels in other neurons, as detailed in the

following paragraph.

Knock-out of TASK-3 channels in cerebellar granule neurons increased excitability at low

input currents, but decreased excitability at high input currents and led to failure of sustained

high-frequency firing (Brickley et al., 2007). In addition, knock-out of TASK-3 decreased the

maximum firing frequency and decreased action-potential height while increasing action

potential width through decease in the rates of both action-potential rise and fall. Similarly,

double knock-out of TASK-1 and TASK-3 in hypothalamic hypocretin/orexin neurons inhibited

high-frequency action potential firing, reduced the rates of action-potential rise and fall, and

decreased the depth of the AHP (Gonzalez et al., 2009). The connection between our results

and these knock-out studies is supported by experiments in a heterologous model system

consisting of HEK cells transfected with TREK-1 and TASK-3 (with endogenous K_V channels

blocked) and Na_V channels simulated by dynamic clamp (MacKenzie et al., 2015). In this model

system, high levels of K2P expression were necessary for repetitive action potential firing, and

increased K2P conductance increased the rates of action-potential rise and fall and increased

the threshold current for action-potential firing. (In the context of this model system, a 74%

potentiation of the K2P conductance by halothane produced effects on the rates of rise fall of

the same order as we see here, potentially providing an estimate of the potentiation of K2P

conductance by ultrasound in our experiments. However, caution should be used in

extrapolating from this heterologous model system to neurons expressing a considerably more

complex array of ion channels.) Finally, TREK-1 and TRAAK channels are also necessary for high

frequency firing at the nodes of Ranvier of afferent neurons (Kanda et al., 2019).

Ultrasound neuromodulation has been studied in vertebrate axon preparations, where

it has generally been found that ultrasound inhibits action-potential conduction, with the effect

specifically attributed to heating in some cases (Young and Henneman, 1961; Mihran et al.,

1990; Tsui et al., 2005; Colucci et al., 2009). These results are consistent with the idea that

activation of K2P channels by ultrasound can inhibit action potential firing, but it would be

worthwhile to revisit these experiments to see whether the bidirectional, spike-frequency-

dependent effect that we observe is also present in such preparations. Our results also suggest

an approach to ultrasound neuromodulation in which action potential propagation is the locus

of neuromodulation, targeting white-matter tracts instead of soma-dense gray matter. The idea

that ultrasound-activated K^+ currents can both inhibit and potentiate firing might also help

explain why ultrasound can both inhibit and potentiate neural activity *in vivo* (Min et al., 2011).

Although activation of K2P channels is sufficient to explain our results, we do not rule

out the possibility that ultrasound affects other channels in addition to K2P's; indeed, we think

it is likely that ultrasound does affect other channels to some extent. All ion-channel gating

reactions are sensitive to temperature, with typical Q10 values of ~3, such that their rates

would be expected to increase by about 10% based on the temperature changes in our

simulations (Hille, 2001). Mechanical effects of radiation force could also affect channels

besides K2P channels. The mechanically gated channel Piezo2 is expressed in a subset of CA1

pyramidal neurons (Wang and Hamill, 2020). Piezo1 (Qiu et al., 2019) and TRP channels (Oh et

al., 2020; Yoo et al., 2020) have been experimentally linked to ultrasound neuromodulation

effects. In addition, most ion channels and membrane proteins, while not functioning

physiologically as mechanoreceptors, are sensitive to mechanical stimuli to some extent, either

through the energetics of their interactions with hydrophobic core of the lipid bilayer or

through mechanical interactions with the cytoskeleton or extracellular matrix. In fact, gating of

voltage-dependent Na^+ (Morris and Juranka, 2007), K^+ (Tabarean and Morris, 2002; Laitko and

Morris, 2004; Beyder et al., 2010), and Ca^{2+} (Calabrese et al., 2002) channels, and of NMDA

receptor channels (Kloda et al., 2007), can be modulated by membrane stretch in membrane

patches. However, we previously were unable to detect any mechanical modulation of

heterologously expressed NaV1.2 channels by ultrasound at 43 MHz and 90 W/cm^2 under

conditions where ultrasound activated the mammalian mechanoreceptor channel Piezo1

(Prieto et al., 2018). Neural Na_V channels and KCNQ channels interact with the periodic actin

cytoskeleton of axons through spectrin and ankyrin-G at the axon initial segment and nodes of

Ranvier and (Zhou et al., 1998; Pan et al., 2006; Leterrier, 2018), suggesting that they may be

sensitive to modulation by cytoskeletal tension due to acoustic radiation force. The

concentration of TREK-1 and TRAAK channels at the nodes of Ranvier suggest that some similar

interaction with the cytoskeleton may be involved in the localization of these channels,

although this has not been demonstrated, and the intracellular domains that would facilitate

such interactions are relatively small in K2P channels. Ultrasound has been shown to cause

changes in cytoskeletal structure (Mizrahi et al., 2012), which might also affect the activity of

cytoskeleton-associated channels. Thus investigation of the role of other ion channels in

ultrasound neuromodulation should continue. Notably, several of the channels discussed above

have roles in synaptic transmission, so effects of ultrasound on these channels would not be

revealed by our experiments on somatic excitability in response to injected current.

It is well established that high-intensity light at infrared and shorter wavelengths can

modulate neural activity through tissue heating. As a general principle, thermal

neuromodulation effects in response to optical stimulation would be expected to be very

similar to thermal neuromodulation effects caused by ultrasound, and studies of optical

neuromodulation could therefore provide useful guidance in interpreting our results. However,

optically based thermal neuromodulation experiments are highly heterogeneous in terms of the

neuromodulation effect, the temperature rise required to produce the effect, and the

mechanistic interpretation of the results (Wells et al., 2007; Richter et al., 2011; Shapiro et al.,

2012; Duke et al., 2013; Walsh et al., 2016; Lothet et al., 2017; Paris et al., 2017; Owen et al.,

2019; Zhu et al., 2019), so their usefulness is limited in this respect. Both inhibition and

potentiation of firing have been reported, and the increase in temperature has varied

considerably, ranging from less than 1 C to 10's of degrees C. The temperature rise in our

simulations is on the low end of this range. However, it has been proposed that spatial or

temporal gradients in temperature, rather than the absolute temperature change, may

determine the response to thermal stimuli (Wells et al., 2007; Paris et al., 2017). In addition,

heating can cause phase changes in lipid bilayers, which have been shown to modulate ion

channel activity (Seeger et al., 2010); in this case, the response to heating would also depend

critically on the initial temperature. Interestingly, a recent study demonstrated inhibition of

firing by small (2°C or less) increases in temperature in several different types of neurons, but

not in CA1 pyramidal neurons (Owen et al., 2019). These effects were attributed to inward

rectifier (K_{ir}) potassium channels, which are not expressed in CA1 pyramidal neurons. However,

our results suggest that the lack of an effect in pyramidal neurons could also be explained if the

experiments were performed at a point on the f-i curve where competing inhibitory and

excitatory effects of heat-activated K^+ current result in no net effect on firing frequency. A

recent *in vivo* ultrasound neuromodulation study in a rat model using ultrasound at 3.2 MHz

with exceptionally long ultrasound exposure times (10's of seconds; see below for a more

general discussion of *in vivo* ultrasound neuromodulation studies) also found that inhibition

could be produced by ultrasound-induced temperature rises of 2°C or less (Darrow et al., 2019),

consistent with this result and our results.

Finally, a critical question—to which we cannot yet provide a definitive answer—is

whether the mechanisms underlying the neuromodulatory effects of ultrasound are the same

in our experiments and in *in vivo* experiments using low frequencies. Two results that strongly

argue against similar mechanisms are the much lower intensities that have been reported to

cause neuromodulatory effects in some low-frequency, *in vivo* experiments in small animal

models (Tufail et al., 2010) as compared with our results, and the apparent increase in efficacy

of ultrasound neuromodulation at lower frequencies in *in vivo* experiments. Both thermal and

radiation force effects are proportional to ultrasound intensity, and to the ultrasound

attenuation coefficient, which in tissue is proportional to frequency raised to the power of ~1.1

(Hand, 1998), so these effects would generally be smaller in *in vivo* experiments (especially

small animal experiments) as compared with our experiments. Compounding this issue is the

fact that opposite dependences on the ultrasound frequency have been observed in *in vivo* and

in vitro experiments. In an *in vivo* mouse model of ultrasound neuromodulation, it was

determined that the efficacy of neuromodulation decreased with increasing frequency over the

range 0.25-2.9 MHz (King et al., 2013; Ye et al., 2016). This frequency dependence is the

opposite of what would be expected for either a thermal or radiation force mechanism. In

contrast, Menz et al. (2019) found that the efficacy of neuromodulation increases with

frequency over the range 1.9-43 MHz in the retina *in vitro*, a thin neural tissue preparation

similar to the one used in our experiments. However, they proposed a model to explain this

discrepancy. Lower ultrasound frequencies generally result in a larger stimulated tissue volume,

which could translate into a more effective stimulus for certain structures of circuit-level neural

connectivity, despite a weaker effect of low-frequency ultrasound at the level of an individual

cell. (The model was presented in the context of a radiation force mechanism, but the same

principle could apply for a thermal mechanism). The idea that circuit-level mechanisms can

amplify the response to ultrasound is supported by comparison of our results with the response

to ultrasound in the retina at 43 MHz. In the retina, potentiation of action-potential firing by

ultrasound at 43 MHz saturates at 10 W/cm^2, as measured at the population level in an intact,

active neural circuit. Although we have not performed a detailed investigation of the intensity

dependence, we find that a much higher intensity, 50 W/cm^2, produces relatively moderate

effects on excitability in single cells in the absence of significant network activity. Focusing

solely on the local, cell-level amplitude of thermal and radiation force effects may therefore

overlook important factors related to the global, network-level distribution of these effects.

Such considerations may eliminate the differences in effective ultrasound parameters for *in*

vivo and *in vitro* experiments as an argument against similar physical mechanisms for

neuromodulation by high- and low-frequency ultrasound.

In conclusion, our results demonstrate that high-frequency ultrasound is a viable and

promising modality for neuromodulation applications where frequency is not limited by

transmission through the skull, and our insights into the common molecular mechanisms

underlying both inhibitory and excitatory effects of high-frequency ultrasound pave the way for

rational design and optimization of neuromodulation protocols to consistently produce either

inhibitory or excitatory effects.

ACKNOWLEDGMENTS

We thank the PIs and members of the Baccus, Butts-Pauly and Huguenard laboratories for

helpful discussions. We thank Mike Menz and Stephen Baccus for critically reading the

manuscript. We thank Marianna Kiraly, Dong Li, and Jason Clark for help with hippocampal slice

preparation and recording. We thank the Stephen Boxer lab for sharing their plasma cleaner.

This work was supported by NIH R01 EB019005 (M.M. and B.K.Y.) and NIH BRAIN initiative R01

NS11215 (M.M. and B.K.Y.).

The authors declare no competing financial interests.

AUTHOR CONTRIBUTIONS

Martin Prieto performed experiments, analyzed data, and wrote the manuscript. Martin Prieto

and Kamyar Firouzi performed computational modeling. Merritt Maduke, Daniel Madison, and

Butrus Khuri-Yakub supervised research. All authors contributed to the design and

interpretation of experiments and edited and revised the manuscript.

Table 1. Values of Material Properties Used in Finite-Element Simulations.

	water	polystyrene	brain tissue
density (kg/m^3)	1000[a]	1040[b]	1007[c]
speed of sound (m/s)	1500[a]	2300[b]	1538[c]
attenuation coefficient at 43 MHz (neper/m)	46[d]	160[e]	253[f]
heat capacity (J/kg·K)	4180[a]	1200[g]	3500[h]
thermal conductivity (W/m·K)	0.6[a]	0.1[g]	0.5[h]
Young's modulus (Pa)	(not applicable)	10^9 [i]	500[j]
Poisson's ratio	(not applicable)	0.4[i]	0.4998[k]
shear viscosity (Pa·s)	(not applicable)	(not applicable)	1[l]

Sources for material properties are as follows: [a]standard value; [b]based on typical acoustic

properties of plastics (Selfridge, 1985); [c]following Menz et al. (2019), based on (Thijssen et al.,

1985); [d](Company, 1965); [e]measured (Prieto et al., 2018); [f]following Menz et al. (2019), based

on (de Korte et al., 1994); [g]based on typical thermal properties of plastics (Gaur and

Wunderlich, 1982; Harper, 2006); [h]typical values for soft tissues (Hand, 1998); [i]based on typical

mechanical properties of plastics (Harper, 2006); [j]Menz et al. (2019), from measurements of

ultrasound-induced displacement in the retina; [k]tissue assumed to be incompressible for small

deformations; [l]see text.

Figure 1. Consistent inhibition of action potential firing by high-frequency ultrasound. A.
Diagram of experimental set-up. Ultrasound is applied to 500-micron hippocampal brain slices
resting on a 25-micron film of polystyrene. The 43-MHz focused transducer is located below the
experimental chamber, with ultrasound propagating perpendicular to the bottom of the
recording chamber. **B.** Experimental protocol and example voltage traces showing inhibition of

action potential firing by ultrasound. A 1-s, continuous-wave ultrasound pulse at 43 MHz and 50 W/cm^2 (*red bar*) is applied 500 ms before the start of a 2-s current injection. Voltage traces are shown in the presence (*red*) and absence (*blue*) of the ultrasound stimulus. The *dashed line* indicates the resting membrane voltage. **C.** Example raster plots showing a consistent effect of ultrasound on firing frequency. The results of twenty consecutive trials of the protocol in panel B, alternating between the control (*top*) and ultrasound (*bottom*) conditions, are shown. The voltage traces were divided into 50-ms bins; a solid black bin indicates that an action potential occurred within that particular time bin. Time is relative to the start of the ultrasound pulse. **D.** Spike-time histograms prepared by summing the ten trials for the control (*top, blue bars*) and ultrasound (*bottom, red bars*) conditions from panel C. **E.** Summary of the effects of ultrasound for N = 66 cells. The average firing frequency during the first 500 ms of the current step is shown for the control (*blue*) and ultrasound (*red*) conditions. **F.** Stability of the ultrasound response. Mean (±SEM, N = 10 cells) spike frequencies during the first 500 ms of a current step in the presence (*red circles*) and absence (*blue circles*) of ultrasound for the protocol shown in panel B, as a function of time relative to break-in (establishment of whole-cell recording configuration). Spike frequencies were measured at various time points between 0 and 10, 10 and 20, 20 and 30, and 30 and 40 minutes after break-in. The x-values represent the mean start time for the protocol to measure spike frequencies (which comprised 2 minutes of recording time). The amplitude of the current step was adjusted over time to maintain spiking behavior as close as possible to that at the start of the experiment.

Figure 2. Ultrasound can inhibit or potentiate action-potential firing. A. Example frequency-input curve showing average firing frequency during the first 500 ms of a current step as a function of input current, with (*red*) and without (*blue*) a 1-s ultrasound pulse starting before 500 ms before the start of the current step. Each point represents the average of three trials on the same cell. **B.** Example voltage traces for the cell in A showing action potential firing during the first 500 ms of current steps to either +30, +40, +80, or +250 pA, with (*red*) and without (*blue*) ultrasound. The *dashed lines* indicate the approximate resting membrane voltage of -60 mV. **C.** Mean (±SEM, N = 9 cells) change in spike frequency in response to ultrasound as a function of input current. **D.** Inhibition is strongest near threshold. Mean (±SEM, N = 7 cells) change in spike frequency in response to ultrasound as a function of input current relative to the threshold current for action potential firing, for near-threshold currents.

Figure 3. Ultrasound can also inhibit or potentiate action-potential firing when applied late in a current step. A. Experimental protocol and example voltage traces with (*red*) and without (*blue*) ultrasound. Ultrasound is applied 1 s after the start of a 3-s current step. The *dashed line* indicates the resting membrane voltage. **B.** Example frequency-input curve showing average firing frequency during the ultrasound application for the protocol in A (*red*) and during the same time window without ultrasound (*blue*), as a function of input current. Each point represents the average of three trials on one cell. **C.** Example voltage traces for the cell in B showing action potential firing during the ultrasound application (*red*) and during the same time window without ultrasound (*blue*) in response to current steps to either +50, +100, +250, or +450 pA. The *dashed lines* indicate the approximate resting membrane voltage of -60 mV. **D.** Mean (±SEM) change in spike frequency in response to ultrasound as a function of input current (N = 3 cells at +50 pA; N = 6 cells at all other input currents). (In 3 cells, the firing frequency at +50 pA was zero for both the control and ultrasound conditions).

Figure 4. Effects of ultrasound on action potential timing. A-C. Mean (±SEM) latency between

the start of the current step and the first action potential, and mean intervals between the first

and second, and second and third, etc. action potentials, for cells firing at average frequencies

of approximately 5 Hz (N = 13 cells), 10 Hz (N = 15 cells), and 20 Hz (N = 13 cells) during the first

500 ms of the current step, with (*red*) or without (*blue*) a 1-s ultrasound pulse starting 500 ms

before the current step. The actual spike frequencies were 5.6 ± 0.1, 10.5 ± 0.2, and 20.4 ± 0.3

Hz, and the injected currents were 70 ± 7, 115 ± 9, and 292 ± 28 pA for the 5, 10, and 20 Hz

conditions. **D.** Example voltage traces showing decreased interval between the first and second

action potentials at low average firing frequency. The top panel shows the first two action

potentials for the control (*blue*) and ultrasound (*red*) conditions. The *dashed line* indicates the

approximate resting membrane voltage of -60 mV. The bottom panel shows the same data,

aligned to the action potential threshold on a zoomed-in time scale. **E.** Same as A-C, for an

average firing frequency of 5 Hz, except that ultrasound was applied 1-s after the start of a 3-s

current step, and the average firing frequency and intervals/latency were determined during

ultrasound stimulus or during the same time period without ultrasound (N = 6). The actual spike

frequency was 5.2 ± 0.2 Hz, and the injected currents was 183 ± 21 pA.

Figure 5. Effects of ultrasound on resting membrane potential and membrane capacitance. A.

Six individual voltage traces (*pink*) and the average of these voltage traces (*red*) showing the

effect of ultrasound (*red bar*) on resting membrane potential. The *black arrows* indicate

transients due to changes in membrane capacitance. The *dashed line* indicates the mean resting

membrane voltage. **B.** Zoomed-in timescale showing the fast voltage transients (*black arrows*

in A) for ultrasound onset (*top left*) and offset (*bottom right*). Exponential fits (*black lines*) to the

rise and fall of the voltage transients give amplitudes and time constants of 0.45 mV and 16.3

ms for the onset and 0.22 mV and 6.7 ms for the offset, for the example shown; mean values

(±SEM) were 0.41 ± 0.04 mV and 10.4 ± 1.2 ms for the onset and 0.35 ± 0.04 mv and 9.7 ± 1.7

ms for the offset (N = 15). No significant differences were found between the time constants (P

= 0.73) or the amplitude (P = 0.087) of the transients (paired, two-tailed Student's t-tests) **C.**

Slow membrane hyperpolarization in response to ultrasound from the average trace in panel A

on a zoomed-in scale, along with an exponential fit (*black line*) to the initial hyperpolarization.

The amplitude and time constant of the exponential fit were 1.55 mV and 132 ms for the

example shown; mean values (±SEM) were 2.4 ± 0.3 mV and 173 ± 18 ms (N = 15).

Figure 6. Ultrasound decreases action-potential width. A-B. Mean (±SEM, N = 13 cells) action-potential widths (A) and example action-potential waveforms aligned to the action-potential threshold (B) as function of action-potential number in the presence (*red*) and absence (*blue*) of a 1-s ultrasound pulse starting 500 ms before the current step, for cells firing at an average firing frequency of approximately 5 Hz (as measured during the first 500 ms of the current step)

in the control condition. **C-D.** As in A-B, but for cells firing at an average firing frequency of
approximately 10 Hz in the control condition (N = 15). **E-F.** As in A-B, but for cells firing at an
average firing frequency of approximately 20 Hz in the control condition (N = 13). **G-H.** As in A-
B, but with ultrasound applied 1 s after the start of a 3-s current step, for cells firing at an
average firing frequency of approximately 5 Hz in the control condition, with firing frequency
determined in a 1-s window starting 1 s after the current step (corresponding to the time
period of the ultrasound stimulus), and action potential number relative to the start of the
ultrasound stimulus (N = 6).
879

Figure 7. Ultrasound reduces action-potential broadening. Example data showing action-

potential width as a function of action-potential number (indicated in *grayscale*, scale bar at far

left) and input current level, for the first 500 ms of the current step, for currents from 0 to +450

pA in 50 pA steps, with (*right*) or without (*left*) a 1-s ultrasound pulse starting 500 ms before the

current step. The vertical lines indicate the approximate location of the transition between

inhibitory and potentiating effects of ultrasound.

Figure 8. Effects of ultrasound on depolarization and repolarization rates. A. Example traces

showing the membrane voltage (*top*) and its first derivative (*bottom*) for the first three action

potentials in response to a +100 pA current step in the presence (*red*) and absence (*blue*) of a 1-

s ultrasound pulse starting 500 ms before the current step, aligned to the action potential

threshold. **B.** Phase plots for the action potentials shown in panel A. **C-D.** Maximum rates of

depolarization (*left)* and repolarization (*right*) during the action potential (mean ± SEM, N = 13

cells), as a function of action potential number, in the presence (*red*) and absence (*blue*) of a 1-s

ultrasound pulse starting 500 ms before the current step, for cells firing at an average firing

frequency of 5 Hz (**C**) or 20 Hz (**D**) in the control condition.

◆ prepulse, control (N = 6)
● pulse, control (N = 6, AP # 1-5; N = 3, AP # 6)
■ postpulse, control (N = 6, AP # 1-2; N = 5, AP # 3; N = 3, AP # 4)
◆ prepulse, ultrasound (N = 6)
● pulse, ultrasound (N = 6, AP # 1-3; N = 4, AP # 4-6)
■ postpulse, ultrasound (N = 6)

Figure 9. Ultrasound increases the depth of the afterhyperpolarization. A. Example voltage
traces comparing voltage minima between action potentials in response to a 3-s, 250-pA
current step with (*red voltage trace, right*) and without (*blue voltage trace, left*) a 1-s
ultrasound pulse (*red bar*) starting 1 s after the start of the current step. The *solid black lines*
connect the voltage minima between action potentials before, during, or after the ultrasound
pulse, or during the corresponding time periods for the control condition. The dashed lines
indicate a reference voltage level of -40 mV. The resting membrane voltage for this cell was -63
mV. **B.** Same as panel A, on a zoomed-in voltage scale. The *red diamonds* indicate voltage
minima before the ultrasound pulse, *red circles* indicate voltage minima during the ultrasound
pulse, and *red squares* indicate voltage minima following the ultrasound pulse; *blue symbols*
indicate voltage minima for the corresponding time periods for the control condition. The
dashed lines indicate a reference voltage level of -44 mV. **C.** Mean (±SEM, N = 3-6 cells, see
figure panel for details) values of the voltage minimum, as a function of action potential (AP)
number for the first four to six action potentials before, during, and after the ultrasound pulse,
along with the equivalent mean values for the control condition, following the symbolism
indicated in panel B. The means were determined for cells firing at the same average frequency

(5 Hz) during a 1-s window starting 1 s after the start of the current step (corresponding to the
period of the ultrasound stimulus) in the control condition. For clarity, the results are shown
separately for the control group only (*left*), the ultrasound group only (*middle*), and for both
groups simultaneously (*right*). Significant differences between groups were only found in the
presence of ultrasound (P = 1.0 x 10^{-4}, 1.4 x 10^{-5}, and 0.16 for before versus during, during
versus after, and before versus after the ultrasound pulse; P = 0.92, 0.39, and 0.35 for
comparisons of the same time periods in the control condition (unpaired, two-tailed Student's
t-tests, unequal variance).
924

Figure 10. Simulated tissue heating and displacement in response to ultrasound. A. Spatial

profile of temperature after one second of ultrasound exposure at 43 MHz and 50 W/cm^2, as a

function of axial distance from the transducer surface and radial distance from the ultrasound

beam axis, in a 500-micron thick brain slice and 25-micron thick polystyrene film (*middle two*

layers) and the surrounding fluid (external solution, *top layer*, and distilled water, *bottom*

layer). **B.** Spatial profile of the static total displacement in response to acoustic radiation force

in the brain slice and polystyrene film. **C.** Normalized values of the acoustic intensity (*solid black*

line), temperature rise after 1 second of ultrasound exposure (*solid gray line*), and total

displacement (*dashed line*), at a depth of 250 microns in the brain slice, as a function of radial

distance. **D.** Time course of the temperature rise in response at ultrasound at a depth of 250

microns in the brain slice on the axis of the ultrasound beam. The time course of the

temperature change can be described by two exponential components with amplitudes and

time constants of -0.56 C and 30 ms, and -0.57 C and 295 ms, for a weighted time constant of

164 ms. **E.** Time course of the displacement in response at ultrasound at a depth of 250 microns

in the brain slice on the axis of the ultrasound beam. The time course of the displacement

change can be described by two exponential components with amplitudes and time constants

of -1.2 microns and 6 ms, and -0.13 microns and 344 ms, for a weighted time constant of 39 ms.

Note that the steady-state displacement is slightly smaller than in the static displacement

simulation due to the inclusion of the fluid loading in the dynamic displacement simulation.

Abolfathi, N., A. Naik, M. Sotudeh Chafi, G. Karami, and M. Ziejewski. 2009. A micromechanical
procedure for modelling the anisotropic mechanical properties of brain white matter.
Comput Methods Biomech Biomed Engin. 12:249-262.
Alvarez, O., and R. Latorre. 1978. Voltage-dependent capacitance in lipid bilayers made from
monolayers. *Biophys J*. 21:1-17.
Arbogast, K.B., and S.S. Margulies. 1999. A fiber-reinforced composite model of the viscoelastic
behavior of the brainstem in shear. *J Biomech*. 32:865-870.
Bean, B.P. 2007. The action potential in mammalian central neurons. *Nat Rev Neurosci*. 8:451-
465.
Beyder, A., J.L. Rae, C. Bernard, P.R. Strege, F. Sachs, and G. Farrugia. 2010. Mechanosensitivity
of Nav1.5, a voltage-sensitive sodium channel. *J Physiol*. 588:4969-4985.
Blackmore, J., S. Shrivastava, J. Sallet, C.R. Butler, and R.O. Cleveland. 2019. Ultrasound
Neuromodulation: A Review of Results, Mechanisms and Safety. *Ultrasound Med Biol*.
45:1509-1536.
Blanton, M.G., J.J. Lo Turco, and A.R. Kriegstein. 1989. Whole cell recording from neurons in
slices of reptilian and mammalian cerebral cortex. *J Neurosci Methods*. 30:203-210.
Brickley, S.G., M.I. Aller, C. Sandu, E.L. Veale, F.G. Alder, H. Sambi, A. Mathie, and W. Wisden.
2007. TASK-3 two-pore domain potassium channels enable sustained high-frequency
firing in cerebellar granule neurons. *J Neurosci*. 27:9329-9340.
Brohawn, S.G., Z. Su, and R. MacKinnon. 2014. Mechanosensitivity is mediated directly by the
lipid membrane in TRAAK and TREK1 K+ channels. *Proc Natl Acad Sci U S A*. 111:3614-
3619.
Brohawn, S.G., W. Wang, A. Handler, E.B. Campbell, J.R. Schwarz, and R. MacKinnon. 2019. The
mechanosensitive ion channel TRAAK is localized to the mammalian node of Ranvier.
Elife. 8.
Bystritsky, A., A.S. Korb, P.K. Douglas, M.S. Cohen, W.P. Melega, A.P. Mulgaonkar, A. DeSalles,
B.K. Min, and S.S. Yoo. 2011. A review of low-intensity focused ultrasound pulsation.
Brain Stimul. 4:125-136.
Calabrese, B., I.V. Tabarean, P. Juranka, and C.E. Morris. 2002. Mechanosensitivity of N-type
calcium channel currents. *Biophys J*. 83:2560-2574.
Calhoun, M.A., S.A. Bentil, E. Elliott, J.J. Otero, J.O. Winter, and R.B. Dupaix. 2019. Beyond linear
elastic modulus: viscoelastic models for brain and brain mimetic hydrogels. *ACS
Biomaterials Science & Engineering*. 5:3964-3973.
Castaneda-Castellanos, D.R., A.C. Flint, and A.R. Kriegstein. 2006. Blind patch clamp recordings
in embryonic and adult mammalian brain slices. *Nat Protoc*. 1:532-542.
Colucci, V., G. Strichartz, F. Jolesz, N. Vykhodtseva, and K. Hynynen. 2009. Focused ultrasound
effects on nerve action potential in vitro. *Ultrasound Med Biol*. 35:1737-1747.
Company, C.R. 1965. Handbook of Chemistry and Physics. 45th.ed. Chemical Rubber Compnay,
Cleveland, OH.
Cotero, V., Y. Fan, T. Tsaava, A.M. Kressel, I. Hancu, P. Fitzgerald, K. Wallace, S. Kaanumalle, J.
Graf, W. Rigby, T.J. Kao, J. Roberts, C. Bhushan, S. Joel, T.R. Coleman, S. Zanos, K.J.
Tracey, J. Ashe, S.S. Chavan, and C. Puleo. 2019. Noninvasive sub-organ ultrasound
stimulation for targeted neuromodulation. *Nat Commun*. 10:952.

Darrow, D.P., P. O'Brien, T.J. Richner, T.I. Netoff, and E.S. Ebbini. 2019. Reversible
neuroinhibition by focused ultrasound is mediated by a thermal mechanism. *Brain*
Stimul. 12:1439-1447.
de Korte, C.L., A.F. van der Steen, and J.M. Thijssen. 1994. Acoustic velocity and attenuation of
eye tissues at 20 MHz. *Ultrasound Med Biol*. 20:471-480.
Downs, M.E., S.A. Lee, G. Yang, S. Kim, Q. Wang, and E.E. Konofagou. 2018. Non-invasive
peripheral nerve stimulation via focused ultrasound in vivo. *Phys Med Biol*. 63:035011.
Duck, F.A. 1998. Radiation pressure and acoustic streaming. *In* Ultrasound in Medicine. F. Duck,
A. Baker and H. Starritt, editors. Institute of Physics Publishing, Philadelphia, PA.
Duke, A.R., M.W. Jenkins, H. Lu, J.M. McManus, H.J. Chiel, and E.D. Jansen. 2013. Transient and
selective suppression of neural activity with infrared light. *Sci Rep*. 3:2600.
Evans, E.A., R. Waugh, and L. Melnik. 1976. Elastic area compressibility modulus of red cell
membrane. *Biophys J*. 16:585-595.
Fomenko, A., C. Neudorfer, R.F. Dallapiazza, S.K. Kalia, and A.M. Lozano. 2018. Low-intensity
ultrasound neuromodulation: An overview of mechanisms and emerging human
applications. *Brain Stimul*. 11:1209-1217.
Fry, F.J., H.W. Ades, and W.J. Fry. 1958. Production of reversible changes in the central nervous
system by ultrasound. *Science*. 127:83-84.
Gaur, U., and B. Wunderlich. 1982. Heat-Capacity and Other Thermodynamic Properties of
Linear Macromolecules .5. Polystyrene. *J Phys Chem Ref Data*. 11:313-325.
Gavrilov, L.R., E.M. Tsirulnikov, and I.A. Davies. 1996. Application of focused ultrasound for the
stimulation of neural structures. *Ultrasound Med Biol*. 22:179-192.
Giese, K.P., J.F. Storm, D. Reuter, N.B. Fedorov, L.R. Shao, T. Leicher, O. Pongs, and A.J. Silva.
1998. Reduced K+ channel inactivation, spike broadening, and after-hyperpolarization in
Kvbeta1.1-deficient mice with impaired learning. *Learn Mem*. 5:257-273.
Gonzalez, J.A., L.T. Jensen, S.E. Doyle, M. Miranda-Anaya, M. Menaker, L. Fugger, D.A. Bayliss,
and D. Burdakov. 2009. Deletion of TASK1 and TASK3 channels disrupts intrinsic
excitability but does not abolish glucose or pH responses of orexin/hypocretin neurons.
Eur J Neurosci. 30:57-64.
Gu, N., K. Vervaeke, and J.F. Storm. 2007. BK potassium channels facilitate high-frequency firing
and cause early spike frequency adaptation in rat CA1 hippocampal pyramidal cells. *J*
Physiol. 580:859-882.
Hand, J.W. 1998. Ultrasound hyperthermia and the prediction of heating. *In* Ultrasound in
Medicine. F. Duck, A. Baker and H. Starritt, editors. Institute of Physics Publishing,
Philadelphia, PA.
Harper, C.A. 2006. Handbook of plastics technologies: the complete guide to properties and
performance. *In*. McGraw-Hill, New York, NY.
Hille, B. 2001. Ion Channels of Excitable Membranes. 3rd
.ed. Sinauer Associates, Inc., Sunderland, MA.
Kanda, H., J. Ling, S. Tonomura, K. Noguchi, S. Matalon, and J.G. Gu. 2019. TREK-1 and TRAAK
Are Principal K(+) Channels at the Nodes of Ranvier for Rapid Action Potential
Conduction on Mammalian Myelinated Afferent Nerves. *Neuron*. 104:960-971 e967.
Kang, D., C. Choe, and D. Kim. 2005. Thermosensitivity of the two-pore domain K+ channels
TREK-2 and TRAAK. *J Physiol*. 564:103-116.

Khaliq, Z.M., and B.P. Bean. 2010. Pacemaking in dopaminergic ventral tegmental area neurons:
depolarizing drive from background and voltage-dependent sodium conductances. *J*
Neurosci. 30:7401-7413.
Kim, J., D.S. Wei, and D.A. Hoffman. 2005. Kv4 potassium channel subunits control action
potential repolarization and frequency-dependent broadening in rat hippocampal CA1
pyramidal neurones. *J Physiol*. 569:41-57.
King, R.L., J.R. Brown, W.T. Newsome, and K.B. Pauly. 2013. Effective parameters for
ultrasound-induced in vivo neurostimulation. *Ultrasound Med Biol*. 39:312-331.
Kloda, A., L. Lua, R. Hall, D.J. Adams, and B. Martinac. 2007. Liposome reconstitution and
modulation of recombinant N-methyl-D-aspartate receptor channels by membrane
stretch. *Proc Natl Acad Sci U S A*. 104:1540-1545.
Kole, M.H. 2011. First node of Ranvier facilitates high-frequency burst encoding. *Neuron*.
71:671-682.
Krasovitski, B., V. Frenkel, S. Shoham, and E. Kimmel. 2011. Intramembrane cavitation as a
unifying mechanism for ultrasound-induced bioeffects. *Proc Natl Acad Sci U S A*.
108:3258-3263.
Kubanek, J., J. Shi, J. Marsh, D. Chen, C. Deng, and J. Cui. 2016. Ultrasound modulates ion
channel currents. *Sci Rep*. 6:24170.
Kwok, R., and E. Evans. 1981. Thermoelasticity of large lecithin bilayer vesicles. *Biophys J*.
35:637-652.
Laitko, U., and C.E. Morris. 2004. Membrane tension accelerates rate-limiting voltage-
dependent activation and slow inactivation steps in a Shaker channel. *J Gen Physiol*.
123:135-154.
Lee, S.A., H.A.S. Kamimura, M.T. Burgess, and E.E. Konofagou. 2020. Displacement imaging for
focused ultrasound peripheral nerve neuromodulation. *IEEE Trans Med Imaging*.
Leighton, T.G. 1998. An introduction to acoustic cavitation. *In* Ultrasound in Medicine. F. Duck,
A. Baker and H. Starritt, editors. Institute of Physics Publishing, Philadelphia, PA.
Leterrier, C. 2018. The Axon Initial Segment: An Updated Viewpoint. *J Neurosci*. 38:2135-2145.
Lien, C.C., and P. Jonas. 2003. Kv3 potassium conductance is necessary and kinetically optimized
for high-frequency action potential generation in hippocampal interneurons. *J Neurosci*.
23:2058-2068.
Liu, P.W., and B.P. Bean. 2014. Kv2 channel regulation of action potential repolarization and
firing patterns in superior cervical ganglion neurons and hippocampal CA1 pyramidal
neurons. *J Neurosci*. 34:4991-5002.
Lothet, E.H., K.M. Shaw, H. Lu, J. Zhuo, Y.T. Wang, S. Gu, D.B. Stolz, E.D. Jansen, C.C. Horn, H.J.
Chiel, and M.W. Jenkins. 2017. Selective inhibition of small-diameter axons using
infrared light. *Sci Rep*. 7:3275.
MacKenzie, G., N.P. Franks, and S.G. Brickley. 2015. Two-pore domain potassium channels
enable action potential generation in the absence of voltage-gated potassium channels.
Pflugers Arch. 467:989-999.
Madison, D.V., and R.A. Nicoll. 1984. Control of the repetitive discharge of rat CA 1 pyramidal
neurones in vitro. *J Physiol*. 354:319-331.

Maingret, F., I. Lauritzen, A.J. Patel, C. Heurteaux, R. Reyes, F. Lesage, M. Lazdunski, and E.
Honore. 2000. TREK-1 is a heat-activated background K(+) channel. *EMBO J*. 19:2483-
2491.
Malinow, R., and R.W. Tsien. 1990. Presynaptic enhancement shown by whole-cell recordings
of long-term potentiation in hippocampal slices. *Nature*. 346:177-180.
Marinc, C., C. Derst, H. Pruss, and R.W. Veh. 2014. Immunocytochemical localization of TASK-3
protein (K2P9.1) in the rat brain. *Cell Mol Neurobiol*. 34:61-70.
Menz, M.D., O. Oralkan, P.T. Khuri-Yakub, and S.A. Baccus. 2013. Precise neural stimulation in
the retina using focused ultrasound. *J Neurosci*. 33:4550-4560.
Menz, M.D., P. Ye, K. Firouzi, A. Nikoozadeh, K.B. Pauly, P. Khuri-Yakub, and S.A. Baccus. 2019.
Radiation Force as a Physical Mechanism for Ultrasonic Neurostimulation of the Ex Vivo
Retina. *J Neurosci*. 39:6251-6264.
Mihran, R.T., F.S. Barnes, and H. Wachtel. 1990. Temporally-specific modification of myelinated
axon excitability in vitro following a single ultrasound pulse. *Ultrasound Med Biol*.
16:297-309.
Min, B.K., A. Bystritsky, K.I. Jung, K. Fischer, Y. Zhang, L.S. Maeng, S.I. Park, Y.A. Chung, F.A.
Jolesz, and S.S. Yoo. 2011. Focused ultrasound-mediated suppression of chemically-
induced acute epileptic EEG activity. *BMC Neurosci*. 12:23.
Mizrahi, N., E.H. Zhou, G. Lenormand, R. Krishnan, D. Weihs, J.P. Butler, D.A. Weitz, J.J.
Fredberg, and E. Kimmel. 2012. Low intensity ultrasound perturbs cytoskeleton
dynamics. *Soft Matter*. 8:2438-2443.
Morris, C.E., and P.F. Juranka. 2007. Nav channel mechanosensitivity: activation and
inactivation accelerate reversibly with stretch. *Biophys J*. 93:822-833.
Needham, D., and R.S. Nunn. 1990. Elastic deformation and failure of lipid bilayer membranes
containing cholesterol. *Biophys J*. 58:997-1009.
Oh, S.J., J.M. Lee, H.B. Kim, J. Lee, S. Han, J.Y. Bae, G.S. Hong, W. Koh, J. Kwon, E.S. Hwang, D.H.
Woo, I. Youn, I.J. Cho, Y.C. Bae, S. Lee, J.W. Shim, J.H. Park, and C.J. Lee. 2020. Ultrasonic
Neuromodulation via Astrocytic TRPA1. *Curr Biol*. 30:948.
Owen, S.F., M.H. Liu, and A.C. Kreitzer. 2019. Thermal constraints on in vivo optogenetic
manipulations. *Nat Neurosci*. 22:1061-1065.
Pan, Z., T. Kao, Z. Horvath, J. Lemos, J.Y. Sul, S.D. Cranstoun, V. Bennett, S.S. Scherer, and E.C.
Cooper. 2006. A common ankyrin-G-based mechanism retains KCNQ and NaV channels
at electrically active domains of the axon. *J Neurosci*. 26:2599-2613.
Paris, L., I. Marc, B. Charlot, M. Dumas, J. Valmier, and F. Bardin. 2017. Millisecond infrared
laser pulses depolarize and elicit action potentials on in-vitro dorsal root ganglion
neurons. *Biomed Opt Express*. 8:4568-4578.
Plaksin, M., E. Kimmel, and S. Shoham. 2016. Cell-Type-Selective Effects of Intramembrane
Cavitation as a Unifying Theoretical Framework for Ultrasonic Neuromodulation.
eNeuro. 3.
Prieto, M.L., K. Firouzi, B.T. Khuri-Yakub, and M. Maduke. 2018. Activation of Piezo1 but Not
NaV1.2 Channels by Ultrasound at 43 MHz. *Ultrasound Med Biol*. 44:1217-1232.
Prieto, M.L., O. Oralkan, B.T. Khuri-Yakub, and M.C. Maduke. 2013. Dynamic response of model
lipid membranes to ultrasonic radiation force. *PLoS One*. 8:e77115.

Qiu, Z., J. Guo, S. Kala, J. Zhu, Q. Xian, W. Qiu, G. Li, T. Zhu, L. Meng, R. Zhang, H.C. Chan, H.
Zheng, and L. Sun. 2019. The Mechanosensitive Ion Channel Piezo1 Significantly
Mediates In Vitro Ultrasonic Stimulation of Neurons. *iScience*. 21:448-457.
Rashid, B., M. Destrade, and M.D. Gilchrist. 2012. Mechanical characterization of brain tissue in
compression at dynamic strain rates. *J Mech Behav Biomed Mater*. 10:23-38.
Rashid, B., M. Destrade, and M.D. Gilchrist. 2013. Mechanical characterization of brain tissue in
simple shear at dynamic strain rates. *J Mech Behav Biomed Mater*. 28:71-85.
Ricca, B.L., G. Venugopalan, and D.A. Fletcher. 2013. To pull or be pulled: parsing the multiple
modes of mechanotransduction. *Curr Opin Cell Biol*. 25:558-564.
Richter, C.P., A.I. Matic, J.D. Wells, E.D. Jansen, and J.T. Walsh, Jr. 2011. Neural stimulation with
optical radiation. *Laser Photon Rev*. 5:68-80.
Sarvazyan, A.P., O.V. Rudenko, and W.L. Nyborg. 2010. Biomedical applications of radiation
force of ultrasound: historical roots and physical basis. *Ultrasound Med Biol*. 36:1379-
1394.
Schewe, M., E. Nematian-Ardestani, H. Sun, M. Musinszki, S. Cordeiro, G. Bucci, B.L. de Groot,
S.J. Tucker, M. Rapedius, and T. Baukrowitz. 2016. A Non-canonical Voltage-Sensing
Mechanism Controls Gating in K2P K(+) Channels. *Cell*. 164:937-949.
Seeger, H.M., L. Aldrovandi, A. Alessandrini, and P. Facci. 2010. Changes in single K(+) channel
behavior induced by a lipid phase transition. *Biophys J*. 99:3675-3683.
Selfridge, A.R. 1985. Approximate Material Properties in Isotropic Materials. *Ieee T Son*
Ultrason. 32:381-394.
Shah, M.M., M. Migliore, I. Valencia, E.C. Cooper, and D.A. Brown. 2008. Functional significance
of axonal Kv7 channels in hippocampal pyramidal neurons. *Proc Natl Acad Sci U S A*.
105:7869-7874.
Shao, L.R., R. Halvorsrud, L. Borg-Graham, and J.F. Storm. 1999. The role of BK-type Ca2+-
dependent K+ channels in spike broadening during repetitive firing in rat hippocampal
pyramidal cells. *J Physiol*. 521 Pt 1:135-146.
Shapiro, M.G., K. Homma, S. Villarreal, C.P. Richter, and F. Bezanilla. 2012. Infrared light excites
cells by changing their electrical capacitance. *Nat Commun*. 3:736.
Tabarean, I.V., and C.E. Morris. 2002. Membrane stretch accelerates activation and slow
inactivation in Shaker channels with S3-S4 linker deletions. *Biophys J*. 82:2982-2994.
Talley, E.M., G. Solorzano, Q. Lei, D. Kim, and D.A. Bayliss. 2001. Cns distribution of members of
the two-pore-domain (KCNK) potassium channel family. *J Neurosci*. 21:7491-7505.
Taverna, S., T. Tkatch, A.E. Metz, and M. Martina. 2005. Differential expression of TASK
channels between horizontal interneurons and pyramidal cells of rat hippocampus. *J*
Neurosci. 25:9162-9170.
Taylor, R.E. 1965. Impedance of the squid axon membrane. *J Cell Comp Physiol*. 66:21-25.
Thijssen, J.M., H.J. Mol, and M.R. Timmer. 1985. Acoustic parameters of ocular tissues.
Ultrasound Med Biol. 11:157-161.
Tsui, P.H., S.H. Wang, and C.C. Huang. 2005. In vitro effects of ultrasound with different
energies on the conduction properties of neural tissue. *Ultrasonics*. 43:560-565.
Tufail, Y., A. Matyushov, N. Baldwin, M.L. Tauchmann, J. Georges, A. Yoshihiro, S.I. Tillery, and
W.J. Tyler. 2010. Transcranial pulsed ultrasound stimulates intact brain circuits. *Neuron*.
66:681-694.

Tyler, W.J., S.W. Lani, and G.M. Hwang. 2018. Ultrasonic modulation of neural circuit activity.
Curr Opin Neurobiol. 50:222-231.
Tyler, W.J., Y. Tufail, M. Finsterwald, M.L. Tauchmann, E.J. Olson, and C. Majestic. 2008. Remote
excitation of neuronal circuits using low-intensity, low-frequency ultrasound. *PLoS One*.
3:e3511.
Walsh, A.J., G.P. Tolstykh, S. Martens, B.L. Ibey, and H.T. Beier. 2016. Action potential block in
neurons by infrared light. *Neurophotonics*. 3:040501.
Wang, J., and O.P. Hamill. 2020. Piezo2, a pressure sensitive channel is expressed in select
neurons of
the mouse brain: a putative mechanism for synchronizing neural
networks by transducing intracranial pressure pulses. *bioRxiv*.
Wells, J., C. Kao, P. Konrad, T. Milner, J. Kim, A. Mahadevan-Jansen, and E.D. Jansen. 2007.
Biophysical mechanisms of transient optical stimulation of peripheral nerve. *Biophys J*.
93:2567-2580.
White, S.H., and T.E. Thompson. 1973. Capacitance, area, and thickness variations in thin lipid
films. *Biochim Biophys Acta*. 323:7-22.
Wu, J., and W.L. Nyborg. 2008. Ultrasound, cavitation bubbles and their interaction with cells.
Adv Drug Deliv Rev. 60:1103-1116.
Yamada-Hanff, J., and B.P. Bean. 2015. Activation of Ih and TTX-sensitive sodium current at
subthreshold voltages during CA1 pyramidal neuron firing. *J Neurophysiol*. 114:2376-
2389.
Ye, P.P., J.R. Brown, and K.B. Pauly. 2016. Frequency Dependence of Ultrasound
Neurostimulation in the Mouse Brain. *Ultrasound Med Biol*. 42:1512-1530.
Yoo, S., D.R. Mittelstein, R. Hurt, J. Lacroix, and M.G. Shapiro. 2020. Focused ultrasound excites
neurons via mechanosensitive calcium accumulation and ion channel amplification.
bioRxiv.
Young, R.R., and E. Henneman. 1961. Functional effects of focused ultrasound on mammalian
nerves. *Science*. 134:1521-1522.
Yue, C., and Y. Yaari. 2004. KCNQ/M channels control spike afterdepolarization and burst
generation in hippocampal neurons. *J Neurosci*. 24:4614-4624.
Zachs, D.P., S.J. Offutt, R.S. Graham, Y. Kim, J. Mueller, J.L. Auger, N.J. Schuldt, C.R.W. Kaiser,
A.P. Heiller, R. Dutta, H. Guo, J.K. Alford, B.A. Binstadt, and H.H. Lim. 2019. Noninvasive
ultrasound stimulation of the spleen to treat inflammatory arthritis. *Nat Commun*.
10:951.
Zhou, D., S. Lambert, P.L. Malen, S. Carpenter, L.M. Boland, and V. Bennett. 1998. AnkyrinG is
required for clustering of voltage-gated Na channels at axon initial segments and for
normal action potential firing. *J Cell Biol*. 143:1295-1304.
Zhu, X., J.W. Lin, and M.Y. Sander. 2019. Infrared inhibition and waveform modulation of action
potentials in the crayfish motor axon. *Biomed Opt Express*. 10:6580-6594.
1204

1205

1206 **SUPPLEMENTAL MATERIALS**

1207

Supplemental Figure 1. Effects of ultrasound at different intensities on firing rate. A. Mean (±

SEM) action potential firing rate with (*red*) or without (*blue*) a 1-s ultrasound pulse at various

intensities starting 200 ms after the start of a 2-s current step, as measured during the period of

overlap between the ultrasound and current stimuli, or during the same time period in the

absence of ultrasound. N = 4 cells, except for 6 W/cm^2, where N = 3 cells. **B.** As in panel A, but

showing the difference in firing rate between the ultrasound and control conditions.

1214

Supplemental Figure 2. An ATP-regenerating internal solution stabilizes the response to

ultrasound. A-B. Mean (±SEM) spike rates during the first 500 ms of a current step in the

presence (*red circles*) and absence (*blue circles*) of a 1-s ultrasound application starting 500 ms

before the current step, as a function of time relative to break-in (establishment of whole-cell

recording configuration). The internal solution contained 10 Na-phosphocreatine with (A) or

without (B) 50 U/mL creatine phosphokinase to provide an ATP-regenerating system. Spike

rates were measured at various time points between 0 and 10, 10 and 20, 20 and 30, and 30

and 40 minutes after break-in. The x-values represent the mean start time for the protocol to

measure spike rates (which comprised 2 minutes of recording time) with horizontal error bars

(in some cases smaller than the symbol size) representing the SEM. The amplitude of the

current step was adjusted over time to maintain spiking behavior as close as possible to that at

the start of the experiment. **C.** Mean (±SEM) difference in spike rate between the control and

ultrasound conditions for measurements at 1-10 minutes after break-in and 30-40 minutes

after break-in, with (*left*) or without (*right*) the ATP-regenerating system. The difference was

only statistically significant without the ATP-regenerating system (P = 0.11, with; and P =

0.0064, without). **D-F.** Same as A-B, but for latency to the first action potential following the

start of the current step. N = 10 cells with and N = 6 cells without the ATP-regenerating system.

The difference was not statistically significant for either group (P = 0.21, with; and P = 0.089,

without).

1234

1235

Supplemental Figure 3. Effects of ultrasound on action potential height. A-C. Mean (±SEM)

action potential heights as a function of action potential number in the presence (*red*) and

absence (*blue*) of a 1-s ultrasound pulse starting 500 ms before the current step, for cells firing

at an average firing rates (as measured during the first 500 ms of the current step) of

approximately 5 Hz (N = 13), 10 Hz (N = 15), and 20 Hz (N = 13) in the control condition. **D.** As in

A-C, but with ultrasound applied 1 s after the start of a 3-s current step, for cells firing at an

average firing rate of approximately 5 Hz in the control condition, with firing rate determined in

a 1-s window starting 1 s after the current step (corresponding to the time period of the

ultrasound stimulus), and action potential number relative to the start of the ultrasound

stimulus (N = 6). Note that the y-axes do not begin at zero.

1246

Supplemental Figure 4. Effects of ultrasound on action potential firing and waveform at near-physiological temperature. A. Example voltage traces showing inhibition of action potential firing by ultrasound at 30°C. The response to a 300-pA current step is shown with (*red voltage trace*) and without (*blue voltage trace*) a 1-s ultrasound pulse (*red bar*) applied 500 ms before the start of the current step. The *dashed line* indicates the resting membrane voltage. **B.** Mean (± SEM, N =3) spike frequency during the first 500 ms of the current step (corresponding to the period of overlap between the current and ultrasound stimuli, or the equivalent time period in the absence of ultrasound) for the protocol shown in panel A for the control (*blue*) and ultrasound (*red*) conditions, for cells firing at an average spike frequency of approximately 5 Hz in the absence of ultrasound. Data points for the individual cells are shown connected by *dashed lines*. (Compare Figure 1E.) **C.** Latency between the start of the current step and the first action potential, and between the first and second, and second and third action potentials, for the control (*blue*) and ultrasound (*red*) conditions for three individual cells. (Compare Figure

1261   4A.) **D.** As in panel C, but for effects of ultrasound on action potential width. (Compare Figure
1262   6A.)