Spike-frequency dependent inhibition and potentiation of neural activity by ultrasound

Martin Loynaz Prieto[1*], Kamyar Firouzi[2], Butrus T. Khuri-Yakub[2], Daniel V. Madison[1], and Merritt Maduke[1*]

[1]Department of Molecular and Cellular Physiology, Stanford University
[2]E. L. Ginzton Laboratory, Stanford University

*Corresponding authors

Condensed title: Spike-Rate-Dependent Neuromodulation by Ultrasound

SUMMARY Prieto et al. describe how ultrasound can either inhibit or potentiate action potential firing in hippocampal pyramidal neurons and demonstrate that these effects can be explained by increased potassium conductance.

ABSTRACT

Ultrasound can modulate action potential firing *in vivo* and *in vitro*, but the mechanistic basis of this phenomenon is not well understood. To address this problem, we used patch-clamp recording to quantify the effects of focused, high-frequency (43 MHz) ultrasound on evoked action potential firing in CA1 pyramidal neurons in acute rodent hippocampal brain slices. We find that ultrasound can either inhibit or potentiate firing in a spike-frequency-dependent manner: at low (near-threshold) input currents and low firing frequencies, ultrasound inhibits firing, while at higher input currents and higher firing frequencies, ultrasound potentiates firing. The net result of these two competing effects is that ultrasound increases the threshold current for action potential firing, the slope of frequency-input curves, and the maximum firing frequency. In addition, ultrasound slightly hyperpolarizes the resting membrane potential, decreases action potential width, and increases the depth of the afterhyperpolarization. All of these results can be explained by the hypothesis that ultrasound activates a sustained potassium conductance. According to this hypothesis, increased outward potassium currents hyperpolarize the resting membrane potential and inhibit firing at near-threshold input currents, but potentiate firing in response to higher input currents by limiting inactivation of voltage-dependent sodium channels during the action potential. This latter effect is a consequence of faster action-potential repolarization, which limits inactivation of voltage-dependent sodium channels, and deeper (more negative) afterhyperpolarization, which increases the rate of recovery from inactivation. Based on these results we propose that ultrasound activates thermosensitive and mechanosensitive, voltage-insensitive two-pore-domain potassium (K2P) channels, through heating or mechanical effects of acoustic radiation force. Finite-element modelling of the effects of ultrasound on brain tissue suggests that the effects of ultrasound on firing frequency are caused by a small (less than 2°C) increase in temperature, with possible additional contributions from mechanical effects.

INTRODUCTION

Ultrasound can non-invasively modulate action potential activity in neurons *in vivo* and *in vitro*,

with improved depth penetration and spatial resolution relative to other non-invasive

neuromodulation modalities, and it may therefore become an important new technology in basic

and clinical neuroscience (Fry et al., 1958; Gavrilov et al., 1996; Tufail et al., 2010; Bystritsky et

al., 2011; Fomenko et al., 2018; Blackmore et al., 2019). Investigation of this phenomenon has

predominantly focused on low-frequency ultrasound (defined here as less than 3 MHz, although

there is no firmly defined boundary between "high" and "low" frequency in the neuromodulation

field), but higher ultrasound frequencies have also been shown to modulate action potential firing

in vitro (Menz et al., 2013; Menz et al., 2019) and to directly modulate ion channel activity in

heterologous systems (Kubanek et al., 2016; Prieto et al., 2018). The focus on lower frequency

ultrasound is understandable, since envisioned clinical applications involving transcranial focused

ultrasound have been a primary motivation for research on ultrasound neuromodulation, and loss

of ultrasound power due to attenuation in the skull limits these applications to low frequency

ultrasound. For applications in which transmission through the skull does not impose limits on

frequency, such as *in vitro* studies, neuromodulation in the peripheral nervous system (Downs et

al., 2018; Cotero et al., 2019; Zachs et al., 2019), neuromodulation using subcranial implants, or

neuromodulation in experimental animal model systems involving craniotomies or acoustically

transparent cranial windows, high frequencies have a distinct advantage in terms of the greater

spatial resolution that can be achieved.

These applications motivate investigation of the fundamental physical, cellular, and

molecular mechanisms underlying neuromodulation by high-frequency ultrasound, which are all

not well understood. It remains an open question to what extent these mechanisms overlap for

high- and low-frequency ultrasound neuromodulation. In terms of the basic physical mechanism

by which acoustic energy is transduced into effects on biological tissue, most proposed

mechanisms for ultrasound neuromodulation involve either heating due to absorption of acoustic

energy (Hand, 1998), mechanical effects of acoustic radiation force (Duck, 1998; Sarvazyan et

al., 2010), or effects of cavitation (the nucleation, growth, oscillation, and sometimes collapse, of

microscopic gas bubbles) (Leighton, 1998; Wu and Nyborg, 2008; Krasovitski et al., 2011; Plaksin

et al., 2016). Of these, the first two increase with acoustic frequency, while the probability of

cavitation decreases with acoustic frequency. There are also many unanswered questions at the

cellular level. Both excitatory and inhibitory effects of ultrasound have been observed using direct

or indirect measures of neural activity at the population level (Bystritsky et al., 2011; Blackmore

et al., 2019), but it is unclear whether the direct effect of ultrasound at the cellular level is excitatory

or inhibitory. Of course, the answer to this question could depend on any number of possible

relevant biological and experimental variables, such as species, tissue, specific neural subtype,

ultrasound stimulus parameters, or whether effects on intrinsic or evoked activity are measured.

For example, a cellular-level excitatory effect, specific to GABAergic interneurons, could produce

an inhibitory effect at the population level. This leads to the question of whether certain

subpopulations of neurons are more sensitive to ultrasound than others, and if so, what are the

specific molecular mechanisms underlying the differences in sensitivity. Do certain ion channels

respond directly to ultrasound? What biophysical properties might account for the sensitivity of

these channels to ultrasound, and how might cell-type specific differences in the density and

localization of these channels, and the way in which they interact with other ion channels to

regulate action potential firing, produce differences in the response to ultrasound?

One reason there are so many outstanding questions regarding ultrasound

neuromodulation is that patch-clamp recordings of the effects of ultrasound on action potential

firing in neurons have been unavailable. At low ultrasound frequencies, we (Prieto et al., 2018)

and others (Tyler et al., 2008) have found that patch-clamp seals are extremely unstable in the

presence of ultrasound at low frequencies, precluding detailed, mechanistic studies of ultrasound

neuromodulation with this technique, which provides quantitative information on action potential

timing and dynamics unobtainable with other techniques. However, we have previously shown

that stable patch-clamp recordings can be achieved using ultrasound at the frequency of 43 MHz

(Prieto et al., 2018). Here, we use patch-clamp recording to measure the effects of ultrasound at

43 MHz and 50 W/cm^2 on action-potential firing in response to injected current in pyramidal

neurons of the CA1 layer of the hippocampus in acute rodent brain slices. We find that ultrasound

has a bidirectional, spike-frequency dependent effect on excitability, and that this and other

observed neurophysiological effects of ultrasound can be explained by activation by ultrasound

of a steady K$^+$ current, resembling that of two pore domain potassium channels (K2P channels).

MATERIALS AND METHODS

Slice preparation. Brain slices were prepared from male Sprague-Dawley rats, 35-50 days old.

Rats were anesthetized with isofluorane and decapitated, and the brain was immediately removed

and placed in ice-cold artificial cerebral spinal fluid (ACSF), bubbled with 95% O_2 and 5% CO_2.

The hippocampus was dissected out and placed on the slicing apparatus, consisting of a manual

micrometer and a gravity-driven vertical slicing mechanism, with the CA1 layer oriented

approximately parallel to the slicing blade. Slices (500 microns thick) were prepared and then

stored in a humidified chamber with an atmosphere of 95% O_2/5% CO_2, resting on a square of

filter paper placed on a dish of ACSF. Slices were used within 1-6 hours of slice preparation.

Animals were handled in accordance with protocols approved by Stanford University's Institutional

Animal Care and Use Committee.

Ultrasound. Continuous wave ultrasound at 43 MHz and 50 W/cm^2 was applied to brain slices

using a set-up similar to that we previously used for our experiments on cultured cells (Prieto et

al., 2018), except that the tissue was visualized with a dissecting microscope at low magnification.

The bottom of the experimental chamber was a 25-micron film of polystyrene, plasma-treated with

a Harrick plasma cleaner (Harrick Plasma, Ithaca, NY). Ultrasound was transmitted from below

(through the polystyrene film) with the sound beam perpendicular to the bottom of the chamber.

The 43 MHz transducer was a custom-built device, calibrated as described previously (Prieto et

al., 2013), excited using an ENI 403LA (37 dB) amplifier (ENI, Rochester, NY). The focal volume

of the transducer is approximately a cylinder 90 microns in diameter by 500 microns long, and the

focal distance is approximately 4.2 mm. The set-up was based on the stage from a Zeiss

Axioskop-2 microscope (Zeiss Microscopes, Jena, Germany), with the housing for the sub-stage

condenser modified to accommodate the transducer, such that the position of the transducer

could be adjusted using the controls for alignment of the condenser, and the position of the tissue

sample relative to the transducer could be adjusted with the microscope stage. The transducer

was coupled to the polystyrene film at the bottom of the experimental chamber using a small

volume of distilled water held in place by a rubber O-ring attached to the tip of the transducer with

silicone grease. The focal volume of the transducer was aligned along the z-axis using a pulse-

echo protocol, adjusting the height of the transducer to maximize the echo signal from the bottom

of the empty chamber. The focus was aligned in the x-y plane by adding to the chamber a small

volume of ACSF, barely sufficient to cover the bottom of the chamber, such that a thin layer of

solution was spread over the bottom of the chamber. Ultrasound pulses, one second in length,

were then applied, raising a mound of fluid at the focus of the transducer (due to the radiation

force produced by reflection of the acoustic wave at the interface between the solution and the air

above it (Duck, 1998)). The mound of fluid was then aligned in the x-y plane to the center of a

reticle in one eyepiece of the dissecting microscope, and, after adding additional ACSF and the

tissue sample to the chamber, the center of the reticle was aligned with the region of the tissue

targeted for patch-clamp recording. The ultrasound intensity (50 W/cm^2) is the spatial peak, pulse

average intensity for the free field. The interval between ultrasound applications was at least 12

seconds.

Electrophysiology. Current clamp recordings were performed using an Axon Instruments

Axoclamp-2B amplifier operating in "Bridge" mode and Digidata 1330A digitizer with pClamp

software (Molecular Devices, Sunnyvale, CA), except for the preliminary experiments in

Supplemental Figure 1, which were performed with an Axon Instruments Axopatch 200B amplifier

(Molecular Devices, Sunnyvale, CA). Patch-clamp recording was performed using a "blind-patch"

approach (Blanton et al., 1989; Malinow and Tsien, 1990; Castaneda-Castellanos et al., 2006), in

which the recording pipette was positioned above the CA1 layer of the hippocampus, as identified

visually at low magnification, and then slowly lowered into the tissue while applying positive

pressure to the pipette and monitoring the pipette tip resistance in voltage-clamp mode. In the

blind-patch approach, a small decrease in tip resistance is used to indicate possible contact of

the pipette tip with a neuron in the absence of the usual visual cues. Typically, the first two

instances of possible cell contact were not used, to avoid patching on cells at the surface of the

tissue that may have been damaged during the slicing procedure. Gigaseals and the subsequent

whole-cell recording configuration were obtained following the standard procedure in voltage-

clamp mode before switching to current-clamp mode. In most experiments, slices were held in

place with a Warner Instruments RC-22 slice anchor (Harvard Bioscience, Hamden, CT); the

experiments in Supplemental Figure 1 and some of the experiments in Figure 1E were performed

without a slice anchor. No obvious effects of the slice anchor on the ultrasound response were

noted. Series resistance, monitored and compensated throughout the recording, was between 30

and 100 M• . All of the neurons used for experiments could be unambiguously identified as

pyramidal cells by their distinct adaptive firing patterns in response to 2-s current steps. Current

records were low-pass filtered at 10 kHz and sampled at 100 kHz. Brain slices were continuously

perfused with ACSF (in mM: 119 NaCl, 2.5 KCl, 1.3 MgSO$_4$, 2.5 CaCl$_2$, 1 NaH$_2$PO$_4$, 26.2 NaHCO$_3$,

11 glucose), bubbled with 95% O$_2$/5% CO$_2$, at ~100-250 mL/hour. The internal solution was (in

mM): 126 K-gluconate, 4 KCl, 10 HEPES, 4 Mg-ATP, 0.3 Na$_2$GTP, 10 Na-phosphocreatine, 10

sucrose, and 50 U/mL creatine phosphokinase (porcine), pH 7.2 (KOH). This internal solution

contains an ATP-regenerating system (phosphocreatine and creatine phosphokinase) because

we found that the strength of the response to ultrasound was unstable, gradually declining over

the course of a recording unless the ATP-regenerating system was included (Supplemental

Figure 2). Na-phosphocreatine was obtained from Abcam and creatine phosphokinase was

obtained from EMD Millipore. All other salts and chemicals were obtained from either Sigma-

Aldrich or Fisher Scientific. (For some of the preliminary recordings shown in Figure 1E, the

creatine phosphokinase was omitted from the internal solution, or a different internal solution,

containing 120 K-gluconate, 40 HEPES, 5 $MgCl_2$, 0.3 Na_2GTP, 2 Na_2ATP, pH 7.2 (KOH) was

used. Other than the instability of the ultrasound response over time in the absence of creatine

phosphokinase, no obvious differences in recordings with different internal solutions were noted.)

Because creatine phosphokinase increases the viscosity of the solution, making it difficult to

obtain gigaseals, a small volume of internal solution without the enzyme was added to the tip of

the pipette (enough to fill approximately the first 3 mm of the tip) before back-filling the pipette

with the enzyme-containing solution. Pipettes were pulled from thick walled glass and had

resistances between 5 and 10 M• • Recordings were performed at room temperature (21-23°C).

Data analysis. Current records were analyzed in Igor Pro (Wavemetrics, Lake Oswego, OR) with

user-written procedures. Action-potential threshold was defined as the point at which the first

derivative of the voltage reached 4% of its peak value during the rising phase of the action

potential. This quantitative criterion was previously found to correspond with action-potential

thresholds as identified visually (Khaliq and Bean, 2010; Yamada-Hanff and Bean, 2015), and we

found that it also works well with our data, using phase plots to visually confirm the threshold

value. Action-potential height was defined as the difference between the action-potential peak

and the action-potential threshold voltage. Action-potential width was measured at 50% of action

potential height defined in this manner. Threshold current levels for action-potential firing were

estimated based on a series of current steps in 10-pA increments. Frequency-input plots and

action-potential parameters (height, width, latency, and interspike intervals) were determined from

the average values of at least 3 trials each for the control and ultrasound conditions. Frequency-

input trials were performed alternatingly for the control and ultrasound conditions, with the first

condition tested varying randomly on a cell-by-cell basis.

Average traces for analysis of the effects of ultrasound on membrane resting potential and

membrane capacitance were derived from at least 3 voltage traces.

Statistical significance was assessed using paired or unpaired two-tailed Student's t-tests, with P

< 0.05 defined as significant. Statistical analysis was performed in Microsoft Excel.

Finite-element simulations. Finite-element were performed in COMSOL (COMSOL Inc., Palo Alto,

CA, USA). The simulation domain had radially symmetric geometry and was 6 mm in the axial

direction. The simulation domain contained four layers of different materials: a lower layer of water

(4.2 mm thick in the axial direction), followed by a layer of polystyrene (25 microns thick), followed

by a layer of brain tissue (500 microns thick), followed by an upper layer of water (1.275 mm

thick). The width of the simulation domain in the axial direction was 1 mm (for simulation for

acoustic pressure and heating) or 5 mm (for simulation of mechanical deformation) in the axial

direction. A 940-micron diameter by 100-micron height arc on the lower axial boundary of the

simulation domain represented the quartz lens of the transducer.

Simulations of acoustic pressure, heating, and static displacement in response to radiation

force were performed as described previously (Prieto et al., 2018). For simulation of dynamic

tissue displacement in response to radiation force, the brain slice was modeled as a

incompressible, linear viscoelastic material (Calhoun et al., 2019), characterized by Young's

modulus, Poisson's ratio, and shear viscosity, loaded by the fluid layer above it. The polystyrene

was modeled as a linear elastic material, because we determined in a series of simulations that

including viscosity of the polystyrene had no effect on the tissue displacement. A time step of 0.1

ms was used for simulation of the dynamic tissue displacement. Material properties used for

water, polystyrene, and brain tissue used in the simulation and sources for these values are given

in Table 1. Additional details on mesh size, boundary conditions, and solver configurations are

available in Prieto et al. (2018).

SUMMARY OF SUPPLEMENTAL MATERIAL

Supplemental materials include three figures showing: the effect of ultrasound at different

intensities on action potential firing frequency (Supplemental Figure 1); stabilization of the

response to ultrasound by an ATP-regenerating system in the internal solution (Supplemental

Figure 2); and effects of ultrasound on action potential height (Supplemental Figure 3).

RESULTS

We measured the effects of ultrasound on action-potential firing in hippocampal CA1 pyramidal

cells using the set-up shown in Figure 1A (described in *Materials and Methods*). Throughout the

experiments reported here, ultrasound was applied at 43 MHz and 50 W/cm^2 as a 1-s, continuous-

wave pulse. In an initial exploration of the effects of intensity on the ultrasound response, we

determined that 50 W/cm^2 had a sufficiently robust effect on firing frequency to permit quantitative

investigation of this effect (Supplemental Figure 1), but we did not perform a detailed investigation

of the intensity dependence. We chose to use continuous-wave ultrasound (without additional

low-frequency modulation within the pulse) because continuous-wave ultrasound was previously

found to be optimal for ultrasound neuromodulation of retinal ganglion cells at 43 MHz (Menz et

al., 2013). With these ultrasound parameters, we found robust, reproducible inhibition of action-

potential firing by ultrasound using the protocol illustrated in Figure 1B. In these experiments, a

current-injection amplitude sufficient to induce firing at an average frequency of ~4-12 Hz during

the first 500 ms of the current step (corresponding to the overlap between the ultrasound stimulus

and current step) was used. This range of firing frequencies is physiologically relevant and

sufficient to detect either inhibition or potentiation of firing. With these experimental conditions,

we established that the response to ultrasound is highly reproducible, both on a trial-by-trial basis

within the same cell (Figure 1C-D), and between cells, with similar effects seen in over fifty cells

(Figure 1E). In addition, the response to ultrasound was stable over the course of recordings

lasting over 30 minutes (Figure 1F, Supplemental Figure 2); in a few cases where the patch seal

lasted for over 90 minutes, the ultrasound response remained stable.

Effects of ultrasound on frequency-input curves. To explore the effects of ultrasound on excitability

over a wider range of firing frequencies, we generated frequency-input (f-i) curves comparing

average firing frequencies as a function of input current in the presence and absence of

ultrasound. An example f-i curve generated with the protocol illustrated in Figure 1B is shown in

Figure 2A, along with example voltage traces in Figure 2B. The average spike frequency during

the first 500 ms of the current step was compared to the spike frequency in the same time window

in the absence of ultrasound. To compare the effects of ultrasound across neurons, we converted

the f-i curves into plots of the relative increase or decrease in firing frequency as a function of the

input current (Figure 2C-D). These data reveal two distinct regimes with contrasting inhibitory and

excitatory ultrasound effects. At relatively low input currents, near the threshold for action-

potential firing under this current stimulation protocol, ultrasound decreases the average firing

frequency; while at relatively high input currents, well above the action-potential threshold,

ultrasound increases the average firing frequency. Between these two regimes, there is a

transitional region where there is little or no effect on average firing frequency, presumably due

to the balance between competing inhibitory and excitatory effects. Other notable effects of

ultrasound on the f-i curves are an increase in the threshold current for action-potential firing, an

increase in the slope of the f-i relationship in the approximately linear region of the f-i curve, and

an increase in the maximum firing frequency in the sublinear "plateau" region of the curve (Figure

2A). The mean (± SEM) slope of linear region of the f-i curve increased from 0.108 ± 0.007 Hz/pA

in the control condition to 0.145 ± 0.012 Hz/pA in the ultrasound condition; and the mean

maximum firing frequency increased from 23 ± 1 Hz in the control condition to 30 ± 2 Hz/pA in the

ultrasound condition (N = 9).

234   Action-potential firing behavior is determined by the interaction between numerous K^+,

Na^+, and other ionic currents (Madison and Nicoll, 1984; Bean, 2007). Some of these currents are

clearly identified with specific ion channel subtypes, while the molecular identity of others is still

uncertain. Thus, f-i curves are a complicated function of the density, localization, conductance,

and kinetic properties of these channels/currents. Some currents inactivate relatively rapidly and

only influence firing frequency during the initial response to a current step, while others continue

to influence firing frequency throughout the step. To explore the molecular basis of the response

to ultrasound, we therefore generated a second set of f-i curves with ultrasound applied 1 s after

the start of a 3-s current step (Figure 3A).

243   Ultrasound also has a bidirectional, spike-frequency-dependent effect on excitability when

it was applied 1 second after the start of a current step (Figure 3). As with ultrasound applied

before the start of the current step, ultrasound applied in the middle of the current step decreased

firing frequency in the low-firing-frequency, near-threshold region of the f-i curve, and increased

spike rate in the high-firing-frequency, suprathreshold region of the curve (Figure 3B-D). However,

the excitatory effect was more pronounced than we observed when the ultrasound pulse started

500 ms before the start of the current step. Here, ultrasound potentiated firing frequency by

several hundred percent for high input currents (Figure 3D), as compared with a maximum

potentiation of 49 ± 16% at 450 pA seen with the earlier ultrasound application (Figure 2C). This

effect can be explained by the fact that prolonged injection of high amplitude currents causes

accumulation of inactivation in many voltage-gated channels, which can drive pyramidal cells into

a refractory state where spiking is infrequent and irregular or entirely absent. (For example, the

voltage trace for the control condition at +450 pA in Figure 3C shows a steep decline in action

potential height (due to voltage-gated Na_V channel inactivation) followed by a period of no action

potential activity). If neurons are in this refractory state during the ultrasound application,

ultrasound can "rescue" firing (as in the example voltage trace for the ultrasound condition at +450

pA in Figure 3C). A hint of this rescue phenomenon is also seen when the ultrasound application

starts before the current step, as seen in the abrupt increase in the potentiation effect at +450 pA

(Figure 2C).

Effects of ultrasound on interspike intervals. To examine the effects of ultrasound on action-

potential firing in more detail we compared the latency to the first spike, and the intervals between

subsequent spikes (interspike intervals), in the control and ultrasound conditions. Figure 4A

shows raster plots summarizing action potential timing for individual current steps, ranging from

0 pA to +450 pA in 50 pA intervals, for the control (*left*) and ultrasound (*right*) conditions. To

summarize these results, and to account for the variability in intrinsic excitability between cells,

we averaged *instantaneous* firing frequencies (latency and interspike intervals) across cells firing

at approximately the same *average* firing frequency (either 5, 10, or 20 Hz) in the control condition

at whatever input current was necessary to achieve these average firing frequencies, and at the

same input current in the ultrasound condition (Figure 4B-E). We note, however, that this

averaging procedure can obscure some of the details of ultrasound's effects. At 5 Hz the effect of

ultrasound applied 500 ms before the start of the current step is predominantly inhibitory (as seen

in the longer average latency and interspike intervals in the ultrasound as compared with the

control condition (Figure 4B), but the interval between the first and second spikes was actually

shorter in the ultrasound condition than the control condition in some cells (6 out of 13 cells in this

data set). This effect occurs because, even at relatively low average firing frequencies, pyramidal

cells will occasionally fire "doublets" or high-frequency bursts of two action potentials, in which a

second action potential is triggered by the after-depolarization of the initial action potential. When

this occurs, ultrasound decreases the interval between spikes (Figure 4E). This result indicates

that the mechanism by which ultrasound potentiates firing at high average firing frequencies is

also active under conditions of low overall average firing frequency, during localized periods of

high-frequency firing. A similar combination of inhibitory and excitatory effects can be observed

at 10 Hz and even 20 Hz (Figure 4C-D). At 20 Hz, ultrasound still increased the latency to the first

spike (16 ± 2 ms versus 12 ± 1 ms in the control condition; N = 12, P = 5.2 x 10^{-4}, paired, two-

tailed Student's t-test), despite decreasing the interspike interval for all subsequent spikes (Figure

4D). The effect of ultrasound was also mixed when ultrasound was applied 1 s after the start of

the current step (Figure 4F).

Effects of ultrasound on resting membrane potential. Ultrasound also has effects on resting

membrane potential, which can be observed by averaging several voltage traces aligned to the

onset of the ultrasound pulse, in the absence of injected current. As shown in Figure 5A,

ultrasound has a slight hyperpolarizing effect on resting membrane potential. The average voltage

traces also show another interesting effect of ultrasound. In addition to the relatively constant

hyperpolarization, there is a transient *depolarization* of the resting membrane potential, preceding

the hyperpolarization effect and acting on a faster time scale, at the onset of the ultrasound pulse;

this transient depolarization is matched by a roughly symmetrical transient hyperpolarization at

the offset of the pulse (Figure 5A, *arrows*). The symmetrical, on/off nature of these transients

suggests that they are caused by changes in membrane capacitance, as does the fact that they

occur much faster than the steady-state changes in resting membrane potential (Figure 5A-C).

(Changes in membrane potential due to changes in capacitance can occur much faster than those

due to ionic currents because they do not involve actual redistribution of charges across the

membrane and are therefore not limited by membrane conductance). Thus, the results in Figure

5 can be described by three distinct steps: 1) ultrasound rapidly increases membrane capacitance

(C) at the onset of the ultrasound pulse, which causes the membrane voltage (V) to become less

negative (due to an increase in the denominator in the equation V=Q/C, where Q is the negative

total charge on the membrane); 2) ionic currents then slowly change the membrane voltage to a

steady-state value determined by the total ionic current (one or more ion channels having been

affected by ultrasound) resulting in membrane hyperpolarization; 3) at the offset of the ultrasound

pulse, capacitance rapidly relaxes back to its initial value, producing a transient decrease in

membrane voltage, through essentially the same mechanism as in step 1. We investigate the

physical basis of these capacitance changes and their relationship to the effects of ultrasound on

excitability further below, but one point worth mentioning here is that the change in capacitance

is too small for its effect on the rate of membrane charging (less than 1% change in membrane

time constant) to have a significant effect on excitability in and of itself.

The K2P channel hypothesis. What ion channels might be responsible for the effects of ultrasound

on excitability and resting membrane potential? A compelling hypothesis—able to explain all of

our data—is that ultrasound activates a non-inactivating, voltage-insensitive potassium channel,

such as members of the K2P potassium channel family. A primary reason for suspecting such

channels is that the effects of ultrasound are similar regardless of whether ultrasound is presented

500 ms before or 1 s after the start of the current step (compare Figures 2 and 3), consistent with

the idea that ultrasound affects a channel whose activation/inactivation status does not change

during that time period. Related to this point, the effects of ultrasound are not diminished by

repetitive, high-frequency action potential firing (for example, in Figure 4D effects of ultrasound

are clearly present throughout the entire 20-Hz, 10-spike train). Further, a striking feature of the

effects of ultrasound on spike intervals is that ultrasound always increases the latency to the first

spike (Figure 4B-D), regardless of the input current or intrinsic spike frequency. This observation

is also consistent with the idea that ultrasound affects a voltage-insensitive channel, whose

activation is not modulated by input current or spike frequency.

CA1 pyramidal neurons express a variety of K2P channel subunits. Expression of TASK-

1 and TASK-3, TREK-1 and TREK-2, TRAAK, and TWIK has been shown at the mRNA level

(Talley et al., 2001), while expression at the protein level has been shown has been shown for

TASK-3 in CA1 pyramidal neurons specifically (Marinc et al., 2014), and for TRAAK throughout

the central nervous system (Brohawn et al., 2019). In addition, functional expression of TASK-like

currents has been shown in CA1 pyramidal neurons using patch-clamp recording (Taverna et al.,

2005). TREK and TRAAK channels are particularly interesting in the present context since they

are exceptionally sensitive to mechanical force and to increases in temperature between

approximately 20 and 40°C (Maingret et al., 2000; Kang et al., 2005). Thus, TREK and TRAAK

channels are responsive to the two leading candidate mechanisms by which ultrasound at 43

MHz could modulate ion channel activity.

Activation of K2P channels by ultrasound could account for all of the neurophysiological

effects of ultrasound described so far: hyperpolarization of resting membrane potential; inhibition

of action potential firing in response to near-threshold current injections; and—although this last

point may seem counter-intuitive—potentiation of action potential firing at high firing frequencies.

Hyperpolarization of resting membrane potential by increased outward K^+ current is

straightforward, as is the idea that K^+ current can inhibit firing, but K^+ current can also potentiate

firing by its effects on action potential repolarization and afterhyperpolarization (AHP). By

accelerating the rate of membrane repolarization following the peak of an action potential (thereby

reducing action potential width) K^+ current can reduce inactivation of voltage-dependent Na^+

channels during the action potential, and by increasing the depth of the AHP, it can accelerate

the voltage-dependent recovery of Na_V channels from inactivation. Both of these effects would

tend to increase the population of Na_V channels available for activation in response to depolarizing

current and would increase the maximum action potential firing frequency, as we in fact see in

response to ultrasound (Figures 2 and 3). This mechanism is well-known and widespread in

neurophysiology, with several K^+ channels, including both K2P channels and voltage   -

dependent K^+ channels (K_V channels), having been shown to facilitate high frequency firing (Lien

and Jonas, 2003; Brickley et al., 2007; Gu et al., 2007; Liu and Bean, 2014; Kanda et al., 2019).

The idea that the potentiation of firing by ultrasound is due to effects on action-potential

repolarization and AHP is supported by the results in Figure 4D. For neurons firing at an average

firing frequency of 20 Hz in the absence of ultrasound, ultrasound increases the latency to the

first spike, while it decreases the intervals between all subsequent spikes. The lack of a

potentiating effect on the first spike, despite the otherwise strongly potentiating effects of

ultrasound, indicates that the potentiating effect acts through a process (such as action-potential

repolarization and afterhyperpolarization) that occurs *after* the initiation of the first action potential.

The idea that ultrasound can potentiate firing by activating K^+ current makes specific predictions

about the effects of ultrasound on action potential waveform: ultrasound should accelerate

repolarization, decrease action potential width, and increase the depth of the

afterhyperpolarization.

Effects of ultrasound on action-potential waveform. To test the idea that potentiation of firing by

ultrasound is due to activation of K^+ channels, we examined the effects of ultrasound on action-

potential waveform in our recordings. The effects of ultrasound on action-potential waveform are

consistent with the idea that ultrasound facilitates high-frequency firing by accelerating action-

potential repolarization. Figure 6A-F shows the effect of ultrasound applied 500 ms before the

start of the current step on action-potential width for cells firing at average frequencies of 5, 10,

and 20 Hz. (Ultrasound also had effects on action-potential height, although these were less

pronounced than the effects on width; effects on height are detailed in Supplemental Figure 3.)

Ultrasound decreased action-potential width for every action potential at all firing frequencies. As

shown in Figure 6G-H, ultrasound also decreased action-potential width when applied 1 s after

the start of a current step, again indicating that the channels responsible for these effects continue

to influence firing frequency and remain responsive to ultrasound throughout sustained

depolarizing current steps.

The effects of ultrasound on action-potential width tended to counteract the broadening of

action-potential width that occurs during high-frequency firing. Figure 7 plots action potential width

as a function of action potential number and input current for the control and ultrasound conditions.

In the control condition, there are dramatic differences in width between the first action potential

and subsequent action potentials at high input currents, while in the ultrasound condition these

differences are much less pronounced. To quantify this effect, we measured the difference in

width between the first and third, first and fifth, and first and last action potentials during the

ultrasound stimulus, in response to a +450 pA current step for the control and ultrasound

conditions. These differences (mean ± SEM, control vs. ultrasound, N = 9) were 2.6 ± 0.9 vs. 0.7

± 0.1 ms, 1.5 ± 0.1 vs. 0.5 ± 0.1 ms, and 1.2 ± 0.2 vs 0.5 ± 0.1 ms for the third, fifth, and last action

potentials (P = 0.064, 1.4 x 10^{-5}, and 0.0013, paired, two-tailed Student's t-test). One plausible

interpretation of this result is that K2P channels activated by ultrasound take over the role of K_V

channels that would otherwise contribute to action-potential repolarization. Since time-dependent

activation and inactivation of K_V channels causes action-potential broadening during repetitive

firing (Giese et al., 1998; Shao et al., 1999; Yue and Yaari, 2004; Kim et al., 2005; Gu et al.,

2007), replacement of these channels by K2P channels lacking time-dependent activation and

inactivation would reduce time- and frequency-dependent action potential broadening.

Although the effects of ultrasound on action-potential widths are predominantly due to

acceleration of the repolarization phase, we also noted effects on the rising phase of the action

potential. These effects are readily apparent in the first derivative of membrane voltage during the

action potential (Figure 8A) or in action potential phase plots (plots of the first derivative of voltage

versus voltage, Figure 8B). In fact, the maximum rates of voltage rise and fall during the action

potential were both consistently increased by ultrasound throughout a spike train (Figure 8C-D).

The effect on the falling phase is to be expected based on the observed decrease in spike width

and the hypothesis that ultrasound activates K2P channels. The effect on the rising phase is also

consistent with this hypothesis, as activation of K2P leading to reduced Na_V channel inactivation

would increase the number of Na_V channels available to activate during the rising phase of the

action potential.

In addition to effects of ultrasound on the rising and falling phases of the action potential,

we also found effects on the AHP. To quantify these effects, we measured the voltage minimum

between action potentials during repetitive firing. Because measurements of this parameter are

very sensitive to changes in resting membrane voltage and series resistance that can occur over

long recording times, we compared voltage minimums before, during, and after ultrasound

application within the same voltage trace (Figure 9A-B) and made a similar comparison for control

recordings. We performed these comparisons for ultrasound applied 1 s after the start of a current,

for cells firing at an average frequency of 5 Hz in the control condition. This firing frequency is

near the transition region between the inhibitory and potentiating effects of ultrasound on spike

frequency, such that the spike frequency is similar for the control and ultrasound conditions,

allowing us to compare a similar number of interspike voltage minima for the control and

ultrasound conditions (Figure 9C). This analysis demonstrates that the depth of the AHP is greater

during the ultrasound application than before or after it, or during the same time windows for the

control condition. Together with the effects of ultrasound on spike waveform (Figure 6), this result

supports the idea that ultrasound activates a sustained outward current, which limits Na_V channel

inactivation and thereby potentiates high-frequency firing. Removal of Na_V channel inactivation

by membrane hyperpolarization also explains how ultrasound can rescue spiking in neurons that

have entered a refractory state due to accumulation of Na_V channel inactivation (Figure 3C,

bottom right).

Physical mechanism of neuromodulation by high-frequency ultrasound. The idea that ultrasound

acts on K2P channels is also consistent with the physical effects of ultrasound on biological tissue.

At 43 MHz, two plausible mechanisms through which ultrasound might modulate the activity of

ion channels are heating and mechanical stress due to acoustic radiation force. Absorption of

acoustic energy by biological tissue as heat can increase its temperature, with effects on ion

channel gating and all other biological reactions. Absorption also results in attenuation of

ultrasound intensity as the wave propagates, creating spatial gradients in intensity that give rise

to radiation force, which in turn produces tissue displacement and strain. At the microscopic scale,

this displacement and strain may involve increased tension in the cell membrane, cytoskeleton,

and extracellular matrix, all of which may affect excitability through mechanical effects on ion

channel proteins. Among the K2P channels that may be expressed by CA1 pyramidal cells, TREK

and TRAAK channels are especially sensitive to thermal and mechanical stimuli.

To gain further insight into these physical mechanisms, we performed finite-element

simulations of the effects of ultrasound on brain slices in the context of our experimental set-up.

The simulated spatial profiles of ultrasound-induced heating and macroscopic tissue

displacement in response to radiation force are shown in Figure 10A-B. Notably, the spatial

profiles of heating and displacement effects are significantly wider than the 90-micron diameter

of the focal volume of the ultrasound beam, with significant heating and displacement occurring

several hundreds of microns away from the beam axis (Figure 10C). This is an important result

because the thermo- and mechanosensitive K2P channels TREK-1 and TRAAK are expressed

at high density at the nodes of Ranvier of vertebrate neurons (Brohawn et al., 2019; Kanda et al.,

2019). The first node of Ranvier is located approximately 100 microns from the axon initial

segment (Kole, 2011). In our experiments, the soma of the patched neuron is located

approximately on the axis of the ultrasound beam (see *Materials and Methods*) so the first node

of Ranvier is probably within the region of the tissue exposed to thermal and mechanical effects

of ultrasound. Axonal K^+ channels play important roles in regulating excitability (Shah et al., 2008;

Kole, 2011; Kanda et al., 2019). It is therefore plausible that a subpopulation of TREK-1 or TRAAK

channels at the nodes of Ranvier could contribute to the neurophysiological effects of ultrasound.

The magnitude of the temperature change is also consistent with a thermal mechanism for the

effects of ultrasound. The maximum temperature change in the simulation is 1.3 C; temperature

changes of this size have previously been shown to affect neural excitability (Owen et al., 2019).

The maximum value of the simulated displacement (1.7 microns) is similar to the displacement

measured in the retina during ultrasound neuromodulation with stimulus parameters similar to

those used here (Menz et al., 2019).

It is instructive to consider the amplitude and time course of the membrane capacitance

change in response to ultrasound (Figure 5) in the context of possible thermal and mechanical

mechanisms. As described above (under *Effects of ultrasound on resting membrane potential*),

the capacitance change in response to ultrasound is fast relative to the membrane time constant,

so we can assume that the total charge on the membrane is constant during the initial transient

depolarization in response to ultrasound (Figure 5A, *left arrow*). In other words, the numerator in

the equation $V = Q/C$ is constant, so for small changes in voltage the relative change in voltage

is approximately inversely proportional to the relative change in capacitance. Empirically,

membrane capacitance increases by approximately 1% per degree C (Taylor, 1965). This is

consistent with the size of the simulated temperature rise (peak simulated temperature rise of

1.3°C compared with the measured amplitude of the initial decrease in voltage of 0.7 ± 0.1%).

However, the time course of the temperature rise (Figure 10D) is much slower than the time

course of the voltage transient (which again, assuming constant Q, is identical to the time course

of the capacitance change). The time course of the change in resting membrane potential (173 ±

18 ms, Figure 5), however, parallels that of the temperature rise. (A capacitance change on the

time course of the simulated temperature rise would not have a significant effect on the membrane

voltage, as it would be counteracted by ionic currents.) Thus, the simulated ultrasound heating

results strongly suggest that ultrasound affects action-potential firing in our experiments at least

in part through a thermal effect on ion channels, but do not explain the presence of the capacitive

transients.

We considered whether the time course of the capacitive transients could instead be

explained by the dynamics of the tissue mechanical response to acoustic radiation force. We

sought to determine whether, having already modeled the static displacement of the tissue, we

could, without retroactively changing any of the tissue material properties, obtain a time course

for tissue displacement similar to that of the capacitive transients using a simple viscoelastic

model with reasonable tissue viscous properties (see *Materials and Methods*). We found that this

could be achieved using a shear viscosity (m) of 1 Pa·s (Figure 10E). Since the tissue is

essentially incompressible in our model (Poisson's ratio (n) = 0.4998), this is equivalent to a

relaxation time of $2m \cdot (1+ n)/E$ = 6 ms (where E is Young's modulus). Biological tissue is a highly

heterogeneous material that displays a variety of active and passive mechanical responses to

force, spanning time scales from milliseconds to hours (Ricca et al., 2013), and as a result its

viscous properties are highly sensitive to the time scale of the measurement and even complex

viscoelastic models encompassing multiple relaxation times may not fully describe the

viscoelastic behavior of tissue. Nonetheless, the shear viscosity/relaxation time in our model is

reasonable for a soft, gel-like material, and is comparable to fast relaxation times observed

experimentally in brain tissue (Arbogast and Margulies, 1999; Abolfathi et al., 2009; Rashid et al.,

2012, 2013). Moreover, the simulated time course of displacement is consistent with experimental

measurements of the tissue displacement in response to ultrasound at 43 MHz and 40 W/cm^2 in

the salamander retina, which was found to be complete in less than 10 ms (Menz et al., 2019).

We can therefore make the reasonable assumption that the capacitive transients are due

to a mechanical effect on membrane properties, and we can estimate the size of the potential ion-

channel gating effects that would occur as a result of this mechanical effect. The capacitance of

a lipid bilayer membrane is given by C = $\varepsilon \cdot \varepsilon_0 \cdot A/L$, where ε is the dielectric constant of the

hydrophobic core of the lipid bilayer, ε_0 is the permittivity (polarizability) of free space, A is

membrane area, and L is the thickness of the hydrophobic core of the membrane. For small

strains like those under consideration here, lipid bilayer membranes can be considered

incompressible, such that a 1% increase in capacitance corresponds to a 0.5% increase in area

and a 0.5% decrease in thickness (White and Thompson, 1973; Alvarez and Latorre, 1978). An

increase in membrane area can be converted to an increase in membrane tension (γ) according

to $\gamma = \Delta A \cdot K_A$, where ΔA is the relative change in area and K_A is the area elastic constant of the

membrane. Area elastic constants measured for lipid membranes are on the order of 100's of

mN/m (Evans et al., 1976; Kwok and Evans, 1981; Needham and Nunn, 1990). If the capacitance

transients are due to membrane strain, the resulting membrane tension is on the order of a few

0.1 mN/n to a few mN/m. These values are similar to the tension thresholds for activation of

mechanosensitive K2P channels (estimated as 0.5-4 mN/m for activation of TREK-1 and TRAAK

(Brohawn et al., 2014)), which are low relative to other known mammalian mechanosensitive

channels. Notably, a recent *in vivo* ultrasound neuromodulation study of the murine sciatic nerve

at 4 MHz, found that tissue displacement *in vivo* was highly correlated with the neuromodulation

effects (Lee et al., 2020). Nonetheless, additional data or theoretical advances would be required

to firmly associate these capacitive transients with changes in membrane tension. If such an

association could be made, it would provide strong evidence that ultrasound modulates action

potential firing through mechanical effects of radiation force in our experiments. At present, our

results do not rule out this idea, but the case for mechanical effects remains speculative, while

the role of thermal effects seems highly plausible. Nonetheless, our simulation results support the

conclusion that both inhibitory and excitatory effects of high-frequency ultrasound on action-

potential firing are due to activation of thermo- and mechanosensitive K2P channels.

DISCUSSION

To summarize, the neurophysiological effects of ultrasound that we have described here can all

be explained by activation of a sustained outward current. We argue that the molecular basis of

this outward current is most likely one or more of the K2P channels expressed by CA1 pyramidal

neurons. Although a variety of voltage-dependent K+ currents shape the action-potential

waveform and regulate excitability in these neurons, several arguments suggest that K2P

channels are the molecular basis of the ultrasound-activated outward conductance. First, the K2P

channels TREK and TRAAK, being strongly mechanosensitive and thermosensitive, have

biophysical properties that make them especially sensitive to physical effects of ultrasound.

Second, the fact that ultrasound has similar effects on firing frequency whether it is applied 500

ms before or 1 s after the start of a current step suggests that ultrasound affects firing through a

channel whose properties are not strongly modulated by voltage. Finally, the neurophysiological

effects of ultrasound are, strikingly, essentially the opposite of those caused by knock-out of

TASK-3 channels in cerebellar granule neurons (Brickley et al., 2007), which increased excitability

at low input currents, but decreased excitability at high input currents and led to failure of

sustained high-frequency firing. In addition, knock-out of TASK-2 decreased the maximum firing

frequency and decreased action-potential height while increasing action potential width. TREK-1

and TRAAK channels are also necessary for high frequency firing at the nodes of Ranvier of

afferent neurons (Kanda et al., 2019).

Ultrasound neuromodulation has been studied in vertebrate axon preparations, where it

has generally been found that ultrasound inhibits action-potential conduction, with the effect

specifically attributed to heating in some cases (Young and Henneman, 1961; Mihran et al., 1990;

Tsui et al., 2005; Colucci et al., 2009). These results are consistent with the idea that activation

of K2P channels by ultrasound can inhibit action potential firing, but it would be worthwhile to

revisit these experiments to see whether the bidirectional, spike-frequency-dependent effect that

we observe is also present in such preparations. Our results also suggest an approach to

ultrasound neuromodulation in which action potential propagation is the locus of neuromodulation,

targeting white-matter tracts instead of soma-dense gray matter. The idea that ultrasound-

activated K^+ currents can both inhibit and potentiate firing might also help explain why ultrasound

can both inhibit and potentiate neural activity *in vivo* (Min et al., 2011).

Although activation of K2P channels is sufficient to explain our results, we do not rule out

the possibility that ultrasound affects other channels in addition to K2P's; indeed, we think it is

likely that ultrasound does affect other channels to some extent. All ion-channel gating reactions

are sensitive to temperature, with typical Q10 values of ~3, such that their rates would be

expected to increase by about 10% based on the temperature changes in our simulations (Hille,

2001). Mechanical effects of radiation force could also affect channels besides K2P channels.

The mechanically gated channel Piezo2 is expressed in a subset of CA1 pyramidal neurons

(Wang and Hamill, 2020). In addition, most ion channels and membrane proteins, while not

functioning physiologically as mechanoreceptors, are sensitive to mechanical stimuli to some

extent, either through the energetics of their interactions with hydrophobic core of the lipid bilayer

or through mechanical interactions with the cytoskeleton or extracellular matrix. In fact, gating of

voltage-dependent Na^+, K^+, and Ca^{2+} channels can be modulated by membrane stretch in

membrane patches (Calabrese et al., 2002; Tabarean and Morris, 2002; Laitko and Morris, 2004;

Morris and Juranka, 2007; Beyder et al., 2010). However, we previously were unable to detect

any mechanical modulation of heterologously expressed NaV1.2 channels by ultrasound at 43

MHz and 90 W/cm^2 under conditions where ultrasound activated the mammalian

mechanoreceptor channel Piezo1 (Prieto et al., 2018). Neural Na_V channels and KCNQ channels

interact with the periodic actin cytoskeleton of axons through spectrin and ankyrin-G at the axon

initial segment and nodes of Ranvier and (Zhou et al., 1998; Pan et al., 2006; Leterrier, 2018),

suggesting that they may be sensitive to modulation by cytoskeletal tension. The concentration

of TREK-1 and TRAAK channels at the nodes of Ranvier suggest that some similar interaction

with the cytoskeleton may be involved in the localization of these channels, although this has not

been demonstrated, and the intracellular domains that would facilitate such interactions are

relatively small in K2P channels.

It is well established that high-intensity light at infrared and shorter wavelengths can

modulate neural activity through tissue heating. As a general principle, thermal neuromodulation

effects in response to optical stimulation would be expected to be very similar to thermal

neuromodulation effects caused by ultrasound, and studies of optical neuromodulation could

therefore provide useful guidance in interpreting our results. However, optically based thermal

neuromodulation experiments are highly heterogeneous in terms of the neuromodulation effect,

the temperature rise required to produce the effect, and the mechanistic interpretation of the

results (Wells et al., 2007; Richter et al., 2011; Shapiro et al., 2012; Duke et al., 2013; Walsh et

al., 2016; Lothet et al., 2017; Paris et al., 2017; Owen et al., 2019; Zhu et al., 2019), so their

usefulness is limited in this respect. Both inhibition and potentiation of firing have been reported,

and the increase in temperature has varied considerably, ranging from less than 1 C to 10's of

degrees C. The temperature rise in our simulations is on the low end of this range. However, it

has been proposed that spatial or temporal gradients in temperature, rather than the absolute

temperature change, may determine the response to thermal stimuli (Wells et al., 2007; Paris et

al., 2017). In addition, heating can cause phase changes in lipid bilayers, which have been shown

to modulate ion channel activity (Seeger et al., 2010); in this case, the response to heating would

also depend critically on the initial temperature. Interestingly, a recent study demonstrated

inhibition of firing by small (2°C or less) increases in temperature in several different types of

neurons, but *not* in CA1 pyramidal neurons (Owen et al., 2019). These effects were attributed to

inward rectifier (K_{ir}) potassium channels, which are not expressed in CA1 pyramidal neurons.

However, our results suggest that the lack of an effect in pyramidal neurons could also be

explained if the experiments were performed at a point on the f-i curve where competing inhibitory

and excitatory effects of heat-activated K^+ current result in no net effect on firing frequency. A

recent *in vivo* ultrasound neuromodulation study in a rat model using ultrasound at 3.2 MHz with

exceptionally long ultrasound exposure times (10's of seconds; see below for a more general

discussion of *in vivo* ultrasound neuromodulation studies) also found that inhibition could be

produced by ultrasound-induced temperature rises of 2°C or less (Darrow et al., 2019), consistent

with this result and our results.

Finally, a critical question—to which we cannot yet provide a definitive answer—is whether

the mechanisms underlying the neuromodulatory effects of ultrasound are the same in our

experiments and in *in vivo* experiments using low frequencies. Two results that strongly argue

against similar mechanisms are the apparent frequency-dependence of neuromodulation in *in*

vivo experiments, and the much lower intensities that have been reported to cause

neuromodulatory effects in some low-frequency, *in vivo* experiments as compared with our

results. In an *in vivo* mouse model of ultrasound neuromodulation, it was determined that the

efficacy of neuromodulation decreased with increasing frequency over the range 0.25-2.9 MHz

(King et al., 2013; Ye et al., 2016). This frequency dependence is the opposite of what would be

expected for either a thermal or radiation force mechanism, since both of these are proportional

to the ultrasound attenuation coefficient, which in tissue is proportional to frequency raised to the

power of ~1.1 (Hand, 1998). In contrast, Menz et al. (2019) found that the efficacy of

neuromodulation increases with frequency over the range 1.9-43 MHz in the retina *in vitro*, a thin

neural tissue preparation similar to the one used in our experiments. However, they proposed a

model to explain this discrepancy. Lower ultrasound frequencies generally result in a larger

stimulated tissue volume, which could translate into a more effective stimulus for certain

structures of circuit-level neural connectivity, despite a weaker effect of low-frequency ultrasound

at the level of an individual cell. (The model was presented in the context of a radiation force

mechanism, but the same principle could apply for a thermal mechanism). The idea that circuit-

level mechanisms can amplify the response to ultrasound is supported by comparison of our

results with the response to ultrasound in the retina at 43 MHz. In the retina, potentiation of action-

potential firing by ultrasound at 43 MHz saturates at 10 W/cm^2, as measured at the population

level in an intact, active neural circuit. Although we have not performed a detailed investigation of

the intensity dependence, we find that a much higher intensity, 50 W/cm^2, produces relatively

moderate effects on excitability in single cells in the absence of significant network activity.

Such considerations may mitigate frequency-dependence as an argument against similar

physical mechanism for neuromodulation by high- and low-frequency ultrasound. The other major

argument against similar mechanisms for high and low frequencies is the difference in the size of

the potential thermal of radiation force effects at different intensities and frequencies. One point

specific to the K2P channel hypothesis is worth mentioning in this context. The midpoints of the

temperature-activation curves for TREK and TRAAK channels are near 37°C (in other words,

near body temperature in mammals) so that the increase in K2P current in response to increased

temperature is near maximal at physiological temperature (Maingret et al., 2000; Kang et al.,

2005), whereas our experiments were performed at room temperature (21-23°C). Room

temperature is near the threshold for temperature activation of these channels, such that they are

mainly inactive at room temperature in the absence of other stimuli such as membrane stretch,

lipid agonists, or acidic pH.

In conclusion, our results demonstrate that high-frequency ultrasound is a viable and

promising modality for neuromodulation applications where frequency is not limited by

transmission through the skull, and our insights into the common molecular mechanisms

underlying both inhibitory and excitatory effects of high-frequency ultrasound pave the way for

rational design and optimization of neuromodulation protocols to consistently produce either

inhibitory or excitatory effects.

ACKNOWLEDGMENTS

We thank the PIs and members of the Baccus, Butts-Pauly and Huguenard laboratories for helpful

discussions. We thank Mike Menz and Stephen Baccus for critically reading the manuscript. We

thank Marianna Kiraly, Dong Li, and Jason Clark for help with hippocampal slice preparation and

recording. We thank the Stephen Boxer lab for sharing their plasma cleaner.

This work was supported by NIH R01 EB019005 (M.M. and B.K.Y.), NIH BRAIN initiative R01

NS11215 (M.M. and B.K.Y.), NIH R01 MH111768 (D.M) and by the Harold and Leila Y. Mathers

Charitable Foundation (M.M and D.M).

The authors declare no competing financial interests.

AUTHOR CONTRIBUTIONS

Martin Prieto performed experiments, analyzed data, and wrote the manuscript. Martin Prieto and

Kamyar Firouzi performed computational modeling. Merritt Maduke, Daniel Madison, and Butrus

Khuri-Yakub supervised research. All authors contributed to the design and interpretation of

experiments and edited and revised the manuscript.

Table 1. Values of Material Properties Used in Finite-Element Simulations.

	water	polystyrene	brain tissue
density (kg/m^3)	1000[a]	1040[b]	1007[c]
speed of sound (m/s)	1500[a]	2300[b]	1538[c]
attenuation coefficient at 43 MHz (neper/m)	46[d]	160[e]	253[f]
heat capacity (J/kg·K)	4180[a]	1200[g]	3500[h]
thermal conductivity (W/m·K)	0.6[a]	0.1[g]	0.5[h]
Young's modulus (Pa)	(not applicable)	10^9 [i]	500[j]
Poisson's ratio	(not applicable)	0.4[i]	0.4998[k]
shear viscosity (Pa·s)	(not applicable)	(not applicable)	1[l]

Sources for material properties are as follows: [a]standard value; [b]based on typical acoustic properties of plastics (Selfridge, 1985); [c]following Menz et al. (2019), based on (Thijssen et al., 1985); [d] (Company, 1965); [e]measured (Prieto et al., 2018); [f]following Menz et al. (2019), based on (de Korte et al., 1994); [g]based on typical thermal properties of plastics (Gaur and Wunderlich, 1982; Harper, 2006); [h]typical values for soft tissues (Hand, 1998); [i]based on typical mechanical properties of plastics (Harper, 2006); [j]Menz et al. (2019), from measurements of ultrasound-induced displacement in the retina; [k]tissue assumed to be incompressible for small deformations; [l]see text.

Figure 1. Consistent inhibition of action potential firing by high-frequency ultrasound. A. Diagram of experimental set-up. Ultrasound is applied to 500-micron hippocampal brain slices resting on a 25-micron film of polystyrene. The 43-MHz focused transducer is located below the experimental chamber, with ultrasound propagating perpendicular to the bottom of the recording chamber. **B.** Experimental protocol and example voltage traces showing inhibition of action potential firing by ultrasound. A 1-s, continuous-wave ultrasound pulse at 43 MHz and 50 W/cm^2 (*red bar*) is applied 500 ms before the start of a 2-s current injection. Voltage traces are shown in the presence (*red*) and absence (*blue*) of the ultrasound stimulus. **C.** Example raster plots showing a consistent effect of ultrasound on firing frequency. The results of twenty consecutive trials of the protocol in panel B, alternating between the control (*top*) and ultrasound (*bottom*) conditions, are shown. The voltage traces were divided into 50-ms bins; a solid black bin indicates that an action potential occurred within that particular time bin. Time is relative to the start of the ultrasound pulse. **D.** Spike-time histograms prepared by summing the ten trials for the control and ultrasound conditions from panel C. **E.** Summary of the effects of ultrasound for N = 66 cells. The average firing frequency during the first 500 ms of the current step is shown for the control and ultrasound conditions. **F.** Stability of the ultrasound response. Mean (±SEM, N = 10 cells) spike frequencies during the first 500 ms of a current step in the presence (*solid circles*) and absence (*open circles*) of ultrasound for the protocol shown in panel B, as a function of time relative to break-in (establishment of whole-cell recording configuration). Spike frequencies were measured at various time points between 0 and 10, 10 and 20, 20 and 30, and 30 and 40 minutes after break-in. The x-values represent the mean start time for the protocol to measure spike frequencies (which comprised 2 minutes of recording time). The amplitude of the current step was adjusted over time to maintain spiking behavior as close as possible to that at the start of the experiment.

Figure 2. Ultrasound can inhibit or potentiate action-potential firing. A. Example frequency-input curve showing average firing frequency during the first 500 ms of a current step as a function of input current, with (*red*) and without (*blue*) a 1-s ultrasound pulse starting before 500 ms before the start of the current step. Each point represents the average of three trials on the same cell. **B.** Example voltage traces for the cell in A showing action potential firing during the first 500 ms of current steps to either +30, +40, +80, or +250 pA, with (*red*) and without (*blue*) ultrasound. **C.** Mean (±SEM, N = 9 cells) change in spike frequency in response to ultrasound as a function of input current. **D.** Inhibition is strongest near threshold. Mean (±SEM, N = 7 cells) change in spike frequency in response to ultrasound as a function of input current relative to the threshold current for action potential firing, for near-threshold currents.

Figure 3. Ultrasound can also inhibit or potentiate action-potential firing when applied late in a current step. A. Experimental protocol and example voltage traces with (*red*) and without (*blue*) ultrasound. Ultrasound is applied 1 s after the start of a 3-s current step. **B.** Example frequency-input curve showing average firing frequency during the ultrasound application for the protocol in A (*red*) and during the same time window without ultrasound (*blue*), as a function of input current. Each point represents the average of three trials on one cell. **C.** Example voltage traces for the cell in B showing action potential firing during the ultrasound application (*red*) and during the same time window without ultrasound (*blue*) in response to current steps to either +50, +100, +250, or +450 pA. **D.** Mean (±SEM) change in spike frequency in response to ultrasound as a function of input current (N = 3 cells at +50 pA; N = 6 cells at all other input currents). (In 3 cells, the firing frequency at +50 pA was zero for both the control and ultrasound conditions).

Figure 4. Effects of ultrasound on action potential timing. A-C. Mean (±SEM) intervals between the start of the current step and the first action potential, and between the first and second, and second and third, etc. action potentials, for cells firing at average frequencies of approximately 5 Hz (N = 13 cells), 10 Hz (N = 15 cells), and 20 Hz (N = 13 cells) during the first 500 ms of the current step, with (*red*) or without (*blue*) a 1-s ultrasound pulse starting 500 ms before the current step. The actual spike frequencies were 5.6 ± 0.1, 10.5 ± 0.2, and 20.4 ± 0.3 Hz, and the injected currents were 70 ± 7, 115 ± 9, and 292 ± 28 pA for the 5, 10, and 20 Hz conditions. **D.** Example voltage traces showing decreased interval between the first and second action potentials at low average firing frequency. The top panel shows the first two action potentials for the control (*blue*) and ultrasound (*red*) conditions. The bottom panel shows the same data, aligned to the action potential threshold on zoomed-in time scale. **E.** Same as A-C, for an currents was 183 ± 21 pA.

Figure 5. Effects of ultrasound on resting membrane potential and membrane capacitance. A. Six individual voltage traces (*pink*) and the average of these voltage traces (*red*) showing the effect of ultrasound at (*red bar*) on resting membrane potential. The *black arrows* indicate transients due to changes in membrane capacitance. **B.** Zoomed-in timescale showing the fast voltage transients (*black arrows* in A) for ultrasound onset (*top left*) and offset (*bottom right*). Exponential fits (*black lines*) to the rise and fall of the voltage transients give amplitudes and time constants of 0.45 mV and 16.3 ms for the onset and 0.22 mV and 6.7 ms for the offset, for the example shown; mean values (±SEM) were 0.41 ± 0.04 mV and 10.4 ± 1.2 ms for the onset and 0.35 ± 0.04 mv and 9.7 ± 1.7 ms for the offset (N = 15). No significant differences were found between the time constants (P = 0.73) or the amplitude (P = 0.087) of the transients (paired, two-tailed Student's t-tests) **C.** Slow membrane hyperpolarization in response to ultrasound from the average trace in panel A on a zoomed-in scale, along with an exponential fit (*black line*) to the initial hyperpolarization. The amplitude and time constant of the exponential fit were 1.55 mV and 132 ms for the example shown; mean values (±SEM) were 2.4 ± 0.3 mV and 173 ± 18 ms (N = 15).

Figure 6. Ultrasound decreases action-potential width. A-B. Mean (±SEM, N = 13 cells) action potential-widths (A) and example action-potential waveforms aligned to the action-potential threshold (B) as function of action potential number in the presence (*red*) and absence (*blue*) of a 1-s ultrasound pulse starting 500 ms before the current step, for cells firing at an average firing frequency of approximately 5 Hz (as measured during the first 500 ms of the current step) in the control condition. **C-D.** As in A-B, but for cells firing at an average firing frequency of approximately 10 Hz in the control condition (N = 15). **E-F.** As in A-B, but for cells firing at an average firing frequency of approximately 20 Hz in the control condition (N = 13). **G-H.** As in A-B, but with ultrasound applied 1 s after the start of a 3-s current step, for cells firing at an average firing frequency of approximately 5 Hz in the control condition, with firing frequency determined in a 1-s window starting 1 s after the current step (corresponding to the time period of the ultrasound stimulus), and action potential number relative to the start of the ultrasound stimulus (N = 6).

Figure 7. Ultrasound reduces action-potential broadening. Example data showing action-potential width as a function of action-potential number (indicated in *grayscale*, scale bar at far left) and input current level, for the first 500 ms of the current step, for currents from 0 to +450 pA in 50 pA steps, with (*right*) or without (*left*) a 1-s ultrasound pulse starting 500 ms before the current step. The vertical lines indicate the approximate location of the transition between inhibitory and potentiating effects of ultrasound.

Figure 8. Effects of ultrasound on depolarization and repolarization rates. A. Example traces showing the membrane voltage (*top*) and its first derivative (*bottom*) for the first three action potentials in response to a +100 pA current step in the presence (*red*) and absence (*blue*) of a 1-s ultrasound pulse starting 500 ms before the current step, aligned to the action potential threshold. **B.** Phase plots for the action potentials shown in panel A. **C-D.** Maximum rates of depolarization (*left*) and repolarization (*right*) during the action potential (mean ± SEM, N = 13 cells), as a function of action potential number, in the presence (*red*) and absence (*blue*) of a 1-s ultrasound pulse starting 500 ms before the current step, for cells firing at an average firing frequency of 5 Hz (**C**) or 20 Hz (**D**) in the control condition.

Figure 9. Ultrasound increases the depth of the afterhyperpolarization. A. Example voltage trace comparing voltage minima between action potentials in response to a 3-s, 250-pA current step with a 1-s ultrasound pulse (*red bar*) starting 1 s after the start of the current step. The *black lines* connect the voltage minima between action potentials before, during, or after the ultrasound pulse. **B.** Same as panel A, on a zoomed-in voltage scale.

C. Mean (±SEM, N = 3-6 cells, see figure panel for details) values of the voltage minimum, as a function of action potential number, for the first four to six action potentials before (*solid diamonds*, 5 action potentials), during (*solid circles*, 6 action potentials), and after (*solid squares*, 5 action potentials) the ultrasound pulse, along with the equivalent mean values for the control condition (*open symbols*, 4, 6, and 5 action potentials). The means were determined for cells firing at the same average frequency (5 Hz) during a 1-s window starting 1 s after the start of the current step (corresponding to the period of the ultrasound stimulus) in the control condition. Significant differences between groups were only found in the presence of ultrasound (P = 1.0 x 10^{-4}, 1.4 x 10^{-5}, and 0.16 for before versus during, during versus after, and before versus after the ultrasound pulse; P = 0.92, 0.39, and 0.35 for comparisons of the same time periods in the absence of ultrasound (unpaired, two-tailed Student's t-tests, unequal variance).

Figure 10. Simulated tissue heating and displacement in response to ultrasound. A. Spatial profile of temperature after one second of ultrasound exposure at 43 MHz and 50 W/cm², as a function of axial distance from the transducer surface and radial distance from the ultrasound beam axis, in a 500-micron thick brain slice and 25-micron thick polystyrene film (*middle two layers*) and the surrounding fluid (external solution, *top layer*, and distilled water, *bottom layer*). **B.** Spatial profile of the static total displacement in response to acoustic radiation force in the brain slice and polystyrene film. **C.** Normalized values of the acoustic intensity (*solid black line*), temperature rise after 1 second of ultrasound exposure (*solid gray line*), and total displacement (*dashed line*), at a depth of 250 microns in the brain slice, as a function of radial distance. **D.** Time course of the temperature rise in response at ultrasound at a depth of 250 microns in the brain slice on the axis of the ultrasound beam. The time course of the temperature change can be described by two exponential components with amplitudes and time constants of -0.56 C and 30 ms, and -0.57 C and 295 ms, for a weighted time constant of 164 ms. **E.** Time course of the displacement in response at ultrasound at a depth of 250 microns in the brain slice on the axis of the ultrasound beam. The time course of the displacement change can be described by two exponential components with amplitudes and time constants of -1.2 microns and 6 ms, and -0.13 microns and 344 ms, for a weighted time constant of 39 ms. Note that the steady-state displacement is slightly smaller than in the static displacement simulation due to the inclusion of the fluid loading in the dynamic displacement simulation.

REFERENCES

Abolfathi, N., A. Naik, M. Sotudeh Chafi, G. Karami, and M. Ziejewski. 2009. A micromechanical procedure for modelling the anisotropic mechanical properties of brain white matter. *Comput Methods Biomech Biomed Engin*. 12:249-262.

Alvarez, O., and R. Latorre. 1978. Voltage-dependent capacitance in lipid bilayers made from monolayers. *Biophys J*. 21:1-17.

Arbogast, K.B., and S.S. Margulies. 1999. A fiber-reinforced composite model of the viscoelastic behavior of the brainstem in shear. *J Biomech*. 32:865-870.

Bean, B.P. 2007. The action potential in mammalian central neurons. *Nat Rev Neurosci*. 8:451-465.

Beyder, A., J.L. Rae, C. Bernard, P.R. Strege, F. Sachs, and G. Farrugia. 2010. Mechanosensitivity of Nav1.5, a voltage-sensitive sodium channel. *J Physiol*. 588:4969-4985.

Blackmore, J., S. Shrivastava, J. Sallet, C.R. Butler, and R.O. Cleveland. 2019. Ultrasound Neuromodulation: A Review of Results, Mechanisms and Safety. *Ultrasound Med Biol*. 45:1509-1536.

Blanton, M.G., J.J. Lo Turco, and A.R. Kriegstein. 1989. Whole cell recording from neurons in slices of reptilian and mammalian cerebral cortex. *J Neurosci Methods*. 30:203-210.

Brickley, S.G., M.I. Aller, C. Sandu, E.L. Veale, F.G. Alder, H. Sambi, A. Mathie, and W. Wisden. 2007. TASK-3 two-pore domain potassium channels enable sustained high-frequency firing in cerebellar granule neurons. *J Neurosci*. 27:9329-9340.

Brohawn, S.G., Z. Su, and R. MacKinnon. 2014. Mechanosensitivity is mediated directly by the lipid membrane in TRAAK and TREK1 K+ channels. *Proc Natl Acad Sci U S A*. 111:3614-3619.

Brohawn, S.G., W. Wang, A. Handler, E.B. Campbell, J.R. Schwarz, and R. MacKinnon. 2019. The mechanosensitive ion channel TRAAK is localized to the mammalian node of Ranvier. *Elife*. 8.

Bystritsky, A., A.S. Korb, P.K. Douglas, M.S. Cohen, W.P. Melega, A.P. Mulgaonkar, A. DeSalles, B.K. Min, and S.S. Yoo. 2011. A review of low-intensity focused ultrasound pulsation. *Brain Stimul*. 4:125-136.

Calabrese, B., I.V. Tabarean, P. Juranka, and C.E. Morris. 2002. Mechanosensitivity of N-type calcium channel currents. *Biophys J*. 83:2560-2574.

Calhoun, M.A., S.A. Bentil, E. Elliott, J.J. Otero, J.O. Winter, and R.B. Dupaix. 2019. Beyond linear elastic modulus: viscoelastic models for brain and brain mimetic hydrogels. *ACS Biomaterials Science & Engineering*. 5:3964-3973.

Castaneda-Castellanos, D.R., A.C. Flint, and A.R. Kriegstein. 2006. Blind patch clamp recordings in embryonic and adult mammalian brain slices. *Nat Protoc*. 1:532-542.

Colucci, V., G. Strichartz, F. Jolesz, N. Vykhodtseva, and K. Hynynen. 2009. Focused ultrasound effects on nerve action potential in vitro. *Ultrasound Med Biol*. 35:1737-1747.

Company, C.R. 1965. Handbook of Chemistry and Physics. 45th.ed. Chemical Rubber Compnay, Cleveland, OH.

Cotero, V., Y. Fan, T. Tsaava, A.M. Kressel, I. Hancu, P. Fitzgerald, K. Wallace, S. Kaanumalle, J. Graf, W. Rigby, T.J. Kao, J. Roberts, C. Bhushan, S. Joel, T.R. Coleman, S. Zanos, K.J. Tracey, J. Ashe, S.S. Chavan, and C. Puleo. 2019. Noninvasive sub-organ ultrasound stimulation for targeted neuromodulation. *Nat Commun*. 10:952.

Darrow, D.P., P. O'Brien, T.J. Richner, T.I. Netoff, and E.S. Ebbini. 2019. Reversible neuroinhibition by focused ultrasound is mediated by a thermal mechanism. *Brain Stimul*. 12:1439-1447.

de Korte, C.L., A.F. van der Steen, and J.M. Thijssen. 1994. Acoustic velocity and attenuation of eye tissues at 20 MHz. *Ultrasound Med Biol*. 20:471-480.

Downs, M.E., S.A. Lee, G. Yang, S. Kim, Q. Wang, and E.E. Konofagou. 2018. Non-invasive peripheral nerve stimulation via focused ultrasound in vivo. *Phys Med Biol*. 63:035011.

Duck, F.A. 1998. Radiation pressure and acoustic streaming. *In* Ultrasound in Medicine. F. Duck, A. Baker and H. Starritt, editors. Institute of Physics Publishing, Philadelphia, PA.

Duke, A.R., M.W. Jenkins, H. Lu, J.M. McManus, H.J. Chiel, and E.D. Jansen. 2013. Transient and selective suppression of neural activity with infrared light. *Sci Rep*. 3:2600.

Evans, E.A., R. Waugh, and L. Melnik. 1976. Elastic area compressibility modulus of red cell membrane. *Biophys J*. 16:585-595.

Fomenko, A., C. Neudorfer, R.F. Dallapiazza, S.K. Kalia, and A.M. Lozano. 2018. Low-intensity ultrasound neuromodulation: An overview of mechanisms and emerging human applications. *Brain Stimul*. 11:1209-1217.

Fry, F.J., H.W. Ades, and W.J. Fry. 1958. Production of reversible changes in the central nervous system by ultrasound. *Science*. 127:83-84.

Gaur, U., and B. Wunderlich. 1982. Heat-Capacity and Other Thermodynamic Properties of Linear Macromolecules .5. Polystyrene. *J Phys Chem Ref Data*. 11:313-325.

Gavrilov, L.R., E.M. Tsirulnikov, and I.A. Davies. 1996. Application of focused ultrasound for the stimulation of neural structures. *Ultrasound Med Biol*. 22:179-192.

Giese, K.P., J.F. Storm, D. Reuter, N.B. Fedorov, L.R. Shao, T. Leicher, O. Pongs, and A.J. Silva. 1998. Reduced K+ channel inactivation, spike broadening, and after-hyperpolarization in Kvbeta1.1-deficient mice with impaired learning. *Learn Mem*. 5:257-273.

Gu, N., K. Vervaeke, and J.F. Storm. 2007. BK potassium channels facilitate high-frequency firing and cause early spike frequency adaptation in rat CA1 hippocampal pyramidal cells. *J Physiol*. 580:859-882.

Hand, J.W. 1998. Ultrasound hyperthermia and the prediction of heating. *In* Ultrasound in Medicine. F. Duck, A. Baker and H. Starritt, editors. Institute of Physics Publishing, Philadelphia, PA.

Harper, C.A. 2006. Handbook of plastics technologies: the complete guide to properties and performance. *In*. McGraw-Hill, New York, NY.

Hille, B. 2001. Ion Channels of Excitable Membranes. 3rd

.ed. Sinauer Associates, Inc., Sunderland, MA.

Kanda, H., J. Ling, S. Tonomura, K. Noguchi, S. Matalon, and J.G. Gu. 2019. TREK-1 and TRAAK Are Principal K(+) Channels at the Nodes of Ranvier for Rapid Action Potential Conduction on Mammalian Myelinated Afferent Nerves. *Neuron*. 104:960-971 e967.

Kang, D., C. Choe, and D. Kim. 2005. Thermosensitivity of the two-pore domain K+ channels TREK-2 and TRAAK. *J Physiol*. 564:103-116.

Khaliq, Z.M., and B.P. Bean. 2010. Pacemaking in dopaminergic ventral tegmental area neurons: depolarizing drive from background and voltage-dependent sodium conductances. *J Neurosci*. 30:7401-7413.

Kim, J., D.S. Wei, and D.A. Hoffman. 2005. Kv4 potassium channel subunits control action potential repolarization and frequency-dependent broadening in rat hippocampal CA1 pyramidal neurones. *J Physiol*. 569:41-57.

King, R.L., J.R. Brown, W.T. Newsome, and K.B. Pauly. 2013. Effective parameters for ultrasound-induced in vivo neurostimulation. *Ultrasound Med Biol*. 39:312-331.

Kole, M.H. 2011. First node of Ranvier facilitates high-frequency burst encoding. *Neuron*. 71:671-682.

Krasovitski, B., V. Frenkel, S. Shoham, and E. Kimmel. 2011. Intramembrane cavitation as a unifying mechanism for ultrasound-induced bioeffects. *Proc Natl Acad Sci U S A*. 108:3258-3263.

Kubanek, J., J. Shi, J. Marsh, D. Chen, C. Deng, and J. Cui. 2016. Ultrasound modulates ion channel currents. *Sci Rep*. 6:24170.

Kwok, R., and E. Evans. 1981. Thermoelasticity of large lecithin bilayer vesicles. *Biophys J*. 35:637-652.

Laitko, U., and C.E. Morris. 2004. Membrane tension accelerates rate-limiting voltage-dependent activation and slow inactivation steps in a Shaker channel. *J Gen Physiol*. 123:135-154.

Lee, S.A., H.A.S. Kamimura, M.T. Burgess, and E.E. Konofagou. 2020. Displacement imaging for focused ultrasound peripheral nerve neuromodulation. *IEEE Trans Med Imaging*.

Leighton, T.G. 1998. An introduction to acoustic cavitation. *In* Ultrasound in Medicine. F. Duck, A. Baker and H. Starritt, editors. Institute of Physics Publishing, Philadelphia, PA.

Leterrier, C. 2018. The Axon Initial Segment: An Updated Viewpoint. *J Neurosci*. 38:2135-2145.

Lien, C.C., and P. Jonas. 2003. Kv3 potassium conductance is necessary and kinetically optimized for high-frequency action potential generation in hippocampal interneurons. *J Neurosci*. 23:2058-2068.

Liu, P.W., and B.P. Bean. 2014. Kv2 channel regulation of action potential repolarization and firing patterns in superior cervical ganglion neurons and hippocampal CA1 pyramidal neurons. *J Neurosci*. 34:4991-5002.

Lothet, E.H., K.M. Shaw, H. Lu, J. Zhuo, Y.T. Wang, S. Gu, D.B. Stolz, E.D. Jansen, C.C. Horn, H.J. Chiel, and M.W. Jenkins. 2017. Selective inhibition of small-diameter axons using infrared light. *Sci Rep*. 7:3275.

Madison, D.V., and R.A. Nicoll. 1984. Control of the repetitive discharge of rat CA 1 pyramidal neurones in vitro. *J Physiol*. 354:319-331.

Maingret, F., I. Lauritzen, A.J. Patel, C. Heurteaux, R. Reyes, F. Lesage, M. Lazdunski, and E. Honore. 2000. TREK-1 is a heat-activated background K(+) channel. *EMBO J*. 19:2483-2491.

Malinow, R., and R.W. Tsien. 1990. Presynaptic enhancement shown by whole-cell recordings of long-term potentiation in hippocampal slices. *Nature*. 346:177-180.

Marinc, C., C. Derst, H. Pruss, and R.W. Veh. 2014. Immunocytochemical localization of TASK-3 protein (K2P9.1) in the rat brain. *Cell Mol Neurobiol*. 34:61-70.

Menz, M.D., O. Oralkan, P.T. Khuri-Yakub, and S.A. Baccus. 2013. Precise neural stimulation in the retina using focused ultrasound. *J Neurosci*. 33:4550-4560.

Menz, M.D., P. Ye, K. Firouzi, A. Nikoozadeh, K.B. Pauly, P. Khuri-Yakub, and S.A. Baccus. 2019. Radiation Force as a Physical Mechanism for Ultrasonic Neurostimulation of the Ex Vivo Retina. *J Neurosci*. 39:6251-6264.

Mihran, R.T., F.S. Barnes, and H. Wachtel. 1990. Temporally-specific modification of myelinated axon excitability in vitro following a single ultrasound pulse. *Ultrasound Med Biol*. 16:297-309.

Min, B.K., A. Bystritsky, K.I. Jung, K. Fischer, Y. Zhang, L.S. Maeng, S.I. Park, Y.A. Chung, F.A. Jolesz, and S.S. Yoo. 2011. Focused ultrasound-mediated suppression of chemically-induced acute epileptic EEG activity. *BMC Neurosci*. 12:23.

Morris, C.E., and P.F. Juranka. 2007. Nav channel mechanosensitivity: activation and inactivation accelerate reversibly with stretch. *Biophys J*. 93:822-833.

Needham, D., and R.S. Nunn. 1990. Elastic deformation and failure of lipid bilayer membranes containing cholesterol. *Biophys J*. 58:997-1009.

Owen, S.F., M.H. Liu, and A.C. Kreitzer. 2019. Thermal constraints on in vivo optogenetic manipulations. *Nat Neurosci*. 22:1061-1065.

Pan, Z., T. Kao, Z. Horvath, J. Lemos, J.Y. Sul, S.D. Cranstoun, V. Bennett, S.S. Scherer, and E.C. Cooper. 2006. A common ankyrin-G-based mechanism retains KCNQ and NaV channels at electrically active domains of the axon. *J Neurosci*. 26:2599-2613.

Paris, L., I. Marc, B. Charlot, M. Dumas, J. Valmier, and F. Bardin. 2017. Millisecond infrared laser pulses depolarize and elicit action potentials on in-vitro dorsal root ganglion neurons. *Biomed Opt Express*. 8:4568-4578.

Plaksin, M., E. Kimmel, and S. Shoham. 2016. Cell-Type-Selective Effects of Intramembrane Cavitation as a Unifying Theoretical Framework for Ultrasonic Neuromodulation. *eNeuro*. 3.

Prieto, M.L., K. Firouzi, B.T. Khuri-Yakub, and M. Maduke. 2018. Activation of Piezo1 but Not NaV1.2 Channels by Ultrasound at 43 MHz. *Ultrasound Med Biol*. 44:1217-1232.

Prieto, M.L., O. Oralkan, B.T. Khuri-Yakub, and M.C. Maduke. 2013. Dynamic response of model lipid membranes to ultrasonic radiation force. *PLoS One*. 8:e77115.

Rashid, B., M. Destrade, and M.D. Gilchrist. 2012. Mechanical characterization of brain tissue in compression at dynamic strain rates. *J Mech Behav Biomed Mater*. 10:23-38.

Rashid, B., M. Destrade, and M.D. Gilchrist. 2013. Mechanical characterization of brain tissue in simple shear at dynamic strain rates. *J Mech Behav Biomed Mater*. 28:71-85.

Ricca, B.L., G. Venugopalan, and D.A. Fletcher. 2013. To pull or be pulled: parsing the multiple modes of mechanotransduction. *Curr Opin Cell Biol*. 25:558-564.

Richter, C.P., A.I. Matic, J.D. Wells, E.D. Jansen, and J.T. Walsh, Jr. 2011. Neural stimulation with optical radiation. *Laser Photon Rev*. 5:68-80.

Sarvazyan, A.P., O.V. Rudenko, and W.L. Nyborg. 2010. Biomedical applications of radiation force of ultrasound: historical roots and physical basis. *Ultrasound Med Biol*. 36:1379-1394.

Seeger, H.M., L. Aldrovandi, A. Alessandrini, and P. Facci. 2010. Changes in single K(+) channel behavior induced by a lipid phase transition. *Biophys J*. 99:3675-3683.

Selfridge, A.R. 1985. Approximate Material Properties in Isotropic Materials. *Ieee T Son Ultrason*. 32:381-394.

Shah, M.M., M. Migliore, I. Valencia, E.C. Cooper, and D.A. Brown. 2008. Functional significance of axonal Kv7 channels in hippocampal pyramidal neurons. *Proc Natl Acad Sci U S A*. 105:7869-7874.

Shao, L.R., R. Halvorsrud, L. Borg-Graham, and J.F. Storm. 1999. The role of BK-type Ca2+-dependent K+ channels in spike broadening during repetitive firing in rat hippocampal pyramidal cells. *J Physiol*. 521 Pt 1:135-146.

Shapiro, M.G., K. Homma, S. Villarreal, C.P. Richter, and F. Bezanilla. 2012. Infrared light excites cells by changing their electrical capacitance. *Nat Commun*. 3:736.

Tabarean, I.V., and C.E. Morris. 2002. Membrane stretch accelerates activation and slow inactivation in Shaker channels with S3-S4 linker deletions. *Biophys J*. 82:2982-2994.

Talley, E.M., G. Solorzano, Q. Lei, D. Kim, and D.A. Bayliss. 2001. Cns distribution of members of the two-pore-domain (KCNK) potassium channel family. *J Neurosci*. 21:7491-7505.

Taverna, S., T. Tkatch, A.E. Metz, and M. Martina. 2005. Differential expression of TASK channels between horizontal interneurons and pyramidal cells of rat hippocampus. *J Neurosci*. 25:9162-9170.

Taylor, R.E. 1965. Impedance of the squid axon membrane. *J Cell Comp Physiol*. 66:21-25.

Thijssen, J.M., H.J. Mol, and M.R. Timmer. 1985. Acoustic parameters of ocular tissues. *Ultrasound Med Biol*. 11:157-161.

Tsui, P.H., S.H. Wang, and C.C. Huang. 2005. In vitro effects of ultrasound with different energies on the conduction properties of neural tissue. *Ultrasonics*. 43:560-565.

Tufail, Y., A. Matyushov, N. Baldwin, M.L. Tauchmann, J. Georges, A. Yoshihiro, S.I. Tillery, and W.J. Tyler. 2010. Transcranial pulsed ultrasound stimulates intact brain circuits. *Neuron*. 66:681-694.

Tyler, W.J., Y. Tufail, M. Finsterwald, M.L. Tauchmann, E.J. Olson, and C. Majestic. 2008. Remote excitation of neuronal circuits using low-intensity, low-frequency ultrasound. *PLoS One*. 3:e3511.

Walsh, A.J., G.P. Tolstykh, S. Martens, B.L. Ibey, and H.T. Beier. 2016. Action potential block in neurons by infrared light. *Neurophotonics*. 3:040501.

Wang, J., and O.P. Hamill. 2020. Piezo2, a pressure sensitive channel is expressed in select neurons of the mouse brain: a putative mechanism for synchronizing neural networks by transducing intracranial pressure pulses. *bioRxiv*.

Wells, J., C. Kao, P. Konrad, T. Milner, J. Kim, A. Mahadevan-Jansen, and E.D. Jansen. 2007. Biophysical mechanisms of transient optical stimulation of peripheral nerve. *Biophys J*. 93:2567-2580.

White, S.H., and T.E. Thompson. 1973. Capacitance, area, and thickness variations in thin lipid films. *Biochim Biophys Acta*. 323:7-22.

Wu, J., and W.L. Nyborg. 2008. Ultrasound, cavitation bubbles and their interaction with cells. *Adv Drug Deliv Rev*. 60:1103-1116.

Yamada-Hanff, J., and B.P. Bean. 2015. Activation of Ih and TTX-sensitive sodium current at subthreshold voltages during CA1 pyramidal neuron firing. *J Neurophysiol*. 114:2376-2389.

Ye, P.P., J.R. Brown, and K.B. Pauly. 2016. Frequency Dependence of Ultrasound Neurostimulation in the Mouse Brain. *Ultrasound Med Biol*. 42:1512-1530.

Young, R.R., and E. Henneman. 1961. Functional effects of focused ultrasound on mammalian nerves. *Science*. 134:1521-1522.

Yue, C., and Y. Yaari. 2004. KCNQ/M channels control spike afterdepolarization and burst generation in hippocampal neurons. *J Neurosci*. 24:4614-4624.

Zachs, D.P., S.J. Offutt, R.S. Graham, Y. Kim, J. Mueller, J.L. Auger, N.J. Schuldt, C.R.W. Kaiser, A.P. Heiller, R. Dutta, H. Guo, J.K. Alford, B.A. Binstadt, and H.H. Lim. 2019. Noninvasive ultrasound stimulation of the spleen to treat inflammatory arthritis. *Nat Commun*. 10:951.

Zhou, D., S. Lambert, P.L. Malen, S. Carpenter, L.M. Boland, and V. Bennett. 1998. AnkyrinG is required for clustering of voltage-gated Na channels at axon initial segments and for normal action potential firing. *J Cell Biol*. 143:1295-1304.

Zhu, X., J.W. Lin, and M.Y. Sander. 2019. Infrared inhibition and waveform modulation of action potentials in the crayfish motor axon. *Biomed Opt Express*. 10:6580-6594.

Supplemental Figure 1. Effects of ultrasound at different intensities on firing rate. A. Mean (± SEM) action potential firing rate with (*red*) or without (*blue*) a 1-s ultrasound pulse at various intensities starting 200 ms after the start of a 2-s current step, as measured during the period of overlap between the ultrasound and current stimuli, or during the same time period in the absence of ultrasound. N = 4 cells, except for 6 W/cm^2, where N = 3 cells. **B.** As in panel A, but showing the difference in firing rate between the ultrasound and control conditions.

Supplemental Figure 2. An ATP-regenerating internal solution stabilizes the response to ultrasound. A-B. Mean (±SEM) spike rates during the first 500 ms of a current step in the presence (*solid circles*) and absence (*open circles*) of a 1-s ultrasound application starting 500 ms before the current step, as a function of time relative to break-in (establishment of whole-cell recording configuration). The internal solution contained 10 Na-phosphocreatine with (A) or without (B) 50 U/mL creatine phosphokinase to provide an ATP-regenerating system. Spike rates were measured at various time points between 0 and 10, 10 and 20, 20 and 30, and 30 and 40 minutes after break-in. The x-values represent the mean start time for the protocol to measure spike rates (which comprised 2 minutes of recording time) with horizontal error bars (in some cases smaller than the symbol size) representing the SEM. The amplitude of the current step was adjusted over time to maintain spiking behavior as close as possible to that at the start of the experiment. **C.** Mean (±SEM) difference in spike rate between the control and ultrasound conditions for measurements at 1-10 minutes after break-in and 30-40 minutes after break-in, with (*left*) or without (*right*) the ATP-regenerating system. The difference was only statistically significant without the ATP-regenerating system (P = 0.11, with; and P = 0.0064, without). **D-F.** Same as A-B, but for latency to the first action potential following the start of the current step. N = 10 cells with and N = 6 cells without the ATP-regenerating system. The difference was not statistically significant for either group (P = 0.21, with; and P = 0.089, without).

Supplemental Figure 3. Effects of ultrasound on action potential height. A-C. Mean (±SEM) action potential heights as a function of action potential number in the presence (*red*) and absence (*blue*) of a 1-s ultrasound pulse starting 500 ms before the current step, for cells firing at an average firing rates (as measured during the first 500 ms of the current step) of approximately 5 Hz (N = 13), 10 Hz (N = 15), and 20 Hz (N = 13) in the control condition. **D.** As in A-C, but with ultrasound applied 1 s after the start of a 3-s current step, for cells firing at an average firing rate of approximately 5 Hz in the control condition, with firing rate determined in a 1-s window starting 1 s after the current step (corresponding to the time period of the ultrasound stimulus), and action potential number relative to the start of the ultrasound stimulus (N = 6). Note that the y-axes do not begin at zero.