TITLE

Genome-wide analysis in *Escherichia coli* unravels an unprecedented level of genetic

homoplasy associated with cefotaxime resistance.

Jordy P.M. Coolen[1#], Evert P.M. den Drijver[2,3#], Jaco J. Verweij[3], Jodie A. Schildkraut[1], Kornelia

Neveling[4], Willem J.G. Melchers[1], Eva Kolwijck[1], Heiman F.L. Wertheim[1^], Jan A.J.W.

Kluytmans[2,5,6^], Martijn A. Huynen[7]

[1] Department of Medical Microbiology and Radboudumc Center for Infectious Diseases, Radboud University

Medical Center, Nijmegen, The Netherlands.

[2] Department of Infection Control, Amphia Ziekenhuis, Breda, The Netherlands.

[3] Laboratory for Medical Microbiology and Immunology, Elisabeth-Tweesteden Hospital, Tilburg, The Netherlands.

[4]Department of Human Genetics, Radboud University Medical Center, Nijmegen, The Netherlands.

[5] Laboratory for Microbiology, Microvida, Location Breda, The Netherlands.

[6] Julius Center for Health Sciences and Primary Care, UMCU, Utrecht, The Netherlands.

[7] Centre for Molecular and Biomolecular Informatics, Radboud University Medical Center, Nijmegen, The

Netherlands.

[#]These authors share first authorship

[^]These authors contributed equally

ABSTRACT

Cefotaxime (CTX) is a commonly used third-generation cephalosporin (3GC) to treat infections

caused by *Escherichia coli*. Two genetic mechanisms have been associated with 3GC resistance in

E. coli. The first is the conjugative transfer of a plasmid harboring antibiotic resistance genes. The

second is the introduction of mutations in the promoter region of the *ampC* β-lactamase gene

that cause chromosomal-encoded β-lactamase hyperproduction. A wide variety of promoter

mutations related to AmpC hyperproduction have been described. However, their link to a

specific 3GC such as CTX resistance has not been reported. Here, we measured CTX MICs in 172

cefoxitin resistant *E. coli* isolates and performed genome-wide analysis of homoplastic mutations

associated with CTX resistance by comparing Illumina whole-genome sequencing data of all

isolates to a PacBio tailored-made reference chromosome. We mapped the mutations on the

reference chromosome and determined their occurrence in the phylogeny, revealing extreme

homoplasy at the -42 position of the *ampC* promoter. The 24 occurrences of a "T" at the -42

position rather than the wild type "C", resulted from 18 independent C>T mutations in 5

phylogroups. The -42 C>T mutation was only observed in *E. coli* lacking a plasmid-encoded *ampC*

gene. The association of the -42 C>T mutation with CTX resistance was confirmed to be significant

(FDR < 0.05). To conclude, genome-wide analysis of homoplasy in combination with CTX

resistance identifies the -42 C>T mutation of the *ampC* promotor as significantly associated with

CTX resistance and underline the role of recurrent mutations in the spread of antibiotics

resistance.

Keywords

Escherichia Coli, Genomics, Whole genome sequencing, *ampC,* Bioinformatics

Abbreviations

3GC, third-generation cephalosporin; *campC*, chromosomal-mediated ampC; CAT,

computerized adaptive testing; CTX, cefotaxime; DNA, Deoxyribonucleic acid;EHEC,

enterohemorrhagic *Escherichia Coli*; ESBL, extended-spectrum β-lactamases; FOX, cefoxitin;

gDNA, genomic DNA; MICs, minimal inhibitory concentrations;MLST, multilocus sequence typing;

pampC, plasmid-mediated ampC; PG, peptidoglycan; qRT-PCR, quantitative reverse transcriptase

polymerase chain reaction; SMRT, Single-molecule real-time sequencing; SNP, single-nucleotide

polymorphism; ST, sequence type; UPEC, uropathogene *Escherichia Coli*; WGS, whole genome

sequencing

Impact Statement

In the past decades, the worldwide spread of extended spectrum beta-lactamases (ESBLs) has

led to a substantial increase in the prevalence of resistant common pathogens, thereby

restricting available treatment options. Although acquired resistance genes, e.g. ESBLs, get most

attention, chromosome-encoded resistance mechanisms may play an important role as well. In

E. coli chromosome-encoded β-lactam resistance can be caused by alterations in the promoter

region of the *ampC* gene. To improve our understanding of how frequently these alterations

occur, a comprehensive interpretation of the evolution of these mutations is essential. This study

is the first to apply genome-wide homoplasy analysis to better perceive adaptation of the E. coli

genome to antibiotics. Thereby, this study grants insights into how chromosomal-encoded

antibiotic resistance evolves and, by combining genome-wide association studies with homoplasy

analyses, provides potential strategies for future association studies into the causes of antibiotics

resistance.

Data summary

All data is available under BioProject: PRJNA592140. Raw Illumina sequencing data and metadata

of all 171 *E. coli* isolates used in this study is available from the Sequence Read Archive database

under accession no. SAMN15052485 to SAMN15052655. Full reference chromosome of

ampC_0069 is available via GenBank accession no. CP046396.1 and NCBI Reference Sequence:

NZ_CP046396.1.

Introduction

Escherichia coli is an important pathogen in both community and healthcare-associated

infections [1,2]. In the past decades, a substantial increase in resistance to third-generation

cephalosporin (3GC) antibiotics in *E. coli* has been observed worldwide, mainly caused by the

production of extended-spectrum β-lactamases (ESBL) and AmpC β-lactamases, restricting

available treatment options for common infections [3]. AmpC β-lactamases differ from ESBL as

they not only hydrolyze broad-spectrum penicillins and cephalosporins, but also cephamycins.

Moreover, AmpC β-lactamases are not inhibited by ESBL-inhibitors like clavulanic acid [3], limiting

antibiotic treatment options even further. A widely used screening method for AmpC production

is the use of cefoxitin (FOX) susceptibility, a member of the cephamycins [4].

Although *ampC* β-lactamase genes can be plasmid-encoded (*pampC*), they are also

encoded on the chromosomes of numerous Enterobacterales. *E. coli* naturally carries a

chromosomal-mediated *ampC* (*campC*) gene, but unlike most other Enterobacterales this gene

is non-inducible due to the absence of the *ampR* regulator gene [3]. Consequently, chromosomal

AmpC production in *E. coli* is exclusively regulated by promoter and attenuator mechanisms. This

results in constitutive low-level c*ampC* expression that still allows the use of 3GC antibiotics, such

as cefotaxime (CTX), to treat *E. coli* infections [3]. However, various mutations in the

promoter/attenuator region of *E. coli* may cause constitutive hyperexpression of *campC* [5,6],

thereby increasing the Minimal Inhibitory Concentrations (MICs) for broad-spectrum penicillins

and cephalosporins and limiting appropriate treatment options.

A wide variety of promoter and attenuator mutations have been related to AmpC

hyperproduction [6]. AmpC hyperproduction is primarily caused by alterations of the *ampC*

promoter region, leading to a promoter sequence that more closely resembles the *E. coli*

consensus sigma 70 promoter with a TTGACA -35 box separated by 17 bp from a TATAAT -10 box.

These alterations can be divided into different variants associated with e.g. an alternate displaced

promoter box, a promotor box mutation or an alternate spacer length due to insertions [6].

Furthermore, mutations of the attenuator sequence can lead to changes in the hairpin structure

that strengthen the effect of promoter alterations on AmpC hyperproduction. In a study on

cefoxitin-resistant *E. coli* isolated from Canadian hospitals, Tracz *et al.* described 52 variants of

the promoter and attenuator region [6]. In this study a two-step quantitative reverse

transcriptase (qRT-)PCR was used to determine the effect of promoter/attenuator variants on

ampC expression. Various mutations were related to different delta–delta cycle threshold values

in the RT-PCR and corresponding variations in FOX resistance. An interesting observation that

emerged from this study was that the -32T>A and the -42C>T mutation were the major

alterations that strengthened the *ampC* promoter. Both result in a consensus -35 box. Although

it is known that AmpC hyperproduction leads to FOX resistance as studied by Tracz *et al.,* the

effect of various mutations on resistance to a 3GC antibiotic such as CTX have not been explored.

This is relevant because CTX is commonly used in the treatment of patients with severe *E. coli*

infections, often in combination with selective digestive tract decontamination in Intensive Care

Units [7,8].

While previous research mainly focused on the chromosomal AmpC resistance

mechanism and the impact of AmpC hyperproduction, there is a lack in knowledge and

understanding of the evolutionary origin of these promoter/attenuator variants. Notably, it is

unexplored how the two most prominent promoter mutations -32T>A and -42C>T are distributed

over the *E. coli* phylogeny and therewith how often they occur. More precisely, literature shows

selective pressure can lead to convergent evolution that results in the reoccurrence of a mutation

in multiple isolates independently and in separate lineages [9]. This phenomenon is named

homoplasy [10]. A Consistency Index can be calculated to quantify homoplasy by dividing the

minimum number of changes on the phylogeny by the number of different nucleotides observed

at that site minus one [11], effectively quantifying how often the same mutation occurred in a

phylogenetic tree. One can use the Consistency Index to recognize genomic locations subjected

to homoplasy, and relate the single-nucleotide polymorphism (SNP) positions that are

inconsistent with the phylogeny to antibiotic resistance, as has e.g. been done in multiple studies

on *Mycobacteria* spp [12–15]*.*

In the present study, we hypothesize that some of the mutations in the *ampC*

promoter/attenuator region are homoplastic and are associated with CTX resistance. To test our

hypothesis, we performed genome-wide homoplasy analysis and combined it with a genome-

wide analysis of polymorphisms associated with CTX resistance by constructing a tailored *E. coli*

reference chromosome and combining it with WGS data of 172 FOX resistant *E. coli* isolates

previously collected by our research group [16].

Methods

Isolate selection, DNA isolation, library preparation and DNA sequencing

One hundred seventy-two *Escherichia coli* isolates previously used by our study group [16] were

selected in the present study (see Table S1 in supplemental material). To summarize the method;

DNA isolation was performed as previously described [16], library preparations were performed

using Illumina Nextera XT library preparation kit (Illumina, San Diego, CA, USA), and DNA

sequencing was performed using an Illumina NextSeq 500 (Illumina, San Diego, CA, USA) to

generate 2x 150bp paired-end reads or 2x 300 bp reads on an Illumina MiSeq (Illumina, San Diego,

CA, USA). *De novo* assembly was also performed identical to the method as described in Coolen

et al. 2019 [16] using SPAdes version 3.11.1 [17].

Phylogroup and MLST

Phylogroup stratification was performed using ClermonTyping version 1.4.0 [18]. MLST STs were

derived from the contigs using mlst version 2.5 pubMLST, 31 October 2017 [19,20].

Obtaining the *ampC* promoter/attenuator region

To detect the promoter/attenuator region a custom blast database [21] was created using the

271 bp fragment as described by Peter-Getzlaff *et al.*[22] using *Escherichia coli* K-12 strain ER3413

(accession: CP009789.1) ABRicate version 0.8.9 [23] was used to locate matching regions per

sample and were extracted and converted into multi-fasta format using a custom python script.

Strains were labelled AmpC hyperproducer when promoter mutations were found, as reported

by Caroff *et al.* [24] and Tracz *et al.* [6].

Plasmid-mediated ampC detection

Detection of *pampC* genes was performed by using ABRicate version 0.5 [23] and ResFinder

database (2018-02-16) as described by Coolen *et al.* [16].

PacBio single molecule real-time (SMRT) sequencing of *E. coli* isolate

For PacBio SMRT sequencing, genomic DNA (gDNA) was extracted using the Bacterial gDNA

Isolation Kit (Norgen Biotek Corp., CAN, ON, Thorold). A single *E. coli* isolate was subjected for

DNA shearing using Covaris g-TUBEs (Covaris Inc, US, MA, Woburn) for 30 seconds on 11,000

RPM (*g*). Each DNA sample was separated into two aliquots. Size selection was performed using

a 0.75% agarose cassette and marker S1 on the BluePippin (Sage Science Inc, US, MA, Beverly) to

obtain either 4-8 kb or 4-12 kb DNA fragments. This size selection was chosen to maintain all DNA

fragments including these originating from plasmids (data not used in this study). Library

preparation was performed using the SMRTbell Template prep kit 1.0 (Pacific Biosciences, US,

CA, Menlo Park). For cost-effectiveness, samples were barcoded and pooled with other samples

that are not relevant for this study. Sequencing was conducted using the PacBio Sequel I (Pacific

Biosciences, US, CA, Menlo Park) on a Sequel SMRT Cell 1M v2 (Pacific Biosciences, US, CA, Menlo

Park) with a movie time of 10 h (and 186 min pre-extension time). Subreads per sample were

obtained by extracting the bam files using SMRT Link version 5.1.0.26412 (Pacific Biosciences, US,

CA, Menlo Park).

Chromosomal reconstruction using *de novo* hybrid assembly

To obtain a full-length chromosome, Unicycler version 0.4.7 [25] (settings: --mode bold) was

used, combining Illumina NextSeq 500 2x 150 bp paired-end reads with PacBio Sequel SMRT

subreads. Because unicycler requires fasta reads as input, the subreads in bam format were

converted   to   fasta   by   using   bam2fasta   version   1.1.1   from   pbbioconda

(https://github.com/PacificBiosciences/pbbioconda) prior to *de novo* hybrid assembly. The full

circular chromosome was uploaded to NCBI and annotated using the NCBI Prokaryotic Genome

Annotation Pipeline (PGAP) version 4.10 [26,27].

SNP analysis using *E. coli* reference ampC_0069

Alignment of Illumina reads and SNP calling was performed for all isolates to the reference

chromosome   of   *E.   coli*   isolate   ampC_0069   using   Snippy   version   4.3.6

(https://github.com/tseemann/snippy). A full-length alignment (fullSNP) and a coreSNP

alignment containing SNP positions shared among all isolates were generated by using snippy-

core version 4.3.6 (https://github.com/tseemann/snippy).

Inferring of phylogeny

A phylogenetic tree was inferred by using the coreSNP alignment as input for FastTree(MP)

version 2.1.3 SSE3 (settings: -nt -gtr) [28].

Detection of Homoplasy

The Consistency Index for all nucleotide positions on the chromosome was calculated using

HomoplasyFinder version 0.0.0.9000 [10]. The coreSNP phylogeny was used as true phylogeny

and the Consistency Index was calculated using the multiple sequence alignments fullSNP

alignment as input.

Relate mutations to CTX resistance

To assess if certain mutations were linked to CTX resistance all non-plasmid harboring *ampC E.*

coli isolates were used. CTX resistance was defined using EUCAST guidelines standards of CTX

MIC > 2 mg/L [29]. CTX MIC results were obtained from our previous study [16]. For each

nucleotide position on the reference chromosome the number of Resistant and Sensitive isolates

were counted and tested for adenine vs all other nucleotides, thymine vs all other nucleotides,

cytosine vs all other nucleotides, and guanine vs all other nucleotides creating a contingency

table and performing a Fisher Exact Test in R 3.6.1 [30]. To correct for multiple testing, *P* values

were adjusted using FDR [31].

Selection of genomic positions of interest

By combining previous metrics most relevant genomic positions were selected. Criteria for

selection are, FDR <= 0.05 to CTX and Consistency Index of <= 0.05882353. Annotation of

mutation positions was obtained by using the genome annotation of reference chromosome

(GenBank accession no. CP046396) and applying snpEff (version 4.3t) [32]. The EnteroBase core-

genome MLST and whole-genome MLST schemes were used to distinguish core and accessory

genes [33].

Recombination analysis

Gubbins version 2.4.1. (settings: -f 30) was used to detect recombination regions with coreSNP

alignment and tree as input [34].

Visualization of data

The interactive Tree of Life web-based tool iTOL version 5.3 was used to visualize the

phylogenetic tree [35]. Information about CTX resistance, presence of the *pampC* gene, *campC*

hyperproduction as defined, MLST and phylogroup, as well as alignments of promoter and

attenuator region were incorporated into visualization. The sequence logo of the promoter and

the attenuator alignment were generated using the web-based application Weblogo version 3.7

[36] (http://weblogo.threeplusone.com). A chromosome ideogram of the *E. coli* isolate

ampC_0069 reference chromosome was visualized using CIRCOS software package version 0.69-

8 [37]. Consistency Index scores and significant mutations associated with CTX resistance were

plotted in the ideogram. Gubbins results were displayed by using phandango [38].

Overview of method

A workflow graph of the methods is visualized in Fig 1 using the web-based application yEd Live

version 4.4.2 (https://www.yworks.com/yed-live/).

RESULTS

E. coli **collection**

To study genetic homoplasy events in suspected AmpC producing *Escherichia coli,* FOX MIC > 8

ug/ml and ESBL phenotype negative *E. coli* isolates (*n*=172) were selected as previously described

by Coolen *et al.* [16] (see Table S1 in supplemental material). The entire collection was subjected

to whole-genome sequencing followed by *de novo* assembly of the sequence reads to obtain

contigs.

MLST and phylogroup variants

To access the genetic diversity of our *E. coli* collection, we identified both multi-locus sequence

typing (MLST) and phylogroup variants of each of the 172 *E. coli* isolates. Seventy-five different

sequence types (STs) were identified, of which ST 131 (8.1% *n*=14), ST 38 (7.0% *n*=12), and ST 73

(7.0% *n*=12) were the most prevalent. The sequence types of 13 isolates are unknown.

Phylogroup stratification revealed that the isolates belonged to eight different phylogroups

(Table 1). Phylogroup B1, B2, and D were the most prevalent. One isolate belonged to *Escherichia*

clade IV (st. no. ampC_0128).

ampC **promoter and attenuator variants**

We examined the whole *E. coli* genome. However, we firstly focused on mutations in the *ampC*

promoter and attenuator region. Previously described mutations in the *ampC* promoter region

that according to Tracz *et al.* lead to "hyperproduction" of AmpC were detected in 59 (34.4%) of

the isolates [6]. These isolates were therefore labelled as hyperproducers. Analysis of the

promotor area (-42 to -8) resulted in 21 different variants and the wild type (see Table 1). In the

attenuator region (+17 to +37), 18 different variants were identified (see Table 1). One isolate

(ampC_0128) showed an unusual promoter variant, a four-nucleotide deletion (-45_-42delATCC).

Moreover, an insertion (21_22insG) of unknown function was detected in the attenuator (see

Table S1 in supplemental material).

Plasmid-mediated *ampC* variants

As we aim to associate chromosomal mutations with CTX resistance, differentiation of *pampC*

harboring isolates from non-*pampC* harboring isolates was required. Genomic analysis showed

that 90 (52.3%) of the isolates harbored a *pampC* gene of which bla_{CMY-2} was the most prevalent

(*n*=78). One isolate harbored two different *pampC* genes (bla_{CMY-4} and bla_{DHA-1}) (ampC_0119).

One isolate contained a $bla_{CTX-M-27}$ gene combined with a bla_{CMY-2} gene (ampC_0114) but was ESBL

disc test negative (see Table S1 in supplemental material). In 23 (13.4%) of the isolates neither

pampC nor described mutations related to AmpC hyperproduction were detected and are noted

as low-level AmpC producers.

Tailored reference chromosome

To be able to reconstruct an accurate phylogeny we selected *E. coli* isolate ampC_0069, one of

the strains of the study, to use as reference chromosome for SNP calling. The tailored reference

chromosome was constructed through a hybrid assembly of *n*=4,423,109 2x 150 bp Illumina

NextSeq 500 paired-end reads together with *n*=218,475 PacBio Sequel SMRT subreads (median

5,640 bp). This resulted in a high-quality full circular chromosome of *E. coli* isolate ampC_0069,

with a size of 5,056,572 bp. This isolate belongs to ST648 and contains a plasmid-encoded *bla*$_{CMY-42}$.

The full circular chromosome was uploaded to GenBank accession no. CP046396 and was used

for further analysis. Genome annotation with the NCBI Prokaryotic Genome Annotation Pipeline

(PGAP) identified 4,720 Coding Sequences.

SNP calling

To be able to reconstruct the phylogeny and obtain SNP positions, we mapped reads of all isolates

to the reference chromosome *E. coli* ampC_0069, (accession no. CP046396) resulting in a

coreSNP alignment containing 314,200 variable core SNP positions. For further details per isolate

see Table S2 in supplemental material. To validate our SNP calling method we compared the

ampC_0069 Illumina NextSeq 500 paired-end reads to the reference chromosome of

ampC_0069, resulting in 0 SNPs detected, supporting that the SNP calling data and method

produce no false positives.

Phylogenetic tree based on coreSNP

The coreSNP alignment was used for further analysis. Figure 2 illustrates the approximately

maximum-likelihood phylogenetic tree of all 172 isolates based on the coreSNP alignment. The

tree has a robust topology as indicated by computerized adaptive testing (CAT) likelihood

calculations, resulting in only three positions with a value ≤60% [28]. When focusing on the *ampC*

promoter mutations, they were most prevalent in phylogroups B1, B2, and C, although they were

present in all phylogroups except phylogroup E that lacked mutations in either the promoter or

attenuator region. Interestingly, two positions previously highlighted by Tracz *et al.,* -42 and -32,

are only mutated in the absence of a *pampC* gene, even in isolates with a similar MLST (ST12,

ST88, and ST131). The -42C>T mutation, which results in an alternate displaced promoter box

and therefore leads to increased resistance [6], is present in 24 isolates in 5 distinct phylogroups

and in 17 separate phylogenetic branches, indicating that this mutation is homoplastic.

Additionally, the -32T>A mutation in the promoter, previously also associated to resistance [6],

is present in 20 isolates in 3 distinct phylogroups and in 14 separate phylogenetic branches. To

quantify the level of homoplasy we calculated the Consistency Index.

Genomic homoplastic mutations

We calculated the Consistency Index for all positions on the *E. coli* reference chromosome. A low

Consistency Index value for a position indicates a high degree of inconsistency with the

chromosomal phylogeny and can be calculated by HomoplasyFinder as described in earlier

studies [10,39,40]. As can be observed in Figure 3, results clearly indicate that notwithstanding

multiple other low scoring Consistency Index positions in the promoter and attenuator, position

-42 (4,470,140) and -32 (4,470,150) are the lowest scoring, respectively 0.05882353 and

0.07142857 (see also Fig. S1 in the supplemental material). To access how extreme these

Consistency Index values are, we calculated the Consistency Index for all positions in the

chromosome (see Fig. S2 in the supplemental material). All Consistency Indexes < 1.0 are plotted

in the outer ring (ring A) of Figure 4. Results show that only 9,640 out of 5,056,572 positions

(0.19%) had a Consistency Index ≤ 0.07142857 (see Figure 4 ring A, cutoff is indicated by black

circle). This clearly indicates that positions with low Consistency Indexes are rare, but not unique.

Although these 9,640 positions have a low Consistency Index, we do not yet know their relation

to CTX resistance.

CTX resistance measurements

Cefotaxime MIC measurements from Coolen *et al.* in relation to the genotype of the *E. coli*

isolates are shown in supplementary table 1. Eighty-four of ninety (93.3%) *pampC* harboring *E.*

coli were CTX resistant (MIC >2 mg/L) based on EUCAST clinical breakpoints. Twenty-one of fifty-

nine (35.6%) isolates categorized as hyperproducers based on Trazc *et al.* were CTX resistant,

primarily isolates with the -42 (*n*=15) or -32 mutation (*n*=2). The *pampC* genes never occurred

simultaneously with the -42 or -32 mutations in any of these isolates. One of twenty-three (4.3%)

isolates categorized as a low-level AmpC producer (no *pampC* gene and no known ampC

promoter mutation) tested CTX resistant and contained an insertion in the spacer region (-16_-

15insT), which has not been described by Tracz *et al* [6]. As depicted in Figure 2 the non-*pampC*

strains with a phenotype of CTX > 2 mg/L were present in all phylogroups, although CTX resistant

isolates with the -42 or -32 mutation were predominantly present in phylogroups B1, B2, and C.

Geno- to phenotype

To be able to link *E. coli* chromosomal mutations to CTX resistance we excluded all *E. coli* isolates

with a plasmid containing an *ampC* β-lactamase gene. The association of SNPs to CTX resistance

phenotype (MIC > 2 mg/L) was tested in the remaining 82 isolates using Fisher's Exact Test. After

FDR correction to 0.05, 45,998 significant positions were found (see Figure 4 ring B). Mutation

C>T on position -42 of the *ampC* promoter was found to be significantly associated to CTX

resistance (FDR = 0.034). However, position -32 A>T was not significantly associated to CTX

resistance (FDR = 1).

Homoplasy-based association analysis

Combining the outcome of the homoplasy analysis with the significant CTX resistance associated

positions results in genomic positions associated to CTX resistance that have evolved multiple

times independently. After selecting the lowest scoring Consistency Index positions, ≤

0.05882353, 24 relevant genomic positions were identified that had both a low Consistency Index

and a significant association with CTX resistance. Most notably, one of these 24 positions is

position -42. Only two mutations of those 24 that were located in genes were non-synonymous:

a (conservative) missense mutation in the type II secretion system protein L (*gspL*) gene leading

to Ser330Thr alteration and a mutation in the hydroxyethylthiazole kinase (*thiM*) gene resulting

in a Thr122Ala alteration according to the annotation of *E. coli* strain ampC_0069 (accession no.

CP046396). In addition to the non-synonymous mutation found on the *gspL* gene, eight

synonymous mutations are also located in genes annotated as being part of the type II secretion

system. A complete overview is presented in Table 2.

Recombination analysis

To verify if the level of homoplasy could be a result of recombination, we used Genealogies

Unbiased By recomBinations In Nucleotide Sequences (Gubbins) algorithm to predict

recombination events in our isolate collection [34]. This analysis showed frequent recombination

events in our 172 *E. coli* isolates (see Fig. S3 in the supplemental material). Results illustrate that

recombination blocks cover the region of the *gspL* and the *thiM* gene and their high homoplasy

levels could thus be due to recurrent recombination rather than independent mutations.

Nonetheless, position -42 in the *ampC* promoter is not located in a region effected by

recombination as shown in Fig. S3. Moreover, when inferring the phylogenetic tree corrected for

recombination events as obtained from the Gubbins analysis, the -42C>T mutation actually

occurred in 18 independent branches rather than the 17 branches in the uncorrected tree. This

supports our previous results that this mutation is homoplastic, and not the results of a recurrent

recombination event.

Discussion

We present a genome-wide analysis in which homoplastic mutations are associated with

antibiotic resistance in *E. coli*. By comparing whole-genome sequencing data of 172 *E. coli* isolates

to a tailored reference chromosome we were able to reconstruct the evolution of the genomes

and therewith map recurrent events, allowing us to detect homoplasy associated to CTX

resistance.

Our foremost finding is the significant association of the -42C>T mutation, in the *ampC*

promoter, to CTX resistance that evolved independently at least 17 times in 5 distinct

phylogroups. The -42C>T mutation has been confirmed in former studies to result in AmpC

hyperproduction in *E. coli*. Nelson *et al.* demonstrated an 8 to 18 times increase in activity of

AmpC when cloning the promoter upstream a *lac* operon [41]. Vice versa, Caroff *et al.* found a

decrease in expression of AmpC when cloning the promoter with a -42T>C mutation in a pKK232-

8 reporter plasmid with chloramphenicol acetyltransferase gene [24]. Tracz *et al*. confirmed that

the -42C>T mutation has the strongest effect on the *ampC* promoter, resulting in a high

expression of the *ampC* gene as detected by RT-qPCR [6]. Despite the fact that the -42C>T

mutation has such a strong effect on AmpC production the effect of the mutation on CTX MICs

had not been confirmed. Moreover, the contribution of convergent evolution on this position

relative to the role of the expansion of a clone with a beneficial mutation at this position has not

been determined. That being the case, this study provides evidence that this -42 C>T mutation is

not a result of a recombination event and most likely evolved many times independently.

Remarkably, we observed that the -42C>T mutation never occurs in the presence of a *pampC*

gene (in zero out of twenty-four cases). This was even noticed in isolates with the same MLST,

i.e. ST88 -42C>T (*n*=3) and *pampC* (*n*=1), suggesting preferred exclusivity for one of the resistance

mechanisms. One study mentioned the co-occurrence of the -42C>T mutation and a *pampC* gene

in only one out of thirty-six strains [42]. One could speculate that the exclusivity is a matter of

what arrives first, the plasmid or the mutation, after which there is no selective advantage for

the second mechanism, or that there is actually a fitness cost to having both the mutation and

the plasmid relative to having only the mutation or the plasmid.

The study performed by Tracz *et al.* showed that position -32T>A on the promotor of

ampC associates with AmpC hyperproduction that results in elevated MIC levels for FOX [6].

Surprisingly, in the current study no significant association of -32T>A with CTX resistance was

noticed despite its low Consistency Index. Only two out of twenty isolates with the -32T>A were

CTX resistant, four out of twenty showed an intermediate elevated CTX MIC, and fourteen were

susceptible for CTX. Although we do not know under which conditions this mutation did arise, it

can be speculated that the high level of homoplasy at the -32 position is associated with a

different trait, e.g. resistance against another antibiotic.

Prior studies discovered the importance of mutations in the promoter elements. Random

sequences can even evolve expression comparable to the wild-type promoter elements after only

a single mutation [43]. Furthermore, these promotor elements evolve to only a few forms

indicating convergent evolution [44], as also observed in the present study. All encountered

variants seem to result in a sequence that resembles the *E. coli* consensus sigma 70 promoter

more than the wild type sequence they are derived from [6].

Next to the -42C>T promoter mutation we detected twenty-three other positions in our

analysis that are associated with CTX resistance and have extreme high levels of homoplasy. Most

of these are synonymous mutations, with only two missense mutations (*thiM* and *gspL*) found. It

is remarkable that one missense mutation (p.Ser330Thr) is located in *gspL* that encodes for a

protein of the type II secretion system. The type II secretion system is used by many gram-

negative bacteria to translocate folded proteins from the periplasm, through the outer

membrane, into the extracellular milieu [45]. The system is composed of 12–15 different general

secretory pathway (Gsp) proteins and is related to virulence of various pathogenic *E. coli,* e.g.

EHEC and UPEC [46–48]. It could be that in our selection of mainly clinical samples a certain

predilection has occurred towards isolates with particular virulence traits. The *gspL* gene has

been described as being part of the accessory genome of *E. coli* [49]. Our study supports this

finding as some strains did not harbor this gene. Additionally, we found evidence that

recombination events in the type II secretion system could be the underlying cause of the

extreme homoplasy levels. Still, it is remarkable that missense mutation p.Ser330Thr in the *gspL*

gene correlates with the CTX resistance trait even though it is most likely caused by a

recombination event. To the best of our knowledge no relationship between type II secretion

system and CTX resistance has been observed before. One could hypothesize that the mutation

is a secondary adaptation needed to cope with the elevated AmpC production, as the

peptidoglycan (PG) layer is effected by AmpC hyperproduction and the type II secretion system

contains proteins that are partly localized in the periplasm [50,51].

The use of genomic data to detect homoplasy events is not an uncommon scientific

technique [52–54]. In *Mycobacterium tuberculosis*, it is a well-known method to identify

advantageous mutations, as they are likely to be associated with phenotypes such as drug

resistance, heightened transmissibility, or host adaptation [12–15]. A similar approach was taken

recently by Benjak *et al.* to screen for highly polymorphic genes and genomic regions of

Mycobacterium leprae [55]. Homoplasy-based association analysis limits phylogenetic bias by

correcting for genetic relatedness of strains with the same phenotype, thereby increasing

statistical power to find true associations [14]. Taking this into account, the use of homoplasy-

based association analysis seems viable to relate polymorphic sites to phenotypic traits in

bacteria. Still, studies on other genera than mycobacteria are scarce. To our knowledge, no

homoplasy studies have used this method on *E. coli.*

The increase of 3GC resistance imposes a clinical threat by restricting treatment options

and it is essential to understand the underlying resistance mechanisms. To be able to explore

these mechanisms we selected primarily clinical *E. coli* strains. The current study is directed on

exploring AmpC mediated CTX resistance. Therefore, we included isolates that are already

suspected for increased AmpC production based on elevated FOX resistance. Since a random

sample of *E. coli* would limit finding homoplasy-based associated promoter mutations with CTX

resistance. A downside of these selection criteria might be that we over-estimated certain

genetic variants associated with the trait, as we do not know the frequency of these variants in

the general population. Despite the fact that the spontaneous mutation rate in *E. coli* is relatively

low [56]*,* it is still likely that this particular mutation occurs often in the general population, given

the vast amounts of *E. coli* in the environment [57], providing ample opportunities for adaptation

to antibiotics and arguing for antibiotics of which genomic adaptation requires multiple

mutations in order to develop resistance.

Findings of this study have a number of implications for future practice. This study not

only grants insights into how chromosomal-encoded antibiotic resistance evolves, but also

provides potential strategies for future homoplasy-based association studies. Furthermore, the

use of genome-wide homoplasy-based analysis could be applied to optimize outbreak analysis.

Prior studies have optimized outbreak analysis by removing recombinant regions [58,59].

Homoplasy events disturbs the true phylogeny, hence, removing genomic positions which are

heavily affected by homoplasy could improve tree topology, thereby refining outbreak analysis,

although this strategy is still under debate [60].

Conclusions

To conclude, our method demonstrates extreme levels of homoplasy in *E. coli* that are

significantly associated with CTX resistance. Greater access to WGS data provides new

opportunities to perform large-scale genome-wide analysis. Homoplasy-based methods can have

a potential role in future studies as they constitute an effective strategy to relate phenotypic

traits to variable genomic positions.

Funding information

Not applicable

Author contributions

HW, JK, and MH conceived and supervised the study. JC, ED, EK, JS, JV, WM, and KN performed

the data acquisition. JC and ED performed the data analysis. JC performed bioinformatic

analysis. JC, ED, and MH performed the data interpretation and wrote the manuscript. All

authors read and approved the final manuscript. All authors read and approved the final

manuscript.

Acknowledgements

Special thanks to A. C. J. Soer (Department of Medical Microbiology and Radboudumc Center for

Infectious Diseases, Radboudumc, Nijmegen, the Netherlands), B. A. Lamberts (Department of

Medical Microbiology and Immunology, Rijnstate, Arnhem, the Netherlands) and C. Verhulst

(Department of Infection Control, Amphia Ziekenhuis, Breda, The Netherlands and Laboratory

for Microbiology, Microvida, Location Breda, The Netherlands) for handling the samples on the

lab and creating the Illumina sequence libraries.

M. P. Kwint and R. Derks (Department of Human Genetics, Radboudumc, Nijmegen, the

Netherlands) for helping with SMRT sequencing on the PacBio Sequel I.

Many thanks to M. Janssens (Laboratory for Medical Microbiology and Immunology, Elisabeth-

Tweesteden Hospital, Tilburg, the Netherlands), S. Van Leest (Laboratory for Microbiology,

Microvida, location Bravis Hospital, the Netherlands), K. T. Veldman and D. J. Mevius (department

of Bacteriology and Epidemiology, Wageningen Bioveterinary Research, Lelystad, the

Netherlands, E.A. Reuland (Medical Microbiology and Infection Control, Amsterdam UMC

location VUmc, Amsterdam, the Netherlands), W. H. F. Goessens (Erasmus University Medical

Center, Rotterdam, Netherlands), R. W. Bosboom (Department of Medical Microbiology and

Immunology, Rijnstate, Arnhem, the Netherlands) and P. Vos (Check-Points, Wageningen, the

Netherlands) for providing samples of which some are included in this study.

Conflicts of interest

The authors declare that there are no conflicts of interest.

References

1. Weinstein RA, Gaynes R, Edwards JR. Overview of Nosocomial Infections Caused by Gram-

Negative Bacilli. Clin Infect Dis. 2005;41:848–54.

2. Pitout JDD. Extraintestinal pathogenic Escherichia coli: A combination of virulence with

antibiotic resistance. Front. Microbiol. 2012.

3. Jacoby GA. AmpC B-Lactamases. Clin Microbiol Rev [Internet]. 2009;22:161–82. Available

from: http://cmr.asm.org/cgi/doi/10.1128/CMR.00036-08

4. Martinez- L, Simonsen GS. EUCAST_detection_of_resistance_mechanisms_170711. 2017;1–

43. Available from:

http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Resistance_mechanisms/EUC

AST_detection_of_resistance_mechanisms_170711.pdf.

5. Tracz DM, Boyd DA, Bryden L, Hizon R, Giercke S, Caeseele P V., et al. Increase in ampC

promoter strength due to mutations and deletion of the attenuator in a clinical isolate of

cefoxitin-resistant Escherichia coli as determined by RT-PCR. J Antimicrob Chemother.

2005;55:768–72.

6. Tracz DM, Boyd DA, Hizon R, Bryce E, McGeer A, Ofner-Agostini M, et al. ampC gene

expression in promoter mutants of cefoxitin-resistant Escherichia coli clinical isolates. FEMS

Microbiol Lett [Internet]. 2007;270:265–71. Available from:

https://academic.oup.com/femsle/article-lookup/doi/10.1111/j.1574-6968.2007.00672.x

7. De Smet AMGA, Kluytmans JAJW, Cooper BS, Mascini EM, Benus RFJ, Van Der Werf TS, et al.

Decontamination of the digestive tract and oropharynx in ICU patients. N Engl J Med [Internet].

Massachussetts Medical Society; 2009 [cited 2020 May 14];360:20–31. Available from:

http://www.nejm.org/doi/abs/10.1056/NEJMoa0800394

8. Aardema H, Bult W, Van Hateren K, Dieperink W, Touw DJ, Alffenaar JWC, et al. Continuous

versus intermittent infusion of cefotaxime in critically ill patients: A randomized controlled trial

comparing plasma concentrations. J Antimicrob Chemother [Internet]. 2020 [cited 2020 May

14];75:441–8. Available from: https://academic.oup.com/jac/article/75/2/441/5614359

9. Wake DB. Homoplasy: the result of natural selection, or evidence of design limitations? Am

Nat. 1991;138:543–67.

10. Crispell J, Balaz D, Gordon SV. Homoplasyfinder: A simple tool to identify homoplasies on a

phylogeny. Microb Genomics [Internet]. 2019 [cited 2019 Aug 7];5. Available from:

http://www.danielwilson.me.uk/

11. Kluge AG, Farris JS. Quantitative phyletics and the evolution of anurans. Syst Biol.

1969;18:1–32.

12. Farhat MR, Shapiro BJ, Kieser KJ, Sultana R, Jacobson KR, Victor TC, et al. Genomic analysis

identifies targets of convergent positive selection in drug-resistant Mycobacterium

tuberculosis. Nat Genet. 2013;45:1183–9.

13. Mortimer TD, Weber AM, Pepperell CS. Signatures of Selection at Drug Resistance Loci in

Mycobacterium tuberculosis. Gilbert JA, editor. mSystems [Internet]. 2018;3:1–9. Available

from: https://msystems.asm.org/lookup/doi/10.1128/mSystems.00108-17

14. Ruesen C, Chaidir L, van Laarhoven A, Dian S, Ganiem AR, Nebenzahl-Guimaraes H, et al.

Large-scale genomic analysis shows association between homoplastic genetic variation in

Mycobacterium tuberculosis genes and meningeal or pulmonary tuberculosis. BMC Genomics.

BioMed Central Ltd.; 2018;19.

15. Miotto P, Cabibbe AM, Feuerriegel S, Casali N, Drobniewski F, Rodionova Y, et al.

Mycobacterium tuberculosis pyrazinamide resistance determinants: A multicenter study. Nacy

CA, editor. MBio [Internet]. 2014;5:1–10. Available from:

https://mbio.asm.org/lookup/doi/10.1128/mBio.01819-14

16. Coolen JPM, Den Drijver EPM, Kluytmans JAJW, Verweij JJ, Lamberts BA, Soer JACJ, et al.

Development of an algorithm to discriminate between plasmid- and chromosomal-mediated

AmpC β-lactamase production in Escherichia coli by elaborate phenotypic and genotypic

characterization. J Antimicrob Chemother [Internet]. 2019;74:3481–8. Available from:

https://academic.oup.com/jac/advance-article/doi/10.1093/jac/dkz362/5554444

17. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A new

genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol

[Internet]. 2012;19:455–77. Available from:

http://www.liebertpub.com/doi/10.1089/cmb.2012.0021

18. Beghain J, Bridier-Nahmias A, Nagard H Le, Denamur E, Clermont O. ClermonTyping: An

easy-to-use and accurate in silico method for Escherichia genus strain phylotyping. Microb

Genomics. 2018;4:1–8.

19. Seemann T. mlst. Github https://github.com/tseemann/mlst.

20. Jolley KA, Maiden MCJ. BIGSdb: Scalable analysis of bacterial genome variation at the

population level. BMC Bioinformatics. 2010;11.

21. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol

Biol [Internet]. 1990;215:403–10. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0022283605803602

22. Peter-Getzlaff S, Polsfuss S, Poledica M, Hombach M, Giger J, Böttger EC, et al. Detection of

AmpC beta-lactamase in Escherichia coli: Comparison of three phenotypic confirmation assays

and genetic analysis. J Clin Microbiol [Internet]. 2011;49:2924–32. Available from:

http://jcm.asm.org/cgi/doi/10.1128/JCM.00091-11

23. Seemann T. Abricate [Internet]. Github; Available from:

https://github.com/tseemann/abricate

24. Caroff N, Espaze E, Gautreau D, Richet H, Reynaud A. Analysis of the effects of -42 and -32

ampC promoter mutations in clinical isolates of Escherichia coli hyperproducing AmpC. J

Antimicrob Chemother. 2000;45:783–8.

25. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome assemblies

from short and long sequencing reads. Phillippy AM, editor. PLoS Comput Biol [Internet].

2017;13:e1005595. Available from: https://dx.plos.org/10.1371/journal.pcbi.1005595

26. Tatusova T, Dicuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, et al. NCBI

prokaryotic genome annotation pipeline. Nucleic Acids Res [Internet]. 2016;44:6614–24.

Available from: https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gkw569

27. Haft DH, DiCuccio M, Badretdin A, Brover V, Chetvernin V, O'Neill K, et al. RefSeq: An

update on prokaryotic genome annotation and curation. Nucleic Acids Res [Internet].

2018;46:D851–60. Available from: http://academic.oup.com/nar/article/46/D1/D851/4588110

28. Price MN, Dehal PS, Arkin AP. FastTree 2 - Approximately maximum-likelihood trees for

large alignments. Poon AFY, editor. PLoS One [Internet]. 2010;5:e9490. Available from:

https://dx.plos.org/10.1371/journal.pone.0009490

29. European Committee on Antimicrobial Susceptibility Testing (EUCAST). The European

Committee on Antimicrobial Susceptibility Testing. Breakpoint tables for interpretation of MICs

and zone diameters. Version 10.0, 2020 [Internet]. 2020. Available from:

http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_10.0_Bre

akpoint_Tables.pdf

30. Mehta CR, Patel NR. A network algorithm for performing fisher's exact test in r × c

contingency tables. J Am Stat Assoc [Internet]. Taylor & Francis; 1983;78:427–34. Available

from: https://doi.org/10.1080/01621459.1983.10477989

31. Benjamini Y, Hochberg Y. <Benjamini&Hochberg1995_FDR.pdf>. J R Stat Soc Ser B

[Internet]. 1995;57:289–300. Available from: http://www.jstor.org/stable/2346101

32. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, et al. A program for annotating

and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of

Drosophila melanogaster strain w1118; iso-2; iso-3. Fly (Austin). 2012;6:80–92.

33. Zhou Z, Alikhan NF, Mohamed K, Fan Y, Achtman M. The EnteroBase user's guide, with case

studies on Salmonella transmissions, Yersinia pestis phylogeny, and Escherichia core genomic

diversity. Genome Res. 2020;30:138–52.

34. Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, et al. Rapid

phylogenetic analysis of large samples of recombinant bacterial whole genome sequences using

Gubbins. Nucleic Acids Res [Internet]. 2015;43:e15. Available from:

http://academic.oup.com/nar/article/43/3/e15/2410982/Rapid-phylogenetic-analysis-of-large-

samples-of

35. Letunic I, Bork P. Interactive Tree of Life (iTOL) v4: Recent updates and new developments.

Nucleic Acids Res [Internet]. 2019;47:W256–9. Available from:

https://academic.oup.com/nar/advance-article/doi/10.1093/nar/gkz239/5424068

36. Crooks GE, Hon G, Chandonia JM, Brenner SE. WebLogo: A sequence logo generator.

Genome Res [Internet]. 2004;14:1188–90. Available from:

ftp://ftp.ncbi.nih.gov/genomes/Bacteria

37. Connors J, Krzywinski M, Schein J, Gascoyne R, Horsman D, Jones SJ, et al. Circos : An

information aesthetic for comparative genomics. Genome Res. 2009;19:1639–45.

38. Hadfield J, Croucher NJ, Goater RJ, Abudahab K, Aanensen DM, Harris SR. Phandango: An

interactive viewer for bacterial population genomics. Kelso J, editor. Bioinformatics [Internet].

2018;34:292–3. Available from:

https://academic.oup.com/bioinformatics/article/34/2/292/4212949

39. Crispell J, Benton CH, Balaz D, De Maio N, Ahkmetova A, Allen A, et al. Combining genomics

and epidemiology to analyse bi-directional transmission of mycobacterium bovis in a multi-host

system. Elife. 2019;8:1–36.

40. Van Dorp L, Gelabert P, Rieux A, De Manuel M, De-Dios T, Gopalakrishnan S, et al.

Plasmodium vivax Malaria Viewed through the Lens of an Eradicated European Strain. Mol Biol

Evol [Internet]. 2020;37:773–85. Available from:

https://www.biorxiv.org/content/10.1101/736702v1

41. Nelson EC, Gay Elisha B. Molecular basis of ampC hyperproduction in clinical isolates of

Escherichia coli [Internet]. Antimicrob. Agents Chemother. 1999. Available from:

http://aac.asm.org/

42. Mulvey MR, Bryce E, Boyd DA, Ofner-Agostini M, Land AM, Simor AE, et al. Molecular

characterization of cefoxitin-resistant Escherichia coli from Canadian hospitals. Antimicrob

Agents Chemother. 2005;49:358–65.

43. Yona AH, Alm EJ, Gore J. Random sequences rapidly evolve into de novo promoters. Nat

Commun [Internet]. 2018;9:1530. Available from: http://www.nature.com/articles/s41467-018-

04026-w

44. Liu S, Libchaber A. Some aspects of E. coli promoter evolution observed in a molecular

evolution experiment. J Mol Evol. 2006;62:536–50.

45. Korotkov K V., Sandkvist M, Hol WGJ. The type II secretion system: Biogenesis, molecular

architecture and mechanism. Nat. Rev. Microbiol. 2012. p. 336–51.

46. Ho TD, Davis BM, Ritchie JM, Waldor MK. Type 2 secretion promotes enterohemorrhagic

Escherichia coli adherence and intestinal colonization. Infect Immun. 2008;76:1858–65.

47. Baldi DL, Higginson EE, Hocking DM, Praszkier J, Cavaliere R, James CE, et al. The type II

secretion system and its ubiquitous lipoprotein substrate, SslE, are required for biofilm

formation and virulence of enteropathogenic Escherichia coli. Infect Immun. 2012;80:2042–52.

48. Kulkarni R, Dhakal BK, Slechta ES, Kurtz Z, Mulvey MA, Thanassi DG. Roles of putative type II

secretion and type IV pilus systems in the virulence of uropathogenic Escherichia coli. PLoS One.

2009;4.

49. Dunne KA, Chaudhuri RR, Rossiter AE, Beriotto I, Browning DF, Squire D, et al. Sequencing a

piece of history: Complete genome sequence of the original Escherichia coli strain. Microb

Genomics. 2017;3.

50. Vanderlinde EM, Strozen TG, Hernández SB, Cava F, Howard SP. Alterations in peptidoglycan

cross-linking suppress the secretin assembly defect caused by mutation of GspA in the type II

secretion system. J Bacteriol [Internet]. 2017 [cited 2020 Mar 25];199. Available from:

http://jb.asm.org/

51. Juan C, Torrens G, Barceló IM, Oliver A. Interplay between Peptidoglycan Biology and

Virulence in Gram-Negative Pathogens. Microbiol Mol Biol Rev [Internet]. 2018 [cited 2020 Mar

25];82. Available from: http://mmbr.asm.org/

52. Read TD, Massey RC. Characterizing the genetic basis of bacterial phenotypes using

genome-wide association studies: A new direction for bacteriology. Genome Med. BioMed

Central Ltd.; 2014.

53. Chen PE, Shapiro BJ. The advent of genome-wide association studies for bacteria. Curr.

Opin. Microbiol. Elsevier Ltd; 2015. p. 17–24.

54. Shapiro BJ, David LA, Friedman J, Alm EJ. Looking for Darwin's footprints in the microbial

world. Trends Microbiol. 2009;17:196–204.

55. Benjak A, Avanzi C, Singh P, Loiseau C, Girma S, Busso P, et al. Phylogenomics and

antimicrobial resistance of the leprosy bacillus Mycobacterium leprae. Nat Commun. Nature

Publishing Group; 2018;9.

56. Lee H, Popodi E, Tang H, Foster PL. Rate and molecular spectrum of spontaneous mutations

in the bacterium Escherichia coli as determined by whole-genome sequencing. Proc Natl Acad

Sci U S A. 2012;109.

57. Tenaillon O, Skurnik D, Picard B, Denamur E. The population genetics of commensal

Escherichia coli. Nat Rev Microbiol [Internet]. Nature Publishing Group; 2010;8:207–17.

Available from: http://dx.doi.org/10.1038/nrmicro2298

58. Escobar-Páramo P, Sabbagh A, Darlu P, Pradillon O, Vaury C, Denamur E, et al. Decreasing

the effects of horizontal gene transfer on bacterial phylogeny: The Escherichia coli case study.

Mol Phylogenet Evol [Internet]. Academic Press Inc.; 2004;30:243–50. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S1055790303001817

59. Price LB, Johnson JR, Aziz M, Clabots C, Johnston B, Tchesnokova V, et al. The epidemic of

extended-spectrum-β-lactamase-producing Escherichia coli ST131 is driven by a single highly

pathogenic subclone, H30-Rx. Parkhill J, editor. MBio [Internet]. 2013;4:1–10. Available from:

https://mbio.asm.org/lookup/doi/10.1128/mBio.00377-13

60. Hedge J, Wilson DJ. Bacterial phylogenetic reconstruction from whole genomes is robust to

recombination but demographic inference is not. Vidaver AK, editor. MBio [Internet]. American

Society for Microbiology; 2014;5. Available from:

https://mbio.asm.org/lookup/doi/10.1128/mBio.02158-14

LEGENDS

FIG 1 Schematic of workflow used to perform the homoplasy-based association analysis.

Starting from the top A) the *de novo* assembly of the NextSeq/MiSeq reads and B) the hybrid

assembly of the reference chromosome ampC_069. On the left side C) the alignment of

promoter/attenuator region. In the middle D) the coreSNP analysis for the phylogeny used in E)

the homoplasy analysis combined with F) the fullSNP data on the right, which was also used for

G) the statistics (Fisher Exact & FDR) to relate cefotaxime (CTX) resistance to SNP positions. H)

Inferring recombination events using Gubbins.

FIG 2   Approximately maximum-likelihood phylogenetic tree of all 172 *E. coli* isolates based on

the coreSNP alignment with the resistance for cefotaxime (CTX), *pampC* gene presence, MLSTs,

phylogroups, and the alignments of the promoter and the attenuator region. Positions with a CAT

likelihood score ≤60% are indicated as red dots.

FIG 3   Sequence logo with probability score for the promoter and the attenuator region. The

Consistency Index and the minimum number of changes on the tree per position are represented

below the sequence logos.

FIG 4   Circos plot for the full chromosome of ampC_0069 (accession no. CP046396) with per

position the various metrics used. A) The blue colored ring represents the Consistency Index

results per genomic position. The two red dots indicate the -42 and -32 position on the promoter.

The black circle line indicates the 0.07142857 Consistency Index value. B) The ring with a red

background shows all positions that were significantly associated to cefotaxime (CTX) resistance

in all non-*pampC* harboring *E. coli* isolates. Larger bars pointing outwards indicate multiple

significant associated positions in a small genomic region. C) The ring with the green background

shows all 24 positions that have a low Consistency Index of ≤ 0.05882353 and are significantly

associated with CTX resistance in all non-*pampC* harboring *E. coli* isolates.

TABLE 1   Table of the distribution of AmpC promoter and attenuator variants as well as the

amount of different MLST and phylogroups per grouped genotype (*pampC*, hyperproducers and

low-level AmpC producers).

TABLE 2   The *n*=24 positions with a significant association with cefotaxime resistance

(FDR ≤0.05) and with a consistency index ≤ 0.05882353.

FIG S1   Violin plots of the log10 Consistency Indexes of the promoter and attenuator.

FIG S2   Distribution of the log10 Consistency Indexes of all genomic position based on the *E. coli*

ampC_0069 reference chromosome, compared to the log10 Consistency Indexes of the

promoter and attenuator region.

FIG S3   Recombination events inferred from all 172 *E. coli* isolates by Gubbins displayed along the

approximately maximum-likelihood phylogenetic tree based on the coreSNP alignment.

Phylogroups are depicted as in FIG 2. Gubbins blocks are colored red if they are ancestral, and

blue if they only affect one isolate. The line graph represents the recombination prevalence along

the sequence. The 24 positions with a significant association with cefotaxime resistance (FDR

≤0.05) and a consistency index ≤ 0.05882353 are indicated on the top of the figure. The two

missense mutations and *ampC* promoter region are displayed in blue.

Table S1   Classification of n=172 *E. coli* isolates in the three genotypes (*pampC*,

hyperproducer, low-level AmpC producers), with the results of the MLST and phylogroups

stratification and the different mutations in the promoter and attenuator per isolate. Isolates

with a CTX MIC >2 mg/L without a confirmed *pampC* gene are depicted in **bold**.

Table S2   SNP analysis for n=172 *E. coli* isolates according to snippy statistics.

FIG 1 Schematic of workflow used to perform the homoplasy-based association analysis. Starting from the top A) the *de novo* assembly of the NextSeq/MiSeq reads and B) the hybrid assembly of the reference chromosome ampC_069. On the left side C) the alignment of promoter/attenuator region. In the middle D) the coreSNP analysis for the phylogeny used in E) the homoplasy analysis combined with F) the fullSNP data on the right, which was also used for G) the statistics (Fisher Exact & FDR) to relate cefotaxime (CTX) resistance to SNP positions. H) Inferring recombination events using Gubbins.

FIG 2 Approximately maximum-likelihood phylogenetic tree of all 172 *E. coli* isolates based on the coreSNP alignment with the resistance for cefotaxime (CTX), *pampC* gene presence, MLSTs, phylogroups, and the alignments of the promoter and the attenuator region. Positions with a CAT likelihood score ≤60% are indicated as red dots.

FIG 3   Sequencelogo with probability score for the promoter and the attenuator region. The Consistency Index and the minimum number of changes on the tree per position are represented below the sequence logos.

FIG 4 Circos plot for the full chromosome of ampC_0069 (accession no. CP046396) with per position the various metrics used. A) The blue colored ring represents the Consistency Index results per genomic position. The two red dots indicate the -42 and -32 position on the promoter. The black circle line indicates the 0.07142857 Consistency Index value. B) The ring with a red background shows all positions that were significantly associated to cefotaxime (CTX) resistance in all non-*pampC* harboring *E. coli* isolates. Larger bars pointing outwards indicate multiple significant associated positions in a small genomic region. C) The ring with the green background shows all 24 positions that have a low Consistency Index of ≤ 0.05882353 and are significantly associated with CTX resistance in all non-*pampC* harboring *E. coli* isolates.

TABLE 1

Table of the distribution of AmpC promoter and attenuator variants as well as the amount of different MLST and phylogroups per grouped genotype (*pampC*, hyperproducers and low-level AmpC producers)

	Isolates	Promoter variants	Attenuator variants	MLST	Phylogroups
pampC	*n*=90	*n*=3	*n*=6	44 STs & 4 unknown	A (11.1%), B1 (13.3%), B2 (27.8%), C (2.2%), D (31.1%), E (3.3%), F (11.1%)
Hyperproducers	*n*=59	*n*=12	*n*=14	30 STs & 5 unknown	A (1.7%), B1 (30.5%), B2 (50.9%), C (8.5%), D (5.1%), F (1.7%), clade IV (1.7%)
Low-level AmpC producers	*n*=23	*n*=10	*n*=5	14 STs & 4 unknown	A (4.4%), B1 (13.0%), B2 (39.1%), C (8.7%), D (30.4%), E (4.4%)
Total	*n*=172	*n*=22	*n*=18	75 STs and 13 unknown	A (7.0%), B1 (19.2%), B2 (37.2%), C (5.2%), D (22.1%), E (2.3%), F (6.4%), clade IV (0.6%)

TABLE 2 The *n*=24 positions with a significant association with cefotaxime resistance (FDR ≤0.05) and with a consistency index ≤ 0.05882353.

Genomic position	Consistency Index	A FDR	C FDR	G FDR	T FDR	Counts ACGT	# strains	Min No Changes on Tree	Gene name	Enterobase core / accessory gene	Variant	HGVS annotation	Genomic start position	Genomic stop position	Product	Locus_tag
810581	0.05555556	0.06837015	1.00000000	0.03945241	1.00000000	41:0:122:0	163	18	*glcE*	accessory	synonymous		810312	811364	Glycolate oxidase subunit glcE	GNX12_03825
810791	0.05882353	1.00000000	1.00000000	0.04099492	0.03471297	0:0:54:109	163	17	*glcE*	accessory	synonymous		810312	811364	Glycolate oxidase subunit glcE	GNX12_03825
815680	0.05555556	1.00000000	0.14069566	1.00000000	0.04226791	0:69:0:93	162	18	*glcA*	not available	synonymous		815555	817237	Glycolate permease glcA	GNX12_03845
824522	0.05555556	1.00000000	0.03648231	1.00000000	0.14069566	0:82:0:73	155	18	*gspC*	accessory	synonymous		823719	824678	Type II secretion system protein gspC	GNX12_03865
828695	0.05555556	0.04226791	1.00000000	0.03394030	1.00000000	79:0:74:0	153	18	*gspF*	accessory	synonymous		828261	829484	Type II secretion system protein gspF	GNX12_03880
830684	0.05555556	1.00000000	0.03394030	1.00000000	0.03648231	0:80:0:73	153	18	*gspI*	accessory	synonymous		830520	830891	Type II secretion system protein gspI	GNX12_03895
830708	0.05263158	0.04099492	1.00000000	0.03648231	1.00000000	68:0:85:0	153	19	*gspI*	accessory	synonymous		830520	830891	Type II secretion system protein gspI	GNX12_03895
830732	0.05882353	1.00000000	0.03648231	1.00000000	0.04099492	0:82:0:71	153	17	*gspI*	accessory	synonymous		830520	830891	Type II secretion system protein gspI	GNX12_03895
831564	0.05882353	0.03648231	1.00000000	0.08198761	1.00000000	71:0:83:0	154	17	*gspK*	accessory	synonymous		831490	832467	General secretion pathway protein gspK	GNX12_03905
832152	0.05555556	1.00000000	0.03394030	1.00000000	0.05525251	0:86:0:72	158	18	*gspK*	accessory	synonymous		831490	832467	General secretion pathway protein gspK	GNX12_03905
832287	0.05882353	0.05525251	1.00000000	1.00000000	0.03394030	81:0:0:79	160	17	*gspK*	accessory	synonymous		831490	832467	General secretion pathway protein gspK	GNX12_03905
833451	0.05000000	0.04099492	1.00000000	1.00000000	0.03394030	64:0:0:98	162	20	*gspL*	accessory	missense	p.Ser330Thr	832464	833642	Type II secretion system protein gspL	GNX12_03910
843887	0.05263158	0.03648231	1.00000000	1.00000000	1.00000000	64:0:40:0	104	19	unnamed	accessory	synonymous		842822	844849	Capsular polysaccharide biosynthesis protein	GNX12_03950
1863004	0.05555556	0.03648231	1.00000000	0.03648231	1.00000000	126:0:46:0	172	18	*thiM*	core	missense	p.Thr122Ala	1862641	1863429	Hydroxyethylthiazole kinase	GNX12_08695
1911551	0.05882353	1.00000000	0.03648231	1.00000000	0.45554848	0:116:0:44	160	17	unnamed	accessory	synonymous		1910691	1911830	Polysaccharide export protein Wza	GNX12_08905
1946016	0.04347826	0.67473250	1.00000000	0.04099492	1.00000000	58:0:91:0	149	23	*ugd*	accessory	synonymous		1944883	1946049	UDP-glucose 6-dehydrogenase	GNX12_09060
1946067	0.04545455	1.00000000	1.00000000	1.00000000	0.04962540	0:0:65:84	149	22	non-coding region	not available	upstream		1946049	1946196	None	GNX12_09065
1946072	0.04545455	0.04962540	1.00000000	1.00000000	1.00000000	84:0:0:65	149	22	non-coding region	not available	upstream		1946049	1946196	None	GNX12_09065
1952745	0.03571429	0.05907658	1.00000000	0.03732584	1.00000000	131:0:40:0	171	28	*hisD*	accessory	synonymous		1952169	1953473	Histidinol dehydrogenase	GNX12_09100
2051214	0.05263158	0.04099492	1.00000000	1.00000000	0.09170046	137:0:0:34	171	19	unnamed	accessory	synonymous		2050846	2052204	putative sensor-like heavy metal sensor histidine kinase	GNX12_09550
2051220	0.05263158	0.04099492	1.00000000	1.00000000	0.09170046	137:0:0:34	171	19	unnamed	accessory	synonymous		2050846	2052204	heavy metal sensor histidine kinase	GNX12_09550
2057518	0.05263158	0.04099492	1.00000000	0.03648231	1.00000000	111:0:60:0	171	19	*dcm*	core	synonymous		2056604	2058022	DNA-cytosine methyltransferase	GNX12_09575
2068593	0.05882353	0.03732584	1.00000000	0.11254604	1.00000000	45:0:123:0	168	17	*fliM*	accessory	synonymous		2067810	2068814	Flagellar motor switch protein fliM	GNX12_09650
4470140	0.05882353	1.00000000	0.03394030	1.00000000	0.03394030	0:147:0:24	171	17	*ampC* promoter	core	upstream		4470140	4470174	None	GNX12_21360